



**HAL**  
open science

# GENERALIZED $k$ -FRACTIONAL CONFORMABLE INTEGRALS AND RELATED INEQUALITIES

Siddra Habib, Shahid Mubeen, Muhammad Nawaz Naeem, Feng Qi

► **To cite this version:**

Siddra Habib, Shahid Mubeen, Muhammad Nawaz Naeem, Feng Qi. GENERALIZED  $k$ -FRACTIONAL CONFORMABLE INTEGRALS AND RELATED INEQUALITIES: Generalized  $k$ -fractional conformable integrals. 2018. hal-01788916v1

**HAL Id: hal-01788916**

**<https://hal.science/hal-01788916v1>**

Preprint submitted on 9 May 2018 (v1), last revised 19 Jan 2020 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# GENERALIZED $k$ -FRACTIONAL CONFORMABLE INTEGRALS AND RELATED INEQUALITIES

SIDDRA HABIB, SHAHID MUBEEN, MUHAMMAD NAWAZ NAEEM, AND FENG QI

ABSTRACT. In the paper, the authors introduce the generalized  $k$ -fractional conformable integrals, which are the  $k$ -analogues of the recently introduced fractional conformable integrals and can be reduced to other fractional integrals under specific values of the parameters involved, prove the existence of  $k$ -fractional conformable integrals, and generalize some integral inequalities to ones for generalized  $k$ -fractional conformable integrals.

## 1. INTRODUCTION

Fractional calculus is the study of derivatives and integrals of non-integer order and is the generalized form of classical derivatives and integrals. It is as dated as classical calculus, but it acquires more importance in recent two decades, this is due to its applications in various fields such as physics, biology, fluid dynamics, control theory, image processing, signal processing, and computer networking. See [5, 6, 7, 8, 9, 10, 11, 12, 15, 16, 21, 22, 24, 45, 50, 51, 52, 53]. In recent years, the research has been proceeded to generalize the existing inequalities through innovative ideas and approaches of fractional calculus. One of the trendiest approaches among researchers is the use of fractional integral operators. Due to their potentials to be expended for the existence of nontrivial and positive solutions of several classes of fractional differential equations, the integral inequalities involving fractional integrals are considerably important.

A large bulk of existing literature consists of generalizations of numerous inequalities via fractional integral operators and their applications [3, 27, 32, 46, 49]. Mubeen and Iqbal [28] contributed the ongoing research by presenting the improved version of generalized Grüss type integral inequalities for  $k$ -Riemann–Liouville fractional integrals. Agarwal et al. [2] obtained certain Hermite–Hadamard type inequalities for generalized  $k$ -fractional integrals. Set et al. [39] presented an integral identity and generalized Hermite–Hadamard type inequalities for Riemann–Liouville fractional integral. Mubeen et al. [29] established integral inequalities of Ostrowski type for  $k$ -fractional Riemann–Liouville integrals. Sarikaya and Budak [37] utilized local fractional integrals to derive a generalized inequality. Khan et al. [25] produced some important generalized inequalities for a finite class of positive decreasing functions for fractional conformable integrals. Jleli et al. [18] determined a Hartman–Winter type inequality involving fractional derivative with

---

2010 *Mathematics Subject Classification.* Primary 26A33; Secondary 26D10, 35A23, 47A63.

*Key words and phrases.* left  $k$ -fractional conformable integral; right  $k$ -fractional conformable integral; fractional integral inequality.

This paper was typeset using  $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$ .

respect to another function. In the papers [43, 44, 48] and closely related references therein, there are more information on this topic.

The main object of this paper is to develop a new notion ‘‘generalized  $k$ -fractional conformable integral’’ which is the generalized form of fractional operators reported in [17]. Hereafter, we also generalize some integral inequalities given in [25] for a finite class of positive and decreasing functions to ones involving our newly introduced  $k$ -fractional conformable integrals. For details of those inequalities, their applications, and their stability, we refer readers to [23, 26, 41, 42].

## 2. NOTATIONS

The notion of left and right fractional conformable derivatives for a differentiable function  $f$ , introduced by Abdeljawad [1], can be expressed as

$$\mathfrak{T}_{a+}^{\alpha} f(\mathcal{T}) = (\mathcal{T} - a)^{1-\alpha} f'(\mathcal{T}) \quad \text{and} \quad \mathfrak{T}_{b-}^{\alpha} f(\mathcal{T}) = (b - \mathcal{T})^{1-\alpha} f'(\mathcal{T}).$$

Correspondingly, left and right fractional conformable integrals for  $0 < \alpha < 1$  can be represented by

$$\mathcal{H}_{a+}^{\alpha} f(\mathcal{T}) = \int_a^{\mathcal{T}} \frac{f(x)}{(x-a)^{1-\alpha}} dx \quad \text{and} \quad \mathcal{H}_{b-}^{\alpha} f(\mathcal{T}) = \int_{\mathcal{T}}^b \frac{f(x)}{(b-x)^{1-\alpha}} dx.$$

Let  $\Gamma(z)$  for  $\Re(z) > 0$  denote the classical gamma function. The left and right fractional conformable integral (LFCI and RFCI) operators of order  $\beta \in \mathbb{C}$  for  $\Re(\beta) > 0$  can be defined [17] respectively by

$${}^{\beta}\mathcal{H}_{a+}^{\alpha} f(x) = \frac{1}{\Gamma(\beta)} \int_a^x \left[ \frac{(x-a)^{\alpha} - (t-a)^{\alpha}}{\alpha} \right]^{\beta-1} \frac{f(t)}{(t-a)^{1-\alpha}} dt$$

and

$${}^{\beta}\mathcal{H}_{b-}^{\alpha} f(x) = \frac{1}{\Gamma(\beta)} \int_x^b \left[ \frac{(b-x)^{\alpha} - (b-t)^{\alpha}}{\alpha} \right]^{\beta-1} \frac{f(t)}{(b-t)^{1-\alpha}} dt.$$

Díaz and Pariguan [13] generalized the classical Pochhammer symbol  $(\lambda)_n$ , the classical gamma function  $\Gamma(z)$ , and the classical beta function  $B(u, v)$  respectively as

$$(\lambda)_{n,k} = \begin{cases} 1, & n = 0; \\ \lambda(\lambda+k) \cdots (\lambda+(n-1)k), & n \in \mathbb{N}, \end{cases}$$

$$\Gamma_k(x) = \lim_{n \rightarrow \infty} \frac{n!k^n (nk)^{x/k-1}}{(x)_{n,k}},$$

and

$$B_k(u, v) = \frac{1}{k} \int_0^1 t^{u/k-1} (1-t)^{v/k-1} dt.$$

See also [30, 31, 33, 34, 35]. It is not difficult to see that the  $k$ -gamma function  $\Gamma_k(x)$  and the  $k$ -beta function  $B_k(u, v)$  satisfy

$$\Gamma_k(x) = \int_0^{\infty} u^{x-1} e^{-u^k/k} du, \quad \Gamma(x) = \lim_{k \rightarrow 1} \Gamma_k(x),$$

$$\Gamma_k(x) = k^{x/k-1} \Gamma\left(\frac{x}{k}\right), \quad \Gamma_k(x+k) = x\Gamma_k(x),$$

and

$$B_k(u, v) = \frac{1}{k} B\left(\frac{u}{k}, \frac{v}{k}\right), \quad B_k(u, v) = \frac{\Gamma_k(u)\Gamma_k(v)}{\Gamma_k(u+v)}.$$

3. GENERALIZED  $k$ -FRACTIONAL CONFORMABLE INTEGRALS

In this section, we introduce the generalized left and right  $k$ -fractional conformable integrals which generalize the Riemann–Liouville fractional integrals [36, p. 44], Hadamard fractional integrals [4], Katugampola fractional integrals [19], and generalized fractional integrals [38].

**Definition 3.1.** Let  $f$  be a continuous function on a finite real interval  $[a, b]$ . Then the generalized left and right  $k$ -fractional conformable integrals ( $k$ -FCI) of order  $\beta \in \mathbb{C}$  for  $\Re(\beta) > 0$  are respectively defined as

$${}^{\beta}_k\mathcal{H}_{a^+}^{\alpha}f(x) = \frac{1}{k\Gamma_k(\beta)} \int_a^x \left[ \frac{(x-a)^{\alpha} - (t-a)^{\alpha}}{\alpha} \right]^{\beta/k-1} \frac{f(t)}{(t-a)^{1-\alpha}} dt$$

and

$${}^{\beta}_k\mathcal{H}_{b^-}^{\alpha}f(x) = \frac{1}{k\Gamma_k(\beta)} \int_x^b \left[ \frac{(b-x)^{\alpha} - (b-t)^{\alpha}}{\alpha} \right]^{\beta/k-1} \frac{f(t)}{(b-t)^{1-\alpha}} dt,$$

where  $k > 0$  and  $\alpha \in \mathbb{R} \setminus \{0\}$ .

**Theorem 3.1.** Let  $f \in L_1[a, b]$ ,  $\alpha \in \mathbb{R} \setminus \{0\}$ , and  $k > 0$ . Then both  ${}^{\beta}_k\mathcal{H}_{a^+}^{\alpha}f(x)$  and  ${}^{\beta}_k\mathcal{H}_{b^-}^{\alpha}f(x)$  exist for all  $x \in [a, b]$  and  $\Re(\beta) > 0$ .

*Proof.* Let  $\Delta' = [a, b] \times [a, b]$  and  $P' : \Delta' \rightarrow \mathbb{R}$  such that

$$P'(x, t) = [(x-a)^{\alpha} - (t-a)^{\alpha}]^{\beta/k-1} (t-a)^{\alpha-1}.$$

It is clear that  $P' = P'_+ + P'_-$ , where

$$P'_+(x, t) = \begin{cases} [(x-a)^{\alpha} - (t-a)^{\alpha}]^{\beta/k-1} (t-a)^{\alpha-1}, & a \leq t \leq x \leq b; \\ 0, & a \leq x \leq t \leq b \end{cases}$$

and

$$P'_-(x, t) = \begin{cases} [(t-a)^{\alpha} - (x-a)^{\alpha}]^{\beta/k-1} (x-a)^{\alpha-1}, & a \leq t \leq x \leq b; \\ 0, & a \leq x \leq t \leq b. \end{cases}$$

Since  $P'$  is measurable on  $\Delta'$ , we can write

$$\begin{aligned} \int_a^b P'(x, t) dt &= \int_a^x P'(x, t) dt \\ &= \int_a^x [(x-a)^{\alpha} - (t-a)^{\alpha}]^{\beta/k-1} (t-a)^{\alpha-1} dt = \frac{\alpha k}{\beta} (x-a)^{\alpha\beta/k}. \end{aligned}$$

Therefore, we obtain

$$\begin{aligned} \int_a^b \left[ \int_a^b P'(x, t) |f(x)| dt \right] dx &= \int_a^b |f(x)| \left[ \int_a^b P'(x, t) dt \right] dx \\ &= \frac{\alpha k}{\beta} \int_a^b (x-a)^{\alpha\beta/k} |f(x)| dx \leq \frac{\alpha k}{\beta} (b-a)^{\alpha\beta/k} \int_a^b |f(x)| dx \\ &= \frac{\alpha k}{\beta} (b-a)^{\alpha\beta/k} \|f(x)\|_{L_1[a, b]} < \infty. \end{aligned}$$

So, by Tonelli's theorem for iterated integrals [14, p. 147], the function  $Q' : \Delta' \rightarrow \mathbb{R}$  such that  $Q'(x, t) = P'(x, t)f(x)$  is integrable over  $\Delta'$ . Hence, by Fubini's theorem,

it follows that  $\int_a^b P'(x, t)f(x) dx$  is integrable over  $[a, b]$  as a function of  $t \in [a, b]$ . This implies that  ${}_{k}^{\beta}\mathcal{H}_{a+}^{\alpha}f(x)$  exists.

The existence of the right  $k$ -fractional conformable integral  ${}_{k}^{\beta}\mathcal{H}_{b-}^{\alpha}f(x)$  can be proved in a similar manner. The proof of Theorem 3.1 is complete.  $\square$

#### 4. INEQUALITIES FOR GENERALIZED $k$ -FRACTIONAL CONFORMABLE INTEGRALS

Fractional integral inequalities have been analyzed for many useful purposes. One of the most useful applications of such inequalities is the existence of nontrivial solutions of fractional differential equations. Many applications find in the literature for the existence of nontrivial solution eigenvalue problems by inequalities, see [32, 49]. Generalizing pre-existing inequalities by applying fractional integral operators is the most popular trend in the research field nowadays.

In this section, we present some  $k$ -analogues of inequalities in [40, 46, 47] for generalized  $k$ -fractional conformable integrals.

**Theorem 4.1.** *Let  $\{g_i, 1 \leq i \leq n\}$  be a sequence of continuous positive decreasing functions on the interval  $[a, b]$ . Let  $a < x \leq b$ ,  $\eta > 0$ ,  $\xi \geq \gamma_p > 0$  for  $1 \leq p \leq n$ . Then the left  $k$ -FCI operator  ${}_{k}^{\beta}\mathcal{H}_{a+}^{\alpha}$  satisfies*

$$\frac{{}_{k}^{\beta}\mathcal{H}_{a+}^{\alpha}(\prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x))}{{}_{k}^{\beta}\mathcal{H}_{a+}^{\alpha}(\prod_{i=1}^n g_i^{\gamma_i}(x))} \geq \frac{{}_{k}^{\beta}\mathcal{H}_{a+}^{\alpha}((x-a)^{\eta} \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x))}{{}_{k}^{\beta}\mathcal{H}_{a+}^{\alpha}((x-a)^{\eta} \prod_{i=1}^n g_i^{\gamma_i}(x))}. \quad (4.1)$$

*Proof.* Under given conditions, we have

$$[(\rho - a)^{\eta} - (\tau - a)^{\eta}][g_p^{\xi - \gamma_p}(\tau) - g_p^{\xi - \gamma_p}(\rho)] \geq 0.$$

Let us define a function

$$\begin{aligned} {}_{k}^{\beta}\mathfrak{J}_{a+}^{\alpha}(x, \rho, \tau) &= \frac{1}{k\Gamma_k(\beta)} \left[ \frac{(x-a)^{\alpha} - (\tau-a)^{\alpha}}{\alpha} \right]^{\beta/k-1} \\ &\times \frac{\prod_{i=1}^n g_i^{\gamma_i}(\tau)}{(\tau-a)^{1-\alpha}} [(\rho-a)^{\eta} - (\tau-a)^{\eta}][g_p^{\xi - \gamma_p}(\tau) - g_p^{\xi - \gamma_p}(\rho)]. \end{aligned} \quad (4.2)$$

Under given assumptions, the function  ${}_{k}^{\beta}\mathfrak{J}_{a+}^{\alpha}(x, \rho, \tau)$  is positive for all  $\tau \in (a, b]$ . Integrating on both sides of the above equation (4.2) with respect to  $\tau$  from  $a$  to  $x$  gives

$$\begin{aligned} 0 &\leq \int_a^x {}_{k}^{\beta}\mathfrak{J}_{a+}^{\alpha}(x, \rho, \tau) d\tau = \frac{1}{k\Gamma_k(\beta)} \int_a^x \left[ \frac{(x-a)^{\alpha} - (\tau-a)^{\alpha}}{\alpha} \right]^{\beta/k-1} \\ &\times \prod_{i=1}^n g_i^{\gamma_i}(\tau) [(\rho-a)^{\eta} - (\tau-a)^{\eta}][g_p^{\xi - \gamma_p}(\tau) - g_p^{\xi - \gamma_p}(\rho)] \frac{d\tau}{(\tau-a)^{1-\alpha}} \\ &= (\rho-a)^{\eta} \left[ {}_{k}^{\beta}\mathcal{H}_{a+}^{\alpha} \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] + g_p^{\xi - \gamma_p}(\rho) \left[ {}_{k}^{\beta}\mathcal{H}_{a+}^{\alpha} \left( (x-a)^{\eta} \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \\ &- (\rho-a)^{\eta} g_p^{\xi - \gamma_p}(\rho) \left[ {}_{k}^{\beta}\mathcal{H}_{a+}^{\alpha} \left( \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] - \left[ {}_{k}^{\beta}\mathcal{H}_{a+}^{\alpha} \left( (x-a)^{\eta} \prod_{i \neq p}^n g_i^{\gamma_i}(x) \right) \right]. \end{aligned} \quad (4.3)$$

Multiplying the relation (4.3) by

$$\frac{1}{k\Gamma_k(\beta)} \left[ \frac{(x-a)^\alpha - (\rho-a)^\alpha}{\alpha} \right]^{\beta/k-1} \frac{\prod_{i=1}^n g_i^{\gamma_i}(\rho)}{(\rho-a)^{1-\alpha}} \quad (4.4)$$

and integrating on both sides with respect to  $\rho$  from  $a$  to  $x$  yield

$$0 \leq \left[ {}_k^{\beta} \mathcal{H}_{a^+}^{\alpha} \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \left[ {}_k^{\beta} \mathcal{H}_{a^+}^{\alpha} \left( (x-a)^\eta \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \\ - \left[ {}_k^{\beta} \mathcal{H}_{a^+}^{\alpha} \left( (x-a)^\eta \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \left[ {}_k^{\beta} \mathcal{H}_{a^+}^{\alpha} \left( \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right]. \quad (4.5)$$

Dividing on both sides of relation (4.5) by

$$\left[ {}_k^{\beta} \mathcal{H}_{a^+}^{\alpha} \left( (x-a)^\eta \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \left[ {}_k^{\beta} \mathcal{H}_{a^+}^{\alpha} \left( \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right]$$

results in (4.1). The proof of Theorem 4.1 is complete.  $\square$

**Corollary 4.1.** *Let  $\{g_i, 1 \leq i \leq n\}$  be a sequence of continuous positive decreasing functions on the interval  $[a, b]$ . Let  $a < x \leq b$ ,  $\eta > 0$ ,  $\xi \geq \gamma_p > 0$  for  $1 \leq p \leq n$ . Then the left  $k$ -FCI operator  ${}_k^{\beta} \mathcal{H}_{a^+}^{\alpha}$  satisfies*

$$\left[ {}_k^{\theta} \mathcal{H}_{a^+}^{\alpha} \left( (x-a)^\eta \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \left[ {}_k^{\beta} \mathcal{H}_{a^+}^{\alpha} \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \\ + \left[ {}_k^{\theta} \mathcal{H}_{a^+}^{\alpha} \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \left[ {}_k^{\beta} \mathcal{H}_{a^+}^{\alpha} \left( (x-a)^\eta \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \\ \geq \left[ {}_k^{\theta} \mathcal{H}_{a^+}^{\alpha} \left( (x-a)^\eta \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \left[ {}_k^{\beta} \mathcal{H}_{a^+}^{\alpha} \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \\ + \left[ {}_k^{\theta} \mathcal{H}_{a^+}^{\alpha} \left( \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \left[ {}_k^{\beta} \mathcal{H}_{a^+}^{\alpha} \left( (x-a)^\eta \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right]. \quad (4.6)$$

*Proof.* Multiplying on both sides of the relation (4.3) by

$$\frac{1}{k\Gamma_k(\theta)} \left[ \frac{(x-a)^\alpha - (\rho-a)^\alpha}{\alpha} \right]^{\theta/k-1} \frac{\prod_{i=1}^n g_i^{\gamma_i}(\rho)}{(\rho-a)^{1-\alpha}} \quad (4.7)$$

and integrating on both sides with respect to  $\rho$  from  $a$  to  $x$  arrive at

$$0 \leq \left[ {}_k^{\theta} \mathcal{H}_{a^+}^{\alpha} \left( (x-a)^\eta \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \left[ {}_k^{\beta} \mathcal{H}_{a^+}^{\alpha} \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \\ + \left[ {}_k^{\theta} \mathcal{H}_{a^+}^{\alpha} \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \left[ {}_k^{\beta} \mathcal{H}_{a^+}^{\alpha} \left( (x-a)^\eta \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \\ - \left[ {}_k^{\theta} \mathcal{H}_{a^+}^{\alpha} \left( (x-a)^\eta \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \left[ {}_k^{\beta} \mathcal{H}_{a^+}^{\alpha} \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right]$$

$$- \left[ \theta {}_k\mathcal{H}_{a^+}^\alpha \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \left[ \beta {}_k\mathcal{H}_{a^+}^\alpha \left( (x-a)^\eta \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right]. \quad (4.8)$$

Dividing on both sides of (4.8) by

$$\begin{aligned} & \left[ \theta {}_k\mathcal{H}_{a^+}^\alpha \left( (x-a)^\eta \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \left[ \beta {}_k\mathcal{H}_{a^+}^\alpha \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \\ & + \left[ \theta {}_k\mathcal{H}_{a^+}^\alpha \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \left[ \beta {}_k\mathcal{H}_{a^+}^\alpha \left( (x-a)^\eta \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \end{aligned}$$

leads to (4.6). The proof of Corollary 4.1 is complete.  $\square$

**Theorem 4.2.** *Let  $h(x)$  be a continuous increasing function and  $\{g_i, 1 \leq i \leq n\}$  be a sequence of continuous positive decreasing functions on the interval  $[a, b]$ . Let  $a < x \leq b$ ,  $\eta > 0$ ,  $\xi \geq \gamma_p > 0$  for  $1 \leq p \leq n$ . Then the left  $k$ -FCI operator  ${}^\beta_k\mathcal{H}_{a^+}^\alpha$  satisfies*

$$\frac{{}^\beta_k\mathcal{H}_{a^+}^\alpha \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right)}{{}^\beta_k\mathcal{H}_{a^+}^\alpha \left( \prod_{i=1}^n g_i^{\gamma_i}(x) \right)} \geq \frac{{}^\beta_k\mathcal{H}_{a^+}^\alpha \left( h^\eta(x) \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right)}{{}^\beta_k\mathcal{H}_{a^+}^\alpha \left( h^\eta(x) \prod_{i=1}^n g_i^{\gamma_i}(x) \right)}. \quad (4.9)$$

*Proof.* Under given conditions, we have

$$[h^\eta(\rho) - h^\eta(\tau)] [g_p^{\xi-\gamma_p}(\tau) - g_p^{\xi-\gamma_p}(\rho)] \geq 0.$$

Let us define a function

$$\begin{aligned} {}^\beta_k\mathcal{J}_{a^+}^\alpha(x, \rho, \tau) &= \frac{1}{k\Gamma_k(\beta)} \left[ \frac{(x-a)^\alpha - (\tau-a)^\alpha}{\alpha} \right]^{\beta/k-1} \\ &\times \frac{\prod_{i=1}^n g_i^{\gamma_i}(\tau)}{(\tau-a)^{1-\alpha}} [h^\eta(\rho) - h^\eta(\tau)] [g_p^{\xi-\gamma_p}(\tau) - g_p^{\xi-\gamma_p}(\rho)]. \quad (4.10) \end{aligned}$$

Accordingly, the function  ${}^\beta_k\mathcal{J}_{a^+}^\alpha(x, \rho, \tau)$  is positive for all  $\tau \in (a, b]$ . Integrating on both sides of the above equation (4.10) with respect to  $\tau$  from  $a$  to  $x$  shows

$$\begin{aligned} 0 &\leq \int_a^x {}^\beta_k\mathcal{J}_{a^+}^\alpha(x, \rho, \tau) d\tau = \frac{1}{k\Gamma_k(\beta)} \int_a^x \left[ \frac{(x-a)^\alpha - (\tau-a)^\alpha}{\alpha} \right]^{\beta/k-1} \\ &\quad \times \prod_{i=1}^n g_i^{\gamma_i}(\tau) [h^\eta(\rho) - h^\eta(\tau)] [g_p^{\xi-\gamma_p}(\tau) - g_p^{\xi-\gamma_p}(\rho)] \frac{d\tau}{(\tau-a)^{1-\alpha}} \\ &= h^\eta(\rho) \left[ \beta {}_k\mathcal{H}_{a^+}^\alpha \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] + g_p^{\xi-\gamma_p}(\rho) \left[ \beta {}_k\mathcal{H}_{a^+}^\alpha \left( h^\eta(x) \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \\ &\quad - h^\eta(\rho) g_p^{\xi-\gamma_p}(\rho) \left[ \beta {}_k\mathcal{H}_{a^+}^\alpha \left( \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] - \left[ \beta {}_k\mathcal{H}_{a^+}^\alpha \left( h^\eta(x) \prod_{i \neq p}^n g_i^{\gamma_i}(x) \right) \right]. \quad (4.11) \end{aligned}$$

Multiplying on both sides of the relation (4.11) by (4.4) and integrating on both sides with respect to  $\rho$  from  $a$  to  $x$  give

$$0 \leq \left[ \beta {}_k\mathcal{H}_{a^+}^\alpha \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \left[ \beta {}_k\mathcal{H}_{a^+}^\alpha \left( h^\eta(x) \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right]$$

$$- \left[ {}_{k}^{\beta} \mathcal{H}_{a^{+}}^{\alpha} \left( h^{\eta}(x) \prod_{i \neq p}^{n} g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \left[ {}_{k}^{\beta} \mathcal{H}_{a^{+}}^{\alpha} \left( \prod_{i=1}^{n} g_i^{\gamma_i}(x) \right) \right]. \quad (4.12)$$

Dividing on both sides of (4.12) by

$${}_{k}^{\beta} \mathcal{H}_{a^{+}}^{\alpha} \left( h^{\eta}(x) \prod_{i=1}^{n} g_i^{\gamma_i}(x) \right) \left[ {}_{k}^{\beta} \mathcal{H}_{a^{+}}^{\alpha} \left( \prod_{i=1}^{n} g_i^{\gamma_i}(x) \right) \right]$$

leads to (4.9). The proof of Theorem 4.2 is complete.  $\square$

**Corollary 4.2.** *Let  $h(x)$  be a continuous increasing function and  $\{g_i, 1 \leq i \leq n\}$  be a sequence of continuous positive decreasing functions on the interval  $[a, b]$ . Let  $a < x < b$ ,  $\eta > 0$ ,  $\xi \geq \gamma_p > 0$  for  $1 \leq p \leq n$ . Then the left  $k$ -FCI operator  ${}_{k}^{\beta} \mathcal{H}_{a^{+}}^{\alpha}$  satisfies*

$$\begin{aligned} & \left[ {}_{k}^{\theta} \mathcal{H}_{a^{+}}^{\alpha} \left( h^{\eta}(x) \prod_{i=1}^{n} g_i^{\gamma_i}(x) \right) \right] \left[ {}_{k}^{\beta} \mathcal{H}_{a^{+}}^{\alpha} \left( \prod_{i \neq p}^{n} g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \\ & + \left[ {}_{k}^{\theta} \mathcal{H}_{a^{+}}^{\alpha} \left( \prod_{i \neq p}^{n} g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \left[ {}_{k}^{\beta} \mathcal{H}_{a^{+}}^{\alpha} \left( h^{\eta}(x) \prod_{i=1}^{n} g_i^{\gamma_i}(x) \right) \right] \\ & \geq \left[ {}_{k}^{\theta} \mathcal{H}_{a^{+}}^{\alpha} \left( h^{\eta}(x) \prod_{i \neq p}^{n} g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \left[ {}_{k}^{\beta} \mathcal{H}_{a^{+}}^{\alpha} \left( \prod_{i \neq p}^{n} g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \\ & + \left[ {}_{k}^{\theta} \mathcal{H}_{a^{+}}^{\alpha} \left( \prod_{i=1}^{n} g_i^{\gamma_i}(x) \right) \right] \left[ {}_{k}^{\beta} \mathcal{H}_{a^{+}}^{\alpha} \left( h^{\eta}(x) \prod_{i=1}^{n} g_i^{\gamma_i}(x) \right) \right]. \quad (4.13) \end{aligned}$$

*Proof.* Multiplying on both sides of the relation (4.11) by (4.7) and integrating on both sides with respect to  $\rho$  from  $a$  to  $x$  derive

$$\begin{aligned} 0 & \leq \left[ {}_{k}^{\theta} \mathcal{H}_{a^{+}}^{\alpha} \left( h^{\eta}(x) \prod_{i=1}^{n} g_i^{\gamma_i}(x) \right) \right] \left[ {}_{k}^{\beta} \mathcal{H}_{a^{+}}^{\alpha} \left( \prod_{i \neq p}^{n} g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \\ & + \left[ {}_{k}^{\theta} \mathcal{H}_{a^{+}}^{\alpha} \left( \prod_{i \neq p}^{n} g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \left[ {}_{k}^{\beta} \mathcal{H}_{a^{+}}^{\alpha} \left( h^{\eta}(x) \prod_{i=1}^{n} g_i^{\gamma_i}(x) \right) \right] \\ & - \left[ {}_{k}^{\theta} \mathcal{H}_{a^{+}}^{\alpha} \left( h^{\eta}(x) \prod_{i \neq p}^{n} g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \left[ {}_{k}^{\beta} \mathcal{H}_{a^{+}}^{\alpha} \left( \prod_{i \neq p}^{n} g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \\ & - \left[ {}_{k}^{\theta} \mathcal{H}_{a^{+}}^{\alpha} \left( \prod_{i \neq p}^{n} g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \left[ {}_{k}^{\beta} \mathcal{H}_{a^{+}}^{\alpha} \left( h^{\eta}(x) \prod_{i=1}^{n} g_i^{\gamma_i}(x) \right) \right]. \quad (4.14) \end{aligned}$$

Dividing on both sides of (4.14) by

$$\begin{aligned} & \left[ {}_{k}^{\theta} \mathcal{H}_{a^{+}}^{\alpha} \left( h^{\eta}(x) \prod_{i \neq p}^{n} g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \left[ {}_{k}^{\beta} \mathcal{H}_{a^{+}}^{\alpha} \left( \prod_{i \neq p}^{n} g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \\ & + \left[ {}_{k}^{\theta} \mathcal{H}_{a^{+}}^{\alpha} \left( \prod_{i \neq p}^{n} g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \left[ {}_{k}^{\beta} \mathcal{H}_{a^{+}}^{\alpha} \left( h^{\eta}(x) \prod_{i=1}^{n} g_i^{\gamma_i}(x) \right) \right] \end{aligned}$$

reveals (4.13). The proof of Corollary 4.2 is complete.  $\square$



**Theorem 4.3.** *Let  $\{g_i, 1 \leq i \leq n\}$  be a sequence of continuous positive decreasing functions on the interval  $[a, b]$ . Let  $a < x \leq b$ ,  $\eta > 0$ ,  $\xi \geq \gamma_p > 0$  for  $1 \leq p \leq n$ . Then the right  $k$ -FCI operator  ${}^\beta_k\mathcal{H}_{b-}^\alpha$  satisfies*

$$\frac{{}^\beta_k\mathcal{H}_{b-}^\alpha\left(\prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x)\right)}{{}^\beta_k\mathcal{H}_{b-}^\alpha\left(\prod_{i=1}^n g_i^{\gamma_i}(x)\right)} \geq \frac{{}^\beta_k\mathcal{H}_{b-}^\alpha\left((b-x)^\eta \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x)\right)}{{}^\beta_k\mathcal{H}_{b-}^\alpha\left((b-x)^\eta \prod_{i=1}^n g_i^{\gamma_i}(x)\right)}. \quad (4.15)$$

*Proof.* Under given conditions, we have

$$[(b-\rho)^\eta - (b-\tau)^\eta][g_p^{\xi-\gamma_p}(\tau) - g_p^{\xi-\gamma_p}(\rho)] \geq 0.$$

Let us define a function

$$\begin{aligned} {}^\beta_k\mathfrak{J}_{b-}^\alpha(x, \rho, \tau) &= \frac{1}{k\Gamma_k(\beta)} \left[ \frac{(b-x)^\alpha - (b-\tau)^\alpha}{\alpha} \right]^{\beta/k-1} \\ &\quad \times \frac{\prod_{i=1}^n g_i^{\gamma_i}(\tau)}{(b-\tau)^{1-\alpha}} [(b-\rho)^\eta - (b-\tau)^\eta][g_p^{\xi-\gamma_p}(\tau) - g_p^{\xi-\gamma_p}(\rho)]. \end{aligned} \quad (4.16)$$

Consequently, the function  ${}^\beta_k\mathfrak{J}_{b-}^\alpha(x, \rho, \tau)$  is positive for all  $\tau \in (a, b]$ . Integrating on both sides of the above equation (4.16) with respect to  $\tau$  from  $x$  to  $b$  gives

$$\begin{aligned} 0 &\leq \int_x^b {}^\beta_k\mathfrak{J}_{b-}^\alpha(x, \rho, \tau) d\tau = \frac{1}{k\Gamma_k(\beta)} \int_x^b \left[ \frac{(b-x)^\alpha - (b-\tau)^\alpha}{\alpha} \right]^{\beta/k-1} \\ &\quad \times \prod_{i=1}^n g_i^{\gamma_i}(\tau) [(b-\rho)^\eta - (b-\tau)^\eta][g_p^{\xi-\gamma_p}(\tau) - g_p^{\xi-\gamma_p}(\rho)] \frac{d\tau}{(b-\tau)^{1-\alpha}} \\ &= (b-\rho)^\eta \left[ {}^\beta_k\mathcal{H}_{b-}^\alpha\left(\prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x)\right) \right] + g_p^{\xi-\gamma_p}(\rho) \left[ {}^\beta_k\mathcal{H}_{b-}^\alpha\left((b-x)^\eta \prod_{i=1}^n g_i^{\gamma_i}(x)\right) \right] \\ &\quad - (b-\rho)^\eta g_p^{\xi-\gamma_p}(\rho) \left[ {}^\beta_k\mathcal{H}_{b-}^\alpha\left(\prod_{i=1}^n g_i^{\gamma_i}(x)\right) \right] - \left[ {}^\beta_k\mathcal{H}_{b-}^\alpha\left((b-x)^\eta \prod_{i \neq p}^n g_i^{\gamma_i}(x)\right) \right]. \end{aligned} \quad (4.17)$$

Multiplying the relation (4.17) by

$$\frac{1}{k\Gamma_k(\beta)} \left[ \frac{(b-x)^\alpha - (b-\rho)^\alpha}{\alpha} \right]^{\beta/k-1} \frac{\prod_{i=1}^n g_i^{\gamma_i}(\rho)}{(b-\rho)^{1-\alpha}} \quad (4.18)$$

and integrating on both sides with respect to  $\rho$  from  $x$  to  $b$  produce

$$\begin{aligned} 0 &\leq \left[ {}^\beta_k\mathcal{H}_{b-}^\alpha\left(\prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x)\right) \right] \left[ {}^\beta_k\mathcal{H}_{b-}^\alpha\left((b-x)^\eta \prod_{i=1}^n g_i^{\gamma_i}(x)\right) \right] \\ &\quad - \left[ {}^\beta_k\mathcal{H}_{b-}^\alpha\left((b-x)^\eta \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x)\right) \right] \left[ {}^\beta_k\mathcal{H}_{b-}^\alpha\left(\prod_{i=1}^n g_i^{\gamma_i}(x)\right) \right]. \end{aligned} \quad (4.19)$$

Dividing on both sides of (4.19) by

$${}^\beta_k\mathcal{H}_{b-}^\alpha\left((b-x)^\eta \prod_{i=1}^n g_i^{\gamma_i}(x)\right) \left[ {}^\beta_k\mathcal{H}_{b-}^\alpha\left(\prod_{i=1}^n g_i^{\gamma_i}(x)\right) \right]$$

yields (4.15). The proof of Theorem 4.3 is complete.  $\square$

**Corollary 4.3.** *Let  $\{g_i, 1 \leq i \leq n\}$  be a sequence of continuous positive decreasing functions on the interval  $[a, b]$ . Let  $a < x \leq b$ ,  $\eta > 0$ ,  $\xi \geq \gamma_p > 0$  for  $1 \leq p \leq n$ . Then the right  $k$ -FCI operator  ${}^\beta_k\mathcal{H}_{b-}^\alpha$  satisfies*

$$\begin{aligned} & \left[ {}^\theta_k\mathcal{H}_{b-}^\alpha \left( (b-x)^\eta \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \left[ {}^\beta_k\mathcal{H}_{b-}^\alpha \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \\ & + \left[ {}^\theta_k\mathcal{H}_{b-}^\alpha \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \left[ {}^\beta_k\mathcal{H}_{b-}^\alpha \left( (b-x)^\eta \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \\ & \geq \left[ {}^\theta_k\mathcal{H}_{b-}^\alpha \left( (b-x)^\eta \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \left[ {}^\beta_k\mathcal{H}_{b-}^\alpha \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \\ & + \left[ {}^\theta_k\mathcal{H}_{b-}^\alpha \left( \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \left[ {}^\beta_k\mathcal{H}_{b-}^\alpha \left( (b-x)^\eta \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right]. \quad (4.20) \end{aligned}$$

*Proof.* Multiplying the relation (4.17) by

$$\frac{1}{k\Gamma_k(\theta)} \left[ \frac{(b-x)^\alpha - (b-\rho)^\alpha}{\alpha} \right]^{\theta/k-1} \frac{\prod_{i=1}^n g_i^{\gamma_i}(\rho)}{(b-\rho)^{1-\alpha}} \quad (4.21)$$

and integrating on both sides with respect to  $\rho$  from  $x$  to  $b$  procure

$$\begin{aligned} 0 & \leq \left[ {}^\theta_k\mathcal{H}_{b-}^\alpha \left( (b-x)^\eta \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \left[ {}^\beta_k\mathcal{H}_{b-}^\alpha \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \\ & + \left[ {}^\theta_k\mathcal{H}_{b-}^\alpha \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \left[ {}^\beta_k\mathcal{H}_{b-}^\alpha \left( (b-x)^\eta \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \\ & - \left[ {}^\theta_k\mathcal{H}_{b-}^\alpha \left( (x-a)^\eta \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \left[ {}^\beta_k\mathcal{H}_{b-}^\alpha \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \\ & - \left[ {}^\theta_k\mathcal{H}_{b-}^\alpha \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \left[ {}^\beta_k\mathcal{H}_{b-}^\alpha \left( (b-x)^\eta \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right]. \quad (4.22) \end{aligned}$$

Dividing on both sides of (4.22) by

$$\begin{aligned} & \left[ {}^\theta_k\mathcal{H}_{b-}^\alpha \left( (b-x)^\eta \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \left[ {}^\beta_k\mathcal{H}_{b-}^\alpha \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \\ & + \left[ {}^\theta_k\mathcal{H}_{b-}^\alpha \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \left[ {}^\beta_k\mathcal{H}_{b-}^\alpha \left( (b-x)^\eta \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \end{aligned}$$

demonstrates (4.20). The proof of Corollary 4.3 is complete.  $\square$

**Theorem 4.4.** *Let  $h(x)$  be a continuous increasing function and  $\{g_i, 1 \leq i \leq n\}$  be a sequence of continuous positive decreasing functions on the interval  $[a, b]$ . Let  $a < x \leq b$ ,  $\eta > 0$ ,  $\xi \geq \gamma_p > 0$  for  $1 \leq p \leq n$ . Then the right  $k$ -FCI operator  ${}^\beta_k\mathcal{H}_{b-}^\alpha$  satisfies*

$$\frac{{}^\beta_k\mathcal{H}_{b-}^\alpha \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right)}{{}^\beta_k\mathcal{H}_{b-}^\alpha \left( \prod_{i=1}^n g_i^{\gamma_i}(x) \right)} \geq \frac{{}^\beta_k\mathcal{H}_{b-}^\alpha \left( h^\eta(x) \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right)}{{}^\beta_k\mathcal{H}_{b-}^\alpha \left( h^\eta(x) \prod_{i=1}^n g_i^{\gamma_i}(x) \right)}. \quad (4.23)$$

*Proof.* Under given conditions, we have

$$[h^\eta(\rho) - h^\eta(\tau)][g_p^{\xi-\gamma_p}(\tau) - g_p^{\xi-\gamma_p}(\rho)] \geq 0.$$

Let us define a function

$$\begin{aligned} {}^\beta_k \mathcal{J}_{b^-}^\alpha(x, \rho, \tau) &= \frac{1}{k\Gamma_k(\beta)} \left[ \frac{(b-x)^\alpha - (b-\tau)^\alpha}{\alpha} \right]^{\beta/k-1} \\ &\quad \times \frac{\prod_{i=1}^n g_i^{\gamma_i}(\tau)}{(b-\tau)^{1-\alpha}} [h^\eta(\rho) - h^\eta(\tau)][g_p^{\xi-\gamma_p}(\tau) - g_p^{\xi-\gamma_p}(\rho)]. \end{aligned} \quad (4.24)$$

Thus, the function  ${}^\beta_k \mathcal{J}_{b^-}^\alpha(x, \rho, \tau)$  is positive for all  $\tau \in (a, b]$ . Integrating on both sides of the above equation (4.24) with respect to  $\tau$  from  $x$  to  $b$  results in

$$\begin{aligned} 0 &\leq \int_x^b {}^\beta_k \mathfrak{J}_{b^-}^\alpha(x, \rho, \tau) d\tau = \frac{1}{k\Gamma_k(\beta)} \int_x^b \left[ \frac{(b-x)^\alpha - (b-\tau)^\alpha}{\alpha} \right]^{\beta/k-1} \\ &\quad \times \prod_{i=1}^n g_i^{\gamma_i}(\tau) [h^\eta(\rho) - h^\eta(\tau)][g_p^{\xi-\gamma_p}(\tau) - g_p^{\xi-\gamma_p}(\rho)] \frac{d\tau}{(b-\tau)^{1-\alpha}} \\ &= h^\eta(\rho) \left[ {}^\beta_k \mathcal{H}_{b^-}^\alpha \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] + g_p^{\xi-\gamma_p}(\rho) \left[ {}^\beta_k \mathcal{H}_{b^-}^\alpha \left( h^\eta(x) \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \\ &\quad - h^\eta(\rho) g_p^{\xi-\gamma_p}(\rho) \left[ {}^\beta_k \mathcal{H}_{b^-}^\alpha \left( \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] - \left[ {}^\beta_k \mathcal{H}_{b^-}^\alpha \left( h^\eta(x) \prod_{i \neq p}^n g_i^{\gamma_i}(x) \right) \right]. \end{aligned} \quad (4.25)$$

Multiplying the relation (4.25) by (4.18) and integrating on both sides with respect to  $\rho$  from  $x$  to  $b$  yield

$$\begin{aligned} 0 &\leq \left[ {}^\beta_k \mathcal{H}_{b^-}^\alpha \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \left[ {}^\beta_k \mathcal{H}_{b^-}^\alpha \left( h^\eta(x) \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \\ &\quad - \left[ {}^\beta_k \mathcal{H}_{b^-}^\alpha \left( h^\eta(x) \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \left[ {}^\beta_k \mathcal{H}_{b^-}^\alpha \left( \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right]. \end{aligned} \quad (4.26)$$

Dividing on both sides of (4.26) by

$${}^\beta_k \mathcal{H}_{b^-}^\alpha \left( h^\eta(x) \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \left[ {}^\beta_k \mathcal{H}_{b^-}^\alpha \left( \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right]$$

leads to (4.23). The proof of Theorem 4.4 is complete.  $\square$

**Corollary 4.4.** *Let  $h(x)$  be a continuous increasing function and  $\{g_i, 1 \leq i \leq n\}$  be a sequence of continuous positive decreasing functions on the interval  $[a, b]$ . Let  $a < x \leq b$ ,  $\eta > 0$ ,  $\xi \geq \gamma_p > 0$  for  $1 \leq p \leq n$ . Then the right  $k$ -FCI operator  ${}^\beta_k \mathcal{H}_{b^-}^\alpha$  satisfies*

$$\begin{aligned} &\left[ {}^\theta_k \mathcal{H}_{b^-}^\alpha \left( h^\eta(x) \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \left[ {}^\beta_k \mathcal{H}_{b^-}^\alpha \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \\ &\quad + \left[ {}^\theta_k \mathcal{H}_{b^-}^\alpha \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^\xi(x) \right) \right] \left[ {}^\beta_k \mathcal{H}_{b^-}^\alpha \left( h^\eta(x) \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \end{aligned}$$

$$\begin{aligned} &\geq \left[ {}_{\theta}^{\alpha} \mathcal{H}_{b^{-}}^{\alpha} \left( h^{\eta}(x) \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \left[ {}_{\beta}^{\alpha} \mathcal{H}_{b^{-}}^{\alpha} \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \\ &\quad + \left[ {}_{\theta}^{\alpha} \mathcal{H}_{b^{-}}^{\alpha} \left( \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \left[ {}_{\beta}^{\alpha} \mathcal{H}_{b^{-}}^{\alpha} \left( h^{\eta}(x) \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right]. \end{aligned} \quad (4.27)$$

*Proof.* Multiplying the relation (4.25) by (4.21) and integrating on both sides with respect to  $\rho$  from  $x$  to  $b$  give

$$\begin{aligned} 0 &\leq \left[ {}_{\theta}^{\alpha} \mathcal{H}_{b^{-}}^{\alpha} \left( h^{\eta}(x) \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \left[ {}_{\beta}^{\alpha} \mathcal{H}_{b^{-}}^{\alpha} \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \\ &\quad + \left[ {}_{\theta}^{\alpha} \mathcal{H}_{b^{-}}^{\alpha} \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \left[ {}_{\beta}^{\alpha} \mathcal{H}_{b^{-}}^{\alpha} \left( h^{\eta}(x) \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \\ &\quad - \left[ {}_{\theta}^{\alpha} \mathcal{H}_{b^{-}}^{\alpha} \left( h^{\eta}(x) \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \left[ {}_{\beta}^{\alpha} \mathcal{H}_{b^{-}}^{\alpha} \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \\ &\quad - \left[ {}_{\theta}^{\alpha} \mathcal{H}_{b^{-}}^{\alpha} \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \left[ {}_{\beta}^{\alpha} \mathcal{H}_{b^{-}}^{\alpha} \left( h^{\eta}(x) \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right]. \end{aligned} \quad (4.28)$$

Dividing on both sides of (4.28) by

$$\begin{aligned} &\left[ {}_{\theta}^{\alpha} \mathcal{H}_{b^{-}}^{\alpha} \left( h^{\eta}(x) \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \left[ {}_{\beta}^{\alpha} \mathcal{H}_{b^{-}}^{\alpha} \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \\ &\quad + \left[ {}_{\theta}^{\alpha} \mathcal{H}_{b^{-}}^{\alpha} \left( \prod_{i \neq p}^n g_i^{\gamma_i} g_p^{\xi}(x) \right) \right] \left[ {}_{\beta}^{\alpha} \mathcal{H}_{b^{-}}^{\alpha} \left( h^{\eta}(x) \prod_{i=1}^n g_i^{\gamma_i}(x) \right) \right] \end{aligned}$$

concludes (4.27). The proof of Corollary 4.4 is complete. □

### 5. CONCLUSIONS

In this paper, we have presented the left and right  $k$ -fractional conformable integrals and generalized some important integral inequalities to ones for our newly introduced  $k$ -FCI operators related to a finite sequence of positive and decreasing functions. Our work produces  $k$ -analogues of many pre-existing results in the literature. Further, many special cases for other integral operators can be derived from our generalizations. The results obtained can be employed to confirm the existence of nontrivial solutions of fractional differential equations of different classes. The  $k$ -FCI operators in this paper are different from those introduced by Katugampola [20] as their kernels depend on the boundary points  $a$  and  $b$  and need a different convolution theory under conformable Laplace. Our  $k$ -fractional conformable integrals in this paper generalize well-known fractional integral operators such as Caputo integral operators [36, p. 44], Riemann–Liouville integral operators [36, p. 44], Hadamard integral operators [4], and their  $k$ -analogues.

### REFERENCES

[1] T. Abdeljawad, *On conformable fractional calculus*, J. Comput. Appl. Math. **279** (2015), 57–66; Available online at <https://doi.org/10.1016/j.cam.2014.10.016>.

- [2] P. Agarwal, M. Jleli, and M. Tomar. *Certain Hermite–Hadamard type inequalities via generalized  $k$ -fractional integrals*, J. Inequal. Appl. **2017**, Paper No. 55, 10 pp. Available online at <https://doi.org/10.1186/s13660-017-1318-y>.
- [3] R. P. Agarwal and A. Özbekler, *Lyapunov type inequalities for mixed nonlinear Riemann–Liouville fractional differential equations with a forcing term*, J. Comput. Appl. Math. **314** (2017), 69–78; Available online at <https://doi.org/10.1016/j.cam.2016.10.009>.
- [4] A. A. Kilbas, *Hadamard-type fractional calculus*, J. Korean Math. Soc. **38** (2001), no. 6, 1191–1204.
- [5] D. Băleanu, R. P. Agarwal, H. Khan, R. A. Khan, and H. Jafari, *On the existence of solution for fractional differential equations of order  $3 < \delta_1 \leq 4$* , Adv. Difference Equ. **2015**, 2015:362, 9 pp. Available online at <https://doi.org/10.1186/s13662-015-0686-1>.
- [6] D. Băleanu, R. P. Agarwal, H. Mohammadi, and S. Rezapour, *Some existence results for a nonlinear fractional differential equation on partially ordered Banach spaces*, Bound. Value Probl. **2013**, 2013:112, 8 pp. Available online at <https://doi.org/10.1186/1687-2770-2013-112>.
- [7] D. Băleanu, H. Khan, H. Jafari, R. A. Khan, and M. Alipour, *On existence results for solutions of a coupled system of hybrid boundary value problems with hybrid conditions*, Adv. Difference Equ. **2015**, 2015:318, 14 pp. Available online at <https://doi.org/10.1186/s13662-015-0651-z>.
- [8] D. Băleanu and O. G. Mustafa, *On the global existence of solutions to a class of fractional differential equations*, Comput. Math. Appl. **59** (2010), no. 5, 1835–1841; Available online at <https://doi.org/10.1016/j.camwa.2009.08.028>.
- [9] D. Băleanu, O. G. Mustafa, and R. P. Agarwal, *An existence result for a superlinear fractional differential equation*, Appl. Math. Lett. **23** (2010), no. 9, 1129–1132; Available online at <https://doi.org/10.1016/j.aml.2010.04.049>.
- [10] D. Băleanu, O. G. Mustafa, and R. P. Agarwal, *On the solution set for a class of sequential fractional differential equations*, J. Phys. A **43** (2010), no. 38, 385209, 7 pp. Available online at <https://doi.org/10.1088/1751-8113/43/38/385209>.
- [11] A. Bolandtalat, E. Babolian, H. Jafari, *Numerical solutions of multi-order fractional differential equations by Boubaker polynomials*, Open Phys. **14** (2016), no. 1, 226–230; Available online at <https://doi.org/10.1515/phys-2016-0028>.
- [12] A. Debbouche and V. Antonov, *Finite-dimensional diffusion models of heat transfer in fractal mediums involving local fractional derivatives*, Nonlinear Stud. **24** (2017), no. 3, 527–535.
- [13] R. Diaz and E. Pariguan, *On hypergeometric functions and Pochhammer  $k$ -symbol*, Divulg. Mat. **15** (2007), no. 2, 179–192.
- [14] P. R. Halmos, *Measure Theory*, D. Van Nostrand Company, Inc., New York, N. Y., 1950.
- [15] H. Jafari, H. K. Jassim, S. P. Moshokoa, V. M. Ariyan, and F. Tchier, *Reduced differential transform method for partial differential equations within local fractional derivative operators*, Adv. Mech. Eng. **8** (2016), no. 4, 1–6; Available online at <https://doi.org/10.1177/1687814016633013>.
- [16] H. Jafari, H. K. Jassim, F. Tchier, and D. Băleanu, *On the approximate solutions of local fractional differential equations with local fractional operators*, Entropy **18** (2016), no. 4, Paper No. 150, 12 pp. Available online at <https://doi.org/10.3390/e18040150>.
- [17] F. Jarad, E. Uğurlu, T. Abdeljawad, and D. Băleanu. *On a new class of fractional operators*, Adv. Difference Equ. **2017**, Paper No. 247, 16 pp. Available online at <https://doi.org/10.1186/s13662-017-1306-z>.
- [18] M. Jleli, M. Kirane, and B. Samet, *Hartman–Wintner-type inequality for a fractional boundary value problem via a fractional derivative with respect to another function*, Discrete Dyn. Nat. Soc. **2017**, Art. ID 5123240, 8 pp. Available online at <https://doi.org/10.1155/2017/5123240>.
- [19] U. N. Katugampola, *New approach to a generalized fractional integral*, Appl. Math. Comput. **218** (2011), no. 3, 860–865; Available online at <https://doi.org/10.1016/j.amc.2011.03.062>.
- [20] U. N. Katugampola, *New fractional integral unifying six existing fractional integrals*, arXiv preprint (2016), available online at <https://arxiv.org/abs/1612.08596>.
- [21] H. Khan, H. Jafari, D. Băleanu, R. A. Khan, and A. Khan, *On iterative solutions and error estimations of a coupled system of fractional order differential-integral equations with*

- initial and boundary conditions*, Differ. Equ. Dyn. Syst. (2018), in press; Available online at <https://doi.org/10.1007/s12591-017-0365-7>.
- [22] H. Khan, Y.-J. Li, W. Chen, D. Baleanu, and A. Khan, *Existence theorems and Hyers-Ulam stability for a coupled system of fractional differential equations with  $p$ -Laplacian operator*, Bound. Value Probl. **2017**, Paper No. 157, 16 pp. Available online at <https://doi.org/10.1186/s13661-017-0878-6>.
- [23] A. Khan, Y.-J. Li, K. Shah, and T. S. Khan, *On coupled  $p$ -Laplacian fractional differential equations with nonlinear boundary conditions*, Complexity **2017**, Art. ID 8197610, 9 pp. Available online at <https://doi.org/10.1155/2017/8197610>.
- [24] H. Khan, Y.-J. Li, H.-G. Sun, and A. Khan, *Existence of solution and Hyers-Ulam stability for a coupled system of fractional differential equations with  $p$ -Laplacian operator*, J. Nonlinear Sci. Appl. **10** (2017), no. 10, 5219–5229.
- [25] H. Khan, H.-G. Sun, W. Chen, and D. Baleanu, *Inequalities for new class of fractional integral operators*, J. Nonlinear Sci. Appl. **10** (2017), no. 12, 6166–6176.
- [26] Y.-J. Li, K. Shah, and R. A. Khan, *Iterative technique for coupled integral boundary value problem of non-integer order differential equations*, Adv. Difference Equ. **2017**, Paper No. 251, 14 pp. Available online at <https://doi.org/10.1186/s13662-017-1308-x>.
- [27] W.-J. Liu, Q.-A. Ngô, and V. N. Huy, *Several interesting integral inequalities*, J. Math. Inequal. **3** (2009), no. 2, 201–212; Available online at <https://doi.org/10.7153/jmi-03-20>.
- [28] S. Mubeen and S. Iqbal, *Grüss type integral inequalities for generalized Riemann–Liouville  $k$ -fractional integrals*, J. Inequal. Appl. **2016**, Paper No. 109, 13 pp. Available online at <https://doi.org/10.1186/s13660-016-1052-x>.
- [29] S. Mubeen, S. Iqbal, and Z. Iqbal, *On Ostrowski type inequalities for generalized  $k$ -fractional integrals*, J. Inequal. Spec. Funct. **8** (2017), no. 3, 107–118.
- [30] K. S. Nisar and F. Qi, *On solutions of fractional kinetic equations involving the generalized  $k$ -Bessel function*, Note Mat. **37** (2017) no. 2, 11–20; Available online at <https://doi.org/10.1285/i15900932v37n2p11>.
- [31] K. S. Nisar, F. Qi, G. Rahman, S. Mubeen, and M. Arshad, *Some inequalities involving the extended gamma and beta functions*, HAL archives (2018), available online at <https://hal.archives-ouvertes.fr/hal-01701647>.
- [32] D. O'Regan and B. Samet, *Lyapunov-type inequalities for a class of fractional differential equations*, J. Inequal. Appl. **2015**, 2015:247, 10 pp. Available online at <https://doi.org/10.1186/s13660-015-0769-2>.
- [33] F. Qi, A. Akkurt, and H. Yildirim, *Catalan numbers,  $k$ -gamma and  $k$ -beta functions, and parametric integrals*, J. Comput. Anal. Appl. **25** (2018), no. 6, 1036–1042.
- [34] F. Qi and K. S. Nisar, *Some integral transforms of the generalized  $k$ -Mittag-Leffler function*, Publ. Inst. Math. (Beograd) (N.S.) **103(117)** (2018), in press; Preprints **2016**, 2016100020, 8 pages; Available online at <https://doi.org/10.20944/preprints201610.0020.v1>.
- [35] F. Qi, G. Rahman, and K. S. Nisar, *Convexity and inequalities related to extended beta and confluent hypergeometric functions*, HAL archives (2018), available online at <https://hal.archives-ouvertes.fr/hal-01703900>.
- [36] S. G. Samko, A. A. Kilbas, and O. I. Marichev, *Fractional Integrals and Derivatives: Theory and Applications*, Edited and with a foreword by S. M. Nikol'skiĭ. Translated from the 1987 Russian original. Revised by the authors. Gordon and Breach Science Publishers, Yverdon, 1993.
- [37] M. Z. Sarikaya and H. Budak, *Generalized Ostrowski type inequalities for local fractional integrals*, Proc. Amer. Math. Soc. **145** (2017), no. 4, 1527–1538; Available online at <https://doi.org/10.1090/proc/13488>.
- [38] M. Z. Sarikaya, Z. Dahmani, M. E. Kiris, and F. Ahmad,  *$(k, s)$ -Riemann–Liouville fractional integral and applications*, Hacet. J. Math. Stat. **45** (2016), no. 1, 77–89.
- [39] E. Set, M. A. Noor, M. U. Awan, and A. Gözpinar, *Generalized Hermite–Hadamard type inequalities involving fractional integral operators*, J. Inequal. Appl. **2017**, Paper No. 169, 10 pp. Available online at <https://doi.org/10.1186/s13660-017-1444-6>.
- [40] E. Set, M. Tomar, and M. Z. Sarikaya, *On generalized Grüss type inequalities for  $k$ -fractional integrals*, Appl. Math. Comput. **269** (2015), 29–34; Available online at <https://doi.org/10.1016/j.amc.2015.07.026>.

- [41] K. Shah, H. Khalil, and R. A. Khan, *Upper and lower solutions to a coupled system of nonlinear fractional differential equations*, Progress Frac. Differ. Equ. Appl. **2** (2016), no. 1, 31–39; Available online at <https://doi.org/10.18576/pfda/020104>.
- [42] K. Shah and R. A. Khan, *Study of solution to a toppled system of fractional differential equations with integral boundary conditions*, Int. J. Appl. Comput. Math. **3** (2017), no. 3, 2369–2388; Available online at <https://doi.org/10.1007/s40819-016-0243-y>.
- [43] D.-P. Shi, B.-Y. Xi, and F. Qi, *Hermite–Hadamard type inequalities for  $(m, h_1, h_2)$ -convex functions via Riemann–Liouville fractional integrals*, Turkish J. Anal. Number Theory **2** (2014), no. 1, 22–27; Available online at <https://doi.org/10.12691/tjant-2-1-6>.
- [44] D.-P. Shi, B.-Y. Xi, and F. Qi, *Hermite–Hadamard type inequalities for Riemann–Liouville fractional integrals of  $(\alpha, m)$ -convex functions*, Fract. Diff. Calculus **4** (2014), no. 2, 33–43; Available online at <https://doi.org/10.7153/fdc-04-02>.
- [45] H.-G. Sun, Y. Zhang, W. Chen, and D. M. Reeves, *Use of a variable-index fractional-derivative model to capture transient dispersion in heterogeneous media*, J. Contaminant Hydrology **157** (2014), 47–58; Available online at <https://doi.org/10.1016/j.jconhyd.2013.11.002>.
- [46] Y.-Z. Tian, M. Fan, and Y.-G. Sun, *Certain nonlinear integral inequalities and their applications*, Discrete Dyn. Nat. Soc. **2017**, Art. ID 8290906, 8 pp. Available online at <https://doi.org/10.1155/2017/8290906>.
- [47] M. Tunç *On new inequalities for  $h$ -convex functions via Riemann–Liouville fractional integration*, Filomat **27** (2013), no. 4, 559–565; Available online at <https://doi.org/10.2298/FIL1304559T>.
- [48] S.-H. Wang and F. Qi, *Hermite–Hadamard type inequalities for  $s$ -convex functions via Riemann–Liouville fractional integrals*, J. Comput. Anal. Appl. **22** (2017), no. 6, 1124–1134.
- [49] X. Yang and K. Lo, *Lyapunov-type inequality for a class of even-order differential equations*, Appl. Math. Comput. **215** (2010), no. 11, 3884–3890; Available online at <https://doi.org/10.1016/j.amc.2009.11.032>.
- [50] X.-J. Yang, J. A. T. Machado, and J. J. Nieto, *A new family of the local fractional PDEs*, Fund. Inform. **151** (2017), no. 1-4, 63–75; Available online at <https://doi.org/10.3233/FI-2017-1479>.
- [51] X.-J. Yang, *Fractional derivatives of constant and variable orders applied to anomalous relaxation models in heat-transfer problems*, Therm. Sci. **21** (2017), no. 3, 1161–1171; Available online at <https://doi.org/10.2298/TSCI161216326Y>.
- [52] X.-J. Yang, J. A. T. Machao, and D. Băleanu, *Anomalous diffusion models with general fractional derivatives within the kernels of the extended Mittag-Leffler type functions*, Roman. Rep. Phys. **69** (2017), Article no. 115, 20 pp.
- [53] X.-J. Yang, H. M. Srivastava, and J. A. T. Machado, *A new fractional derivative without singular kernel: Application to the modelling of the steady heat flow*, Therm. Sci. **20** (2016), no. 2, 753–756; Available online at <https://doi.org/10.2298/TSCI151224222Y>.

(Habib) DEPARTMENT OF MATHEMATICS, GOVERNMENT COLLEGE UNIVERSITY FAISALABAD, PAKISTAN

*Email address:* [siddrahabib1605@gmail.com](mailto:siddrahabib1605@gmail.com)

(Mubeen) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SARGODHA, SARGODHA, PAKISTAN

*Email address:* [smjhanda@gmail.com](mailto:smjhanda@gmail.com)

(Naeem) DEPARTMENT OF MATHEMATICS, GOVERNMENT COLLEGE UNIVERSITY FAISALABAD, PAKISTAN

*Email address:* [mnawaznaem@yahoo.com](mailto:mnawaznaem@yahoo.com)

(Qi) INSTITUTE OF MATHEMATICS, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO, HENAN, 454010, CHINA; INSTITUTE OF FUNDAMENTAL AND FRONTIER SCIENCES, UNIVERSITY OF ELECTRONIC SCIENCE AND TECHNOLOGY OF CHINA, CHENGDU, SICHUAN, 610054, CHINA; DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCE, TIANJIN POLYTECHNIC UNIVERSITY, TIANJIN, 300387, CHINA

*Email address:* [qifeng618@gmail.com](mailto:qifeng618@gmail.com), [qifeng618@hotmail.com](mailto:qifeng618@hotmail.com), [qifeng618@qq.com](mailto:qifeng618@qq.com)

*URL:* <https://qifeng618.wordpress.com>