



HAL
open science

The red-vented bulbul (*Pycnonotus cafer*): serious pest or understudied invader?

Martin Thibault, Eric Vidal, Murray A. Potter, Ellie Dyer, Fabrice Brescia

► To cite this version:

Martin Thibault, Eric Vidal, Murray A. Potter, Ellie Dyer, Fabrice Brescia. The red-vented bulbul (*Pycnonotus cafer*): serious pest or understudied invader?. *Biological Invasions*, 2018, 20 (1), pp.121-136. 10.1007/s10530-017-1521-2 . hal-01789436

HAL Id: hal-01789436


<https://hal.science/hal-01789436>

Submitted on 2 Apr 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The red-vented bulbul (*Pycnonotus cafer*): serious pest or understudied invader?

Martin Thibault  · Eric Vidal · Murray A. Potter ·
Ellie Dyer · Fabrice Brescia

Received: 28 December 2016 / Accepted: 23 July 2017

Abstract Recently, debate has flourished about inadequacies in the simplistic “worst invasive species” approach and its global scale. Here we investigate the status of the red-vented bulbul (*Pycnonotus cafer*), an Asian passerine bird. This species has been introduced widely across Pacific islands and is commonly blamed for its impacts on agriculture and biodiversity via dispersal of invasive plant seeds and competition with native fauna. This case study evaluates all available data on the impacts and management of this invasive species and identifies priorities

Electronic supplementary material The online version of this article (doi:[10.1007/s10530-017-1521-2](https://doi.org/10.1007/s10530-017-1521-2)) contains supplementary material, which is available to authorized users.

M. Thibault (✉) · F. Brescia
Institut Agronomique néo-Calédonien (IAC), Equipe ARBOREAL (Agriculture BiODiversité Et vAlorisation),
73, 98890 Païta, New Caledonia
e-mail: thibault@iac.nc

E. Vidal
Institut Méditerranéen de Biodiversité et d'Ecologie marine et continentale (IMBE), CNRS, IRD, Avignon Université Centre IRD Nouméa, Aix Marseille Université, BP A5, 98848 Nouméa Cedex, New Caledonia

M. Thibault · M. A. Potter
Ecology Group, Institute of Agriculture and Environment,
Massey University, Palmerston North 4442, New Zealand

E. Dyer
Institute of Zoology, Zoological Society of London,
Regent's Park, London NW1 4RY, UK

for future research. We reviewed the scientific literature and information from three databases (ABBA, GAVIA, eBird) and highlight that the attention paid to this species by scientists and managers varied considerably between islands and contexts and was globally lower than the attention paid to other species on the IUCN-ISSG list. The red-vented bulbul has now established on 37 islands and in seven continental locations outside its native range. We show that three categories of effects are associated with this species: plant damage, seed dispersal and disturbance of fauna. We compiled lists of 110 plant species consumed, 33 plant species dispersed, and 15 species of bird that this bulbul interacts with. However, these lists were mainly made of opportunistic observations rather than specific assessments. Research outputs that focus on better ways to prevent or quantify the impacts of the red-vented bulbul remain scarce. We found very few references exploring potential positive impacts of this species, and only two examples of management actions undertaken against it. The latter are required to inform management actions, especially on sensitive tropical islands where invasions and dispersal of the red-vented bulbul are ongoing. Our analysis of the literature found no clear support for considering this species to be one of the “world’s worst” invasive alien species.

Keywords Invasive alien bird · Islands · Impact · Biodiversity · Conservation

Introduction

Invasive alien species (IAS) are one of the main causes of biodiversity loss (Sala et al. 2000; Keane and Crawley 2002; Pereira et al. 2012; Gren et al. 2016), with associated economic impacts (Bergman et al. 2000; Pimentel et al. 2005; Pimentel 2011) and degradation of ecosystem services (Walsh et al. 2016). The highly ambitious goal of the 2010 Convention for Biological Diversity, Nagoya, Japan, was to ensure that “By 2020, IAS and pathways are identified and prioritized, priority species are controlled or eradicated, and measures are in place to manage pathways to prevent their introduction and establishment” (Secretariat CBD 2010). Concerns about the impacts of IAS have led to the production of several lists of high priority alien species including the “100 of the World’s Worst” from the IUCN Invasive Species Specialist Group (IUCN-ISSG) (Brochier et al. 2010; Burgiel and Perrault 2011; Lowe et al. 2000). Such prioritization attempts have incited intense debate among the scientific community on the definition of an invasive species (e.g. Russell and Blackburn 2017). Some considered invasion as a natural phenomenon and compared the prevention of species dispersal as a kind of racism (Valéry et al. 2013). Others saw the observed impacts of alien species as an important challenge for our developing societies (Richardson and Ricciardi 2013; Simberloff and Vitule. 2014; Blondel et al. 2014; Pereyra 2016). Such debate also applied to species classification methods, as prioritization attempts based on expert assessments is opposed to different classification frameworks based on data analysis and statistics (Donlan and Wilcox 2008; Kumschick et al. 2012; Blackburn et al. 2014; Kumschick et al. 2015). In this study, we consider that an alien species expanding its range in a sensitive territory deserves particular attention from both scientists and managers. For their part, scientists must consider the possibility that a species could be harmless in an alien territory and should produce a local assessment of potential issues associated with that species.

Of the terrestrial vertebrates in the IUCN-ISSG list, 14 are mammals, three are birds and only two are reptiles (Lowe et al. 2000). Unsurprisingly, 10 years after the publication of this “World’s Worst” list, authors have commented on the imbalance in attention paid by scientists and managers to mammals in

contrast to alien birds (Pyšek et al. 2008; Kumschick and Nentwig 2010). Several studies have called for improvements in the way in which impact values of IAS are assessed beyond experts’ “worst” lists, particularly for bird species (Strubbe et al. 2011; Ricciardi et al. 2013; Kumschick et al. 2015; Saxena 2015) and they have stressed that this is vital to better inform management decisions.

The three bird species classified as the world’s worst invasive species in the IUCN-ISSG list are the common starling (*Sturnus vulgaris*), the common myna (*Acridotheres tristis*), and the red-vented bulbul (*Pycnonotus cafer*). A recent review of the impact of alien birds on native ecosystems globally (Martin-Albarracin et al. 2015) identified the three species with the highest global impact as being the mallard (*Anas platyrhynchos*, score = 16), the common myna (score = 13), and the red-whiskered bulbul (*Pycnonotus jocosus*, score = 10) whereas the global impact score of the red-vented bulbul in this study should be only 4. Recently, Kumschick et al. (2015) identified important overlaps in the impacts associated with the common myna and the red-vented bulbul. This raises the question as to whether the red-vented bulbul should be considered as one of the three worst invasive bird species on the planet. Local farmers and environment managers need management frameworks in order to deal with the dispersal/impacts of IAS (Blackburn et al. 2011). Biosecurity protocols at frontiers were demonstrated to be the most useful techniques to prevent biological invasions (Edelaar and Tella 2012). However, controlling a newly established invasive species in a territory depends on economic, ecological and social factors and therefore on specific assessment of these factors at local scales (Mack et al. 2000). A synthesis of management programs toward an alien species offers concrete baselines for managers, and this knowledge also helps understanding how the impacts associated with an invasive species give rise to management operations. It is urgent that we review existing assessments of impacts and management programs implemented against the red-vented bulbul.

We review invasion data to determine the nature and severity of the impacts of this species, whether its impacts are consistent throughout its alien range, whether its status as a major invasive species has led to more research and management programs at local scales, and to assess if its current acknowledged pest

status is deserved. We present an updated assessment of an invasive species nearly 110 years after it was first record outside of its native range (Fiji in 1903, Watling 1978) and identify priorities for future research.

Methods

Species description

The red-vented bulbul (*Pycnonotus cafer* Linnaeus, 1766) is a passerine belonging to the family Pycnonotidae. Earlier names include *Molpastes haemorrhous* J.F. Gmelin, 1789 and *Molpastes cafer* Baker, 1930. The genus *Pycnonotus* comprises 47 species (Delacour 1943; Dickinson and Dekker 2002), among which the red-vented bulbul is represented by eight different subspecies (Dickinson et al. 2002). The Pacific sub-species is *P. c. bengalensis*, Blyth 1845 (Watling 1978).

The red-vented bulbul is native to the Indian Subcontinent, Southeast Asia, and Malay Peninsula (Long 1981). It occurs naturally from Eastern Pakistan to southern China and Vietnam, and from Northern India to Sri Lanka. The species also has an historic presence in Bangladesh, Bhutan, Myanmar and Nepal.

Data collection and analysis

We searched for “*Pycnonotus cafer*” and “red-vented bulbul” keywords on Google Scholar, Web of Science, ScienceDirect and SpringerLink search engines. We looked for information on impacts primarily, and collected references on pathways of invasion, establishment success, and management. We also visited the websites of the Governments, Environment Ministry, Associations and NGOs for each country where the red-vented bulbul was signaled as present. When our searches failed to find the information we sought, we made direct contact with people who had reported the presence of this species in each country. Most of the documents obtained concerned the red-vented bulbul in its alien range. Those that related to this species in its native range were used to extract information on its biology and ecology in countries of origin. We also compared the number of references obtained by searching for each of the

species names listed in the IUCN “World’s Worst” list in Google Scholar.

In order to update distribution maps, we included records from three international databases. We used the 252 quotations with references from the Global AVian Invasion Atlas Project (GAVIA, Dyer et al. 2017), 150 sightings from the Atlas of Breeding Birds of Arabia (ABBA, Ornithological Society of the Middle East, Jennings 2010) and 40,152 sightings from the participative eBird database (eBird, Sullivan et al. 2009). Maps were designed using the following R software packages: maps (Becker et al. 2015a), mapdata (Becker et al. 2015b), and mapproj (McIlroy et al. 2015) and maptools (Bivand and Lewin-Koh 2016).


We classified the reported impacts of the red-vented bulbul into three categories: (1) plant damage; (2) seed dispersal; and (3) disturbance and impact on fauna. We treated each mention of a species-specific plant or animal impact as one “report”. One published article thus often contained several “reports” when listing, for example, species of plants consumed, and the full set of documents potentially contained several reports of impacts on the same species, sometimes at the same location. We chose this index because it facilitates across-taxa comparisons and it is simple to calculate from the large number of references obtained. In addition, the ratio of the number of reports by the number of impacted species provides an informative insight into the attention paid to each impact-category.

Results

Sources of the information

We identified 112 published documents on the red-vented bulbul, and obtained comments from seven ornithologists and environment managers about the bulbul from its alien range. The publications comprised 78 academic articles, 15 books, five conference proceedings, five newsletters and nine professional reports. Details about the information obtained are presented in Fig. 1. Among the collected references, 83 addressed the red-vented bulbul in their alien range 74 from islands and nine from continental areas. Three locations (Fiji, Hawaii and French Polynesia) were the focus of 42 documents. We used 12 references that

Fig. 1 Sources of the collected information. The “General information” scope refers to documents dealing with invasion biology at a global scale



focused on red-vented bulbul in their native range. We also used information from 17 documents dealing with biological invasions at a larger scale. These documents cover a period from 1926 to today, but we focus here on documents from 1975 onwards. The cumulative numbers of publications through time are shown in Fig. 2. A full list of the 112 documents is given in Online Resource 1.

Searching for “*Pycnonotus cafer*” in Google Scholar produced 1370 references. Thus, among the 100 species listed by the IUCN, the red-vented bulbul ranked 11th. In comparison, we found 4880 references for “*Acridotheres tristis*”, and 36,500 for “*Sturnus vulgaris*”, the two other bird species from the list. Searches for “*Pycnonotus jocosus*” and “*Anas platyrhynchos*” resulted 1300 and 24,300 references respectively.

Pathways of transport and introduction

The red-vented bulbul was first reported in Fiji in ~1903 (Parham 1955), corresponding to the

transportation of Indian immigrants from Calcutta harbor to Fiji in the early 1900s (Watling 1978). This species was widely used in bird fights in India (Ali and Ripley 1996) because of its aggressive behavior. Over the following century, the red-vented bulbul was introduced into 19 countries and established in 17 of them (Fig. 3). It is now present in at least 37 islands and seven continental locations, and is anticipated to continue its range expansion in several archipelagos. The first recorded year of observation per country is presented in Table 1. Most introductions of the red-vented bulbul have been in the Pacific and in the Middle East, but the species was recently recorded in southern Europe (Malaga, Spain) and in North America (Houston, Texas, USA) (Fig. 3). The exact reason for introduction is known for only three locations. The red-vented bulbul was deliberately introduced (1940s) to Tongatapu (Tonga) to control unwanted insects (Watling 1978). An American troopship re-routed to Apia took caged birds to Samoa in 1943 instead of New Caledonia that was the intended destination




Fig. 2 Number of alien locations and published references for red-vented bulbul for the period 1903–2013




Fig. 3 Native and alien range of the red vented bulbul

(Watling 1978), and the species was intentionally released in Nouméa (New Caledonia) in 1983 by bird dealers to avoid prosecution (Gill et al. 1995). For 10 other locations, bird trade is most often the suspected reason for introduction. Birds were kept in cages and transported by boat or airplane, with accidental or intentional release occurring commonly around harbors, airports and markets. For the Pacific locations, transportation of caged birds and accidental transport of free birds by boat have been the main introduction pathways, with a few records indicating that some introductions have occurred near airports. The red-vented bulbul remains abundant in Tahiti (French Polynesia) and is expanding its range in the Polynesian archipelago (T. Ghestemme *pers.comm.*). In the Middle East, land and air transport of cage birds between markets is implicated. It is not known how the species got to Houston (Texas, USA), Malaga or Corralejo (Spain).

Establishment success

The red vented bulbul is currently considered established in 36 of the 46 locations where it has been historically recorded. Up-to-date information is lacking for three small Pacific islands (‘Eua, Savai’i and Ailinglaplap). We found mentions of red-vented bulbuls in Melbourne in 1918 and 1942 (Lendon 1952; Watling 1978), but the species has not been

reported there since and it was recorded as “Eradicated” in Australia in the global invasive species database (<http://www.issg.org/database>). It was observed on five islands in the Hawaii archipelago between 1982 and 1989, but it seems that it failed to establish beyond Oahu (Walker 2008). It was eradicated from Auckland, New Zealand, in 1955 (Watling 1978), 3 years after the first observation in 1952 (Turbott 1956).

Recorded impacts

The red-vented bulbul is commonly blamed for three categories of negative impacts, mostly related to its diverse diet that comprises fruits and berries (Islam and Williams 2000; Brooks 2013), and flowers, buds, insects and small reptiles (VanderVelde 2002). We found 165 reports (110 species) of plants that are eaten by the red-vented bulbul. Among these, 50% concerned the degradation of cultivated plants and 35% related to seed dispersal. The remaining 17% (26 species from 17 families) were reports of consumption without consideration of the impacts. In comparison, we found 22 reports of impacts on local fauna in the bulbul’s alien range.

Damage to cultivated plants is the most frequently reported impact of the red-vented bulbul in its alien range (Fig. 4), but these studies were conducted in just four locations. In contrast, the publications reporting

Table 1 The current alien distribution of the red-vented bulbul *Pycnonotus cafer*, year of first observation (Y.F.O), number of colonized islands, current introduction success, status, range trend, and associated references. (+) Scarce (++) Common (+++) Very common

Country	Y.F.O	Colonized Islands	Naturalization success	Current status	Range trend	Number of references	Main references
Island							
Fiji	1903	9	9	++	→↗	10	(1)
Australia	1919	1	0	-	-	4	(2)
Tonga	1928	4	≥3	++		3	(3)
Western Samoa	1943	1	1	++		7	(4)
New Zealand	1952	1	0	-	-	1	(5)
American Samoa	>1957	2	≥1			7	(2)
United States (Hawaii)	1966	6	1	+++	→	20	(7)
French Polynesia	1979	9	9	+++	↗	12	(8)
New Caledonia	1983	1	1	+++	↗	5	(9)
Bahrain	1986	1	1			2	(10)
Spain	1998	1	1	+	↗	1	(11)
Marshall Islands	2000	2	≥1		→	1	(12)
Iran	2007	1	1	+		1	(13)
Continental							
Qatar	1971	-	1	+		1	(14)
United Arab Emirates	1974	-	1	++	↗	3	(15)
Kuwait	1981	-	1	+	↘	1	(16)
Oman	1987	-	1	++	↗	0	(17)
Saudi Arabia	1980's	-	1	+		2	(18)
United States (texas)	1996	-	1	+	→	1	(19)
Spain	na	-	1	+	↗	1	(11)

(1) Watling (1978), (2) Lendon (1952), (3) Carlson (1974), (4) Dhondt (1976), (5) Turbott (1956), (6) Clapp and Sibley (1966), (7) Berger (1975), (8) Meyer (1996), (9) Gill et al. (1995), (10)Khamis (2010), (11) MAAMA (2013), (12) VanderVelde (2002), (13) Azin et al. (2008), (14) Nation et al. (1997), (15) Pederson and Aspinall (2015), (16) Gregory (2005), (17) J. Eriksen com.pers., (18) J. Babbington com.pers, (19) Brooks (2013)

the red-vented bulbul to be a problematic seed disperser come from eight locations (six countries), and faunal impacts are reported for 17 species from 11 locations.

Plant damage

The red-vented bulbul has been reported to cause damage to at least 52 plant species (Table 2) belonging to 25 families with 67% (35 species) being food plants and 33% (17 species) being ornamental plant species. The full list of damaged and dispersed plant

species by family and species is given in Online Resource 2.

The impact of the red-vented bulbul appears to be particularly serious on Oahu (Hawaii), where Walker (2008) reported them consuming several species of fruits, vegetables and flowers, leading to considerable economic losses. The estimated value of the damage to Oahu's Orchid industry in 1 year (1989) was \$300,000 (Fox 2011) when the red-vented bulbul together with the Japanese white-eye (*Zosterops japonicus*) reportedly destroying up to 75% of Hawaiian orchid and anthurium plantations (Cummings et al. 1994). In New

Fig. 4 Representation of the three impact categories associated with the red-vented bulbul *Pycnonotus cafer*. Each axis corresponds to one category and represents the number of reports, species and location




Table 2 Numbers of plant species reported as damaged, dispersed or just consumed by the red-vented bulbul *Pycnonotus cafer* in the literature and corresponding number of reports

Impact	Status	Species	Reports
Damage		52	81
	Food plant	35	61
	Ornamental plant	17	20
Seeds dispersal		33	56
	Endemic	1	1
	Native	8	11
	Alien	10	16
	Invasive	14	28
Consumption only		25	28
Total		110	165

A report corresponds to one mention in one reference. Endemic plants occurred at one location only, native plants are indigenous to the location but also present elsewhere, alien species were introduced in the corresponding location and invasive plants are alien species with negative impacts at the current location

Caledonia, significant impacts have been recorded for some crops and plant nurseries (Metzdorf and Brescia 2008) with up to 35% losses (Caplong and Barjon 2010). Conversely, the red-vented bulbul is not considered an agricultural pest in Fiji (Watling 1979), nor in Houston (Texas, USA) where it was found to consume mainly introduced tropical plant species (Brooks 2013).

Seed dispersal

We found 56 mentions of problematic seed dispersal by the red-vented bulbul (Table 2) from six countries inside its alien range. The red-vented bulbul is able to spread the seeds of at least 33 plant species from 25 families. Among these species, 30% are considered alien (10 species) and 42% invasive (14 species) in the alien locations. We found records of only one endemic (*Coprosma taitensis*, Tahiti) and eight native species that are spread by this bird (Spotswood et al. 2012).

The red-vented bulbul is considered a major vector of the invasive tree *Miconia calyvescens* in Tahiti (Meyer 1996) and can potentially disperse seven other alien plant species in French Polynesia including *Lantana camara* (Spotswood et al. 2012, 2013). Its ability to disperse *Miconia* and *Lantana* is not unique to the red-vented bulbul, and many other species, both alien and native, also disperse seeds of these plants, and the propensity of the red-vented bulbul to disperse seeds of these plants varies from island to island. For example, the introduced silvereye (*Zosterops lateralis*) also disperses these seeds in Tahiti, but in Moorea the endemic fruit dove (*Ptilinopus purpuralis*) disperses seeds of these alien plants. In Fiji, the red-vented bulbul contributes to the spread of primary colonist weeds (Watling 1979). In New Caledonia, the red-vented bulbul is suspected of spreading seeds of another invasive species: *Schinus terebinthifolius*, as it

is often observed feeding on fruits (Spotswood et al. 2012; Thouzeau-Fonseca 2013).

Disturbance and impact on fauna

The list of animal species reported to be impacted by the red-vented bulbul is presented in Table 3. The list comprises 15 species of bird, one reptile and one insect. Only one study addressed the issue of how the aggressive behavior of the red-vented bulbul affected the other bird species (Pernetta and Watling 1978).

On Oahu (Hawaii), direct predation of the monarch butterfly (*Danaus plexippus*) by the red-vented bulbul led to an induced color selection against the orange morph in the monarch (Stimson and Berman 1990). After 10 years, the same authors reported a predation transfer to the larvae, leading to an overall decline in abundance of the butterfly (Stimson and Kasuya 2000). In Tahiti, red-vented bulbuls are considered a threat to the Tahiti monarch (*Pomarea nigra*), an endemic and critically endangered passerine, through competition for nest sites and territory (Blanvillain et al. 2003).

Table 3 List of animal species reported as being impacted by the red-vented bulbul *Pycnonotus cafer*, with associated locations, inter-specific relationship, reported impact, method and references

Species	Countries	Islands	Inter-specific relationship	Reported impact	Method	References
Insects						
<i>Danaus plexippus</i>	H	O'ahu	Predation	Decline	Indirect Obs.	(1)
Birds						
<i>Pomarea nigra</i>	PF	Tahiti	Competition	Decline	Direct Obs.	(2)
<i>Lamprolia victoriae</i>	FJ	Vanua Levu	Competition	Decline	Hypothesis	(3)
<i>Myiagra vanikorensis</i>	FJ	Viti Levu	Aggressivity/competition	Nest parasitism	Monitoring	(4), (5)
<i>Lalage maculosa</i>	FJ	Viti Levu	Aggressivity	NA	Monitoring	(5)
<i>Acridotheres tristis</i>	FJ	Viti Levu	Aggressivity	NA	Monitoring	(5)
<i>Acridotheres fuscus</i>	FJ	Viti Levu	Aggressivity	NA	Monitoring	(5)
<i>Zosterops lateralis</i>	FJ	Viti Levu	Aggressivity	NA	Monitoring	(5)
<i>Amandava amandava</i>	FJ	Viti Levu	Aggressivity	NA	Monitoring	(5)
<i>Streptopelia chinensis</i>	FJ	Viti Levu	Aggressivity	NA	Monitoring	(5)
<i>Foulehaio carnunculata</i>	FJ, AS	Viti Levu; Tutuila	Aggressivity/competition	NA	Monitoring	(5), (6)
<i>Myzomela cardinalis</i>	AS	Tutuila	Aggressivity/competition	NA	Direct Obs.	(6)
<i>Pycnonotus leucogenys</i>	AE, BH		Cross-breeding	Setrile hybrids	Obs./ Hypothesis	(7), (8)
<i>Pycnonotus leucotis</i>	KW, QA, IR	Kish Island	Cross-breeding	NA	Obs./ Hypothesis	(9), (10), (11)
<i>Pycnonotus xanthopygos</i>	UAE		Cross-breeding	NA	Obs./ Hypothesis	(8)
<i>Zosterops xantochroa</i>	NC	Grande-Terre	Competition	NA	Hypothesis	(12)
Reptile						
<i>Hemidactylus frenatus</i>	NC	Grande-Terre	Predation	NA	Direct. Obs	<i>Pers. Obs.</i>

H Hawaii, PF French Polynesia, FJ Fiji, AS American Samoa, AE United Arab Emirates, BH Bahrain, KW Kuwait, QA Qatar, IR Iran, NC New Caledonia

(1) Stimson and Berman 1990, (2) Thibault et al. (2002), (3) Williams (2011), (4) Clunie (1976), (5) Pernetta and Watling (1978), (6) Sherman and Fall (2010), (7) Khamis (2010), (8) Kahn (1993), (9) Azin et al. (2008), (10) Gregory (2005), (11) Nation et al. (1997), (12) Hannecart and Letocart (1980)

In Fiji, several authors have reported red-vented bulbuls displaying aggressive behavior and competition for food resources towards other passerine species (Clunie 1976; Pernetta and Watling 1978; Williams 2011). However, Watling (1979) suspected that the observed confinement of native bird species to forest was mainly due to habitat loss rather than the aggressive behavior of the red-vented bulbul in Fiji. On Tutuila (American Samoa), Sherman and Fall (2010) observed that bulbuls competed for access to food resources with two passerine species. Finally, insect and skink predation by red-vented bulbuls is mentioned in several studies (VanderVelde 2002; Walker 2008; Brooks 2013). In the Middle East, cross-breeding between the exotic red-vented bulbul and the three closely related native species (white-cheeked bulbul, *P. leucogenys*; the white-eared bulbul (*P. leucotis*) and the yellow-vented bulbul, *P. xanthopygos*) is often reported as a potential threat for native bulbuls (Khan 1993; Nation et al. 1997; Gregory 2005; Azin et al. 2008; Khamis 2010).

Dispersal of neither endo- nor ecto-parasites by red-vented bulbul is well documented in its alien range (Table 4). In its native range, the red-vented bulbul is known to host *Isoospora* spp. (Boughton et al. 1938), *Menacanthus eurysternus* (Price 1975), *Bruelia guldum* and *Sturnidoecus guldum* (Ansari 1957) and *Pterotherpus pycnonoti* (Constantinescu et al., unpublished).

Table 4 P Parasite load of the red-vented bulbul *Pycnonotus cafer* in the literature

Type	Species	Host	References
Ecto-	<i>Isoospora</i> sp.	Yes	(1)
	<i>Menacanthus Eurysternus</i>	Yes	(2)
	<i>Bruelia guldum</i>	Yes	(3)
	<i>Sturnidoecus guldum</i>	Yes	(3)
Endo-	<i>Plasmodium</i> sp.	No	(4), (5)
	<i>Trypanosoma</i> sp.	No	(4)
	<i>Atoxoplasma</i> sp.	No	(4)
	<i>Chlamydia</i> sp.	Yes	(6)

Ecto- (Ectoparasites) corresponds to parasites living outside of the animal body. Conversely Endo- (Endoparasites) corresponds to parasites living inside the animal body

(1) Boughton et al. (1938), (2) Price (1975), (3) Ansari (1957), (4) Atkinson et al. (2006), (5) Jarvi et al. (2003), (6) Blanvillain et al. (2013)

In 1996, Jarvi et al. (2003) detected no avian malaria (*Plasmodium* spp.) in blood smears, and Atkinson et al. (2006) found no evidence of *Plasmodium*, *Trypanosoma*, *Atoxoplasma* or microfilaria. Red-vented bulbuls in Tahiti, however, have been found to carry the zoonotic disease *Chlamydia* sp. (Blanvillain et al. 2013).

Positive impacts

Red vented bulbuls feed on a variety of native plant species (Trail 1994; Sherman and Fall 2010), and dispersal of native seeds is the only service that has been explored in the bulbul's alien range (Spotswood et al. 2012). Interestingly, in a village-scale survey led by Daigneault and Brown (2013) in Viti Levu (Fiji), 47% of the respondents reported that the red-vented bulbul was good for their community and highlighted three main reasons. First, the bulbul was effective at insect control. Second, the bulbul reduced mongoose attacks on chickens. Third, village focus groups responded that red-vented bulbuls were occasionally eaten by villagers.

Management

The red-vented bulbul is considered an invasive species and environmental pest under the law in Australia (Tasmanian government 2010), Fiji (Minister of Primary Industries 1985), French Polynesia (Direction de l'environnement de la Polynésie Française 2016), Hawaii (Division of Forestry and Wildlife 2014), New Caledonia (Direction du Développement Economique et de l'Environnement 2008; Direction de l'ENVironnement de la Province Sud 2016), New Zealand (Ministry of Primary Industries 2017), South Africa (Department of Environmental Affairs 2016) and Spain (Ministerio de Agricultura, Alimentacion y Medio Ambiente 2013). In these countries, transportation, trade or possession of this species is forbidden, and hunting is authorized. We found no mention of this species as a pest or invasive species in other countries.

We found only three examples of management action taken against the red-vented bulbul in its alien range. The first one is the successful eradication program implemented in New Zealand between 1952 and 1955 (Turbott 1956). This program allowed the early detection and shooting of bulbuls thanks to a

reward associated with a call for information and led to an announcement of eradication in 1955 (Watling 1978). This management strategy remains in place in New Zealand and it helped prevent establishment following two more recent introduction events (September 2006 and February 2013).

Second, a cage test conducted in Hawaii on bird repellent showed that Ziram, Methiocarb and Methyl anthranilate reduced the consumption of treated papaya mash by red-vented bulbuls (Cummings et al. 1994). In an open-field test, the same authors showed that Methiocarb significantly reduced damages on orchids.

The third location where management actions have been implemented against the red-vented bulbul is the island of Tahiti in French Polynesia. In Tahiti, a management program that was not focused on red-vented bulbul management specifically, but rather on Tahiti monarch conservation, aimed to control alien birds. Pilot control campaigns were implemented twice, in 2012 and 2013 (Saavedra 2012, 2013), against the red-vented-bulbul and the common myna. These actions resulted in 1035 red-vented bulbuls being trapped in 2012, and 849 in 2013 and led to an increase in the breeding success of the Tahiti monarch (Saavedra 2013). Elsewhere in the French Polynesia archipelago, bulbul removal programs are in progress in Bora–Bora, Makatea and Nuku Hiva, three islands where the species is still rare but that are located near uninvaded parts of the archipelago.

In Fiji, a recent cost–benefit analysis of controlling the red-vented bulbul recommended “taking no action against the bulbul until such time as other benefits and or means of control have been field tested” (Daigneault and Brown 2013).

Discussion

The red-vented bulbul is still expanding its range into islands and continental areas across a wide geographic range between latitudes 22°N and 36°S. The number of references associated with this species outside its native range is also growing, but remains low compared to other species listed in the IUCN “100 world’s worst list”. As an example, searching for “*Acridotheres tristis*” in Google Scholar results in a four times larger output than the keywords “*Pycnonotus cafer*”. Based on this metric, the mallard and the red-

whiskered bulbul could have been included in the IUCN list in the same way as suggested in Martin-Albarracin et al. (2015). This reflects the heterogeneity in the attention paid to this “world’s worst invasive species”. In fact, more than half of the information we obtained came from just three island locations: Fiji, where the species was first transported; Hawaii, where it was responsible for huge economic losses; and French Polynesia, where it was considered to contribute to pressures on endemic biodiversity. Dispersal of the red-vented bulbul is strongly linked to human activities, as is the case for other bird species (Cassey et al. 2015). In Assam in the north-east of India, bulbul fights were part of a traditional and religious annual celebration until this was banned in January 2016. Wild bulbuls were trapped, kept in cages and prepared for the fights, and finally released if they won (Shalet 2016). The long and close relationship with humans led to the transportation of caged birds across the Pacific Ocean by Indian migrants from the early 20th century, first by boat, and then by airplane from the 1950s, certainly fostered the bulbul expansion (Hulme 2009). This was also a key period for invasion biology, with the publication of the Elton’s book (1958) marking the start of an increasing scientific interest in this field. While we found just eight references to this species between 1926 and 1966, 15 were published between 1967 and 1978. This species is still sold in local markets in several countries of the Arabic Peninsula (J. Babbington *pers.com.*), and bird trade remains the suspected principal vector of red-vented bulbul in this region.

Precise historical data are lacking regarding the propagule pressure, exact pathways of introduction, and dates associated with each introduction event, and we found very few records of this species being introduced but failing to establish. Globally, the establishment success recorded from Pacific islands to the USA or Europe suggests a better latitudinal plasticity of this species toward climate than expected when looking at the native distribution only. Moreover, its populations are considered to be self-sustaining or increasing in most of the tropical islands to which it has been introduced. Conversely, in most of the alien continental areas, population trends are considered steady or decreasing (ABBA database, Jennings 2004). This global pattern is consistent with the finding of Cassey et al. (2004) who showed that without consideration of the propagule pressure,

islands are significantly associated with introduction success and increased geographical range in birds.

Interest in introduced red-vented bulbuls grew rapidly in response to the considerable damage it caused on orchid production on Oahu, Hawaii, following its arrival in 1966. However, except for a few mentions of the cost associated with this issue (Cummings et al. 1994; Fox 2011), all references that reported damage to plant production referred only to species lists, inducing a lack in quantitative data on this impact category (Martin-Albarracin et al. 2015). Impact scores attributed to the red-vented bulbul in the study of Martin-Albarracin et al. (2015) were based on the two other impact categories. Seeds dispersal was demonstrated in three studies that explored the dispersal pattern of invasive plants such as *M. calvescence*. But these studies were all conducted in French Polynesia, and concluded that seed dispersal networks are complex and the interactions between native and alien plants and birds depend on both the frugivore community and on the relative abundance of available fruit (Spotswood et al. 2012). Negative impacts through competition also gain mention in three studies. Particularly, the aggressive behavior of red-vented bulbul was reported in Fiji and French Polynesia. In Tahiti, its aggressiveness toward adults of Tahiti monarchs (*P. nigra*) combined with predation by black rats (*Rattus rattus*) has contributed substantially to the decline in abundance of the critically endangered monarch species (Thibault et al. 2002). However, the same author reported that the main cause of the Tahiti monarch decline was predation by the black rat. The red-vented bulbul was blamed as a strong competitor because of its aggressive behavior, but rats, cats, and other bird species such as the common myna are also recognized as chick predators or nest competitors (Blanvillain et al. 2003; Ghestemme 2011). According to Saavedra (2012), the combined effects of the myna and red-vented bulbul were responsible for 35% of the nest failing of the Tahiti Monarch in 2012. Except for observed hybridization with its native cousins from the Pycnonotidae family in the Middle-East (Khan 1993; Nation et al. 1997), there are no reported impacts of red-vented bulbuls in continental areas (Khamis 2010; Brooks 2013). However, we reported some other potential impacts of the red-vented bulbul such as predation, hybridization, and dispersal of ecto- and endoparasites that were not included in any previous

impact scoring attempts. This highlights a large knowledge gap about how the inter-specific behavior of the red-vented bulbul impacts other species. Therefore, we believe that the role of the red-vented bulbul in the decline of plant or animal species is still to be demonstrated, or at least quantified, as has been done for other major invasive bird species such as the common myna (Lowe et al. 2011).

Moreover, positive effects or ecosystem services brought by introduced red-vented bulbuls have been poorly studied in its alien range, but may compensate to some degree for noxious impacts at the local scale (Daigneault and Brown 2013). Studies conducted in the bulbul native range confirmed part of this assessment. For example, it was shown that the bulbul was effective at insect control, including eating the widespread and highly polyphagous agricultural pest *Helicoverpa armigera* (Rana et al. 2014, 2017). By doing so, they improved curd and seed yields of cauliflower. The bulbul was also found to be an efficient pollinator of *Erythrina variegata* in India (Raju et al. 2004). Finally, an anti-predator response strategy that relies on eavesdropping of the bulbul's alarm call may also benefit other species such as *Emoia cyanurea*, a species of skink that is widespread throughout Pacific islands (Fuong et al. 2014). These few examples suggest that positive impacts may partly counterbalance the three categories of negative impacts attributed to the red-vented bulbul depending on the environment where the species occur.

For this reason, local-scale surveys led by Daigneault and Brown (2013) are crucial to inform local farmers and environment managers. We found few published studies dealing with the local management of the red-vented bulbul in its alien range. One is the biosecurity protocol currently in place in New Zealand (Watling 1978) that illustrates the efficiency of locally preventing alien species introductions on reducing their dispersal (Edelaar and Tella 2012). A test of bird repellents on Hawaiian orchids and papaya production demonstrated the efficiency of three chemicals (Cummings et al. 1994). In their study exploring the efficiency of bird repellent methods in the bulbul native range, Patyal and Rana (2003) highlighted nets as the most efficient methods although it can be costly to implement on large orchards. In their overview of birds impacts on Indian agriculture, Kale et al. (2012) reviewed the existing repellent techniques used against birds including the

red-vented bulbul, and underlined two main limits to their use being (1) social and ecological issues associated with killing birds and (2) danger of most chemical repellants for the biodiversity. This suggest that preventing damages of the red-vented bulbul on plants is feasible and that the investment intensity and the method used mostly depends on local communities. On the other hand, preventing impacts on seed dispersal and native fauna will rely on bird control programs and we found no feedbacks of such operations from the red-vented bulbul alien range yet. Results of the control programs currently in course in French Polynesia will certainly contribute to fill this gap (Saavedra 2013). In comparison, 13 eradication programs were conducted on islands against the common Myna and two against the red-whiskered bulbul that were mostly successful (DIISE 2015). Thus, more research is needed in the countries where the bulbul was introduced to evaluate threats associated with this species and guide adapted management strategies. Priority should be given to captive and field assessments of its diet and foraging ecology in its alien range. This would allow more accurate determination of the range of resources it uses and its prey (Bhatt and Kumar 2001), its role in seed dispersal (Spotswood et al. 2012), and its interspecific relationships (Bates et al. 2014).

Management strategies often rely on rigorous expert assessment and are mostly “restricted only to species for which there is already some suspicion of a threat, often an agricultural one” (Simberloff 2003). Even for suspected pests, risk assessment is often based on “anecdotal observations relating to small areas only” rather than direct scientific research (Strubbe et al. 2011). The alien range of the red-vented bulbul, mostly consisting of tropical islands, could have also contributed to the negative reputation of the bird as island ecosystems are especially sensitive to the arrival of alien species (Sax and Gaines 2008; Tershy et al. 2015). The high endemism and naivety of insular species accentuates their vulnerability (Gerard et al. 2016; Walsh et al. 2012). This sensitivity of tropical islands towards alien species may also be reinforced with the risk that a newly established population becomes a stepping stone for further introduction events through short-distance colonization (Gillespie et al. 2012). The information we present here supports this claim, with most reported impacts of red-vented bulbul on

biodiversity and plant production being from tropical islands, but even here the bulbul’s reported impacts are heterogeneous and typically non-specific. This work reveal that the red-vented bulbul remains highly understudied considering its invasive and pest status. The species’ long and close associations with people in its native range and subsequent transportation around the world as a cage-bird, coupled with its competitive foraging behavior (Sherman and Fall 2010), have surely contributed to its presence among the IUCN-ISSG list of the world’s worst invasive species, but this may well be overstated. Detailed and specific knowledge of this bulbul’s impacts and the threats it poses is essential, and Kumschick et al. (2015) recently insisted on the need for such information to inform the construction of global prioritization lists. In comparison, the red-whiskered bulbul or the mallard, for example, apparently attracted a more attention from both scientists and managers.

In conclusion, we found few references on the red-vented bulbul, reflecting a less attention paid by scientists to this species compared to the other world’s worst invasive species. The consideration of its negative impacts is largely influenced by few island locations whereas it is considered elsewhere as harmless, which prevent us from considering the bulbul as an absolute pest. Negative impacts led to the implementation of management programs in only one country and crop protection methods exist but are not necessarily used by local communities. Therefore, we suggest that the red-vented may not always be a dangerous pest.

Acknowledgements We thank the Global Avian Invasions Atlas program, the eBird community and the Ornithological Society of the Middle East for giving access to parts of their databases. Thanks to T. Ghestemme, J. Babbington, J. Eriksen, M. Pope, N. Morris, J. Buchan and M. Jennings for their assistance while summarizing information from the Middle East.

Compliance with ethical standards

Conflict of interest The authors state that they have no conflict of interest.

References

- Ali S, Ripley SD (1996) Handbook of the birds of India and Pakistan Together with those of Bangladesh Nepal, Bhutan And Sri Lanka. Oxford University Press, Oxford

- Ansari MAR (1957) Studies on ischnoceran Mallophaga infesting birds in the Panjab. *Indian J Entomol* 20:46–62
- Atkinson CT, Utzurrum RC, Seamon JO et al (2006) Hematozoa of forest birds in American Samoa—evidence for a diverse, indigenous parasite fauna from the South Pacific. *Pacific Conserv Biol* 12:229–237
- Azin F, Nosrati SM, Amini H (2008) Occurrence of the Red-vented Bulbul *Pycnonotus cafer* on Kish Island, North-eastern Persian Gulf, Iran. *Poocoes* 3:105–107
- Bates JH, Spotswood EN, Russell JC (2014) Foraging behaviour and habitat partitioning in sympatric invasive birds in French Polynesia. *Notornis* 61:35–42
- Becker RA, Wilks AR, Brownrigg R, Minka TP, Deckmyn A (2015a) maps: Draw Geographical Maps. R package version 3.0.0-2. <http://CRAN.R-project.org/package=maps>
- Becker RA, Wilks AR, Brownrigg R (2015b) mapdata: Extra Map Databases. R package version 2.2-5. <http://CRAN.R-project.org/package=mapdata>
- Berger AJ (1975) Red-whiskered and Red-vented bulbuls on Oahu. *Elepaio* 36:16–19
- Bergman DL, Chandler MD, Locklear A (2000) The economic impact of invasive species to wildlife services' cooperators. In: Clark L, Hone J, Shivik JA et al (eds) *Human Conflicts with Wildlife: Economic Considerations*. Proceedings of the Third NWRC Special Symposium, August 1–3, 2000, Fort Collins, CO, pp 169–178
- Bhatt D, Kumar A (2001) Foraging ecology of Red-vented Bulbul *Pycnonotus cafer* in Haridwar, India. *Forktail* 17:109–110
- Bivand R, Lewin-Koh N (2016) maptools: Tools for reading hling spatial objects. R package version 0.8-37. <http://CRAN.R-project.org/package=maptools>
- Blackburn TM, Pyšek P, Bacher S et al (2011) A proposed unified framework for biological invasions. *Trends Ecol Evol* 26:333–339. doi:10.1016/j.tree.2011.03.023
- Blackburn TM, Essl F, Evans T et al (2014) A unified classification of alien species based on the magnitude of their environmental impacts. *PLoS Biol* 12:e1001850. doi:10.1371/journal.pbio.1001850
- Blanvillain C, Salducci JM, Tutururai G, Maeura M (2003) Impact of introduced birds on the recovery of the Tahiti Flycatcher (*Pomarea nigra*), a critically endangered forest bird of Tahiti. *Biol Conserv* 109:197–205. doi:10.1016/S0006-3207(02)00147-7
- Blanvillain C, Saavedra S, Withers T (2013) Mission d'enquête sur la contamination de l'avifaune par salmonella enteritidis et d'autre pathogènes des volailles. Rapport SOP/DIREN, Tahiti, French Polynesia
- Blondel J, Hoffmann B, Courchamp F (2014) The end of invasion biology: intellectual debate does not equate to nonsensical science. *Biol Invasions* 16:977–979. doi:10.1007/s10530-013-0560-6
- Boughton DC, Boughton RB, Volk J (1938) Avian hosts of the genus *Isospora* (Coccidiida). *Ohio J Sci* 38:149–163
- Brochier B, Vangeluwe D, van den Berg T (2010) Alien invasive birds. *Rev Sci Tech OIE* 29:217–225
- Brooks DM (2013) Ecology, behavior, and reproduction of an introduced population of Red-vented Bulbuls (*Pycnonotus cafer*) in Houston, Texas. *Wilson J Ornithol* 125:800–808. doi:10.1676/13-037.1
- Burgiel SW, Perrault AM (2011) Black, white gray lists. In: Simberloff D, Rejmánek M (eds) *Encyclopedia of biological invasions*. University of California Press, London, pp 75–77
- Caplong P, Barjon F (2010) Le Bulbul à ventre rouge, une star méconnue en Nouvelle Calédonie. *La Calédonie Agricole* 123:22–25
- Carlson EA (1974) The avifauna of Tonga. Peace Corps, Tonga
- Cassey P, Blackburn TM, Sol D et al (2004) Global patterns of introduction effort and establishment success in birds. *Proc R Soc Lond B Biol Sci* 271:S405–S408. doi:10.1098/rsbl.2004.0199
- Cassey P, Vall-Llosera Camps M, Dyer E, Blackburn TM (2015) The Biogeography of Avian Invasions: history, accident and market trade. In: Canning-Clode J (ed) *Biological invasions in changing ecosystems: vectors, ecological impacts, management and predictions*. De Gruyter, Berlin, pp 37–54
- Clapp RB, Sibley FC (1966) Notes on the birds of Tutuila, American Samoa. *Notornis* 13:157–164
- Clunie F (1976) Red-vented Bulbul uses Vanikoro Broadbill nest. *Notornis* 23:263
- Cummings JL, Mason JR, Otis DL et al (1994) Evaluation of methiocarb, ziram, and methyl anthranilate as bird repellents applied to dendrobium orchids. *Wildl Soc B* 22:633–638
- Daigneault A, Brown P (2013) Invasive species management in the Pacific using survey data and benefit-cost analysis. In: 57th Australian agriculture and resource economics society annual conference. Australian Agriculture and Resource Economics Society, Sydney, Australia, pp 1–28
- Delacour J (1943) A revision of the genera and species of the family Pyconotidae (Bulbuls). *Zoologica* 28(1):17–28
- Department of environmental affairs (2016). Biodiversity (NEMBA) act 10 of 2004, List 4: National List of Invasive Bird Species. Pretoria, South Africa. <http://invasives.org.za/legislation/what-does-the-law-say> accessed on 10 April 2017
- Dhondt A (1976) Bird observations in Western Samoa. *Notornis* 23:29–43
- Dickinson E, Dekker R (2002) Systematic notes on Asian birds. 25. A preliminary review of the Pycnonotidae. *Zoologische Verhandelingen* 340:93–114
- Dickinson EC, Dekker RWRJ, Eck S, Somadikarta S (2002) Systematic notes on Asian birds. 26. Types of the Pycnonotidae. *Zoologische Verhandelingen* 340:115–160
- DIISE (2015) The database of island invasive species eradications, developed by Island Conservation, Coastal Conservation Action Laboratory UCSC, IUCN SSC Invasive Species Specialist Group, University of Auckland and Landcare Research New Zealand. <http://diise.islandconservation.org/>. Accessed 11 April 2017
- Direction de l'environnement de la Polynésie Française (2016) Code de l'environnement de la Polynésie Française, Section 2: Désignation des espèces menaçant la biodiversité. Papeete-Tahiti, Polynésie Française. <http://www.2dattitude.org/ressources/k2d/pdf/1/1D/1D05/1D05-01/1D05-01-01.pdf>. Accessed on 15 April 2017
- Direction de l'environnement de la Province Sud (2016) Code de l'environnement de la Province Sud, Livre II, Titre V, Article 250-2, IV: Liste des espèces animales exotiques envahissantes., Nouméa, Nouvelle-Calédonie. https://eprovince-sud.nc/sites/default/files/2016-04-Code_environne

- [ment_province_Sud_Avril_2016.pdf](#). Accessed on 15 April 2017
- Direction du développement économique et de l'environnement (2008) Code de l'environnement de la Province Nord, Livre II, Titre VI, Chap 1, Annexe à l'article 261-1: Liste des espèces envahissantes en province Nord., Koné, Nouvelle-Calédonie. <http://www.province-nord.nc/documents/deliberations/DDEE/Code%20Environnement.pdf>. Accessed on 15 April 2017
- Division of Forestry and wildlife (2014) Hawai'i administrative rules, Chap 124-Exhibit 4: Introduced wild birds other than game birds which have become established in the wild. Honolulu, Hawaii. <http://dlnr.hawaii.gov/dofaw/files/2013/09/Chap124a-Ex.pdf>. Accessed on 15 April 2017
- Donlan CJ, Wilcox C (2008) Assessing ecological responses to environmental change using statistical models. *J Appl Ecol* 45:1114–1123. doi:10.1111/j.1365-2664.2007.0
- Dyer EE, Redding DW, Blackburn TM (2017) The global avian invasions atlas—a database of alien bird distributions worldwide. *Sci Data* 4:0170041
- Edelaar PIM, Tella JL (2012) Managing non-native species: don't wait until their impacts are proven. *Ibis* 154:635–637
- Elton CS (1958) *The ecology of invasions by plants and animals*. Methuen, London
- Fox L (2011) Red-vented bulbul can alter ecosystems. *Maui News* 6
- Fuong H, Keeley KN, Bulut Y, Blumstein DT (2014) Heterospecific alarm call eavesdropping in nonvocal, white-bellied copper-striped skinks, *Emoia cyanura*. *Anim Behav* 95:129–135. doi:10.1016/j.anbehav.2014.07.005
- Gerard A, Jourdan H, Millon A, Vidal E (2016) Knocking on Heaven's door: are novel invaders necessarily facing naïve native species on islands? *PLoS ONE* 11:1–14. doi:10.1371/journal.pone.0151545
- Ghestemme T (2011) Impact of introduced birds on Tahiti Monarch. *PII Newsletter* 12-2011
- Gill BJ, Hunt GR, Sirgouant S (1995) Red-vented Bulbuls (*Pycnonotus cafer*) in New Caledonia. *Notornis* 42: 214–215
- Gillespie RG, Baldwin BG, Waters JM et al (2012) Long-distance dispersal: a framework for hypothesis testing. *Trends Ecol Evol* 27:47–56
- Gregory G (2005) *The birds of the State of Kuwait*. Cadeby Books, Grimsby
- Gren IM, Campos M, Gustafsson L (2016) Economic development, institutions, and biodiversity loss at the global scale. *Reg Environ Change* 16:445–457. doi:10.1007/s10113-015-0754-9
- Hannecart F, Letocart Y (1980) *Oiseaux de Nouvelle Calédonie et des Loyautés*. Cardinalis, Nouméa
- Hulme PE (2009) Trade, transport and trouble: managing invasive species pathways in an era of globalization. *J Appl Ecol* 46:10–18. doi:10.1111/j.1365-2664.2008.01600.x
- Islam K, Williams RN (2000) Red-vented Bulbul (*Pycnonotus cafer*) Red-whiskered Bulbul (*Pycnonotus jocosus*). In: *The Birds of North America* number 520
- Jarvi SI, Farias MEM, Baker H et al (2003) Detection of avian malaria (*Plasmodium* spp.) in native land birds of American Samoa. *Conserv Genet* 4:629–637. doi:10.1023/A:1025626529806
- Jennings MC (2004) Breeding birds in Central Arabia 1978–2003. *Sandgrouse* 26:35–47
- Jennings MC (2010) Atlas of the breeding birds of Arabia. *Fauna of Arabia* 25:1–751
- Kale M, Balfors B, Mörtberg U, Bhattacharya P, Chakane S (2012) Damage to agricultural yield due to farmland birds, present repelling techniques and its impacts: an insight from the Indian perspective. *J Agric Technol* 8:49–62
- Keane RM, Crawley MJ (2002) Exotic plant invasions and the enemy release hypothesis. *Trends Ecol Evol* 17:164–170. doi:10.1016/S0169-5347(02)02499-0
- Khamis A (2010) Alien species in Bahrain the overseen threats. *Bahrain Natural History Society Newsletter* number 25
- Khan MAR (1993) The introduced but naturalized avifauna of the United Arab Emirates. *J Bombay Nat Hist Soc* 90:437–445
- Kumschick S, Nentwig W (2010) Some alien birds have as severe an impact as the most effectual alien mammals in Europe. *Biol Conserv* 143:2757–2762. doi:10.1016/j.biocon.2010.07.023
- Kumschick S, Bacher S, Dawson W et al (2012) A conceptual framework for prioritization of invasive alien species for management according to their impact. *NeoBiota* 15:69–100. doi:10.3897/neobiota.15.3323
- Kumschick S, Blackburn TM, Richardson DM (2015) Managing alien bird species: time to move beyond '100 of the worst' lists? *Bird Conserv Int* 26:1–10. doi:10.1017/S0959270915000167
- Lendon A (1952) Bulbuls in Melbourne. *Emu* 52:67–68
- Long JL (1981) *Introduced birds of the world: The worldwide history, distribution and influence of birds introduced to new environments*. Universe Books, New York
- Lowe S, Browne M, Boudjelas S, De Poorter M (2000) *100 of the World's Worst Invasive Alien Species A selection from the Global Invasive Species Database*. The Invasive Species Specialist Group (ISSG), Auckland
- Lowe K, Taylor C, Major R (2011) Do common mynas significantly compete with native birds in urban environments? *J Ornithol* 152:1–11
- Mack RN, Simberloff D, Lonsdale MW, Evans H, Clout M, Bazzaz FA (2000) Biotic invasions: causes, epidemiology, global consequences, and control. *Ecol Appl* 10:689–710
- Martin-Albarracín VL, Amico GC, Simberloff D, Nuñez MA (2015) Impact of non-native birds on native ecosystems: a global analysis. *PLoS ONE* 10:e0143070. doi:10.1371/journal.pone.0143070
- McIlroy D, Brownrigg R, Minka TP, Biv R (2015) *mapproj: Map Projections*. R package version 1.2-4. <http://CRAN.R-project.org/package=mapproj>
- Metzdorf N, Brescia F (2008) Impact de l'avifaune et des roussettes sur les productions fruitières en Nouvelle-Calédonie, Etude préliminaire – Etat des lieux et recommandations. IAC/Axe2, Paita Unpublished report
- Meyer J-Y (1993) Dispersion de *Miconia calvescens* par les oiseaux dans les îles de la société. *Seminaire Manu (Connaissance et Protection des Oiseaux en Polynésie orientale)*. Société d'Ornithologie de Polynésie, Papeete
- Meyer J-Y (1996) Status of *Miconia calvescens* (Melastomaceae), a dominant invasive tree in the society Islands (French Polynesia). *Pac Sci* 50:66–79

- Minister of primary industries (1985). Laws of Fiji, Chapter 170: Birds and game protection. Viti-Levu, Fiji. http://www.paclii.org/fj/legis/consol_act_OK/bagpa231/ accessed on 15 April 2017
- Ministerio de Agricultura, Alimentacion y Medio Ambiente (2013) *Pycnonotus cafer*. Catálogo español de especies exóticas invasoras. Ministerio de Agricultura, Alimentación y Medio Ambiente, Madrid
- Ministry of primary industries (2017) Biosecurity Act 1993. Wellington, New Zealand. <http://www.legislation.govt.nz/act/public/1993/0095/latest/whole.html#DLM315361> accessed on 15 April 2017
- Nation B, Nation H, Hooper A (1997) Birds New to Qatar. Sandgrouse 19:56–62
- Parham BEV (1955) Birds as pests in Fiji. J Fijian Agric 25:9–14
- Patyal SK, Rana RS (2003) Damage potential and abundance of avian fauna associated with grapes in Indian Mid Hills of Himachal Pradesh. In: VII international symposium on temperate zone fruits in the tropics. ISHS Acta Horticulturae, Nauni, Solan, India
- Pedersen T, Aspinall SJ (compilers) (2015) EBRC Annotated Checklist of the birds of the United Arab Emirates. Ornithological Society of the Middle East, the Caucasus and Central Asia. Sandy, Bedfordshire, UK
- Pereira HM, Navarro LM, Martins IS (2012) Global biodiversity change: the bad, the good, and the unknown. Annu Rev Environ Resour 37:25–50. doi:10.1146/annurev-environ-042911-093511
- Pereyra PJ (2016) Revisiting the use of the invasive species concept: an empirical approach. Austral Ecol 41:519–528. doi:10.1111/aec.12340
- Pernetta JC, Watling D (1978) the introduced and native terrestrial vertebrates of Fiji. Pac Sci 32:223–244
- Pimentel D (2011) Biological invasions: economic and environmental costs of alien plant, animal, microbe species. CRC Press, Boca Raton
- Pimentel D, Zuniga R, Morrison D (2005) Update on the environmental and economic costs associated with alien-invasive species in the United States. Ecol Econ 52:273–288. doi:10.1016/j.ecolecon.2004.10.002
- Price RD (1975) The *Menacanthus eurysternus* Complex (Mallophaga: menoponidae) of the Passeriformes and Piciformes (Aves). Ann Entomol Soc Am 68:617–622
- Pyšek P, Richardson DM, Pergl J et al (2008) Geographical and taxonomic biases in invasion ecology. Trends Ecol Evol 23:237–244. doi:10.1016/j.tree.2008.02.002
- Raju AJS, Rao SP, Zafar R, Roopkalpana P (2004) Passerine bird-pollination and fruiting behavior in *Erythrina variegata* L. (Fabaceae) in the Eastern Ghats, India. Beitr Biol Pflanzen 73:321–330
- Rana RS, Chand J, Patyal SK (2014) Evaluation of birds as predators of insect pests infesting cauliflower. J Commun Mobil Sustain Dev 9:18–22
- Rana R, Chand J, Patyal S, Sharma K (2017) Studies on the role of insectivorous birds in managing insect pests of cabbage (*Brassica oleracea* var capitata L.). Int J Farm Sci 6:245–253
- Ricciardi A, Hoopes MF, Marchetti MP, Lockwood JL (2013) Progress toward understanding the ecological impacts of nonnative species. Ecol Monogr 83:263–282. doi:10.1890/13-0183.1
- Richardson DM, Ricciardi A (2013) Misleading criticisms of invasion science: a field guide. Divers Distrib 19:1461–1467. doi:10.1111/ddi.12150
- Russell JC, Blackburn TM (2017) The rise of invasive species denialism. Trends Ecol Evol 32(1):3–6
- Saavedra S (2012) First control campaign for Common myna and Red vented bulbul on Tahiti. Internal Report, SOP Manu, Tahiti
- Saavedra S (2013) Second control campaign for Common myna and Red vented bulbul on Tahiti. Internal Report, SOP Manu, Tahiti
- Sala OE, Stuart Chapin F et al (2000) Global biodiversity scenarios for the year 2100. Science 287:1770–1774
- Sax DF, Gaines SD (2008) Species invasions and extinction: the future of native biodiversity on islands. Proc Natl Acad Sci USA 105:11490–11497. doi:10.1073/pnas.0802290105
- Saxena A (2015) Growing concern and threat of invasive alien species on natural ecosystem and native species. Int J Med Pharm Res 3:1246–1248
- Secretariat CBD (2010) The Strategic Plan for Biodiversity 2011–2020 and the Aichi Biodiversity Targets. In: Document UNEP/CBD/COP/DEC/X/2. Secretariat of the Convention on Biological Diversity, Nagoya, Japan
- Shalet J (2016) India: Guwahati High Court stays bulbul fight, protests erupt in Assam. International Business Time 14-01-2016. <http://www.ibtimes.co.uk/india-guwahati-high-court-stays-bulbul-fight-protests-erupt-1537937> accessed on 25 January 2016
- Sherman J, Fall P (2010) Observations on feeding frequencies among native and exotic birds and fruit bats at *Erythrina variegata* and *Dysoxylum* trees on American Samoa. Australasian National University Press, Canberra
- Simberloff D (2003) Confronting introduced species: a form of xenophobia? Biol Invasions 5:179–192. doi:10.1023/A:1026164419010
- Simberloff D, Vitule JRS (2014) A call for an end to calls for the end of invasion biology. Oikos 123:408–413. doi:10.1111/j.1600-0706.2013.01228.x
- Spotswood EN, Meyer J-Y, Bartolome JW (2012) An invasive tree alters the structure of seed dispersal networks between birds and plants in French Polynesia. J Biogeogr 39:2007–2020. doi:10.1111/j.1365-2699.2012.02688.x
- Spotswood EN, Meyer J-Y, Bartolome JW (2013) Preference for an invasive fruit trumps fruit abundance in selection by an introduced bird in the Society Islands, French Polynesia. Biol Invasions 15:2147–2156. doi:10.1007/s10530-013-0441-z
- Stimson J, Berman M (1990) Predator Induced Color Polymorphism in *Danaus plexippus* L (Lepidoptera, Nymphalidae) in Hawaii. Heredity 65:401–406
- Stimson J, Kasuya M (2000) Decline in the frequency of the white morph of the monarch butterfly (*Danaus plexippus plexippus* L., Nymphalidae) on Oahu, Hawaii. J Lepid Soc 54:29–32
- Strubbe D, Schwartz A, Chiron F (2011) Concerns regarding the scientific evidence informing impact risk assessment and management recommendations for invasive birds. Biol Conserv 144:2112–2118
- Sullivan BL, Wood CL, Iliff MJ, Bonney RE, Fink D, Kelling S (2009) eBird: a citizen-based bird observation network in the biological sciences. Biol Conserv 142:2282–2292

- Tasmanian government (2010) Wildlife (General) Regulations, S.R. 2010, NO. 113 – Schedule 5: Restricted animals., Hobart, Tasmania. http://www.austlii.edu.au/au/legis/tas/num_reg/wr20102010n113319/sch5.html accessed on 15 April 2017
- Tershy BR, Shen KW, Newton KM et al (2015) The importance of islands for the protection of biological and linguistic diversity. *Bioscience* 65:592–597. doi:[10.1093/biosci/biv031](https://doi.org/10.1093/biosci/biv031)
- Thibault JC, Martin JL, Penloup A, Meyer JY (2002) Understanding the decline and extinction of monarchs (Aves) in Polynesian Islands. *Biol Conserv* 108:161–174. doi:[10.1016/S0006-3207\(02\)00102-7](https://doi.org/10.1016/S0006-3207(02)00102-7)
- Thouzeau-Fonseca C (2013) Contribution à la gestion d'un oiseau introduit envahissant en Nouvelle-Calédonie, le bulbul à ventre rouge (*Pycnonotus cafer* L.): étude des modalités de piégeage et du comportement alimentaire en milieu naturel. IAC Production, Paita, Nouvelle-Calédonie
- Trail PW (1994) The Phenology of Rainforest plants in Tutuila, American Samoa. Department of Marine and Wildlife Resources Biological Report Series, Pago Pago, American Samoa
- Turbott EG (1956) Bulbuls in Auckland. *Notornis* 6:185–192
- Valéry L, Fritz H, Lefeuvre JC (2013) Another call for the end of invasion biology. *Oikos* 122:1143–1146. doi:[10.1111/j.1600-0706.2013.00445.x](https://doi.org/10.1111/j.1600-0706.2013.00445.x)
- VanderVelde N (2002) The Red-vented bulbul has come to Micronesia. *Aliens* 16:13–14
- Walker R (2008) The Red-vented bulbul, Superbird? *Elepaio* 68:71–78
- Walsh JC, Venter O, Watson JEM et al (2012) Exotic species richness and native species endemism increase the impact of exotic species on islands. *Glob Ecol Biogeogr* 21:841–850. doi:[10.1111/j.1466-8238.2011.00724.x](https://doi.org/10.1111/j.1466-8238.2011.00724.x)
- Walsh JR, Carpenter SR, Vander Zanden MJ (2016) Invasive species triggers a massive loss of ecosystem services through a trophic cascade. *Proc Natl Acad Sci USA* 113:201600366. doi:[10.1073/pnas.1600366113](https://doi.org/10.1073/pnas.1600366113)
- Watling D (1978) Observation on the naturalised distribution of the red-vented bulbul in the Pacific, with special reference to the Fiji islands. *Notornis* 25:109–117
- Watling D (1979) The Bulbul gets a clean bill. *New Sci* 81:963–965
- Williams G (2011) 100 alien invaders. Bradt Travel Guides, Buckinghamshire
- Wood CA, Wetmore A (1926) A collection of birds from the Fiji Islands. *Ibis* 68:91–136