



HAL
open science

Effect of concentrate feeding sequence on equine hindgut fermentation parameters

Sophie Sadet-Bourgeteau, Christelle Philippeau, Véronique Julliand

► To cite this version:

Sophie Sadet-Bourgeteau, Christelle Philippeau, Véronique Julliand. Effect of concentrate feeding sequence on equine hindgut fermentation parameters. *Animal*, 2017, 11 (07), pp.1146 - 1152. 10.1017/S1751731116002603 . hal-01795068

HAL Id: hal-01795068

<https://hal.science/hal-01795068>

Submitted on 13 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Effect of concentrate feeding sequence on equine hindgut fermentation parameters

S. Sadet-Bourgeteau[†], C. Philippeau and V. Julliand

AgroSup Dijon, URANIE, USC1335 Nutrition du cheval athlète, 26 Bd Docteur Petitjean, F-21079 Dijon, France

(Received 29 January 2016; Accepted 3 November 2016)

Feeding a diet with a small amount of hay and a high proportion of concentrate given in large meals entails a risk of colic to horses as this can impact the hindgut microbial ecosystem. To counteract this potential negative effect, one feeding strategy is to modify the distribution sequence of concentrate and hay. The purpose of the present study was to assess the impact of feeding the concentrate meal before or with the hay meal on fermentative parameters postprandial variations in caecum and right ventral colon contents and faeces. We focused on the comparison of pH value and volatile fatty acid (VFA) concentrations between the three compartments. Four horses fitted with two cannulas in the caecum and right ventral colon were fed a 100% hay diet (H) for a 3-week period, and then a 40% concentrate–60% hay diet for two consecutive periods of 3 weeks each. Concentrate and hay were offered simultaneously in two equal meals at 0800 and 1700 h (HC1) during one period, or separately in two equal meals at 0800 and 1600 h for the concentrate and at 1000 and 1730 h for hay (HC2) during the second one. During the 2nd week of each diet period (H, HC1, HC2), caecal, right ventral colonic and faecal samples were collected before (0) and at 2, 4, 6 and 8 h after the morning meal. pH values and VFA concentrations were measured. Adding concentrate to the hay was associated with a decrease of pH value in the caecum, right ventral colon and faeces and an increase of faecal VFA concentrations ($P < 0.01$). An impact of the concentrate feeding sequence was observed on caecal pH and faecal propionate concentration ($P < 0.001$) but none on the other parameters investigated. In the caecum and right ventral colon, pH values decreased and VFA concentrations increased within the range of 4 to 8 h after the morning meal. Our data suggested that the variation of faecal pH could be an indicator of those of the caecal and right ventral colonic pH with a delay of 2 h. The present study showed that hay and commercial pelleted could be safely distributed simultaneously, thus making easier every day horse feeding.

Keywords: concentrate feeding sequence, fermentative parameters, hindgut, horse, postprandial evolution

Implications

In riding centres horses are fed low quantity of hay, with high amounts of concentrate in a limited number of meals that can increase the incidence of colic. To limit these potential negative effects, the distribution sequence of a concentrate meal containing 33% starch before or with the hay meal was studied and compared with a hay diet. Our results suggested that to simplify feeding practice, hay and concentrate could be distributed simultaneously without provoking digestive disturbances.

Introduction

Horses possess a great number and variety of microbes in their voluminous hindgut. These microbes are essential

for utilizing dietary plant fibre and converting them into volatile fatty acids (VFAs) that provide energy to the host and for preserving a healthy environment of the intestinal ecosystem. To sustain the efficient functioning of the hindgut microbes, it has long been recommended to feed at least 10 g of hay dry matter (DM)/kg BW per day (Miyaji *et al.*, 2008). Recent recommendation proposed increasing the minimal amount of roughage to 20 g (84% to 90% DM)/kg BW per day (Coenen *et al.*, 2011). In most riding centres in France, the quantity of hay distributed to horses is often lower. Combined with large meals of concentrate fed to meet horses' exercise energy requirements, such diets may be associated with increased risk of colic (Goncalves *et al.*, 2002; Archer and Proudman, 2006). To limit this potential negative impact, one strategy is to change the distribution sequence of concentrate and hay. Indeed, compared with feeding concentrate before hay, distributing hay first has been reported to lessen the hindgut physicochemical

[†] E-mail: sophie.bourgeteau-sadet@agrosupdijon.fr

environment alterations which may disturb the caecal (Jensen *et al.*, 2012), colonic (Philippeau *et al.*, 2009) and faecal (Zeyner *et al.*, 2004) microbial ecosystems. The impact of feeding both hay and concentrate simultaneously has not been studied on the equine gastrointestinal health, although in the field, for practical issues, this is a common practice. The purpose of the present study was thus to assess the impact of distributing the concentrate meal before or with the hay meal on postprandial variations in pH and VFA from different hindgut compartments and faeces. These fermentation parameters can serve as an indicator for microbial activity (Santos *et al.*, 2011).

Material and methods

The experiment was conducted at the horse research facilities of AgroSup Dijon under approval from the Ethics Committee for Animal Well-Being of the University of Burgundy, Dijon, France (agreement no. C1311).

Animals, management and diets

Four adult cross-bred geldings (average BW of 500.0 ± 35.7 kg and mean age 11.5 ± 3.2 years) were individually housed in 3×3 m free stalls bedded with wood shavings. They were dewormed (Eraquell-Virbac SA, Glattbrugg, Switzerland) before the beginning of the trial. Except on collection days, horses had access to a sand paddock for free exercise 5 h/day (1200 to 1700 h), 6 days each week and were exercised with an automatic walker for 1 h/day (1 h after the morning meal), five times per week. Each animal was fitted with caecum and right ventral colon polyvinyl chloride cannulas (30 mm).

Horses were fed a hay diet (100% meadow hay) during the first 3 weeks (H) of the experimental trial, a hay–concentrate diet (40% concentrate and 60% meadow hay; DM basis) with both concentrate and hay distributed at the same time (HC1 diet) during the following 3 weeks, and finally with concentrate offered before hay (HC2 diet) during the last 3 weeks of the study. The concentrate was a commercial pelleted feed (Hippo 63; Royal Horse, Vannes, France). The chemical composition of the diets is presented in Table 1. The diets H, HC1 and HC2 were fed at 1.8 kg DM/100 kg BW. For H and HC1 diets, hay or hay plus concentrate were offered daily in two equal meals at 0800 and 1700 h. For the diet HC2, animals were fed concentrate at 0800 and 1600 h and hay at 1000 and 1730 h. Each experimental period (H, HC1 and HC2) was separated by 10 days, where horses were fed the hay diet, followed by 10 days where horses were progressively adapted to the next diet. Salt blocks and water were available *ad libitum*.

Measurements and analyses

For each period on day 15, digestive contents were collected before (0) and at 2, 4, 6 and 8 h after the first morning meal was given (hay or hay plus concentrate or concentrate morning meal for periods H, HC1 and HC2, respectively).

Table 1 Ingredients and mean chemical composition of the diets

Items	Diets ¹		
	H	HC1	HC2
Ingredient (DM) (%)			
Hay	100	60	60
Concentrate (Hippo 63; Royal Horse)	–	40	40
Chemical composition (DM basis)			
DM (%)	91.3	89.9	89.9
OM (%)	94.4	92.0	92.0
NDF (%)	59.5	33.3	33.3
ADF (%)	35.1	17.4	17.4
ADL (%)	5.1	4.1	4.1
CP (%)	6.3	14.7	14.7
Starch (%)	0	12.9	12.9
Daily consumption			
DM intake (g/100 kg BW)	1.8	1.8	1.8

DM = dry matter; OM = organic matter.

¹Diet H: 100% forage diet; diet HC: 40% concentrate/60% forage; for diets H and HC1, feeds were offered daily in two equal meals at 0800 and 1700 h; for the diet HC2, concentrate was offered at 0800 and 1600 h and hay at 1000 and 1730 h.

Caecal and right ventral colonic digesta samples were collected by gravity via the cannulas. Faecal samples were grabbed in the rectum. All digestive contents were filtered (Blutax nylon filter 100 μ m; Tripette and Renaud, Paris, France) and immediately frozen (-20°C) for further analysis of VFA (1 ml added with 0.1 ml of a preservative mixture of 5% H_3PO_4 + 1% HgCl_2).

The pH of caecal, colonic and filtered faecal content was measured immediately after each collection, using an electronic pH meter (pH 340i/SET; Wissenschaftlich-Technische Werkstätten, Weilheim, Germany). Volatile fatty acid concentrations were determined using GLC (Gas chromatograph model Clarus 500; PerkinElmer, Courtaboeuf, France) (Jouany, 1982).

Statistical analyses

Data were processed by ANOVA using the MIXED procedure of SAS v9.3 (SAS Institute Inc., Cary, NC, USA). For each digestive segment, the statistical model included horse, diet, sampling time and diet \times sampling time. Horse was considered as random effect. A repeated-measures ANOVA was performed to compare profiles of fermentation parameters (VFA concentrations, proportions and pH) using the repeated time option of SAS v9.3. All statistical procedures were performed separately for the caecum, colon and faeces with horse as experimental unit. Least-squares means were compared using a Tukey's test, and effects declared significant at $P < 0.05$.

Volatile fatty acid concentrations and pH were also analyzed using a principal component analysis (PCA) performed on the covariance matrix of the samples (caecum, right ventral colon and faeces). Principal component analysis and the Monte Carlo test (10 000 permutations) were performed using the ADE-4 software (Thioulouse *et al.*, 1997).

Volatile fatty acid proportions and pH were also analyzed using a PCA; however, the same effects than those observed on concentrations were highlighted (data not shown). Thus, in the present study only PCA analysis carried on VFA concentrations and pH was presented. Each PCA analysis has been made with 180 data (3 sites \times 3 diets \times 4 animals \times 5 h of sampling).

Results

All horses remained clinically healthy throughout the study.

Postprandial evolution in the caecum, colon and faeces

Irrespective of the digestive segments, no significant effect of time of sampling, diet and their interaction were found on VFA proportions nor on the ratio ((acetate + butyrate)/propionate) (data not shown).

In the caecum, no interaction of the time of sampling and the diet was detected for pH values and VFA concentrations (Table 2, $P > 0.05$). pH values and butyrate concentration depended on the diet ($P < 0.001$). Regardless of the time of sampling, caecal pH value was in mean significantly lower when horses were fed diet HC2 than diets H or HC1 ($P < 0.001$). It was significantly greater in horses fed H diet compared with HC1 diet. Butyrate concentration was significantly lower in horses receiving diet H than diets HC1

and HC2 ($P < 0.001$). With the concentrate diets there was a high trend for acetate ($P = 0.051$) and propionate ($P = 0.052$) decrease.

pH values, total VFA, acetate and propionate concentrations differed significantly with sampling time ($P < 0.001$). Whatever the diet, caecal pH decreased significantly 4 h after the morning meal and stayed at the minimum value until 8 h postprandial ($P < 0.001$). Caecal, total VFA, acetate and propionate concentrations increased after feeding to reach a maximum from 4 to 8 h postprandial ($P < 0.001$).

In the right ventral colon, no interaction was found between the time of sampling and the diet (Table 3, $P > 0.05$). pH values and butyrate concentrations depended on the diet ($P < 0.001$). The colonic pH value was significantly lower when horses were fed diets HC1 and HC2 than diet H ($P < 0.001$). Butyrate concentration was significantly lower in horses fed diet H than diets HC1 and HC2 ($P < 0.001$). Butyrate concentrations remained stable over time ($P = 0.121$). Colonic pH decreased within the first 4 h postprandial and remained minimum between 4 and 8 h after the meal ($P < 0.001$). Colonic acetate, propionate and total VFA concentrations increased after feeding to reach a maximum between 4 and 8 h postprandial ($P < 0.01$).

In the faeces, no interaction between the time of sampling and the diet was found (Table 4, $P = 0.882$). pH values and all VFA concentrations depended on the diet (Table 4, $P < 0.001$). The faecal pH value was significantly lower when

Table 2 Postprandial evolution of caecal pH and volatile fatty acid (VFA) concentrations of horses given three different diets

Diets ¹	Time of sampling (h)					SEM	P-value		
	0	2	4	6	8		D	T	D \times T
pH									
H	7.0 ^{Aa}	6.9 ^a	6.7 ^c	6.7 ^c	6.7 ^c	0.02	<0.001	<0.001	0.253
HC1	6.9 ^B	6.8	6.5	6.5	6.5				
HC2	6.7 ^C	6.8	6.5	6.4	6.4				
Total VFA (mmol/l)									
H	62.2 ^{Ac}	62.1 ^c	78.1 ^{ab}	81.4 ^a	73.0 ^{ab}	1.64	0.075	<0.001	0.616
HC1	61.9 ^A	53.4	70.3	75.9	78.9				
HC2	50.4 ^A	55.4	69.9	73.2	76.8				
Acetate (mmol/l)									
H	48.3 ^{Ac}	45.2 ^c	59.0 ^{ab}	62.6 ^a	55.8 ^{ab}	1.29	0.051	<0.001	0.685
HC1	45.8 ^A	40.6	51.3	54.7	57.4				
HC2	36.7 ^A	42.2	50.5	56.7	57.3				
Propionate (mmol/l)									
H	11.5 ^{Ac}	11.9 ^c	15.7 ^{ab}	15.3 ^a	13.9 ^{ab}	0.37	0.052	<0.001	0.202
HC1	11.4 ^A	8.7	14.8	16.6	17.3				
HC2	10.3 ^A	9.9	13.9	11.4	14.5				
Butyrate (mmol/l)									
H	0.5 ^{Ba}	4.1 ^a	1.2 ^a	1.3 ^a	1.5 ^a	0.23	<0.001	0.583	0.051
HC1	3.6 ^A	2.6	3.8	4.3	3.7				
HC2	2.5 ^A	2.3	4.5 ^{Aa}	3.6	3.6				

D = diet; T = time of sampling.

^{A,B}Within a column, means without common superscript letters differ ($P < 0.05$) and are representative of diet effect.

^{a,b,c}Within a row, means without common superscript letters differ ($P < 0.05$) and are representative of time of sampling effect.

¹Diet H: 100% forage diet; diet HC: 40% concentrate/60% forage; for diets H and HC1, feeds were offered daily in two equal meals at 0800 and 1700 h; for the diet HC2, concentrate was offered at 0800 and 1600 h and hay at 1000 and 1730 h.

Table 3 Postprandial evolution of colonic pH and volatile fatty acid (VFA) concentrations of horses given three different diets

Diets ¹	Time of sampling (h)					SEM	P-value		
	0	2	4	6	8		D	T	D × T
pH									
H	7.0 ^{Aa}	6.9 ^{ab}	6.7 ^c	6.7 ^c	6.7 ^c				
HC1	6.8 ^B	6.8	6.6	6.5	6.5	0.02	<0.001	<0.001	0.390
HC2	6.8 ^B	6.8	6.5	6.4	6.5				
Total VFA (mmol/l)									
H	70.2 ^{Abc}	79.7 ^c	84.9 ^{abc}	77.4 ^{ab}	88.2 ^a				
HC1	67.2 ^A	55.4	73.3	85.3	84.2	2.45	0.418	<0.001	0.455
HC2	70.0 ^A	51.7	79.1	91.5	89.8				
Acetate (mmol/l)									
H	54.2 ^{Aab}	61.6 ^b	65.7 ^{ab}	61.2 ^a	68.7 ^a				
HC1	49.5 ^A	41.4	53.2	62.6	61.6	1.95	0.140	<0.01	0.615
HC2	53.1 ^A	40.2	59.2	70.3	67.2				
Propionate (mmol/l)									
H	12.7 ^{Abc}	13.4 ^c	15.3 ^{ab}	13.0 ^{ab}	15.5 ^a				
HC1	11.5 ^A	9.8	14.6	16.9	16.5	0.45	0.551	<0.001	0.323
HC2	11.8 ^A	7.9	14.3	14.1	15.7				
Butyrate (mmol/l)									
H	0.6 ^{Ba}	0.8 ^a	1.4 ^a	0.6 ^a	1.4 ^a				
HC1	4.3 ^A	3.1	4.5	4.8	4.9	0.29	<0.001	0.135	0.891
HC2	3.4 ^A	2.6	4.0	4.7	5.0				

D = diet; T = time of sampling.

^{A,B}Within a column, means without common superscript letters differ ($P < 0.05$) and are representative of diet effect.^{a,b,c}Within a row, means without common superscript letters differ ($P < 0.05$) and are representative of time of sampling effect.¹Diet H: 100% forage diet; diet HC: 40% concentrate/60% forage; for diets H and HC1, feeds were offered daily in two equal meals at 0800 and 1700 h; for the diet HC2, concentrate was offered at 0800 and 1600 h and hay at 1000 and 1730 h.**Table 4** Postprandial evolution of faecal pH and volatile fatty acid (VFA) concentrations of horses given three different diets

Diets ¹	Time of sampling (h)					SEM	P-value		
	0	2	4	6	8		D	T	D × T
pH									
H	6.8 ^{Aa}	6.7 ^{ab}	6.8 ^{abc}	6.8 ^{bc}	6.7 ^c				
HC1	6.6 ^B	6.6	6.4	6.2	6.1	0.05	<0.01	0.048	0.889
HC2	6.6 ^B	6.6	6.5	6.2	6.3				
Total VFA (mmol/l)									
H	43.3 ^{Ba}	49.2 ^a	44.4 ^a	53.6 ^a	50.8 ^a				
HC1	60.8 ^A	61.3	77.1	81.7	82.2	2.68	<0.001	0.150	0.887
HC2	59.6 ^A	57.2	58.2	79.6	66.7				
Acetate (mmol/l)									
H	33.9 ^{Ba}	37.2 ^a	34.1 ^a	41.5 ^a	38.5 ^a				
HC1	45.0 ^A	46.1	57.4	61.0	61.1	2.03	<0.001	0.182	0.907
HC2	46.3 ^A	44.5	44.8	61.4	51.0				
Propionate (mmol/l)									
H	7.6 ^{Ba}	9.0 ^a	8.0 ^a	9.6 ^a	9.1 ^a				
HC1	10.8 ^A	10.6	13.7	14.0	14.3	0.47	<0.001	0.213	0.935
HC2	9.0 ^{AB}	8.9	9.4	12.4	11.0				
Butyrate (mmol/l)									
H	0.5 ^{Ba}	1.5 ^a	1.0 ^a	1.1 ^a	1.5 ^a				
HC1	3.2 ^A	3.0	3.9	4.4	4.3	0.21	<0.001	0.226	0.801
HC2	2.7 ^A	2.5	2.7	3.8	3.3				

D = diet; T = time of sampling.

^{A,B}Within a column, means without common superscript letters differ ($P < 0.05$) and are representative of diet effect.^{a,b,c}Within a row, means without common superscript letters differ ($P < 0.05$) and are representative of time of sampling effect.¹Diet H: 100% forage diet; diet HC: 40% concentrate/60% forage; for diets H and HC1, feeds were offered daily in two equal meals at 0800 and 1700 h; for the diet HC2, concentrate was offered at 0800 and 1600 h and hay at 1000 and 1730 h.




Figure 1 Comparison of pH values and volatile fatty acid concentration variability performed by principal component (PC) analysis. Data were obtained from caecum (CA), right ventral colon (CN) and faeces (FC) of horses ($n = 4$) given three different diets (H: 100% forage diet; HC: 40% concentrate/60% forage; for diets H and HC1, feeds were offered daily in two equal meals at 0800 and 1700 h; for the diet HC2, concentrate was offered at 0800 and 1600 h and hay at 1000 and 1730 h). Each line represents one observation. Principal component analysis was performed taking into account the digestive segment (a) or the diet (b).

horses were fed diets HC1 and HC2 than diet H ($P < 0.001$). The concentration of acetate, butyrate and total VFA were significantly higher when horses received diets HC1 and HC2 than diet H ($P < 0.0001$). No differences were observed between diets HC1 and HC2 ($P > 0.05$). The concentration of propionate was higher when horses were fed diet HC1 than diet H ($P < 0.001$). No differences were observed between diet H and diet HC2 ($P > 0.05$). Faecal pH decreased regularly during the 8 h postprandial (Table 4, $P = 0.048$).

Global appraisal of the effects of gut location and diet

Principal component analysis data for the pH values and VFA concentrations of caecal, right ventral colonic and faecal samples are presented according to the gut location (Figure 1a) or the diet (Figure 1b). In both figures, the first axis (PC1) explained 63.7% of the total variability and the second axis (PC2) explained 19.9% of the total variability. pH values and VFA concentrations showed no differences

between the three gut locations (Figure 1a, Monte Carlo test: $P > 0.05$). However, irrespective of digestive segment considered, samples from H diet were separated from those of HC diets on PC1 and PC2 axes (Figure 1b). This discrimination on PC1 and PC2 was explained by acetate, propionate and total VFA concentrations on PC1 axis and by pH and butyrate concentration on PC2 axis (Figure 2) (Monte Carlo test: $P < 0.001$).

Discussion

Considering the small number of data reported in the literature on *in vivo* measurements of fermentation parameters using horses fitted with two cannulas in the hindgut (de Fombelle *et al.*, 2001; Julliand *et al.*, 2001; Medina *et al.*, 2002), the present work using horses fistulated in both caecum and ventral colon for studying the impact of feeding




Figure 2 Principal component analysis correlation circle of pH values and volatile fatty acid concentration profiles obtained from caecum, right ventral colon and faeces of horses ($n = 4$) given three different diets (H: 100% forage diet; HC: 40% concentrate/60% forage; for diets H and HC1, feeds were offered daily in two equal meals at 0800 and 1700 h; for the diet HC2, concentrate was offered at 0800 and 1600 h and hay at 1000 and 1730 h).

sequence on postprandial variations of microbial fermentations is an original point.

Adding concentrate in the diet was associated with an overall alteration of the pH values and VFA concentrations in caecum, right ventral colon and faeces compared with the hay diet. In agreement with the literature, caecal, colonic and faecal pH values decreased (Goodson *et al.*, 1988; McLean *et al.*, 2000; Medina *et al.*, 2002), total faecal VFA concentration increased (Hussein *et al.*, 2004) as well as faecal acetate and propionate concentrations (Hussein *et al.*, 2004; Swyers *et al.*, 2008) and caecal, colonic and faecal butyrate concentrations (de Fombelle *et al.*, 2003; Hussein *et al.*, 2004). Most of these data are in line with a change towards an acidogenic environment which in large extent can be at risk and disrupt the balance of the hindgut microbial ecosystem. On the contrary, some of our results were not in compliance with studies reporting the impact of adding concentrate to a forage diet on caecal acetate and propionate concentrations (Kern *et al.*, 1973; Medina *et al.*, 2002). Compared with these previous studies, the percentage of concentrate in our ration was lower; 40% *v.* >50% of the DM intake. In addition, the concentrate tested in the present work was low in starch (32.3% DM) compared with cereals. When starch amounts >0.2% BW are fed, undigested starch can flow into the hindgut and be rapidly fermented, thus leading to an increase in VFA production and a decrease in pH and cellulolytic activity (Julliand *et al.*, 2001).

The overall analysis carried on all pH values and VFA concentrations from caecum, right ventral colon and faeces, showed no impact on the concentrate feeding sequence. In the caecum and colon as in the faeces, we found little impact of the concentrate feeding sequence on the fermentation parameters. Only caecal pH values were significantly lower

when the concentrate meal was offered 2 h before the hay meal instead of at the same time. However, pH value did not decrease intensely (6.4 *v.* 6.7 in average), suggesting a moderate impact of feeding concentrate before hay on equine gastrointestinal health. This was in accordance with Jensen *et al.* (2012), who observed that feeding hay before or after oats did not affect minimum pH in the caecum. No impact of the concentrate feeding sequence was observed on the colonic pH values and fermentation parameters in the present work. This could be related to the low starch content of the commercial pelleted concentrate we fed to our horses. In colon digesta, Philippeau *et al.* (2009) observed indeed a higher magnitude of the pH drop when a ground barley concentrate was offered 2 h before the forage meal compared with 2 h after, whereas no effect of the concentrate feeding sequence was observed when horses were fed a fibrous commercial pelleted concentrate. Measuring faecal fermentation parameters, Zeyner *et al.* (2004) reported that offering the forage meal before the crushed oat meal seemed to protect against the acidification due to the diet. Our results appeared to be in disagreement as no effect of concentrate feeding sequence was observed on pH and VFA concentrations.

In the present study, pH and VFA postprandial responses to diet changes were also compared between different hindgut segments. Taking into account all sampling times, we found no difference regarding pH values and VFA concentrations between the caecum, right ventral colon and faeces. This was not in accordance with previous data reporting that microbial fermentation parameters were different between caecum, right dorsal colon and faeces (Dougal *et al.*, 2012). The dissimilarities of the results could be due to differences in experimental procedures; whereas horses were euthanized for collecting a single digestive sample at the end point of the experimental trial, we collected five samples of digestive contents on alive horses from each intestinal compartment and from faeces.

When pH and VFA postprandial responses were studied in each digestive segment separately, the response differed according to the hindgut location. To our knowledge, this postprandial response had never been compared between the caecum, right ventral colon and faeces of the horse within the same trial. In the caecum, the pH decrease and the VFA concentrations increase observed from 4 to 8 h after the morning meal were consistent with data previously reported in the literature (Moore-Colyer *et al.*, 2000; Medina *et al.*, 2002; Zeyner *et al.*, 2004; Jensen *et al.*, 2012). Similarly to the caecum, pH values decreased and VFA concentrations increased in the colon within the range of 4 to 8 h after the morning meal. In their recent chapter, Merritt and Julliand (2013) summarized that the transit rate of digesta through the small intestine of the horse is very rapid for liquids and in average between 4 and 7 h for particulate matter. This means that 4 to 8 h after feed consumption, most digesta have already reached the caecum and ventral colon (Van Weyenberg *et al.*, 2006), and are subjected to a maximal microbial fermentation (Medina *et al.*, 2002; Philippeau

et al., 2009). In the faeces, the postprandial response of the microbial ecosystem was observed on pH values and appeared between 6 and 8 h after the morning meal. This suggested that the variation of faecal pH could be an indicator of those of the caecal and colonic pH with a 2 h delay. This is consistent with the average total tract mean retention time of particulate matter that was estimated to vary from 22.3 to 29.7 h in the digestive tract of horses fed forage-only diets (Merritt and Julliard, 2013).

The present study showed that with a 33% starch commercial pelleted, concentrate feeding sequence had minor effect on hindgut microbial fermentation parameters, and thus on equine gastrointestinal function and health. In practice, this suggests that hay and concentrate could be distributed simultaneously, therefore making easier every day horse feeding. However, more studies are needed to confirm whether similar conclusion could be drawn with other type of concentrate such as different sources or forms of cereals.

Acknowledgement

The authors are grateful to the staff of AgroSup Dijon's horse research facilities and laboratory for their technical assistance.

References

- Archer D and Proudman C 2006. Epidemiological clues to preventing colic. *The Veterinary Journal* 172, 29–39.
- Coenen M, Kienzle E, Vervuert I and Zeyner A 2011. Recent German developments in the formulation of energy and nutrient requirements in horses and the resulting feeding recommendations. *Journal of Equine Veterinary Science* 31, 219–229.
- de Fombelle A, Julliard V, Drogoul C and Jacotot E 2001. Feeding and microbial disorders in horses: 1-effects of an abrupt incorporation of two levels of barley in a hay diet on microbial profile and activities. *Journal of Equine Veterinary Science* 21, 439–445.
- de Fombelle A, Varloud M, Goachet A-G, Jacotot E, Philippeau C, Drogoul C and Julliard V 2003. Characterisation of the microbial and biochemical profile of the different segments of the digestive tract in horses fed two distinct diets. *Animal Science* 77, 293–304.
- Dougal K, Harris PA, Edwards A, Pachebat JA, Blackmore TM, Worgan HJ and Newbold CJ 2012. A comparison of the microbiome and the metabolome of different regions of the equine hindgut. *FEMS Microbiology Ecology* 82, 642–652.
- Goncalves S, Julliard V and Leblonde A 2002. Risk factors associated with colic in horses. *Veterinary Research* 33, 641–652.
- Goodson J, Tyznik WJ, Cline JH and Dehority BA 1988. Effects of an abrupt diet change from hay to concentrate on microbial numbers and physical environment in the cecum of the pony. *Applied and Environmental Microbiology* 54, 1946–1950.
- Hussein HS, Vogedes LA, Fernandez GCJ and Frankeny RL 2004. Effects of cereal grain supplementation on apparent digestibility of nutrients and concentrations of fermentation end-products in the feces and serum of horses consuming alfalfa cubes. *Journal of Animal Science* 82, 1986–1996.
- Jensen RB, Austbø D and Tauson AH 2012. Feeding forage before or after oats affects caecum pH profiles of the horse. In *Forages and grazing in horse nutrition* (ed. M Saastamoinen, M Fradinho, A Santos and N Miraglia), pp. 327–330. Wageningen Academic Publishers, Wageningen, The Netherlands.
- Jouany J-P 1982. Volatile fatty acid and alcohol determination in digestive contents, silage juices, bacterial cultures and anaerobic fermentor contents. *Sciences des Aliments* 2, 131–144.
- Julliard V, de Fombelle A, Drogoul C and Jacotot E 2001. Feeding and microbial disorders in horses: part 3 – effects of three hay:grain ratios on microbial profile and activities. *Journal of Equine Veterinary Science* 21, 543–546.
- Kern DL, Slyter LL, Weaver JM, Leffel EC and Samuelsons G 1973. Pony cecum vs. steer rumen: the effect of oats and hay on the microbial ecosystem. *Journal of Animal Science* 37, 463–469.
- McLean BML, Hyslop JJ, Longland AC, Cuddeford D and Hollands T 2000. Physical processing of barley and its effects on intra-caecal fermentation parameters in ponies. *Animal Feed Science and Technology* 85, 79–87.
- Medina B, Girard ID, Jacotot E and Julliard V 2002. Effect of a preparation of *Saccharomyces cerevisiae* on microbial profiles and fermentation patterns in the large intestine of horses fed a high fiber or a high starch diet. *Journal of Animal Science* 80, 2600–2609.
- Merritt AM and Julliard V 2013. Gastrointestinal physiology. In *Equine applied and clinical nutrition: health, welfare and performance* (ed. RJ Geor, PA Harris and M Coenen), pp. 3–32. W.B. Saunders.
- Miyaji M, Ueda K, Nakatsuji H, Tomioka T, Kobayashi Y, Hata H and Kondo S 2008. Mean retention time of digesta in the different segments of the equine hindgut. *Animal Science Journal* 79, 89–96.
- Moore-Colyer MJS, Hyslop JJ, Longland AC and Cuddeford D 2000. Intra-caecal fermentation parameters in ponies fed botanically diverse fibre based diets. *Animal Feed Science and Technology* 84, 183–197.
- Philippeau C, Faubladiere C, Goachet AG and Julliard V 2009. Is there an impact of feeding concentrate before or after forage on colonic pH and redox potential in horses? In *Equine Nutrition and TRAINING Conference (ENUTRACO)*, 26–29 September 2009, Madrid, Spain.
- Santos AS, Rodrigues MAM, Bessa RJB, Ferreira LM and Martin-Rosset W 2011. Understanding the equine cecum-colon ecosystem: current knowledge and future perspectives. *Animal* 5, 48–56.
- Swyers KL, Burk AO, Hartssock TG, Ungerfeld EM and Shelton JL 2008. Effects of direct-fed microbial supplementation on digestibility and fermentation end-products in horses fed low- and high-starch concentrates. *Journal of Animal Science* 86, 2596–2608.
- Thioulouse J, Chessel D, Dolédec S and Olivier JM 1997. ADE-4: a multivariate analysis and graphical display software. *Statistics and Computing* 7, 75–83.
- Van Weyenberg S, Sales J and Janssens GPJ 2006. Passage rate of digesta through the equine gastrointestinal tract: a review. *Livestock Science* 99, 3–12.
- Zeyner A, Geißler C and Dittrich A 2004. Effects of hay intake and feeding sequence on variables in faeces and faecal water (dry matter, pH value, organic acids, ammonia, buffering capacity) of horses. *Journal of Animal Physiology and Animal Nutrition* 88, 7–19.