



HAL
open science

Embeddings of finite groups in $B_n/\Gamma_k(P_n)$ for $k = 2, 3$

Daciberg Lima Gonçalves, John Guaschi, Oscar Ocampo

► **To cite this version:**

Daciberg Lima Gonçalves, John Guaschi, Oscar Ocampo. Embeddings of finite groups in $B_n/\Gamma_k(P_n)$ for $k = 2, 3$. 2018. hal-01800722v1

HAL Id: hal-01800722

<https://hal.science/hal-01800722v1>

Preprint submitted on 27 May 2018 (v1), last revised 28 Oct 2018 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Embeddings of finite groups in $B_n/\Gamma_k(P_n)$ for $k = 2, 3$

DACIBERG LIMA GONÇALVES

Departamento de Matemática - IME-USP,
Rua do Matão 1010 CEP: 05508-090 - São Paulo - SP - Brazil.
e-mail: dlgoncal@ime.usp.br

JOHN GUASCHI

Normandie Univ., UNICAEN, CNRS,
Laboratoire de Mathématiques Nicolas Oresme UMR CNRS 6139,
CS 14032, 14032 Cedex Cedex 5, France.
e-mail: john.guaschi@unicaen.fr

OSCAR OCAMPO

Universidade Federal da Bahia, Departamento de Matemática - IME,
Av. Adhemar de Barros S/N CEP: 40170-110 - Salvador - BA - Brazil.
e-mail: oscaro@ufba.br

27th May 2018

Abstract

Let $n \geq 3$. In this paper, we study the problem of whether a given finite group G embeds in a quotient of the form $B_n/\Gamma_k(P_n)$, where B_n is the n -string Artin braid group, $k \in \{2, 3\}$, and $\{\Gamma_l(P_n)\}_{l \in \mathbb{N}}$ is the lower central series of the n -string pure braid group P_n . Previous results show that a necessary condition for such an embedding to exist is that $|G|$ is odd (resp. is relatively prime with 6) if $k = 2$ (resp. $k = 3$). We show that any finite group G of odd order (resp. of order relatively prime with 6) embeds in $B_{|G|}/\Gamma_2(P_{|G|})$ (resp. in $B_{|G|}/\Gamma_3(P_{|G|})$), where $|G|$ denotes the order of G . The result in the case of $B_{|G|}/\Gamma_2(P_{|G|})$ has been proved independently by Beck and Marin. One may then ask whether G embeds in a quotient of the form $B_n/\Gamma_k(P_n)$, where $n < |G|$ and $k \in \{2, 3\}$. If G is of the form $\mathbb{Z}_{p^r} \rtimes_{\theta} \mathbb{Z}_d$, where p is an odd prime (resp. $p \geq 5$ is prime) and d is odd (resp. d is relatively prime with 6) and divides $p - 1$, we show that G embeds in $B_{p^r}/\Gamma_2(P_{p^r})$ (resp. in $B_{p^r}/\Gamma_3(P_{p^r})$). This generalises a result of Marin concerning the embedding of the Frobenius groups in $B_n/\Gamma_2(P_n)$. Finally, we prove that the two non-Abelian groups of order 27, namely the semi-direct product $\mathbb{Z}_9 \rtimes \mathbb{Z}_3$, where the action is given by multiplication by 4, and the Heisenberg group mod 3, both embed in $B_9/\Gamma_2(P_9)$.

1 Introduction

If $n \in \mathbb{N}$, let B_n denote the (*Artin*) *braid group* on n strings. It is well known that B_n admits a presentation with generators $\sigma_1, \dots, \sigma_{n-1}$ that are subject to the relations $\sigma_i \sigma_j = \sigma_j \sigma_i$ for all $1 \leq i < j \leq n-1$ for which $|i-j| \geq 2$, and $\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}$ for all $1 \leq i \leq n-2$. Let $\sigma: B_n \rightarrow S_n$ denote the surjective homomorphism onto the symmetric group S_n defined by $\sigma(\sigma_i) = (i, i+1)$ for all $1 \leq i \leq n-1$. The *pure braid group* P_n on n strings is defined to be the kernel of σ , from which we obtain the following short exact sequence:

$$1 \longrightarrow P_n \longrightarrow B_n \xrightarrow{\sigma} S_n \longrightarrow 1. \quad (1)$$

If G is a group, recall that its *lower central series* $\{\Gamma_k(G)\}_{k \in \mathbb{N}}$ is defined by $\Gamma_1(G) = G$, and $\Gamma_k(G) = [\Gamma_{k-1}(G), G]$ for all $k \geq 2$ (if H and K are subgroups of G , $[H, K]$ is defined to be the subgroup of G generated by the commutators of the form $[h, k] = hkh^{-1}k^{-1}$, where $h \in H$ and $k \in K$). Note that $\Gamma_2(G)$ is the commutator subgroup of G , and that $\Gamma_k(G)$ is a normal subgroup of G for all $k \in \mathbb{N}$. In our setting, since P_n is normal in B_n , it follows that $\Gamma_k(P_n)$ is also normal in B_n , and the extension (1) induces the following short exact sequence:

$$1 \longrightarrow P_n/\Gamma_k(P_n) \longrightarrow B_n/\Gamma_k(P_n) \xrightarrow{\bar{\sigma}} S_n \longrightarrow 1, \quad (2)$$

obtained by taking the quotient of P_n and B_n by $\Gamma_k(P_n)$. It follows from results of Falk and Randell [FR] and Kohno [K] that the kernel of (2) is torsion free (see Proposition 5 for more information). The quotient groups of the form $B_n/\Gamma_k(P_n)$ have been the focus of several recent papers. First, the quotient $B_n/\Gamma_2(P_n)$ belongs to a family of groups known as *enhanced symmetric groups* [M1, page 201] that were analysed in [T]. Secondly, in their study of pseudo-symmetric braided categories, Panaite and Staic showed that this quotient is isomorphic to the quotient of B_n by the normal closure of the set $\left\{ \sigma_i \sigma_{i+1}^{-1} \sigma_i \sigma_{i+1}^{-1} \sigma_i \sigma_{i+1}^{-1} \mid i = 1, 2, \dots, n-2 \right\}$ [PS]. Thirdly, in [GGO1], we showed that $B_n/\Gamma_2(P_n)$ is a crystallographic group, and that up to isomorphism, its finite Abelian subgroups are the Abelian subgroups of S_n of odd order. In particular, the torsion of $B_n/\Gamma_2(P_n)$ is the odd torsion of S_n . We also gave an explicit embedding in $B_7/\Gamma_2(P_7)$ of the Frobenius group $\mathbb{Z}_7 \rtimes \mathbb{Z}_3$ of order 21, which is the smallest finite non-Abelian group of odd order. As far as we know, this is the first example of a finite non-Abelian group that embeds in a quotient of the form $B_n/\Gamma_2(P_n)$. Almost all of the results of [GGO1] were subsequently extended to the generalised braid groups associated to an arbitrary complex reflection group by Marin [M2]. If $p > 3$ is a prime number for which $p \equiv 3 \pmod{4}$, he showed that the Frobenius group $\mathbb{Z}_p \rtimes \mathbb{Z}_{(p-1)/2}$ embeds in $B_p/\Gamma_2(P_p)$. Observe that this group cannot be embedded in $B_n/\Gamma_2(P_n)$ for any $n < p$, since \mathbb{Z}_p cannot be embedded in S_n in this case. In another direction, the authors studied some aspects of the quotient $B_n/\Gamma_k(P_n)$ for all $n, k \geq 3$, and proved that it is an almost-crystallographic group [GGO2]. For the case $k = 3$, it was shown that the torsion of $B_n/\Gamma_3(P_n)$ is the torsion of S_n that is relatively prime with 6. For future reference, we summarise some of these results in the following theorem.

THEOREM 1 ([GGO1, Corollary 4], [GGO2, Theorems 2 and 3]). *Let $n \geq 3$.*

(a) *The torsion of the quotient $B_n/\Gamma_2(P_n)$ is equal to the odd torsion of S_n .*

(b) The group $B_n/\Gamma_3(P_n)$ has no elements of order 2 or 3, and if $m \in \mathbb{N}$ is relatively prime with 6 then $B_n/\Gamma_3(P_n)$ possesses elements of order m if and only if S_n does.

Almost nothing is known about the torsion and the finite subgroups of $B_n/\Gamma_k(P_n)$ in the case where $k > 3$.

Suppose that $n \geq 3$ and $k \geq 2$. The results of [GGO1, GGO2, M2] lead to a number of interesting problems involving the quotients $B_n/\Gamma_k(P_n)$. Given a finite group G , a natural question in our setting is whether it can be embedded in some $B_n/\Gamma_k(P_n)$. In order to formulate some of these problems, we introduce the following notation. Let $|G|$ denote the order of G , let $m(G)$ denote the least positive integer r for which G embeds in the symmetric group S_r , and if $k \geq 2$, let $\ell_k(G)$ denote the least positive integer s , if such an integer exists, for which G embeds in the group $B_s/\Gamma_k(P_s)$. The integer $\ell_k(G)$ is not always defined. For example, if G is of even order, Theorem 1(a) implies that G does not embed in any group of the form $B_n/\Gamma_k(P_n)$. However, if $\ell_k(G)$ is defined, then $m(G) \leq \ell_k(G)$ using (2) and the fact that $P_n/\Gamma_k(P_n)$ is torsion free.

The main aim of this paper is to study the embedding of finite groups in the two quotients $B_n/\Gamma_k(P_n)$, where $k \in \{2, 3\}$. In Section 2, we start by recalling some results from [GGO1, GGO2] about the action of S_n on certain bases of the free Abelian groups $P_n/\Gamma_2(P_n)$ and $\Gamma_2(P_n)/\Gamma_3(P_n)$, which we use to obtain information about the cycle structure of elements of S_n that fix elements of these bases. In Proposition 7, using cohomological arguments, we show that a short exact sequence splits if its quotient is a finite group G and its kernel is a free $\mathbb{Z}[G]$ -module. This provides a fundamental tool for embedding G in our quotients. In Section 3, we prove the following result.

THEOREM 2. *Let G be a finite group, and let $k \in \{2, 3\}$. Then the group G embeds in $B_{|G|}/\Gamma_k(P_{|G|})$ if and only if $\gcd(|G|, k!) = 1$.*

The result of Theorem 2 has recently been proved independently by Beck and Marin in the case $k = 2$ [BM]. Theorem 2 may be viewed as a Cayley-type result for $B_n/\Gamma_k(P_n)$ since the proof makes use of the embedding of G in the symmetric group $S_{|G|}$, as well as Proposition 8 that provides sufficient conditions on the fixed points in the image of an embedding of G in S_m for G to embed in $B_n/\Gamma_k(P_n)$. If $\gcd(|G|, k!) = 1$, it follows from this theorem that $\ell_k(G) \leq |G|$ by Theorem 2, from which we obtain:

$$m(G) \leq \ell_k(G) \leq |G|. \quad (3)$$

The analysis of the inequalities of (3) is itself an interesting problem. Using Theorem 1, if G is a cyclic group of prime order at least 5 and k is equal to either 2 or 3 then $m(G) = \ell_k(G) = |G|$. We do not currently know whether there exist groups for which $m(G) < \ell_k(G)$.

In Section 4, we study the embedding of certain finite groups in $B_n/\Gamma_k(P_n)$, where $k \in \{2, 3\}$. In Section 4.1, we consider certain semi-direct products of the form $\mathbb{Z}_n \rtimes_{\theta} \mathbb{Z}_m$ for which the action θ is injective, and we analyse their possible embedding in $B_n/\Gamma_k(P_n)$. Our main result in this direction is the following.

THEOREM 3. *Let $m, n \geq 3$, let $G = \mathbb{Z}_n \rtimes_{\theta} \mathbb{Z}_m$, where $\theta: \mathbb{Z}_m \rightarrow \mathbb{Z}_n$ is the associated action, and let $1 \leq t < n$ be such that $\theta(1_m)$ is multiplication by t in \mathbb{Z}_n . Assume that $\gcd(t^l - 1, n) = 1$ for all $1 \leq l \leq m - 1$. If mn is odd (resp. $\gcd(mn, 6) = 1$) then G embeds in $B_n/\Gamma_2(P_n)$ (resp. in $B_n/\Gamma_3(P_n)$).*

Using Lemma 11(a), we remark that the hypotheses of Theorem 3 imply that the action $\theta: \mathbb{Z}_m \rightarrow \text{Aut}(\mathbb{Z}_n)$ is injective. As an application of this theorem, we obtain the following corollary.

COROLLARY 4. *Let p be an odd prime, let $p - 1 = 2^j d$, where d is odd, let d_1 be a divisor of d , and let G be a group of the form $\mathbb{Z}_{p^r} \rtimes_{\theta} \mathbb{Z}_{d_1}$, where $\theta: \mathbb{Z}_{d_1} \rightarrow \text{Aut}(\mathbb{Z}_{p^r})$ is injective.*

(a) *If $p \geq 3$ then G embeds in $B_{p^r}/\Gamma_2(P_{p^r})$.*

(b) *If $p \geq 5$ and d_1 satisfies $\gcd(d_1, 3) = 1$ then G embeds in $B_{p^r}/\Gamma_3(P_{p^r})$.*

Since the group \mathbb{Z}_{p^r} cannot be embedded in S_m for any $m < p^r$, the groups of Corollary 4 satisfy $m(G) = \ell_k(G) = p^r$, where $k \in \{2, 3\}$, so the results of this corollary are sharp in this sense. Further, the groups that appear in [M2, Corollary 3.11] correspond to the case of where $r = 1$, $p \equiv 3 \pmod{4}$, and $d_1 = (p - 1)/2$ is odd. Hence Corollary 4 generalises Marin’s result to the case where p is any odd prime and d_1 is the greatest odd divisor of $p - 1$, and more generally, in the case $k = 2$, the family of groups obtained in Theorem 3 extends even further that of the Frobenius groups of [M2, Corollary 3.11].

At the end of the paper, in Section 4.2, we show that the two non-Abelian groups of order 27 embed in $B_9/\Gamma_2(P_9)$. Neither of these groups satisfies the hypotheses of Theorem 3, and their embedding is not a consequence of any known results. Within our framework, it is natural to study these two groups, first because with the exception of the Frobenius group of order 21 analysed in [GG01], they are the smallest non-Abelian groups of odd order, and secondly because they are of order 27, so are related to the discussion in Section 4.1 on groups whose order is a prime power. The direct embedding of these groups in $B_9/\Gamma_2(P_9)$ is computationally difficult due to the fact that the kernel $P_9/\Gamma_2(P_9)$ of (2) is of rank 36, but we get round this problem by first considering an embedding in a quotient where the corresponding kernel is free of rank 9, and then by applying Proposition 7. We believe that this technique will prove to be useful for other groups.

Very little is known about the classification of all finite subgroups of $B_n/\Gamma_2(P_n)$ for fixed n . The study of other finite subgroups of $B_n/\Gamma_k(P_n)$ for $k \in \{2, 3\}$ constitutes work in progress.

Acknowledgements

This work took place during several visits to the Departamento de Matemática, Universidade Federal de Bahia, the Departamento de Matemática do IME – Universidade de São Paulo and to the Laboratoire de Mathématiques Nicolas Oresme UMR CNRS 6139, Université de Caen Normandie. The first and the third authors were partially supported by the FAPESP Projeto Temático “Topologia Algébrica, Geométrica e Diferencial” 2016/24707-4 (Brazil), and the first two authors were partially supported by the CNRS/FAPESP PRC project n° 275209 (France) and n° 2016/50354-1 (Brazil). The third author was initially supported by a project grant n° 151161/2013-5 from CNPq. We would also like to thank Vincent Beck and Ivan Marin for sharing their results with us.

2 Preliminaries

In this section, we recall several results concerning the torsion of the groups $B_n/\Gamma_k(P_n)$, where $k \in \{2, 3\}$, as well as some group-cohomological facts from [Br] that will be used in this paper. We first state the following result from [GGO2] that we will require.

PROPOSITION 5. [GGO2, Lemma 11]

- (a) Let $n, k \geq 2$. Then the group $P_n/\Gamma_k(P_n)$ is torsion free.
 (b) Let $n \geq 3$, let $k \geq l \geq 1$, and let G be a finite group. If $B_n/\Gamma_k(P_n)$ possesses a (normal) subgroup isomorphic to G then $B_n/\Gamma_l(P_n)$ possesses a (normal) subgroup isomorphic to G . In particular, if p is prime, and if $B_n/\Gamma_l(P_n)$ has no p -torsion then $B_n/\Gamma_k(P_n)$ has no p -torsion.

Note that the first part of Proposition 5 follows from papers by Falk and Randell [FR, Theorem 4.2] and Kohno [K, Theorem 4.5] who proved independently that for all $n \geq 2$ and $k \geq 1$, the group $\Gamma_k(P_n)/\Gamma_{k+1}(P_n)$ is free Abelian of finite rank, the rank being related to the Poincaré polynomial of certain hyperplane complements.

It is well known that a set of generators for P_n is given by the set $\{A_{i,j}\}_{1 \leq i < j \leq n}$ [H]. If $j > i$ then we take $A_{j,i} = A_{i,j}$. By abuse of notation, for $k \geq 2$ and $1 \leq i < j \leq n$, we also denote the image of $A_{i,j}$ under the canonical projection $P_n \rightarrow P_n/\Gamma_k(P_n)$ by $A_{i,j}$. The groups $P_n/\Gamma_2(P_n)$ and $\Gamma_2(P_n)/\Gamma_3(P_n)$ are free Abelian groups of finite rank $n(n-1)/2$ and $n(n-1)(n-2)/6$ respectively [FR, Theorem 4.2]. By [GGO1, Section 3, p. 399] (resp. [GGO2, equation (17)]), a basis for $P_n/\Gamma_2(P_n)$ (resp. $\Gamma_2(P_n)/\Gamma_3(P_n)$) is given by:

$$\mathcal{B} = \{A_{i,j} \mid 1 \leq i < j \leq n\} \text{ (resp. by } \mathcal{B}' = \{\alpha_{i,j,k} \mid 1 \leq i < j < k \leq n\}), \quad (4)$$

where $\alpha_{i,j,k} = [A_{i,j}, A_{j,k}]$. If $\tau \in S_n$, $A_{i,j} \in \mathcal{B}$ and $\alpha_{i,j,k} \in \mathcal{B}'$ then by [GGO1, Proposition 12] and [GGO2, equation (8)], we have:

$$\tau \cdot A_{i,j} = A_{\tau^{-1}(i), \tau^{-1}(j)} \text{ and } \tau \cdot \alpha_{i,j,k} = \tau \cdot [A_{i,j}, A_{j,k}] = [A_{\tau^{-1}(i), \tau^{-1}(j)}, A_{\tau^{-1}(j), \tau^{-1}(k)}]. \quad (5)$$

The following lemma implies that S_n acts on \mathcal{B} and $\widehat{\mathcal{B}}'$ respectively, where $\widehat{\mathcal{B}}' = \mathcal{B}' \cup \mathcal{B}'^{-1}$. In each case, the nature of the action gives rise by linearity to an action of S_n on the whole group. We also obtain some information about the stabilisers of the elements of \mathcal{B} and $\widehat{\mathcal{B}}'$. This will play a crucial rôle in the proof of Proposition 8.

LEMMA 6. Let $n \geq 2$, and let $\tau \in S_n$.

- (a) Let $A_{i,j}$ be an element of the basis \mathcal{B} of $P_n/\Gamma_2(P_n)$, where $1 \leq i < j \leq n$. Then the element $\tau \cdot A_{i,j}$ given by the action of S_n on $P_n/\Gamma_2(P_n)$ belongs to \mathcal{B} . Further, if $\tau \cdot A_{i,j} = A_{i,j}$ then the cycle decomposition of τ either contains a transposition, or at least two fixed elements.
 (b) Let $\alpha_{i,j,k}$ be an element of the basis \mathcal{B}' of $\Gamma_2(P_n)/\Gamma_3(P_n)$, where $1 \leq i < j < k \leq n$. Then the element $\tau \cdot \alpha_{i,j,k}$ given by the action of S_n on $\Gamma_2(P_n)/\Gamma_3(P_n)$ belongs to $\widehat{\mathcal{B}}'$. Further, if $\tau \cdot \alpha_{i,j,k} \in \{\alpha_{i,j,k}, \alpha_{i,j,k}^{-1}\}$ then the cycle decomposition of τ contains either a transposition, or a 3-cycle, or at least three fixed elements.

Proof.

- (a) The first part follows from (5). If $1 \leq i < j \leq n$ and $\tau \in S_n$ are such that $\tau \cdot A_{i,j} = A_{i,j}$ then $\tau(\{i, j\}) = \{\tau(i), \tau(j)\} = \{i, j\}$, which implies the second part of the statement.

(b) The first part is a consequence of [GGO2, equation (16)]. Now suppose that $\tau \cdot \alpha_{i,j,k} \in \{\alpha_{i,j,k}, \alpha_{i,j,k}^{-1}\}$, where $1 \leq i < j < k \leq n$ and $\tau \in S_n$. By [GGO2, equation (18)], we have $\tau(\{i, j, k\}) = \{\tau(i), \tau(j), \tau(k)\} = \{i, j, k\}$, from which we deduce the second part. \square

Lemma 6 implies that if $G = S_n$ then \mathcal{B} and $\widehat{\mathcal{B}}'$ are G -sets, and the action of G on each of these sets extends to a \mathbb{Z} -linear action of G on the free \mathbb{Z} -modules $\mathbb{Z}\mathcal{B}$ and $\mathbb{Z}\mathcal{B}'$ respectively, with respect to the embedding of $\widehat{\mathcal{B}}'$ in $\mathbb{Z}\mathcal{B}'$ given by $\alpha_{i,j,k} \mapsto \alpha_{i,j,k}$ and $\alpha_{i,j,k}^{-1} \mapsto (-1) \cdot \alpha_{i,j,k}$, the underlying free Abelian groups being naturally identified with $P_n/\Gamma_2(P_n)$ and $\Gamma_2(P_n)/\Gamma_3(P_n)$ respectively. We conclude that $P_n/\Gamma_2(P_n)$ and $\Gamma_2(P_n)/\Gamma_3(P_n)$ each admit a G -module structure inherited by the action of S_n on \mathcal{B} and $\widehat{\mathcal{B}}'$ respectively.

Given a group G , let $\mathbb{Z}[G]$ denote its group ring. The underlying Abelian group, also denoted by $\mathbb{Z}[G]$, may be regarded as a $\mathbb{Z}[G]$ -module (or as a G -module) via the multiplication in the ring $\mathbb{Z}[G]$ (see [Br, Chapter I, Sections 2 and 3] for more details), namely:

$$g \cdot \sum_{i=1}^m n_i g_i = \sum_{i=1}^m n_i (g g_i) \text{ for all } m \in \mathbb{N}, g, g_1, \dots, g_m \in G \text{ and } n_1, \dots, n_m \in \mathbb{Z}. \quad (6)$$

The cohomology of the group G with coefficients in $\mathbb{Z}[G]$ regarded as a G -module is well understood. In the case that G is finite, we have the following result concerning its embedding in certain extensions whose kernel is a free $\mathbb{Z}[G]$ -module. First recall that if M is an Abelian group that fits into an extension of the following form:

$$1 \longrightarrow M \longrightarrow E \longrightarrow G \longrightarrow 1, \quad (7)$$

then M is also a $\mathbb{Z}[G]$ -module, the action being given by (7).

PROPOSITION 7. *Let G be a finite group. Given an extension of the form (7), suppose that M is a free $\mathbb{Z}[G]$ -module. Then the short exact sequence (7) splits. In particular, G embeds in E as a subgroup, and the restriction of the projection $E \longrightarrow G$ to the embedded copy of G is an isomorphism.*

Proof. First suppose that $M \cong \mathbb{Z}[G]$. By [Br, equation 6.5, p. 73], the group $H^*(G, \mathbb{Z}[G])$ is trivial for all $* \geq 1$. In particular, $H^2(G, \mathbb{Z}[G]) = 0$, which implies that any extension of the form (7) with $M = \mathbb{Z}[G]$ is split [Br, Chapter IV, Theorem 3.12], where the action of the quotient on the kernel turns $\mathbb{Z}[G]$ into a $\mathbb{Z}[G]$ -module that is isomorphic to the one-dimensional free $\mathbb{Z}[G]$ -module. Now suppose that M is an arbitrary free $\mathbb{Z}[G]$ -module whose $\mathbb{Z}[G]$ -module structure is defined by (7). So $M \cong \bigoplus_J \mathbb{Z}[G]$ as a $\mathbb{Z}[G]$ -module for some set J , and $H^2(G, M) \cong H^2(G, \bigoplus_J \mathbb{Z}[G]) \cong \bigoplus_J H^2(G, \mathbb{Z}[G]) = 0$ by the first part of the proof. The short exact sequence (7) splits as in the case $M \cong \mathbb{Z}[G]$. \square

3 Cayley-type results for subgroups of $B_n/\Gamma_k(P_n)$, $k = 2, 3$

Let $k \in \{2, 3\}$. In this section, we prove Theorem 2 that may be viewed as an analogue of Cayley's theorem for $B_n/\Gamma_k(P_n)$. The following proposition will be crucial in the proofs of Theorems 2 and 3.

PROPOSITION 8. Let $k \in \{2, 3\}$, let G be a finite group whose order is relatively prime with $k!$, let $m \geq 3$, and let $\varphi: G \rightarrow S_m$ be an embedding. Assume that for all $g \in G \setminus \{e\}$, the cycle decomposition of $\varphi(g)$ contains at most $k - 1$ fixed elements. Then the group G embeds in $B_m/\Gamma_k(P_m)$.

Proof. Assume first that $k = 2$, so $|G|$ is odd. Let \tilde{G} be the (isomorphic) image of G by φ in S_m . Taking the inverse image by $\bar{\sigma}$ of \tilde{G} in (2) with $n = m$ and $k = 2$ gives rise to the following short exact sequence:

$$1 \longrightarrow P_m/\Gamma_2(P_m) \longrightarrow \bar{\sigma}^{-1}(\tilde{G}) \xrightarrow{\bar{\sigma}|_{\bar{\sigma}^{-1}(\tilde{G})}} \tilde{G} \longrightarrow 1. \quad (8)$$

From Section 2, G , and hence \tilde{G} , acts on the free Abelian group $P_m/\Gamma_2(P_m)$ of rank $m(m-1)/2$, and the restriction of this action to the basis \mathcal{B} is given by (5). Let $1 \leq i < j \leq m$, and let $g \in G$ be such that $\varphi(g) \cdot A_{i,j} = A_{i,j}$. Since $|G|$ is odd, the cycle decomposition of $\varphi(g)$ contains no transposition, and by Lemma 6(a) and the hypothesis on the fixed points of $\varphi(g)$, we see that $g = e$. So for all $1 \leq i < j \leq m$, the orbit of $A_{i,j}$ contains exactly $|G|$ elements. In particular, $|G|$ divides $m(m-1)/2$, and \mathcal{B} may be decomposed as the disjoint union of the form $\coprod_{k=1}^{m(m-1)/2|G|} \mathcal{O}_k$, where each \mathcal{O}_k is an orbit of length $|G|$. For $k = 1, \dots, m(m-1)/2|G|$, let $e_k \in \mathcal{O}_k$, and let H_k denote the subgroup of $P_m/\Gamma_2(P_m)$ generated by \mathcal{O}_k . Then $P_m/\Gamma_2(P_m) \cong \bigoplus_{k=1}^{m(m-1)/2|G|} H_k$, and for all $x \in \mathcal{O}_k$, there exists a unique element $g \in G$ such that $\varphi(g) \cdot e_k = x$. Thus the map that to x associates g defines a bijection between \mathcal{O}_k and G . Since \mathcal{O}_k is a basis of H_k , if $h \in H_k$, there exists a unique family of integers $\{q_g\}_{g \in G}$ such that $h = \prod_{g \in G} (\varphi(g) \cdot e_k)^{q_g}$, and the map $\Phi: H_k \rightarrow \mathbb{Z}[G]$ defined by $\Phi(h) = \sum_{g \in G} q_g g$ may be seen to be an isomorphism. Further, via Φ , the action of G on H_k corresponds to the usual action of G on $\mathbb{Z}[G]$. More precisely, if $\gamma \in G$, then:

$$\begin{aligned} \Phi(\varphi(\gamma) \cdot h) &= \Phi\left(\varphi(\gamma) \cdot \prod_{g \in G} (\varphi(g) \cdot e_k)^{q_g}\right) = \Phi\left(\prod_{g \in G} (\varphi(\gamma g) \cdot e_k)^{q_g}\right) \\ &= \sum_{g \in G} q_g \gamma g = \gamma \cdot \Phi(h), \end{aligned}$$

the action of γ on $\Phi(h)$ being given by (6). Hence $P_m/\Gamma_2(P_m) \cong \bigoplus_1^{m(m-1)/2|G|} \mathbb{Z}[G]$ as $\mathbb{Z}[G]$ -modules, and by Proposition 7, we conclude that the extension (8) splits. Thus G is isomorphic to a subgroup \widehat{G} of $\bar{\sigma}^{-1}(\tilde{G})$, which in turn is a subgroup of $B_m/\Gamma_2(P_m)$, and this proves the result in the case $k = 2$. Now suppose that $k = 3$. Since $\gcd(|G|, 6) = 1$, $|G|$ is odd, and as above, G is isomorphic to the subgroup \widehat{G} of $B_m/\Gamma_2(P_m)$. Consider the following extension:

$$1 \longrightarrow \Gamma_2(P_m)/\Gamma_3(P_m) \longrightarrow B_m/\Gamma_3(P_m) \xrightarrow{\rho} B_m/\Gamma_2(P_m) \longrightarrow 1,$$

where $\rho: B_m/\Gamma_3(P_m) \rightarrow B_m/\Gamma_2(P_m)$ denotes the canonical projection. Taking the inverse image of \widehat{G} by ρ gives rise to the following short exact sequence:

$$1 \longrightarrow \Gamma_2(P_m)/\Gamma_3(P_m) \longrightarrow \rho^{-1}(\widehat{G}) \xrightarrow{\rho|_{\rho^{-1}(\widehat{G})}} \widehat{G} \longrightarrow 1. \quad (9)$$

Let $\varphi': \widehat{G} \rightarrow S_m$ denote the embedding of \widehat{G} in S_m given by composing φ by an isomorphism between \widehat{G} and G . Then \widehat{G} acts on the kernel $\Gamma_2(P_m)/\Gamma_3(P_m)$ of (9) via (5). Since $\gcd(|\widehat{G}|, 6) = 1$, for all $\widehat{g} \in \widehat{G} \setminus \{e\}$, the cycle decomposition of $\varphi'(\widehat{g})$ contains neither a transposition nor a 3-cycle, and by hypothesis, $\varphi'(\widehat{g})$ contains at most 2 fixed elements. It follows from Lemma 6(b) that if $\varphi'(\widehat{g}) \cdot \alpha_{i,j,k} \in \{\alpha_{i,j,k}, \alpha_{i,j,k}^{-1}\}$, where $1 \leq i < j < k \leq m$ and $\widehat{g} \in \widehat{G}$, then $\widehat{g} = e$. In particular, the orbits of $\alpha_{i,j,k}$ and $\alpha_{i,j,k}^{-1}$ are disjoint, every orbit contains exactly $|G|$ elements, and thus $|G|$ divides $m(m-1)(m-2)/6$. So there exists a basis of $\Gamma_2(P_m)/\Gamma_3(P_m)$ that is the disjoint union of $m(m-1)(m-2)/6|G|$ orbits of elements of \widehat{B}' , and that for all $1 \leq i < j < k \leq m$, contains exactly one element of $\{\alpha_{i,j,k}, \alpha_{i,j,k}^{-1}\}$. As in the case $k = 2$, we conclude that the short exact sequence (9) splits, and that G embeds in $B_m/\Gamma_3(P_m)$. \square

REMARK 9. The most efficient way to use Proposition 8 is as follows. Let $\varphi: G \rightarrow S_n$ be an embedding, and for an order-preserving inclusion $\iota: \{1, 2, \dots, m\} \hookrightarrow \{1, 2, \dots, n\}$, where $m < n$, consider the embedding $S_m \rightarrow S_n$. Suppose that the homomorphism φ factors through S_m , and let $\varphi': G \rightarrow S_m$ be the factorisation. It may happen that the hypotheses of Proposition 8 hold for φ' but not for φ . In this case, we may apply Proposition 8 to φ' to conclude the existence of an embedding of G in $B_m/\Gamma_k(P_m)$, which in turn implies that G embeds in $B_n/\Gamma_k(P_n)$.

We are now able to prove Theorem 2.

Proof of Theorem 2. Let G be a finite group, and let $k \in \{2, 3\}$. If G embeds in $B_{|G|}/\Gamma_k(P_{|G|})$ then Theorem 1 implies that $\gcd(|G|, k!) = 1$. Conversely, suppose that $\gcd(|G|, k!) = 1$. Consider the classical embedding of G in $S_{|G|}$ that is used in the proof of Cayley's theorem (note that we identify S_G with $S_{|G|}$). More precisely, let $\psi: G \times G \rightarrow G$ denote the action of G on itself given by left multiplication. For all $g \in G$, the map $\psi_g: G \rightarrow G$ defined by $\psi_g(h) = \psi(g, h) = gh$ is a permutation of G , and the map $\Psi: G \rightarrow S_{|G|}$ defined by $\Psi(g) = \psi_g$ is an injective homomorphism, so $\widetilde{G} = \{\psi_g \mid g \in G\}$ is a subgroup of $S_{|G|}$ that is isomorphic to G . The action ψ is free: if $h \in G$ then $h = \psi_g(h)$ if and only if $g = e$. In particular, if $g \neq e$ then ψ_g is fixed-point free, and so the permutation $\Psi(g)$ is fixed-point free for all $g \in G \setminus \{e\}$. Taking $m = |G|$, the hypotheses of Proposition 8 are satisfied for the embedding $\Psi: G \rightarrow S_{|G|}$, and we conclude that G embeds in $B_{|G|}/\Gamma_k(P_{|G|})$. \square

4 Embeddings of some semi-direct products in $B_n/\Gamma_k(P_n)$, $k \in \{2, 3\}$

Let $m, n \in \mathbb{N}$, and let $k \in \{2, 3\}$. In this section, we study the problem of embedding groups of the form $\mathbb{Z}_n \rtimes_{\theta} \mathbb{Z}_m$ in $B_n/\Gamma_k(P_n)$, where the representation $\theta: \mathbb{Z}_m \rightarrow \text{Aut}(\mathbb{Z}_n)$ is taken to be injective. With additional conditions on θ , in Section 4.1, we prove Theorem 3. In Section 4.2, we study the two non-Abelian groups of order 27. The first such group is of the form $\mathbb{Z}_n \rtimes_{\theta} \mathbb{Z}_m$, where θ is injective, but the additional conditions of Theorem 3 are not satisfied. The second such group is not of the form $\mathbb{Z}_n \rtimes_{\theta} \mathbb{Z}_m$. We prove that both of these groups embed in $B_9/\Gamma_2(P_9)$. In the case of the first group, this

shows that the hypotheses of Theorem 3 are sufficient to embed $\mathbb{Z}_n \rtimes_{\theta} \mathbb{Z}_m$ in $B_n/\Gamma_k(P_n)$, but not necessary. With respect to (3), these groups also satisfy $m(G) = \ell_k(G) < |G|$.

4.1 Proof of Theorem 3

Let $m, n \in \mathbb{N}$. In this section, G will be a group of the form $\mathbb{Z}_n \rtimes_{\theta} \mathbb{Z}_m$. We study the question of whether G embeds in $B_n/\Gamma_k(P_n)$, where $k \in \{2, 3\}$. By Theorem 1, for such an embedding to exist, $\gcd(|G|, k!) = 1$, and so we shall assume from now on that this is the case. In order to apply Proposition 8, we will make use of a specific embedding of G in S_n studied by Marin in the case where n is prime and $m = (n - 1)/2$ [M2], as well as the restriction to G of the action of S_n on $P_n/\Gamma_2(P_n)$ and $\Gamma_2(P_n)/\Gamma_3(P_n)$ described by equation (5). If $q \in \mathbb{N}$, we will denote the image of an integer r under the canonical projection $\mathbb{Z} \rightarrow \mathbb{Z}_q$ by r_q , or simply by r if no confusion is possible.

Following [M2, proof of Corollary 3.11], we start by describing a homomorphism from G to S_n . Let $(u, v) \in G$, where the elements of G are written with respect to the semi-direct product $\mathbb{Z}_n \rtimes_{\theta} \mathbb{Z}_m$, let $\varphi_{(u,v)}: \mathbb{Z}_n \rightarrow \mathbb{Z}_n$ be the affine transformation defined by:

$$\varphi_{(u,v)}(z) = \theta(v)(z) + u \text{ for all } z \in \mathbb{Z}_n, \quad (10)$$

and let $S_{\mathbb{Z}_n}$ denote the symmetric group on the set \mathbb{Z}_n .

LEMMA 10.

- (a) For all $(u, v) \in G$, the map $\varphi_{(u,v)}: \mathbb{Z}_n \rightarrow \mathbb{Z}_n$ defined in (10) is a bijection.
- (b) Let $\varphi: G \rightarrow S_{\mathbb{Z}_n}$ be the map defined by $\varphi(u, v) = \varphi_{(u,v)}$ for all $(u, v) \in G$. Then φ is a homomorphism. If the action $\theta: \mathbb{Z}_m \rightarrow \text{Aut}(\mathbb{Z}_n)$ is injective then φ is too.

Proof.

(a) If $(u, v) \in G$, the statement follows from the fact that $\theta(v)$ is an automorphism of \mathbb{Z}_n .

(b) Part (a) implies that the map φ is well defined. We now prove that φ is a homomorphism. If $(u_1, v_1), (u_2, v_2) \in G$, then for all $z \in \mathbb{Z}_n$, we have:

$$\begin{aligned} (\varphi_{(u_2,v_2)} \circ \varphi_{(u_1,v_1)})(z) &= \varphi_{(u_2,v_2)}(\theta(v_1)(z) + u_1) = \theta(v_2)(\theta(v_1)(z) + u_1) + u_2 \\ &= \theta(v_2)(\theta(v_1)(z)) + \theta(v_2)(u_1) + u_2 \\ &= \theta(v_2 + v_1)(z) + \theta(v_2)(u_1) + u_2 \\ &= \varphi_{(u_2+\theta(v_2)(u_1), v_2+v_1)}(z) = \varphi_{(u_2,v_2)(u_1,v_1)}(z), \end{aligned}$$

so $\varphi_{(u_2,v_2)} \circ \varphi_{(u_1,v_1)} = \varphi_{(u_2,v_2)(u_1,v_1)}$, and φ is a homomorphism. Finally, suppose that θ is injective, and let $(u, v) \in \text{Ker}(\varphi)$. Then $z = \varphi(u, v)(z) = \varphi_{(u,v)}(z) = \theta(v)(z) + u$ for all $z \in \mathbb{Z}_n$. Taking $z = 0_n$ yields $u = 0_n$. Hence $\theta(v) = \text{Id}_{\mathbb{Z}_n}$, and it follows that $v = 0_m$ by the injectivity of θ , which completes the proof of the lemma. \square

As above, we consider the group $G = \mathbb{Z}_n \rtimes_{\theta} \mathbb{Z}_m$, where $\theta: \mathbb{Z}_m \rightarrow \text{Aut}(\mathbb{Z}_n)$ is the associated action. The element 1_m generates the additive group \mathbb{Z}_m , so $\theta(1_m)$ is an automorphism of \mathbb{Z}_n whose order divides m , and since any automorphism of \mathbb{Z}_n is multiplication by an integer that is relatively prime with n , there exists $1 \leq t < n$ such that $\gcd(t, n) = 1$, $\theta(1_m)$ is multiplication by t , and $t^m \equiv 1 \pmod{n}$. If θ is injective, the order of the automorphism $\theta(1_m)$ is equal to m , and so $t^l \not\equiv 1 \pmod{n}$ for all $1 \leq l < m$,

but this does not imply that $t^l - 1$ is relatively prime with n . However, the condition that $\gcd(t^l - 1, n) = 1$ for all $1 \leq l < m$ is the hypothesis that we require in order to prove Theorem 3, and as we shall now see, implies that θ is injective.

LEMMA 11. Let $n, m \in \mathbb{N}$, let G be a semi-direct product of the form $\mathbb{Z}_n \rtimes_{\theta} \mathbb{Z}_m$, and let $1 \leq t < n$ be such that $\theta(1_m)$ is multiplication in \mathbb{Z}_n by t . Suppose that $\gcd(t^l - 1, n) = 1$ for all $1 \leq l \leq m - 1$.

(a) The action $\theta: \mathbb{Z}_m \rightarrow \text{Aut}(\mathbb{Z}_n)$ is injective, and the homomorphism $\varphi: G \rightarrow S_{\mathbb{Z}_n}$ defined in Lemma 10 is injective.

(b) For all $(u, v) \in G \setminus \{(0_n, 0_m)\}$, the permutation $\varphi(u, v)$ fixes at most one element, and if $v \neq 0_m$ then $\varphi(u, v)$ fixes precisely one element.

Proof.

(a) To prove the first part, we argue by contraposition. Suppose that θ is not injective. Then there exists $1 \leq l < m$ such that $\theta(l_m) = \text{Id}_{\mathbb{Z}_n}$. Now $\theta(1_m)$ is multiplication by t , so $\theta(l_m)$ is multiplication by t^l , and thus $t^l \equiv 1 \pmod{n}$, which implies that $\gcd(t^l - 1, n) \neq 1$. The second part of the statement follows from Lemma 10(b).

(b) Let $(u, v) \in G \setminus \{(0_n, 0_m)\}$. If $v = 0_m$ then $u \neq 0_n$, so $\varphi_{(u, 0_m)}(z) = z + u \neq z$ for all $z \in \mathbb{Z}_n$, hence $\varphi_{(u, 0_m)}$ is fixed-point free. So suppose that $v \neq 0_m$. Since $\theta(v)$ is multiplication by t^v , the corresponding affine transformation $\varphi_{(u, v)}: \mathbb{Z}_n \rightarrow \mathbb{Z}_n$ is given by $\varphi_{(u, v)}(z) = t^v z + u$ for all $z \in \mathbb{Z}_n$. By hypothesis, $\gcd(t^v - 1, n) = 1$, so $t^v - 1$ is invertible in \mathbb{Z}_n , and if $z \in \mathbb{Z}_n$, we have:

$$\varphi_{(u, v)}(z) = z \iff t^v z + u = z \iff (t^v - 1)z = -u \iff z = -u(t^v - 1)^{-1}$$

in \mathbb{Z}_n . Hence $\varphi_{(u, v)}$ possesses a unique fixed point. \square

The above framework enables us to prove Theorem 3 using Proposition 8.

Proof of Theorem 3. Let $\varphi: G \rightarrow S_{\mathbb{Z}_n}$ be the embedding of G in $S_{\mathbb{Z}_n}$ given in Lemma 10(b). By Lemma 11(b), for all $g \in G \setminus \{e\}$, the cycle decomposition of the permutation $\varphi(g)$ contains at most one fixed point. So the embedding φ satisfies the hypotheses of Proposition 8, from which we conclude that G embeds in $B_n/\Gamma_2(P_n)$ (resp. in $B_n/\Gamma_3(P_n)$) if mn is odd (resp. if $\gcd(mn, 6) = 1$). \square

As an application of Theorem 3, we consider groups of the form $\mathbb{Z}_n \rtimes_{\theta} H$, where H is finite, $n = p^r$ is a power of an odd prime p , where $r \in \mathbb{N}$, and the homomorphism $\theta: H \rightarrow \text{Aut}(\mathbb{Z}_{p^r})$ is injective. Recall from [Z, p. 146, lines 16–17] that:

$$\text{Aut}(\mathbb{Z}_{p^r}) \cong \mathbb{Z}_{(p-1)p^{r-1}} \cong \mathbb{Z}_{p-1} \oplus \mathbb{Z}_{p^{r-1}}, \quad (11)$$

where the isomorphisms of (11) are described in [Z, pp. 145–146]. We now prove Corollary 4 by studying injective actions $\theta: H \rightarrow \mathbb{Z}_{(p-1)p^{r-1}}$, where H is a cyclic group whose order is an odd divisor of $p - 1$.

Proof of Corollary 4. Let $p > 2$ be prime, let $p - 1 = 2^j d$, where d is odd, and let d_1 be a divisor of d . So identifying $\text{Aut}(\mathbb{Z}_{p^r})$ with $\mathbb{Z}_{p-1} \oplus \mathbb{Z}_{p^{r-1}}$ via (11), and taking the subgroup H in the above discussion to be \mathbb{Z}_{d_1} , there exists an injective homomorphism

$\theta: \mathbb{Z}_{d_1} \longrightarrow \mathbb{Z}_{p-1} \oplus \mathbb{Z}_{p^{r-1}}$, where $\theta(1_{d_1})$ is an automorphism of \mathbb{Z}_{p^r} given by multiplication by an integer t that is relatively prime with p , and the order d_1 of this automorphism is also relatively prime with p . In particular, the order of t in the group $\mathbb{Z}_{p^r}^*$ is equal to d_1 . We claim that $\gcd(t^l - 1, p) = 1$ for all $0 < l < d_1$. Suppose on the contrary that $t^l - 1$ is divisible by p for some $0 < l < d_1$. Then $t^l = 1 + kp$, where $k \in \mathbb{N}$, and [Z, p. 146, line 12] implies that the order of t^l in $\mathbb{Z}_{p^r}^*$ is a power of p , which contradicts the fact that t is of order d_1 and $\gcd(d_1, p) = 1$. This proves the claim. Part (a) (resp. part (b)) follows using the fact that the order of $\mathbb{Z}_{p^r} \rtimes_{\theta} \mathbb{Z}_{d_1}$ is odd (resp. relatively prime with 6) and by applying Theorem 3. \square

REMARK 12. As we mentioned in the introduction, the results of Corollary 4 are sharp in the sense that if $k \in \{2, 3\}$, the groups $\mathbb{Z}_{p^r} \rtimes_{\theta} \mathbb{Z}_d$ satisfy $\ell_k(G) = m(G) = p^r$ if $d > 1$, where $k \in \{2, 3\}$. It is unlikely that this will be the case in general, in other words, if n is not a prime power, it may be possible to embed groups of the form $\mathbb{Z}_n \rtimes_{\theta} H$ in $B_r/\Gamma_k(P_r)$ for some $r < n$.

4.2 Further examples

In this final section, we give examples of two semi-direct products of the form $\mathbb{Z}_9 \rtimes_{\theta} \mathbb{Z}_3$ and $(\mathbb{Z}_3 \oplus \mathbb{Z}_3) \rtimes_{\theta} \mathbb{Z}_3$ respectively that do not satisfy the hypotheses of Theorem 3, but that embed in $B_9/\Gamma_2(P_9)$. We start with some general comments. If p is an odd prime, consider the group $G = \mathbb{Z}_{p^r} \rtimes_{\theta} \mathbb{Z}_{p^{r-1}(p-1)}$, where with respect to the notation of [Z, p. 146, line 8], the homomorphism $\theta: \mathbb{Z}_{p^{r-1}(p-1)} \longrightarrow \text{Aut}(\mathbb{Z}_{p^r})$ sends $1_{p^{r-1}(p-1)}$ to the element of $\text{Aut}(\mathbb{Z}_{p^r})$ given by multiplication in \mathbb{Z}_{p^r} by t' , where $t' = (1+p)g_1$, and where the order of g_1 (resp. $1+p$) is equal to $p-1$ (resp. p^{r-1}) in the multiplicative group $\mathbb{Z}_{p^r}^*$. If d_1 is an odd divisor of $p^{r-1}(p-1)$, let $q = p^{r-1}(p-1)/d_1$, and consider the subgroup $\mathbb{Z}_{p^r} \rtimes_{\theta'} \mathbb{Z}_{d_1}$ of G , where \mathbb{Z}_{d_1} is the subgroup of $\mathbb{Z}_{p^{r-1}(p-1)}$ of order d_1 , and $\theta': \mathbb{Z}_{d_1} \longrightarrow \text{Aut}(\mathbb{Z}_{p^r})$ is the restriction of θ to \mathbb{Z}_{d_1} . Then $\theta'(1_d) = \theta(q_{p^{r-1}(p-1)})$ is multiplication by $t = t'^q$ in \mathbb{Z}_{p^r} , and by injectivity, t is of order d_1 in $\mathbb{Z}_{p^r}^*$. If further d_1 is divisible by p then $t^{d_1/p}$ is of order p in $\mathbb{Z}_{p^r}^*$, and by [Z, p. 146, line 12], $t^{d_1/p} \equiv (1+p)^{\lambda p^{r-2}} \pmod{p^r}$, where $\lambda \in \mathbb{N}$ and $\gcd(\lambda, p) = 1$. It follows that $t^{d_1/p} - 1$ is divisible by p , and since $0 < d_1/p < d_1$, the hypotheses of Theorem 3 are not satisfied for the group $\mathbb{Z}_{p^r} \rtimes_{\theta'} \mathbb{Z}_{d_1}$. As Example 13(a) below shows, if $p = 3$ and $r = 2$, such a group may nevertheless embed in $B_{p^r}/\Gamma_2(P_{p^r})$. In Example 13(b), we show that the other non-Abelian group of order 27, which is of the form $(\mathbb{Z}_p \oplus \mathbb{Z}_p) \rtimes \mathbb{Z}_p$, also embeds in $B_9/\Gamma_2(P_9)$. It does not satisfy the hypotheses of Theorem 3 either.

EXAMPLES 13. Suppose that G is a group of order 27 that embeds in S_9 . If $k = 2$ and $n = 9$ in (2), taking the preimage of G by $\bar{\sigma}$ leads to the following short exact sequence:

$$1 \longrightarrow P_9/\Gamma_2(P_9) \longrightarrow \bar{\sigma}^{-1}(G) \xrightarrow{\bar{\sigma}|_{\bar{\sigma}^{-1}(G)}} G \longrightarrow 1, \quad (12)$$

where the rank of the free Abelian group $P_9/\Gamma_2(P_9)$ is equal to 36. As we shall now see, if G is one of the two non-Abelian groups of order 27, the action of S_9 on the basis \mathcal{B} of $P_9/\Gamma_2(P_9)$ given by (5) restricts to an action of G on \mathcal{B} for which there are two orbits,

that of $A_{1,2}$, which contains 9 elements, and is given by:

$$\mathcal{O} = \{A_{1,2}, A_{8,9}, A_{5,6}, A_{2,3}, A_{7,8}, A_{4,5}, A_{1,3}, A_{7,9}, A_{4,6}\}, \quad (13)$$

and that of $A_{5,9}$, which contains the remaining 27 elements of \mathcal{B} . Let H be the subgroup of $P_9/\Gamma_2(P_9)$ generated by the orbit of $A_{5,9}$. Then $H \cong \mathbb{Z}^{27}$, and H is not normal in $B_9/\Gamma_2(P_9)$, but it is normal in the subgroup $\bar{\sigma}^{-1}(G)$ of $B_9/\Gamma_2(P_9)$ since the basis $\mathcal{B} \setminus \mathcal{O}$ of H is invariant under the action of G . We thus have an extension:

$$1 \longrightarrow H \longrightarrow \bar{\sigma}^{-1}(G) \xrightarrow{\pi} \tilde{H} \longrightarrow 1, \quad (14)$$

where $\tilde{H} = \bar{\sigma}^{-1}(G)/H$, and $\pi: \bar{\sigma}^{-1}(G) \rightarrow \tilde{H}$ is the canonical projection. Equation (12) induces the following short exact sequence:

$$1 \longrightarrow (P_9/\Gamma_2(P_9))/H \longrightarrow \tilde{H} \xrightarrow{\tilde{\sigma}} G \longrightarrow 1, \quad (15)$$

where $\tilde{\sigma}: \tilde{H} \rightarrow G$ is the surjective homomorphism induced by $\bar{\sigma}$, and we also have an extension:

$$1 \longrightarrow H \longrightarrow P_9/\Gamma_2(P_9) \xrightarrow{\rho} (P_9/\Gamma_2(P_9))/H \longrightarrow 1, \quad (16)$$

obtained from the canonical projection $\rho: P_9/\Gamma_2(P_9) \rightarrow (P_9/\Gamma_2(P_9))/H$. From the construction of H , and using the fact that $P_9/\Gamma_2(P_9)$ (resp. H) is the free Abelian group of rank 36 (resp. 27) for which \mathcal{B} (resp. $\mathcal{B} \setminus \mathcal{O}$) is a basis, the kernel of (15) is isomorphic to \mathbb{Z}^9 and a basis is given by the H -cosets of the nine elements of \mathcal{O} . Further, the restriction of ρ to the subgroup of $P_9/\Gamma_2(P_9)$ generated by \mathcal{O} is an isomorphism, and thus the set $\rho(\mathcal{O})$ is a basis of $(P_9/\Gamma_2(P_9))/H$, which we denote by \mathcal{O}' . In the examples below, we shall construct an explicit embedding $\iota: G \rightarrow \tilde{H}$ of G in \tilde{H} . This being the case, using (14), we thus obtain the following short exact sequence:

$$1 \longrightarrow H \longrightarrow \pi^{-1}(\iota(G)) \xrightarrow{\pi|_{\pi^{-1}(\iota(G))}} \iota(G) \longrightarrow 1. \quad (17)$$

Now H is isomorphic to $\mathbb{Z}[G]$, and the action of $\iota(G)$ on H is given via (5). It follows from Proposition 7 that (17) splits. Hence G embeds in $\pi^{-1}(\iota(G))$, which is a subgroup of $\bar{\sigma}^{-1}(G)$, and so it embeds in $B_9/\Gamma_2(P_9)$. We could have attempted to embed G in $B_9/\Gamma_2(P_9)$ directly via (12). However one of the difficulties with this approach is that the rank of the kernel is 36, whereas that of the kernel of (15) is much smaller, and this decreases greatly the number of calculations needed to show that G embeds in \tilde{H} . We now give the details of the computations of this embedding in the two cases.

(a) Consider the semi-direct product $\mathbb{Z}_9 \rtimes_{\theta'} \mathbb{Z}_3$, where $\theta': \mathbb{Z}_3 \rightarrow \text{Aut}(\mathbb{Z}_9)$ is as defined in the second paragraph of this subsection. We have $g_1 = 8$, so $\theta'(1_3)$ is multiplication by $(1+3) \cdot (8) = 32 \equiv 5 \pmod{9}$. It will be more convenient for us to work with a different automorphism, but that gives rise to a semi-direct product that is isomorphic to $\mathbb{Z}_9 \rtimes_{\theta'} \mathbb{Z}_3$ as follows. Let $\varphi: \mathbb{Z}_9 \rightarrow \mathbb{Z}_9$ be the automorphism given by multiplication by 4, let $\theta: \mathbb{Z}_3 \rightarrow \text{Aut}(\mathbb{Z}_9)$ be such that $\theta(1_3) = \varphi$, and let $G = \mathbb{Z}_9 \rtimes_{\theta} \mathbb{Z}_3$. Then $\theta'(1_3) = \varphi \circ \theta(1_3)$, which implies that the groups G and $\mathbb{Z}_9 \rtimes_{\theta'} \mathbb{Z}_3$ are isomorphic, see [AB, Chap. 1.2, Proposition 12]. It is clear that θ is injective, but that the hypothesis of Theorem 3 that $\gcd(t^l - 1, n) = 1$ for all $1 \leq l < d$ is not satisfied if $l = 1$.

Nevertheless, the group G embeds in $B_9/\Gamma_2(P_9)$. To see this, consider the elements $\alpha = (1, 2, 3)(4, 5, 6)(7)(8)(9)$ and $\beta = (1, 4, 7, 3, 5, 8, 2, 6, 9)$ of S_9 . Then:

$$\begin{aligned}\alpha\beta\alpha^{-1} &= (1, 2, 3)(4, 5, 6)(7)(8)(9) \cdot (1, 4, 7, 3, 5, 8, 2, 6, 9) \cdot ((1, 2, 3)(4, 5, 6)(7)(8)(9))^{-1} \\ &= (1, 5, 9, 3, 6, 7, 2, 4, 8) = (1, 4, 7, 3, 5, 8, 2, 6, 9)^4 = \beta^4,\end{aligned}$$

and an embedding of G in S_9 is realised by sending 1_3 (resp. 1_9) to α (resp. β). From now on, we identify G with its image in S_9 under this embedding. One may check that \mathcal{O} and $\mathcal{B} \setminus \mathcal{O}$ are the two orbits arising from the action of G on \mathcal{B} given by (5) (for future reference, note that the order of the elements of \mathcal{O} is that obtained by the action of successive powers of β). We define the map $\iota: G \rightarrow \tilde{H}$ on the generators of G by $\iota(\alpha) = \hat{\alpha}$ and $\iota(\beta) = \hat{\beta}$, where $\hat{\alpha} = \sigma_2\sigma_1^{-1}\sigma_5\sigma_4^{-1}$ and $\hat{\beta} = A_{1,2}A_{8,9}^{-1}w\sigma_8\sigma_7\sigma_6\sigma_5\sigma_4^{-1}\sigma_3^{-1}\sigma_2^{-1}\sigma_1^{-1}w^{-1}$, and where $w = \sigma_3\sigma_6\sigma_2\sigma_3\sigma_4\sigma_5\sigma_4\sigma_3\sigma_7\sigma_6$. By abuse of notation, we will denote an element of $B_9/\Gamma_2(P_9)$ in the same way as its projection on the quotient $(B_9/\Gamma_2(P_9))/H$. We claim that $\hat{\alpha}\hat{\beta}\hat{\alpha}^{-1}\hat{\beta}^{-4} = 1$ in \tilde{H} , from which we may conclude that ι extends to a group homomorphism. To prove the claim, using [GG01, Proposition 12] and the action of β on the orbit of $A_{1,2}$ mentioned above, first note that:

$$\begin{aligned}\hat{\alpha}\hat{\beta}\hat{\alpha}^{-1}\hat{\beta}^{-4} &= \hat{\alpha}A_{1,2}A_{8,9}^{-1}\hat{\alpha}b\hat{\alpha}^{-1}(A_{1,2}A_{8,9}^{-1}b)^{-4} \\ &= A_{1,3}A_{8,9}^{-1}\hat{\alpha}b\hat{\alpha}^{-1}b^{-4} \cdot b^3A_{1,2}^{-1}A_{8,9}b^{-3} \cdot b^2A_{1,2}^{-1}A_{8,9}b^{-2} \cdot bA_{1,2}^{-1}A_{8,9}b^{-1} \cdot A_{1,2}^{-1}A_{8,9} \\ &= A_{1,3}A_{8,9}^{-1} \cdot \hat{\alpha}b\hat{\alpha}^{-1}b^{-4} \cdot A_{2,3}^{-1}A_{7,8} \cdot A_{5,6}^{-1}A_{2,3} \cdot A_{8,9}^{-1}A_{5,6} \cdot A_{1,2}^{-1}A_{8,9} \\ &= A_{1,2}^{-1}A_{7,8}A_{1,3}A_{8,9}^{-1}\hat{\alpha}b\hat{\alpha}^{-1}b^{-4}.\end{aligned}\tag{18}$$

To obtain the final equality, we have also used the fact that $\hat{\alpha}b\hat{\alpha}^{-1}b^{-4}$ belongs to the quotient $(P_9/\Gamma_2(P_9))/H$, so commutes with the other terms in the expression. To compute $\hat{\alpha}b\hat{\alpha}^{-1}b^{-4}$ in terms of the basis \mathcal{O}' of $(P_9/\Gamma_2(P_9))/H$, we use the method of crossing numbers given in [GG01, Proposition 15], except that since we are working in $(P_9/\Gamma_2(P_9))/H$, using the isomorphism $\rho|_{\langle \mathcal{O} \rangle}: \langle \mathcal{O} \rangle \rightarrow (P_9/\Gamma_2(P_9))/H$ induced by (16), it suffices to compute the crossing numbers of the pairs of strings corresponding to the elements of \mathcal{O} given in (13). Using the braid $\hat{\alpha}b\hat{\alpha}^{-1}b^{-4}$ illustrated in Figure 1, one may verify that:

$$\hat{\alpha}b\hat{\alpha}^{-1}b^{-4} = A_{1,2}A_{1,3}^{-1}A_{7,8}^{-1}A_{8,9}\tag{19}$$

in the quotient $(P_9/\Gamma_2(P_9))/H$. It follows from equations (18) and (19) that $\hat{\alpha}\hat{\beta}\hat{\alpha}^{-1}\hat{\beta}^{-4} = 1$ in \tilde{H} , which proves the claim. Thus $\langle \hat{\alpha}, \hat{\beta} \rangle$ is a quotient of G , but since it is non-Abelian, and the only non-Abelian quotient of G is itself, we conclude that $\langle \hat{\alpha}, \hat{\beta} \rangle \cong G$, and hence ι is an embedding. It follows from the discussion at the beginning of these examples that G embeds in $\pi^{-1}(\iota(G))$, and therefore in $B_9/\Gamma_2(P_9)$.

(b) We now give an explicit example of a non-Abelian group G of the form $A \rtimes \mathbb{Z}_m$ that embeds in $B_{|A|}/\Gamma_2(P_{|A|})$, where A is a non-cyclic finite Abelian group. To our knowledge, this is the first explicit example of a finite group that embeds in such a quotient but that is not a semi-direct product of two cyclic groups. In particular, this subgroup does not satisfy the hypothesis of Theorem 3 either. Let G be the Heisenberg group mod p of order p^3 , where p is an odd prime. There exists an extension of the form:

$$1 \rightarrow \mathbb{Z}_p \rightarrow G \rightarrow \mathbb{Z}_p \oplus \mathbb{Z}_p \rightarrow 1,\tag{20}$$






Figure 1: The braid $\hat{a}b\hat{a}^{-1}b^{-4}$



(a) The braid $[\hat{\alpha}, \hat{\beta}] \cdot \hat{\gamma}^{-1}$



(b) The braid $[\hat{\alpha}, \hat{\gamma}]$

Figure 2: The braids $[\hat{\alpha}, \hat{\beta}] \cdot \hat{\gamma}^{-1}$ and $[\hat{\alpha}, \hat{\gamma}]$

and a presentation of G is given by:

$$\langle a, b, c \mid a^p = b^p = c^p = 1, c = [a, b] \text{ and } [a, c] = [b, c] = 1 \rangle, \quad (21)$$

where c is an element of G emanating from a generator of the kernel \mathbb{Z}_p of the extension (20), and a and b are elements of G that project to the generators of the summands of the quotient. This group is also isomorphic to $(\mathbb{Z}_p \oplus \mathbb{Z}_p) \rtimes_{\theta} \mathbb{Z}_p$, where the action $\theta: \mathbb{Z}_p \rightarrow \text{Aut}(\mathbb{Z}_p \oplus \mathbb{Z}_p)$ is given by $\theta(1_p) = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. From now on, we assume that $p = 3$. Consider the map from G to S_9 given by sending a to $\alpha = (1, 4, 7)(2, 5, 8)(3, 6, 9)$ and b to $\beta = (1)(2)(3)(4, 5, 6)(7, 9, 8)$, so c is sent to $\gamma = [\alpha, \beta] = (1, 2, 3)(4, 5, 6)(7, 8, 9)$. The relations of (21) hold for the elements α, β and γ , and since the only non-Abelian quotient of G is G itself, it follows that this map extends to an embedding of G in S_9 . Once more, one may check that \mathcal{O} and $\mathcal{B} \setminus \mathcal{O}$ are the orbits arising from the action of G on \mathcal{B} . It remains to show that G embeds in \tilde{H} . Let $\iota: G \rightarrow \tilde{H}$ be the map defined by $\iota(\alpha) = \hat{\alpha}$, $\iota(\beta) = \hat{\beta}$ and $\iota(\gamma) = \hat{\gamma}$, where:

$$\hat{\alpha} = w' \sigma_2 \sigma_1^{-1} \sigma_5 \sigma_4^{-1} \sigma_8 \sigma_7^{-1} w'^{-1}, \hat{\beta} = \sigma_5 \sigma_4^{-1} \sigma_7 \sigma_8^{-1} \text{ and } \hat{\gamma} = \sigma_2 \sigma_1^{-1} \sigma_5 \sigma_4^{-1} \sigma_8 \sigma_7^{-1},$$

and where $w' = \sigma_3 \sigma_2 \sigma_4 \sigma_6 \sigma_5 \sigma_4 \sigma_3 \sigma_7 \sigma_6$. Using the notation of [GG01, equations (14) and (16)], $\hat{\alpha} = w' \delta_{0,3} \delta_{3,3} \delta_{6,3} w'^{-1} = w' \delta(0, 3, 3, 3) w'^{-1}$, $\hat{\beta} = \delta_{3,3} \delta_{6,3}^{-1}$ and $\hat{\gamma} = w' \hat{\alpha} w'^{-1}$. So these three elements are of order 3 in $B_9/\Gamma_2(P_9)$ by the argument of [GG01, line 4, p. 412], and hence they satisfy the first three relations of (21) in \tilde{H} , a, b and c being taken to be $\hat{\alpha}, \hat{\beta}$ and $\hat{\gamma}$ respectively. One may check in a straightforward manner that $[\hat{\beta}, \hat{\gamma}] = 1$ as elements of B_9 , hence $[\hat{\beta}, \hat{\gamma}] = 1$ in \tilde{H} . To see that the two remaining relations of (21) hold, as in the first example, we use the method of crossing numbers of the strings given in [GG01, Proposition 15], but in \tilde{H} rather than $P_9/\Gamma_2(P_9)$. In this way, we see from Figures (2)(a) and (b) that $[\hat{\alpha}, \hat{\beta}] \cdot \hat{\gamma}^{-1} = 1$ and $[\hat{\alpha}, \hat{\gamma}] = 1$ in \tilde{H} . It thus follows that $\langle \hat{\alpha}, \hat{\beta}, \hat{\gamma} \rangle$ is a quotient of G , but since this subgroup is non-Abelian, and the only non-Abelian quotient of G is itself, we conclude that $\langle \hat{\alpha}, \hat{\beta} \rangle \cong G$, and hence ι is an embedding. Once more, it follows from the discussion at the beginning of these examples that G embeds in $\pi^{-1}(\iota(G))$, and therefore in $B_9/\Gamma_2(P_9)$.

REMARKS 14.

(a) Let G be one of the two groups of order 27 analysed in Examples 13. With the notation introduced at the beginning of Section 4, the fact that G embeds in $B_9/\Gamma_2(P_9)$ implies that $\ell_2(G) \leq 9$. On the other hand, if G embeds in S_r then $r \geq 9$ by Lagrange's Theorem. Hence $m(G) \geq 9$, and it follows from (3) that $m(G) = \ell_2(G) = 9$.

(b) We would like to be able to determine the finite groups of the form $A \rtimes_{\theta} \mathbb{Z}_m$, where A is a finite Abelian group, that embed in $B_{|A|}/\Gamma_2(P_{|A|})$.

References

- [AB] J. L. Alperin and R. B. Bell, Groups and representations, Graduate Texts in Mathematics **162**, Springer-Verlag, New York (1995).
- [BM] V. Beck and I. Marin, Torsion subgroups of quasi-Abelianized braid groups, [arXiv:1709.01853](https://arxiv.org/abs/1709.01853).

- [Br] K. Brown, *Cohomology of groups*, Graduate Texts in Mathematics **87**, Springer-Verlag, New York-Berlin (1982).
- [FR] M. Falk and R. Randell, The lower central series of a fiber-type arrangement. *Invent. Math.* **82** (1985), 77–88.
- [GGO1] D. L. Gonçalves, J. Guaschi and O. Ocampo, A quotient of the Artin braid groups related to crystallographic groups, *J. Algebra* **474** (2017), 393–423.
- [GGO2] D. L. Gonçalves, J. Guaschi and O. Ocampo, Almost-crystallographic groups as quotients of Artin braid groups, preprint (2017).
- [H] V. L. Hansen, *Braids and Coverings: selected topics*, London Math. Soc. Student Text **18**, Cambridge University Press, 1989.
- [K] T. Kohno, Série de Poincaré-Koszul associée aux groupes de tresses pures, *Invent. Math.* **82** (1985), 57–75.
- [M1] I. Marin, On the representation theory of braid groups, *Ann. Math. Blaise Pascal* **20** (2013), 193–260.
- [M2] I. Marin, Crystallographic groups and flat manifolds from complex reflection groups, *Geom. Dedicata* **182** (2016), 233–247.
- [PS] F. Panaite and M. Staic, A quotient of the braid group related to pseudosymmetric braided categories, *Pacific J. Math.* **244** (2010), 155–167.
- [T] J. Tits, Normalisateurs de tores I : groupes de Coxeter étendus, *J. Algebra* **4** (1966), 96–116.
- [Z] H. Zassenhaus, *The theory of groups*, Reprint of the second (1958) edition, Dover Publications Inc., Mineola, NY, 1999.