

Supporting Information

Low-lying excited states of model proteins:

Performances of the CC2 method versus multireference methods

Nadia Ben Amor^{1,2}, Sophie Hoyau², Daniel Maynau^{1,2}, Valérie Brenner³

¹*CNRS, UPS, LCPQ (Laboratoire de Chimie et Physique Quantiques), IRSAMC,
118, rte de Narbonne F-31062 Toulouse Cedex. France.*

²*Université de Toulouse, UPS, LCPQ (Laboratoire de Chimie et Physique
Quantiques), IRSAMC, 118, rte de Narbonne F-31062 Toulouse Cedex. France.*

³*Laboratoire Interactions, Dynamiques et Lasers, LIDYL, CEA, CNRS, Université
Paris-Saclay, 91191 Gif-sur-Yvette Cedex, France*

Appendix S1: Cartesian coordinates of all the geometries of the NAPA B conformer.

Table S1-1: Cartesian coordinates of the B97-D2/TZVPP optimized ground state geometry (Å).

| | | | |
|---|------------|------------|------------|
| C | 2.1012809 | 2.8012753 | -0.7962777 |
| H | 2.6829063 | 2.8102244 | -1.7245541 |
| H | 2.5283087 | 3.5605002 | -0.1314497 |
| C | 2.2623085 | 1.4411635 | -0.1341555 |
| O | 3.3725270 | 0.9987098 | 0.1749704 |
| N | 1.1086496 | 0.7565672 | 0.0869590 |
| H | 0.2393554 | 1.1443544 | -0.2536865 |
| C | 1.0243877 | -0.5231547 | 0.7911005 |
| H | 1.8269094 | -0.5134581 | 1.5407075 |
| C | 1.3400483 | -1.7462516 | -0.1295981 |
| O | 0.5981846 | -2.7152051 | -0.2351044 |
| C | -0.3343449 | -0.6562608 | 1.5214627 |
| H | -0.4399299 | -1.7064709 | 1.8095437 |
| H | -0.2778540 | -0.0512285 | 2.4333322 |
| C | -1.5369592 | -0.2132956 | 0.7138013 |
| C | -2.1280013 | -1.0502808 | -0.2454252 |
| H | -1.7207727 | -2.0451963 | -0.3957499 |
| C | -2.0613549 | 1.0770206 | 0.8942465 |
| H | -1.6167840 | 1.7278717 | 1.6457609 |
| C | -3.2139170 | -0.6036792 | -1.0020174 |
| H | -3.6629927 | -1.2651954 | -1.7394558 |
| C | -3.1467592 | 1.5269869 | 0.1356274 |
| H | -3.5416242 | 2.5279347 | 0.2943963 |
| C | -3.7254001 | 0.6854571 | -0.8175615 |
| H | -4.5712026 | 1.0289782 | -1.4085182 |
| N | 2.5389948 | -1.6466053 | -0.7580640 |
| H | 3.1694248 | -0.8769566 | -0.5480749 |
| H | 1.0590375 | 3.0589417 | -1.0161458 |
| H | 2.8460937 | -2.4248093 | -1.3223449 |

Table S1-2: Cartesian coordinates of the CC2/cc-pVDZ optimized M_{ππ*} excited state geometry (Å).

| | | | |
|---|------------|------------|------------|
| C | -2.1252717 | -1.1546159 | -0.2226251 |
| C | -1.4713441 | -0.2826615 | 0.7293417 |
| C | -1.9349089 | 1.0883775 | 0.8621547 |
| C | -3.0483759 | 1.5583897 | 0.0760321 |
| C | -3.6857968 | 0.6814654 | -0.8690932 |
| C | -3.2248664 | -0.6720585 | -1.0206385 |
| C | -0.2951474 | -0.7452156 | 1.5369858 |
| C | 1.0498089 | -0.5759332 | 0.7973685 |
| C | 1.2975704 | -1.7314188 | -0.2001655 |
| O | 0.5385732 | -2.7046127 | -0.3087644 |
| N | 1.1010772 | 0.7390793 | 0.1722200 |

| | | | |
|---|------------|------------|------------|
| C | 2.2692556 | 1.4061498 | -0.0572559 |
| C | 2.1185112 | 2.8033248 | -0.6258816 |
| N | 2.4460980 | -1.5724348 | -0.9158344 |
| O | 3.3857242 | 0.8992651 | 0.1719711 |
| H | 0.2109399 | 1.1618115 | -0.0921014 |
| H | 1.8863566 | -0.6115673 | 1.5215295 |
| H | -0.2436729 | -0.1626483 | 2.4735101 |
| H | -0.3952562 | -1.8139177 | 1.7858947 |
| H | -1.4967595 | 1.7307338 | 1.6337328 |
| H | -3.4106788 | 2.5826595 | 0.2023664 |
| H | -4.5254285 | 1.0463043 | -1.4671186 |
| H | -3.7060896 | -1.3379560 | -1.7416317 |
| H | -1.7538018 | -2.1751941 | -0.3353733 |
| H | 2.7395599 | -2.3517642 | -1.4963761 |
| H | 3.1136449 | -0.8520683 | -0.6304265 |
| H | 2.5028856 | 2.8104427 | -1.6577124 |
| H | 1.0781531 | 3.1613156 | -0.6236561 |
| H | 2.7442935 | 3.4879346 | -0.0348836 |

Table S1-3: Cartesian coordinates of the CC2/cc-pVDZ CI_{ππ*/nπ*co(2)} geometry connecting the two CC2/cc-pVDZ optimized equilibrium geometries of M_{ππ*} and M_{nπ*co(2)} excited states (Å).

| | | | |
|---|------------|------------|------------|
| C | 0.0000000 | 0.0000000 | 0.0000000 |
| C | 0.0000000 | 0.0000000 | 1.4312773 |
| C | 1.2391763 | 0.0000000 | 2.1599976 |
| C | 2.4796286 | -0.0080348 | 1.4262933 |
| C | 2.4748325 | 0.0057958 | -0.0058128 |
| C | 1.2359834 | 0.0033726 | -0.7178467 |
| C | 1.2578136 | -0.0536430 | 3.6614461 |
| C | 1.1350762 | -1.4933553 | 4.2003383 |
| N | 2.0850246 | -2.3611405 | 3.5209949 |
| C | 2.5855665 | -3.4925577 | 4.1062350 |
| O | 2.2510507 | -3.8589728 | 5.2486638 |
| C | -0.3098479 | -2.0014229 | 4.0938291 |
| N | -0.4697529 | -3.3215271 | 4.4758722 |
| O | -1.2970638 | -1.1985291 | 3.9749218 |
| C | 3.5697526 | -4.2744402 | 3.2584141 |
| H | 2.2817155 | -2.1512616 | 2.5424468 |
| H | 1.3995750 | -1.5163068 | 5.2760373 |
| H | 2.2023650 | 0.3755308 | 4.0378391 |
| H | 0.4154747 | 0.5224126 | 4.0786254 |
| H | 3.4271446 | 0.0508287 | 1.9734036 |
| H | 3.4216053 | 0.0229433 | -0.5538155 |
| H | 1.2363642 | 0.0103754 | -1.8115672 |
| H | -0.9492037 | -0.0031297 | -0.5429853 |
| H | -0.9385859 | -0.0125117 | 1.9915155 |
| H | -1.4361344 | -3.6306501 | 4.5583056 |
| H | 0.2581229 | -3.7212198 | 5.0779238 |
| H | 3.0752606 | -5.1881297 | 2.8924693 |
| H | 3.9457162 | -3.6994674 | 2.3989321 |
| H | 4.4119887 | -4.5817866 | 3.8945851 |

Table S1-4: Cartesian coordinates of the CC2/cc-pVDZ optimized $M_{n\pi^*co}(2)$ excited state geometry (\AA).

| | | | |
|---|------------|------------|------------|
| C | -2.0710079 | -1.0323881 | -0.2022098 |
| C | -1.5678491 | -0.1272217 | 0.7592886 |
| C | -2.1114537 | 1.1725455 | 0.8201167 |
| C | -3.1397784 | 1.5646740 | -0.0575728 |
| C | -3.6349389 | 0.6568083 | -1.0086910 |
| C | -3.0996247 | -0.6434017 | -1.0756759 |
| C | -0.4173462 | -0.5278243 | 1.6501587 |
| C | 0.9184000 | -0.4323210 | 0.8960325 |
| C | 0.9368428 | -1.4163059 | -0.2450883 |
| O | 0.7860441 | -2.7549114 | 0.0447886 |
| N | 1.1586968 | 0.9208466 | 0.4339294 |
| C | 2.4444299 | 1.3482369 | 0.1783934 |
| C | 2.5338368 | 2.6961717 | -0.5110960 |
| N | 2.0533078 | -1.3447090 | -1.1472043 |
| O | 3.4441780 | 0.6827201 | 0.4898223 |
| H | 0.3913168 | 1.3706418 | -0.0635912 |
| H | 1.7523961 | -0.6911897 | 1.5819539 |
| H | -0.3695385 | 0.1180135 | 2.5422971 |
| H | -0.5315591 | -1.5738024 | 1.9843057 |
| H | -1.7336468 | 1.8775548 | 1.5700987 |
| H | -3.5562588 | 2.5750893 | 0.0071629 |
| H | -4.4376838 | 0.9568293 | -1.6893331 |
| H | -3.4857796 | -1.3567780 | -1.8107803 |
| H | -1.6542363 | -2.0450905 | -0.2529732 |
| H | 2.0729734 | -2.1822416 | -1.7380872 |
| H | 2.9431410 | -1.2352645 | -0.6336886 |
| H | 2.5656000 | 2.5342368 | -1.6010837 |
| H | 1.6751686 | 3.3427689 | -0.2752528 |
| H | 3.4679020 | 3.1886045 | -0.2095565 |

Table S1-5: Cartesian coordinates of the CC2/cc-pVDZ partially optimized M'_{CT} excited state geometry with the NH(1) bond length constrained to 1.216 \AA (\AA).

| | | | |
|---|------------|------------|------------|
| C | -2.4030317 | -0.8977387 | 0.1859705 |
| C | -1.4668265 | -0.1348730 | 0.8751197 |
| C | -1.2910596 | 1.3273539 | 0.6039599 |
| C | -2.3001929 | 1.8955076 | -0.3096010 |
| C | -3.2288935 | 1.1064093 | -0.9659611 |
| C | -3.3164430 | -0.3027639 | -0.7247173 |
| C | -0.3567674 | -0.7512422 | 1.6775734 |
| C | 0.9596593 | -0.6674531 | 0.8698808 |
| C | 0.8992366 | -1.8114093 | -0.3029468 |
| O | 0.0960084 | -2.7301264 | -0.2093568 |
| N | 1.0911097 | 0.6132648 | 0.3090830 |
| C | 2.3376747 | 1.1563660 | -0.0575722 |
| C | 2.2772910 | 2.6462432 | -0.2817687 |
| N | 1.8202098 | -1.6192598 | -1.2693289 |

| | | | |
|---|------------|------------|------------|
| O | 3.3655615 | 0.4684626 | -0.1771118 |
| H | 0.0037979 | 1.1547614 | 0.2609467 |
| H | 1.8786188 | -0.9756284 | 1.4151467 |
| H | -0.1812596 | -0.1971147 | 2.6205915 |
| H | -0.5406372 | -1.8128072 | 1.9123216 |
| H | -1.1385370 | 1.9258629 | 1.5373266 |
| H | -2.2913194 | 2.9786670 | -0.4954559 |
| H | -3.9311051 | 1.5736053 | -1.6679871 |
| H | -4.0738825 | -0.9109348 | -1.2296265 |
| H | -2.4360459 | -1.9815602 | 0.3579954 |
| H | 1.9668960 | -2.3988960 | -1.9076839 |
| H | 2.6122350 | -1.0095121 | -1.0416360 |
| H | 3.2489672 | 3.0094273 | -0.6402510 |
| H | 1.4841348 | 2.8743662 | -1.0115153 |
| H | 2.0083962 | 3.1503138 | 0.6600804 |

Appendix S2: Dependence of the excitation energy and reference weight of the zeroth-order wave function on the level shift in CASPT2 calculations of the excited states




Figure S2-1: CASPT2 excitation energy as a function of the level shift (IPEA default value 0.25).




Figure S2-2: Reference weight of the zeroth-order wave function as a function of the level shift (IPEA default value 0.25).

Appendix S3: Diffuse orbital effects on the excitation energy in CC2 and MS-CASPT2 calculations.

| ΔE (eV) | CC2 <i>{cc-pVDZ/aug-ccpVDZ}</i> | CC2 <i>{cc-pVTZ/aug-ccpVTZ}</i> | MS-CASPT2 <i>20R/{cc-pVDZ/aug-ccpVDZ}</i> |
|---|---------------------------------------|--|---|
| $\pi_{\text{cycle}} \rightarrow \pi^*_{\text{cycle}}$ | 5.22 [0.89] | 5.14 [0.85] | 5.17 [0.74] |
| $n \rightarrow \pi^*_{\text{CO}}$ | 5.50 [0.63] | 5.44 [0.53] + CT [0.10] | 5.81 [0.54] + LE $\pi_{\text{CO}}\pi^*_{\text{CO}}$ [0.19] |
| $n \rightarrow \pi^*_{\text{CO}}$ | 5.79 [0.73] | 5.72 [0.73] | 6.07 [0.501] + LE $\pi_{\text{CO}}\pi^*_{\text{CO}}$ [0.18] |
| $\pi_{\text{cycle}} \rightarrow \pi^*_{\text{cycle}}$ | 6.36 [0.74] | 6.27 [0.72] | 6.45 [0.86] |
| <i>MR</i> | | | 6.69 LEπ [0.22] + LE _{pep} [0.18] + CT [0.11] |
| <i>MR</i> | | | 6.86 CT [0.28] + LE $\pi_{\text{CO}}\pi^*_{\text{CO}}$ [0.24] + LE _{pep} [0.21] + LE π [0.12] |
| $\pi_{\text{cycle}} \rightarrow \pi^*_{\text{cycle}}$ | 6.44[0.48]+ CT[0.33] | | 6.95 [0.63] + CT [0.10] |
| $n \rightarrow \pi^*_{\text{cycle}}$ | 6.53 [0.43]+ LE _{pep} [0.20] | 6.34 [0.29] + LE π [0.27] + LE _{pep} [0.13] | 7.15 [0.53] + LE π [0.17] |
| $\pi_{\text{cycle}} \rightarrow \pi^*_{\text{cycle}}$ | 6.59 [0.38]+ CT[0.37] | | |
| $n \rightarrow \pi^*_{\text{cycle}}$ | 6.69 [0.48] + LE π [0.11] | 6.44 [0.38] + LE _{pep} [0.15] + LE π [0.14] | 7.21 [0.34] + LEπ [0.39] |

Table S3: CC2 and MS-CASPT2 excitation energy (eV) of the excited states of the NAPA B conformer at its DFT-D optimized ground state geometry (Figure 1(a) and Table S1-1) calculated within adapted basis sets (see text). All states up to the two first CT states are reported. For the definition of the values in bracket, see captions of Tables 2 and 3.

Appendix S4: First CASSCF excited states for the ground state geometry as a function of the number of roots.

| 14 roots | 20 roots | |
|---|---|--|
| Ground state energy: -683.673225 a.u. | Ground state energy: -683.670317 a.u. | |
| state 2: 4.81 eV $\pi \rightarrow \pi^*$ 0.71 | state 2: 4.82 eV $\pi \rightarrow \pi^*$ 0.57 | |
| state 3: 6.08 eV $n_{pept1} \rightarrow \pi^*_{pept1}$ 0.57 $n_{pept1} \rightarrow \pi^*_{pept2}$ 0.12 | state 3: 5.66 eV $n_{pept2} \rightarrow \pi^*_{pept2}$ 0.52 $n_{pept2} \rightarrow \pi^*$ 0.14 | |
| state 4: 6.11 eV $n_{pept2} \rightarrow \pi^*_{pept2}$ 0.40 $\pi_{pept1} \rightarrow \pi^*_{pept2}$ 0.11 $n_{pept2} \rightarrow \pi^*_{pept1}$ 0.18 | state 4: 6.14 eV $n_{pept1} \rightarrow \pi^*_{pept1}$ 0.36 $\pi \rightarrow \pi^*_{pept1}$ 0.24 | |
| state 5: 7.87 eV $\pi \rightarrow \pi^*$ 0.77 | state 5: 7.91 eV $\pi \rightarrow \pi^*$ 0.72 | |
| state 6: 8.00 eV $\pi \rightarrow \pi^*$ 0.42 $\pi \pi \rightarrow \pi^* \pi^*$ 0.26 | state 6: 8.00 eV $\pi \rightarrow \pi^*$ 0.31 $\pi \pi \rightarrow \pi^* \pi^*$ 0.34 | |

Table S4: CASSCF total and excitation energy of the NAPA B conformer at its DFT-D optimized ground state geometry (Figure 1(a) and Table S1-1) and weights of the different excitations in the wave function as a function of the number of roots.

Appendix S5: Characteristic structural parameters of the different studied NAPA B geometries.

| Geometry | Ac-Phe-NH2 | | NAPA B | | | |
|-----------------------------|------------|--------|--------|------------|---------------|------------------|
| | Φ | Ψ | χ | ω_2 | $d_{NH(2)-O}$ | $d_{NH(1)-C\pi}$ |
| Ground state | -83 | 55 | 44 | -172 | 2.02 | 2.57 |
| $M_{\pi\pi^*}$ | -83 | 53 | 45 | -167 | 1.95 | 2.35 |
| $CI_{\pi\pi^*/n\pi^*co(2)}$ | -82 | 52 | 49 | -143 | 2.00 | 2.42 |
| $M_{n\pi^*co(2)}$ | -80 | 49 | 60 | -90 | 2.28 | 2.66 |
| M'_{CT} | -85 | 44 | 45 | -159 | 1.87 | 1.35 |

Table S5-1: Characteristic structural parameters of the ground state, $M_{\pi\pi^*}$, $M_{n\pi^*co(2)}$, $CI_{\pi\pi^*/n\pi^*co(2)}$ and M'_{CT} geometries. Ramachandran Φ , Ψ , side chain orientation χ and second peptide bond flatness ω_2 angles are in degrees and bond length in Å.

| Geometry | Ac-Phe-NH2 | | | NAPA B | | | | $\Sigma\theta_{Ci}$ | d_{CO} |
|-----------------------------|---------------|---------------|---------------|---------------------|---------------|---------------|---------------|---------------------|-------------|
| | θ_{N1} | θ_{N2} | θ_{N3} | $\Sigma\theta_{Ni}$ | θ_{C1} | θ_{C2} | θ_{C3} | | |
| Ground state | 118 | 119 | 121 | 358 | 124 | 113 | 123 | 360 | 1.23 |
| $M_{\pi\pi^*}$ | 117 | 119 | 121 | 357 | 124 | 113 | 123 | 360 | 1.24 |
| $CI_{\pi\pi^*/n\pi^*co(2)}$ | 115 | 117 | 121 | 353 | 122 | 114 | 122 | 358 | 1.28 |
| $M_{n\pi^*co(2)}$ | 110 | 111 | 111 | 332 | 105 | 117 | 118 | 340 | 1.38 |
| M'_{CT} | 116 | 117 | 119 | 352 | 128 | 113 | 120 | 361 | 1.22 |

Table S5-2: Characteristic geometrical parameters of the second peptide bond deformation. θ_{Ni} and θ_{Ci} ($i=1-3$) angles (see Figure S5-3 for the definition) are in degrees and bond length in Å.




Figure S5-3: Definition of the angles characteristic of the pyramidalization of the NH₂ group and the COCH group.