



HAL
open science

A LOGARITHMICALLY COMPLETELY MONOTONIC FUNCTION INVOLVING THE q -GAMMA FUNCTION

Feng Qi

► **To cite this version:**

Feng Qi. A LOGARITHMICALLY COMPLETELY MONOTONIC FUNCTION INVOLVING THE q -GAMMA FUNCTION: A monotonic function involving q -gamma function. 2018. hal-01803352

HAL Id: hal-01803352

<https://hal.science/hal-01803352>

Preprint submitted on 30 May 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A LOGARITHMICALLY COMPLETELY MONOTONIC FUNCTION INVOLVING THE q -GAMMA FUNCTION

FENG QI

ABSTRACT. In the paper, the author proves the logarithmically complete monotonicity of a function involving the q -gamma function and derives some inequalities for the q -multinomial coefficient and the q -multivariate beta function. These conclusions generalize corresponding ones for the gamma function, the multinomial coefficient, and the multivariate beta function respectively.

CONTENTS

1.	Preliminaries	1
2.	Motivations	3
3.	A logarithmically completely monotonic function	4
4.	Inequalities for the q -multinomial coefficient	5
5.	Inequalities for the q -multivariate beta function	7
	Acknowledgements	8
	References	8

1. PRELIMINARIES

Suppose that $f : I \rightarrow \infty$ is an infinite differentiable function on an interval I . If

$$0 \leq (-1)^{n-1} f^{(n-1)}(x) < \infty, \quad n \in \mathbb{N},$$

then we call f a completely monotonic function; if $f : I \rightarrow (0, \infty)$ is positive and

$$(-1)^n [\ln f(x)]^{(n)} \geq 0, \quad n \in \mathbb{N},$$

then we call f a logarithmically completely monotonic function; if $f : (0, \infty) \rightarrow [0, \infty)$ can be written in the form

$$f(x) = \frac{a}{x} + b + \int_0^\infty \frac{1}{s+x} d\mu(s),$$

where a, b are non-negative constants and μ is a measure on $(0, \infty)$ such that $\int_0^\infty \frac{1}{1+s} d\mu(s) < \infty$, then we call f a Stieltjes transform. For more details on definitions and properties of these kinds of functions, please see [15, Chapter XIII], [32, Chapter 1], and [33, Chapter IV]. Among these three kinds of functions, there are the following relations:

2010 *Mathematics Subject Classification*. Primary 44A10; Secondary 05A20, 26A48, 26D07, 33B15.

Key words and phrases. logarithmically completely monotonic function; q -gamma function; inequality; q -multinomial coefficient; q -multivariate beta function.

This paper was typeset using $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$.

- (1) a function f is completely monotonic on $(0, \infty)$ if and only if it is a Laplace transform, that is, there is a positive measure μ on $[0, \infty)$ such that

$$f(x) = \int_0^\infty e^{-xt} d\mu(t);$$

- (2) the set of all logarithmically completely monotonic functions is a strict subset of all completely monotonic functions;
(3) the set of all Stieltjes transforms is a strict subset of all logarithmically completely monotonic functions on $(0, \infty)$.

For details on these relations, please refer to [5, 8, 24, 25, 32] and closely related references therein. For information on new developments of these kinds of functions, please refer to [7, 9, 23, 26, 30, 31, 32] and closely related references therein.

The classical Euler's gamma function $\Gamma(z)$ can be defined [22] by

$$\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt, \quad \Re(z) > 0$$

or by

$$\Gamma(z) = \lim_{n \rightarrow \infty} \frac{n! n^z}{\prod_{k=0}^n (z+k)}, \quad z \in \mathbb{C} \setminus \{0, -1, -2, \dots\}.$$

The logarithmic derivative of $\Gamma(z)$, denoted by $\psi(z) = \frac{\Gamma'(z)}{\Gamma(z)}$, and the derivatives $\psi^{(i)}(x)$ for $i \in \mathbb{N}$ are respectively called the digamma (or psi) function and polygamma functions. See [1, Chapter 6] and [16, Chapter 5].

The q -analogue $\Gamma_q(x)$ of the gamma function $\Gamma(x)$ for $q > 0$ and $x > 0$, called the q -gamma function, can be defined [3, pp. 493–496] and [6, Section 1.10] by

$$\Gamma_q(x) = \begin{cases} (1-q)^{1-x} \prod_{i=0}^{\infty} \frac{1-q^{i+1}}{1-q^{i+x}}, & 0 < q < 1; \\ (q-1)^{1-x} q^{\binom{x}{2}} \prod_{i=0}^{\infty} \frac{1-q^{-(i+1)}}{1-q^{-(i+x)}}, & q > 1; \\ \Gamma(x), & q = 1. \end{cases} \quad (1.1)$$

They satisfy

$$\lim_{q \rightarrow 1^+} \Gamma_q(z) = \lim_{q \rightarrow 1^-} \Gamma_q(z) = \Gamma(z) \quad \text{and} \quad \Gamma_q(x) = q^{\binom{x-1}{2}} \Gamma_{1/q}(x).$$

The q -analogue $\psi_q(x)$ of the digamma function $\psi(x)$ for $q > 0$ and $x > 0$, called the q -digamma function, is defined by

$$\psi_q(x) = \begin{cases} \frac{\Gamma'_q(x)}{\Gamma_q(x)}; & q \neq 1, \\ \psi(x), & q = 1. \end{cases}$$

The q -analogues $\psi_q^{(k)}(x)$ of the polygamma functions $\psi^{(k)}(x)$ for $k \in \mathbb{N}$ are called the q -polygamma functions. From (1.1), we obtain that,

- (1) when $0 < q < 1$ and $x \in (0, \infty)$,

$$\psi_q(x) = -\ln(1-q) + (\ln q) \sum_{k=0}^{\infty} \frac{q^{k+x}}{1-q^{k+x}} = -\ln(1-q) + (\ln q) \sum_{k=1}^{\infty} \frac{q^{kx}}{1-q^k},$$

- (2) when $q > 1$ and $x \in (0, \infty)$,

$$\psi_q(x) = -\ln(q-1) + (\ln q) \left(x - \frac{1}{2} - \sum_{i=1}^{\infty} \frac{q^{-ix}}{1-q^{-i}} \right).$$

The formula (1.11) in the paper [11] and its corrected version [12] reads that

$$\psi_q(x) = -\ln(1-q) - \int_0^\infty \frac{e^{-xt}}{1-e^{-t}} d\gamma_q(t) \quad (1.2)$$

for $0 < q < 1$ and $x > 0$, where

$$\gamma_q(t) = \begin{cases} -\ln q \sum_{k=1}^{\infty} \delta(t+k \ln q), & 0 < q < 1 \\ t, & q = 1 \end{cases} \quad (1.3)$$

and $\delta(t)$ represents the Dirac delta function, that is, $d\gamma_q(t)$ is a discrete measure with positive masses $|\ln q|$ at the positive points $k|\ln q|$ for $k \in \mathbb{N}$. Accordingly, we obtain

$$\int_0^\infty e^{-xt} d\gamma_q(t) = -\frac{q^x \ln q}{1-q^x}$$

and

$$\int_0^\infty t e^{-xt} d\gamma_q(t) = \frac{q^x (\ln q)^2}{(1-q^x)^2}$$

for $0 < q < 1$ and $x > 0$. Differentiating with respect to x on both sides of (1.2) yields

$$\psi_q^{(k)}(x) = (-1)^{k+1} \int_0^\infty \frac{t^k e^{-xt}}{1-e^{-t}} d\gamma_q(t), \quad 0 < q < 1, \quad k \in \mathbb{N}. \quad (1.4)$$

In [10, p. 1245, Theorem 4.4, (4.15)], [18, Lemma 2.3 and Remark 2.1], [19, Theorem 7.2, (7.5)], and [28, p. 152, Theorem 4.22, (4.20)], it was presented that, when $0 < q < 1$, the identity

$$\psi_q^{(k-1)}(x) - \psi_q^{(k-1)}(x+1) = (\ln q) \frac{d^{k-1}}{dx^{k-1}} \left(\frac{q^x}{1-q^x} \right)$$

is valid for $x \in (0, \infty)$ and $k \in \mathbb{N}$. One can also find these knowledge in [18, 19, 20, 21, 27] and closely related references.

2. MOTIVATIONS

Let $\mathbf{a} = (a_1, a_2, \dots, a_m)$ with $a_i > 0$ for $1 \leq i \leq m$, let $\mathbf{p} = (p_1, p_2, \dots, p_m)$ with $\sum_{i=1}^m p_i = 1$ and $p_i \in (0, 1)$ for $1 \leq i \leq m$, let

$$\binom{\sum_{i=1}^m a_i}{a_1, a_2, \dots, a_m} = \frac{\Gamma(1 + \sum_{i=1}^m a_i)}{\Gamma(1 + a_1) \Gamma(1 + a_2) \cdots \Gamma(1 + a_m)}$$

denote the multinomial coefficient, and let

$$B(a_1, a_2, \dots, a_m) = \frac{\Gamma(a_1) \Gamma(a_2) \cdots \Gamma(a_m)}{\Gamma(a_1 + a_2 + \cdots + a_m)}$$

be the multivariate beta function. In [17] and [29], the authors considered the function

$$\begin{aligned} \mathcal{Q}(x) &= \mathcal{Q}_{\mathbf{a}, \mathbf{p}; m}(x) = \frac{\Gamma(1 + x \sum_{i=1}^m a_i)}{\prod_{i=1}^m \Gamma(1 + x a_i)} \prod_{i=1}^m p_i^{x a_i} = \binom{x \sum_{i=1}^m a_i}{x a_1, x a_2, \dots, x a_m} \prod_{i=1}^m p_i^{x a_i} \\ &= \frac{\sum_{i=1}^m a_i}{\prod_{i=1}^m a_i} \frac{\prod_{i=1}^m p_i^{x a_i}}{x^{m-1} B(x a_1, x a_2, \dots, x a_m)}, \quad x \in (0, \infty), \quad m \in \mathbb{N} \end{aligned} \quad (2.1)$$

and, among other things, proved the following theorem.

Theorem 2.1 ([17, Theorem 2.1] and [29, Theorem 2.2]). *The function $\mathcal{Q}(x) = \mathcal{Q}_{\mathbf{a}, \mathbf{p}; m}(x)$ defined in (2.1) is logarithmically completely monotonic on $(0, \infty)$.*

For more information on the backgrounds, history, motivations, origins, and applications of the function $\mathcal{Q}(x) = \mathcal{Q}_{\mathbf{a}, \mathbf{p}; m}(x)$, please refer to [2, 13, 17, 29] and closely related references therein.

In this paper, we consider the q -analogue of the function $\mathcal{Q}(x) = \mathcal{Q}_{\mathbf{a}, \mathbf{p}; m}(x)$ and investigate its logarithmically complete monotonicity.

3. A LOGARITHMICALLY COMPLETELY MONOTONIC FUNCTION

Let $\mathbf{a} = (a_1, a_2, \dots, a_m)$ with $a_i > 0$ for $1 \leq i \leq m$ and let $\mathbf{p} = (p_1, p_2, \dots, p_m)$ with $\sum_{i=1}^m p_i = 1$ and $p_i \in (0, 1)$ for $1 \leq i \leq m$. Define

$$\mathcal{Q}_q(x) = \mathcal{Q}_{q; \mathbf{a}, \mathbf{p}; m}(x) = \frac{\Gamma_q(1 + x \sum_{i=1}^m a_i)}{\prod_{i=1}^m \Gamma_q(1 + x a_i)} \prod_{i=1}^m p_i^{x a_i} \quad (3.1)$$

for $q \in (0, 1)$, $x \in (0, \infty)$, and $m \in \mathbb{N}$. It is clear that

$$\lim_{q \rightarrow 1^-} \mathcal{Q}_q(x) = \lim_{q \rightarrow 1^-} \mathcal{Q}_{q; \mathbf{a}, \mathbf{p}; m}(x) = \mathcal{Q}(x) = \mathcal{Q}_{\mathbf{a}, \mathbf{p}; m}(x).$$

Theorem 3.1. *The function $\mathcal{Q}_q(x) = \mathcal{Q}_{q; \mathbf{a}, \mathbf{p}; m}(x)$ defined in (3.1) is logarithmically completely monotonic on $(0, \infty)$.*

Proof. From (1.3) for $0 < q < 1$ and the well-known property $\delta(at) = \frac{1}{|a|} \delta(t)$, it follows that

$$\gamma_q\left(\frac{t}{\tau}\right) = -\ln q \sum_{k=1}^{\infty} \delta\left(\frac{t + k\tau \ln q}{\tau}\right) = -\tau \ln q \sum_{k=1}^{\infty} \delta(t + k\tau \ln q)$$

for $\tau > 0$. This implies that

$$d\gamma_q\left(\frac{t}{\tau}\right) = \tau d\gamma_q(t). \quad (3.2)$$

Direct calculation gives

$$\begin{aligned} \ln \mathcal{Q}_q(x) &= \ln \Gamma_q\left(1 + x \sum_{i=1}^m a_i\right) - \sum_{i=1}^m \ln \Gamma_q(1 + a_i x) + x \sum_{i=1}^m a_i \ln p_i, \\ [\ln \mathcal{Q}_q(x)]' &= \left(\sum_{i=1}^m a_i\right) \psi_q\left(1 + x \sum_{i=1}^m a_i\right) - \sum_{i=1}^m a_i \psi_q(1 + a_i x) + \sum_{i=1}^m a_i \ln p_i, \end{aligned}$$

and

$$[\ln \mathcal{Q}_q(x)]'' = \left(\sum_{i=1}^m a_i\right)^2 \psi_q'\left(1 + x \sum_{i=1}^m a_i\right) - \sum_{i=1}^m a_i^2 \psi_q'(1 + a_i x).$$

Making use of (1.4) for $k = 1$ and the equality (3.2), it follows that

$$\begin{aligned} \psi_q'(\tau x + 1) &= \int_0^{\infty} \frac{t}{1 - e^{-t}} e^{-(\tau x + 1)t} d\gamma_q(t) = \int_0^{\infty} \frac{t}{e^t - 1} e^{-\tau x t} d\gamma_q(t) \\ &= \int_0^{\infty} h\left(\frac{v}{\tau}\right) e^{-v x} d\gamma_q\left(\frac{v}{\tau}\right) = \tau \int_0^{\infty} h\left(\frac{v}{\tau}\right) e^{-v x} d\gamma_q(v), \end{aligned}$$

where $\tau > 0$ and $h(t) = \frac{t}{e^t - 1}$. Hence, we have

$$[\ln \mathcal{Q}_q(x)]'' = \int_0^{\infty} \left[\left(\sum_{i=1}^m a_i\right)^3 h\left(\frac{v}{\sum_{i=1}^m a_i}\right) - \sum_{i=1}^m a_i^3 h\left(\frac{v}{a_i}\right) \right] e^{-x v} d\gamma_q(v). \quad (3.3)$$

By calculus, we have

$$\frac{d}{dt} \left[t^3 h\left(\frac{1}{t}\right) \right] = \frac{d}{dt} \left(\frac{t^2}{e^{1/t} - 1} \right) = \frac{e^{1/t}(2t + 1) - 2t}{(e^{1/t} - 1)^2},$$

$$\begin{aligned} \frac{d^2}{dt^2} \left[t^3 h\left(\frac{1}{t}\right) \right] &= \frac{2t^2 - e^{1/t}(4t^2 + 2t - 1) + e^{2/t}(2t^2 + 2t + 1)}{(e^{1/t} - 1)^3 t^2} \rightarrow 0, \quad t \rightarrow 0^+, \\ \frac{d^3}{dt^3} \left[t^3 h\left(\frac{1}{t}\right) \right] &= \frac{e^{1/t}(4e^{1/t} + e^{2/t} + 1)}{(e^{1/t} - 1)^4 t^4} > 0 \end{aligned}$$

on $(0, \infty)$. As a result, the function $t^3 h\left(\frac{1}{t}\right)$ is strictly convex on $(0, \infty)$. Recall from [14, p. 650] that

- (1) a function $\phi : [0, \infty) \rightarrow \mathbb{R}$ is said to be star-shaped if $\phi(\alpha x) \leq \alpha\phi(x)$ for all $\alpha \in [0, 1]$ and $x \geq 0$;
- (2) a real function ϕ defined on a set $S \subset \mathbb{R}^n$ is said to be super-additive if $x, y \in S$ implies $x + y \in S$ and $\phi(x + y) \geq \phi(x) + \phi(y)$;
- (3) if ϕ is a real function defined on $[0, \infty)$, $\phi(0) \leq 0$, and ϕ is convex, then ϕ is star-shaped, but convexity is not a property of all star-shaped functions;
- (4) if $\phi : [0, \infty) \rightarrow \mathbb{R}$ is star-shaped, then ϕ is super-additive.

Consequently, since $\lim_{t \rightarrow 0^+} \frac{t^2}{e^{1/t} - 1} = 0$, the function $t^3 h\left(\frac{1}{t}\right)$ is star-shaped, and then super-additive, on $(0, \infty)$. As a result, it follows inductively that

$$\left(\sum_{i=1}^m \frac{a_i}{v} \right)^3 h\left(\frac{1}{\sum_{i=1}^m (a_i/v)} \right) \geq \sum_{i=1}^m \left(\frac{a_i}{v} \right)^3 h\left(\frac{1}{a_i/v} \right)$$

which can be rearranged as

$$\left(\sum_{i=1}^m a_i \right)^3 h\left(\frac{v}{\sum_{i=1}^m a_i} \right) \geq \sum_{i=1}^m a_i^3 h\left(\frac{v}{a_i} \right).$$

Combining this with (3.3) yields that the second derivative $[\ln \mathcal{Q}_q(x)]''$ is completely monotonic on $(0, \infty)$.

Complete monotonicity of $[\ln \mathcal{Q}_q(x)]''$ implies that the first derivative $[\ln \mathcal{Q}_q(x)]'$ is strictly increasing on $(0, \infty)$. Hence, from (1.2), it follows that

$$\begin{aligned} [\ln \mathcal{Q}_q(x)]' &\leq \lim_{x \rightarrow \infty} \left[\left(\sum_{i=1}^m a_i \right) \psi_q \left(1 + x \sum_{i=1}^m a_i \right) - \sum_{i=1}^m a_i \psi_q(1 + a_i x) \right] + \sum_{i=1}^m a_i \ln p_i \\ &= \left[\left(\sum_{i=1}^m a_i \right) \lim_{x \rightarrow \infty} \psi_q \left(1 + x \sum_{i=1}^m a_i \right) - \sum_{i=1}^m a_i \lim_{x \rightarrow \infty} \psi_q(1 + a_i x) \right] + \sum_{i=1}^m a_i \ln p_i \\ &= \left[- \left(\sum_{i=1}^m a_i \right) \ln(1 - q) + \sum_{i=1}^m a_i \ln(1 - q) \right] + \sum_{i=1}^m a_i \ln p_i = \sum_{i=1}^m a_i \ln p_i < 0. \end{aligned}$$

By definition, the function $\mathcal{Q}_q(x)$ is logarithmically completely monotonic on $(0, \infty)$. The proof of Theorem 3.1 is complete. \square

4. INEQUALITIES FOR THE q -MULTINOMIAL COEFFICIENT

Let $\mathbf{a} = (a_1, a_2, \dots, a_m)$ with $a_i > 0$ for $1 \leq i \leq m$. Then the q -multinomial coefficient for $q \in (0, 1)$ is

$$\binom{\sum_{i=1}^m a_i}{a_1, a_2, \dots, a_m}_q = \frac{\Gamma_q(1 + \sum_{i=1}^m a_i)}{\Gamma_q(1 + a_1) \Gamma_q(1 + a_2) \dots \Gamma_q(1 + a_m)}. \quad (4.1)$$

With the aid of logarithmically complete monotonicity of $\mathcal{Q}_q(x) = \mathcal{Q}_{q;\mathbf{a},\mathbf{p};m}(x)$, we now present some inequalities for the q -multinomial coefficient.

Theorem 4.1. *Let $q \in (0, 1)$, $\ell, m \in \mathbb{N}$, and $\mathbf{a} = (a_1, a_2, \dots, a_m)$ with $a_i > 0$ for $1 \leq i \leq m$. If $x_j > 0$ for $1 \leq j \leq \ell$ and $\lambda_j \in (0, 1)$ with $\sum_{j=1}^{\ell} \lambda_j = 1$, then*

$$\begin{aligned} & \left(a_1 \sum_{j=1}^{\ell} \lambda_j x_j, a_2 \sum_{j=1}^{\ell} \lambda_j x_j, \dots, a_m \sum_{j=1}^{\ell} \lambda_j x_j \right)_q \\ & \leq \prod_{j=1}^{\ell} \left(x_j \sum_{i=1}^m a_i, x_j a_1, x_j a_2, \dots, x_j a_m \right)_q^{\lambda_j} \end{aligned} \quad (4.2)$$

and the equality in (4.2) holds if and only if $x_1 = x_2 = \dots = x_{\ell}$.

Proof. The logarithmically complete monotonicity in Theorem 3.1 indicates that the function $\mathcal{Q}_q(x)$ is logarithmically convex on $(0, \infty)$. Hence, one acquires

$$\mathcal{Q}_q \left(\sum_{j=1}^{\ell} \lambda_j x_j \right) \leq \prod_{j=1}^{\ell} \mathcal{Q}_q^{\lambda_j}(x_j).$$

Using the expression

$$\mathcal{Q}_q(x) = \left(x \sum_{i=1}^m a_i, x a_1, x a_2, \dots, x a_m \right)_q \prod_{i=1}^m p_i^{x a_i}$$

leads to

$$\begin{aligned} & \left(a_1 \sum_{j=1}^{\ell} \lambda_j x_j, a_2 \sum_{j=1}^{\ell} \lambda_j x_j, \dots, a_m \sum_{j=1}^{\ell} \lambda_j x_j \right)_q \prod_{i=1}^m p_i^{a_i \sum_{j=1}^{\ell} \lambda_j x_j} \\ & \leq \prod_{j=1}^{\ell} \left[\left(x_j \sum_{i=1}^m a_i, x_j a_1, x_j a_2, \dots, x_j a_m \right)_q \prod_{i=1}^m p_i^{a_i x_j} \right]^{\lambda_j} \end{aligned}$$

which can be rearranged as (4.2). The proof of Theorem 4.1 is complete. \square

Theorem 4.2. Let $q \in (0, 1)$, $\ell, m \in \mathbb{N}$, and $\mathbf{a} = (a_1, a_2, \dots, a_m)$ with $a_i > 0$ for $1 \leq i \leq m$. If $x_j > 0$ for $1 \leq j \leq \ell$, then

$$\begin{aligned} & \prod_{j=1}^{\ell} \left(x_j \sum_{i=1}^m a_i, x_j a_1, x_j a_2, \dots, x_j a_m \right)_q \\ & < \left(\sum_{j=1}^{\ell} x_j \sum_{i=1}^m a_i, a_1 \sum_{j=1}^{\ell} x_j, a_2 \sum_{j=1}^{\ell} x_j, \dots, a_m \sum_{j=1}^{\ell} x_j \right)_q. \end{aligned} \quad (4.3)$$

Proof. In [2, Lemma 3], it was proved that, if $g : [0, \infty) \rightarrow (0, 1]$ is differentiable and $\frac{g'}{g}$ is strictly increasing on $(0, \infty)$, then the inequality $g(x)g(y) < g(x+y)$ is valid for $x, y \in (0, \infty)$. As a result, we can inductively deduce

$$\prod_{j=1}^{\ell} g(x_j) < g \left(\sum_{j=1}^{\ell} x_j \right).$$

Applying this inequality to the logarithmic convexity of the function $\mathcal{Q}_q(x)$ reveals

$$\begin{aligned} & \prod_{j=1}^{\ell} \left[\left(x_j \sum_{i=1}^m a_i, x_j a_1, x_j a_2, \dots, x_j a_m \right)_q \prod_{i=1}^m p_i^{a_i x_j} \right] \\ & < \left(\sum_{j=1}^{\ell} x_j \sum_{i=1}^m a_i, a_1 \sum_{j=1}^{\ell} x_j, a_2 \sum_{j=1}^{\ell} x_j, \dots, a_m \sum_{j=1}^{\ell} x_j \right)_q \prod_{i=1}^m p_i^{a_i \sum_{j=1}^{\ell} x_j} \end{aligned}$$

which can be rewritten as (4.3). The proof of Theorem 4.2 is complete. \square

Theorem 4.3. Let $q \in (0, 1)$, $m \in \mathbb{N}$, and $\mathbf{a} = (a_1, a_2, \dots, a_m)$ with $a_i > 0$ for $1 \leq i \leq m$. If $0 < a \leq c$ and $x > 0$, then

$$\begin{aligned} & \left(\begin{array}{c} (a+x) \sum_{i=1}^m a_i \\ (a+x)a_1, (a+x)a_2, \dots, (a+x)a_m \end{array} \right)_q \left(\begin{array}{c} c \sum_{i=1}^m a_i \\ ca_1, ca_2, \dots, ca_m \end{array} \right)_q \\ & \leq \left(\begin{array}{c} a \sum_{i=1}^m a_i \\ aa_1, aa_2, \dots, aa_m \end{array} \right)_q \left(\begin{array}{c} (c+x) \sum_{i=1}^m a_i \\ (c+x)a_1, (c+x)a_2, \dots, (c+x)a_m \end{array} \right)_q \end{aligned} \quad (4.4)$$

and the equality in (4.4) holds if and only if $a = c$.

Proof. For $0 < a < c$, set

$$V(x) = \ln \mathcal{Q}_q(a+x) + \ln \mathcal{Q}_q(c) - \ln \mathcal{Q}_q(a) - \ln \mathcal{Q}_q(c+x).$$

Because

$$V'(x) = \frac{\mathcal{Q}'_q(a+x)}{\mathcal{Q}_q(a+x)} - \frac{\mathcal{Q}'_q(c+x)}{\mathcal{Q}_q(c+x)}$$

and the logarithmically complete monotonicity of $\mathcal{Q}_q(x)$ hints that the function $\frac{\mathcal{Q}'_q(x)}{\mathcal{Q}_q(x)}$ is strictly increasing on $(0, \infty)$, we see that $V'(x) < 0$ and $V(x) < V(0) = 0$. Consequently, it follows that

$$\ln \mathcal{Q}_q(a+x) + \ln \mathcal{Q}_q(c) \leq \ln \mathcal{Q}_q(a) + \ln \mathcal{Q}_q(c+x)$$

which is equivalent to

$$\begin{aligned} & \ln \left[\left(\begin{array}{c} (a+x) \sum_{i=1}^m a_i \\ (a+x)a_1, (a+x)a_2, \dots, (a+x)a_m \end{array} \right)_q \prod_{i=1}^m p_i^{a_i(a+x)} \right] \\ & + \ln \left[\left(\begin{array}{c} c \sum_{i=1}^m a_i \\ ca_1, ca_2, \dots, ca_m \end{array} \right)_q \prod_{i=1}^m p_i^{a_i c} \right] \leq \ln \left[\left(\begin{array}{c} a \sum_{i=1}^m a_i \\ aa_1, aa_2, \dots, aa_m \end{array} \right)_q \prod_{i=1}^m p_i^{a_i a} \right] \\ & + \ln \left[\left(\begin{array}{c} (c+x) \sum_{i=1}^m a_i \\ (c+x)a_1, (c+x)a_2, \dots, (c+x)a_m \end{array} \right)_q \prod_{i=1}^m p_i^{a_i(c+x)} \right]. \end{aligned}$$

This can be simplified as (4.4). The proof of Theorem 4.3 is complete. \square

5. INEQUALITIES FOR THE q -MULTIVARIATE BETA FUNCTION

Let $\mathbf{a} = (a_1, a_2, \dots, a_m)$ with $a_i > 0$ for $1 \leq i \leq m$. Then the q -multivariate beta function for $q \in (0, 1)$ is

$$B_q(a_1, a_2, \dots, a_m) = \frac{\Gamma_q(a_1)\Gamma_q(a_2)\cdots\Gamma_q(a_m)}{\Gamma_q(a_1 + a_2 + \cdots + a_m)}.$$

In [3, p. 544], it was proved that

$$\Gamma_q(x+1) = \frac{1-q^x}{1-q} \Gamma_q(x)$$

with $\Gamma_q(1) = 1$ for $q \in (0, 1)$. Applying this recurrence relation to (4.1) yields

$$\left(\begin{array}{c} \sum_{i=1}^m a_i \\ a_1, a_2, \dots, a_m \end{array} \right)_q = \frac{(1-q)^{m-1} (1-q^{\sum_{i=1}^m a_i})}{\prod_{i=1}^m (1-q^{a_i})} \frac{1}{B_q(a_1, a_2, \dots, a_m)}. \quad (5.1)$$

Substituting (5.1) into those inequalities in Theorems 4.1 to 4.3, we can conclude several inequalities for the q -multivariate beta function $B_q(a_1, a_2, \dots, a_m)$. Equivalently speaking, Theorems (4.1) to (4.3) can be respectively reformulated as follows.

Let $q \in (0, 1)$, $\ell, m \in \mathbb{N}$, and $\mathbf{a} = (a_1, a_2, \dots, a_m)$ with $a_i > 0$ for $1 \leq i \leq m$. If $x_j > 0$ for $1 \leq j \leq \ell$ and $\lambda_j \in (0, 1)$ with $\sum_{j=1}^{\ell} \lambda_j = 1$, then

$$\begin{aligned} & \frac{\prod_{i=1}^m (1 - q^{a_i \sum_{j=1}^{\ell} \lambda_j x_j})}{1 - q^{\sum_{j=1}^{\ell} \lambda_j x_j \sum_{i=1}^m a_i}} B_q \left(a_1 \sum_{j=1}^{\ell} \lambda_j x_j, a_2 \sum_{j=1}^{\ell} \lambda_j x_j, \dots, a_m \sum_{j=1}^{\ell} \lambda_j x_j \right) \\ & \geq \prod_{j=1}^{\ell} \left[\frac{\prod_{i=1}^m (1 - q^{x_j a_i})}{1 - q^{x_j \sum_{i=1}^m a_i}} B_q(a_1 x_j, a_2 x_j, \dots, a_m x_j) \right]^{\lambda_j} \quad (5.2) \end{aligned}$$

and the equality in (5.2) holds if and only if $x_1 = x_2 = \dots = x_{\ell}$.

The inequality (5.2) implies that the function

$$\frac{\prod_{i=1}^m (1 - q^{a_i x})}{1 - q^{x \sum_{i=1}^m a_i}} B_q(a_1 x, a_2 x, \dots, a_m x)$$

for $m \in \mathbb{N}$ is logarithmically concave with respect to $x \in (0, \infty)$. Generally, we claim that the function

$$\frac{1 - q^{x \sum_{i=1}^m a_i}}{\prod_{i=1}^m (1 - q^{a_i x})} B_q(a_1 x, a_2 x, \dots, a_m x)$$

for $m \in \mathbb{N}$ is logarithmically completely monotonic with respect to $x \in (0, \infty)$.

Let $q \in (0, 1)$, $\ell, m \in \mathbb{N}$, and $\mathbf{a} = (a_1, a_2, \dots, a_m)$ with $a_i > 0$ for $1 \leq i \leq m$. If $x_j > 0$ for $1 \leq j \leq \ell$, then

$$\begin{aligned} & \frac{1}{(1 - q)^{(\ell-1)(m-1)}} \prod_{j=1}^{\ell} \left[\frac{\prod_{i=1}^m (1 - q^{a_i x_j})}{1 - q^{x_j \sum_{i=1}^m a_i}} B_q(a_1 x_j, a_2 x_j, \dots, a_m x_j) \right] \\ & > \frac{\prod_{i=1}^m (1 - q^{a_i \sum_{j=1}^{\ell} x_j})}{1 - q^{\sum_{j=1}^{\ell} x_j \sum_{i=1}^m a_i}} B_q \left(a_1 \sum_{j=1}^{\ell} x_j, a_2 \sum_{j=1}^{\ell} x_j, \dots, a_m \sum_{j=1}^{\ell} x_j \right). \end{aligned}$$

Let $q \in (0, 1)$, $m \in \mathbb{N}$, and $\mathbf{a} = (a_1, a_2, \dots, a_m)$ with $a_i > 0$ for $1 \leq i \leq m$. If $0 < a \leq c$ and $x > 0$, then

$$\begin{aligned} & \frac{1 - q^{(a+x) \sum_{i=1}^m a_i}}{1 - q^{(c+x) \sum_{i=1}^m a_i}} \frac{\prod_{i=1}^m (1 - q^{(c+x) a_i})}{\prod_{i=1}^m (1 - q^{(a+x) a_i})} B_q((c+x)a_1, (c+x)a_2, \dots, (c+x)a_m) \\ & \leq \frac{1 - q^{c \sum_{i=1}^m a_i}}{1 - q^{a \sum_{i=1}^m a_i}} \frac{\prod_{i=1}^m (1 - q^{a a_i})}{\prod_{i=1}^m (1 - q^{c a_i})} B_q(ca_1, ca_2, \dots, ca_m) \quad (5.3) \end{aligned}$$

and the equality in (5.3) holds if and only if $a = c$.

Acknowledgements. The author is thankful to Dr. Frédéric Ouimet (Université de Montréal, Canada) and Dr. Li Yin (Binzhou University, China) for their careful corrections to and valuable comments on the original version of this paper.

REFERENCES

- [1] M. Abramowitz and I. A. Stegun (Eds), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards, Applied Mathematics Series **55**, 10th printing, Dover Publications, New York and Washington, 1972.
- [2] H. Alzer, *Complete monotonicity of a function related to the binomial probability*, J. Math. Anal. Appl. **459** (2018), no. 1, 10–15; Available online at <https://doi.org/10.1016/j.jmaa.2017.10.077>.
- [3] G. E. Andrews, R. Askey, and R. Roy, *Special Functions*, Encyclopedia of Mathematics and its Applications **71**, Cambridge University Press, Cambridge, 1999; Available online at <http://dx.doi.org/10.1017/CB09781107325937>.
- [4] R. D. Atanassov and U. V. Tsoukrovski, *Some properties of a class of logarithmically completely monotonic functions*, C. R. Acad. Bulgare Sci. **41** (1988), no. 2, 21–23.
- [5] C. Berg, *Integral representation of some functions related to the gamma function*, Mediterr. J. Math. **1** (2004), no. 4, 433–439; Available online at <http://dx.doi.org/10.1007/s00009-004-0022-6>.

- [6] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, 2nd ed., Encyclopedia of Mathematics and its Applications **96**, Cambridge University Press, Cambridge, 2004; Available online at <http://dx.doi.org/10.1017/CB09780511526251>.
- [7] B.-N. Guo and F. Qi, *A completely monotonic function involving the tri-gamma function and with degree one*, Appl. Math. Comput. **218** (2012), no. 19, 9890–9897; Available online at <http://dx.doi.org/10.1016/j.amc.2012.03.075>.
- [8] B.-N. Guo and F. Qi, *A property of logarithmically absolutely monotonic functions and the logarithmically complete monotonicity of a power-exponential function*, Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys. **72** (2010), no. 2, 21–30.
- [9] B.-N. Guo and F. Qi, *On the degree of the weighted geometric mean as a complete Bernstein function*, Afr. Mat. **26** (2015), no. 7, 1253–1262; Available online at <http://dx.doi.org/10.1007/s13370-014-0279-2>.
- [10] B.-N. Guo and F. Qi, *Properties and applications of a function involving exponential functions*, Commun. Pure Appl. Anal. **8** (2009), no. 4, 1231–1249; Available online at <http://dx.doi.org/10.3934/cpaa.2009.8.1231>.
- [11] M. E. H. Ismail and M. E. Muldoon, *Inequalities and monotonicity properties for gamma and q -gamma functions*, in: R.V.M. Zahar (Ed.), Approximation and Computation: A Festschrift in Honour of Walter Gautschi, ISNM, Vol. **119**, BirkhRauser, Basel, 1994, 309–323; Available online at http://dx.doi.org/10.1007/978-1-4684-7415-2_19.
- [12] M. E. H. Ismail and M. E. Muldoon, *Inequalities and monotonicity properties for gamma and q -gamma functions*, arXiv preprint (2013), available online at <http://arxiv.org/abs/1301.1749>.
- [13] A. Leblanc and B. C. Johnson, *On a uniformly integrable family of polynomials defined on the unit interval*, J. Inequal. Pure Appl. Math. **8** (2007), no. 3, Article 67, 5 pp. Available online at <https://www.emis.de/journals/JIPAM/article878.html>.
- [14] A. W. Marshall, I. Olkin, and B. C. Arnold, *Inequalities: Theory of Majorization and its Applications*, 2nd Ed., Springer Verlag, New York-Dordrecht-Heidelberg-London, 2011; Available online at <http://dx.doi.org/10.1007/978-0-387-68276-1>.
- [15] D. S. Mitrinović, J. E. Pečarić, and A. M. Fink, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht-Boston-London, 1993; Available online at <http://dx.doi.org/10.1007/978-94-017-1043-5>.
- [16] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark (eds.), *NIST Handbook of Mathematical Functions*, Cambridge University Press, New York, 2010; Available online at <http://dlmf.nist.gov/>.
- [17] F. Ouimet, *Complete monotonicity of multinomial probabilities and its application to Bernstein estimators on the simplex*, arXiv preprint (2018), available online at <https://arxiv.org/abs/1804.02108>.
- [18] F. Qi, *A completely monotonic function related to the q -trigamma function*, Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys. **76** (2014), no. 1, 107–114.
- [19] F. Qi, *Bounds for the ratio of two gamma functions*, J. Inequal. Appl. **2010** (2010), Article ID 493058, 84 pages; Available online at <http://dx.doi.org/10.1155/2010/493058>.
- [20] F. Qi, *Certain logarithmically N -alternating monotonic functions involving gamma and q -gamma functions*, Nonlinear Funct. Anal. Appl. **12** (2007), no. 4, 675–685.
- [21] F. Qi, *Complete monotonicity of functions involving the q -trigamma and q -tetragamma functions*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM **109** (2015), no. 2, 419–429; Available online at <http://dx.doi.org/10.1007/s13398-014-0193-3>.
- [22] F. Qi, *Limit formulas for ratios between derivatives of the gamma and digamma functions at their singularities*, Filomat **27** (2013), no. 4, 601–604; Available online at <http://dx.doi.org/10.2298/FIL1304601Q>.
- [23] F. Qi, *Properties of modified Bessel functions and completely monotonic degrees of differences between exponential and trigamma functions*, Math. Inequal. Appl. **18** (2015), no. 2, 493–518; Available online at <http://dx.doi.org/10.7153/mia-18-37>.
- [24] F. Qi and C.-P. Chen, *A complete monotonicity property of the gamma function*, J. Math. Anal. Appl. **296** (2004), 603–607; Available online at <http://dx.doi.org/10.1016/j.jmaa.2004.04.026>.
- [25] F. Qi, B.-N. Guo, and C.-P. Chen, *Some completely monotonic functions involving the gamma and polygamma functions*, J. Aust. Math. Soc. **80** (2006), 81–88; Available online at <http://dx.doi.org/10.1017/S1446788700011393>.
- [26] F. Qi and W.-H. Li, *Integral representations and properties of some functions involving the logarithmic function*, Filomat **30** (2016), no. 7, 1659–1674; Available online at <https://doi.org/10.2298/FIL1607659Q>.
- [27] F. Qi, F.-F. Liu, and X.-T. Shi, *Comments on two completely monotonic functions involving the q -trigamma function*, J. Inequal. Spec. Funct. **7** (2016), no. 4, 211–217.

- [28] F. Qi and Q.-M. Luo, *Bounds for the ratio of two gamma functions—From Wendel’s and related inequalities to logarithmically completely monotonic functions*, Banach J. Math. Anal. **6** (2012), no. 2, 132–158; Available online at <http://dx.doi.org/10.15352/bjma/1342210165>.
- [29] F. Qi, D.-W. Niu, D. Lim, and B.-N. Guo, *Some logarithmically completely monotonic functions and inequalities for multinomial coefficients and multivariate beta functions*, HAL archives (2018), available online at <https://hal.archives-ouvertes.fr/hal-01769288>.
- [30] F. Qi and S.-H. Wang, *Complete monotonicity, completely monotonic degree, integral representations, and an inequality related to the exponential, trigamma, and modified Bessel functions*, Glob. J. Math. Anal. **2** (2014), no. 3, 91–97; Available online at <http://dx.doi.org/10.14419/gjma.v2i3.2919>.
- [31] F. Qi, X.-J. Zhang, and W.-H. Li, *Lévy-Khintchine representations of the weighted geometric mean and the logarithmic mean*, Mediterr. J. Math. **11** (2014), no. 2, 315–327; Available online at <http://dx.doi.org/10.1007/s00009-013-0311-z>.
- [32] R. L. Schilling, R. Song, and Z. Vondraček, *Bernstein Functions—Theory and Applications*, 2nd ed., de Gruyter Studies in Mathematics **37**, Walter de Gruyter, Berlin, Germany, 2012; Available online at <http://dx.doi.org/10.1515/9783110269338>.
- [33] D. V. Widder, *The Laplace Transform*, Princeton University Press, Princeton, 1946.

INSTITUTE OF MATHEMATICS, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO, HENAN, 454010, CHINA; COLLEGE OF MATHEMATICS, INNER MONGOLIA UNIVERSITY FOR NATIONALITIES, TONGLIAO, INNER MONGOLIA, 028043, CHINA; DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCE, TIANJIN POLYTECHNIC UNIVERSITY, TIANJIN, 300387, CHINA; INSTITUTE OF FUNDAMENTAL AND FRONTIER SCIENCES, UNIVERSITY OF ELECTRONIC SCIENCE AND TECHNOLOGY OF CHINA, CHENGDU, SICHUAN, 610054, CHINA

Email address: qifeng618@gmail.com, qifeng618@hotmail.com, qifeng618@qq.com

URL: <https://qifeng618.wordpress.com>