



HAL
open science

Impact of Thermodiffusion on the Initial Vertical Distribution of Species in Hydrocarbon Reservoirs

Guillaume Galliero, Henri Bataller, Fabrizio Croccolo, Romain Vermorel, P.-A. Artola, B. Rousseau, V. Vesovic, M. Bou-Ali, J.M. Ortiz De Zárate, S. Xu, et al.

► **To cite this version:**

Guillaume Galliero, Henri Bataller, Fabrizio Croccolo, Romain Vermorel, P.-A. Artola, et al.. Impact of Thermodiffusion on the Initial Vertical Distribution of Species in Hydrocarbon Reservoirs. *Microgravity Science and Technology*, 2016, 28 (2), pp.79-86. 10.1007/s12217-015-9465-6 . hal-01804384

HAL Id: hal-01804384

<https://hal.science/hal-01804384>

Submitted on 6 Feb 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Impact of thermodiffusion on the initial vertical distribution of species in hydrocarbon reservoirs

Guillaume Galliero¹, Henri Bataller¹, Fabrizio Croccolo¹, Romain Vermorel¹, Pierre-Arnaud Artola², Bernard Rousseau², Velisa Vesovic³, Mounir Bou-Ali⁴, José M. Ortiz de Zárate⁵, François Montel⁶

¹*Laboratoire des Fluides Complexes et leurs Réservoirs, UMR-5150 CNRS-TOTAL, Université de Pau et des Pays de l'Adour, Pau, FR.*

²*Laboratoire de Chimie-Physique, UMR 8000 CNRS, Université Paris-Sud, Orsay, FR.*

³*Department of Earth Science and Engineering, Imperial College London, London, UK.*

⁴*MGEP Mondragon Goi Eskola Politeknikoa, Mechanical and Industrial Manufacturing Department, Loramendi 4 Apdo. 23, 20500 Mondragon, SP.*

⁵*Departamento de Física Aplicada I. Universidad Complutense, 28040 Madrid, SP.*

⁷*TOTAL E.P., Pau, FR.*

Abstract

In this work we propose a methodology, based on molecular dynamics simulations, to quantify the influence of segregation and thermodiffusion on the initial state distribution of the fluid species in hydrocarbon reservoirs. This convection-free approach has been applied to a synthetic oil composed of three normal alkanes and to a real acid gas. It has been found that the thermodiffusion effect induced by the geothermal gradient is similar (but opposite in sign) to that due to segregation for both mixtures. In addition, because of the combined effect of thermal expansion and thermodiffusion, it has been observed that the density gradient can be reversed, in the presence of a geothermal gradient. These numerical results emphasize the need of improving our quantification of thermodiffusion in multicomponent mixtures. The SCCO-SJ10 experiments will be a crucial step towards this goal.

Keywords: Thermodiffusion, segregation, multicomponent mixtures, oil and gas, molecular dynamics.

Introduction

Compositional grading has a strong impact on the estimation of the initial (before production) hydrocarbon in place and is important for selecting the best reservoir development scenario. In a closed, convection free reservoir, the compositional variations are mainly determined by gravitational segregation (Høier and Whitson, 2001). However, other phenomena may affect the compositional grading in real reservoirs (Holt et al., 1983; Whitson and Belery, 1994; Montel et al., 2007). Among them is thermodiffusion, or Soret effect, induced by the geothermal gradient (Ghorayeb et al., 2003; Montel et al., 2007), which is a phenomenon that couples heat and mass fluxes (de Groot and Mazur, 1984). The contribution of thermodiffusion is difficult to quantify, mainly due to a lack of experimental data as well as accurate modelling for multi-component mixtures. Although noticeable progresses have been made during the last twenty years, see Assael et al., 2014 and references therein, especially on ternary mixtures both theoretically (Firoozabadi et al., 2000; Kempers, 2001; Galliero et al., 2003) and experimentally (Leahy-Dios et al., 2005; Bou-Ali et al., 2015) more work is necessary. Micro-gravity experiments are one possible way to provide further data on thermodiffusion in multicomponent mixtures (Van Vaerenbergh et al., 2009; Touzet et al., 2011). They are particularly interesting because they provide, by default, experimental conditions free of convection effect.

The need for further advancement in the field motivated the development of a project named Soret Coefficient measurements of Crude Oil in SJ10 (SCCO-SJ10). This project aims at measuring thermodiffusion in multicomponent fluid mixtures of petroleum interest using a microgravity set-up. The project is made possible as a result of a partnership between European Space Agency and China's National Space Science Center (Hu et al., 2014). The micro-gravity experiment is scheduled to fly on China's SJ-10 Shi Jian spacecraft towards the end of 2015 or later (Hu et al., 2014). In addition to the two space agencies, this project

involves academics from France (Université de Pau et des Pays de l'Adour, Université de Paris-Sud), Spain (Mondragon Unibertsitatea, Universidad Complutense), United Kingdom (Imperial College London), China (Chinese Academy of Sciences) and industrials from France (Total) and China (RIPED). The microgravity set-up will consist of six cells designed by Sanchez Technology (Georis et al., 1998) and already employed during the previous SCCO experiments that have flown on Russian Foton M3 missions (Van Vaerenbergh et al., 2009, Touzet et al., 2011). Each cell will contain a different mixture (3 of “academic” interest and 3 of “industrial” interest) under high pressures, up to 40 MPa, at an average temperature of 50°C, chosen to mimic reservoir conditions. The filling, as well as the post-flight analysis, of the cells will be performed by RIPED.

To complement this micro-gravity experiment, various on-ground activities will be carried on.

They will consist of:

- ground experiments (Croccolo et al., 2012; Urteaga et al., 2012; Giraudet et al., 2014; Larrañaga et al., 2015) on thermodiffusion in ternary mixtures that will take advantage of an extension of the theory of non-equilibrium fluctuations in ternary mixtures (Ortiz de Zárate et al., 2014),
- Molecular Dynamics (MD) simulations of thermodiffusion in multicomponent mixtures in bulk (Galliero et al., 2003; Galliero et al., 2009; Artola and Rousseau 2013) and confined situations (Galliero et al., 2006; Hannaoui et al., 2013),
- developing models to take into account thermodiffusion when describing the initial state of a reservoir (Montel et al., 2007; Touzet et al., 2011).

Within this framework, the purpose of the current article is to describe a numerical approach emphasizing the influence of thermodiffusion on the vertical distribution of species at the initial state of hydrocarbon reservoirs. The approach relies on MD simulations and on classical thermodynamic modelling based on an Equation of State (EoS) as described in

Galliero and Montel, 2008. The main focus of this article is on the influence of gravitational and geothermal fields in a convection free configuration. We have neglected the influence of the porous media on segregation and thermodiffusion which is a reasonable assumption at the steady state providing the permeability is not too low (Shapiro and Stenby, 2000; Platten and Costeseque, 2004; Montel et al., 2007; Hannoui et al., 2013). To apply this approach, two mixtures have been selected, a synthetic one composed of methane, n-butane and n-dodecane (C_1 - nC_4 - nC_{12}), and one representing a realistic acid gas, rich in carbon dioxide and lean in hydrogen sulphide.

Theory and Methods

Thermo-gravitation

A gravitational field applied to a convection free fluid mixture leads to a partial separation of the species along the vertical axis (de Groot and Mazur 1984). This phenomenon is usually named gravitational segregation and its dynamics is controlled by mass diffusion (Galliero and Montel, 2008). At equilibrium, the isothermal vertical distribution (assuming a one dimensional reservoir) of the species i can be deduced from (Sage and Lacey, 1939):

$$\left(\frac{d\mu_i}{dz} \right)_T = M_i g \quad (1)$$

where μ_i and M_i are the chemical potential and the molecular weight of species i , respectively, z is the vertical position, T the temperature and g the gravitational acceleration. This equation, given a composition at a reference point z^0 , can be used to obtain the variation of composition at the stationary state as a function of depth. However, except for ideal mixtures, such as isotope ones (Galliero and Montel, 2008), eq. (1) cannot be solved analytically and a thermodynamic modeling combined with a numerical procedure has to be employed (Montel and Gouel, 1985; Halldórsson and Stenby, 2000).

When a vertical geothermal gradient is added (the fluid column is heated from below), at the stationary state the distribution of the species i can be obtained from an extension of the Gibbs equation (Galliero and Montel, 2008):

$$\left(\frac{d\mu_i}{dz}\right)_T = M_i g - \frac{\alpha_{T_i}}{T} \frac{dT}{dz} \quad (2)$$

where α_{T_i} is the “generalized” thermal diffusion factor of component i (Kempers, 2001). Such an equation also cannot be solved analytically, except when dealing with ideal mixtures.

Molecular Dynamics Simulations

The classical molecular dynamics method consists in numerically estimating the positions, velocities and acceleration of a number of selected molecules evolving in space and time. The motions of the molecules are described through a numerical integration of the classical Newton’s law equations while the interactions between molecules are described by effective potentials (Allen and Tildesley, 1987). This method can be used to study both equilibrium and transport properties of model fluids, mimicking real ones.

Fluid models

To model the species present in the selected alkane mixture (C_1 , nC_4 , nC_{12}), the simple Lennard-Jones (LJ) chain model has been employed (Galliero, 2014). The molecules are represented by N freely jointed tangent spheres and non-bonded interaction between spheres i and j is described by the usual Lennard-Jones 12-6 potential:

$$U_{ij}^{LJ} = 4\epsilon_{ij} \left[\left(\frac{\sigma_{ij}}{r_{ij}}\right)^{12} - \left(\frac{\sigma_{ij}}{r_{ij}}\right)^6 \right] \quad (3)$$

where σ_{ij} is the distance at which the potential is equal to zero (the “sphere diameter”), ϵ_{ij} the potential depth and r_{ij} the distance between spheres i and j . Concerning bonded interaction, two adjacent spheres are connected by a constrained bond of a length equal to σ .

To describe the polar compounds, i.e. H₂S and CO₂, a single LJ sphere combined with a simple isotropic multipolar potential (IMP) has been employed (Galliero et al., 2007a):

$$U_{ij}^{IMP} = U_{ij}^{LJ} - \frac{1}{k_B T} \left[\frac{p_i^2 p_j^2}{3r_{ij}^6} + \frac{p_i^2 Q_j^2 + p_j^2 Q_i^2}{2r_{ij}^8} + \frac{7Q_i^2 Q_j^2}{5r_{ij}^{10}} \right] \quad (4)$$

where p_i is the dipole moment of molecule i and Q_i the quadrupole moment of molecule i .

A cutoff radius equal to $3.5\sigma_{ij}$ has been used during simulations and long range corrections have been included (Allen and Tildesley, 1987). Cross parameters between molecules of different species have been deduced using the classical Lorentz-Berthelot combining rules.

The molecular parameters, provided in Table 1, have been adjusted using the pure component properties, including equilibrium vapour-liquid phase envelope and viscosities (Galliero et al., 2007a; Touzet et al., 2011).

Table 1: Molecular parameters employed in this work

Species	N	ϵ (J.mol ⁻¹)	σ (Å)	μ (D)	Q (D.Å)
C ₁	1	1246.5	3.7327	0	0
nC ₄	2	2075	3.979	0	0
nC ₁₂	5	2490	4.042	0	0
H ₂ S	1	2320	3.688	0.9	0
CO ₂	1	1619	3.758	0	4.0982

Thermo-gravitation MD algorithm

To simulate a one dimensional gravitational segregation process using MD, the simulation box is divided into N_s slabs along the vertical direction. Then, an external potential equal to -

$M_i g z$ is applied on each molecule belonging to slabs 1 to $N_s/2$, whereas an external potential equal to $+M_i g z$ is applied to each molecule belonging to slabs $N_s/2+1$ to N_s , see Fig. 1. By doing so, a bi-periodical pressure gradient is induced in the simulation box and the segregation process can take place.

The height, h , of the MD simulation box used in this work is typically of the order of 10 nm. So, to mimic the real size of a petroleum reservoir the amplitude of the “gravitational” acceleration is chosen so that the product gh is equal to that of a real reservoir (where h is of the order of hundreds of meters), i.e. in the simulations $g \approx 10^{11} \text{ m.s}^{-2}$ depending on the system studied. This approach is consistent as long as the response remains linear and if there are no fluid-fluid interfaces appearing in the system.




Figure 1: A sketch of the scheme employed to study thermogravitation using MD simulations (only half of the simulation box is shown).

To impose a geothermal gradient (taken here to be around $0.025\text{-}0.03 \text{ K.m}^{-1}$ as typically found in real petroleum reservoirs), a bi-periodical thermal gradient is applied to the simulation box by using the non-equilibrium algorithm proposed by Müller-Plathe, 1997. This scheme consists in exchanging kinetic energy between the edge layers and the central layers of the simulation box as shown in Fig. 1. As for the gravity field the temperature gradient applied

has been chosen to represent the one applied on the simulated reservoir column, resulting in huge thermal gradients of the order of 10^9 K.m⁻¹. However, in line with previous studies (Perronace et al., 2002; Galliero and Montel, 2008), the response of the system was found to be always linear. In addition, in a previous work (Galliero and Montel, 2008), a similar approach has been validated on “isotopic” mixtures for which an analytical solution exists.

It is important to note that because the fluid column is heated from below, which corresponds to a Rayleigh-Bénard configuration, the system may be unstable and convection may occur at the reservoir scale (Montel et al. 2007). However, during MD simulations, the Rayleigh number remains always below its critical threshold because of the small size (nm) of the simulation domain (Galliero and Montel 2008). Thus, the fluid column simulated by this approach is always stable (convection free) and only diffusive processes take place.

Simulations details

Using a homemade MD code, systems composed of 2000 to 4000 molecules have been simulated. The velocity Verlet algorithm has been applied to integrate the equation of motion (Allen and Tildesley, 1987). A reduced timestep, δt^* , equal to 0.003 has been used. To maintain the desired average temperature during simulations, a Berendsen thermostat with a large time constant equal to $1000\delta t^*$ has been utilized (Berendsen et al., 1984). Classical periodic boundary conditions combined with a Verlet neighbors list have been applied (Allen and Tildesley, 1987). Once the stationary state is reached, data have been collected during runs of $5 \cdot 10^7$ time steps. Slabs 1, 2, $N_s/2-1$, $N_s/2$, $N_s/2+1$, $N_s/2+2$, N_s-1 and N_s have been discarded when performing the analysis of the MD concentrations, temperature and density profiles. Error bars, computed using the sub-block method (Allen and Tildesley, 1987), are of the order of ± 0.01 on mole fraction, ± 0.1 K on temperature and ± 0.2 kg/m³ on density. They have been omitted in the figures, for sake of clarity.

Results and Discussion

Synthetic mixture composed of normal alkanes

As a first example we have studied the behavior of a C_1 - nC_4 - nC_{12} synthetic mixture in which the mole fractions are respectively 0.2, 0.4 and 0.4 (Van Vaerenbergh et al., 2009). Reference conditions at the mid column depth are a temperature equal to 333.15K and a pressure equal to 35 MPa. The study has been performed at a hypothetical depth between 2000 and 2600 m, consistent with the chosen thermodynamic conditions. Such a system is not intended to represent a real reservoir fluid, but it allows for studying a simple ternary asymmetric oil mixture composed of typical hydrocarbons at reservoir conditions.

Two cases have been studied:

- an isothermal case, in which only segregation occurs,
- a thermogravitation case in which the system is subject both to the gravity field and a geothermal gradient equals to 0.027 K.m^{-1} .

Using the scheme described previously, MD simulations have been performed for both cases. In addition, a conventional thermodynamic modeling using the Peng-Robinson EoS with volume shifts and cross interactions parameters has been carried out for the pure segregation case (Montel and Gouel, 1985). At the reference conditions, EoS and MD simulations yield results on density in excellent agreement with each other, i.e. $\rho_{\text{EoS}}=675.8 \text{ kg/m}^3$ and $\rho_{\text{MD}}=676\pm 0.1 \text{ kg/m}^3$.

As expected, when only the gravity field is active (segregation), C_1 is enriched at the top of the fluid column (Fig 2a) and nC_{12} is enriched at the bottom of the fluid column (Fig 2c). Concerning nC_4 , one can notice a small enrichment (of about 0.001 in mole fraction) at the top of the fluid column (Fig 2b). The concentration gradients obtained from the thermodynamic modeling are in good agreement with those coming from MD simulations

despite quite different approaches. It is worth pointing out that the thermodynamic approach yields gradients that are slightly larger than those coming from the MD simulations, see Fig. 2.




Figure 2: Mole fractions versus depth in a ternary C_1 - nC_4 - nC_{12} mixture subject to segregation (*isoT*) and thermogravitation (ΔT) using MD simulations and classical thermodynamic modelling.

When the geothermal gradient is added to segregation (thermogravitation) the MD simulation results are completely modified, see Figure 2. Mole fraction gradients of C_1 , nC_4 and nC_{12} are even reversed, compared to the purely gravitational segregation case, because of the

thermodiffusion effect. More precisely, nC_{12} is slightly enriched at the top of the reservoir whereas the opposite is true for C_1 . This indicates that, at least for the C_1 - nC_4 - nC_{12} mixture, thermodiffusion has a larger impact than gravitational segregation on the vertical distribution of the components. Such a result confirms quantitatively the importance of thermodiffusion, in addition to gravitational segregation, when trying to estimate the initial vertical distribution of the components in a petroleum reservoir.




Figure 3: Densities versus depth in a ternary C_1 - nC_4 - nC_{12} mixture subject to segregation (*isoT*) and thermogravitation (ΔT).

Another very interesting point concerns the density profile shown in Fig. 3. It appears that the density gradient is reversed when the reverse thermal gradient is present as compared to the gravitational-only segregation case. This is due to the thermal expansion effect and to thermodiffusion leading to an enrichment of the heavier compounds (nC_{12}) at the top of the fluid column, as seen in Fig. 2c. This reverse density gradient indicates that such a system would be unstable at the reservoir scale and would lead to convection. At the MD scale the system is, as already alluded to, always stable as the Rayleigh number remains always small and well below the critical threshold (Galliero and Montel, 2008).

In this respect, we note that in real petroleum reservoirs, since hydrocarbon mixtures are within a porous matrix, permeability impacts the dynamic of the system, e.g. the Rayleigh-Darcy number should be considered instead of the Rayleigh number (Montel et al., 2007).

Realistic acid gas mixture

As a second example, we have applied the approach to a real Far-East CO₂ rich reservoir. The reservoir fluid is an acid gas mixture composed of C₁, CO₂ and H₂S with an average mole fraction equal to 0.28, 0.71 and 0.01, respectively. Reference conditions at the mid column depth are a temperature of 443.15 K and a pressure of 40 MPa. The fluid column possesses an extremely high vertical extension equal to 1600m.

A methodology similar to that applied to the n-alkane mixture (see previous section) has been used to deal with the segregation and the thermogravitation processes using a geothermal gradient equal to 0.03 K.m⁻¹. At the reference conditions, EoS and MD simulations yield densities consistent with each other, i.e. $\rho_{\text{EoS}}=408.9 \text{ kg/m}^3$ and $\rho_{\text{MD}}= 407.3\pm 0.2 \text{ kg/m}^3$.

Concerning the thermogravitation profiles, illustrated in Fig. 4, the thermal gradient induces a noticeable modification of the molar fraction profiles as a direct consequence of the thermodiffusion process. However, contrary to the n-alkane mixture discussed previously, the molar fraction gradients are not reversed as compared to the purely gravitational segregation case. This implies that the gravity segregation is dominant over thermodiffusion for this acid gas mixture. This is expected as thermodiffusion is usually smaller in gases than in liquids (Wiegand, 2004; Galliero et al., 2007b). Nevertheless, it is clear from these MD results that neglecting thermodiffusion could lead to large errors when estimating the initial state of the reservoir.




Figure 4: Mole fractions versus depth in a ternary CO_2 - C_1 - H_2S mixture subject to segregation (*isoT*) and thermogravitation (ΔT) using MD simulations and thermodynamic modelling.

Similarly to what was found on the n-alkane mixture, the chosen thermal gradient, see Fig. 5, leads to an unstable fluid column (slightly reverse density gradient). However, after more detailed investigation of field data, the thermal gradient has been reevaluated and the current consensus is that $0.02 \text{ K}\cdot\text{m}^{-1}$ is a more likely value. With such thermal gradient, the fluid column remains stable.




Figure 5: *Densities versus depth in an acid gas mixture subject to segregation (isoT) and thermogravitation (ΔT).*

Conclusions

We have presented in this work a methodology to estimate the initial vertical distribution of the components in an idealized hydrocarbon reservoir subject to gravity alone (segregation) or combined with a geothermal gradient (thermogravitation). The presented approach made use of Molecular Dynamics simulations and can be used to test and complement the usual methods based on classical EoS. Furthermore, it can provide thermodiffusion coefficients and shed light on some of the underlying physical mechanism (stability) of the thermogravitational process.

This methodology has been applied to two mixtures, a synthetic oil composed of C_1 , nC_4 and nC_{12} and a real acid gas composed of CO_2 , C_1 and H_2S . For the segregation case, an excellent agreement has been found between the proposed methodology and the results obtained by classical thermodynamic model (EoS). For thermogravitation case, MD simulations have shown that the thermodiffusion effect induced by the geothermal gradient is similar in magnitude (but opposite in sign) to that of gravitational segregation one. In addition, it has been observed that the density gradient can be reversed resulting in a higher density at the top

of the fluid column than at the bottom, (i.e. an unstable situation) as a consequence of thermal expansion and thermodiffusion due to a presence of a geothermal gradient.

These numerical results confirm quantitatively the fact that thermodiffusion in multi-component mixtures noticeably impacts the initial vertical distribution of the species in a petroleum reservoir and should be taken into account to determine accurately the initial state of a reservoir. Although molecular dynamics and classical EoS can provide important information on such systems, it remains paramount to validate these modeling tools by actual measurements. The SCCO-SJ10 experiments will be a crucial step towards this goal.

Acknowledgements

We gratefully acknowledge computational facilities provided by the University of Pau and the University of Bordeaux. This work has been supported by ESA through the SCCO project. The authors thank Total S.A. for permission to publish this paper. The Spanish teams (Mondragon Unibertsitatea and Universidad Complutense) are thankful to the TERDISOMEZ (FIS2014-58950-C2-1-P and FIS2014-58950-C2-2-P) of MINECO.

References

- Allen, M.P., Tildesley, D.J.: Computer simulation of liquids. Oxford: Clarendon Press (1987)
- Artola, P.A., Rousseau, B.: Thermal diffusion in simple liquid mixtures: What have we learnt from molecular dynamics simulations?, *Molec. Phys.* 111, 3394-3403 (2013).
- Assael, M.J., Goodwin, A.R.H., Vesovic, V., Wakeham, W.A.: Experimental Thermodynamics Volume IX: Advances in Transport Properties of Fluids, Royal Society of Chemistry, London, (2014).
- Berendsen, H.J.C., Postam, J.P.M., Van Gunsteren, W.F., DiNola, A., Haak, J.R.: Molecular Dynamics with coupling to an external bath, *J. Chem. Phys.* 81, 3684-3690 (1984).
- Bou-Ali, M.M., Ahadi, A., Alonso de Mezquia, D., Galand, Q., Gebhardt, M., Khlybov, O., Köhler, W., Larrañaga, M., Legros, J.C., Lyubimova, T., Mialdun, A., Ryzhkov, I., Saghir, M.Z., Shevtsova, V., Van Vaerenbergh, S.: Benchmark values for the Soret, thermodiffusion and molecular diffusion coefficients of the ternary mixture tetralin+isobutylbenzene+n-dodecane with 0.8-0.1-0.1 mass fraction, *Eur. Phys. J. E* 38, 30 (2015).
- Croccolo, F., Bataller, H., Scheffold, F.: A light scattering study of non equilibrium fluctuations in liquid mixtures to measure the Soret and mass diffusion coefficient, *J. Chem. Phys.* 137, 234202 (2012).
- de Groot, S.R., Mazur, P.: *Nonequilibrium Thermodynamics*. Dover, New York (1984).
- Firoozabadi, A., Ghorayeb, K., Shukla, K.: Theoretical model of thermal diffusion factors in multicomponent mixtures, *AIChE Journal* 46, 892-900, 2000.
- Galliero, G., Duguay, B., Caltagirone, J.P., Montel, F.: On thermal diffusion in binary and ternary mixtures by non-equilibrium molecular dynamics, *Phil. Mag.* 83, 2097-2108 (2003).
- Galliero, G., Colombani, J., Bopp, P.A., Duguay, B., Caltagirone, J.P., Montel, F.: Thermal diffusion in micropores by molecular dynamics computer simulations, *Physica A* 361, 494-510 (2006).

Galliero, G., Nieto-Draghi, C., Boned, C., Avalos, J.B., Mackie, A.D., Baylaucq, A., Montel, F.: Molecular dynamics simulation of acid gas mixtures: A comparison between several approximations, *Ind. Eng. Chem. Res.* 46, 5238-5244 (2007a).

Galliero, G., Bugel, M., Duguay, B., Montel, F.: Mass effect on thermodiffusion using molecular dynamics, *J. Non-Equi. Thermodyn.* 32, 251-258 (2007b)

Galliero, G., Montel, F.: Nonisothermal gravitational segregation by molecular dynamics simulations, *Phys. Rev. E* 78, 041203 (2008).

Galliero, G., Srinivasan, S., Saghir, M.Z.: Estimation of thermodiffusion in ternary alkane mixtures using molecular dynamics and an irreversible thermodynamic theory, *High Temp. – High Press* 38, 315-328 (2009).

Galliero G.: Equilibrium, interfacial and transport properties of n-alkanes: Towards the simplest coarse grained molecular model, *Chem. Eng. Res. Des.* 92, 2031-2037 (2014).

Georis, P., Montel, F., Van Vaerenbergh, S., Decoly, Y., Legros, J.C.: Proceedings of the European Petroleum Conference, 1, pp. 57-62 (1998).

Ghorayeb, K., Firoozabadi, A., Anraku, T.: Interpretation of the unusual fluid distribution in the Yufutsu gas-condensate field, *SPE Journal* 8, 114-123 (2003).

Giraudet, C., Bataller, H., Croccolo, F.: High-pressure mass transport properties measured by dynamic near-field scattering of non-equilibrium fluctuations, *Eur. Phys. J. E* 37, 107 (2014).

Halldórsson, S., Stenby, E.H.: Isothermal gravitational segregation: Algorithms and specifications, *Fluid Phase Equil.* 175 (1-2): 175-183 (2000).

Hannaoui, R., Galliero, G., Hoang, H., Boned, C.: Influence of confinement on thermodiffusion, *J. Chem. Phys.* 139, 114704 (2013).

Høier, L., Whitson, C.H.: Compositional grading-theory and practice, *SPE Reservoir Evaluation and Engineering* 4, 525-535 (2001).

Holt, T., Lindeberg E., Ratkje, K.S.: The effect of gravity and temperature gradients on Methane distribution in Oil reservoirs, *SPE Paper* 11761 (1983).

Hu W.R. J. F. Zhao, M. Long, X. W. Zhang, Q. S. Liu, M. Y. Hou, Q. Kang, Y. R. Wang, S. H. Xu, W. J. Kong, H. Zhang, S. F. Wang, Y. Q. Sun, H. Y. Hang, Y. P. Huang, W. M. Cai, Y. Zhao, J. W. Dai, H. Q. Zheng, E. K. Duan, J. F. Wang: Space Program SJ10 of Microgravity Research, *Microgravity Sci. Tec.*, 26, 156-169 (2014).

Kempers L.J.T.M.: A comprehensive thermodynamic theory of the Soret effect in a multicomponent gas, liquid, or solid, *J. Chem. Phys.* 115, 6330-6341 (2001).

Larrañaga, M., Bou-Ali, M.M., Alonso de Mezquía, D., Rees, D.A.S., Madariaga, J.A., Santamaría, C., Platten, J.K.: Contribution to the benchmark for ternary mixtures: Determination of Soret coefficients by the thermogravitational and the sliding symmetric tubes techniques, *Eur. Phys. J. E* 38: 28 (2015).

Leahy-Dios, A., Bou-Ali, M.M., Platten, J.K., Firoozabadi, A.: Measurements of molecular and thermal diffusion coefficients in ternary mixtures, *J. Chem. Phys.* 122, 234502 (2005).

Montel, F., Gouel, P.L.: Prediction of compositional grading in a reservoir fluid column, *SPE Paper* 14410 (1985).

Montel, F., Bickert, J., Lagisquet, A., Galliero, G.: Initial state of petroleum reservoirs: A comprehensive approach, *J. Pet. Sci. Eng.* 58, 391-402 (2007).

Müller-Plathe, F.: A simple nonequilibrium molecular dynamics method for calculating the thermal conductivity, *J. Chem. Phys.* 106, 6082-6085 (1997).

Ortiz de Zárate, J.M., Giraudet, C., Bataller, H., Croccolo, F.: Non-equilibrium fluctuations induced by the Soret effect in a ternary mixture, *Eur. Phys. J. E* 37, 77 (2014).

Perronace, A., Leppla, C., Leroy, F., Rousseau, B., Wiegand, S.: Soret and mass diffusion measurements and molecular dynamics simulations of n-pentane-n-decane mixtures, *J. Chem. Phys.* 116, 3718-3729 (2002).

Platten, J.K., Costeseque, P.: The Soret coefficient in porous media, *J. Porous Media* 7, 317-329 (2004).

Sage, B.H., Lacey, W.N.: Gravitational concentration gradients in static column of hydrocarbon fluids, *Trans. AIME* 132, 120-131 (1939).

Shapiro, A., Stenby, E.H.: Factorization of transport coefficients in macroporous media, *Transp. Porous Media* 41, 305-323 (2000).

Touzet, M., Galliero, G., Lazzeri, V., Saghir, M.Z., Montel, F., Legros, J.C.: Thermodiffusion: from microgravity experiments to the initial state of petroleum reservoirs, *Comptes Rendus - Mécanique* 339, 318-323 (2011).

Urteaga, P., Bou-Ali, M.M., Alonso de Mezquía, D., Santamaría, J., Santamaría, C., Madariaga, J.A., Bataller, H.: Measurement of thermodiffusion coefficient of hydrocarbon binary mixtures under pressure with the thermogravitational technique, *Rev. Sci. Instrum.* 83, 074903 (2012).

Van Vaerenbergh, S., Srinivasan, S., Saghir M.Z.: Thermodiffusion in multicomponent hydrocarbon mixtures: Experimental investigations and computational analysis, *J. Chem. Phys.* 131, 114505 (2009).

Whitson, C.H., Belery, P.: Compositional Gradients in Petroleum Reservoirs, *SPE Paper* 28000 (1994).

Wiegand, S.: Thermal diffusion in liquid mixtures and polymer solutions, *J. Phys.: Cond. Matter* 16, R357-R379 (2004).