

Evidence for the rare decay $\Sigma^+ \to p \mu^+ \mu^-$

Roel Aaij, Bernardo Adeva, Marco Adinolfi, Ziad Ajaltouni, Simon Akar, Johannes Albrecht, Federico Alessio, Michael Alexander, Alejandro Alfonso Albero, Suvayu Ali, et al.

▶ To cite this version:

Roel Aaij, Bernardo Adeva, Marco Adinolfi, Ziad Ajaltouni, Simon Akar, et al.. Evidence for the rare decay $\Sigma^+ \rightarrow p \mu^+ \mu^-$. Phys.Rev.Lett., 2018, 120 (22), pp.221803. 10.1103/PhysRevLett.120.221803. hal-01807288

HAL Id: hal-01807288 https://hal.science/hal-01807288

Submitted on 12 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH (CERN)



CERN-EP-2017-319 LHCb-PAPER-2017-049 June 4, 2018

Evidence for the rare decay $\Sigma^+ ightarrow p \mu^+ \mu^-$

The LHCb collaboration[†]

Abstract

A search for the rare decay $\Sigma^+ \to p\mu^+\mu^-$ is performed using pp collision data recorded by the LHCb experiment at centre-of-mass energies $\sqrt{s} = 7$ and 8 TeV, corresponding to an integrated luminosity of 3 fb⁻¹. An excess of events is observed with respect to the background expectation, with a signal significance of 4.1 standard deviations. No significant structure is observed in the dimuon invariant mass distribution, in contrast with a previous result from the HyperCP experiment. The measured $\Sigma^+ \to p\mu^+\mu^-$ branching fraction is $(2.2^{+1.8}_{-1.3}) \times 10^{-8}$, where statistical and systematic uncertainties are included, which is consistent with the Standard Model prediction.

Published in Phys. Rev. Lett. 120 (2018) 221803.

© CERN on behalf of the LHCb collaboration, licence CC-BY-4.0.

[†]Authors are listed at the end of this Letter.

The $\Sigma^+ \to p \mu^+ \mu^-$ decay is an $s \to d$ quark-flavour-changing neutral-current process, allowed only at loop level in the standard model (SM). The process is dominated by long-distance contributions for a predicted branching fraction of $1.6 \times 10^{-8} < \mathcal{B}(\Sigma^+ \rightarrow \Sigma^+)$ $p\mu^+\mu^-) < 9.0 \times 10^{-8}$ [1], while the short-distance SM contributions are suppressed and contribute to the branching fraction at the level of about 10^{-12} . Evidence for this decay was reported by the HyperCP collaboration [2] with a measured branching fraction $\mathcal{B}(\Sigma^+ \to p\mu^+\mu^-) = (8.6^{+6.6}_{-5.4} \pm 5.5) \times 10^{-8}$, which is compatible with the SM prediction. HyperCP observed three candidates; remarkably, all of them have almost the same dimuon invariant mass of $m_{X^0} = 214.3 \pm 0.5 \,\text{MeV}/c^2$, close to the lower kinematic limit. Such a distribution, if confirmed, would point towards a process with an intermediate particle X^0 coming from the Σ^+ baryon and decaying into two muons, *i.e.* a $\Sigma^+ \to pX^0 (\to \mu^+ \mu^-)$ decay, which would constitute evidence for physics beyond the SM (BSM). Various BSM theories have been proposed to explain the HyperCP result. The intermediate X^0 particle could be, for example, a light pseudoscalar Higgs boson [3, 4] or a sgoldstino [5, 6] in various supersymmetric models. Other interpretations and implications can be found in Refs. [7–13]; in general a pseudoscalar particle is favoured over a scalar particle and a lifetime of the order of 10^{-14} s is estimated for the former case. Attempts to confirm the existence of this X^0 particle have been made at several experiments in various initial and final states without finding any signal [14–21]; these null results include studies of the decays $B^0_{(s)} \to \mu^+ \mu^- \mu^+ \mu^-$ [22], $B^0 \to K^{*0} \mu^+ \mu^-$ [23], $B^+ \to K^+ \mu^+ \mu^-$ [24], and a search for photon-like particles [25] by the LHCb experiment. However, the search for the $\Sigma^+ \to p \mu^+ \mu^-$ decay has not been repeated due to the lack of experiments with large hyperon production rates and to the experimental difficulty of reconstructing soft and long-lived hadrons.

Hyperons are produced copiously in high-energy proton-proton collisions at the Large Hadron Collider. A search for $\Sigma^+ \to p\mu^+\mu^-$ decays at the LHCb experiment, as also suggested in Ref. [26], could therefore confirm or disprove the HyperCP evidence, and the branching fraction can be measured. This Letter presents a search for the $\Sigma^+ \to p\mu^+\mu^$ decay performed using pp collision data recorded by the LHCb experiment at centre-ofmass energies $\sqrt{s} = 7$ and 8 TeV, corresponding to an integrated luminosity of 3 fb⁻¹. The inclusion of charge-conjugated processes is implied throughout this Letter.

This search follows a strategy similar to that of other studies of rare decays in LHCb, although with differences due to the relatively low transverse momenta of the final-state particles. First, a loose selection is applied based on geometric and kinematic variables. The final sample is obtained rejecting the background with requirements on the output of a multivariate selection, based on a boosted decision tree algorithm (BDT) [27,28], and on particle identification variables. The signal yield is obtained from a fit to the $p\mu^+\mu^$ invariant-mass spectrum and is converted into a branching fraction by normalising to the $\Sigma^+ \to p\pi^0$ control channel. The analysis is designed in order to search for possible peaks in the dimuon invariant-mass distribution, in view of the possible existence of unknown intermediate particles.

The LHCb detector is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, described in detail in Refs. [29, 30]. It includes a high-precision tracking system consisting of a silicon-strip vertex detector surrounding the *pp* interaction region, a large-area silicon-strip detector located upstream of a dipole magnet with a bending power of about 4 Tm, and three stations of silicon-strip detectors and straw drift tubes placed downstream of the magnet. Particle identification is provided by two ring-imaging Cherenkov detectors, an electromagnetic and a hadronic calorimeter, and a muon system composed of alternating layers of iron and multiwire proportional chambers.

The online event selection is performed by a trigger system, which consists of a hardware stage, based on information from the calorimeter and muon systems, followed by two software stages. The first software stage performs a preliminary event reconstruction based on partial information while the second applies a full event reconstruction. Each of the three trigger stages is divided into many trigger selections dedicated to various types of signal. The final-state particles from the signal decay involved in this analysis typically have insufficient transverse momenta to satisfy the requirements of one or more trigger stages. Nevertheless, given the large production rate of Σ^+ baryons in pp collisions. the present search can be performed with data selected at one or more trigger stages by other particles in the event. In the offline processing, trigger decisions are associated with reconstructed candidates. A trigger decision can thus be ascribed to the reconstructed candidate, the rest of the event or a combination of both; events triggered as such are defined respectively as triggered on signal (TOS), triggered independently of signal (TIS). and triggered on both. While all the candidates passing the trigger selection are used in the search for $\Sigma^+ \to p \mu^+ \mu^-$ decays, only the TIS candidates are used in the normalisation channel $\Sigma^+ \to p\pi^0$. Furthermore, control channels with large yields are exploited to estimate the trigger efficiency by measuring the overlap of candidates which are TIS and TOS simultaneously [31].

Simulation is used to devise and optimise the analysis strategy, as well as to estimate reconstruction and selection efficiencies. In the simulation, pp collisions are generated using PYTHIA [32] with a specific LHCb configuration [33]. Decays of hadronic particles are described by EVTGEN [34], in which final-state radiation is generated using PHOTOS [35]. The interaction of the generated particles with the detector, and its response, are implemented using the GEANT4 toolkit [36], as described in Ref. [37]. The signal $\Sigma^+ \to p\mu^+\mu^$ decay is generated according to a phase-space model.

Candidate $\Sigma^+ \to p \mu^+ \mu^-$ decays are selected by combining two good-quality oppositely charged tracks identified as muons with a third track identified as a proton. The three tracks are required to form a secondary vertex (SV) with a good vertex-fit quality. The short lifetime estimated for the X^0 particle would result in a prompt signal in this search, hence no attempt is made to distinguish the dimuon origin vertex from the SV of the Σ^+ baryon. The measured Σ^+ candidate proper decay time is required to be greater than 6 ps, ensuring that the SV is displaced from any pp interaction vertex (primary vertex, PV). The final-state particles are required to be inconsistent with originating from any PV in the event. Only Σ^+ candidates with transverse momentum $p_{\rm T} > 0.5 \,{\rm GeV}/c$ and a decay topology consistent with a particle originating from the PV are retained. A candidate $\Sigma^+ \to p\mu^+\mu^-$ decay is considered only if its invariant mass, $m_{p\mu^+\mu^-}$, satisfies $|m_{p\mu^+\mu^-} - m_{\Sigma^+}| < 500 \text{ MeV}/c^2$, where m_{Σ^+} is the known mass of the Σ^+ particle [38]. The background component due to $\Lambda \to p\pi^-$ decays is vetoed by discarding candidates having a $p\mu^-$ pair invariant mass, calculated with the $p\pi^-$ mass hypothesis, within $10 \,\mathrm{MeV}/c^2$ from the known A mass [38]. Possible backgrounds from decays peaking in the $p\mu^+\mu^$ invariant mass have been examined, including $K^+ \to \pi^+ \pi^- \pi^+$, $K^+ \to \pi^+ \mu^- \mu^+$, and various hyperon decays, and none has been found to contribute significantly. After all selection requirements, no retained event contains more than one candidate.

Candidate $\Sigma^+ \to p\pi^0$ decays are selected by combining one good-quality track identified as a proton with a π^0 reconstructed in the $\pi^0 \to \gamma\gamma$ mode from two clusters in the electromagnetic calorimeter. Given the impossibility to reconstruct the Σ^+ decay SV with the proton track only, the momentum direction of the π^0 is calculated assuming the π^0 is produced at the PV. The selection of $\Sigma^+ \to p\pi^0$ decays is similar to that of the signal, with tighter requirements applied, in order to reduce the large combinatorial background, on the proton identification and on the transverse momenta of the final-state particles ($p_T > 0.5 \text{ GeV}/c$ for the proton and $p_T > 0.7 \text{ GeV}/c$ for the π^0). Finally, candidate $K^+ \to \pi^+\pi^-\pi^+$ decays, selected as control channel for various parts of the analysis, are required to pass a selection similar to that of the signal, starting from three good-quality tracks, with total charge equal to ± 1 , and which are assigned the pion mass hypothesis without requirements on the identification of the particle.

The sample of $\Sigma^+ \to p \mu^+ \mu^-$ candidates in data after the initial selection is dominated by combinatorial background, part of which is due to misidentified particles. This background is rejected by placing requirements on the BDT output variable and on multivariate particle identification variables [30] on the muons and on the proton. The BDT combines information from the following input variables: the angle between the Σ^+ reconstructed momentum and the vector joining the PV to the SV, the flight distance significance of the Σ^+ candidate, the distance of closest approach among the final-state particles, the transverse momenta of the final-state particles, the impact parameter χ^2 $(\chi_{\rm IP}^2)$ of the final-state particles, defined as the difference between the vertex-fit χ^2 of a PV formed with and without the particle in question, the $\chi_{\rm IP}^2$ of the Σ^+ candidate, the χ^2 of the SV, and an isolation variable constructed from the number of tracks within an angular cone around each of the final-state particles. These variables are chosen so that the dependence on the $p\mu^+\mu^-$ invariant mass and on the dimuon invariant mass is small and linear to minimise potential biases. The BDT is optimised using simulated samples of $\Sigma^+ \to p \mu^+ \mu^-$ events for the signal and $\overline{p} \mu^+ \mu^+$ candidates in data for the background. The selection for the control $\overline{p}\mu^+\mu^+$ sample is identical to that of the signal but considering muons of identical charge. The final selection criteria are chosen in order to optimise the potential to obtain evidence for a signal with a branching fraction as small as possible [39]. No BDT selection is applied to the normalisation and control channels.

The number of signal candidates is converted into a branching fraction with the formula

$$\mathcal{B}(\Sigma^+ \to p\mu^+\mu^-) = \frac{\varepsilon_{\Sigma^+ \to p\pi^0}}{\varepsilon_{\Sigma^+ \to p\mu^+\mu^-}} \frac{N_{\Sigma^+ \to p\mu^+\mu^-}}{N_{\Sigma^+ \to p\pi^0}} \mathcal{B}(\Sigma^+ \to p\pi^0)$$
$$= \alpha \cdot N_{\Sigma^+ \to p\mu^+\mu^-} ,$$

where ε , N and \mathcal{B} are the efficiency, candidate yield and branching fraction of the corresponding channel, respectively, and α is the single-event sensitivity. The ratio of signal and normalisation channel efficiencies, which includes the acceptance, the trigger efficiency, the reconstruction efficiency of the final-state particles and the selection efficiency, is computed with samples of simulated events corrected to take into account known differences between data and simulation. The reconstruction efficiency for the π^0 is calibrated using the ratio of $B^+ \to J/\psi K^{*+}(\to K^+\pi^0)$ and $B^+ \to J/\psi K^+$ decays reconstructed in data [40]. The particle-identification efficiencies of protons and muons are calibrated with control channels in data. Residual differences between data and simulation are treated as sources of systematic uncertainty. The ratio of the trigger efficiencies for the signal and normalisation channels is estimated with simulated samples and cross-checked in data: the trigger efficiency is obtained for selected trigger lines from the overlap of TIS and TOS events in the normalisation channel and is compared between data and

Source	Uncertainty
Selection efficiency	1%
BDT efficiency	6%
PID efficiency ratio	28%
π^0 efficiency	10%
Trigger efficiency ratio	40%
Total	50%

Table 1: Relative systematic uncertainties associated with the normalisation.



Figure 1: Distribution of the corrected mass m_{Σ}^{corr} , defined as in the text, for $\Sigma^+ \to p\pi^0$ candidates superimposed with the fit to data.

simulation [31]. The small size of this overlap induces a 40% relative systematic uncertainty associated with the trigger efficiency ratio. The ratio of the trigger efficiencies is of the order of 0.09, owing to the use of all events for the signal, while TIS-only events are used for the normalisation channel. Possible differences in the BDT selection efficiency for the $\Sigma^+ \to p\mu^+\mu^-$ signal in data and in simulation are calibrated using the $K^+ \to \pi^+\pi^-\pi^+$ control channel. The sources of systematic uncertainties associated with the normalisation are reported in Table 1.

The observed number of $\Sigma^+ \to p\pi^0$ candidates is $(1171 \pm 9) \times 10^3$, as obtained from a binned extended maximum likelihood fit to the corrected invariant mass distribution m_{Σ}^{corr} . The corrected invariant mass is defined as $m_{\Sigma}^{\text{corr}} = m_{p\gamma\gamma} - m_{\gamma\gamma} + m_{\pi^0}$, where m_{π^0} is the known mass of the π^0 meson [38], to account for the limited precision in the reconstructed invariant mass of the two photons $(m_{\gamma\gamma})$. The $\Sigma^+ \to p\pi^0$ distribution is described as a Gaussian function with a power-law tail on the higher-mass side, while the background is described by a modified ARGUS function [41], where the power parameter is allowed to vary as in Ref. [42]. The distribution is shown in Fig. 1, superimposed with the fit.

The single-event sensitivity is $\alpha = (2.2 \pm 1.2) \times 10^{-9}$, where the uncertainty is dominated by the systematic contribution. This sensitivity corresponds to about $10^{14} \Sigma^+$ baryons



Figure 2: Invariant mass distribution of $\Sigma^+ \to p \mu^+ \mu^-$ candidates in data.

produced in the LHCb acceptance in the considered dataset. The number of expected signal $\Sigma^+ \to p \mu^+ \mu^-$ candidates is 23 ± 20 assuming a branching fraction of $(5 \pm 4) \times 10^{-8}$, to cover the range predicted by the SM.

The observed number of signal $\Sigma^+ \to p\mu^+\mu^-$ decays is obtained with a fit to the $p\mu^+\mu^$ invariant-mass distribution in the range 1149.6 $< m_{p\mu^+\mu^-} < 1409.6 \text{ MeV}/c^2$. The signal distribution is described by an Hypatia function [43]. The peak position and resolution are calibrated using the control channel $K^+ \to \pi^+\pi^-\pi^+$ and by comparing distributions in data and simulation. No bias is seen in the peak position, while a relative positive correction of 25% with respect to the simulation is applied to the resolution. A resolution of $4.28 \pm 0.19 \text{ MeV}/c^2$ is obtained for the signal $\Sigma^+ \to p\mu^+\mu^-$ distribution and is used in the fit to define a Gaussian constraint to the width of the signal distribution. The combinatorial background is described as a modified ARGUS function with all parameters left free with the exception of the threshold, which is fixed to the kinematic limit. The shape of this background is also cross-checked with that of $\overline{p}\mu^+\mu^+$ candidates in data.

The invariant mass distribution of the $\Sigma^+ \to p\mu^+\mu^-$ candidates in data is shown in Fig. 2. The significance of the signal is 4.1σ , obtained from a comparison of the likelihood value of the nominal fit with that of a background-only fit [44], and with the relevant systematic uncertainties included as Gaussian constraints to the likelihood. A signal yield of $10.2^{+3.9}_{-3.5}$ is observed. The corresponding branching fraction is $\mathcal{B}(\Sigma^+ \to p\mu^+\mu^-) = (2.2^{+0.9}_{-0.8} + 1.5) \times 10^{-8}$, where the first uncertainty is statistical and the second is systematic, consistent with the SM prediction. As a cross-check, the fit is repeated with tighter or looser requirements on the BDT or on the particle identification variables, and the signal yield is found to vary consistently with the signal efficiency. The fit is also repeated assuming a linear function for the background, in place of an ARGUS function, and the signal yield and significance are found to be stable. Candidates in data are composed of about $48\% \overline{\Sigma^+}$ anti-baryons in the final sample.

The distribution of the dimuon invariant mass after background subtraction, performed with the *sPlot* method [45], is shown in Fig. 3. A scan for a possible resonant structure in the dimuon invariant mass is performed, considering a region within two times the resolution in the $p\mu^+\mu^-$ invariant mass around the known Σ^+ mass. The distribution of



Figure 3: Background-subtracted distribution of the dimuon invariant mass for $\Sigma^+ \to p\mu^+\mu^-$ candidates, superimposed with the distribution from the simulated phase-space (PS) model. Uncertainties on data points are calculated as the square root of the sum of squared weights.

these candidates as a function of the dimuon invariant mass is shown in the supplemental material to this Letter [46]. Steps of half the resolution on the dimuon invariant mass, $\sigma(m_{\mu^+\mu^-})$, are considered in this scan, following the method outlined in Ref. [47]. The value of $\sigma(m_{\mu^+\mu^-})$ varies in the range [0.3, 2.3] MeV/ c^2 depending on the dimuon invariant mass as shown in Ref. [46]. For each step the putative signal is estimated in a window of $\pm 1.5 \times \sigma(m_{\mu^+\mu^-})$ around the considered particle mass, while the background is estimated from the lower and upper sidebands contained in the range $[1.5 - 4.0] \times \sigma(m_{\mu^+\mu^-})$ from the same mass. Only one of the two sidebands is considered when the second is outside the allowed kinematic range. The local p-value of the background-only hypothesis as a function of the dimuon mass is shown in Ref. [46], and no significant signal is found. The fit to the $p\mu^+\mu^-$ invariant mass is then repeated restricting the sample to events within 1.5 times the resolution from the putative particle $(m_{\mu^+\mu^-} \in [214.3 \pm 0.75] \text{ MeV}/c^2)$. No significant signal is found and a yield of $3.0^{+1.7}_{-1.4}$ is measured corresponding to 30% of the $\Sigma^+ \to p\mu^+\mu^-$ yield. An upper limit on the branching fraction of the resonant channel is thus set with the CL_S method [48] at $\mathcal{B}(\Sigma^+ \to pX^0(\to \mu^+\mu^-)) < 1.4 \times 10^{-8} (1.7 \times 10^{-8})$ at 90% (95%) confidence level.

In summary, a search for the $\Sigma^+ \to p\mu^+\mu^-$ rare decay is performed by the LHCb experiment using pp collisions at centre-of-mass energies $\sqrt{s} = 7$ and 8 TeV, corresponding to an integrated luminosity of 3 fb⁻¹. Evidence for the $\Sigma^+ \to p\mu^+\mu^-$ decay is found with a significance of 4.1 standard deviations, including systematic uncertainties. A branching fraction $\mathcal{B}(\Sigma^+ \to p\mu^+\mu^-) = (2.2^{+1.8}_{-1.3}) \times 10^{-8}$ is measured, consistent with the SM prediction. No significant peak consistent with an intermediate particle is found in the dimuon invariant-mass distribution of the signal candidates.

Acknowledgements

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (The Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MinES and FASO (Russia); MinECo (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (The Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (USA). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from AvH Foundation (Germany), EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union), ANR, Labex P2IO and OCEVU, and Région Auvergne-Rhône-Alpes (France), RFBR, RSF and Yandex LLC (Russia), GVA, XuntaGal and GENCAT (Spain), Herchel Smith Fund, the Royal Society, the English-Speaking Union and the Leverhulme Trust (United Kingdom).

References

- [1] X.-G. He, J. Tandean, and G. Valencia, $Decay \Sigma^+ \to p\ell^+\ell^-$ within the standard model, Phys. Rev. **D72** (2005) 074003, arXiv:hep-ph/0506067.
- [2] HyperCP collaboration, H. Park *et al.*, Evidence for the decay $\Sigma^+ \to p\mu^+\mu^-$, Phys. Rev. Lett. **94** (2005) 021801, arXiv:hep-ex/0501014.
- [3] X.-G. He, J. Tandean, and G. Valencia, Does the HyperCP evidence for the decay Σ⁺ → pμ⁺μ⁻ indicate a light pseudoscalar Higgs boson?, Phys. Rev. Lett. 98 (2007) 081802, arXiv:hep-ph/0610362.
- [4] X.-G. He, J. Tandean, and G. Valencia, Light Higgs production in hyperon decay, Phys. Rev. D74 (2006) 115015, arXiv:hep-ph/0610274.
- [5] D. S. Gorbunov and V. A. Rubakov, Sgoldstino interpretation of HyperCP events, Phys. Rev. D73 (2006) 035002, arXiv:hep-ph/0509147.
- [6] S. V. Demidov and D. S. Gorbunov, More about the sgoldstino interpretation of HyperCP events, JETP Lett. 84 (2007) 479, arXiv:hep-ph/0610066.
- [7] X.-G. He, J. Tandean, and G. Valencia, *Implications of a new particle from the HyperCP data on* $\Sigma^+ \rightarrow p\mu^+\mu^-$, Phys. Lett. **B631** (2005) 100, arXiv:hep-ph/0509041.
- [8] C. Q. Geng and Y. K. Hsiao, Constraints on the new particle in $\Sigma^+ \to p\mu^+\mu^-$, Phys. Lett. **B632** (2006) 215, arXiv:hep-ph/0509175.
- [9] N. G. Deshpande, G. Eilam, and J. Jiang, On the possibility of a new boson X^0 (214 MeV) in $\Sigma^+ \to p\mu^+\mu^-$, Phys. Lett. **B632** (2006) 212, arXiv:hep-ph/0509081.

- [10] C.-H. Chen, C.-Q. Geng, and C.-W. Kao, U-boson and the HyperCP exotic events, Phys. Lett. B663 (2008) 400, arXiv:0708.0937.
- [11] G. Xiangdong, C. S. Li, Z. Li, and H. Zhang, Contributions from SUSY-FCNC couplings to the interpretation of the HyperCP events for the decay $\Sigma^+ \to p\mu^+\mu^-$, Eur. Phys. J. C55 (2008) 317, arXiv:0712.0257.
- [12] M. L. Mangano and P. Nason, Radiative quarkonium decays and the NMSSM Higgs interpretation of the HyperCP $\Sigma^+ \to p\mu^+\mu^-$ events, Mod. Phys. Lett. A22 (2007) 1373, arXiv:0704.1719.
- [13] M. Pospelov, Secluded U(1) below the weak scale, Phys. Rev. D80 (2009) 095002, arXiv:0811.1030.
- [14] CLEO collaboration, W. Love et al., Search for very light CP-odd Higgs boson in radiative decays of $\Upsilon(1S)$, Phys. Rev. Lett. **101** (2008) 151802, arXiv:0807.1427.
- [15] E391a collaboration, Y. C. Tung *et al.*, Search for a light pseudoscalar particle in the decay $K_L^0 \to \pi^0 \pi^0 X$, Phys. Rev. Lett. **102** (2009) 051802, arXiv:0810.4222.
- [16] D0 collaboration, V. M. Abazov *et al.*, Search for Next-to-Minimal Supersymmetric Higgs bosons in the $h \to aa \to \mu\mu\mu\mu$, $\mu\mu\tau\tau$ channels using $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \ TeV$, Phys. Rev. Lett. **103** (2009) 061801, arXiv:0905.3381.
- [17] BaBar collaboration, B. Aubert *et al.*, Search for dimuon decays of a light scalar boson in radiative transitions $\Upsilon \rightarrow \gamma A^0$, Phys. Rev. Lett. **103** (2009) 081803, arXiv:0905.4539.
- [18] Belle collaboration, H. J. Hyun *et al.*, Search for a low mass particle decaying into $\mu^+\mu^-$ in $B^0 \to K^{*0}X$ and $B^0 \to \rho^0 X$ at Belle, Phys. Rev. Lett. **105** (2010) 091801, arXiv:1005.1450.
- [19] KTeV collaboration, E. Abouzaid *et al.*, Search for the rare decays $K_L \rightarrow \pi^0 \pi^0 \mu^+ \mu^-$ and $K_L \rightarrow \pi^0 \pi^0 X^0 \rightarrow \pi^0 \pi^0 \mu^+ \mu^-$, Phys. Rev. Lett. **107** (2011) 201803, arXiv:1105.4800.
- [20] BaBar collaboration, J. P. Lees et al., Search for a dark photon in e⁺e⁻ collisions at BaBar, Phys. Rev. Lett. 113 (2014) 201801, arXiv:1406.2980.
- BESIII, M. Ablikim, Search for a light CP-odd Higgs boson in radiative decays of J/, Phys. Rev. D93 (2016), no. 5 052005, arXiv:1510.01641.
- [22] LHCb collaboration, R. Aaij *et al.*, Search for rare $B^0_{(s)} \rightarrow \mu^+ \mu^- \mu^+ \mu^-$ decays, Phys. Rev. Lett. **110** (2013) 211801, arXiv:1303.1092.
- [23] LHCb collaboration, R. Aaij *et al.*, Search for hidden-sector bosons in $B^0 \to K^{*0} \mu^+ \mu^$ decays, Phys. Rev. Lett. **115** (2015) 161802, arXiv:1508.04094.
- [24] LHCb collaboration, R. Aaij *et al.*, Search for long-lived scalar particles in $\mathcal{B}^+ \to K^+\chi(\mu^+\mu^-)$ decays, Phys. Rev. **D95** (2017) 071101, arXiv:1612.07818.

- [25] LHCb collaboration, R. Aaij et al., Search for dark photons produced in 13 TeV pp collisions, Phys. Rev. Lett. 120 (2018) 061801, arXiv:1710.02867.
- [26] H.-K. Park, Search for the HyperCP event at the LHCb experiment, JHEP 10 (2010) 052.
- [27] L. Breiman, J. H. Friedman, R. A. Olshen, and C. J. Stone, *Classification and regression trees*, Wadsworth international group, Belmont, California, USA, 1984.
- [28] Y. Freund and R. E. Schapire, A decision-theoretic generalization of on-line learning and an application to boosting, J. Comput. Syst. Sci. 55 (1997) 119.
- [29] LHCb collaboration, A. A. Alves Jr. et al., The LHCb detector at the LHC, JINST 3 (2008) S08005.
- [30] LHCb collaboration, R. Aaij et al., LHCb detector performance, Int. J. Mod. Phys. A30 (2015) 1530022, arXiv:1412.6352.
- [31] R. Aaij et al., The LHCb trigger and its performance in 2011, JINST 8 (2013) P04022, arXiv:1211.3055.
- [32] T. Sjöstrand, S. Mrenna, and P. Skands, *PYTHIA 6.4 physics and manual*, JHEP 05 (2006) 026, arXiv:hep-ph/0603175; T. Sjöstrand, S. Mrenna, and P. Skands, A brief introduction to PYTHIA 8.1, Comput. Phys. Commun. 178 (2008) 852, arXiv:0710.3820.
- [33] I. Belyaev et al., Handling of the generation of primary events in Gauss, the LHCb simulation framework, J. Phys. Conf. Ser. **331** (2011) 032047.
- [34] D. J. Lange, The EvtGen particle decay simulation package, Nucl. Instrum. Meth. A462 (2001) 152.
- [35] P. Golonka and Z. Was, PHOTOS Monte Carlo: A precision tool for QED corrections in Z and W decays, Eur. Phys. J. C45 (2006) 97, arXiv:hep-ph/0506026.
- [36] Geant4 collaboration, J. Allison et al., Geant4 developments and applications, IEEE Trans. Nucl. Sci. 53 (2006) 270; Geant4 collaboration, S. Agostinelli et al., Geant4: A simulation toolkit, Nucl. Instrum. Meth. A506 (2003) 250.
- [37] M. Clemencic et al., The LHCb simulation application, Gauss: Design, evolution and experience, J. Phys. Conf. Ser. 331 (2011) 032023.
- [38] Particle Data Group, C. Patrignani et al., Review of particle physics, Chin. Phys. C40 (2016) 100001.
- [39] G. Punzi, Sensitivity of searches for new signals and its optimization, in Statistical Problems in Particle Physics, Astrophysics, and Cosmology (L. Lyons, R. Mount, and R. Reitmeyer, eds.), p. 79, 2003. arXiv:physics/0308063.
- [40] E. Govorkova, Study of π^0/γ efficiency using B meson decays in the LHCb experiment, Phys. Atom. Nucl. **79** (2016) 1474, arXiv:1505.02960.

- [41] ARGUS collaboration, H. Albrecht *et al.*, Search for hadronic $b \rightarrow u$ decays, Phys. Lett. **B241** (1990) 278.
- [42] CLEO collaboration, S. Dobbs et al., Measurement of absolute hadronic branching fractions of D mesons and e⁺e⁻ → DD̄ cross-sections at the ψ(3770), Phys. Rev. D76 (2007) 112001, arXiv:0709.3783.
- [43] D. Martínez Santos and F. Dupertuis, Mass distributions marginalized over per-event errors, Nucl. Instrum. Meth. A764 (2014) 150, arXiv:1312.5000.
- [44] S. S. Wilks, The large-sample distribution of the likelihood ratio for testing composite hypotheses, Ann. Math. Stat. 9 (1938) 60.
- [45] M. Pivk and F. R. Le Diberder, sPlot: A statistical tool to unfold data distributions, Nucl. Instrum. Meth. A555 (2005) 356, arXiv:physics/0402083.
- [46] See Supplemental Material to this Letter for additional plots and tables.
- [47] M. Williams, Searching for a particle of unknown mass and lifetime in the presence of an unknown non-monotonic background, JINST 10 (2015) P06002, arXiv:1503.04767.
- [48] A. L. Read, Presentation of search results: The CL_S technique, J. Phys. **G28** (2002) 2693.

LHCb collaboration

R. Aaij⁴⁰, B. Adeva³⁹, M. Adinolfi⁴⁸, Z. Ajaltouni⁵, S. Akar⁵⁹, J. Albrecht¹⁰, F. Alessio⁴⁰, M. Alexander⁵³, A. Alfonso Albero³⁸, S. Ali⁴³, G. Alkhazov³¹, P. Alvarez Cartelle⁵⁵, A.A. Alves Jr⁵⁹, S. Amato², S. Amerio²³, Y. Amhis⁷, L. An³, L. Anderlini¹⁸, G. Andreassi⁴¹, M. Andreotti^{17,g}, J.E. Andrews⁶⁰, R.B. Appleby⁵⁶, F. Archilli⁴³, P. d'Argent¹², J. Arnau Romeu⁶, A. Artamonov³⁷, M. Artuso⁶¹, E. Aslanides⁶, M. Atzeni⁴², G. Auriemma²⁶, M. Baalouch⁵, I. Babuschkin⁵⁶, S. Bachmann¹², J.J. Back⁵⁰, A. Badalov^{38,m}, C. Baesso⁶², S. Baker⁵⁵, V. Balagura^{7,b}, W. Baldini¹⁷, A. Baranov³⁵, R.J. Barlow⁵⁶, C. Barschel⁴⁰, S. Barsuk⁷, W. Barter⁵⁶, F. Baryshnikov³², V. Batozskaya²⁹, V. Battista⁴¹, A. Bay⁴¹, L. Beaucourt⁴, J. Beddow⁵³, F. Bedeschi²⁴, I. Bediaga¹, A. Beiter⁶¹, L.J. Bel⁴³, N. Beliv⁶³, V. Bellee⁴¹, N. Belloli^{21,i}, K. Belous³⁷, I. Belyaev^{32,40}, E. Ben-Haim⁸, G. Bencivenni¹⁹, S. Benson⁴³, S. Beranek⁹, A. Berezhnoy³³, R. Bernet⁴², D. Berninghoff¹², E. Bertholet⁸, A. Bertolin²³, C. Betancourt⁴², F. Betti¹⁵, M.O. Bettler⁴⁰, M. van Beuzekom⁴³, Ia. Bezshyiko⁴², S. Bifani⁴⁷, P. Billoir⁸, A. Birnkraut¹⁰, A. Bizzeti^{18,u}, M. Bjørn⁵⁷, T. Blake⁵⁰, F. Blanc⁴¹, S. Blusk⁶¹, V. Bocci²⁶, T. Boettcher⁵⁸, A. Bondar^{36,w}, N. Bondar³¹, I. Bordyuzhin³², S. Borghi^{56,40}, M. Borisyak³⁵, M. Borsato³⁹, F. Bossu⁷, M. Boubdir⁹, T.J.V. Bowcock⁵⁴, E. Bowen⁴², C. Bozzi^{17,40}, S. Braun¹², J. Brodzicka²⁷, D. Brundu¹⁶, E. Buchanan⁴⁸, C. Burr⁵⁶, A. Bursche^{16, f}, J. Buytaert⁴⁰, W. Byczynski⁴⁰, S. Cadeddu¹⁶, H. Cai⁶⁴, R. Calabrese^{17, g}, R. Calladine⁴⁷, M. Calvi^{21,i}, M. Calvo Gomez^{38,m}, A. Camboni^{38,m}, P. Campana¹⁹, D.H. Campora Perez⁴⁰, L. Capriotti⁵⁶, A. Carbone^{15,e}, G. Carboni^{25,j}, R. Cardinale^{20,h}, A. Cardini¹⁶, P. Carniti^{21,i}, L. Carson⁵², K. Carvalho Akiba², G. Casse⁵⁴, L. Cassina²¹, M. Cattaneo⁴⁰, G. Cavallero^{20,40,h}, R. Cenci^{24,t}, D. Chamont⁷, M.G. Chapman⁴⁸, M. Charles⁸, Ph. Charpentier⁴⁰, G. Chatzikonstantinidis⁴⁷, M. Chefdeville⁴, S. Chen¹⁶, S.F. Cheung⁵⁷, S.-G. Chitic⁴⁰, V. Chobanova³⁹, M. Chrzaszcz⁴², A. Chubykin³¹, P. Ciambrone¹⁹, X. Cid Vidal³⁹, G. Ciezarek⁴⁰, P.E.L. Clarke⁵², M. Clemencic⁴⁰, H.V. Cliff⁴⁹, J. Closier⁴⁰, V. Coco⁴⁰, J. Cogan⁶, E. Cogneras⁵, V. Cogoni^{16,f}, L. Cojocariu³⁰, P. Collins⁴⁰, T. Colombo⁴⁰, A. Comerma-Montells¹², A. Contu¹⁶, G. Coombs⁴⁰, S. Coquereau³⁸, G. Corti⁴⁰, M. Corvo^{17,g}, C.M. Costa Sobral⁵⁰, B. Couturier⁴⁰, G.A. Cowan⁵², D.C. Craik⁵⁸, A. Crocombe⁵⁰, M. Cruz Torres¹, R. Currie⁵², C. D'Ambrosio⁴⁰, F. Da Cunha Marinho², C.L. Da Silva⁷³, E. Dall'Occo⁴³, J. Dalseno⁴⁸, A. Davis³, O. De Aguiar Francisco⁴⁰, K. De Bruyn⁴⁰, S. De Capua⁵⁶, M. De Cian¹², J.M. De Miranda¹, L. De Paula², M. De Serio¹⁴, d, P. De Simone¹⁹, C.T. Dean⁵³, D. Decamp⁴, L. Del Buono⁸, H.-P. Dembinski¹¹, M. Demmer¹⁰, A. Dendek²⁸, D. Derkach³⁵, O. Deschamps⁵, F. Dettori⁵⁴, B. Dey⁶⁵, A. Di Canto⁴⁰, P. Di Nezza¹⁹, H. Dijkstra⁴⁰, F. Dordei⁴⁰, M. Dorigo⁴⁰, A. Dosil Suárez³⁹, L. Douglas⁵³, A. Dovbnya⁴⁵, K. Dreimanis⁵⁴, L. Dufour⁴³, G. Dujany⁸, P. Durante⁴⁰, J.M. Durham⁷³, D. Dutta⁵⁶, R. Dzhelyadin³⁷, M. Dziewiecki¹², A. Dziurda⁴⁰, A. Dzyuba³¹, S. Easo⁵¹, U. Egede⁵⁵, V. Egorychev³², S. Eidelman^{36,w}, S. Eisenhardt⁵², U. Eitschberger¹⁰, R. Ekelhof¹⁰, L. Eklund⁵³, S. Ely⁶¹, S. Esen¹², H.M. Evans⁴⁹, T. Evans⁵⁷, A. Falabella¹⁵, N. Farley⁴⁷, S. Farry⁵⁴, D. Fazzini^{21,i}, L. Federici²⁵, D. Ferguson⁵², G. Fernandez³⁸, P. Fernandez Declara⁴⁰, A. Fernandez Prieto³⁹, F. Ferrari¹⁵, L. Ferreira Lopes⁴¹, F. Ferreira Rodrigues², M. Ferro-Luzzi⁴⁰, S. Filippov³⁴, R.A. Fini¹⁴, M. Fiorini^{17,g}, M. Firlej²⁸, C. Fitzpatrick⁴¹, T. Fiutowski²⁸, F. Fleuret^{7,b}, M. Fontana^{16,40}, F. Fontanelli^{20,h}, R. Forty⁴⁰, V. Franco Lima⁵⁴, M. Frank⁴⁰, C. Frei⁴⁰, J. Fu^{22,q}, W. Funk⁴⁰, E. Furfaro^{25,j}, C. Färber⁴⁰, E. Gabriel⁵², A. Gallas Torreira³⁹, D. Galli^{15,e}, S. Gallorini²³, S. Gambetta⁵², M. Gandelman², P. Gandini²², Y. Gao³, L.M. Garcia Martin⁷¹, J. García Pardiñas³⁹, J. Garra Tico⁴⁹, L. Garrido³⁸, D. Gascon³⁸, C. Gaspar⁴⁰, L. Gavardi¹⁰, G. Gazzoni⁵, D. Gerick¹², E. Gersabeck⁵⁶, M. Gersabeck⁵⁶, T. Gershon⁵⁰, Ph. Ghez⁴, S. Giani⁴¹, V. Gibson⁴⁹, O.G. Girard⁴¹, L. Giubega³⁰, K. Gizdov⁵², V.V. Gligorov⁸, D. Golubkov³², A. Golutvin^{55,69}, A. Gomes^{1,a}, I.V. Gorelov³³, C. Gotti^{21,i}, E. Govorkova⁴³, J.P. Grabowski¹², R. Graciani Diaz³⁸,

L.A. Granado Cardoso⁴⁰, E. Graugés³⁸, E. Graverini⁴², G. Graziani¹⁸, A. Grecu³⁰, R. Greim⁹, P. Griffith¹⁶, L. Grillo⁵⁶, L. Gruber⁴⁰, B.R. Gruberg Cazon⁵⁷, O. Grünberg⁶⁷, E. Gushchin³⁴, Yu. Guz³⁷, T. Gys⁴⁰, C. Göbel⁶², T. Hadavizadeh⁵⁷, C. Hadjivasiliou⁵, G. Haefeli⁴¹, C. Haen⁴⁰, S.C. Haines⁴⁹, B. Hamilton⁶⁰, X. Han¹², T.H. Hancock⁵⁷, S. Hansmann-Menzemer¹², N. Harnew⁵⁷, S.T. Harnew⁴⁸, C. Hasse⁴⁰, M. Hatch⁴⁰, J. He⁶³, M. Hecker⁵⁵, K. Heinicke¹⁰, A. Heister⁹, K. Hennessy⁵⁴, P. Henrard⁵, L. Henry⁷¹, E. van Herwijnen⁴⁰, M. Heß⁶⁷, A. Hicheur², D. Hill⁵⁷, P.H. Hopchev⁴¹, W. Hu⁶⁵, W. Huang⁶³, Z.C. Huard⁵⁹, W. Hulsbergen⁴³, T. Humair⁵⁵, M. Hushchyn³⁵, D. Hutchcroft⁵⁴, P. Ibis¹⁰, M. Idzik²⁸, P. Ilten⁴⁷, R. Jacobsson⁴⁰, J. Jalocha⁵⁷, E. Jans⁴³, A. Jawahery⁶⁰, F. Jiang³, M. John⁵⁷, D. Johnson⁴⁰, C.R. Jones⁴⁹, C. Joram⁴⁰, B. Jost⁴⁰, N. Jurik⁵⁷, S. Kandybei⁴⁵, M. Karacson⁴⁰, J.M. Kariuki⁴⁸, S. Karodia⁵³, N. Kazeev³⁵, M. Kecke¹², F. Keizer⁴⁹, M. Kelsey⁶¹, M. Kenzie⁴⁹, T. Ketel⁴⁴, E. Khairullin³⁵, B. Khanji¹², C. Khurewathanakul⁴¹, K.E. Kim⁶¹, T. Kirn⁹, S. Klaver¹⁹, K. Klimaszewski²⁹, T. Klimkovich¹¹, S. Koliiev⁴⁶, M. Kolpin¹², R. Kopecna¹², P. Koppenburg⁴³, A. Kosmyntseva³², S. Kotriakhova³¹, M. Kozeiha⁵, L. Kravchuk³⁴, M. Kreps⁵⁰, F. Kress⁵⁵, P. Krokovny^{36,w}, W. Krzemien²⁹, W. Kucewicz^{27,l}, M. Kucharczyk²⁷, V. Kudryavtsev^{36,w}, A.K. Kuonen⁴¹, T. Kvaratskheliya^{32,40}, D. Lacarrere⁴⁰, G. Lafferty⁵⁶, A. Lai¹⁶, G. Lanfranchi¹⁹, C. Langenbruch⁹, T. Latham⁵⁰, C. Lazzeroni⁴⁷, R. Le Gac⁶, A. Leflat^{33,40}, J. Lefrançois⁷, R. Lefèvre⁵, F. Lemaitre⁴⁰, E. Lemos Cid³⁹, O. Leroy⁶, T. Lesiak²⁷, B. Leverington¹², P.-R. Li⁶³, T. Li³, Y. Li⁷, Z. Li⁶¹, X. Liang⁶¹, T. Likhomanenko⁶⁸, R. Lindner⁴⁰, F. Lionetto⁴², V. Lisovskyi⁷, X. Liu³, D. Loh⁵⁰, A. Loi¹⁶, I. Longstaff⁵³, J.H. Lopes², D. Lucchesi^{23,o}, M. Lucio Martinez³⁹, H. Luo⁵², A. Lupato²³, E. Luppi^{17,g}, O. Lupton⁴⁰, A. Lusiani²⁴, X. Lyu⁶³, F. Machefert⁷, F. Maciuc³⁰, V. Macko⁴¹, P. Mackowiak¹⁰, S. Maddrell-Mander⁴⁸, O. Maev^{31,40}, K. Maguire⁵⁶, D. Maisuzenko³¹, M.W. Majewski²⁸, S. Malde⁵⁷, B. Malecki²⁷, A. Malinin⁶⁸, T. Maltsev^{36,w}, G. Manca^{16,f}, G. Mancinelli⁶, D. Marangotto^{22,q}, J. Maratas^{5,v}, J.F. Marchand⁴, U. Marconi¹⁵, C. Marin Benito³⁸, M. Marinangeli⁴¹, P. Marino⁴¹, J. Marks¹², G. Martellotti²⁶, M. Martin⁶, M. Martinelli⁴¹, D. Martinez Santos³⁹, F. Martinez Vidal⁷¹, A. Massafferri¹, R. Matev⁴⁰, A. Mathad⁵⁰, Z. Mathe⁴⁰, C. Matteuzzi²¹, A. Mauri⁴², E. Maurice^{7,b}, B. Maurin⁴¹, A. Mazurov⁴⁷, M. McCann^{55,40}, A. McNab⁵⁶, R. McNulty¹³, J.V. Mead⁵⁴, B. Meadows⁵⁹, C. Meaux⁶, F. Meier¹⁰, N. Meinert⁶⁷, D. Melnychuk²⁹, M. Merk⁴³, A. Merli^{22,40,q}, E. Michielin²³, D.A. Milanes⁶⁶, E. Millard⁵⁰, M.-N. Minard⁴, L. Minzoni¹⁷, D.S. Mitzel¹², A. Mogini⁸, J. Molina Rodriguez¹, T. Mombächer¹⁰, I.A. Monroy⁶⁶, S. Monteil⁵, M. Morandin²³, M.J. Morello^{24,t}, O. Morgunova⁶⁸, J. Moron²⁸, A.B. Morris⁵², R. Mountain⁶¹, F. Muheim⁵², M. Mulder⁴³, D. Müller⁴⁰, J. Müller¹⁰, K. Müller⁴², V. Müller¹⁰, P. Naik⁴⁸, T. Nakada⁴¹, R. Nandakumar⁵¹, A. Nandi⁵⁷, I. Nasteva², M. Needham⁵², N. Neri^{22,40}, S. Neubert¹², N. Neufeld⁴⁰, M. Neuner¹², T.D. Nguyen⁴¹, C. Nguyen-Mau^{41,n}, S. Nieswand⁹, R. Niet¹⁰, N. Nikitin³³, T. Nikodem¹², A. Nogay⁶⁸, D.P. O'Hanlon⁵⁰, A. Oblakowska-Mucha²⁸, V. Obraztsov³⁷, S. Ogilvy¹⁹, R. Oldeman^{16,f}, C.J.G. Onderwater⁷², A. Ossowska²⁷, J.M. Otalora Goicochea², P. Owen⁴², A. Oyanguren⁷¹, P.R. Pais⁴¹, A. Palano¹⁴, M. Palutan^{19,40}, G. Panshin⁷⁰, A. Papanestis⁵¹, M. Pappagallo⁵², L.L. Pappalardo^{17,g}, W. Parker⁶⁰, C. Parkes⁵⁶, G. Passaleva^{18,40}, A. Pastore^{14,d}, M. Patel⁵⁵, C. Patrignani^{15,e}, A. Pearce⁴⁰, A. Pellegrino⁴³, G. Penso²⁶, M. Pepe Altarelli⁴⁰, S. Perazzini⁴⁰, D. Pereima³², P. Perret⁵, L. Pescatore⁴¹, K. Petridis⁴⁸, A. Petrolini^{20,h}, A. Petrov⁶⁸, M. Petruzzo^{22,q}, E. Picatoste Olloqui³⁸, B. Pietrzyk⁴, G. Pietrzyk⁴¹, M. Pikies²⁷, D. Pinci²⁶, F. Pisani⁴⁰, A. Pistone^{20,h}, A. Piucci¹², V. Placinta³⁰, S. Playfer⁵², M. Plo Casasus³⁹, F. Polci⁸, M. Poli Lener¹⁹, A. Poluektov⁵⁰, I. Polyakov⁶¹, E. Polycarpo², G.J. Pomery⁴⁸, S. Ponce⁴⁰, A. Popov³⁷, D. Popov^{11,40}, S. Poslavskii³⁷, C. Potterat², E. Price⁴⁸, J. Prisciandaro³⁹, C. Prouve⁴⁸, V. Pugatch⁴⁶, A. Puig Navarro⁴², H. Pullen⁵⁷, G. Punzi^{24,p}, W. Qian⁵⁰, J. Qin⁶³, R. Quagliani⁸, B. Quintana⁵, B. Rachwal²⁸, J.H. Rademacker⁴⁸, M. Rama²⁴, M. Ramos Pernas³⁹, M.S. Rangel², I. Raniuk^{45,†}, F. Ratnikov^{35,x}, G. Raven⁴⁴, M. Ravonel Salzgeber⁴⁰, M. Reboud⁴, F. Redi⁴¹, S. Reichert¹⁰,

A.C. dos Reis¹, C. Remon Alepuz⁷¹, V. Renaudin⁷, S. Ricciardi⁵¹, S. Richards⁴⁸, M. Rihl⁴⁰,

K. Rinnert⁵⁴, P. Robbe⁷, A. Robert⁸, A.B. Rodrigues⁴¹, E. Rodrigues⁵⁹, J.A. Rodriguez Lopez⁶⁶, A. Rogozhnikov³⁵, S. Roiser⁴⁰, A. Rollings⁵⁷, V. Romanovskiy³⁷, A. Romero Vidal^{39,40}, M. Rotondo¹⁹, M.S. Rudolph⁶¹, T. Ruf⁴⁰, P. Ruiz Valls⁷¹, J. Ruiz Vidal⁷¹, J.J. Saborido Silva³⁹, E. Sadykhov³², N. Sagidova³¹, B. Saitta^{16, f}, V. Salustino Guimaraes⁶², C. Sanchez Mayordomo⁷¹, B. Sanmartin Sedes³⁹, R. Santacesaria²⁶, C. Santamarina Rios³⁹, M. Santimaria¹⁹, E. Santovetti^{25,j}, G. Sarpis⁵⁶, A. Sarti^{19,k}, C. Satriano^{26,s}, A. Satta²⁵, D.M. Saunders⁴⁸, D. Savrina^{32,33}, S. Schael⁹, M. Schellenberg¹⁰, M. Schiller⁵³, H. Schindler⁴⁰, M. Schmelling¹¹, T. Schmelzer¹⁰, B. Schmidt⁴⁰, O. Schneider⁴¹, A. Schopper⁴⁰, H.F. Schreiner⁵⁹, M. Schubiger⁴¹, M.H. Schune⁷, R. Schwemmer⁴⁰, B. Sciascia¹⁹, A. Sciubba^{26,k}, A. Semennikov³², E.S. Sepulveda⁸, A. Sergi⁴⁷, N. Serra⁴², J. Serrano⁶, L. Sestini²³, P. Seyfert⁴⁰, M. Shapkin³⁷, Y. Shcheglov³¹, T. Shears⁵⁴, L. Shekhtman^{36,w}, V. Shevchenko⁶⁸, B.G. Siddi¹⁷, R. Silva Coutinho⁴², L. Silva de Oliveira², G. Simi^{23,o}, S. Simone^{14,d}, M. Sirendi⁴⁹, N. Skidmore⁴⁸, T. Skwarnicki⁶¹, I.T. Smith⁵², J. Smith⁴⁹, M. Smith⁵⁵, l. Soares Lavra¹, M.D. Sokoloff⁵⁹, F.J.P. Soler⁵³, B. Souza De Paula², B. Spaan¹⁰, P. Spradlin⁵³, F. Stagni⁴⁰, M. Stahl¹², S. Stahl⁴⁰, P. Stefko⁴¹, S. Stefkova⁵⁵, O. Steinkamp⁴², S. Stemmle¹², O. Stenyakin³⁷, M. Stepanova³¹, H. Stevens¹⁰, S. Stone⁶¹, B. Storaci⁴², S. Stracka^{24,p}, M.E. Stramaglia⁴¹, M. Straticiuc³⁰, U. Straumann⁴², S. Strokov⁷⁰, J. Sun³, L. Sun⁶⁴, K. Swientek²⁸, V. Syropoulos⁴⁴, T. Szumlak²⁸, M. Szymanski⁶³, S. T'Jampens⁴, A. Tayduganov⁶, T. Tekampe¹⁰, G. Tellarini^{17,g}, F. Teubert⁴⁰, E. Thomas⁴⁰, J. van Tilburg⁴³, M.J. Tilley⁵⁵, V. Tisserand⁵, M. Tobin⁴¹, S. Tolk⁴⁹, L. Tomassetti^{17,g}, D. Tonelli²⁴, R. Tourinho Jadallah Aoude¹, E. Tournefier⁴, M. Traill⁵³, M.T. Tran⁴¹, M. Tresch⁴², A. Trisovic⁴⁹, A. Tsaregorodtsev⁶, P. Tsopelas⁴³, A. Tully⁴⁹, N. Tuning^{43,40}, A. Ukleja²⁹, A. Usachov⁷, A. Ustvuzhanin³⁵, U. Uwer¹², C. Vacca^{16,f}, A. Vagner⁷⁰, V. Vagnoni^{15,40}, A. Valassi⁴⁰, S. Valat⁴⁰, G. Valenti¹⁵, R. Vazquez Gomez⁴⁰, P. Vazquez Regueiro³⁹, S. Vecchi¹⁷, M. van Veghel⁴³, J.J. Velthuis⁴⁸, M. Veltri^{18,r}, G. Veneziano⁵⁷, A. Venkateswaran⁶¹, T.A. Verlage⁹, M. Vernet⁵, M. Vesterinen⁵⁷, J.V. Viana Barbosa⁴⁰, D. Vieira⁶³, M. Vieites Diaz³⁹, H. Viemann⁶⁷, X. Vilasis-Cardona^{38,m}, M. Vitti⁴⁹, V. Volkov³³, A. Vollhardt⁴², B. Voneki⁴⁰, A. Vorobyev³¹, V. Vorobyev^{36,w}, C. Voß⁹, J.A. de Vries⁴³, C. Vázquez Sierra⁴³, R. Waldi⁶⁷, J. Walsh²⁴, J. Wang⁶¹, Y. Wang⁶⁵, D.R. Ward⁴⁹, H.M. Wark⁵⁴, N.K. Watson⁴⁷, D. Websdale⁵⁵, A. Weiden⁴², C. Weisser⁵⁸, M. Whitehead⁴⁰, J. Wicht⁵⁰, G. Wilkinson⁵⁷, M. Wilkinson⁶¹, M. Williams⁵⁶, M. Williams⁵⁸, T. Williams⁴⁷, F.F. Wilson^{51,40}, J. Wimberley⁶⁰, M. Winn⁷, J. Wishahi¹⁰, W. Wislicki²⁹, M. Witek²⁷, G. Wormser⁷, S.A. Wotton⁴⁹, K. Wyllie⁴⁰, Y. Xie⁶⁵, M. Xu⁶⁵, Q. Xu⁶³, Z. Xu³, Z. Xu⁴, Z. Yang³, Z. Yang⁶⁰, Y. Yao⁶¹, H. Yin⁶⁵, J. Yu⁶⁵, X. Yuan⁶¹, O. Yushchenko³⁷, K.A. Zarebski⁴⁷, M. Zavertyaev^{11,c}, L. Zhang³, Y. Zhang⁷, A. Zhelezov¹², Y. Zheng⁶³, X. Zhu³, V. Zhukov^{9,33}, J.B. Zonneveld⁵², S. Zucchelli¹⁵.

¹Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

² Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

³Center for High Energy Physics, Tsinghua University, Beijing, China

⁴ Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France

⁵Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France

⁶Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France

⁷LAL, Univ. Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, Orsay, France

⁸LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France

⁹I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany

¹⁰ Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

¹¹Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

¹²Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

¹³School of Physics, University College Dublin, Dublin, Ireland

¹⁴Sezione INFN di Bari, Bari, Italy

¹⁵Sezione INFN di Bologna, Bologna, Italy

¹⁶Sezione INFN di Cagliari, Cagliari, Italy

¹⁷ Universita e INFN, Ferrara, Ferrara, Italy

¹⁸Sezione INFN di Firenze, Firenze, Italy

¹⁹Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy

²⁰Sezione INFN di Genova, Genova, Italy

²¹Sezione INFN di Milano Bicocca, Milano, Italy

²²Sezione di Milano, Milano, Italy

²³Sezione INFN di Padova, Padova, Italy

²⁴Sezione INFN di Pisa, Pisa, Italy

²⁵Sezione INFN di Roma Tor Vergata, Roma, Italy

²⁶Sezione INFN di Roma La Sapienza, Roma, Italy

²⁷Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland

²⁸AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland

²⁹National Center for Nuclear Research (NCBJ), Warsaw, Poland

³⁰ Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania ³¹ Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia

³²Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia

³³Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia

³⁴Institute for Nuclear Research of the Russian Academy of Sciences (INR RAS), Moscow, Russia

³⁵ Yandex School of Data Analysis, Moscow, Russia

³⁶Budker Institute of Nuclear Physics (SB RAS), Novosibirsk, Russia

³⁷Institute for High Energy Physics (IHEP), Protvino, Russia

³⁸ICCUB, Universitat de Barcelona, Barcelona, Spain

³⁹Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain

⁴⁰ European Organization for Nuclear Research (CERN), Geneva, Switzerland

⁴¹Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland

⁴² Physik-Institut, Universität Zürich, Zürich, Switzerland

⁴³Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands

⁴⁴Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands

⁴⁵NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine

⁴⁶Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine

⁴⁷ University of Birmingham, Birmingham, United Kingdom

⁴⁸H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom

⁴⁹Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom

⁵⁰Department of Physics, University of Warwick, Coventry, United Kingdom

⁵¹STFC Rutherford Appleton Laboratory, Didcot, United Kingdom

⁵²School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom

⁵³School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom

⁵⁴Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom

⁵⁵Imperial College London, London, United Kingdom

⁵⁶School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom

⁵⁷Department of Physics, University of Oxford, Oxford, United Kingdom

⁵⁸ Massachusetts Institute of Technology, Cambridge, MA, United States

⁵⁹ University of Cincinnati, Cincinnati, OH, United States

⁶⁰ University of Maryland, College Park, MD, United States

⁶¹Syracuse University, Syracuse, NY, United States

⁶²Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to ² ⁶³University of Chinese Academy of Sciences, Beijing, China, associated to ³

⁶⁴School of Physics and Technology, Wuhan University, Wuhan, China, associated to ³

⁶⁵Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China, associated to ³

⁶⁶Departamento de Fisica, Universidad Nacional de Colombia, Bogota, Colombia, associated to ⁸

⁶⁷Institut für Physik, Universität Rostock, Rostock, Germany, associated to ¹²

⁶⁸National Research Centre Kurchatov Institute, Moscow, Russia, associated to ³²

⁶⁹National University of Science and Technology MISIS, Moscow, Russia, associated to ³²

⁷⁰National Research Tomsk Polytechnic University, Tomsk, Russia, associated to ³²

⁷¹Instituto de Fisica Corpuscular, Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain, associated to ³⁸

⁷² Van Swinderen Institute, University of Groningen, Groningen, The Netherlands, associated to ⁴³
 ⁷³ Los Alamos National Laboratory (LANL), Los Alamos, United States, associated to ⁶¹

^a Universidade Federal do Triângulo Mineiro (UFTM), Uberaba-MG, Brazil

^bLaboratoire Leprince-Ringuet, Palaiseau, France

^cP.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia

^d Università di Bari, Bari, Italy

^e Università di Bologna, Bologna, Italy

^f Università di Cagliari, Cagliari, Italy

^g Università di Ferrara, Ferrara, Italy

^h Università di Genova, Genova, Italy

ⁱ Università di Milano Bicocca, Milano, Italy

^j Università di Roma Tor Vergata, Roma, Italy

^k Università di Roma La Sapienza, Roma, Italy

¹AGH - University of Science and Technology, Faculty of Computer Science, Electronics and

Telecommunications, Kraków, Poland

^mLIFAELS, La Salle, Universitat Ramon Llull, Barcelona, Spain

ⁿHanoi University of Science, Hanoi, Vietnam

^o Università di Padova, Padova, Italy

^pUniversità di Pisa, Pisa, Italy

^qUniversità degli Studi di Milano, Milano, Italy

^r Università di Urbino, Urbino, Italy

^s Università della Basilicata, Potenza, Italy

^tScuola Normale Superiore, Pisa, Italy

 u Università di Modena e Reggio Emilia, Modena, Italy

 v Iligan Institute of Technology (IIT), Iligan, Philippines

 w Novosibirsk State University, Novosibirsk, Russia

^x National Research University Higher School of Economics, Moscow, Russia

 $^{\dagger}Deceased$