



**HAL**  
open science

# Equivalent partial differential equations of a lattice Boltzmann scheme

François Dubois

► **To cite this version:**

François Dubois. Equivalent partial differential equations of a lattice Boltzmann scheme. *Computers & Mathematics with Applications*, 2008, 55 (7), pp.1441 - 1449. 10.1016/j.camwa.2007.08.003 . hal-01809947

**HAL Id: hal-01809947**

**<https://hal.science/hal-01809947>**

Submitted on 7 Jun 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Equivalent partial differential equations of a lattice Boltzmann scheme

François Dubois<sup>ab</sup>

<sup>a</sup> *Dpt. of Mathematics, University Paris-Sud, Bât. 425, F-91405 Orsay, France.*

<sup>b</sup> *Conservatoire National des Arts et Métiers, LMSSC laboratory, F-75003 Paris, France.*

05 septembre 2007 \*

**Keywords:** Chapman-Enskog expansion, Taylor formula

**AMS (MSC2010) classification:** 76M28.

## Abstract

We show that when we formulate the lattice Boltzmann equation with a small time step  $\Delta t$  and an associated space scale  $\Delta x$ , a Taylor expansion joined with the so-called equivalent equation methodology leads to establish macroscopic fluid equations as a formal limit. We recover the Euler equations of gas dynamics at the first order and the compressible Navier-Stokes equations at the second order.

## 1) Discrete geometry

• We denote by  $d$  the dimension of space and by  $\mathcal{L}$  a regular  $d$ -dimensional lattice. Such a lattice is composed by a set  $\mathcal{L}^0$  of nodes or vertices and a set  $\mathcal{L}^1$  of links or edges between two vertices. From a practical point of view, given a vertex  $x$ , there exists a set  $V(x)$  of neighbouring nodes, including the node  $x$  itself. We consider here that the lattice  $\mathcal{L}$  is parametrized by a space step  $\Delta x > 0$ . For the fundamental example called D2Q9 (see *e.g.* Lallemand and Luo, 2000), the set  $V(x)$  is given with the help of the family of vectors  $(e_j)_{0 \leq j \leq J}$  defined by  $J = 8$ ,

$$(1.1) \quad (e_j) = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ -1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ -1 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right\}$$

and the vicinity

$$(1.2) \quad V(x) = \{x + \Delta x e_j, 0 \leq j \leq J\}.$$

---

\* Contribution published in *Computers & Mathematics with Applications*, volume 55, pages 1441-1449, 2008, doi <https://doi.org/10.1016/j.camwa.2007.08.003>. Edition 07 June 2018.

- In the general case, we still suppose that the equation (1.2) holds but we do not make any precise definition concerning the integer  $J$  and the nondimensionalized vectors  $(e_j)_{0 \leq j \leq J}$ . Nevertheless if  $x$  is a node of the lattice ( $x \in \mathcal{L}^0$ ), then  $y^j = x + \Delta x e_j$  is an other node of the lattice, *i.e.*  $y^j \in \mathcal{L}^0$ .

## 2) Lattice Boltzmann framework

- We introduce a time step  $\Delta t > 0$  and we suppose that the celerity  $\lambda$  defined according to

$$(2.1) \quad \lambda = \frac{\Delta x}{\Delta t}$$

remains fixed. Then we introduce a local velocity  $v_j$  in such a way that

$$(2.2) \quad \Delta t v_j = \Delta x e_j, \quad 0 \leq j \leq J.$$

In this  $d$ -dimensional framework we will denote by  $v_j^\alpha$  ( $1 \leq \alpha \leq d$ ) the Cartesian components of velocities  $v_j$ . Recall that if  $x$  is a node of the lattice, the point  $x + \Delta t v_j$  is also a node of the lattice:

$$(2.3) \quad x \in \mathcal{L}^0 \implies x + \Delta t v_j \in \mathcal{L}^0, \quad \forall j = 0, \dots, J.$$

- According to D’Humières (1992), the lattice Boltzmann scheme describes the dynamics of the density  $f^j(x, t)$  of particles of velocity  $v_j$  at the node  $x$  and for the discrete time  $t$ . We introduce the  $d + 1$  scalar “conservative variables”  $W(x, t)$  composed by the density  $\rho$  and the momentum  $q$ . Note that it is also possible to take into account the conservation of the total energy (see D’Humières’s article for example). We have

$$(2.4) \quad \rho(x, t) = \sum_{j=0}^J f^j(x, t) \equiv W^0(x, t)$$

$$(2.5) \quad q^\alpha(x, t) = \sum_{j=0}^J v_j^\alpha f^j(x, t) \equiv W^\alpha(x, t), \quad 1 \leq \alpha \leq d,$$

and

$$(2.6) \quad W(x, t) = (\rho(x, t), q^1(x, t), \dots, q^d(x, t)).$$

When a state  $W$  is given in space  $\mathbb{R}^{d+1}$ , a Gaussian (or any other choice) equilibrium distribution of particles is defined according to

$$(2.7) \quad f_{\text{eq}}^j = G^j(W), \quad 0 \leq j \leq J$$

in such a way that

$$(2.8) \quad \sum_{j=0}^J G^j(W) \equiv W^0, \quad \sum_{j=0}^J v_j^\alpha G^j(W) \equiv W^\alpha, \quad 1 \leq \alpha \leq d.$$

- Following D’Humières (1992), we introduce the “moment vector”  $m$  according to

$$(2.9) \quad m^k = \sum_{j=0}^J M_j^k f^j, \quad 0 \leq k \leq J.$$

For  $0 \leq i \leq d$ , the moments  $m^i$  are **identical** to the conservative variables:

$$(2.10) \quad m^0 \equiv \rho, \quad m^\alpha \equiv q^\alpha, \quad 1 \leq \alpha \leq d.$$

In other words, the matrix  $M$  satisfies

$$(2.11) \quad M_j^0 \equiv 1, \quad M_j^\alpha \equiv v_j^\alpha, \quad 0 \leq j \leq J, \quad 1 \leq \alpha \leq d.$$

We assume that vectors  $(e_j)_{0 \leq j \leq J}$  are chosen such that the  $(d+1) \times (J+1)$  matrix  $(M_{kj})_{0 \leq k \leq d, 0 \leq j \leq J}$  is of full rank. With this hypothesis, the conservative moments  $W$  introduced in relation (2.6) are independent variables.

- When a particle distribution  $f$  is given, the moments are evaluated according to (2.9). The matrix  $M$  is supposed to be invertible and the inverse relation takes the form:

$$(2.12) \quad f^j = \sum_{k=0}^J (M^{-1})_k^j m^k, \quad 0 \leq j \leq J.$$

When  $f_{\text{eq}}^j$  is determined according to the relation (2.7), the associated equilibrium moments  $m_{\text{eq}}^k$  are given simply according to (2.9), *i.e.* in this case

$$(2.13) \quad m_{\text{eq}}^k = \sum_{j=0}^J M_j^k f_{\text{eq}}^j, \quad 0 \leq k \leq J.$$

We remark also that by construction (relation (2.8)), we have

$$(2.14) \quad m_{\text{eq}}^i = m^i = W^i, \quad 0 \leq i \leq d.$$

### 3) Collision step

- The collision step is local in space and is naturally defined in the space of moments. If  $m^k(x, t)$  denotes the value of the  $k^{\text{th}}$  component of the moment vector  $m$  at position  $x$  and time  $t$ , the same component  $m_*^k(x, t)$  of the moment **after** the collision is trivial by construction for the conservative variables:

$$(3.1) \quad m_*^i(x, t) = m^i(x, t), \quad 0 \leq i \leq d.$$

For the non-conservative components of the moment vector, we **fix** the ratio  $s_k$  ( $k \geq d+1$ ) between the time step  $\Delta t$  and the relaxation time  $\tau_k$  of an underlying process:

$$(3.2) \quad s_k = \frac{\Delta t}{\tau_k}, \quad d+1 \leq k \leq J.$$

- Then  $m_*^k(x, t)$  after the collision is defined according to

$$(3.3) \quad m_*^k(x, t) = (1 - s_k) m^k(x, t) + s_k m_{\text{eq}}^k, \quad d+1 \leq k \leq J.$$

**Proposition 1.** Explicit Euler scheme.

The numerical scheme (3.3) is exactly the explicit Euler scheme relative to the continuous in time relaxation equation

$$(3.4) \quad \frac{d}{dt}(m^k - m_{\text{eq}}^k) + \frac{1}{\tau_k}(m^k - m_{\text{eq}}^k) = 0, \quad d+1 \leq k \leq J.$$

**Proof of Proposition 1.**

Following *e.g.* Strang (1986), we know that the explicit Euler scheme for the evolution (3.4) takes the form

$$(3.5) \quad \frac{1}{\Delta t} \left[ (m^k - m_{\text{eq}}^k)(t + \Delta t) - (m^k - m_{\text{eq}}^k)(t) \right] + \frac{1}{\tau_k} (m^k - m_{\text{eq}}^k)(t) = 0.$$

We have by construction the relation (3.1), that is  $m^i(t + \Delta t) = m^i(t)$  for  $0 \leq i \leq d$  with these notations. Then  $W(t + \Delta t) = W(t)$  and, due to the relation (2.7),  $f_{\text{eq}}^j(t + \Delta t) = f_{\text{eq}}^j(t)$  after the collision step for all the components  $j$  of the particle distribution. Due to (2.13), we deduce that  $m_{\text{eq}}^k(t + \Delta t) = m_{\text{eq}}^k(t)$  for all  $k \leq J$ . Thus the expression (3.5) takes the simpler form

$$(3.6) \quad \frac{1}{\Delta t} [m^k(t + \Delta t) - m^k(t)] + \frac{1}{\tau_k} (m^k - m_{\text{eq}}^k)(t) = 0,$$

which is exactly (3.3), except the change of notations:  $m^k(t + \Delta t)$  is replaced by  $m_*^k$ .  $\square$

- We remark also that the classical stability condition for the explicit Euler scheme (see again *e.g.* the book of Strang) takes the form

$$(3.7) \quad 0 \leq \Delta t \leq 2\tau_k.$$

We will suppose in the following that

$$(3.8) \quad 0 < s_k \leq 2, \quad d + 1 \leq k \leq J.$$

to put in evidence that the moments  $m^k$  are **not** conserved for index  $k$  greater than  $d + 1$ . We remark also that for the physically relevant Boltzmann equation, the relaxation times  $\tau_k$  have a physical sense. With the lattice Boltzmann scheme itself, these physical constants are no longer correctly approximated whereas the **ratios**  $s_k = \frac{\Delta t}{\tau_k}$  are supposed to be fixed in all what follows. Despite the usual ‘‘LBE’’ denomination, a lattice Boltzmann scheme is not a numerical method to approach the Boltzmann equation !

- The particle distribution  $f_*^j$  after the collision step follows the relation (2.12). We have precisely after the collision step

$$(3.9) \quad f_*^j = \sum_{k=0}^J (M^{-1})_k^j m_*^k, \quad 0 \leq j \leq J.$$

#### 4) Advection step

- The advection step of the lattice Boltzmann scheme claims that after the collision step, the particles having velocity  $v_j$  at position  $x$  go in one time step  $\Delta t$  to the  $j^{\text{th}}$  neighbouring vertex. Thus the particle density  $f^j(x + v_j \Delta t, t + \Delta t)$  at the new time step in the neighbouring vertex is equal to the previous particle density  $f_*^j(x, t)$  at the position  $x$  after the collision:

$$(4.1) \quad f^j(x + v_j \Delta t, t + \Delta t) = f_*^j(x, t).$$

We re-write this relation in term of the ‘‘arrival’’ node  $x + v_j \Delta t$ . We set  $\tilde{x} = x + v_j \Delta t$ , then we have  $x = \tilde{x} - v_j \Delta t$  and going back to the notation  $x$ , we write the relation (4.1) in the equivalent manner

$$(4.2) \quad f^j(x, t + \Delta t) = f_*^j(x - v_j \Delta t, t), \quad 0 \leq j \leq J, \quad x \in \mathcal{L}^0.$$

**Proposition 2.** Upwind scheme for the advection equation.

The scheme (4.2) for the advection step of the lattice Boltzmann method is nothing else than the explicit upwind scheme for the advection equation

$$(4.3) \quad \frac{\partial f^j}{\partial t} + v_j \cdot \nabla f^j = 0, \quad 0 \leq j \leq J,$$

with a so-called Courant-Friedrichs-Lewy number  $\sigma_j$  in the  $j^{\text{th}}$  direction of the lattice defined by

$$(4.4) \quad \sigma_j \equiv |v_j| \frac{\Delta t}{\Delta x |e_j|}$$

equal, due to the definition (2.2), to unity:  $\sigma_j = 1$ .

**Proof of Proposition 2.**

When the Courant-Friedrichs-Lewy number  $\sigma_j$  is equal to unity, it is classical (see *e.g.* Strang, 1986) that the upwind scheme is exact for the advection equation.  $\square$

## 5) Equivalent equation at zero order

- The lattice Boltzmann scheme is defined by the relations (2.4) to (2.9), (3.3) and (4.2). It is parametrized by the lattice step  $\Delta x$ , the matrix  $M$  linking the particle distribution  $f$  and the moment vector  $m$ , the choice of the conservative moments, the nonlinear equilibrium function  $G(\bullet)$ , the time step  $\Delta t$  and the ratios  $s_k$  between the time step and the collision time constants for nonequilibrium moments. In what follows, we fix the geometrical and topological structure of the lattice  $\mathcal{L}$ , we fix the matrix  $M$  and the equilibrium function  $G(\bullet)$ , we fix also the ratio  $\lambda$  defined in (2.1) and last but not least, we suppose that the parameters  $s_k$  for  $k \geq d + 1$  have a fixed value. Then the whole lattice Boltzmann scheme depends on the single parameter  $\Delta t$ .

- We explore now formally what are the partial differential equations associated with the Boltzmann numerical scheme, following the so-called “equivalent equation method” introduced and developed by Lerat-Peyret (1974) and Warming-Hyett (1974). This approach is based on the assumption, that a sufficiently smooth function exists which satisfies the difference equation at the grid points. This assumption gives formal responses to put in evidence partial differential equations that minimize the truncation errors of the numerical scheme. Nevertheless, we note here that this method of analysis fails to predict initial layers and boundary effects properly, as discussed by Griffiths and Sanz-Serna (1986) or Chang (1990). The idea of the calculus is to suppose that all the data are sufficiently regular and to expand all the variables with the Taylor formula.

**Proposition 3.** Taylor expansion at zero order.

With the lattice Boltzmann defined previously, we have

$$(5.1) \quad f^j(x, t) = f_{\text{eq}}^j(x, t) + \text{O}(\Delta t), \quad 0 \leq j \leq J,$$

$$(5.2) \quad f_*^j(x, t) = f_{\text{eq}}^j(x, t) + \text{O}(\Delta t), \quad 0 \leq j \leq J,$$

with  $f_{\text{eq}}^j$  defined from the conservative variables  $W$  according to the relation (2.7).

**Proof of Proposition 3.**

The key point is to expand the relation (4.2) relative to the infinitesimal  $\Delta t$ . We have on one hand

$$(5.3) \quad f^j(x, t + \Delta t) = f^j(x, t) + \text{O}(\Delta t)$$

and on the other hand

$$(5.4) \quad f_*^j(x - v_j \Delta t, t) = f_*^j(x, t) + \text{O}(\Delta t)$$

$$\text{Then } m_*^k(x, t) = \sum_{j=0}^J M_j^k f_*^j(x, t) = m^k(x, t) + \text{O}(\Delta t) \quad \text{and}$$

$$(5.5) \quad m_*^k(x, t) - m^k(x, t) = \text{O}(\Delta t).$$

But, due to (3.3), we have

$$(5.6) \quad m_*^k(x, t) - m^k(x, t) = -s_k (m^k(x, t) - m_{\text{eq}}^k(x, t)).$$

From (5.5) and (5.6) we deduce, due to the fact that  $s_k \neq 0$  when  $k \geq d + 1$ :

$$(5.7) \quad m^k(x, t) = m_{\text{eq}}^k(x, t) + \text{O}(\Delta t), \quad k \geq d + 1.$$

We insert (5.7) into (5.5) and we deduce

$$(5.8) \quad m_*^k(x, t) = m_{\text{eq}}^k(x, t) + \text{O}(\Delta t), \quad k \geq d + 1.$$

Taking into account the relations (2.14) and (3.1) on one hand and (2.12) and (3.9) on the other hand, we deduce (5.1) and (5.2) from (5.7) and (5.8).  $\square$

## 6) Taylor expansion at first order

- We expand now the relation (4.2) one step further with respect to the time step  $\Delta t$ . We introduce the second order moment

$$(6.1) \quad F^{\alpha\beta} \equiv \sum_{j=0}^J v_j^\alpha v_j^\beta f_{\text{eq}}^j, \quad 1 \leq \alpha, \beta \leq d.$$

We denote in the following  $\partial_t$  instead of  $\frac{\partial}{\partial t}$  and  $\partial_\beta$  in place of  $\frac{\partial}{\partial x_\beta}$ . Then we have the following result at the first order.

**Proposition 4.** Euler equations of gas dynamics.

With the lattice Boltzmann scheme previously defined, we have the conservation of mass and momentum at the first order:

$$(6.2) \quad \partial_t \rho + \sum_{\beta=1}^d \partial_\beta q^\beta = \text{O}(\Delta t)$$

$$(6.3) \quad \partial_t q^\alpha + \sum_{\beta=1}^d \partial_\beta F^{\alpha\beta} = O(\Delta t).$$

**Proof of Proposition 4.**

We expand both sides of relation (4.2) up to first order:

$$\begin{aligned} f^j(x, t + \Delta t) &= f^j(x, t) + \Delta t \partial_t f^j + O(\Delta t^2) \\ f_*^j(x - v_j \Delta t, t) &= f_*^j(x, t) - \Delta t v_j^\beta \partial_\beta f_*^j + O(\Delta t^2). \end{aligned}$$

We take the moment of order  $k$  of this identity:

$$m^k(x, t) + \Delta t \partial_t m^k + O(\Delta t^2) = m_*^k(x, t) - \Delta t \sum_{j=0}^J M_j^k v_j^\beta \partial_\beta f_*^j + O(\Delta t^2)$$

and we use the previous Taylor expansions (5.1) (5.2) at the order zero:

$$(6.4) \quad m^k(x, t) + \Delta t \partial_t m_{\text{eq}}^k = m_*^k(x, t) - \Delta t \sum_{j=0}^J M_j^k v_j^\beta \partial_\beta f_{\text{eq}}^j + O(\Delta t^2).$$

We take  $k = 0$  inside the relation (6.4). We get (6.2) since  $m^0(x, t) \equiv m_*^0(x, t) \equiv \rho(x, t)$ . Considering now the particular case  $k = \alpha$  with  $1 \leq \alpha \leq d$ , we have also  $m^\alpha(x, t) \equiv m_*^\alpha(x, t) \equiv q^\alpha(x, t)$  and the relation (6.3) is a direct consequence of the definition (6.1) and the property (2.11).  $\square$

**Proposition 5.** Technical lemma.

We introduce the ‘‘conservation defect’’  $\theta^k$  according to the relation

$$(6.5) \quad \theta^k(x, t) = \partial_t m_{\text{eq}}^k + \sum_{j=0}^J M_j^k v_j^\beta \partial_\beta f_{\text{eq}}^j \equiv \sum_{j=0}^J M_j^k (\partial_t f_{\text{eq}}^j + v_j^\beta \partial_\beta f_{\text{eq}}^j).$$

Then we have the following properties:

$$(6.6) \quad m^k(x, t) = m_{\text{eq}}^k(x, t) - \frac{\Delta t}{s_k} \theta^k + O(\Delta t^2), \quad k \geq d+1,$$

$$(6.7) \quad m_*^k(x, t) = m_{\text{eq}}^k(x, t) - \left( \frac{1}{s_k} - 1 \right) \Delta t \theta^k + O(\Delta t^2), \quad k \geq d+1,$$

$$(6.8) \quad \partial_\beta f_*^j = \partial_\beta f_{\text{eq}}^j - \Delta t \sum_{k=d+1}^J \left( \frac{1}{s_k} - 1 \right) (M^{-1})_k^j \partial_\beta \theta^k + O(\Delta t^2).$$

**Proof of Proposition 5.**

We start from the relation (6.4) and we have observed at the previous proposition that

$$(6.9) \quad \theta^i = O(\Delta t), \quad 0 \leq i \leq d.$$

We remark also that from the relation (5.6), we have

$$m^k(x, t) - m_{\text{eq}}^k(x, t) = \frac{1}{s_k} (m^k(x, t) - m_*^k(x, t)) \quad \text{if } k \geq d+1.$$

Then the relation (6.6) is a direct consequence of (6.4) and the definition (6.5). In consequence, the relation (6.7) follows from (6.6) and (6.4). Due to (6.7), (6.9) and (3.9), we have

$$(6.10) \quad f_*^j(x, t) = f_{\text{eq}}^j(x, t) - \Delta t \sum_{k \geq d+1} \left( \frac{1}{s_k} - 1 \right) (M^{-1})_k^j \theta^k + O(\Delta t^2)$$



and the relation (6.8) follows from derivating (6.10) in the direction  $x_\beta$ .  $\square$

## 7) Equivalent equation at second order

- We introduce the tensor  $\Lambda_k^{\alpha\beta}$  according to

$$(7.1) \quad \Lambda_k^{\alpha\beta} \equiv \sum_{j=0}^J v_j^\alpha v_j^\beta (M^{-1})_k^j, \quad 1 \leq \alpha, \beta \leq d, \quad 0 \leq k \leq J.$$

We can now establish the major result of our contribution.

**Proposition 6.** Navier-Stokes equations of gas dynamics.

With the lattice Boltzmann method defined in previous sections and the conservation defect  $\theta^k$  defined in (6.5), we have the following expansions up to second order accuracy:

$$(7.2) \quad \partial_t \rho + \sum_{\beta=1}^d \partial_\beta q^\beta = O(\Delta t^2)$$

$$(7.3) \quad \partial_t q^\alpha + \sum_{\beta=1}^d \partial_\beta \left( F^{\alpha\beta} - \Delta t \sum_{k \geq d+1} \left( \frac{1}{s_k} - \frac{1}{2} \right) \Lambda_k^{\alpha\beta} \theta^k \right) = O(\Delta t^2).$$

- A consequence of relation (7.3) is the fact that a lattice Boltzmann scheme approximates at second order of accuracy a Navier-Stokes type equation with viscosities  $\mu_k$  of the form

$$(7.4) \quad \mu_k = \Delta t \left( \frac{1}{s_k} - \frac{1}{2} \right).$$

We refer for the details to D. D’Humières (1992), Lallemand and Luo (2000) or to our recent survey (2007). The relations (7.4) are known as the “D’Humières relations”. We observe that in practice, the scalar  $\mu_k$  is imposed by the physics and by the parameter  $\Delta t$  is constrained by the space discretization  $\Delta x$  and the relation (2.1). Then the parameter  $s_k$  must be chosen in order to satisfy the D’Humières relations (7.4).

### Proof of Proposition 6.

We start again from the identity (4.2). We expand both terms up to second order accuracy:

$$f^j(x, t + \Delta t) = f^j(x, t) + \Delta t \partial_t f^j + \frac{1}{2} \Delta t^2 \partial_{tt}^2 f^j + O(\Delta t^3)$$

$$f_*^j(x - v_j \Delta t, t) = f_*^j(x, t) - \Delta t v_j^\beta \partial_\beta f_*^j + \frac{1}{2} \Delta t^2 v_j^\beta v_j^\gamma \partial_{\beta\gamma}^2 f_*^j + O(\Delta t^3).$$

We take the moment of order  $i$  ( $0 \leq i \leq d$ ) of this identity. We obtain:

$$(7.5) \quad \left\{ \begin{array}{l} m^i(x, t) + \Delta t \partial_t m^i + \frac{1}{2} \Delta t^2 \partial_{tt}^2 m^i + O(\Delta t^3) = m_*^i(x, t) + \\ - \Delta t \sum_{j=0}^J M_j^i v_j^\beta \partial_\beta f_*^j + \frac{1}{2} \Delta t^2 \sum_{j=0}^J M_j^i v_j^\beta v_j^\gamma \partial_{\beta\gamma}^2 f_*^j + O(\Delta t^3). \end{array} \right.$$

We use the microscopic conservation  $m_*^i(x, t) \equiv m^i(x, t)$  in (7.5) and the previous Taylor expansion at order one, in particular the relation (6.8). We divide by  $\Delta t$  and we deduce:

$$\begin{aligned}
 \partial_t m^i + \frac{1}{2} \Delta t \partial_{tt}^2 m^i &= - \sum_{j=0}^J M_j^i v_j^\beta \partial_\beta f_{\text{eq}}^j + \\
 &+ \Delta t \sum_{j=0}^J \sum_{k \geq d+1} M_j^i v_j^\beta \left( \frac{1}{s_k} - 1 \right) (M^{-1})_k^j \partial_\beta \theta^k + \\
 &+ \frac{1}{2} \Delta t \sum_{j=0}^J M_j^i v_j^\beta v_j^\gamma \partial_{\beta\gamma}^2 f_{\text{eq}}^j + \text{O}(\Delta t^2).
 \end{aligned}$$

Then

$$(7.6) \quad \left\{ \begin{aligned}
 \partial_t m^i + \sum_{\beta=1}^d \sum_{j=0}^J M_j^i v_j^\beta \partial_\beta f_{\text{eq}}^j &= \\
 &= \Delta t \sum_{\beta=1}^d \sum_{j=0}^J \sum_{k \geq d+1} M_j^i v_j^\beta \left( \frac{1}{s_k} - 1 \right) (M^{-1})_k^j \partial_\beta \theta^k + \\
 &+ \frac{\Delta t}{2} \left( - \partial_{tt}^2 m^i + \sum_{\beta=1}^d \sum_{j=0}^J M_j^i v_j^\beta v_j^\gamma \partial_{\beta\gamma}^2 f_{\text{eq}}^j \right) + \text{O}(\Delta t^2).
 \end{aligned} \right.$$

- We set  $i = 0$  in the relation (7.6) and we look for the conservation of mass. Due to the property  $M_j^0 \equiv 1$ , the sum over  $j$  in the second line of (7.6) is null since  $\sum_{j=0}^J v_j^\beta (M^{-1})_k^j$  is equal to zero. We have also the following algebraic calculus:

$$\begin{aligned}
 \partial_{tt}^2 m^0 &= \partial_{tt}^2 \rho = - \sum_{\beta=1}^d \partial_{t\beta}^2 q^\beta + \text{O}(\Delta t) = - \sum_{\beta=1}^d \partial_\beta \partial_t q^\beta + \text{O}(\Delta t) = \\
 &= \sum_{\beta=1}^d \sum_{\gamma=1}^d \partial_{\beta\gamma}^2 F^{\beta\gamma} + \text{O}(\Delta t) = \sum_{\beta=1}^d \sum_{\gamma=1}^d \sum_{j=0}^J v_j^\beta v_j^\gamma \partial_{\beta\gamma}^2 f_{\text{eq}}^j + \text{O}(\Delta t)
 \end{aligned}$$

and the third line of (7.6) is null up to second order accuracy. Thus the conservation of mass (7.2) up to second order accuracy is established.

- We set  $i = \alpha$  with  $1 \leq \alpha \leq d$  and we look for the conservation of momentum. In this particular case, the relation (7.6) takes the form:

$$(7.7) \quad \left\{ \begin{aligned}
 \partial_t q^\alpha + \sum_{\beta=1}^d \sum_{j=0}^J v_j^\alpha v_j^\beta \partial_\beta f_{\text{eq}}^j &= \\
 &= \Delta t \sum_{k \geq d+1} \left( \frac{1}{s_k} - 1 \right) \sum_{\beta=1}^d \left[ \sum_{j=0}^J v_j^\alpha v_j^\beta (M^{-1})_k^j \right] \partial_\beta \theta^k + \\
 &+ \frac{\Delta t}{2} \left( - \partial_{tt}^2 q^\alpha + \sum_{\beta=1}^d \sum_{j=0}^J v_j^\alpha v_j^\beta v_j^\gamma \partial_{\beta\gamma}^2 f_{\text{eq}}^j \right) + \text{O}(\Delta t^2).
 \end{aligned} \right.$$

We have now to play with some algebra:

$$\begin{aligned}
 -\partial_{tt}^2 q^\alpha + \sum_{\beta=1}^d \sum_{j=0}^J v_j^\alpha v_j^\beta v_j^\gamma \partial_{\beta\gamma}^2 f_{\text{eq}}^j &= \\
 &= \sum_{\beta=1}^d \left( \partial_t \partial_\beta F^{\alpha\beta} + \sum_{j=0}^J v_j^\alpha v_j^\beta v_j^\gamma \partial_{\beta\gamma}^2 f_{\text{eq}}^j \right) + \text{O}(\Delta t) \\
 &= \sum_{\beta=1}^d \partial_\beta \left( \sum_{j=0}^J v_j^\alpha v_j^\beta (\partial_t f_{\text{eq}}^j + v_j^\gamma \partial_\gamma f_{\text{eq}}^j) \right) + \text{O}(\Delta t) \\
 &= \sum_{\beta=1}^d \partial_\beta \left( \sum_{j=0}^J v_j^\alpha v_j^\beta \sum_{k=0}^J (M^{-1})_k^j \theta^k \right) + \text{O}(\Delta t) \\
 &= \sum_{\beta=1}^d \partial_\beta \left( \sum_{k \geq d+1} \left[ \sum_{j=0}^J v_j^\alpha v_j^\beta (M^{-1})_k^j \right] \theta^k \right) + \text{O}(\Delta t) \\
 &= \sum_{\beta=1}^d \partial_\beta \left( \sum_{k \geq d+1} \Lambda_k^{\alpha\beta} \theta^k \right) + \text{O}(\Delta t)
 \end{aligned}$$

due to the definition (7.1). We deduce from (6.1), (7.7) and the above calculus:

$$\begin{aligned}
 \partial_t q^\alpha + \sum_{\beta=1}^d \partial_\beta F^{\alpha\beta} &= \Delta t \sum_{k \geq d+1} \left( \frac{1}{s_k} - 1 \right) \sum_{\beta=1}^d \Lambda_k^{\alpha\beta} \partial_\beta \theta^k + \\
 &\quad + \frac{\Delta t}{2} \sum_{\beta=1}^d \partial_\beta \left( \sum_{k \geq d+1} \Lambda_k^{\alpha\beta} \theta^k \right) + \text{O}(\Delta t^2) \\
 &= \Delta t \sum_{\beta=1}^d \sum_{k \geq d+1} \left( \frac{1}{s_k} - \frac{1}{2} \right) \Lambda_k^{\alpha\beta} \partial_\beta \theta^k + \text{O}(\Delta t^2).
 \end{aligned}$$

and the relation (7.3) is established.  $\square$

## 8) Equivalent equation at second order

- The previous propositions establish that the equivalent partial differential equations of a Boltzmann scheme are given up to second order accuracy by the same result as the formal Chapman-Enskog expansion. We find Euler type equation at the first order (Proposition 4) and Navier-Stokes type equation at the second order (Proposition 6). Note that with the above framework no *a priori* formal two-time multiple scaling is necessary to establish the Navier-Stokes equations from a lattice Boltzmann scheme, as done previously in the contribution of D'Humières. We remark also that a so-called diffusive scaling like  $\frac{\Delta t}{\Delta x^2} = \text{constant}$ , instead of our condition (2.1)  $\frac{\Delta t}{\Delta x} = \text{constant}$ , leads to the incompressible Navier-Stokes equations, as proposed by Junk, Klar and Luo (2005). In both cases, we have just to use the Taylor formula for a single infinitesimal parameter.

## 9) Acknowledgments

The author thanks Li-Shi Luo to his kind invitation to present the scientific work of Orsay's team at ICMMES Conference in July 2005. The author thanks also with a great emphasis Pierre Lallemand for very helpfull discussions all along the elaboration of this contribution. Last but not least, the referees transmitted to the author very good remarks that have been incorporated inside the present edition of the article.

## 10) References

- S.C. Chang. "A critical analysis of the modified equation technique of Warming and Hyett", *Journal of Computational Physics*, vol. 86, p. 107-126, 1990.
- D. D'Humières. "Generalized Lattice-Boltzmann Equations", in *Rarefied Gas Dynamics: Theory and Simulations*, vol. 159 of *AIAA Progress in Astronautics and Astronautics*, p. 450-458, 1992.
- F. Dubois. "Une introduction au schéma de Boltzmann sur réseau", *ESAIM Proceedings*, vol. 18, pages 181-215, July 2007.
- D. Griffiths, J. Sanz-Serna. "On the scope of the method of modified equations", *SIAM Journal on Scientific and Statistical Computing*, vol. 7, p. 994-1008, 1986.
- M. Junk, A. Klar, and L.-S. Luo. "Asymptotic analysis of the lattice Boltzmann equation", *Journal of Computational Physics*, vol. 210, p. 676-704, December 2005.
- P. Lallemand, L.-S. Luo. "Theory of the lattice Boltzmann method: Dispersion, dissipation, isotropy, Galilean invariance, and stability", *Physical Review E*, vol. 61, p. 6546-6562, June 2000.
- A. Lerat, R. Peyret, "Noncentered Schemes and Shock Propagation Problems", *Computers and Fluids*, vol. 2, p. 35-52, 1974.
- G. Strang. *An introduction to applied mathematics*, Wellesley-Cambridge press, Wellesley, 1986.
- R.F. Warming, B.J. Hyett, "The modified equation approach to the stability and accuracy analysis of finite difference methods", *Journal of Computational Physics*, vol. 14, p. 159-179, 1974.