



HAL
open science

Atmospheric particle-bound organophosphate ester flame retardants and plasticizers in a North African Mediterranean coastal city (Bizerte, Tunisia)

Javier Castro-Jiménez, Badreddine Barhoumi, Marc Tedetti, Richard Sempere

► To cite this version:

Javier Castro-Jiménez, Badreddine Barhoumi, Marc Tedetti, Richard Sempere. Atmospheric particle-bound organophosphate ester flame retardants and plasticizers in a North African Mediterranean coastal city (Bizerte, Tunisia). *Science of the Total Environment*, 2018, 642, pp.383-393. 10.1016/j.scitotenv.2018.06.010 . hal-01810090v2

HAL Id: hal-01810090

<https://hal.science/hal-01810090v2>

Submitted on 18 Sep 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Atmospheric particle-bound organophosphate ester flame retardants and**
2 **plasticizers in a North African Mediterranean coastal city (Bizerte, Tunisia)**

3
4 Javier Castro-Jiménez* and Richard Sempéré

5 Aix Marseille Univ., University of Toulon, CNRS, IRD, Mediterranean Institute of
6 Oceanography (MIO) UM 110, Marseille, France.

7 *Corresponding author. Phone: +33(0)486090524;

8 E-mail: javier.castro-jimenez@mio.osupytheas.fr

9
10 <https://doi.org/10.1016/j.scitotenv.2018.06.010>

11

12 **Abstract**

13 Organophosphate ester (OPE) flame retardants and plasticizers have been detected at
14 generally high frequencies (70-98%) for the first time in the atmosphere over the NW
15 African coastal Mediterranean. Results from sixty air samples (total suspended
16 particles, TSP) collected between March 2015 and January 2016 in an urban coastal site
17 (Bizerte, Tunisia) revealed Σ_9 OPE concentrations of ~ 100 - 1060 pg m^{-3} (470 pg m^{-3} ,
18 median) with TCPPs, EHDPP and TiBP exhibiting the higher median concentrations (\sim
19 110 , 100 and 85 pg m^{-3} , respectively). Spring generally exhibited the lowest
20 concentrations, probably linked to the influence of local meteorological conditions and
21 air mass trajectories to a lesser extent. Non-chlorinated OPEs generally predominated,
22 in contrast to the most common reported situation in marine environments (i.e. higher
23 abundance of chlorinated OPEs) pointing to the relevance of local OPE sources in the
24 study area. TiBP levels were generally higher than those reported for other
25 marine/coastal environments suggesting this OPE as a good tracer of local sources in
26 Bizerte. Contrarily, the atmospheric levels of other abundant OPEs in the area (e.g.
27 TCPP) seem to be in the range and/or lower than those reported for remote marine
28 environments. These findings point to the interplay of different factors with solar
29 irradiance (potentially enhancing atmospheric photochemical oxidation reactions) and
30 meteorological conditions in the study area likely compensating potential local sources
31 of some OPEs. Not all OPEs presented the same seasonality in terms of atmospheric
32 concentrations and pattern. The estimated atmospheric dry deposition fluxes (Σ_9 OPEs)
33 were 18 - $180 \text{ ng m}^{-2} \text{ d}^{-1}$. Up to $\sim 9 \text{ kg y}^{-1}$ of OPEs ($\sim 1 \text{ kg y}^{-1}$ of new organic
34 anthropogenic phosphorus coming from OPEs) can be loaded to the shallow and
35 enclosed Bizerte lagoon ($\sim 130 \text{ km}^2$), considered as the most important aquaculture area

36 in Tunisia, with yet unknown implications for the environmental exposure and impacts
37 in the ecosystem functioning.


38 **Keywords:**

39 OPEs, plastic additives, marine pollution, dry deposition, environmental risk,
40 anthropogenic organic phosphorus

41

42 **Graphical Abstract**

43



53 **1. Introduction**

54

55 There is no doubt today about the multi-media global scale occurrence of
56 organophosphate esters (OPEs) flame retardants and plasticizers in the marine
57 environment. OPE have been found in air, water, sediment and aquatic organisms from
58 different marine areas of the world (Zhong et al., 2017; Wei et al., 2015; Brandsma et
59 al., 2015; Castro- Jiménez et al., 2014; Van der Veen and de Boer, 2012, Bollmann et
60 al., 2012; Möller et al., 2011; Kim et al., 2011), including remote environments such as
61 the Arctic (Ma et al., 2017; Sühring et al., 2016a; Hallanger et al., 2015; Salamova et
62 al., 2014a, Möller et al., 2012) and the most pristine areas from the tropical and
63 subtropical Atlantic, Pacific and Indian oceans (Castro-Jiménez et al., 2016). In
64 addition, most recent reports point to their environmental persistency, bioaccumulation
65 and adverse effects in aquatic organisms and humans (ECHA 2008; 2009; Van der Veen
66 and de Boer, 2012; Wei et al., 2015; Wang et al., 2015a,b). These findings have raised
67 awareness on their environmental fate and impacts as it already happened in the past for
68 other persistent organic pollutants (POPs). However, the magnitude and spatial
69 distribution of OPE sources in the environment are closely linked to the recent
70 changes/developments in international regulations, with an important expanding
71 production and usage of OPEs from 2009 as a result of the global ban of their
72 “predecessors” polybrominated diphenyl ethers (PBDEs) (UNEP 2009).

73

74 OPE atmospheric levels in remote areas, supposedly far from sources seem to be in
75 many occasions within the range of levels measured in coastal areas close to suspected
76 sources, suggesting complex atmospheric circulation patterns and different
77 environmental processing and dynamics linked to geographic regions (Castro-Jiménez
78 et al., 2016; Li et al., 2018). It is, therefore, crucial to better investigate the airborne

79 occurrence and loadings of OPEs to marine/coastal environments in close relation to
80 their potential impacts. A recent study performed in the Mediterranean Sea supports the
81 hypothesis that atmospheric aerosols can elicit a number of toxic effects in marine
82 organisms due to the presence of legacy POPs such polychlorinated dibenzo-p-dioxins
83 and dibenzofurans (PCDD/Fs), polychlorinated biphenyls (PCBs) and high molecular
84 weight (HMW) polycyclic aromatic hydrocarbons (PAHs) (Mesquita et al., 2016).
85 Other POPs with recognized toxicity too, such as PBDEs and the “emerging” OPEs, can
86 add up to the “contaminant-aerosol cocktail” (De wit, 2002; Vonderheidea et al., 2008).
87 OPEs have been reported in both the atmospheric particle and gaseous phases, but most
88 of the studies focus on their particle phase occurrence. The atmospheric degradation
89 processes (e.g. photodegradation and/or OH radical oxidation) will progress more
90 slowly for particle-bound OPEs as they will be “sheltered”, increasing therefore their
91 long-range atmospheric transport (LRAT) potential (Liu et al., 2014). In addition, OPEs
92 have been described to be enriched in both fine and coarse particles from the
93 atmospheric aerosols in urban environments (Clark et al., 2017; Faiz et al., 2018).
94 The Mediterranean Sea is a semi-enclosed environment of high ecological and socio-
95 economic relevance (The Mermex group, 2011). This ecosystem has been documented
96 to be largely impacted by legacy POPs and other organic contaminants (Paluselli et al.,
97 2018; The Mermex group 2011; Berrojalbiz et al., 2011, 2011b, 2014; Castro-Jiménez
98 et al., 2008, 2010, 2012; Mandalakis et al., 2005) as well as by plastic fragments (Cózar
99 et al., 2015; Schmidt et al., 2018), which may be potential sources of OPEs to the
100 overlying atmosphere in accumulation zones (Cheng et al., 2013). However, few data
101 exist on OPE baseline levels and environmental behaviour in the Mediterranean Sea.
102 Only one study reported surface water and sediment OPE concentrations from rivers
103 draining into the W Mediterranean (Cristale et al., 2013), and the wide-spread OPE

104 atmospheric occurrence and inputs to surface waters far from the coast was only
105 recently proved (Castro-Jiménez et al., 2014). Coastal measurements of OPEs of any
106 kind are very scarce in the Mediterranean Sea, in particular in the African
107 Mediterranean coast, where even less data on the occurrence of toxic organic
108 contaminants are available.

109 The objectives of this work are: 1) to investigate for the first time the atmospheric
110 occurrence of OPEs in the NW African Mediterranean coast; 2) to estimate their
111 atmospheric dry deposition and loading in Bizerte area (Tunisia) and discuss their
112 potential impacts.

113

114 **2. Materials and Methods**

115

116 *2.1. Area description*

117

118 A detailed area description is offered elsewhere (Barhoumi et al., 2018). Briefly, Bizerte
119 is a medium-size city (~127 000 inhabitants), located in the north of Tunisia, between
120 the Mediterranean Sea and the Bizerte lagoon (Figure 1). Average temperature of 22 °C,
121 with hot summer and mild spring, frequent (200 days per year) NW winds (average
122 speed of 6–8 m s⁻¹) and precipitation episodes mostly in autumn and winter months,
123 characterized the climate of Bizerte area. Even if agriculture and fishery activities are
124 important in the area, a considerable number of light and heavy industries (i.e., cement,
125 plastic, textile, mechanic and electronic, iron and steel metallurgy, petroleum refining
126 and lubricants) are present (Barhoumi et al., 2018) inducing a considerable
127 anthropogenic pressure. A certain number of studies focusing on the atmospheric
128 pollution by legacy organic contaminants, like PAHs, PCBs and OCPs, have been

129 performed in the study area (Ben Hassine et al., 2014, Barhoumi et al., 2018). More
130 recent measurements report on dioxin-like POPs, including PCDD/Fs and DL-PCBs and
131 also on PBDEs (Castro-Jiménez et al., 2017) but alternative flame retardants like OPEs
132 have never been studied to the best of our knowledge (nor in the whole NW African
133 coastal Mediterranean).
134



135

136 **Figure 1.** Sampling site location (Bizerte, Tunisia)

137

138 2.2. Sampling strategy

139

140 Integrated (48 h) aerosol samples (total suspended particles, TSP) were collected from
141 March 2015 to January 2016 on pre-combusted quartz fibre filters (QFFs) (Whatman
142 QMA grade, 20.3 x 25.4 cm) using a high volume air sampler (Tisch Environmental
143 Inc., OH, USA). Overall, 60 samples were collected ($1920 \pm 40 \text{ m}^3$, average volume,

144 Table S1). Sampling was conducted on the roof top (~8 m ASL) of the Faculty of
145 Science of Bizerte (37° 16' 0.5802" N, 9° 52' 49.875" E) around 1 km far from the city
146 center, close to the Gulf of Bizerte shoreline and Bizerte lagoon water front (Figure 1).

147

148 2.3. Sample processing

149

150 The samples were processed as described in Barhoumi et al. (2018). Briefly, each QFF
151 was cut into two equal parts. One half was used for organic pollutant analyses and the
152 other half for determination of ancillary data like TSP, total organic carbon (TOC) and
153 organic nitrogen (TON) (Table S1). The corresponding filter halves were cut into small
154 pieces and spiked with 250 ng of deuterated surrogate standards (tri-n-butyl- d_{27} -
155 phosphate, triphenyl- d_{15} -phosphate, tris(2-chloroisopropyl)- d_{18} -phosphate and tris(1,3-
156 dichloro-2-propyl)- d_{15} -phosphate). Filters were extracted with dichloromethane (DCM)
157 (3 cycles of 5 min) using an accelerated solvent extraction system (ASE 350, DIONEX)
158 at 100 °C, 110 bars, 100% of rinsing volume and 60 s purging time. Few grams of
159 activated copper were added to ASE flasks to remove potential traces of sulphur. The
160 obtained extract was rota-evaporated (with a solvent exchange into hexane) to
161 approximately 0.5 ml and then loaded into a silica-alumina chromatography glass
162 column (10 mm internal diameter, i.d.), prepared by adding, from bottom to top; 3 g
163 deactivated alumina (3% w/w), 3 g activated silica, and 1 g dehydrated sodium sulphate.
164 After conditioning with 15 mL of n-hexane, the column was subsequently eluted into
165 three fractions (F): with 20 mL of hexane (F1), 40 mL of hexane/DCM (80:20, v/v) (F2)
166 and 40 mL DCM/Acetone (70:30, v/v) (F3). OPEs were analysed in F3, which was rota-
167 evaporated down to ~0.2 ml, solvent exchanged into isoctane and further concentrated
168 under gentle nitrogen stream up to ~0.1 mL. Prior to injection, 200 ng of deuterated

169 standards (Tri-n-propyl-*d*₂₁-phosphate, Tris(2-chloroethyl)-*d*₁₂-phosphate and
170 Malathion-*d*₇) were added to the extracts to be used as internal standards for
171 quantification.

172

173 2.4. Instrumental Analysis

174

175 OPE analysis was conducted by gas chromatography (Agilent 7820A Series GC)
176 coupled with mass spectrometry (Agilent 5977E MS) (GC-MS) operating in selected
177 ion monitoring (SIM) and electron impact (EI, 70eV) mode. OPEs were quantified by
178 the internal standard procedure. Table S2 shows selected ions for detection and
179 quantification for each compound. The injector temperature was set at 270 °C (splitless).
180 The separation was achieved over a 30m x 0.25mm i.d. x 0.25µm HP-5MS capillary
181 column (Agilent J&W). The oven temperature was programmed from 90°C to 132°C at
182 3°C/min, to 165°C at 10°C/min, to 235°C at 2°C/min, then to 300°C at 25°C/min
183 (holding time 5min). The injection volume was of 2 µL and the helium carrier gas flow
184 was 1 mL min⁻¹. The temperatures of the MS transfer line; ion source, and quadrupole
185 were set at 300, 230 and 150 °C, respectively. Samples were analysed for the following
186 OPEs: tris-(2-chloroethyl)phosphate (TCEP; CAS: 115-96-8), tris[2-chloro-1-
187 (chloromethyl)ethyl]phosphate (TDCP; CAS: 13674-87-8), Tris- (1-chloro-2-
188 propyl)phosphate (TCPPs, mix of isomers; CAS: 13674-84-5), tripropyl phosphate
189 (TPP; CAS: 513-08-6), tri-iso-butyl phosphate (TiBP; CAS: 126-71-6), tri-n-butyl
190 phosphate (TnBP; CAS: 126-73-8), triphenyl phosphate (TPhP; CAS: 115-86-6), 2-
191 ethylhexyl diphenyl phosphate (EHDPP; CAS: 1241-94-7), and tri(2-ethylhexyl)
192 phosphate (TEHP; CAS: 78-42-2). TOC and TON were determined in filter subsamples

193 by high temperature combustion (CHN analyser) (Raimbault et al., 2008). Individual
194 native OPE standards were purchased from Dr. Ehrenstorfer GmbH (Germany),
195 whereas labeled standards tri-n-butyl- d_{27} -phosphate, triphenyl- d_{15} -phosphate, tri-n-
196 propyl- d_{21} -phosphate and malathion- d_7 were obtained from C/D/N Isotopes Inc.
197 (Canada) and tris(2-chloroisopropyl)- d_{18} -phosphate, tris(1,3-dichloro-2-propyl)- d_{15} -
198 phosphate and tris(2-chloroethyl)- d_{12} -phosphate were from Cambridge Isotope
199 Laboratories, Inc. (USA).

200

201 *2.5. Quality assurance / quality control (QA/QC)*

202

203 The high volume air sampler was regularly calibrated (each month) following the
204 manufacturer's procedure (Tisch Environmental, HIGH VOL+). QFFs were
205 individually wrapped in double aluminium foil and baked at 450 °C overnight before
206 sampling. After each sampling event, the filters were wrapped in the same aluminium
207 foil and stored in double sealed plastic bags in a freezer at -20 °C before processing.
208 The pre- and post-sampling QFF weights were determined after they were placed in a
209 desiccator (25 °C) for 24h. Standards (natives + labelled compounds) were introduced
210 in the chromatographic sequence to evaluate possible variations on the detection
211 conditions during the time of analyses. Prior to use, the ASE cells were pre-cleaned by
212 performing an "empty" ASE extraction with DCM. Three field blanks (taken in autumn
213 - summer), consisting on baked QFFs transported to the sampling area, mounted in the
214 sampler, and dismounted, were collected, stored and analysed concurrently with the
215 samples. Three laboratory blanks were also performed. Blanks results are presented in
216 Table S3. An unexpected problem occurred during the processing of one of the field

217 blanks, so results are only reported for the two remaining blanks. OPE levels in the
218 blanks were generally low, ranging from ~1 to 14 ng for most compounds. Relatively
219 higher blank levels were found for TiBP, EHDPP and TCPP (25 to 74 ng). However,
220 these three OPEs exhibited the highest concentrations in the samples. These higher
221 blank levels were suspected to come from the ASE system used for the sample
222 extraction. A high variability was observed, in particular for the most abundant
223 compounds, highlighting the necessity of collecting more field blanks in future long-
224 term OPE monitoring studies. Laboratory blanks showed lower or similar levels to field
225 blanks, so no contamination during sampling or storage occurred. Results were blank
226 corrected by subtracting the filed blank values (autumn). Instrumental limits of
227 detection (LODs), calculated as signal-to-noise (S/N) ratio > 3, and limits of
228 quantification (LOQ), corresponding to a S/N \geq 10, ranged from 5 to 50 and from 10 to
229 100 pg, respectively depending on the compound (Table S4).

230 Average method recoveries (n=60) (extraction-cleanup-analysis) as monitored by the
231 surrogate OPEs varied from 57 ± 11 to $105\pm 11\%$ (Table S5). In addition, spike
232 experiments for native OPEs were conducted. Clean QFF (n=2) were spiked with a
233 mixture of native OPE (250 ng/sample) and extracted as indicated above. Results
234 confirm the good recoveries of our method; with an average native OPE recovery of
235 $101\pm 24\%$ (Table S6). Results were not corrected by recoveries. NIST SRM 2585 was
236 analysed for target OPEs as well. This reference material is not certified for OPEs,
237 however different laboratories have provided concentration of OPEs in this dust
238 reference material, becoming a recommended quality control step (Fan et al., 2014,
239 Castro-Jiménez et al., 2016). The average results for two replicates (blank corrected)
240 proved that the analysis of OPE using the described methodologies provided

241 concentrations generally in good agreement with reported values for most OPEs (Figure
242 S1). Deviations were found for EHDPP and TDCP, only.

243

244 **3. Results and discussion**

245

246 **3.1. Atmospheric occurrence**

247

248 *3.1.1. Detection frequency and concentrations*

249

250 The studied OPEs were detected in the aerosol samples over Bizerte city at generally
251 high frequencies, except for TPP and TDCP (only found in 8% and 12% of the samples,
252 respectively). Low detection frequencies of TDCP in the atmosphere has been
253 previously reported in some environments, like in the Great Lakes (Salamova et al.,
254 2014b). The non-chlorinated OPEs (TPhP, EHDPP, TnBP and TEHP) were generally
255 the most frequently found (92-98%), although closely followed by the chlorinated
256 TCEP and TCPPs (85-90%). The other alkyl-OPE measured (i.e. TiBP) was present in
257 the 70% of the samples (Figure S2). However, the minor frequency observed for TiBP
258 can be attributed to the blank subtraction performed (relatively high field blank levels
259 compared to the other alkyl-OPEs) (Table S3). These results confirm the presence of
260 OPEs at the NW Mediterranean African coast, never reported before.

261 Overall, the $\sum_9\text{OPE}$ particle-bound atmospheric concentrations (considering only
262 detected values) ranged from ~100 to 1060 pg m^{-3} (470 pg m^{-3} , median) along the
263 sampling period with TCPPs, EHDPP and TiBP exhibiting the higher median
264 concentrations of ~ 110, 100 and 85 pg m^{-3} , respectively. Table S7 shows the compound
265 specific concentrations for all aerosol samples. Very few OPE atmospheric

266 measurements in coastal areas have been performed so far. The vast majority of
267 available data in marine environments come from cruise measurements. The measured
268 values in Bizerte are generally lower than those reported across the Mediterranean Sea
269 ($\sum_{14}\text{OPE}$: 400-5000 pg m^{-3}) and the SW Black Sea ($\sum_{14}\text{OPE}$: 1700-6000 pg m^{-3})
270 (Castro-Jiménez et al., 2014), the E China Sea ($\sum_8\text{OPE}$ 230-2900 pg m^{-3}) (Möller et al.,
271 2012) and tropical and subtropical areas from the Atlantic, Pacific and Indian Oceans
272 ($\sum_{14}\text{OPE}$: 360-4400 pg m^{-3} (Castro-Jiménez et al., 2016), but in the range of those from
273 the North Sea ($\sum_8\text{OPE}$: 100-1400 pg m^{-3}) (Möller et al., 2012) and Svalbard coastal area
274 ($\sum_8\text{OPE}$: 30-1450 pg m^{-3}) (Salamova et al., 2014a). However, OPE levels in Bizerte
275 were higher than those reported for other semi-enclosed marine environments like the S
276 China Sea ($\sum_9\text{OPE}$: 50-160 pg m^{-3}) (Lai et al., 2015) and also less impacted areas like
277 the Coral Sea ($\sum_4\text{OPE}$: 90 pg m^{-3}) (Chen et al., 2013). These comparisons should be
278 taken with caution though, due to the consideration of different number (and type, in
279 some cases) of compounds included in the sums and the different seasons and years of
280 the studies. Although it may still high variable, a more meaningful comparison would
281 be considering individual compounds. The atmospheric levels of the predominant OPEs
282 in Bizerte (i.e. TCPPs, EHDPP and TiBP) were individually compared with existing
283 studies. Table 1 provides an updated compilation of atmospheric particle-bound OPE
284 measurements in marine/coastal environments and some reference inland urban
285 locations for comparison. Interestingly, some different trends were found compared to
286 those of $\sum_9\text{OPE}$ s; with TCPPs concentrations in Bizerte higher than in the E China Sea
287 and Svalbard, but lower than those in the North Sea. TiBP levels in Bizerte appear to be
288 generally higher than in other marine/coastal environments studied. This compound is
289 used in a wide variety of industrial applications including building, papers, boards,
290 plastics, rubbers, textile and directly as an additive in various products such as lacquers,

291 hydraulic fluids and floor polishes (ECHA website). Therefore, their relatively high
292 levels measured in this study could be related to the plastic and textile industries located
293 in Menzel Jemil city, 5 km SE of the sampling station (Figure 1) (Barhoumi et al.,
294 2018). EHDPP concentrations in Bizerte were in the range of those reported for the
295 Black Sea but lower than in the W Mediterranean Sea (Castro-Jiménez et al., 2014) and
296 the tropical and sub-tropical Atlantic, Pacific and Indian Oceans (Castro-Jiménez et al.,
297 2016).

298

299 Despite the variability that the mentioned factors may have induced, it is not completely
300 clear why the atmospheric levels measured in Bizerte (a coastal urban city under
301 important anthropogenic pressures) for some of the most abundant OPEs like TCPP
302 (among others, see Table 1), are in the range and/or lower than those reported for
303 remote coastal environments or pristine oceanic regions. Most probably a combination
304 of differential regional emission patterns, meteorological conditions and rates of
305 atmospheric processing can explain these data. For example, the important global solar
306 irradiance in the Mediterranean basin (The Mermex group, 2011, Sempéré et al., 2015),
307 reaching annual average values of $\sim 210 \text{ W m}^{-2}$ in Bizerte city (Chelbi et al., 2015, Ben
308 Othman et al., 2018) could enhance atmospheric degradation due to photochemical
309 oxidation reactions. Even if these processes likely progress faster for gas-phase
310 compounds, the atmospheric photochemical oxidation of particle-bound OPEs has also
311 been described (Liu et al., 2014). These conditions would favour the decrease of OPE
312 atmospheric levels, even if land-based sources could be higher than in other
313 environments (like the Arctic, with less solar irradiation rates and colder temperatures).
314 In addition, the huge expansion of OPE production and usage experienced in EU and
315 other developed countries could have not yet arrived to the NW African edge of the

316 Mediterranean Sea, where OPE consumption (and related emission) may be still
317 moderate. However, no public data on OPE production and consumption exist in this
318 region to the best of our knowledge, highlighting the necessity of OPE emission
319 inventories at the African Mediterranean coast.

320

321 *3.1.2. Atmospheric pattern and seasonality*

322


323 As indicated, TCPPs, EHDPP and TiBP exhibited the higher concentrations and
324 therefore dominated the global OPE atmospheric pattern (n=60) over Bizerte, accounting
325 for the 25±16 %, 23±10 %, and 20±15% of the Σ_9 OPEs, respectively (Figure S3). A
326 general predominance of Cl-OPEs, in particular TCPP, in atmospheric aerosols in open
327 marine environments has been reported (Li et al., 2018; Castro-Jiménez et al., 2016;
328 Möller et al., 2012). However, a higher relative contribution of the alkyl and aryl OPEs
329 have been reported over urban areas or in remote areas under the suspected influence of
330 local sources (Castro-Jiménez et al., 2016; Salamova et al., 2014a). The Cl-OPEs (as
331 sum of TCEP, TCPP and TDCP) and the non-Cl-OPEs (as sum of TPP, TiBP, TnBP,
332 TPhP, EHDPP and TEHP) globally (n=60) accounted for the 28% and the 72% (median
333 values) of the Σ_9 OPEs over Bizerte area, respectively (Table S8), generally pointing to
334 the influence of current local OPE sources in the study area. This relative distribution is
335 also consistent with the suspected non-Cl-OPEs sources, in particular TiBP, from the
336 plastic industry located SE the sampling site. Generally non-Cl-OPEs are used as
337 plasticizers, contrary to Cl-OPEs which are more commonly used as flame retardants
338 (Wei et al., 2015). Interestingly, three samples (Prel_22, 30, 72, Table S8) showed the
339 opposite pattern with a clear predominance of Cl-OPEs, accounting from the 77% to
340 88% of the Σ_9 OPEs, in line with the most common reported situation in marine

341 environments (76% median value of predominance for Cl-OPEs) (Li et al., 2018).
342 Indeed, the air mass back trajectories (BTs), calculated with the HYSPLIT model
343 (Draxler and Rolph, 2011), revealed a clear Atlantic Ocean influence for those samples
344 (Figure S4). All BTs are also available in SI (see discussion below)

345

346 Samples were grouped by seasons based on local climate of Bizerte city as follows:
347 autumn (September, October, and November), winter (December, January), spring
348 (March, April, and May) and summer (June, July and August). When comparing the
349 OPE atmospheric pattern by seasons (Figure 2) with the annual pattern (Figure S3),
350 some differences appeared. Whereas the relative distribution of Cl-OPEs appears to be
351 pretty constant along the year, except for the winter months, when TDCP was not
352 detected at all, some differences in the relative distribution of non-Cl-OPEs were
353 observed. First, TiBP and TnBP showed a similar relative distribution in spring,
354 whereas TiBP dominated over TnBP along the rest of the year, in particular in winter. In
355 addition, EHDPP dominated the atmospheric pattern in all seasons except in winter,
356 when TEHP exhibited higher relative concentrations. These findings point to a temporal
357 variability on the OPE atmospheric pattern in the study area.

358



359


360 **Figure 2.** OPE atmospheric particle-bound pattern in Bizerte area in the different
 361 seasons (2015-2016)

362

363 In order to further investigate the OPE seasonality in the area studied; the differences in
 364 contaminant concentrations were evaluated as well (Figure 3). Non-parametric tests
 365 (STATA/SE 12.1 statistical software) were used in order to investigate possible
 366 significant differences among contaminants. For this data analysis, non-detected values
 367 were imputed to the median value between zero and the LOD for each specific
 368 contaminant. Results from the Kruskal-Wallis test for OPEs (TPP and TDCP were not
 369 considered due to their low detection frequency) showed significant differences (p
 370 ≤ 0.003) among seasons for the Σ OPEs and all individual OPEs, except TPhP (Table
 371 S9). Contaminant concentrations (except for TPhP) were further compared by pairs of
 372 regions (Wilcoxon rank-sum test). Each contaminant level was compared among 4
 373 seasons resulting in a total of 6 pairs of comparisons. The False Discovery Rate (FDR)
 374 correction (Benjamini and Hochberg, 1995; Benjamini and Yekutieli, 2001), with

375 Simes's method (Simes et al., 1986) was used to account for the effect of the multiple
376 comparisons (Table S10).

377 Overall, the two seasons showing more significant differences were spring and summer
378 (Table S10) with all OPEs and the Σ OPEs exhibiting significantly higher concentrations
379 in summer compared to spring. Indeed, the sum of OPEs and most individual OPEs
380 showed the lowest concentrations in spring (i.e. TCPPs, EHDPP, TiBP and TEHP) with
381 median values from 2 to 4-fold significantly lower ($p \leq 0.02$) than those measured in the
382 other months of the year (Figure 3, Table S10). However, only three OPEs showed
383 significant differences between the coldest (winter) and the warmest (summer) seasons,
384 but with contrasting trends. TCEP and TnBP were the only OPEs showing significantly
385 ($p < 0.001$) higher levels (~4-fold) in summer compared to winter, whereas TEHP
386 concentration was ~3-fold higher in winter. In addition, TEHP was the only compound
387 exhibiting significantly ($p \leq 0.01$) higher concentrations in winter (~3-fold) compared to
388 the other three seasons, whereas TnBP showed the opposite trend, with significantly (p
389 ≤ 0.03) lower levels in winter (~2-fold) compared to the other three seasons. These
390 observations are consistent with their different relative distribution observed in the
391 winter pattern (Figure 2).



392

393 **Figure 3.** Box plots of particle-bound atmospheric concentrations (pg m^{-3}) of most
 394 frequently found OPEs by season. Lines within the boxes represent the median
 395 concentrations.

396 *3.1.3. Potential factors affecting seasonality*

397

398 OPE concentrations did not correlate with other parameters simultaneously measured in
 399 the same samples, like TSP, TOC and TON, which ranged from 9.6 – 119.0, 1.0 – 24.5

400 and $0.2 - 3.0 \mu\text{g m}^{-3}$, respectively (Figure S5). The lack of correlation between some
401 OPEs (like TCPP and EHDDP) and the atmospheric TSP and/or TOC levels over urban
402 areas have been previously reported (Clark et al., 2017). However, the authors showed
403 that certain OPE could correlate with other fractions of the atmospheric aerosols, the
404 finest fractions like PM_{2.5}. Indeed, other studies conducted in urban areas confirmed
405 the OPE associations with PM_{2.5} fraction (Faiz et al., 2018; Liu et al., 2016), with
406 strong correlations reported for some OPEs, like TCEP, TnBP and TEHP (Faiz et al.,
407 2018). The form OPEs are emitted to the atmosphere seems to be a key factor
408 determining these associations, with higher probability of enrichment in the finest
409 fractions for OPE directly emitted in the gas phase due to the larger surface area of fine
410 particles compared to coarse particles (Faiz et al., 2018; Clark et al., 2017). However,
411 these associations cannot be evaluated in the study area with available data.

412

413 TCEP, TnBP, EHDPP as well as Σ OPEs showed significant positive correlations with
414 temperature ($p \leq 0.002$), consistent with their higher levels found in summer (average $T =$
415 $26 \pm 2^\circ\text{C}$) compared to winter (average $T = 12 \pm 2^\circ\text{C}$), in particular for TCEP and TnBP.
416 Even if not all OPEs did show correlations with temperature, this could partially explain
417 the lower levels generally found in spring (average $T = 16 \pm 3^\circ\text{C}$) compared to summer.
418 Temperature could play a local effect by increasing the OPE volatilization from land
419 and water and also affecting their environmental partitioning. Interestingly, TCEP and
420 TnBP show also a weak (but statistically significant, $p \leq 0.001$) negative correlation with
421 the relative humidity (RH). Significantly ($p = 0.02$) higher RH values ($70 \pm 7\%$) and
422 slightly higher precipitation rates (9.0 mm) and wind speeds ($17 \pm 7 \text{ km h}^{-1}$) were
423 registered in spring compared to summer ($66 \pm 5\%$, 0.6 mm and $15 \pm 6 \text{ km h}^{-1}$,

424 respectively) (Figure S5, Table S1) pointing to a most efficient atmospheric particle
425 washout in spring, which is consistent with our observation too.

426 The BTs (Figure S6) did not really show a consistent influence on the OPE seasonality,
427 except for punctual cases affecting the atmospheric pattern and/or concentrations (as
428 indicated above). The BTs computed for the samples collected in spring generally had a
429 longer air mass range with Atlantic Ocean influence, compared to those for summer,
430 when air masses generally showed a shorter range and spent more time circulating over
431 the W Mediterranean and continental Europe. In addition, some samples showed a clear
432 SE influence of air mass trajectory in summer (e.g. Prel_27, 28) leading to higher TiBP
433 levels (Tables S7, S8) consistent too with our observation of higher levels in summer
434 compared to spring and suspected influence of plastic industry located at the SE from
435 the sampling site. The local wind regimes could have also an influence in the
436 seasonality observed. Figure S7 shows the wind roses for all seasons. A general
437 predominance of winds blowing in the sector WNW-NW was observed for all seasons
438 except for winter, when a clear predominance for SE winds was registered. This could
439 partially explain as well the different pattern observed in winter for non-Cl-OPEs, with
440 higher TiBP and TEHP levels potentially coming from the plastic factory located SE
441 from the sampling site, as discussed above. In summary, a number of factors, likely
442 acting simultaneously, seem to be affecting the relative abundance and distribution
443 pattern between the Cl- and non-Cl-OPEs and their atmospheric concentrations in the
444 study area, like potential local sources and meteorological conditions, influence of air
445 mass trajectories, the LRAT potential of some OPEs travelling from distant sources
446 and/or the atmospheric degradation and depletion processes in the region.

447

448 **3.2. Atmospheric loading**

449

450 3.2.1. Dry deposition fluxes

451

452 Dry deposition fluxes (F_{DD} , $\text{ng m}^{-2} \text{d}^{-1}$) of target OPEs were calculated as:

453

$$454 \quad F_{DD} = 864 v_d C_A \quad [1]$$

455

456 where C_A (ng m^{-3}) is the specific OPE volumetric concentration in the aerosol, v_d (cm s^{-1})
457 is the particle deposition velocity and 864 is a unit conversion factor. A reference
458 value of 0.2 cm s^{-1} for v_d was adopted for this site. This value is within the range of v_d
459 experimentally measured (Del Vento and Dachs 2007) in coastal NW Mediterranean
460 Sea and has recently been used in the same site for the estimation of dry deposition
461 fluxes of other organic contaminants with similar physical-chemical properties as OPEs
462 (Castro-Jiménez et al., 2017). Even if a degree of uncertainty is associated to this
463 calculation due to the lack of v_d measurements during the sampling events, the selected
464 value enables meaningful comparisons with other estimations in the study area and in
465 the W Mediterranean Sea. Overall, the $\Sigma_9\text{OPEs}$ atmospheric dry deposition fluxes
466 ranged from 18 to $180 \text{ ng m}^{-2} \text{d}^{-1}$ during the sampling period (Table S11). These fluxes
467 were generally lower than previous dry deposition estimations performed in the W
468 Mediterranean Sea and in the SW Black Sea (Castro-Jiménez et al., 2014) but within the
469 range of those reported for the North Sea (Möller et al., 2011) and the Philippine Sea
470 (Möller et al., 2012). However, the presented estimations of dry deposition are higher
471 than those reported for other semi-enclosed environments like the China Sea (Lai et al.,
472 2015) and also less impacted areas like the Coral Sea (Chen et al., 2013).

473 Table 2 presents the dry deposition fluxes for each OPE by season. The highest median
474 atmospheric loading (and associated higher potential exposure) for the assemble of
475 OPEs via atmospheric dry deposition in the study area is estimated to occur in the
476 summer, autumn and winter months (as expected from their relative atmospheric
477 concentrations and the assumptions considered in these calculations) with fluxes
478 ranging from 32 to 38 $\mu\text{g m}^{-2} \text{y}^{-1}$ for these seasons. This loading is 1 to 3 orders of
479 magnitude higher than the reported atmospheric dry deposition fluxes of PBDEs (the
480 OPEs “predecessors”) (0.07-3.4 $\mu\text{g m}^{-2} \text{y}^{-1}$ for $\Sigma_{27}\text{PBDEs}$), and even higher compared to
481 the dry deposition fluxes estimated for other legacy POPs like PCDD/Fs, PCBs and
482 organochlorine pesticides (OCPs) over Bizerte during the same observation period
483 (Castro-Jiménez et al., 2017; Barhoumi et al., 2018). However, the atmospheric dry
484 deposition fluxes of OPEs are in the lower end of those reported for $\Sigma_{34}\text{PAHs}$ (30-1100
485 $\mu\text{g m}^{-2} \text{y}^{-1}$) in Bizerte for the same period (Barhoumi et al., 2018).

486

487 3.2.2. *Environmental relevance and potential impacts*

488

489 According to these calculations, the nearby coastal lagoon of Bizerte ($1.28 \times 10^8 \text{ m}^2$
490 surface, Barhoumi et al., 2016), a well-recognized sensitive environment of ecological
491 and socio-economic importance (the most important aquaculture area in Tunisia,
492 annually producing around 100 tons of Pacific oysters and Mediterranean mussels,
493 Béjaoui-Omri et al., 2014) could receive up to $\sim 9 \text{ kg}$ of OPEs yearly. This loading of
494 OPEs may have, yet unknown, implications for the aquatic organisms and the
495 ecosystem functioning. The input of these hazardous compounds may elicit an effect
496 through environmental exposure, as a single group of contaminants or in combination to
497 other toxic organic chemicals also efficiently incorporated to this aquatic environment

498 *via* the atmosphere like PCDD/Fs, PCBs, PBDEs, OCPs and PAHs (Castro-Jiménez et
499 al., 2017; Barhoumi et al., 2018). In addition, OPE may interfere the natural phosphorus
500 cycle. Recent estimations performed in the Mediterranean Sea and other oceanic
501 environments suggest that atmospheric deposition fluxes of phosphorus due to OPE
502 deposition (and other anthropogenic organic pollutants containing phosphorus) could
503 play an important role in the phosphorus cycle, in particular in oligotrophic
504 environments, affecting the ecosystem structure and functioning, for example modifying
505 primary production rates (Castro-Jiménez et al., 2014, 2016). The dry deposition fluxes
506 of Σ_9 OPEs given as P averaged $\sim 10 \text{ ng}_P \text{ m}^{-2} \text{ d}^{-1}$ over Bizerte lagoon area. The maximum
507 estimated loading of new anthropogenic phosphorus coming from OPEs was of $\sim 1 \text{ kg y}^{-1}$
508 ¹. This input is certainly small, representing only between the 0.02-0.5% of the organic
509 phosphorus dry deposition fluxes recently estimated in the W Mediterranean Sea
510 (Violaki et al., 2018). However, these fluxes accounted for both anthropogenic and non-
511 anthropogenic organic compounds, and our estimates correspond exclusively to nine
512 compounds belonging to an emerging widely diffused anthropogenic contaminant class
513 contain phosphorous. Moreover, the reported organic phosphorous inputs for the
514 Mediterranean (Violaki et al., 2018) correspond to the entire W Mediterranean basin,
515 whereas the organic phosphorous amounts due to OPEs will distribute only on the ~ 130
516 km^2 of lagoon surface, a shallow and enclosed environment with limited water
517 exchanges. In addition, recent studies confirmed that OPE can also exist in the
518 atmospheric gas phase (Li et al., 2018; Sühring et al., 2016b, Castro-Jiménez et al.,
519 2016) and the diffusive deposition fluxes (i.e. the direct input of gas phase OPE to the
520 water dissolved phase) could be 2-3 orders of magnitude higher than the dry deposition,
521 potentially triggering up to 1% of the reported primary production in the most
522 oligotrophic oceanic regions (Castro-Jiménez et al., 2016). OPE could also enter Bizerte

523 lagoon by riverine inputs, a source of new anthropogenic phosphorus associated to OPE
524 not yet evaluated. A global assessment performed with available data before the
525 beginning of the large expansion of OPE production and usage (in 2009) estimated the
526 human interference with the phosphorus cycle very close to a threshold of no return (i.e.
527 irreversible disruptive effect) (Rockström et al., 2009). The not yet accounted new
528 anthropogenic phosphorus coming from OPEs and other P-organic containing
529 contaminants, even if small at a regional/global scale, may be contributing to reach the
530 mentioned threshold and efforts should be devoted to study these links and the
531 integrated impacts of OPEs in aquatic/marine ecosystems.

532

533 **4. Conclusions**

534

535 Organophosphate ester flame retardants and plasticizers have been measured for the
536 first time in the atmosphere of the NW African coastal Mediterranean Sea (Bizerte city
537 area) confirming the general high detection frequency of most common OPEs. A
538 general predominance of non-chlorinated OPEs has been found in the atmospheric
539 aerosols, being different from what is commonly reported in marine environments (i.e.
540 higher predominance of chlorinated OPEs) suggesting current local sources in the area.
541 Some compounds could be regarded as useful tracers of local/regional specific
542 activities/sources, like TiBP as tracer of the potential impact of plastic industries in
543 Bizerte area. The lowest concentrations were generally found in spring, probably linked
544 to local meteorological conditions and the LRAT of OPEs from other areas to a lesser
545 extent. The atmospheric levels of some of the most abundant OPEs (like TCPP) in
546 Bizerte area, a coastal urban city under important anthropogenic pressures, seem to be
547 in the range and/or lower than those reported for remote coastal environments or

548 pristine oceanic regions. These findings suggest a combined effect of differential
549 regional emission patterns, meteorological conditions and rates of atmospheric
550 processing (with high solar irradiance, probably leading to a most efficient atmospheric
551 OPE depletion processes in the area, likely compensating potential local sources). In
552 addition, the influence of the existing regulatory frameworks in different areas of the
553 globe may contribute to this situation. Although OPE atmospheric concentrations as
554 sum of compounds could provide a first useful insight on their environmental levels, the
555 environmental behaviour of OPEs should be better investigated for individual
556 compounds. We found that not all OPEs presented the same seasonality in terms of
557 atmospheric concentrations and pattern. The estimated dry deposition fluxes to Bizerte
558 lagoon point to a yearly input of around ~ 9 kg of OPEs (~1 kg of new anthropogenic
559 organic phosphorus) to this shallow and enclosed aquatic environment (~130 km²)
560 considered as the most important aquaculture area in Tunisia, with yet unknown
561 implications in terms of environmental exposure and impacts in the ecosystem
562 functioning. Further research is needed to elucidate the magnitude and potential impacts
563 of these new organic contaminants and associated sources of anthropogenic
564 phosphorous in the study area.

565

566 **Acknowledgements**

567

568 This work is a contribution to the Labex OT-Med (n° ANR-11-LABX-0061) –
569 MEDPOP- funded by the French Government “Investissements d’Avenir” (ANR)
570 through the A*MIDEX project (no ANR-11-IDEX-0001-02), and takes place in the
571 MERMEX/MISTRALS program. It also received the financial support from the IRD
572 French-Tunisian International Joint Laboratory (LMI) “COSYS-Med” and the PACA
573 region project “Particule”. We thank Badreddine Barhoumi for his work on the
574 sampling, sample processing in the lab and manuscript revision and Marc Tedetti for his
575 logistic support, contribution to the results discussion and revision of the manuscript.
576 Catherine Guigue is acknowledged for her help on the sample preparation for the TOC
577 determinations and Patrick Raimbault for the TOC analyses. Mohamed Driss is kindly
578 acknowledged for allowing sampling on the Bizerte Faculty of Science. The project
579 leading to this publication has received funding from European FEDER Fund under
580 project 1166-39417.

581

582 **Supporting information available**

583

584 Additional data on the sampling and analytical procedures, QA/QC, compound-by-
585 compound atmospheric levels and spatial distribution and deposition fluxes are
586 presented in this section.

587

588 **References**

589

590 Barhoumi, B., Castro-Jiménez, J., Guigue, C., Goutx, M., Sempéré, R., Derouiche, A.,
591 Achouri, A., Touil, S., Driss, M.R., Tedetti, M. 2018. Levels and risk assessment of
592 hydrocarbons and organochlorines in atmospheric suspended particles from a north
593 African coastal city (Bizerte, Tunisia). *Environ. Pollut.* 240, 422-431.

594

595 Barhoumi, B., El Megdiche, Y., Clérandeau, C., Ben Ameer, W., Mekni, S.,
596 Bouabdallah, S., Derouiche, A., Touil, S., Cachot, J., Driss, M. R. 2016. Occurrence of
597 polycyclic aromatic hydrocarbons (PAHs) in mussel (*Mytilus galloprovincialis*) and eel
598 (*Anguilla anguilla*) from Bizerte lagoon, Tunisia, and associated human health risk
599 assessment. *Cont. Shelf Res.* 124, 104–116.

600

601 Béjaoui-Omri, A., Béjaoui, B., Aloui-Bejaoui, N., Harzallah, A., El Bour, M., Aleya, L.
602 2014. Application of a dynamic energy budget model to *Mytilus galloprovincialis* in the
603 Bizerte lagoon (North of Tunisia). *Environ. Sci. Poll. Res.* 21, 13081-13094.

604

605 Ben Hassine, S., Hammami, B., Ben Ameer, W., El Megdiche, Y., Barhoumi, B., Driss,
606 M. R. 2014. Particulate Polycyclic Aromatic Hydrocarbons (PAH) in the atmosphere of
607 Bizerte city, Tunisia. *Bull. Environ. Contam. Toxicol.* 93, 375–382.

608

609 Ben Othman, A., Belkilani, K., Besbes, M. 2018. Global solar radiation on tilted
610 surfaces in Tunisia: Measurement, estimation and gained energy assessments. *Energ.*
611 *Rep.* 4, 101-109.

612 Benjamini, Y., Hochberg, Y. 1995. Controlling the false discovery rate: A practical and
613 powerful approach to multiple testing. *J. Roy. Stat. Soc. B Met.* 57, 289-300.
614

615 Benjamini, Y., Yekutieli, D. 2001 The control of the false discovery rate in multiple
616 testing under dependency. *The Annals of Statistics* 29, 1165-1188.
617

618 Berrojalbiz, N., Dachs, J., Ojeda, M.J., Valle, M.C., Castro-Jiménez, J., Wollgast, J.,
619 Ghiani, M., Hanke, G., Zaldivar, J.M., 2011a Biogeochemical and physical controls on
620 concentrations of polycyclic aromatic hydrocarbons in water and plankton of the
621 Mediterranean and Black Seas. *Global Biogeochem. Cy.* 25,
622 doi:10.1029/2010GB003775
623

624 Berrojalbiz, N., Dachs, J., Ojeda, M.J., Valle, M.C., Del Vento, S., Castro-Jiménez, J.,
625 Mariani, G., Wollgast, J., Hanke G., 2011b. Persistent organic pollutants in
626 Mediterranean seawater and processes affecting their accumulation in plankton.
627 *Environ. Sci. Technol.*, 45, 4315-4322.
628

629 Berrojalbiz, N., Castro-Jiménez, J., Mariani, G., Wollgast, J., Hanke, G., Dachs, J.,
630 2014. Atmospheric occurrence, transport and deposition of polychlorinated biphenyls
631 and hexachlorobenzene in the Mediterranean and Black seas. *Atmos. Chem. Phys.* 14,
632 8947-8959.
633

634 Bollmann, U.E., Möller, A., Xie, Z., Ebinghaus, R., Einax, J.W., 2012. Occurrence and
635 fate of organophosphorus flame retardants and plasticizers in coastal and marine surface
636 waters. *Water Res.* 46, 531-538.

637 Brandsma, S.H., Leonards, P.E.G., Leslie, H.A., de Boer, J. 2015. Tracing
638 organophosphorus and brominated flame retardants and plasticizers in an estuarine food
639 web. *Sci. Total. Environ.* 505, 22-31.

640

641 Castro-Jiménez, J., Deviller, G., Ghiani, M., Loos, R., Mariani, G., Skejo, H., Umlauf,
642 G., Wollgast, J., Laugier, T., Héas-Moisan, K., Léauté, F., Munsch, C., Tixier, C.,
643 Tronczyński, J., 2008. PCDD/F and PCB multi-media ambient concentrations, congener
644 patterns and occurrence in a Mediterranean coastal lagoon (Etang de Thau, France).
645 *Environ. Pollut.* 156, 123-135.

646

647 Castro-Jiménez, J., Eisenreich, S.J., Ghiani, M., Mariani, G., Skejo, H., Umlauf, G.,
648 Wollgast, J., Zaldívar, J.M., Berrojalbiz, N., Reuter, H.I., Dachs, J., 2010. Atmospheric
649 occurrence and deposition of polychlorinated dibenzo-p-dioxins and dibenzofurans
650 (PCDD/Fs) in the open Mediterranean Sea. *Environ. Sci. Technol.* 44, 5456-5463.

651

652 Castro-Jiménez, J., Berrojalbiz, N., Wollgast, J., Dachs, J. 2012. Polycyclic aromatic
653 hydrocarbons (PAHs) in the Mediterranean Sea: atmospheric occurrence, deposition and
654 decoupling with settling fluxes in the water column. *Environ. Pollut.*, 166, 40-47.

655

656 Castro-Jiménez, J., Berrojalbiz, N., Pizarro, M., Dachs, J. 2014. Organophosphate Ester
657 (OPE) flame retardants and plasticizers in the open Mediterranean and Black Seas
658 Atmosphere. *Environ. Sci. Technol.* 48, 3203-3209.

659

660 Castro-Jiménez, J., Casal, P., González Gaya, B., Pizarro, M., Dachs, J., 2016.
661 Organophosphate ester flame retardants and plasticizers in the global oceanic
662 atmosphere. *Environ. Sci. Technol.* 50, 12831–12839
663

664 Castro-Jiménez, J., Barhoumi, B., Paluselli, A., Tedetti, M., Jiménez, B., Muñoz-
665 Arnanz, J., Wortham, H., Driss, M.R., Sempéré, R. 2017. Occurrence, loading and
666 exposure of atmospheric particle-bound POPs at the African and European edges of the
667 western Mediterranean Sea. *Environ. Sci. Technol.* 51, 13180–13189
668

669 Chelbi, M., Gagnon, Y., Waewsak, J. 2015. Solar radiation mapping using sunshine
670 duration-based models and interpolation techniques: Application to Tunisia. *Energ.*
671 *Convers. and Manag.* 101, 203-215.
672

673 Cheng, W., Xie, Z., Blais, J.M., Zhang, P., Li, M., Yang, C., Huang, W., Ding, R., Sun,
674 L. 2013. Organophosphorus esters in the oceans and possible relation with ocean gyres.
675 *Environ. Pollut.* 180, 159-164
676

677 Clark, A.E., Yoon, S., Sheesley, R.J., Usenko, S., 2017. Spatial and temporal
678 distributions of organophosphate ester concentrations from atmospheric particulate
679 matter samples collected across Houston, TX. *Environ. Sci. Technol.* 51, 4239–4247.
680

681 Cózar, A., Sanz-Martín, M., Martí, E., González-Gordillo, J.I., Ubeda, B., Gálvez, J.Á.,
682 Irigoien, X., Duarte, C.M., 2015. Plastic Accumulation in the Mediterranean Sea. *PLoS*
683 *ONE* 10(4), 1-12.
684

685 Cristale, A., García Vázquez, A., Barata, C., Lacorte, S. 2013. Priority and emerging
686 flame retardants in rivers: Occurrence in water and sediment, *Daphnia magna* toxicity
687 and risk assessment. *Environ. Int.* 59, 232-243.

688

689 De Wit, C., 2002. An overview of brominated flame retardants in the environment.
690 *Chemosphere.* 46, 583–624

691

692 Del Vento, S., Dachs, J. 2007. Influence of the surface microlayer on atmospheric
693 deposition of aerosols and polycyclic aromatic hydrocarbons. *Atmos. Environ.* 41,
694 4920–4930.

695

696 Draxler, R.R., Rolph, G.D. 2011. HYSPLIT (HYbrid Single-Particle Lagrangian
697 Integrated Trajectory) Model access via NOAA ARL READY Website,
698 <http://ready.arl.noaa.gov/HYSPLIT.php>. NOAA Air Resources Laboratory, Silver
699 Spring, MD.

700

701 European Chemical Agency (ECHA). 2008. Tris[2-Chloro-1-(Chloromethyl)ethyl]
702 Phosphate (TDCP) CAS No.: 13674-87-8. EINECS No.: 237-159-2. Risk Assessment
703 Report.

704

705 European Chemical Agency (ECHA). 2009. Member State Committee support
706 document for the identification of Tris (2-chloroethyl) phosphate as a substance of very
707 high concern because of its CMR properties.

708

709 European Chemical Agency (ECHA): [https://echa.europa.eu/fr/substance-information/-](https://echa.europa.eu/fr/substance-information/-/substanceinfo/100.004.363)
710 [/substanceinfo/100.004.363](https://echa.europa.eu/fr/substance-information/-/substanceinfo/100.004.363) (accessed on 06/04/2018)
711
712 Faiz, Y., Siddique, N., He, H., Sun, C., Waheed, S., 2018. Occurrence and profile of
713 organophosphorus compounds in fine and coarse particulate matter from two urban
714 areas of China and Pakistan. *Environ. Pollut.* 233, 26-34
715
716 Fan, X., Kubwabo, C., Rasmussen, P.E., Wu, F. 2014. Simultaneous determination of
717 thirteen organophosphate esters in settled indoor house dust and a comparison between
718 two sampling techniques. *Sci. Total Environ.* 491-492, 80–86.
719
720 Green, N.; Schlabach, M.; Bakke, T.; Brevik, E.M.; Dye, C.; Herzke, D.; Huber, S.;
721 Plosz, B.; Remberger, M.; Schøyen, M.; Uggerud, H.T; Vogelsang, C. 2008. Screening
722 of selected metals and new organic contaminants 2007. Norwegian Pollution Control
723 Authority, Report 1014/2008, ISBN- 978-82-577-5304-7.
724
725 Hallanger, I.G., Sagerupb K., Evenset, A., Kovacs, K.M., Leonards, P., Fuglei, E., Routti,
726 H., Aars, J., Strøm, H., Lydersen, C. Gabrielsen G.W., 2015. Organophosphorous flame
727 retardants in biota from Svalbard, Norway. *Mar. Pollut. Bull.* 101, 442–447
728
729 Kim, J-W., Isobe, T., Chang, K-Y., Amano, A., Maneja, R.H., Zamora, P.B., Siringan,
730 F.P., Tanabe, S., 2011. Levels and distribution of organophosphorus flame retardants
731 and plasticizers in fishes from Manila Bay, the Philippines. *Environ. Pollut.* 159, 3653-
732 3659.
733

734 Lai, S., Xie, Z., Song, T., Tang, J., Zhang, Y., Mi, W., Peng, J., Zhao, Y., Zou, S.,
735 Ebinghaus, R. 2015. Occurrence and dry deposition of organophosphate esters in
736 atmospheric particles over the northern South China Sea. *Chemosphere* 127, 195–200.
737

738 Li, J., Tang, J., Mi, W., Tian, C., Emeis, K-C., Ebinghaus, R., Xie. X., 2018. Spatial
739 distribution and seasonal variation of organophosphate esters in air above the Bohai and
740 Yellow Seas, China. *Environ. Sci. Technol.* 52, 89–97.
741

742 Liu, D., Lin, T., Shen, K., Li, J., Yu, Z., Zhang, G., 2016. Occurrence and
743 concentrations of halogenated flame retardants in the atmospheric fine particles in
744 chinese cities. *Environ. Sci. Technol.* 50, 9846–9854.
745

746 Liu, Y., Liggio, J., Harner, T., Jantunen, L., Shoeib, M., Li, S-M., 2014. Heterogeneous
747 OH Initiated Oxidation: A Possible Explanation for the Persistence of Organophosphate
748 Flame Retardants in Air. *Environ. Sci. Technol.* 48, 1041–1048.
749

750 Ma, Y., Xie, Z., Lohmann, R., Mi, W., Gao, G., 2017. Organophosphate Ester Flame
751 Retardants and Plasticizers in Ocean Sediments from the North Pacific to the Arctic
752 Ocean. *Environ. Sci. Technol.* 51, 3809–3815.
753

754 Mandalakis M., Apostolakis M., and Stephanou E.G., 2005. Mass budget and dynamics
755 of polychlorinated biphenyls in the eastern Mediterranean Sea. *Global Biogeochem. Cy.*
756 19, GB 018, 1-16
757

758 Mesquita, S.R., Dachs, J., van Drooge, B.L., Castro-Jiménez, J., Barata, C., Vieira, N.,
759 Guimarães, L., Piña, B., 2016. Toxicity assessment of atmospheric particulate matter in
760 the Mediterranean and Black Seas open waters. *Sci. Total Environ.* 545–546, 163–170.
761

762 Möller, A., Xie, Z., Caba, C., Sturm, R., Ebinghaus, R., 2011. Organophosphorus flame
763 retardants and plasticizers in the atmosphere of the North Sea. *Environ. Pollut.* 159,
764 3660-3665.
765

766 Möller, A., Sturm, R., Xie, Z., Cai, M., He, J., Ebinghaus, R., 2012. Organophosphorus
767 flame retardants and plasticizers in airborne particles over the Northern Pacific and
768 Indian Ocean towards the Polar Regions: Evidence for global occurrence. *Environ. Sci.*
769 *Technol.* 46, 3127-3134.
770

771 Paluselli, A., Fauvelle, V., Schmidt, N., Galgani, F., Net, S., Sempéré, R. 2018.
772 Distribution of phthalates in Marseille Bay (NW Mediterranean Sea). *Sci. Total*
773 *Environ.* 621, 578-587.
774

775 Raimbault P., Garcia, N. Cerrutti, F., 2008. Distribution of inorganic and organic
776 nutrients in the South Pacific Ocean. Evidence for long-term accumulation of organic
777 matter in nitrogen-depleted waters. *Biogeosciences* 5, 281-298.
778

779 Rockström, J., Steffen, W., Noone, K., Persson, Å., Chapin, F.S., Lambin, E.F., Lenton,
780 T.M., Scheffer, M., Folke, C., Schellnhuber, H.J., Nykvist, B., De Wit, C.A., Hughes,
781 T., Van Der Leeuw, S., Rodhe, H., Sörlin, S., Snyder, P.K., Costanza, R., Svedin, U.,
782 Falkenmark, M., Karlberg, L., Corell, R.W., Fabry, V.J., Hansen, J., Walker, B.,

783 Liverman, D., Richardson, K., Crutzen, P., Foley, J.A. 2009. A safe operating space for
784 humanity. *Nature* 461, 472-475.

785

786 Salamova, A., Hermanson, M. H., Hites, R. A., 2014a. Organophosphate and
787 halogenated flame retardants in atmospheric particles from a European Arctic site.
788 *Environ. Sci. Technol.* 48, 6133–6140.

789

790 Salamova, A., Ma, Y., Venter, M., Hites, R.A. 2014b High levels of organophosphate
791 flame retardants in the Great Lakes atmosphere. *Environ. Sci. Technol. Lett.* 1, 8-14.

792

793 Salamova, A.; Peverly A.A.; Venier, M.; Hites, R.A Spatial and Temporal Trends of
794 Particle Phase Organophosphate Ester Concentrations in the Atmosphere of the Great
795 Lakes. 2016. *Environ. Sci. Technol.* 50, 13249–13255

796

797 Schmidt, N., Thibault, D., Paluselli, A., Galgani, F. and R. Sempéré. Microplastics and
798 phthalates release in Mediterranean Sea. *Mermex special issue, Progress in Oceanogr.*
799 <https://doi.org/10.1016/j.pocean.2017.11.010>. In Press.

800

801 Sempéré, R., Para, J., Tedetti, M., Charriere, B., Mallet, M. 2015. Attenuation of UVR
802 and PAR in relation with chromophoric dissolved organic matter in surface coastal
803 waters of the Northwestern Mediterranean Sea. *J. Photochem. Photobiol., A: Chemistry*
804 91, 851-861.

805

806 Simes, R.J. 1986. An improved Bonferroni procedure for multiple tests of significance.
807 *Biometrika*, 73, 751-754.
808
809 Sühling, R., Diamond, M.L., Scheringer, M., Wong, F., Pućko, M., Stern, G., Burt, A.,
810 Hung, H., Fellin, P., Li, H., Jantunen, L.M., 2016a. Organophosphate esters in Canadian
811 Arctic air: Occurrence, Levels and Trends. *Environ. Sci. Technol.* 50, 7409-7415.
812
813 Sühling, R., Wolschke, H., Diamond, M.L., Jantunen, L.M., Scheringer, M. 2016b.
814 Distribution of Organophosphate Esters between the Gas and Particle Phase—Model
815 Predictions vs Measured Data. *Environ. Sci. Technol.* 50, 6644–6651.
816
817 The Mermex Group. 2011. Marine ecosystems' responses to climatic and anthropogenic
818 forcings in the Mediterranean. *Prog. Oceanogr.* 91, 97-166.
819
820 United Nations Environment Programme: Stockholm Convention on Persistent Organic
821 Pollutants: Geneva, 2009. Report of the conference of the parties of the Stockholm
822 Convention on Persistent Organic Pollutants on the work of its fourth meeting, pp 112.
823 van der Veen, I., de Boer, J., 2012. Phosphorus flame retardants: properties, production,
824 environmental occurrence, toxicity and analysis. *Chemosphere.* 88 , 1119-1153.
825
826 Violaki, K., Bourrin, F., Aubert, D., Kouvarakis, G., Delsaut, N., Mihalopoulos, N.
827 2018. Organic phosphorus in atmospheric deposition over the Mediterranean Sea: An
828 important missing piece of the phosphorus cycle. *Prog. Oceanogr.* 163, 50-58.
829

830 Vonderheide, A.P., Mueller, K.E., Meija, M., Welsch, G.L., 2008. Polybrominated
831 diphenyl ethers: Causes for concern and knowledge gaps regarding environmental
832 distribution, fate and toxicity. *Sci. Total. Environ.* 400, 425-436.
833

834 Wang, Q., Lai, N.L-S., Wang, X., Guo, Y., Lam, P. K-S., Lam, J. C-W., Zhou, B.,
835 2015a. Bioconcentration and transfer of the organophosphorus flame retardant 1,3-
836 Dichloro-2-propyl phosphate causes thyroid Endocrine disruption and developmental
837 neurotoxicity in Zebrafish larvae. *Environ. Sci. Technol.* 49, 5123–5132.
838

839 Wang, Q., Lam, J. C-W., Han, J., Wang, X., Guo, Y., Lam, P. K-S., Zhou, B., 2015b
840 Developmental exposure to the organophosphorus flame retardant tris(1,3-dichloro-2-
841 propyl) phosphate: Estrogenic activity, endocrine disruption and reproductive effects on
842 zebrafish. *Aquatic Toxicol.* 160, 163–171.
843

844 Wei, G-L., Li, D-Q., Zhuo, M-N., Liao, Y-S., Xie, Z-Y., Guo, T-L., Li, J-J., Zhang, S-
845 Y., Liang, Z-Q., 2015. Organophosphorus flame retardants and plasticizers: Sources,
846 occurrence, toxicity and human exposure. *Environ. Pollut.* 196, 29-46.
847

848 Wolschke, H.; Sühling, R.; Mi, W.; Möller, A.; Xie, Z.; Ebinghaus, R. Atmospheric
849 occurrence and fate of organophosphorus flame retardants and plasticizer at the German
850 coast. *Atmos. Environ.* 2016, 137, 1-5.
851

852 Zhong, M., Tang, J., Mi, L., Li, F., Wang, W., Huang, G., Wu, H., 2017. Occurrence
853 and spatial distribution of organophosphorus flame retardants and plasticizers in the
854 Bohai and Yellow Seas, China. *Mar. Pollut. Bull.* 121, 331–338

855

856

857

858 **Table 1.** Updated compilation of atmospheric particle-bound OPE measurements in the marine environment and some reference urban areas
 859 (concentrations are in pg m^{-3} , and median, standard deviation, minimum and maximum values are reported when available)
 860

Region	Sampling	Date	Compounds, pg m^{-3} , median [min-max]								Reference [Ⓢ]
			TCEP	TDCP	TCPPs	TIBP	TnBP	TPhP	EHDPP	TEHP	
Northern Hemisphere											
Arctic Ocean	cruise (open sea)	2007-2013	n.d. - 856	n.d. - 13	n.d. - 660	n.r.	n.d. - 97	n.d. - 1930	n.d. - 11	n.d. - 7.5	Sühring et al., 2016
Resolute Bay/Alert (Arctic)	remote (coastal)	2008-2009, 2012	n.d. - 430	n.d. - 46	n.d. - 276	n.r.	n.d. - 2340	1.2 - 96	n.d. - 40	n.r.	Sühring et al., 2016
Arctic Ocean	cruise (open sea)	Jun-September 2010	126 - 585	n.d. - 5	85 - 530	16 - 35	n.d. - 36	10 - 60	n.r.	n.d. - 6	Moller et al., 2012
Ny-Alesund (Arctic)	remote (coastal)	Jun-September 2007	<200 - 270	87 - 250	<200 - 330	<10 - 140	<200	<50	<200 - 260	n.r.	Green et al., 2008
Longyearbyen (Arctic)	remote (coastal)	September 2012-May 2013	40 - 60	2 - 294	10 - 186	n.r.	6 - 1000	1 - 50	6 - 300	1 - 40	Salamova et al., 2014
North Atlantic Ocean	cruise (open sea)	Dec 2010, Jun-Jul 2011	50 [n.d.-1230]	80 [n.d.-425]	770 [n.d.-1310]	40 [5-380]	90 [10-1700]	10 [n.d.-50]	780 [20-1730]	140 [60 - 490]	Castro-Jiménez et al., 2016
North Pacific Ocean	cruise (open sea)	Jun-September 2010	160 - 280	5 - 8	98 - 270	14 - 20	6 - 14	9 - 24	n.r.	1 - 12	Moller et al., 2012
North Pacific Ocean	cruise (open sea)	May-Jun 2011	80 [n.d.-310]	90 [n.d.-500]	640 [100-1460]	30 [3-100]	170 [20-2500]	10 [n.d.-34]	320 [100-1210]	110 [60-380]	Castro-Jiménez et al., 2016
Sea of Japan	cruise (open sea)	Jun-September 2010	237 - 1960	16 - 52	130 - 620	10 - 63	10 - 33	25 - 97	n.r.	5 - 38	Möller et al., 2012
East China Sea	cruise (open sea)	October 2009-March 2010	134	828	9	n.r.	n.r.	n.r.	n.r.	n.r.	Möller et al., 2012 & Chen et al., 2013
South China Sea	cruise (open sea)	September-October 2013	14 - 110	1 - 4	15 - 38	1 - 4	1 - 5	3 - 16	n.r.	2 - 16	Lai et al., 2015
North Sea	cruise (open sea)	March, May, July 2010	6 - 100	7 - 78	30 - 1200	n.d. - 150	n.d. - 150	4 - 150	n.r.	n.d. - 30	Möller et al., 2011
North Sea [Ⓢ]	coastal	August 2011-October 2012	3±2	1±0.8	10±6	9±6	6±11	3±3	n.r.	11±13	Wolschke et al., 2016
Mediterranean Sea	cruise (open sea)	Jun 2006, May 2007	220 [Ⓢ] [70 - 854]	75 [Ⓢ] [n.d. - 460]	884 [Ⓢ] [126 - 2340] [Ⓢ]	240 [Ⓢ] [4 - 650]	310 [Ⓢ] [56 - 600]	21 [Ⓢ] [n.d. - 80]	540 [Ⓢ] [n.d. - 834]	160 [Ⓢ] [56 - 307]	Castro-Jiménez et al., 2014
Black Sea	cruise (open sea)	Jun 2006, May 2007	490 [300 - 2420]	86 [n.d. - 97]	520 [540 - 2720] [Ⓢ]	150 [66 - 190]	310 [200 - 370]	35 [3 - 40]	180 [n.d. - 310]	175 [36 - 190]	Castro-Jiménez et al., 2014
Philippine Sea	cruise (open sea)	November 2010- March 2011	20 - 156	50 - 780	22 - 410	10 - 23	10 - 100	n.d. - 155	n.r.	6 - 92	Möller et al., 2012
Bonhai & Yellow Seas	cruise (open sea)	Jun-July 2016	28 [14-94]	5 [1-13]	66 [19-390] [Ⓢ]	12 [24-29]	5 [1-14]	3 [1-15]	n.r.	2 [n.d.-8.5]	Li et al., 2018
North Huangchen Island	remote (coastal)	May 2015-March 2016	63 [0.4-1000]	4 [n.d.-80]	26 [n.n-160] [Ⓢ]	24 [1-164]	8 [n.d-1100]	11 [n.d-110]	n.r.	1.5 [n.d-180]	Li et al., 2018
Huston [Ⓢ]	urban (inland)	September 2013	90±30 - 270±130	64±20 - 87±50	700±390 - 1200±440	n.r.	120±10 - 560±320	210±70 - 490±580	300±260 - 470±150	38±30 - 56±20	Clark et al., 2017
Great lakes [Ⓢ]	urban (coastal inland)	March 2012 - Dec 2014	270-334	47-100	370-390	n.r.	200-250	130-160	70-80	n.r.	Salamova et al., 2016
Great lakes [Ⓢ]	rural (coastal inland)	March 2012 - Dec 2014	274	15	58	n.r.	32	30	27	n.r.	Salamova et al., 2016
Great lakes [Ⓢ]	remote (coastal inland)	March 2012 - Dec 2014	41-46	7-11	15-36	n.r.	18-50	34-50	15	n.r.	Salamova et al., 2016
Bizerte (Tunisia)	urban (coastal)	March 2015-Jan 2016	31 [2-96]	20 [7-27]	110 [5-347][Ⓢ]	85 [8-528]	36 [1-116]	27 [3-538]	100 [21-429]	45 [20-492]	This study
Southern Hemisphere											
South Atlantic	cruise (open sea)	Jan-Feb 2011	150 [10-540]	130 [n.d.-540]	570 [20-980]	100 [30-280]	330 [120-1180]	10 [n.d.-25]	500 [n.d.-1020]	160 [50-890]	Castro-Jiménez et al., 2016
South Pacific	cruise (open sea)	Feb-April 2011	140 [34-370]	60 [n.d.-1000]	530 [50-800]	50 [15-160]	200 [50-2170]	4 [n.d.-40]	400 [260-800]	160 [40-350]	Castro-Jiménez et al., 2016
Coral Sea	cruise (open sea)	October 2009-March 2010	88	370	7	n.r.	n.r.	n.r.	n.r.	n.r.	Chen et al., 2013
Indian Ocean	cruise (open sea)	November 2010- March 2011	46 - 570	n.d. - 220	37 - 550	7 - 96	7 - 75	n.d. - 74	n.r.	4 - 50	Möller et al., 2012
Indian Ocean	cruise (open sea)	Feb-March 2011	100 [50-620]	20 [n.d.-290]	370 [30-1250]	40 [n.d.-110]	230 [70-940]	8 [n.d.-12]	370 [n.d-630]	180 [n.d-630]	Castro-Jiménez et al., 2016
Southern Ocean	cruise (open sea)	November 2010- March 2011	74	80	55	16	14	20	n.r.	7	Möller et al., 2012
Antarctic Peninsula	cruise (open sea)	October 2009-March 2010	40	76	4	n.r.	n.r.	n.r.	n.r.	n.r.	Chen et al., 2013

[Ⓢ] sum of three isomers; [Ⓢ] sum of two major isomers; [Ⓢ] average±SD; * median values; [Ⓢ] median value correspond only to Western MED, whereas the range correspond to all basins; [Ⓢ] see reference section in the main manuscript for complete citation details, n.r.=not reported

861

862

863
864

Table 2. OPE dry deposition fluxes ($\text{ng m}^{-2} \text{d}^{-1}$, median, average and range) in the different seasons (2015-2016) in Bizerte (Tunisia)

Compound	Spring [n=19]			Summer [n=20]			Autumn [n=13]			Winter [n=8]		
	Median	Average	Range	Median	Average	Range	Median	Average	Range	Median	Average	Range
TPP	0.7	0.6	(0.01 - 1.0)	0.3 ^b	0.3 ^b	/ ^b	n.c	n.c	n.c	1.4 ^b	1.4 ^b	/ ^b
TiBP	4.8	8.6	(1.7 - 30.0)	16.2	29.5	(1.4 - 91.2)	17.1	18.1	(5.8 - 38.3)	13.8	19.3	(3.6 - 46.1)
TnBP	6.1	5.6	(2.5 - 8.7)	9.9	9.8	(2.1 - 20.0)	6.5	7.5	(0.2 - 16.6)	2.8	3.4	(0.9 - 7.6)
TCEP	3.4	3.5	(0.6 - 7.7)	9.7	9.7	(4.8 - 15.8)	7.9	7.7	(1.1 - 16.5)	2.8	2.7	(0.4 - 4.7)
TCPPs ^a	7.3	12.6	(0.8 - 58.9)	18.2	24.2	(6.9 - 56.1)	31.4	31.7	(9.2 - 59.9)	25.5	27.5	(3.5 - 59.9)
TDCP	4.3 ^b	4.3 ^b	/ ^b	3.3	3.3	(1.8 - 4.6)	2.3	2.3	(1.1 - 3.4)	n.c	n.c	n.c
TPhP	4.3	7.1	(0.5 - 24.8)	4.2	5.7	(2.1 - 25.0)	5.4	15.1	(0.9 - 93.0)	7.1	7.4	(2.3 - 14.6)
EHDPP	11.0	11.9	(3.7 - 33.4)	21.4	23.3	(8.6 - 49.0)	20.4	25.4	(6.6 - 74.2)	14.7	18.5	(4.4 - 47.8)
TEHP	6.7	7.0	(3.4 - 13.1)	8.6	8.9	(5.1 - 18.2)	7.2	9.5	(3.5 - 31.8)	21.9	27.4	(5.1 - 85.0)
Σ_9OPEs	48.0	49.5	(17.8 - 109.3)	103.7	107.5	(53.2 - 172.5)	93.3	108.0	(66.8 - 183.3)	89.2	95.0	(37.7 - 162.5)

865

^a sum of the two major TCPP isomers; ^b only one detected value in the season (the values presented under median and mean correspond in this case to the only detected value), n.c = not calculated (aerosol concentrations \leq LOD)