



**HAL**  
open science

# Regularity results for the solutions of a non-local model of traffic

Florent Berthelin, Paola Goatin

► **To cite this version:**

Florent Berthelin, Paola Goatin. Regularity results for the solutions of a non-local model of traffic. Discrete and Continuous Dynamical Systems - Series A, 2019, 39 (6), pp.3197-3213. hal-01813760

**HAL Id: hal-01813760**

**<https://hal.science/hal-01813760v1>**

Submitted on 12 Jun 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Regularity results for the solutions of a non-local model of traffic flow

FLORENT BERTHELIN\*      PAOLA GOATIN†

June 12, 2018

## Abstract

We consider a non-local traffic model involving a convolution product. Unlike other studies, the considered kernel is discontinuous on  $\mathbb{R}$ . We prove Sobolev estimates and prove the convergence of approximate solutions solving a viscous and regularized non-local equation. It leads to weak,  $\mathbf{C}([0, T], \mathbf{L}^2(\mathbb{R}))$ , and smooth,  $\mathbf{W}^{2,2N}([0, T] \times \mathbb{R})$ , solutions for the non-local traffic model.

**Key words:** Scalar conservation laws; Anisotropic non-local flux; Traffic flow models; Viscous approximation; Sobolev estimates.

## 1 Introduction

We consider the non-local traffic model introduced in [4, 8] to account for the reaction of drivers to downstream traffic conditions. It consists in the following scalar conservation law, where the traffic velocity depends on a weighted mean of the density:

$$\partial_t \rho + \partial_x(\rho v(\rho * \omega)) = 0, \quad (1.1)$$

where

$$(\rho * \omega)(t, x) = \int_0^\eta \rho(t, x + y)\omega(y) dy = \int_x^{x+\eta} \rho(t, y)\omega(y - x) dy. \quad (1.2)$$

We make the following assumptions for  $k = 1, 2, 3$ :

( $\mathbf{A}_\omega^k$ )  $\omega \in \mathbf{C}^k([0, \eta])$  is non-negative with support in  $[0, \eta]$  and is non-increasing on  $[0, \eta]$ .

( $\mathbf{A}_v^k$ )  $v \in \mathbf{C}^k(\mathbb{R}^+)$  with  $v', \dots, v^{(k)}$  bounded.

For traffic flow applications, it is reasonable to assume that  $v$  is non-increasing, even if monotonicity is not required in this paper. We also recall that a similar model, considering a weighted mean of downstream speeds, has been recently introduced in [7]. More generally, model (1.1) belongs to the class of conservation laws with non-local flux functions, which appear in several applications, see for example [3, 6, 9, 10, 15]. We remark that most of the available well-posedness results concern equations involving smooth convolution kernels [1, 2], and are based on the construction of finite-volume approximations and the use of Kruřkov's doubling of variable technique [12]. In particular, these results rely on the concept of entropy solutions. Only recently, alternative proofs based on fixed point theorems have been proposed for specific cases [11, 14], allowing to get rid of the entropy requirement.

In general, solutions to non-local equations may be discontinuous [13], despite the expected regularizing effect of the convolution product. Therefore, given any initial datum  $\rho^0 \in \mathbf{L}^\infty(\mathbb{R}) \cap \mathbf{L}^1(\mathbb{R})$ , the solutions to the Cauchy problem for (1.1) are usually intended in the following weak form

---

\*Laboratoire J. A. Dieudonné, UMR 7351 CNRS, Université Côte d'Azur, LJAD, CNRS, Inria, Université de Nice Sophia-Antipolis, Parc Valrose, 06108 Nice cedex 2, France. E-mail: [Florent.Berthelin@unice.fr](mailto:Florent.Berthelin@unice.fr)

†Inria Sophia Antipolis - Méditerranée, Université Côte d'Azur, Inria, CNRS, LJAD, 2004 route des Lucioles - BP 93, 06902 Sophia Antipolis Cedex, France. E-mail: [paola.goatin@inria.fr](mailto:paola.goatin@inria.fr)

**Definition 1.** A function  $\rho \in (\mathbf{L}^\infty \cap \mathbf{L}^1)(\mathbb{R}^+ \times \mathbb{R})$  is a solution of (1.1) with initial datum  $\rho^0$  if

$$\int_0^{+\infty} \int_{-\infty}^{+\infty} (\rho \partial_t \varphi + \rho v(\rho * \omega) \partial_x \varphi)(t, x) dx dt + \int_{-\infty}^{+\infty} \rho^0(x) \varphi(0, x) dx = 0, \quad (1.3)$$

1 for all  $\varphi \in \mathbf{C}_c^\infty(\mathbb{R}^2)$ .

In this paper, we are interested in deriving regularity properties of solutions to (1.1). To this end, we will consider approximate solutions satisfying the viscous and regularized non-local equation

$$\partial_t \rho_\varepsilon + \partial_x(\rho_\varepsilon v(\rho_\varepsilon * \omega_\varepsilon)) = \varepsilon \partial_{xx}^2 \rho_\varepsilon, \quad (1.4)$$

2 where, for any  $\varepsilon \in ]0, 1]$ , the smooth function  $\omega_\varepsilon$  is an extension of  $\omega$  with the following regularities:

( $\mathbf{A}_{\omega_\varepsilon}^k$ )  $\omega_\varepsilon \in \mathbf{C}^k(\mathbb{R})$  is non-negative with a support in  $[-\varepsilon, \eta + \varepsilon]$ , is non-decreasing on  $[-\varepsilon, x_\varepsilon]$ , for some  $x_\varepsilon \in ]-\varepsilon, 0]$ , is non-increasing on  $[x_\varepsilon, \eta + \varepsilon]$  and  $\omega_\varepsilon = \omega$  on  $[0, \eta]$ .

We set  $W_\varepsilon := \omega_\varepsilon(x_\varepsilon)$  and we assume that  $\lim_{\varepsilon \rightarrow 0} W_\varepsilon = \omega(0)$ . Without loss of generality we can assume

$$W_\varepsilon \leq 2\omega(0). \quad (1.5)$$

3 ( $\mathbf{B}_{\omega_\varepsilon}^k$ )  $\omega_\varepsilon^{(j)}(-\varepsilon) = \omega_\varepsilon^{(j)}(\eta + \varepsilon) = 0$  for  $j = 1, \dots, k$  and  $|\omega_\varepsilon'(u)| \leq 2W_\varepsilon/\varepsilon$  on  $[-\varepsilon, 0]$  and  $|\omega_\varepsilon'(u)| \leq$   
4  $2\omega(\eta)/\varepsilon$  on  $[\eta, \eta + \varepsilon]$ .

5 **Remark 1.** Given  $\omega$  satisfying ( $\mathbf{A}_\omega^k$ ), we can construct a function  $\omega_\varepsilon$  satisfying ( $\mathbf{A}_{\omega_\varepsilon}^k$ ) and ( $\mathbf{B}_{\omega_\varepsilon}^k$ ).  
6 To construct such extensions, for example in the simplest case where the derivatives of  $\omega$  vanish at  
7 0 and  $\eta$ , we use the function  $\varphi$  which is zero for  $x \leq -1$ , 1 for  $x \geq 0$ , non-decreasing and of class  
8  $C^\infty$  and we define  $\omega_\varepsilon(x) = \omega(0)\varphi(x/\varepsilon)$  for  $x < 0$  and  $\omega(0)\varphi((\eta - x)/\varepsilon)$  for  $x > \eta$ .

9 Notice that a similar approximation was used in [5] to establish a convergence property for the  
10 singular limit where the (smooth) convolution kernel is replaced by a Dirac delta, in the viscous  
11 case. Here, we will study the properties of smooth solutions  $\rho_\varepsilon$  of this equation corresponding to a  
12 fixed initial datum  $\rho^0$ , and then we will recover properties for  $\rho$  passing to the limit as  $\varepsilon \rightarrow 0$ .

13 We have the following result.

14 **Theorem 1.** We assume ( $\mathbf{A}_\omega^2$ )-( $\mathbf{A}_v^3$ ). Let  $\rho_\varepsilon$  be a solution of (1.4) with initial datum  $\rho^0$ . We  
15 assume  $\rho^0 \in \mathbf{W}^{1,4}(\mathbb{R}) \cap \mathbf{H}^2(\mathbb{R})$ . Then, for  $T > 0$  sufficiently small,  $\rho_\varepsilon$  converges in  $\mathbf{L}_{\text{loc}}^2([0, T] \times \mathbb{R})$   
16 to a solution  $\rho \in \mathbf{C}([0, T], \mathbf{L}^2(\mathbb{R}))$  to equation (1.1) with initial datum  $\rho^0$ . Furthermore, if  $\rho^0 \in$   
17  $\mathbf{W}^{1,2N}(\mathbb{R})$ ,  $N \in \mathbb{N}^*$ , then  $\rho \in \mathbf{W}^{1,2N}([0, T] \times \mathbb{R})$ , and if  $\rho^0 \in \mathbf{W}^{1,4N}(\mathbb{R}) \cap \mathbf{H}^1(\mathbb{R}) \cap \mathbf{W}^{2,2N}(\mathbb{R})$ , then  
18  $\rho \in \mathbf{W}^{2,2N}([0, T] \times \mathbb{R})$ .

19 In particular, this provides an alternative proof of existence of weak solutions, locally in time.  
20 To prove this result, in Section 2 we first establish estimates on the non-local term and we derive  
21  $\mathbf{L}^p(\mathbb{R})$ ,  $p > 1$ , estimates for  $\rho_\varepsilon$ , then we get estimates in  $\mathbf{W}^{1,2N}(\mathbb{R})$  for  $\rho_\varepsilon$  with respect to  $x$ . This  
22 allows to prove that there exists  $T > 0$  such that the sequence  $\rho_\varepsilon$  is uniformly bounded with respect  
23 to  $\varepsilon$  in  $\mathbf{L}^\infty(\mathbb{R})$  on  $[0, T]$ . Then we prove uniform space estimates in  $\mathbf{W}^{2,2N}(\mathbb{R})$  for  $\rho_\varepsilon$ , which allows  
24 to derive estimates on  $\partial_t \rho_\varepsilon$ . The proof of Theorem 1 is deferred to Section 3.

## 2 Estimates

Here and in the following sections, we will denote by

$$\|\rho\|_\infty := \|\rho\|_{\mathbf{L}^\infty([0,T] \times \mathbb{R})}$$

and by

$$\|\rho(t, \cdot)\|_\infty := \|\rho(t, \cdot)\|_{\mathbf{L}^\infty(\mathbb{R})}.$$

Moreover, notice that we have

$$\begin{aligned} (\rho * \omega_\varepsilon)(t, x) &= \int_{\mathbb{R}} \rho(t, x+y) \omega_\varepsilon(y) dy = \int_{\mathbb{R}} \rho(t, y) \omega_\varepsilon(y-x) dy \\ &= \int_{-\varepsilon}^{\eta+\varepsilon} \rho(t, x+y) \omega_\varepsilon(y) dy = \int_{x-\varepsilon}^{x+\eta+\varepsilon} \rho(t, y) \omega_\varepsilon(y-x) dy. \end{aligned}$$

### 2.1 Estimates of the non-local term

We start by proving the following estimates on the non-local term.

**Proposition 1.** 1. We assume  $(\mathbf{A}_{\omega_\varepsilon}^1)$  and that  $\rho$  is a continuous function. For any  $(t, x) \in \mathbb{R}^+ \times \mathbb{R}$ , we have

$$|\partial_x(\rho * \omega_\varepsilon)(t, x)| \leq 2\|\rho(t, \cdot)\|_\infty W_\varepsilon. \quad (2.1)$$

2. We assume  $(\mathbf{A}_{\omega_\varepsilon}^2) - (\mathbf{B}_{\omega_\varepsilon}^1)$  and that  $\rho$  is a  $C^1$  function. For any  $t \geq 0$ ,  $p > 1$ , we have

$$\begin{aligned} \left( \int_{\mathbb{R}} |\partial_{xx}^2(\rho * \omega_\varepsilon)|^p(t, x) dx \right)^{1/p} &\leq \eta^{1/p} \left( \int_0^\eta |\omega''(u)|^{p/(p-1)} du \right)^{1-1/p} \left( \int_{\mathbb{R}} \rho^p(t, y) dy \right)^{1/p} \\ &\quad + (|\omega'(\eta-)| + |\omega'(0+)|) \left( \int_{\mathbb{R}} \rho^p(t, x) dx \right)^{1/p} \\ &\quad + 2(\omega(\eta) + W_\varepsilon) \left( \int_{\mathbb{R}} |\partial_x \rho(t, x)|^p dx \right)^{1/p}. \end{aligned} \quad (2.2)$$

3. We assume  $(\mathbf{A}_\omega^2) - (\mathbf{B}_{\omega_\varepsilon}^2)$  that  $\rho$  is a  $C^2$  function. For any  $t \geq 0$ ,  $p > 1$ , we have

$$\begin{aligned} \left( \int_{\mathbb{R}} |\partial_{xxx}^3(\rho * \omega_\varepsilon)|^p(t, x) dx \right)^{1/p} &\leq \eta^{1/p} \left( \int_0^\eta |\omega''(u)|^{p/(p-1)} du \right)^{1-1/p} \left( \int_{\mathbb{R}} |\partial_x \rho(t, y)|^p dy \right)^{1/p} \\ &\quad + (|\omega'(\eta-)| + |\omega'(0+)|) \left( \int_{\mathbb{R}} |\partial_x \rho(t, x)|^p dx \right)^{1/p} \\ &\quad + 2(\omega(\eta) + W_\varepsilon) \left( \int_{\mathbb{R}} |\partial_{xx}^2 \rho(t, x)|^p dx \right)^{1/p}. \end{aligned} \quad (2.3)$$

*Proof.* 1. From

$$\begin{aligned} \partial_x(\rho * \omega_\varepsilon)(t, x) &= - \int_{x-\varepsilon}^{x+\eta+\varepsilon} \rho(t, y) \omega'_\varepsilon(y-x) dy + \rho(t, x+\eta+\varepsilon) \omega_\varepsilon(\eta+\varepsilon) - \rho(t, x-\varepsilon) \omega_\varepsilon(-\varepsilon) \\ &= - \int_{x-\varepsilon}^{x+\eta+\varepsilon} \rho(t, y) \omega'_\varepsilon(y-x) dy = - \int_{-\varepsilon}^{\eta+\varepsilon} \rho(t, u+x) \omega'_\varepsilon(u) du, \end{aligned}$$

we obtain

$$\begin{aligned}
|\partial_x(\rho * \omega_\varepsilon)(t, x)| &\leq \|\rho(t, \cdot)\|_\infty \int_{-\varepsilon}^{\eta+\varepsilon} |\omega'_\varepsilon(u)| du \\
&\leq \|\rho(t, \cdot)\|_\infty \left( \int_{-\varepsilon}^{x_\varepsilon} \omega'_\varepsilon(u) du - \int_{x_\varepsilon}^{\eta+\varepsilon} \omega'_\varepsilon(u) du \right) \\
&\leq 2\|\rho(t, \cdot)\|_\infty W_\varepsilon.
\end{aligned}$$

2. From

$$\begin{aligned}
\partial_{xx}^2(\rho * \omega_\varepsilon)(t, x) &= \int_{x-\varepsilon}^{x+\eta+\varepsilon} \rho(t, y) \omega''_\varepsilon(y-x) dy - \rho(t, x+\eta+\varepsilon) \omega'_\varepsilon(\eta+\varepsilon) + \rho(t, x-\varepsilon) \omega'_\varepsilon(-\varepsilon) \\
&= \int_{x-\varepsilon}^{x+\eta+\varepsilon} \rho(t, y) \omega''_\varepsilon(y-x) dy = \int_{-\varepsilon}^{\eta+\varepsilon} \rho(t, x+u) \omega''_\varepsilon(u) du \\
&= \int_{-\varepsilon}^0 \rho(t, x+u) \omega''_\varepsilon(u) du + \int_0^\eta \rho(t, x+u) \omega''_\varepsilon(u) du + \int_\eta^{\eta+\varepsilon} \rho(t, x+u) \omega''_\varepsilon(u) du \\
&= \rho(t, x) \omega'_\varepsilon(0) - \rho(t, x-\varepsilon) \omega'_\varepsilon(-\varepsilon) - \int_{-\varepsilon}^0 \partial_x \rho(t, x+u) \omega'_\varepsilon(u) du \\
&\quad + \int_0^\eta \rho(t, x+u) \omega''_\varepsilon(u) du \\
&\quad + \rho(t, x+\eta+\varepsilon) \omega'_\varepsilon(\eta+\varepsilon) - \rho(t, x+\eta) \omega'_\varepsilon(\eta) - \int_\eta^{\eta+\varepsilon} \partial_x \rho(t, x+u) \omega'_\varepsilon(u) du \\
&= \rho(t, x) \omega'(0+) - \rho(t, x+\eta) \omega'(\eta-) + \int_0^\eta \rho(t, x+u) \omega''_\varepsilon(u) du \\
&\quad - \int_{-\varepsilon}^0 \partial_x \rho(t, x+u) \omega'_\varepsilon(u) du - \int_\eta^{\eta+\varepsilon} \partial_x \rho(t, x+u) \omega'_\varepsilon(u) du,
\end{aligned}$$

we have

$$\begin{aligned}
&\left( \int_{\mathbb{R}} |\partial_{xx}^2(\rho * \omega_\varepsilon)|^p(t, x) dx \right)^{1/p} \\
&\leq \left( \int_{\mathbb{R}} \left| \int_{-\varepsilon}^0 \partial_x \rho(t, x+u) \omega'_\varepsilon(u) du \right|^p dx \right)^{1/p} \\
&\quad + \left( \int_{\mathbb{R}} \left| \int_\eta^{\eta+\varepsilon} \partial_x \rho(t, x+u) \omega'_\varepsilon(u) du \right|^p dx \right)^{1/p} \\
&\quad + \left( \int_{\mathbb{R}} \left| \int_0^\eta \rho(t, x+u) \omega''_\varepsilon(u) du \right|^p dx \right)^{1/p} \\
&\quad + \left( \int_{\mathbb{R}} \rho(t, x)^p |\omega'(0+)|^p dx \right)^{1/p} + \left( \int_{\mathbb{R}} \rho(t, x+\eta)^p |\omega'(\eta-)|^p dx \right)^{1/p}.
\end{aligned}$$

Notice that

$$\left( \int_{\mathbb{R}} \left| \int_{-\varepsilon}^0 \partial_x \rho(t, x+u) \omega'_\varepsilon(u) du \right|^p dx \right)^{1/p}$$

$$\begin{aligned}
&\leq \left( \int_{\mathbb{R}} \left( \int_{-\varepsilon}^0 |\partial_x \rho(t, x+u)|^p du \right) \left( \int_{-\varepsilon}^0 |\omega'_\varepsilon(u)|^q dy \right)^{p/q} dx \right)^{1/p} \\
&\leq \left( \int_{-\varepsilon}^0 |\omega'_\varepsilon(u)|^{p/(p-1)} du \right)^{1-1/p} \left( \int_{\mathbb{R}} \left( \int_{-\varepsilon}^{\eta+\varepsilon} |\partial_x \rho(t, x+u)|^p du \right) dx \right)^{1/p} \\
&\leq \left( \int_{-\varepsilon}^0 \left( 2 \frac{W_\varepsilon}{\varepsilon} \right)^{p/(p-1)} du \right)^{1-1/p} \left( \int_{\mathbb{R}} \int_{x-\varepsilon}^x |\partial_x \rho(t, y)|^p dy dx \right)^{1/p} \\
&\leq 2 \frac{W_\varepsilon}{\varepsilon} \varepsilon^{1-1/p} \left( \int_{\mathbb{R}} \int_y^{y+\varepsilon} dx |\partial_x \rho(t, y)|^p dy \right)^{1/p} \\
&\leq 2 \frac{W_\varepsilon}{\varepsilon} \varepsilon^{1-1/p} \varepsilon^{1/p} \left( \int_{\mathbb{R}} |\partial_x \rho(t, y)|^p dy \right)^{1/p} = 2W_\varepsilon \left( \int_{\mathbb{R}} |\partial_x \rho(t, y)|^p dy \right)^{1/p}
\end{aligned}$$

using Hölder's inequality with  $q = p/(p-1)$  the conjugated exponent of  $p$ . Similarly

$$\left( \int_{\mathbb{R}} \left| \int_{\eta}^{\eta+\varepsilon} \partial_x \rho(t, x+u) \omega'_\varepsilon(u) du \right|^p dx \right)^{1/p} \leq 2\omega(\eta) \left( \int_{\mathbb{R}} |\partial_x \rho(t, y)|^p dy \right)^{1/p}.$$

Then we get

$$\begin{aligned}
&\left( \int_{\mathbb{R}} |\partial_{xx}^2(\rho * \omega_\varepsilon)|^p(t, x) dx \right)^{1/p} \\
&\leq 2(\omega(\eta) + W_\varepsilon) \left( \int_{\mathbb{R}} |\partial_x \rho(t, y)|^p dy \right)^{1/p} \\
&\quad + \left( \int_{\mathbb{R}} \left( \int_x^{x+\eta} \rho^p(t, y) dy \right) \left( \int_0^\eta |\omega''_\varepsilon(u)|^q du \right)^{p/q} dx \right)^{1/p} \\
&\quad + (|\omega'(\eta+)| + |\omega'(0-)|) \left( \int_{\mathbb{R}} \rho^p(t, x) dx \right)^{1/p}.
\end{aligned}$$

Furthermore

$$\begin{aligned}
&\int_{\mathbb{R}} \left( \int_x^{x+\eta} \rho^p(t, y) dy \right) \left( \int_0^\eta |\omega''(u)|^q dy \right)^{p-1} dx \\
&\leq \left( \int_0^\eta |\omega''(u)|^{p/(p-1)} du \right)^{p-1} \int_{\mathbb{R}} \int_{y-\eta}^y \rho^p(t, y) dx dy \\
&\leq \eta \left( \int_0^\varepsilon |\omega''(u)|^{p/(p-1)} du \right)^{p-1} \int_{\mathbb{R}} \rho^p(t, y) dy,
\end{aligned}$$

1 then we get the announced formula.

3. Remark that, since  $\omega''_\varepsilon(-\varepsilon) = \omega''_\varepsilon(\eta + \varepsilon) = 0$ , we have

$$\partial_{xxx}^3(\rho * \omega_\varepsilon)(t, x) = - \int_{-\varepsilon}^{\eta+\varepsilon} \rho(t, x+u) \omega_\varepsilon^{(3)}(u) du = \int_{-\varepsilon}^{\eta+\varepsilon} \partial_x \rho(t, x+u) \omega''_\varepsilon(u) du$$

$$= \partial_{xx}^2 (\partial_x \rho * \omega_\varepsilon)(t, x),$$

1 then applying 2., we get

$$\begin{aligned} \left( \int_{\mathbb{R}} |\partial_{xxx}^3 (\rho * \omega)|^p(t, x) dx \right)^{1/p} &\leq \eta^{1/p} \left( \int_0^\eta |\omega''(u)|^{p/(p-1)} du \right)^{1-1/p} \left( \int_{\mathbb{R}} |\partial_x \rho(t, y)|^p dy \right)^{1/p} \\ &\quad + (|\omega'(\eta-)| + |\omega'(0+)|) \left( \int_{\mathbb{R}} |\partial_x \rho(t, x)|^p dx \right)^{1/p} \\ &\quad + 2(\omega(\eta) + W_\varepsilon) \left( \int_{\mathbb{R}} |\partial_{xx}^2 \rho(t, x)|^p dx \right)^{1/p}. \end{aligned}$$

2

□

### 3 2.2 $\mathbf{L}^p$ estimates for the viscous case

4 We turn now to estimates on solutions solving the viscous and regularized non-local equation. First,  
5 we deal with  $\mathbf{L}^p$  estimates.

**Proposition 2.** *We assume  $(\mathbf{A}_\omega^1)$ - $(\mathbf{A}_v^1)$ . Let  $\rho_\varepsilon$  be the solution of (1.4) with initial datum  $\rho^0 \in \mathbf{L}^p(\mathbb{R})$ . If  $\rho_\varepsilon \in \mathbf{L}^\infty([0, T] \times \mathbb{R})$  for some  $T > 0$ , then*

$$\rho_\varepsilon \in \mathbf{L}^\infty([0, T], \mathbf{L}^p(\mathbb{R})) \cap \mathbf{L}^p([0, T] \times \mathbb{R}).$$

*Proof.* The equation (1.4) can be rewritten as

$$\partial_t \rho_\varepsilon + v(\rho_\varepsilon * \omega_\varepsilon) \partial_x \rho_\varepsilon + \rho_\varepsilon v'(\rho_\varepsilon * \omega_\varepsilon) \partial_x (\rho_\varepsilon * \omega_\varepsilon) = \varepsilon \partial_{xx}^2 \rho_\varepsilon. \quad (2.4)$$

6 Multiplying (2.4) by  $\rho_\varepsilon^{p-1}$ , then integrating with respect to  $x$ , we obtain

$$\begin{aligned} \frac{1}{p} \frac{d}{dt} \int_{\mathbb{R}} \rho_\varepsilon^p(t, x) dx &= - \int_{\mathbb{R}} \rho_\varepsilon^{p-1}(t, x) v((\rho_\varepsilon * \omega_\varepsilon)(t, x)) \partial_x \rho_\varepsilon(t, x) dx \\ &\quad - \int_{\mathbb{R}} \rho_\varepsilon^p(t, x) v'((\rho_\varepsilon * \omega_\varepsilon)(t, x)) \partial_x (\rho_\varepsilon * \omega_\varepsilon)(t, x) dx \\ &\quad + \varepsilon \int_{\mathbb{R}} \rho_\varepsilon^{p-1}(t, x) \partial_{xx}^2 \rho_\varepsilon(t, x) dx. \end{aligned}$$

7 We observe that

$$\begin{aligned} \int_{\mathbb{R}} \rho_\varepsilon^{p-1} v(\rho_\varepsilon * \omega_\varepsilon) \partial_x \rho_\varepsilon dx &= \int_{\mathbb{R}} \partial_x \left( \frac{\rho_\varepsilon^p}{p} \right) v(\rho_\varepsilon * \omega_\varepsilon) dx \\ &= - \int_{\mathbb{R}} \frac{\rho_\varepsilon^p}{p} \partial_x (v(\rho_\varepsilon * \omega_\varepsilon)) dx \\ &= - \int_{\mathbb{R}} \frac{\rho_\varepsilon^p}{p} v'(\rho_\varepsilon * \omega_\varepsilon) \partial_x (\rho_\varepsilon * \omega_\varepsilon) dx \end{aligned}$$

8 and

$$\int_{\mathbb{R}} \rho_\varepsilon^{p-1} \partial_{xx}^2 \rho_\varepsilon dx = -(p-1) \int_{\mathbb{R}} \rho_\varepsilon^{p-2} (\partial_x \rho_\varepsilon)^2 dx \leq 0,$$

therefore

$$\frac{d}{dt} \int_{\mathbb{R}} \rho_{\varepsilon}^p(t, x) dx \leq (1-p) \int_{\mathbb{R}} \rho_{\varepsilon}^p(t, x) v'((\rho_{\varepsilon} * \omega_{\varepsilon})(t, x)) \partial_x(\rho_{\varepsilon} * \omega_{\varepsilon})(t, x) dx. \quad (2.5)$$

We use (2.1) to control the right hand side of (2.5) and we get

$$\frac{d}{dt} \int_{\mathbb{R}} \rho_{\varepsilon}^p(t, x) dx \leq C_1^{\varepsilon, p} \int_{\mathbb{R}} \rho_{\varepsilon}^p(t, x) dx, \quad (2.6)$$

which implies

$$\int_{\mathbb{R}} \rho_{\varepsilon}^p(t, x) dx \leq e^{C_1^{\varepsilon, p} t} \int_{\mathbb{R}} \rho_{\varepsilon}^p(0, x) dx, \quad (2.7)$$

with

$$C_1^{\varepsilon, p} = 2(p-1) \|\rho_{\varepsilon}\|_{\infty} W_{\varepsilon} \|v'\|_{\infty}.$$

It gives

$$\sup_{t \in [0, T]} \int_{\mathbb{R}} \rho_{\varepsilon}^p(t, x) dx \leq e^{C_1^{\varepsilon, p} T} \int_{\mathbb{R}} \rho_{\varepsilon}^p(0, x) dx. \quad (2.8)$$

By integration of (2.7) with respect to  $t \in [0, T]$ , we get

$$\int_0^T \int_{\mathbb{R}} \rho_{\varepsilon}^p(t, x) dx dt \leq \frac{1}{C_1^{\varepsilon, p}} \left( e^{C_1^{\varepsilon, p} T} - 1 \right) \int_{\mathbb{R}} \rho_{\varepsilon}^p(0, x) dx. \quad (2.9)$$

1

□

### 2.3 $\mathbf{W}^{1,p}$ estimates for $p = 2N$ in the viscous case

3 We turn now to Sobolev estimates. Let  $N \in \mathbb{N}^*$  and set  $p = 2N$ .

**Proposition 3.** *We assume  $(\mathbf{A}_{\omega}^2)$ - $(\mathbf{A}_{\mathbf{v}}^2)$ . Let  $\rho_{\varepsilon}$  be the solution of (1.4) with initial datum  $\rho^0 \in \mathbf{W}^{1,2N}(\mathbb{R})$ . If  $\rho_{\varepsilon} \in \mathbf{L}^{\infty}([0, T] \times \mathbb{R})$  for some  $T > 0$ , then*

$$\rho_{\varepsilon} \in \mathbf{L}^{\infty}([0, T], \mathbf{W}^{1,2N}(\mathbb{R})) \quad \text{and} \quad \rho_{\varepsilon}, \partial_x \rho_{\varepsilon} \in \mathbf{L}^{\infty}([0, T], \mathbf{L}^{2N}(\mathbb{R})) \cap \mathbf{L}^{2N}([0, T] \times \mathbb{R}).$$

4 *Proof.* We differentiate (2.4) with respect to  $x$ , it gives

$$\begin{aligned} & \partial_t \partial_x \rho_{\varepsilon} + 2v'(\rho_{\varepsilon} * \omega_{\varepsilon}) \partial_x(\rho_{\varepsilon} * \omega_{\varepsilon}) \partial_x \rho_{\varepsilon} + v(\rho_{\varepsilon} * \omega_{\varepsilon}) \partial_{xx}^2 \rho_{\varepsilon} \\ & + \rho_{\varepsilon} v''(\rho_{\varepsilon} * \omega_{\varepsilon}) (\partial_x(\rho_{\varepsilon} * \omega_{\varepsilon}))^2 + \rho_{\varepsilon} v'(\rho_{\varepsilon} * \omega_{\varepsilon}) \partial_{xx}^2(\rho_{\varepsilon} * \omega_{\varepsilon}) = \varepsilon \partial_{xxx}^3 \rho_{\varepsilon}. \end{aligned} \quad (2.10)$$

5 Multiplying this relation by  $(\partial_x \rho_{\varepsilon})^{p-1}$ , then integrating with respect to  $x$ , we have

$$\begin{aligned} & \frac{1}{p} \frac{d}{dt} \int_{\mathbb{R}} (\partial_x \rho_{\varepsilon})^p dx + 2 \int_{\mathbb{R}} v'(\rho_{\varepsilon} * \omega_{\varepsilon}) \partial_x(\rho_{\varepsilon} * \omega_{\varepsilon}) (\partial_x \rho_{\varepsilon})^p dx + \int_{\mathbb{R}} v(\rho_{\varepsilon} * \omega_{\varepsilon}) \partial_{xx}^2 \rho_{\varepsilon} (\partial_x \rho_{\varepsilon})^{p-1} dx \\ & + \int_{\mathbb{R}} \rho_{\varepsilon} v''(\rho_{\varepsilon} * \omega_{\varepsilon}) (\partial_x(\rho_{\varepsilon} * \omega_{\varepsilon}))^2 (\partial_x \rho_{\varepsilon})^{p-1} dx + \int_{\mathbb{R}} \rho_{\varepsilon} v'(\rho_{\varepsilon} * \omega_{\varepsilon}) \partial_{xx}^2(\rho_{\varepsilon} * \omega_{\varepsilon}) (\partial_x \rho_{\varepsilon})^{p-1} dx \\ & = \varepsilon \int_{\mathbb{R}} (\partial_x \rho_{\varepsilon})^{p-1} \partial_{xxx}^3 \rho_{\varepsilon} dx. \end{aligned}$$

6 Notice that

$$\int_{\mathbb{R}} v(\rho_{\varepsilon} * \omega_{\varepsilon}) \partial_{xx}^2 \rho_{\varepsilon} (\partial_x \rho_{\varepsilon})^{p-1} dx = \frac{1}{p} \int_{\mathbb{R}} v(\rho_{\varepsilon} * \omega_{\varepsilon}) \partial_x ((\partial_x \rho_{\varepsilon})^p) dx$$



$$= -\frac{1}{p} \int_{\mathbb{R}} v'(\rho_\varepsilon * \omega_\varepsilon) \partial_x(\rho_\varepsilon * \omega_\varepsilon) (\partial_x \rho_\varepsilon)^p dx$$

1 and, since  $p$  is even,

$$\int_{\mathbb{R}} (\partial_x \rho_\varepsilon)^{p-1} \partial_{xxx}^3 \rho_\varepsilon dx = -(p-1) \int_{\mathbb{R}} (\partial_{xx}^2 \rho_\varepsilon)^2 (\partial_x \rho_\varepsilon)^{p-2} dx \leq 0,$$

2 thus

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}} (\partial_x \rho_\varepsilon)^p dx &\leq (1-2p) \int_{\mathbb{R}} v'(\rho_\varepsilon * \omega_\varepsilon) \partial_x(\rho_\varepsilon * \omega_\varepsilon) (\partial_x \rho_\varepsilon)^p dx \\ &\quad - p \int_{\mathbb{R}} \rho_\varepsilon v''(\rho_\varepsilon * \omega_\varepsilon) (\partial_x(\rho_\varepsilon * \omega_\varepsilon))^2 (\partial_x \rho_\varepsilon)^{p-1} dx \\ &\quad - p \int_{\mathbb{R}} \rho_\varepsilon v'(\rho_\varepsilon * \omega_\varepsilon) \partial_{xx}^2(\rho_\varepsilon * \omega_\varepsilon) (\partial_x \rho_\varepsilon)^{p-1} dx \\ &=: I_1^\varepsilon + I_2^\varepsilon + I_3^\varepsilon. \end{aligned}$$

3 We estimate now each of these terms.

- By (2.1) we get

$$|I_1^\varepsilon| = \left| (1-2p) \int_{\mathbb{R}} v'(\rho_\varepsilon * \omega_\varepsilon) \partial_x(\rho_\varepsilon * \omega_\varepsilon) (\partial_x \rho_\varepsilon)^p dx \right| \leq 2(2p-1) \|v'\|_\infty \|\rho_\varepsilon\|_\infty W_\varepsilon \int_{\mathbb{R}} |\partial_x \rho_\varepsilon|^p dx.$$

- 4 • Again by (2.1) we get

$$\begin{aligned} |I_2^\varepsilon| &= \left| p \int_{\mathbb{R}} \rho_\varepsilon v''(\rho_\varepsilon * \omega_\varepsilon) (\partial_x(\rho_\varepsilon * \omega_\varepsilon))^2 (\partial_x \rho_\varepsilon)^{p-1} dx \right| \\ &\leq p \|v''\|_\infty (2\|\rho_\varepsilon\|_\infty W_\varepsilon)^2 \int_{\mathbb{R}} \rho_\varepsilon |\partial_x \rho_\varepsilon|^{p-1} dx \\ &\leq 4 \|v''\|_\infty \|\rho_\varepsilon\|_\infty^2 W_\varepsilon^2 \left( \int_{\mathbb{R}} \rho_\varepsilon^p dx + (p-1) \int_{\mathbb{R}} |\partial_x \rho_\varepsilon|^p dx \right), \end{aligned}$$

5 where we have used Young's inequality  $uv \leq \frac{1}{p}u^p + \frac{1}{q}v^q$  with  $q = p/(p-1)$ .

- 7 • Similarly,

$$\begin{aligned} |I_3^\varepsilon| &= \left| p \int_{\mathbb{R}} \rho_\varepsilon v'(\rho_\varepsilon * \omega_\varepsilon) \partial_{xx}^2(\rho_\varepsilon * \omega_\varepsilon) (\partial_x \rho_\varepsilon)^{p-1} dx \right| \\ &\leq p \|v'\|_\infty \|\rho_\varepsilon\|_\infty \int_{\mathbb{R}} |\partial_{xx}^2(\rho_\varepsilon * \omega_\varepsilon)| |\partial_x \rho_\varepsilon|^{p-1} dx \\ &\leq \|v'\|_\infty \|\rho_\varepsilon\|_\infty \left( \int_{\mathbb{R}} |\partial_{xx}^2(\rho_\varepsilon * \omega_\varepsilon)|^p dx + (p-1) \int_{\mathbb{R}} |\partial_x \rho_\varepsilon|^p dx \right). \end{aligned}$$

We now observe that

$$(u+v+w)^p \leq 3^p(u^p+v^p+w^p) \quad \text{for any } u, v, w > 0 \text{ and } p > 0. \quad (2.11)$$

8 Indeed, from the binomial expansion we get

$$(u+v+w)^p = (u+(v+w))^p$$

$$\begin{aligned}
&= \sum_{k=0}^p \binom{p}{k} u^k (v+w)^{p-k} \\
&= \sum_{k=0}^p \binom{p}{k} u^k \sum_{l=0}^{p-k} \binom{p-k}{l} v^l w^{p-k-l}.
\end{aligned}$$

1 Observing that  $u^k v^l w^{p-k-l} \leq u^p + v^p + w^p$  and that  $\sum_{l=0}^{p-k} \binom{p-k}{l} = (1+1)^{p-k}$  and

2  $\sum_{k=0}^p \binom{p}{k} 1^k 2^{p-k} = 3^p$ , we get the result.

3 Estimate (2.2) of Proposition 1 and inequality (2.11) give

$$\begin{aligned}
\int_{\mathbb{R}} |\partial_{xx}^2(\rho_\varepsilon * \omega)|^p dx &\leq 3^p \eta \left( \int_0^\eta |\omega''(u)|^{p/(p-1)} du \right)^{p-1} \int_{\mathbb{R}} \rho_\varepsilon^p(t, x) dx \\
&\quad + 3^p (|\omega'(\eta-)| + |\omega'(0+)|)^p \int_{\mathbb{R}} \rho_\varepsilon^p(t, x) dx \\
&\quad + 6^p (\omega(\eta) + W_\varepsilon)^p \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(t, x)|^p dx, \tag{2.12}
\end{aligned}$$

thus

$$\left| p \int_{\mathbb{R}} \rho_\varepsilon v'(\rho * \omega_\varepsilon) \partial_{xx}^2(\rho_\varepsilon * \omega_\varepsilon) (\partial_x \rho_\varepsilon)^{p-1} dx \right| \leq C_2^{\varepsilon, p} \int_{\mathbb{R}} \rho_\varepsilon^p dx + C_3^{\varepsilon, p} \int_{\mathbb{R}} |\partial_x \rho_\varepsilon|^p dx,$$

with

$$C_2^{\varepsilon, p} = \|v'\|_\infty \|\rho_\varepsilon\|_\infty 3^p \left[ \eta \left( \int_0^\eta |\omega''(u)|^{p/(p-1)} du \right)^{p-1} + (|\omega'(\eta-)| + |\omega'(0+)|)^p \right]$$

and

$$C_3^{\varepsilon, p} = \|v'\|_\infty \|\rho_\varepsilon\|_\infty \left[ p-1 + 6^p (\omega(\eta) + W_\varepsilon)^p \right].$$

4 These bounds give finally the estimate

$$\frac{d}{dt} \int_{\mathbb{R}} (\partial_x \rho_\varepsilon(t, x))^p dx \leq C_4^{\varepsilon, p} \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(t, x)|^p dx + C_5^{\varepsilon, p} \int_{\mathbb{R}} \rho_\varepsilon^p(t, x) dx.$$

with

$$C_4^{\varepsilon, p} = 2(2p-1) \|v'\|_\infty \|\rho_\varepsilon\|_\infty W_\varepsilon + 4(p-1) \|v''\|_\infty \|\rho_\varepsilon\|_\infty^2 W_\varepsilon^2 + C_3^\varepsilon$$

and

$$C_5^{\varepsilon, p} = 4 \|v''\|_\infty \|\rho_\varepsilon\|_\infty^2 W_\varepsilon^2 + C_2^\varepsilon.$$

5 With (2.6), we get

$$\begin{aligned}
&\frac{d}{dt} \left( \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(t, x)|^{2N} dx + \int_{\mathbb{R}} \rho_\varepsilon^{2N}(t, x) dx \right) \\
&\leq C_6^{\varepsilon, p} \left( \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(t, x)|^{2N} dx + \int_{\mathbb{R}} \rho_\varepsilon^{2N}(t, x) dx \right), \tag{2.13}
\end{aligned}$$

6 with  $C_6^{\varepsilon, p} = \max(C_4^{\varepsilon, p}, C_5^{\varepsilon, p} + C_1^{\varepsilon, p})$ , which implies

$$\int_{\mathbb{R}} |\partial_x \rho_\varepsilon(t, x)|^{2N} dx + \int_{\mathbb{R}} \rho_\varepsilon^{2N}(t, x) dx$$

$$\leq e^{C_6^{\varepsilon,p}t} \left( \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(0, x)|^{2N} dx + \int_{\mathbb{R}} \rho_\varepsilon^{2N}(0, x) dx \right). \quad (2.14)$$

1 Then

$$\begin{aligned} & \sup_{t \in [0, T]} \left( \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(t, x)|^{2N} dx + \int_{\mathbb{R}} \rho_\varepsilon^{2N}(t, x) dx \right) \\ & \leq e^{C_6^{\varepsilon,p}T} \left( \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(0, x)|^{2N} dx + \int_{\mathbb{R}} \rho_\varepsilon^{2N}(0, x) dx \right). \end{aligned} \quad (2.15)$$

2 Integrating (2.14) with respect to  $t$  on  $[0, T]$ , we get

$$\begin{aligned} & \int_0^T \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(t, x)|^{2N} dx dt + \int_0^T \int_{\mathbb{R}} \rho_\varepsilon^{2N}(t, x) dx dt \\ & \leq \frac{1}{C_6^{\varepsilon,p}} \left( e^{C_6^{\varepsilon,p}T} - 1 \right) \left( \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(0, x)|^{2N} dx + \int_{\mathbb{R}} \rho_\varepsilon^{2N}(0, x) dx \right). \end{aligned} \quad (2.16)$$

3

□

#### 4 2.4 $\mathbf{L}^\infty$ bound on an interval $[0, T]$

5 With the previous estimates, we are now able to prove an  $\mathbf{L}^\infty$  bounds for the sequence  $\{\rho_\varepsilon\}_\varepsilon$  on an  
6 interval  $[0, T]$ .

**Proposition 4.** *We assume  $(\mathbf{A}_w^2)$ - $(\mathbf{A}_v^2)$ . Let  $\rho_\varepsilon$  be the solution of (1.4) with initial datum  $\rho^0 \in \mathbf{H}^1$ . Then there exists a constant  $\bar{T} > 0$  such that  $\rho_\varepsilon \in \mathbf{L}^\infty([0, T] \times \mathbb{R})$  for any  $\varepsilon > 0$ ,  $T < \bar{T}$ . Furthermore*

$$\rho_\varepsilon \in \mathbf{L}^\infty([0, T], \mathbf{W}^{1,2N}(\mathbb{R})) \quad \text{and} \quad \rho_\varepsilon, \partial_x \rho_\varepsilon \in \mathbf{L}^\infty([0, T], \mathbf{L}^{2N}(\mathbb{R})) \cap \mathbf{L}^{2N}([0, T] \times \mathbb{R}) \quad (2.17)$$

7 and this sequence is uniformly bounded in these spaces with respect to  $\varepsilon$ .

8 *Proof.* Let  $\rho_\varepsilon$  be a smooth solution of (1.4) with the same initial datum  $\rho^0 \in \mathbf{H}^1$ . The relation  
9 (2.13) for  $N = 1$  gives

$$\begin{aligned} & \frac{d}{dt} \left( \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(t, x)|^2 dx + \int_{\mathbb{R}} \rho_\varepsilon^2(t, x) dx \right) \\ & \leq C \max \left\{ 1, \|\rho_\varepsilon(t, \cdot)\|_\infty^2 \right\} \left( \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(t, x)|^2 dx + \int_{\mathbb{R}} \rho_\varepsilon^2(t, x) dx \right), \end{aligned}$$

for some constant  $C$  that does not depend on  $\varepsilon$  (since  $W_\varepsilon$  is uniformly bounded). If no uniform  $\mathbf{L}^\infty$ -bound on  $\rho_\varepsilon$  is available, we can use the Sobolev injection of  $\mathbf{H}^1(\mathbb{R})$  in  $\mathbf{L}^\infty(\mathbb{R})$  and get

$$\frac{d}{dt} \|\rho_\varepsilon(t, \cdot)\|_{\mathbf{H}^1}^2 \leq C \|\rho_\varepsilon(t, \cdot)\|_{\mathbf{H}^1}^2 + C \|\rho_\varepsilon(t, \cdot)\|_{\mathbf{H}^1}^4,$$

eventually updating the constant  $C$ . We set  $u_\varepsilon(t) = \|\rho_\varepsilon(t, \cdot)\|_{\mathbf{H}^1}^2$ , then  $u'_\varepsilon \leq C(u_\varepsilon + u_\varepsilon^2)$ , which leads to

$$\frac{u'_\varepsilon}{u_\varepsilon} - \frac{u'_\varepsilon}{1 + u_\varepsilon} \leq C.$$

We obtain

$$u_\varepsilon(t) \leq \frac{C_0 e^{Ct}}{1 - C_0 e^{Ct}}, \quad \text{for any } 0 \leq t < -\frac{\ln C_0}{C},$$

with  $C_0 = \frac{u_0}{1+u_0} < 1$ ,  $u_0 = \|\rho^0\|_{\mathbf{H}^1}^2$ . Notice that the initial datum is the same for all the sequence and then  $u_0$  and  $C_0$  do not depend on  $\varepsilon$ . Setting  $T < \bar{T} := -\frac{\ln C_0}{C}$ , we have

$$\|\rho_\varepsilon(t, \cdot)\|_{\mathbf{H}^1}^2 \leq C \|\rho^0\|_{\mathbf{H}^1}^2, \quad \text{for any } 0 \leq t \leq T, \quad \varepsilon > 0.$$

1 Therefore, by Sobolev injection,  $\rho_\varepsilon \in \mathbf{L}^\infty([0, T] \times \mathbb{R})$ . Using the estimates of Propositions 2 and 3,  
 2 we get (2.17) with bounds independents of  $\varepsilon$ .  $\square$

### 3 **2.5 $\mathbf{W}^{2,p}$ estimate for $p = 2N$**

4 To pass to the limit, we need also estimates in  $\mathbf{W}^{2,p}$ , which will provide, in the next section, with  
 5 the help of the equation, the necessary regularity in time. As in Section 2.3, let  $N \in \mathbb{N}^*$  and set  
 6  $p = 2N$ .

**Proposition 5.** *We assume  $(\mathbf{A}_\omega^2)$ - $(\mathbf{A}_v^3)$ . Let  $\rho_\varepsilon$  be the solution of (1.4) with initial datum  $\rho^0 \in \mathbf{W}^{1,4N}(\mathbb{R}) \cap \mathbf{H}^1(\mathbb{R}) \cap \mathbf{W}^{2,2N}(\mathbb{R})$ . Let  $T > 0$  as in Proposition 4. Then*

$$\rho_\varepsilon \in \mathbf{L}^\infty\left([0, T], \mathbf{W}^{2,2N}(\mathbb{R})\right) \quad \text{and} \quad \rho_\varepsilon, \partial_x \rho_\varepsilon, \partial_{xx}^2 \rho_\varepsilon \in \mathbf{L}^\infty\left([0, T], \mathbf{L}^{2N}(\mathbb{R})\right) \cap \mathbf{L}^{2N}([0, T] \times \mathbb{R})$$

7 and this sequence is bounded in these spaces with respect to  $\varepsilon$ .

8 *Proof.* We differentiate (2.10) with respect to  $x$ , which gives

$$\begin{aligned} & \partial_t \partial_{xx}^2 \rho_\varepsilon + 3v''(\rho_\varepsilon * \omega_\varepsilon) (\partial_x(\rho_\varepsilon * \omega_\varepsilon))^2 \partial_x \rho_\varepsilon + 3v'(\rho_\varepsilon * \omega_\varepsilon) \partial_{xx}^2(\rho_\varepsilon * \omega_\varepsilon) \partial_x \rho_\varepsilon \\ & + 3v'(\rho_\varepsilon * \omega_\varepsilon) \partial_x(\rho_\varepsilon * \omega_\varepsilon) \partial_{xx}^2 \rho_\varepsilon + v(\rho_\varepsilon * \omega_\varepsilon) \partial_{xxx}^3 \rho_\varepsilon + \rho_\varepsilon v^{(3)}(\rho_\varepsilon * \omega_\varepsilon) (\partial_x(\rho_\varepsilon * \omega_\varepsilon))^3 \\ & + 3\rho_\varepsilon v''(\rho_\varepsilon * \omega_\varepsilon) \partial_x(\rho_\varepsilon * \omega_\varepsilon) \partial_{xx}^2(\rho_\varepsilon * \omega_\varepsilon) + \rho_\varepsilon v'(\rho_\varepsilon * \omega_\varepsilon) \partial_{xxx}^3(\rho_\varepsilon * \omega_\varepsilon) = \varepsilon \partial_{xxxx}^4 \rho_\varepsilon. \end{aligned} \quad (2.18)$$

9 Multiplying this relation by  $(\partial_{xx}^2 \rho_\varepsilon)^{p-1}$ , then integrating with respect to  $x$ , we obtain

$$\begin{aligned} & \frac{1}{p} \frac{d}{dt} \int_{\mathbb{R}} (\partial_{xx}^2 \rho_\varepsilon)^p dx + 3 \int_{\mathbb{R}} v''(\rho_\varepsilon * \omega_\varepsilon) (\partial_x(\rho_\varepsilon * \omega_\varepsilon))^2 \partial_x \rho_\varepsilon (\partial_{xx}^2 \rho_\varepsilon)^{p-1} dx \\ & + 3 \int_{\mathbb{R}} v'(\rho_\varepsilon * \omega_\varepsilon) \partial_{xx}^2(\rho_\varepsilon * \omega_\varepsilon) \partial_x \rho_\varepsilon (\partial_{xx}^2 \rho_\varepsilon)^{p-1} dx + 3 \int_{\mathbb{R}} v'(\rho_\varepsilon * \omega_\varepsilon) \partial_x(\rho_\varepsilon * \omega_\varepsilon) (\partial_{xx}^2 \rho_\varepsilon)^p dx \\ & + \int_{\mathbb{R}} v(\rho_\varepsilon * \omega_\varepsilon) \partial_{xxx}^3 \rho_\varepsilon (\partial_{xx}^2 \rho_\varepsilon)^{p-1} dx + \int_{\mathbb{R}} \rho_\varepsilon v^{(3)}(\rho_\varepsilon * \omega_\varepsilon) (\partial_x(\rho_\varepsilon * \omega_\varepsilon))^3 (\partial_{xx}^2 \rho_\varepsilon)^{p-1} dx \\ & + 3 \int_{\mathbb{R}} \rho_\varepsilon v''(\rho_\varepsilon * \omega_\varepsilon) \partial_x(\rho_\varepsilon * \omega_\varepsilon) \partial_{xx}^2(\rho_\varepsilon * \omega_\varepsilon) (\partial_{xx}^2 \rho_\varepsilon)^{p-1} dx \\ & + \int_{\mathbb{R}} \rho_\varepsilon v'(\rho_\varepsilon * \omega_\varepsilon) \partial_{xxx}^3(\rho_\varepsilon * \omega_\varepsilon) (\partial_{xx}^2 \rho_\varepsilon)^{p-1} dx \\ & = \varepsilon \int_{\mathbb{R}} \partial_{xxxx}^4 \rho_\varepsilon (\partial_{xx}^2 \rho_\varepsilon)^{p-1} dx. \end{aligned}$$

10 Now

$$\begin{aligned} \int_{\mathbb{R}} v(\rho_\varepsilon * \omega_\varepsilon) \partial_{xxx}^3 \rho_\varepsilon (\partial_{xx}^2 \rho_\varepsilon)^{p-1} dx &= \frac{1}{p} \int_{\mathbb{R}} v(\rho_\varepsilon * \omega_\varepsilon) \partial_x \left( (\partial_{xx}^2 \rho_\varepsilon)^p \right) dx \\ &= -\frac{1}{p} \int_{\mathbb{R}} v'(\rho_\varepsilon * \omega_\varepsilon) \partial_x(\rho_\varepsilon * \omega_\varepsilon) (\partial_{xx}^2 \rho_\varepsilon)^p dx, \end{aligned}$$

1 therefore

$$\begin{aligned}
\frac{d}{dt} \int_{\mathbb{R}} (\partial_{xx}^2 \rho_\varepsilon)^p dx &= -3p \int_{\mathbb{R}} v''(\rho_\varepsilon * \omega_\varepsilon) (\partial_x(\rho_\varepsilon * \omega_\varepsilon))^2 \partial_x \rho_\varepsilon (\partial_{xx}^2 \rho_\varepsilon)^{p-1} dx \\
&\quad -3p \int_{\mathbb{R}} v'(\rho_\varepsilon * \omega_\varepsilon) \partial_{xx}^2(\rho_\varepsilon * \omega_\varepsilon) \partial_x \rho_\varepsilon (\partial_{xx}^2 \rho_\varepsilon)^{p-1} dx \\
&\quad + (1-3p) \int_{\mathbb{R}} v'(\rho_\varepsilon * \omega_\varepsilon) \partial_x(\rho_\varepsilon * \omega_\varepsilon) (\partial_{xx}^2 \rho_\varepsilon)^p dx \\
&\quad -p \int_{\mathbb{R}} \rho_\varepsilon v^{(3)}(\rho_\varepsilon * \omega_\varepsilon) (\partial_x(\rho_\varepsilon * \omega_\varepsilon))^3 (\partial_{xx}^2 \rho_\varepsilon)^{p-1} dx \\
&\quad -3p \int_{\mathbb{R}} \rho_\varepsilon v''(\rho_\varepsilon * \omega_\varepsilon) \partial_x(\rho_\varepsilon * \omega_\varepsilon) \partial_{xx}^2(\rho_\varepsilon * \omega_\varepsilon) (\partial_{xx}^2 \rho_\varepsilon)^{p-1} dx \\
&\quad -p \int_{\mathbb{R}} \rho_\varepsilon v'(\rho_\varepsilon * \omega_\varepsilon) \partial_{xxx}^3(\rho_\varepsilon * \omega_\varepsilon) (\partial_{xx}^2 \rho_\varepsilon)^{p-1} dx \\
&\quad + \varepsilon p \int_{\mathbb{R}} \partial_{xxxx}^4 \rho_\varepsilon (\partial_{xx}^2 \rho_\varepsilon)^{p-1} dx \\
&=: J_1 + J_2 + J_3 + J_4 + J_5 + J_6 + J_7.
\end{aligned}$$

2 We estimate now each of these terms.

3 • Using (2.1) and Young's inequality, we get

$$\begin{aligned}
|J_1| &= 3p \left| \int_{\mathbb{R}} v''(\rho_\varepsilon * \omega_\varepsilon) (\partial_x(\rho_\varepsilon * \omega_\varepsilon))^2 \partial_x \rho_\varepsilon (\partial_{xx}^2 \rho_\varepsilon)^{p-1} dx \right| \\
&\leq 3p \|v''\|_\infty (2\|\rho_\varepsilon\|_\infty W_\varepsilon)^2 \int_{\mathbb{R}} |\partial_x \rho_\varepsilon| |\partial_{xx}^2 \rho_\varepsilon|^{p-1} dx \\
&\leq 12 \|v''\|_\infty \|\rho_\varepsilon\|_\infty^2 W_\varepsilon^2 \left( \int_{\mathbb{R}} |\partial_x \rho_\varepsilon|^p dx + (p-1) \int_{\mathbb{R}} |\partial_{xx}^2 \rho_\varepsilon|^p dx \right)
\end{aligned}$$

•

$$\begin{aligned}
|J_2| &= 3p \left| \int_{\mathbb{R}} v'(\rho_\varepsilon * \omega_\varepsilon) \partial_{xx}^2(\rho_\varepsilon * \omega_\varepsilon) \partial_x \rho_\varepsilon (\partial_{xx}^2 \rho_\varepsilon)^{p-1} dx \right| \\
&\leq 3p \|v'\|_\infty \int_{\mathbb{R}} |\partial_{xx}^2(\rho_\varepsilon * \omega_\varepsilon)| |\partial_x \rho_\varepsilon| |\partial_{xx}^2 \rho_\varepsilon|^{p-1} dx \\
&\leq 3 \|v'\|_\infty \left( \frac{1}{2} \int_{\mathbb{R}} |\partial_{xx}^2(\rho_\varepsilon * \omega_\varepsilon)|^{2p} dx + \frac{1}{2} \int_{\mathbb{R}} |\partial_x \rho_\varepsilon|^{2p} dx + (p-1) \int_{\mathbb{R}} |\partial_{xx}^2 \rho_\varepsilon|^p dx \right)
\end{aligned}$$

using the inequality

$$uvw \leq \frac{1}{p_1} u^{p_1} + \frac{1}{p_2} v^{p_2} + \frac{1}{p_3} w^{p_3}, \quad \text{with } \frac{1}{p_1} + \frac{1}{p_2} + \frac{1}{p_3} = 1, \quad (2.19)$$

4 setting  $p_1 = 2p = p_2$  and  $p_3 = p/(p-1)$ . Estimate (2.19) can be derived applying twice the  
5 classical Young's inequality to  $uvw = u(vw)$ . Using now the relation (2.12) with  $2p$  at the  
6 place of  $p$ , we get

$$|J_2| \leq \frac{3^{2p+1}}{2} \|v'\|_\infty \eta \left( \int_0^\eta |\omega''(u)|^{2p/(2p-1)} du \right)^{2p-1} \int_{\mathbb{R}} \rho_\varepsilon^{2p} dx$$

$$\begin{aligned}
& + \frac{3^{2p+1}}{2} \|v'\|_\infty (|\omega'(\eta-)| + |\omega'(0+)|)^{2p} \int_{\mathbb{R}} \rho_\varepsilon^{2p} dx \\
& + 3^{2p+1} 2^{2p-1} \|v'\|_\infty (\omega(\eta) + 2\omega(0))^{2p} \int_{\mathbb{R}} |\partial_x \rho_\varepsilon|^{2p} dx \\
& + \frac{3}{2} \|v'\|_\infty \int_{\mathbb{R}} |\partial_x \rho_\varepsilon|^{2p} dx + 3(p-1) \|v'\|_\infty \int_{\mathbb{R}} |\partial_{xx}^2 \rho_\varepsilon|^p dx.
\end{aligned}$$

•

$$\begin{aligned}
|J_3| & = (3p-1) \left| \int_{\mathbb{R}} v'(\rho_\varepsilon * \omega_\varepsilon) \partial_x(\rho_\varepsilon * \omega_\varepsilon) (\partial_{xx}^2 \rho_\varepsilon)^p dx \right| \\
& \leq 2(3p-1) \|v'\|_\infty \|\rho_\varepsilon\|_\infty W_\varepsilon \int_{\mathbb{R}} |\partial_{xx}^2 \rho_\varepsilon|^p dx.
\end{aligned}$$

•

$$\begin{aligned}
|J_4| & = p \left| \int_{\mathbb{R}} \rho_\varepsilon v^{(3)}(\rho_\varepsilon * \omega_\varepsilon) (\partial_x(\rho_\varepsilon * \omega_\varepsilon))^3 (\partial_{xx}^2 \rho_\varepsilon)^{p-1} dx \right| \\
& \leq p \|v^{(3)}\|_\infty (2\|\rho_\varepsilon\|_\infty W_\varepsilon)^3 \int_{\mathbb{R}} \rho_\varepsilon |\partial_{xx}^2 \rho_\varepsilon|^{p-1} dx \\
& \leq 8 \|v^{(3)}\|_\infty \|\rho_\varepsilon\|_\infty^3 W_\varepsilon^3 \left( \int_{\mathbb{R}} \rho_\varepsilon^p dx + (p-1) \int_{\mathbb{R}} |\partial_{xx}^2 \rho_\varepsilon|^p dx \right)
\end{aligned}$$

1 using Young's inequality.

•

$$\begin{aligned}
|J_5| & = 3p \left| \int_{\mathbb{R}} \rho_\varepsilon v''(\rho_\varepsilon * \omega_\varepsilon) \partial_x(\rho_\varepsilon * \omega_\varepsilon) \partial_{xx}^2(\rho_\varepsilon * \omega_\varepsilon) (\partial_{xx}^2 \rho_\varepsilon)^{p-1} dx \right| \\
& \leq 6p \|v''\|_\infty \|\rho_\varepsilon\|_\infty^2 W_\varepsilon \int_{\mathbb{R}} |\partial_{xx}^2(\rho_\varepsilon * \omega_\varepsilon)| |\partial_{xx}^2 \rho_\varepsilon|^{p-1} dx \\
& \leq 6 \|v''\|_\infty \|\rho_\varepsilon\|_\infty^2 W_\varepsilon \left( \int_{\mathbb{R}} |\partial_{xx}^2(\rho_\varepsilon * \omega_\varepsilon)|^p dx + (p-1) \int_{\mathbb{R}} |\partial_{xx}^2 \rho_\varepsilon|^p dx \right) \\
& \leq 6 \|v''\|_\infty \|\rho_\varepsilon\|_\infty^2 W_\varepsilon \left( 3^p \eta \left( \int_0^\eta |\omega''(u)|^{p/(p-1)} du \right)^{p-1} \int_{\mathbb{R}} \rho_\varepsilon^p dx \right. \\
& \quad \left. + 3^p (|\omega'(\eta-)| + |\omega'(0+)|)^p \int_{\mathbb{R}} \rho_\varepsilon^p dx \right. \\
& \quad \left. + 6^p (\omega(\eta) + 2\omega(0))^p \int_{\mathbb{R}} |\partial_x \rho_\varepsilon|^p dx + (p-1) \int_{\mathbb{R}} |\partial_{xx}^2 \rho_\varepsilon|^p dx \right)
\end{aligned}$$

2 using relation (2.12).

•

$$\begin{aligned}
|J_6| & = p \left| \int_{\mathbb{R}} \rho_\varepsilon v'(\rho_\varepsilon * \omega_\varepsilon) \partial_{xxx}^3(\rho_\varepsilon * \omega_\varepsilon) (\partial_{xx}^2 \rho_\varepsilon)^{p-1} dx \right| \\
& \leq p \|\rho_\varepsilon\|_\infty \|v'\|_\infty \int_{\mathbb{R}} |\partial_{xxx}^3(\rho_\varepsilon * \omega_\varepsilon)| |\partial_{xx}^2 \rho_\varepsilon|^{p-1} dx
\end{aligned}$$

$$\leq \|\rho_\varepsilon\|_\infty \|v'\|_\infty \left( \int_{\mathbb{R}} |\partial_{xxx}^3(\rho_\varepsilon * \omega_\varepsilon)|^p dx + (p-1) \int_{\mathbb{R}} |\partial_{xx}^2 \rho_\varepsilon|^p dx \right).$$

1 Estimate (2.3) of Proposition 1 and the inequality (2.11) give

$$\begin{aligned} \int_{\mathbb{R}} |\partial_{xxx}^3(\rho_\varepsilon * \omega_\varepsilon)|^p(t, x) dx &\leq 3^p \eta \left( \int_0^\eta |\omega''(u)|^{p/(p-1)} du \right)^{p-1} \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(t, x)|^p dx \\ &\quad + 3^p (|\omega'(\eta-)| + |\omega'(0+)|)^p \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(t, x)|^p dx \\ &\quad + 6^p (\omega(\eta) + 2\omega(0))^p \int_{\mathbb{R}} |\partial_{xx}^2 \rho_\varepsilon(t, x)|^p dx. \end{aligned}$$

2 Then

$$\begin{aligned} |J_6| &\leq \|\rho_\varepsilon\|_\infty \|v'\|_\infty \left( 3^p \eta \left( \int_0^\eta |\omega''(u)|^{p/(p-1)} du \right)^{p-1} \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(t, x)|^p dx \right. \\ &\quad + 3^p (|\omega'(\eta-)| + |\omega'(0+)|)^p \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(t, x)|^p dx \\ &\quad + 6^p (\omega(\eta) + 2\omega(0))^p \int_{\mathbb{R}} |\partial_{xx}^2 \rho_\varepsilon(t, x)|^p dx \\ &\quad \left. + (p-1) \int_{\mathbb{R}} |\partial_{xx}^2 \rho_\varepsilon(t, x)|^p dx \right). \end{aligned}$$

•

$$J_7 = \varepsilon p \int_{\mathbb{R}} \partial_{xxxx}^4 \rho_\varepsilon (\partial_{xx}^2 \rho_\varepsilon)^{p-1} dx = -\varepsilon p(p-1) \int_{\mathbb{R}} (\partial_{xxx}^3 \rho_\varepsilon)^2 (\partial_{xx}^2 \rho_\varepsilon)^{2(N-1)} dx \leq 0.$$

3 The above estimates give an estimate of the form

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}} (\partial_{xx}^2 \rho_\varepsilon(t, x))^p dx &\leq C_7^{\varepsilon, p} \left( \int_{\mathbb{R}} |\partial_{xx}^2 \rho_\varepsilon(t, x)|^p dx + \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(t, x)|^p dx + \int_{\mathbb{R}} \rho_\varepsilon^p(t, x) dx \right. \\ &\quad \left. + \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(t, x)|^{2p} dx + \int_{\mathbb{R}} \rho_\varepsilon^{2p}(t, x) dx \right), \end{aligned}$$

4 where  $C_7^p = C_7^p(p, \|v'\|_\infty, \|v''\|_\infty, \|v^{(3)}\|_\infty, \sup_\varepsilon \{\|\rho_\varepsilon\|_\infty W_\varepsilon\}, C_\omega^p, C_\omega^{2p})$  and

$$C_\omega^p = \max \left\{ 3^p \eta \left( \int_0^\eta |\omega''(u)|^{p/(p-1)} du \right)^{p-1}, 3^p (|\omega'(\eta-)| + |\omega'(0+)|)^p, 6^p (\omega(\eta) + 2\omega(0))^p \right\}.$$

5 Note that  $C_7^p$  is a constant since  $\|\rho_\varepsilon\|_\infty W_\varepsilon$  is bounded with respect to  $\varepsilon$  thanks to Proposition 4  
6 and 1.5. This estimate, combined with (2.13) and (2.14), give then

$$\frac{d}{dt} \left( \int_{\mathbb{R}} |\partial_{xx}^2 \rho_\varepsilon(t, x)|^{2N} dx + \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(t, x)|^{2N} dx + \int_{\mathbb{R}} \rho_\varepsilon^{2N}(t, x) dx \right)$$

$$\begin{aligned} &\leq C_8^{2N} \left( \int_{\mathbb{R}} |\partial_{xx}^2 \rho_\varepsilon(t, x)|^{2N} dx + \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(t, x)|^{2N} dx + \int_{\mathbb{R}} \rho_\varepsilon^{2N}(t, x) dx \right) \\ &\quad + C_7^{2N} e^{C_9^{2N} t} \left( \int_{\mathbb{R}} |\partial_x \rho(0, x)|^{4N} dx + \int_{\mathbb{R}} |\rho(0, x)|^{4N} dx \right), \end{aligned}$$

where

$$C_8^{2N} = C_7^{2N} + \sup_{\varepsilon} C_6^{\varepsilon, 2N}, \quad C_9^{2N} = \sup_{\varepsilon} C_6^{\varepsilon, 2N}.$$

Note that  $C_6^{\varepsilon, 2N}$  is bounded with respect to  $\varepsilon$  thanks to Proposition 4 and 1.5. Since an inequality of the form

$$u'(t) \leq K_1 u(t) + K_2 e^{K_3 t}$$

implies the estimate

$$u(t) \leq u(0) e^{K_1 t} + K_2 e^{K_1 t} \int_0^t e^{(K_3 - K_1)s} ds \leq u(0) e^{K_1 t} + \frac{K_2}{K_3} \left( e^{(K_1 + K_3)t} - e^{K_1 t} \right),$$

1 we get the estimate

$$\begin{aligned} &\left( \int_{\mathbb{R}} |\partial_{xx}^2 \rho_\varepsilon(t, x)|^{2N} dx + \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(t, x)|^{2N} dx + \int_{\mathbb{R}} \rho_\varepsilon^{2N}(t, x) dx \right) \\ &\leq \left( \int_{\mathbb{R}} |\partial_{xx}^2 \rho(0, x)|^{2N} dx + \int_{\mathbb{R}} |\partial_x \rho(0, x)|^{2N} dx + \int_{\mathbb{R}} \rho^{2N}(0, x) dx \right) e^{C_8^{2N} t} \\ &\quad + \frac{C_7^{2N}}{C_9^{2N}} \left( \int_{\mathbb{R}} |\partial_x \rho(0, x)|^{4N} dx + \int_{\mathbb{R}} |\rho(0, x)|^{4N} dx \right) \left( e^{(C_8^{2N} + C_9^{2N})t} - e^{C_8^{2N} t} \right), \end{aligned}$$

2 which implies

$$\begin{aligned} &\sup_{t \in [0, T]} \left( \int_{\mathbb{R}} |\partial_{xx}^2 \rho_\varepsilon(t, x)|^{2N} dx + \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(t, x)|^{2N} dx + \int_{\mathbb{R}} \rho_\varepsilon^{2N}(t, x) dx \right) \\ &\leq \left( \int_{\mathbb{R}} |\partial_{xx}^2 \rho(0, x)|^{2N} dx + \int_{\mathbb{R}} |\partial_x \rho(0, x)|^{2N} dx + \int_{\mathbb{R}} \rho^{2N}(0, x) dx \right) e^{C_8^{2N} T} \\ &\quad + \frac{C_7^{2N}}{C_9^{2N}} \left( \int_{\mathbb{R}} |\partial_x \rho(0, x)|^{4N} dx + \int_{\mathbb{R}} |\rho(0, x)|^{4N} dx \right) e^{(C_8^{2N} + C_9^{2N})T} \end{aligned}$$

3 and

$$\begin{aligned} &\left( \int_0^T \int_{\mathbb{R}} |\partial_{xx}^2 \rho_\varepsilon(t, x)|^{2N} dx dt + \int_0^T \int_{\mathbb{R}} |\partial_x \rho_\varepsilon(t, x)|^{2N} dx dt + \int_0^T \int_{\mathbb{R}} \rho_\varepsilon^{2N}(t, x) dx dt \right) \\ &\leq \left( \int_{\mathbb{R}} |\partial_{xx}^2 \rho(0, x)|^{2N} dx + \int_{\mathbb{R}} |\partial_x \rho(0, x)|^{2N} dx + \int_{\mathbb{R}} \rho^{2N}(0, x) dx \right) T e^{C_8^{2N} T} \\ &\quad + \frac{C_7^{2N}}{C_9^{2N}} \left( \int_{\mathbb{R}} |\partial_x \rho(0, x)|^{4N} dx + \int_{\mathbb{R}} |\rho(0, x)|^{4N} dx \right) e^{(C_8^{2N} + C_9^{2N})T} T. \end{aligned}$$

4

□



### 1 **3 Proof of Theorem 1**

2 In this section, we pass to the limit as  $\varepsilon \rightarrow 0$  and we show that the limit function  $\rho$  satisfies equation  
3 (1.1).

Using Proposition 2, the sequence  $\{\rho_\varepsilon\}_\varepsilon$  is bounded in  $\mathbf{L}^\infty([0, T], \mathbf{L}^2(\mathbb{R}))$ . Using Proposition 3, the sequence  $\{\partial_x \rho_\varepsilon\}_\varepsilon$  is bounded in  $\mathbf{L}^\infty([0, T], \mathbf{L}^2(\mathbb{R}))$ . Using Propositions 1 and 4, the sequences  $\{v(\rho_\varepsilon * \omega_\varepsilon)\}_\varepsilon$ ,  $\{v'(\rho_\varepsilon * \omega_\varepsilon)\}_\varepsilon$  and  $\{\partial_x(\rho_\varepsilon * \omega_\varepsilon)\}_\varepsilon$  are bounded in  $\mathbf{L}^\infty([0, T] \times \mathbb{R})$ . Then

$$\partial_x(\rho_\varepsilon v(\rho_\varepsilon * \omega_\varepsilon)) = \partial_x \rho_\varepsilon \cdot v(\rho_\varepsilon * \omega_\varepsilon) + \rho_\varepsilon v'(\rho_\varepsilon * \omega_\varepsilon) \partial_x(\rho_\varepsilon * \omega_\varepsilon)$$

is bounded in  $\mathbf{L}^\infty([0, T], \mathbf{L}^2(\mathbb{R}))$ . Using Proposition 5, we also have a bound with respect to  $\varepsilon$  for  $\partial_{xx}^2 \rho_\varepsilon$  in the space  $\mathbf{L}^\infty([0, T], \mathbf{L}^2(\mathbb{R}))$ , then

$$\partial_t \rho_\varepsilon = \varepsilon \partial_{xx}^2 \rho_\varepsilon - \partial_x(\rho_\varepsilon v(\rho_\varepsilon * \omega_\varepsilon)) \in \mathbf{L}^\infty([0, T], \mathbf{L}^2(\mathbb{R}))$$

4 uniformly with respect to  $\varepsilon$ . In particular,  $\rho_\varepsilon \in \mathbf{C}([0, T], \mathbf{L}^2(\mathbb{R}))$  and the sequence is bounded in  
5 this space. Since  $\partial_t \rho_\varepsilon, \partial_x \rho_\varepsilon \in \mathbf{L}^\infty([0, T], \mathbf{L}^2(\mathbb{R}))$  with uniform bounds with respect to  $\varepsilon$ , then  $\{\rho_\varepsilon\}_\varepsilon$   
6 is bounded in  $\mathbf{H}_{\text{loc}}^1([0, T] \times \mathbb{R})$ . Up to the extraction of a subsequence, the sequence  $\{\rho_\varepsilon\}_\varepsilon$  converges  
7 to some  $\rho$  in  $\mathbf{L}_{\text{loc}}^2([0, T] \times \mathbb{R})$  and a.e. We have now to prove that the limit  $\rho \in \mathbf{C}([0, T], \mathbf{L}^2(\mathbb{R}))$  is  
8 a solution of (1.1). Since

$$\begin{aligned} (\rho_\varepsilon * \omega_\varepsilon)(t, x) - (\rho * \omega)(t, x) &= \int_{-\varepsilon}^0 \rho_\varepsilon(t, x + y) \omega_\varepsilon(y) dy + \int_0^\eta (\rho_\varepsilon - \rho)(t, x + y) \omega(y) dy \\ &\quad + \int_\eta^{\eta+\varepsilon} \rho_\varepsilon(t, x + y) \omega_\varepsilon(y) dy \end{aligned}$$

tends to 0 when  $\varepsilon$  goes to zero, we have

$$\rho_\varepsilon v(\rho_\varepsilon * \omega_\varepsilon) \rightarrow \rho v(\rho * \omega) \text{ a.e.}$$

9 Therefore using dominated convergence Theorem, we get  $\rho_\varepsilon v(\rho_\varepsilon * \omega_\varepsilon) \rightarrow \rho v(\rho * \omega)$  in  $\mathbf{L}_{\text{loc}}^1([0, T] \times \mathbb{R})$ ,  
10 implying that  $\rho$  is a solution of (1.1).

### 11 **References**

- 12 [1] A. Aggarwal, R. M. Colombo, and P. Goatin. Nonlocal systems of conservation laws in several space  
13 dimensions. *SIAM J. Numer. Anal.*, 53(2):963–983, 2015.
- 14 [2] P. Amorim, R. Colombo, and A. Teixeira. On the numerical integration of scalar nonlocal conservation  
15 laws. *ESAIM M2AN*, 49(1):19–37, 2015.
- 16 [3] F. Betancourt, R. Bürger, K. H. Karlsen, and E. M. Tory. On nonlocal conservation laws modelling  
17 sedimentation. *Nonlinearity*, 24(3):855–885, 2011.
- 18 [4] S. Blandin and P. Goatin. Well-posedness of a conservation law with non-local flux arising in traffic  
19 flow modeling. *Numer. Math.*, 132(2):217–241, 2016.
- 20 [5] M. Colombo, G. Crippa, and L. V. Spinolo. On the singular local limit for conservation laws with  
21 nonlocal fluxes. *ArXiv e-prints*, Oct. 2017.
- 22 [6] R. M. Colombo, M. Garavello, and M. Lécureux-Mercier. A class of nonlocal models for pedestrian  
23 traffic. *Mathematical Models and Methods in Applied Sciences*, 22(04):1150023, 2012.
- 24 [7] J. Friedrich, O. Kolb, and S. Göttlich. A Godunov type scheme for a class of LWR traffic flow models  
25 with non-local flux. *ArXiv e-prints*, Feb. 2018.

- 1 [8] P. Goatin and S. Scialanga. Well-posedness and finite volume approximations of the LWR traffic flow  
2 model with non-local velocity. *Netw. Heterog. Media*, 11(1):107–121, 2016.
- 3 [9] S. Göttlich, S. Hoher, P. Schindler, V. Schleper, and A. Verl. Modeling, simulation and validation of  
4 material flow on conveyor belts. *Applied Mathematical Modelling*, 38(13):3295 – 3313, 2014.
- 5 [10] M. Gröschel, A. Keimer, G. Leugering, and Z. Wang. Regularity theory and adjoint-based optimal-  
6 ity conditions for a nonlinear transport equation with nonlocal velocity. *SIAM J. Control Optim.*,  
7 52(4):2141–2163, 2014.
- 8 [11] A. Keimer and L. Pflug. Existence, uniqueness and regularity results on nonlocal balance laws. *J.*  
9 *Differential Equations*, 263(7):4023–4069, 2017.
- 10 [12] S. N. Kružkov. First order quasilinear equations with several independent variables. *Mat. Sb. (N.S.)*,  
11 81 (123):228–255, 1970.
- 12 [13] D. Li and T. Li. Shock formation in a traffic flow model with Arrhenius look-ahead dynamics. *Networks*  
13 *and Heterogeneous Media*, 6(4):681–694, 2011.
- 14 [14] E. Rossi and R. M. Colombo. Non Local Conservation Laws in Bounded Domains. *ArXiv e-prints*, Nov.  
15 2017.
- 16 [15] A. Sopasakis and M. A. Katsoulakis. Stochastic modeling and simulation of traffic flow: asymmetric  
17 single exclusion process with Arrhenius look-ahead dynamics. *SIAM J. Appl. Math.*, 66(3):921–944  
18 (electronic), 2006.