



Drinfeld double of quantum groups, tilting modules and Z -modular data associated to complex reflection groups

Abel Lacabanne

► To cite this version:

Abel Lacabanne. Drinfeld double of quantum groups, tilting modules and Z -modular data associated to complex reflection groups. *Journal of Combinatorial Algebra*, 2020, 4 (2020) (no. 3), pp. 269-323. 10.4171/JCA/45 . hal-01827849v3

HAL Id: hal-01827849

<https://hal.science/hal-01827849v3>

Submitted on 22 Oct 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DRINFELD DOUBLE OF QUANTUM GROUPS, TILTING MODULES AND \mathbb{Z} -MODULAR DATA ASSOCIATED TO COMPLEX REFLECTION GROUPS

by

ABEL LACABANNE

In this article, we construct a categorification of some Fourier matrices associated to complex reflections groups by Malle [Mal95]. To any spetsial imprimitive complex reflection group W , he attached a set of unipotent characters, which is in bijection with the set of unipotent characters of the corresponding finite reductive group when W is a Weyl group. He also defined a partition of these characters into families, the analogue of Lusztig's nonabelian Fourier transform and eigenvalues of the Frobenius. In [Cun07], Cuntz showed that for each family, this defines a so-called \mathbb{Z} -modular datum. Therefore, one can associate a \mathbb{Z} -algebra, free of finite rank over \mathbb{Z} . If the structure constants of such an algebra R are positive, it is a classical problem to find a tensor category, with some extra structure, whose Grothendieck ring is precisely R . However, these structure constants belong in general to \mathbb{Z} and the categorification is more involved.

In the case of cyclic groups, Bonnafé and Rouquier [BR17] constructed a tensor triangulated category which categorifies the modular datum attached to the non-trivial family of the cyclic spets. In the present article, we construct fusion categories with symmetric center equivalent to $\text{Rep}(G, z)$ for G a cyclic group and $z \in G$ satisfying $z^2 = 1$. If the order of G is even, we can therefore construct a new category, which is enriched over superspaces, so that its Grothendieck group has negative structure constants in general. As shown in [Lac18], this defines a \mathbb{Z} -modular datum.

Our construction is as follows: we consider the representation theory of a quantum double associated to the complex simple Lie algebra \mathfrak{sl}_{n+1} at a root of unity of even order. Similarly to the usual construction for $\mathcal{U}_q(\mathfrak{g})$ at a root of unity ξ , we consider the semisimplification of the full subcategory of tilting modules. A partial modularization of an integral subcategory gives then the category with symmetric center $\text{Rep}(G, z)$, which is denoted by $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$.

We investigate this construction in full generality considering any simple complex Lie algebra \mathfrak{g} and any root of unity. For certain values of ξ , we are able to compute explicitly the symmetric center of the category $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$, see Theorem 3.14. When \mathfrak{g} is a simple Lie algebra of type A and ξ is a root of unity of even order, the category $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ gives rise to a \mathbb{Z} -modular datum which we relate to some of the Fourier matrices constructed by Malle. The main result of this paper is given by the following theorem:

Theorem A. — *Let $n \geq 1$ and $d \geq n$ be two integers, and $\mathfrak{g} = \mathfrak{sl}_{n+1}$.*

If n is even, the symmetric center of $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ is equal to the category $\text{Rep}(\mathbb{Z}/(n+1)\mathbb{Z})$. If n is odd, the symmetric center of $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ is equal to the category $\text{Rep}(\mathbb{Z}/(n+1)\mathbb{Z}, \frac{n+1}{2})$.

In both cases, there exists a family \mathcal{F} of unipotent characters for the complex reflection group $G(d, 1, \frac{n(n+1)}{2})$ such that the Malle's \mathbb{Z} -modular data coincide with the \mathbb{Z} -modular data arising from the category $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$.

Moreover, some modular data constructed by Broué, Malle and Michel for spetsial exceptional complex reflection groups can be categorified by considering similar categories in type B .

This article is organized as follows: in the first Section, we study the algebra $\mathcal{D}_q(\mathfrak{g})$ when q is a generic parameter. Then, in the second Section, we study some finite dimensional representations of $\mathcal{D}_q(\mathfrak{g})$ and extra structure arising from the Hopf algebra $\mathcal{D}_q(\mathfrak{g})$: tensor product, braiding, pivotality. The third Section contains the main constructions of this paper: we specialize the algebra $\mathcal{D}_q(\mathfrak{g})$ at a root of unity ξ and consider tilting modules for this algebra, which allows us to construct the category $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$. In the fourth Section, we recall the Malle's \mathbb{Z} -modular datum and compare it with the modular datum arising from the category $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$. Finally, in the last Section, we study some exceptional examples arising from complex reflection groups.

Acknowledgements. — I warmly thank my advisor C. Bonnafé for many fruitful discussions and his constant support and R. Rouquier for suggesting me to consider supercategories in this work. I also thank G. Malle for many valuable comments on a first version of this paper. The paper is partially based upon work supported by the NSF under grant DMS-1440140 while the author was in residence at the Mathematical Sciences Research Institute in Berkeley, California, during the Spring 2018 semester.

1. Quantum double of Borel algebras

In this section, we recall the construction of the quantum enveloping algebra $\mathcal{U}_q(\mathfrak{g})$ associated to a simple complex Lie algebra \mathfrak{g} . Our main object of study will be the quantum double of a Borel algebra \mathfrak{b} of \mathfrak{g} , which will be denoted by $\mathcal{D}_q(\mathfrak{g})$.

1.1. Notations. — Let \mathfrak{g} be a simple complex Lie algebra of rank n . We fix a Cartan subalgebra \mathfrak{h} of \mathfrak{g} . Let $\Phi \subseteq \mathfrak{h}^*$ be the set of roots of \mathfrak{g} relative to \mathfrak{h} , $\Pi = \{\alpha_1, \dots, \alpha_n\}$ be a base of Φ , Φ^+ be the positive roots relative to Π . Let \mathfrak{b}^+ (resp. \mathfrak{b}^-) be the Borel subalgebra of \mathfrak{g} relative to Π (resp. $-\Pi$). We denote by h the Coxeter number of \mathfrak{g} and by h^\vee the dual Coxeter number of \mathfrak{g} .

Fix a symmetric bilinear form $\langle \cdot, \cdot \rangle$ on \mathfrak{h}^* normalized such that $\langle \alpha, \alpha \rangle = 2$ for short roots. For $\alpha \in \Phi$, we define $\alpha^\vee \in \mathfrak{h}^*$ by

$$\alpha^\vee = \frac{2\alpha}{\langle \alpha, \alpha \rangle}.$$

Let Q be the root lattice, P the weight lattice, P^+ the cone of dominant weights, Q^\vee the coroot lattice and P^\vee the coweight lattice

$$\begin{aligned} P &= \{\lambda \in \mathfrak{h}^* \mid \langle \lambda, \alpha^\vee \rangle \in \mathbb{Z}, \forall \alpha \in \Pi\} \\ P^+ &= \{\lambda \in \mathfrak{h}^* \mid \langle \lambda, \alpha^\vee \rangle \in \mathbb{N}, \forall \alpha \in \Pi\} \\ P^\vee &= \{\lambda \in \mathfrak{h}^* \mid \langle \lambda, \alpha \rangle \in \mathbb{Z}, \forall \alpha \in \Pi\} \end{aligned}$$

Let $(\varpi_i)_{1 \leq i \leq n}$ be the dual basis of $(\alpha_i^\vee)_{1 \leq i \leq n}$ with respect to the form $\langle \cdot, \cdot \rangle$ and ρ be the half sum of positive roots. We call the ϖ_i 's the fundamental weights. The element ρ is in P and is equal to the sum of fundamental weights [Bou68, VI.1.10 Proposition 29].

We denote by Q^+ the monoid spanned by Π . We denote by \leq the usual partial order on P : $\lambda \leq \mu$ if and only if $\mu - \lambda \in Q^+$.

For any $\alpha \in \Phi$, denote by s_α the reflection in the hyperplane orthogonal to α

$$s_\alpha(v) = v - \langle v, \alpha^\vee \rangle \alpha = v - \langle v, \alpha \rangle \alpha^\vee, \quad \forall v \in \mathfrak{h}^*.$$

For $1 \leq i \leq n$ denote by s_i the reflection s_{α_i} . All these reflections generate the Weyl group W of \mathfrak{g} and for any $w \in W$, $w(\Phi) = \Phi$. The form $\langle \cdot, \cdot \rangle$ is then invariant with respect to W . We denote by $l(w)$ the length of $w \in W$ relative to the generating set $(s_i)_{1 \leq i \leq n}$.

For q an indeterminate, define the following elements of $\mathbb{Z}[q, q^{-1}]$

$$[n]_q = \frac{q^n - q^{-n}}{q - q^{-1}}, n \in \mathbb{Z}, \quad [n]_q! = \prod_{k=1}^n [k]_q, n \in \mathbb{N} \quad \text{and} \quad \begin{bmatrix} n \\ k \end{bmatrix}_q = \prod_{i=1}^k \frac{[n+1-i]_q}{[i]_q}, n \in \mathbb{Z}, k \in \mathbb{N}.$$

We denote by $[n]_\xi$ (resp. $[n]_\xi!$, resp. $\begin{bmatrix} n \\ k \end{bmatrix}_\xi$) the evaluation of $[n]_q$ (resp. $[n]_q!$, resp. $\begin{bmatrix} n \\ k \end{bmatrix}_q$) at an invertible element ξ of a ring. We will work over the field $\mathbb{Q}(q)$ and the ring $\mathcal{A} = \mathbb{Z}[q, q^{-1}]$. We define $q_i = q^{\frac{\langle \alpha_i, \alpha_i \rangle}{2}}$.

1.2. Positive and negative parts of the usual quantum enveloping algebra. — We start by defining the usual positive and negative part of the quantum group $\mathcal{U}_q(\mathfrak{g})$.

Definition 1.1. — Let $\mathcal{U}_q(\mathfrak{b}^+)$ be the associative unital $\mathbb{Q}(q)$ -algebra with generators $K_i^{\pm 1}$, E_i , $1 \leq i \leq n$, and defining relations

$$\begin{aligned} K_i K_j &= K_j K_i, & K_i K_i^{-1} &= 1 = K_i^{-1} K_i, & \forall 1 \leq i, j \leq n, \\ K_i E_j &= q^{\langle \alpha_i, \alpha_j \rangle} E_j K_i, & \forall 1 \leq i, j \leq n, \\ \sum_{r=0}^{1-\langle \alpha_i, \alpha_j^\vee \rangle} (-1)^r \begin{bmatrix} 1-\langle \alpha_i, \alpha_j^\vee \rangle \\ r \end{bmatrix}_{q_i} E_i^{1-\langle \alpha_i, \alpha_j^\vee \rangle-r} E_j E_i^r &= 0, & \forall 1 \leq i \neq j \leq n. \end{aligned}$$

There exist several Hopf algebra structures on $\mathcal{U}_q(\mathfrak{b}^+)$, and we choose the following comultiplication Δ , counit ε and antipode S :

$$\begin{aligned} \Delta(K_i) &= K_i \otimes K_i & \varepsilon(K_i) &= 1 & S(K_i) &= K_i^{-1}, \\ \Delta(E_i) &= 1 \otimes E_i + E_i \otimes K_i & \varepsilon(E_i) &= 0 & S(E_i) &= -E_i K_i^{-1}. \end{aligned}$$

Definition 1.2. — Let $\mathcal{U}_q(\mathfrak{b}^-)$ be the associative unital $\mathbb{Q}(q)$ -algebra with generators $L_i^{\pm 1}$, F_i , $1 \leq i \leq n$, and defining relations

$$\begin{aligned} L_i L_j &= L_j L_i, & L_i L_i^{-1} &= 1 = L_i^{-1} L_i, & \forall 1 \leq i, j \leq n, \\ L_i F_j &= q^{-\langle \alpha_i, \alpha_j \rangle} F_j L_i, & \forall 1 \leq i, j \leq n, \\ \sum_{r=0}^{1-\langle \alpha_i, \alpha_j^\vee \rangle} (-1)^r \begin{bmatrix} 1-\langle \alpha_i, \alpha_j^\vee \rangle \\ r \end{bmatrix}_{q_i} F_i^{1-\langle \alpha_i, \alpha_j^\vee \rangle-r} F_j F_i^r &= 0, & \forall 1 \leq i \neq j \leq n. \end{aligned}$$

There exist several Hopf algebra structures on $\mathcal{U}_q(\mathfrak{b}^-)$, and we choose the following comultiplication Δ , counit ε and antipode S :

$$\begin{aligned} \Delta(L_i) &= L_i \otimes L_i & \varepsilon(L_i) &= 1 & S(L_i) &= L_i^{-1}, \\ \Delta(F_i) &= L_i^{-1} \otimes F_i + F_i \otimes 1 & \varepsilon(F_i) &= 0 & S(F_i) &= -L_i F_i. \end{aligned}$$

For $\lambda = \sum_{i=1}^n \lambda_i \alpha_i \in Q$, we denote by K_λ (resp. L_λ) the element $\prod_{i=1}^n K_i^{\lambda_i}$ (resp. $\prod_{i=1}^n L_i^{\lambda_i}$).

1.3. A Hopf pairing between $\mathcal{U}_q(\mathfrak{b}^+)$ and $\mathcal{U}_q(\mathfrak{b}^-)$. — The following is due to Drinfeld [Dri87, Section 13] and has been studied by Tanisaki [Tan92].

Proposition 1.3. — *There exists a unique bilinear form $(\cdot, \cdot): \mathcal{U}_q(\mathfrak{b}^+) \times \mathcal{U}_q(\mathfrak{b}^-) \rightarrow \mathbb{Q}(q)$ satisfying for all $x, x' \in \mathcal{U}_q(\mathfrak{b}^+)$, $y, y' \in \mathcal{U}_q(\mathfrak{b}^-)$ and $1 \leq i \leq n$*

1. $(x, y y') = (\Delta(x), y \otimes y')$,
2. $(x x', y) = (x' \otimes x, \Delta(y))$,
3. $(x, 1) = \varepsilon(x)$, $(1, y) = \varepsilon(y)$,
4. $(K_i, L_j) = q^{\langle \alpha_i, \alpha_j \rangle}$,
5. $(K_i, F_j) = 0 = (E_i, L_j)$,
6. $(E_i, F_j) = \frac{\delta_{i,j}}{q_i - q_i^{-1}}$.

This bilinear form endows the tensor product $\mathcal{U}_q(\mathfrak{b}^-) \otimes \mathcal{U}_q(\mathfrak{b}^+)$ with a Hopf algebra structure, which we denote by $\mathcal{D}_q(\mathfrak{g})$, see [KS97, Section 8.2]. Using Sweedler notation for the coproduct, the product of $y_1 \otimes x_1$ and $y_2 \otimes x_2$ in $\mathcal{U}_q(\mathfrak{b}^-) \otimes \mathcal{U}_q(\mathfrak{b}^+)$ is given by

$$(y_1 \otimes x_1)(y_2 \otimes x_2) = \sum_{(x_1)(y_2)} (x'_1, y'_2)(x''_1, S(y''_2)) y_1 y'_2 \otimes x'_1 x_2.$$

Consequently, both $\mathcal{U}_q(\mathfrak{b}^-) \otimes 1$ and $1 \otimes \mathcal{U}_q(\mathfrak{b}^+)$ are Hopf subalgebras of $\mathcal{D}_q(\mathfrak{g})$. In the following, we identify $\mathcal{U}_q(\mathfrak{b}^-)$ and $\mathcal{U}_q(\mathfrak{b}^+)$ with their images in $\mathcal{D}_q(\mathfrak{g})$ and will write $y x$ instead of $y \otimes x$.

Proposition 1.4. — *The algebra $\mathcal{D}_q(\mathfrak{g})$ is the associative unital $\mathbb{Q}(s)$ -algebra with generators $K_i^{\pm 1}$, $L_i^{\pm 1}$, E_i and F_i , $1 \leq i \leq n$ and defining relations*

$$\begin{aligned} K_i K_j &= K_j K_i, & K_i K_i^{-1} &= 1 = K_i^{-1} K_i, \\ L_i L_j &= L_j L_i, & L_i L_i^{-1} &= 1 = L_i^{-1} L_i, \\ K_i L_j &= L_j K_i, \\ K_i E_j &= q^{\langle \alpha_i, \alpha_j \rangle} E_j K_i, & K_i F_j &= q^{-\langle \alpha_i, \alpha_j \rangle} F_j K_i, \\ L_i E_j &= q^{\langle \alpha_i, \alpha_j \rangle} E_j L_i, & L_i F_j &= q^{-\langle \alpha_i, \alpha_j \rangle} F_j L_i, \\ [E_i, F_j] &= \delta_{i,j} \frac{K_i - L_i^{-1}}{q_i - q_i^{-1}}, \end{aligned}$$

$$\begin{aligned} \sum_{r=0}^{1-\langle \alpha_i, \alpha_j^\vee \rangle} (-1)^r \begin{bmatrix} 1-\langle \alpha_i, \alpha_j^\vee \rangle \\ r \end{bmatrix}_{q_i} E_i^{1-\langle \alpha_i, \alpha_j^\vee \rangle-r} E_j E_i^r &= 0, & 1 \leq i \neq j \leq n, \\ \sum_{r=0}^{1-\langle \alpha_i, \alpha_j^\vee \rangle} (-1)^r \begin{bmatrix} 1-\langle \alpha_i, \alpha_j^\vee \rangle \\ r \end{bmatrix}_{q_i} F_i^{1-\langle \alpha_i, \alpha_j^\vee \rangle-r} F_j F_i^r &= 0, & 1 \leq i \neq j \leq n. \end{aligned}$$

The elements $z_i = K_i L_i^{-1}$ are central and $\mathcal{U}_q(\mathfrak{g})$ is the quotient of $\mathcal{D}_q(\mathfrak{g})$ by the Hopf ideal generated by $(z_i - 1)_{1 \leq i \leq n}$.

Denote by $\mathcal{D}_q(\mathfrak{g})^{<0}$ (resp. $\mathcal{D}_q(\mathfrak{g})^0$, resp. $\mathcal{D}_q(\mathfrak{g})^{>0}$) the subalgebra of $\mathcal{D}_q(\mathfrak{g})$ generated by $(E_i)_{1 \leq i \leq n}$ (resp. $(K_i, L_i)_{1 \leq i \leq n}$, resp. $(F_i)_{1 \leq i \leq n}$). Multiplication yields an isomorphism of vector spaces $\mathcal{D}_q(\mathfrak{g})^{<0} \otimes \mathcal{D}_q(\mathfrak{g})^0 \otimes \mathcal{D}_q(\mathfrak{g})^{>0} \simeq \mathcal{D}_q(\mathfrak{g})$. It is worth mentioning that the coproduct and antipode of $\mathcal{D}_q(\mathfrak{g})$ do not restrict to $\mathcal{D}_q(\mathfrak{g})^{<0}$ or $\mathcal{D}_q(\mathfrak{g})^{>0}$.

The square of the antipode is given by the conjugation by any element of the form $L_\lambda K_{2\rho-\lambda}$, $\lambda \in Q$ as it is easily checked on the generators.

1.4. Graduation. — There exists a Q -graduation on $\mathcal{D}_q(\mathfrak{g})$ given by $\deg(K_i) = 0 = \deg(L_i)$, $\deg(E_i) = \alpha_i$ and $\deg(F_i) = -\alpha_i$. For $\lambda \in Q$, we denote by $\mathcal{D}_q(\mathfrak{g})_\lambda$ the homogeneous elements of degree λ . We have

$$\mathcal{D}_q(\mathfrak{g})_\lambda = \{v \in \mathcal{D}_q(\mathfrak{g}) \mid K_i v = q^{\langle \lambda, \alpha_i \rangle} v K_i, \forall 1 \leq i \leq n\}.$$

The coproduct, the counit and the antipode respect the grading. Moreover, the same lemma as [Jan96, lemma 4.12] shows that for $\mu \in Q$, $\mu \geq 0$

$$\Delta(\mathcal{D}_q(\mathfrak{g})_\mu^{>0}) \subseteq \bigoplus_{0 \leq \nu \leq \mu} \mathcal{D}_q(\mathfrak{g})_\nu^{>0} \otimes K_\nu \mathcal{D}_q(\mathfrak{g})_{\mu-\nu}^{>0}$$

and

$$\Delta(\mathcal{D}_q(\mathfrak{g})_{-\mu}^{<0}) \subseteq \bigoplus_{0 \leq \nu \leq \mu} L_\nu^{-1} \mathcal{D}_q(\mathfrak{g})_{-(\mu-\nu)}^{<0} \otimes \mathcal{D}_q(\mathfrak{g})_{-\nu}^{<0}.$$

Therefore, for $x \in \mathcal{D}_q(\mathfrak{g})_\mu^{>0}$, there are elements $r_i(x)$ and $r'_i(x)$ in $\mathcal{D}_q(\mathfrak{g})_{\mu-\alpha_i}^{>0}$ such that

$$(1) \quad \Delta(x) \in 1 \otimes x + \sum_{i=1}^n E_i \otimes K_i r_i(x) + \sum_{\substack{0 < \nu \leq \mu \\ \nu \notin \Pi}} \mathcal{D}_q(\mathfrak{g})_\nu^{>0} \otimes K_\nu \mathcal{D}_q(\mathfrak{g})_{\mu-\nu}^{>0},$$

and

$$(2) \quad \Delta(x) \in x \otimes K_\mu + \sum_{i=1}^n r'_i(x) \otimes K_{\mu-\alpha_i} E_i + \sum_{\substack{0 < \nu \leq \mu \\ \mu-\nu \notin \Pi}} \mathcal{D}_q(\mathfrak{g})_\nu^{>0} \otimes K_\nu \mathcal{D}_q(\mathfrak{g})_{\mu-\nu}^{>0}.$$

Similarly, we can define $\rho_i(y)$ and $\rho'_i(y)$ for $y \in \mathcal{D}_q(\mathfrak{g})_{-\mu}^{<0}$ by the following

$$\Delta(y) \in L_\mu^{-1} \otimes y + \sum_{i=1}^n L_{\mu-\alpha_i}^{-1} F_i \otimes \rho_i(y) + \sum_{\substack{0 < \nu \leq \mu \\ \mu-\nu \notin \Pi}} L_\nu^{-1} \mathcal{D}_q(\mathfrak{g})_{-(\mu-\nu)}^{<0} \otimes \mathcal{D}_q(\mathfrak{g})_{-\nu}^{<0},$$

and

$$\Delta(y) \in y \otimes 1 + \sum_{i=1}^n L_{\alpha_i}^{-1} \rho'_i(y) \otimes F_i + \sum_{\substack{0 < \nu \leq \mu \\ \nu \notin \Pi}} L_\nu^{-1} \mathcal{D}_q(\mathfrak{g})_{-(\mu-\nu)}^{<0} \otimes \mathcal{D}_q(\mathfrak{g})_{-\nu}^{<0}.$$

The values of r_i , r'_i , ρ_i and ρ'_i can be computed by induction. Compare with [Jan96, 6.14, 6.15].

Lemma 1.5. — Let $\mu, \mu' \in Q$, $\mu \geq 0$ and $\mu' \geq 0$.

1. For all $x \in \mathcal{D}_q(\mathfrak{g})_\mu^{>0}$ and $x' \in \mathcal{D}_q(\mathfrak{g})_{\mu'}^{>0}$,

$$r_i(x x') = q^{-(\alpha_i, \mu)} x r_i(x') + r_i(x) x', \quad r'_i(x x') = x r'_i(x') + q^{-(\alpha_i, \mu')} r'_i(x) x'.$$

2. For all $y \in \mathcal{D}_q(\mathfrak{g})_{-\mu}^{<0}$ and $y' \in \mathcal{D}_q(\mathfrak{g})_{-\mu'}^{<0}$,

$$\rho_i(y y') = y \rho_i(y') + q^{-(\alpha_i, \mu')} \rho_i(y) y', \quad \rho'_i(y y') = q^{-(\alpha_i, \mu)} y \rho'_i(y') + \rho'_i(y) y'.$$

These elements can also be used to compute some commutators.

Lemma 1.6. — Let $\mu \in Q$, $\mu \geq 0$. Let $x \in \mathcal{D}_q(\mathfrak{g})_\mu^{>0}$ and $y \in \mathcal{D}_q(\mathfrak{g})_{-\mu}^{\leq 0}$. Then

$$xF_i - F_i x = \frac{K_i r_i(x) - r'_i(x) L_i^{-1}}{q_i - q_i^{-1}}$$

and

$$E_i y - y E_i = \frac{\rho_i(y) K_i - L_i^{-1} \rho'_i(y)}{q_i - q_i^{-1}}.$$

Proof. — We only show the first formula, the second one is proven similarly. If $x = 1$ or $x = E_i$, the formula is satisfied. Now, supposing that it is true for x and x' , we show it for xx' :

$$\begin{aligned} xx'F_i - F_i xx' &= x(x'F_i - F_i x') + (xF_i - F_i x)x' \\ &= (q_i - q_i^{-1})^{-1}(x(K_i r_i(x') - r'_i(x') L_i^{-1}) + (K_i r_i(x) - r'_i(x) L_i^{-1})x') \\ &= (q_i - q_i^{-1})^{-1}(K_i(q^{-(a_i, \mu)} x r_i(x') + r_i(x)x') - (x r'_i(x') + q^{-(a_i, \mu')} r'_i(x)x')) \\ &= (q_i - q_i^{-1})^{-1}(K_i r_i(xx') - r'_i(xx') L_i^{-1}), \end{aligned}$$

where we used Lemma 1.5 in the last equality. \square

1.5. Some properties of the pairing. — We gather here some well known properties of the pairing.

Proposition 1.7. — Let $x \in \mathcal{D}_q(\mathfrak{g})^{>0}$, $y \in \mathcal{D}_q(\mathfrak{g})^{\leq 0}$ and $\lambda, \mu \in Q$. Then

1. $(K_\lambda x, L_\mu y) = (K_\lambda, L_\mu)(x, y)$.
2. Suppose $\lambda, \mu \geq 0$. If x is of weight λ and y of weight $-\mu$ then $(x, y) = 0$ if $\lambda \neq \mu$.

Proof. — For the first equality, it suffices to show it for x and y homogeneous. It then follows easily from (1), (2) and the fact that $(K_\lambda, y') = 0 = (x', L_\mu)$ for any $x' \in \mathcal{D}_q(\mathfrak{g})^{>0}$ and $y' \in \mathcal{D}_q(\mathfrak{g})^{\leq 0}$.

For the second assertion, we proceed by induction on $\text{ht}(\lambda) = \sum_{i=1}^n \lambda_i$, where $\lambda = \sum_{i=1}^n \lambda_i \alpha_i$. We can suppose that x is a product of the generators E_i . The case $\text{ht}(\lambda) = 1$, i.e. $\lambda \in \Pi$ is easily proved by induction on $\text{ht}(\mu)$. Then, writing $x = E_i x'$ and

$$\Delta(y) = \sum_{0 \leq v \leq \mu} L_v^{-1} y'_{-(\mu-v)} \otimes y''_{-v}, \quad y'_{-v}, y''_{-v} \in \mathcal{D}_q(\mathfrak{g})_{-v}^{\leq 0},$$

one has

$$(E_i x', y) = \sum_{0 \leq v \leq \mu} (x', L_v^{-1} y'_{-(\mu-v)})(E_i, y''_{-v}) = (x', y'_{-(\mu-\alpha_i)})(E_i, y''_{-\alpha_i}).$$

As $\lambda \neq \mu$, the two terms can not be simultaneously non-zero. \square

Using these two facts and an easy induction, one can show that, for $1 \leq i \leq n$, and $r \in \mathbb{N}$,

$$(E_i^r, F_i^r) = \frac{q_i^{-\frac{r(r-1)}{2}} [r]_{q_i}!}{(q_i - q_i^{-1})^r}.$$

We now turn to the compatibility between the pairing and the r_i , r'_i , ρ_i and ρ'_i defined in section 1.4.

Lemma 1.8. — Let $x \in \mathcal{D}_q(\mathfrak{g})_\mu^{>0}$ and $y \in \mathcal{D}_q(\mathfrak{g})_{-\mu}^{\leq 0}$. One has

$$\begin{aligned} (x, F_i y) &= (E_i, F_i)(r_i(x), y), & (x, y F_i) &= (E_i, F_i)(r'_i(x), y), \\ (E_i x, y) &= (E_i, F_i)(x, \rho'_i(y)), & (x E_i, y) &= (E_i, F_i)(x, \rho_i(y)). \end{aligned}$$

Proof. — It follows easily from (1), (2) and Proposition 1.7. \square

We end this section with [Tan92, Proposition 2.1.4].

Proposition 1.9. — The restriction of the pairing to $\mathcal{D}_q(\mathfrak{g})^{>0} \times \mathcal{D}_q(\mathfrak{g})^{\leq 0}$ is non-degenerate.

1.6. Basis for $\mathcal{D}_q(\mathfrak{g})$. — We introduce the divided power $E_i^{(r)}$ and $F_i^{(r)}$ for $1 \leq i \leq n$ and $r \leq 0$:

$$E_i^{(r)} = \frac{E_i^r}{[r]_{q_i}} \quad \text{and} \quad F_i^{(r)} = \frac{F_i^r}{[r]_{q_i}}.$$

The following is the analogue of [Lus90, Theorem 3.1] for $\mathcal{D}_q(\mathfrak{g})$.

Proposition 1.10. — For all $1 \leq i \leq n$, there exists a unique $\mathbb{Q}(q)$ -algebra isomorphism T_i such that:

$$T_i(K_\lambda) = K_{s_i(\lambda)} \quad T_i(L_\lambda) = L_{s_i(\lambda)}, \quad \forall \lambda \in Q,$$

and, for $1 \leq j \leq n$, setting $r = -\langle \alpha_j, \alpha_i^\vee \rangle$,

$$\begin{aligned} T_i(E_j) &= \begin{cases} -F_i L_i & \text{if } i = j, \\ \sum_{k=0}^r (-1)^k q_i^{-k} E_i^{(r-k)} E_j E_i^{(k)} & \text{otherwise,} \end{cases} \\ T_i(F_j) &= \begin{cases} -K_i^{-1} E_i & \text{if } i = j, \\ \sum_{k=0}^r (-1)^k q_i^k F_i^{(k)} F_j F_i^{(r-k)} & \text{otherwise,} \end{cases} \end{aligned}$$

We also have the analogue of [Lus90, Theorem 3.2] for $\mathcal{D}_q(\mathfrak{g})$.

Proposition 1.11. — The $(T_i)_{1 \leq i \leq n}$ satisfy the braid relations and therefore define a morphism from the braid group of W to the algebra automorphisms of $\mathcal{D}_q(\mathfrak{g})$.

Choose now a reduced decomposition $w_0 = s_{i_1} s_{i_2} \cdots s_{i_r}$ of the longest element w_0 of W . The positive roots are then

$$\Phi^+ = \{s_{i_1} s_{i_2} \cdots s_{i_k}(\alpha_{i_{k+1}}) \mid 0 \leq k \leq r-1\}.$$

For any $\alpha = s_{i_1} s_{i_2} \cdots s_{i_k}(\alpha_{i_{k+1}}) \in \Phi^+$ define

$$E_\alpha = T_{i_1} T_{i_2} \cdots T_{i_k}(E_{i_{k+1}}) \quad \text{and} \quad F_\alpha = T_{i_1} T_{i_2} \cdots T_{i_k}(F_{i_{k+1}}).$$

Note that E_α (resp F_α) is homogeneous of degree α (resp. $-\alpha$). The choice of a reduced decomposition of w_0 gives an order on Φ^+ , namely $s_{i_1} s_{i_2} \cdots s_{i_k}(\alpha_{i_{k+1}}) \preccurlyeq s_{i_1} s_{i_2} \cdots s_{i_{k+1}}(\alpha_{i_{k+2}})$ for every k . All products will be ordered with respect to this order.

Proposition 1.12 ([Lus90, Theorem 6.7]). — The elements

$$\prod_{\alpha \in \Phi^+} E_\alpha^{(n_\alpha)}, \quad n_\alpha \in \mathbb{N}$$

form a $\mathbb{Q}(q)$ -basis of $\mathcal{D}_q(\mathfrak{g})^{>0}$.

The elements

$$\prod_{\alpha \in \Phi^+} F_\alpha^{(n_\alpha)}, \quad n_\alpha \in \mathbb{N}$$

form a $\mathbb{Q}(q)$ -basis of $\mathcal{D}_q(\mathfrak{g})^{<0}$.

We compute now the dual basis of $\mathcal{D}_q(\mathfrak{g})^{>0}$ with respect to (\cdot, \cdot) . The following can be found in [Jan96, 8.29].

Proposition 1.13. — For any $\alpha \in \Phi^+$, let $a_\alpha, b_\alpha \in \mathbb{N}$. We have

$$\left(\prod_{\alpha \in \Phi^+} E_\alpha^{a_\alpha}, \prod_{\alpha \in \Phi^+} F_\alpha^{b_\alpha} \right) = \prod_{\alpha \in \Phi^+} \delta_{a_\alpha, b_\alpha} \frac{q_\alpha^{\frac{a_\alpha(a_\alpha-1)}{2}} [a_\alpha]_{q_\alpha}!}{(q_\alpha - q_\alpha^{-1})^{a_\alpha}}.$$

Therefore, the dual basis of $\left(\prod_{\alpha \in \Phi^+} E_\alpha^{(n_\alpha)} \right)_{(n_\alpha) \in \mathbb{N}^{\Phi^+}}$ is

$$\left(\prod_{\alpha \in \Phi^+} q_\alpha^{\frac{n_\alpha(n_\alpha-1)}{2}} [n_\alpha]_{q_\alpha}! (q_\alpha - q_\alpha^{-1})^{n_\alpha} F_\alpha^{(n_\alpha)} \right)_{(n_\alpha) \in \mathbb{N}^{\Phi^+}}.$$

1.7. Quasi- R -matrix. — The algebra $\mathcal{D}_q(\mathfrak{g})$ is not quasi-triangular. Nevertheless, we construct a quasi- R -matrix which will endow the usual category of modules with a braiding (see section 2.3). We adapt the exposition of Jantzen [Jan96, Chapter 7] to the case of $\mathcal{D}_q(\mathfrak{g})$.

For any $\mu \in Q$, $\mu \geq 0$, fix a basis $(u_i^\mu)_{i \in I_\mu}$ of $\mathcal{D}_q(\mathfrak{g})_\mu^{>0}$ and let $(v_i^\mu)_{i \in I_\mu}$ the dual basis in $\mathcal{D}_q(\mathfrak{g})_\mu^{<0}$ with respect to the pairing (\cdot, \cdot) . Set

$$\Theta_\mu = \sum_{i \in I_\mu} u_i^\mu \otimes v_i^\mu \in \mathcal{D}_q(\mathfrak{g}) \otimes \mathcal{D}_q(\mathfrak{g}).$$

Note that this does not depend on the choice of the basis of $\mathcal{D}_q(\mathfrak{g})_\mu$. A homogeneous basis of $\mathcal{D}_q(\mathfrak{g})^{>0}$ and its dual with respect to (\cdot, \cdot) have already been computed above. Following [Ros90, A.1], we define an algebra automorphism Ψ of $\mathcal{D}_q(\mathfrak{g}) \otimes \mathcal{D}_q(\mathfrak{g})$ by

$$\begin{aligned} \Psi(K_i \otimes 1) &= K_i \otimes 1, & \Psi(1 \otimes K_i) &= 1 \otimes K_i, \\ \Psi(L_i \otimes 1) &= L_i \otimes 1, & \Psi(1 \otimes L_i) &= 1 \otimes L_i, \\ \Psi(E_i \otimes 1) &= E_i \otimes L_i^{-1}, & \Psi(1 \otimes E_i) &= K_i^{-1} \otimes E_i, \\ \Psi(F_i \otimes 1) &= F_i \otimes L_i, & \Psi(1 \otimes F_i) &= K_i \otimes F_i. \end{aligned}$$

Proposition 1.14. — Let $\mu \in Q$, $\mu \geq 0$. We have

$$\begin{aligned} \Theta_\mu(K_i \otimes K_i) &= \Psi(K_i \otimes K_i) \Theta_\mu, \\ \Theta_\mu(L_i \otimes L_i) &= \Psi(L_i \otimes L_i) \Theta_\mu, \\ \Theta_\mu(1 \otimes E_i) + \Theta_{\mu-\alpha_i}(E_i \otimes K_i) &= \Psi(E_i \otimes 1) \Theta_{\mu-\alpha_i} + \Psi(K_i \otimes E_i) \Theta_\mu, \\ \Theta_\mu(F_i \otimes 1) + \Theta_{\mu-\alpha_i}(L_i^{-1} \otimes F_i) &= \Psi(1 \otimes F_i) \Theta_{\mu-\alpha_i} + \Psi(F_i \otimes L_i^{-1}) \Theta_\mu. \end{aligned}$$

Proof. — We follow closely the proof of [Jan96, Lemma 7.1]. The first two equations are trivial. We will use the following fact: for any $\mu \in Q$, $\mu \geq 0$, and any $x \in \mathcal{D}_q(\mathfrak{g})_\mu^{>0}$ and $y \in \mathcal{D}_q(\mathfrak{g})_\mu^{<0}$ we have

$$x = \sum_i (x, v_i^\mu) u_i^\mu \quad \text{and} \quad y = \sum_i (u_i^\mu, y) v_i^\mu.$$

We set $c_i = (q_i - q_i^{-1})^{-1}$, and we have

$$\begin{aligned}
 (1 \otimes E_i)\Theta_\mu - \Theta_\mu(1 \otimes E_i) &= \sum_j u_j^\mu \otimes (E_i v_j^\mu - v_j^\mu E_i) \\
 &= c_i \sum_j u_j^\mu \otimes (\rho_i(v_j^\mu) K_i - L_i^{-1} \rho'_i(v_j^\mu)) && \text{by Lemma 1.6} \\
 &= c_i \sum_{j,k} u_j^\mu \otimes [(u_k^{\mu-\alpha_i}, \rho_i(v_j^\mu)) v_k^{\mu-\alpha_i} K_i - (u_k^{\mu-\alpha_i}, \rho'_i(v_j^\mu)) L_i^{-1} v_k^{\mu-\alpha_i}] \\
 &= c_i \sum_{j,k} u_j^\mu \otimes [(u_k^{\mu-\alpha_i} E_i, v_j^\mu) v_k^{\mu-\alpha_i} K_i - (E_i u_k^{\mu-\alpha_i}, v_j^\mu) L_i^{-1} v_k^{\mu-\alpha_i}] && \text{by Lemma 1.8} \\
 &= \sum_k (u_k^{\mu-\alpha_i} E_i \otimes v_k^{\mu-\alpha_i} K_i - E_i u_k^{\mu-\alpha_i} \otimes L_i^{-1} v_k^{\mu-\alpha_i}) \\
 &= \Theta_{\mu-\alpha_i}(E_i \otimes K_i) - (E_i \otimes L_i^{-1})\Theta_{\mu-\alpha_i},
 \end{aligned}$$

as expected because $\Psi(E_i \otimes 1) = E_i \otimes L_i^{-1}$ and $\Psi(K_i \otimes E_i) = 1 \otimes E_i$. A similar calculation shows the fourth formula. \square

Define, as in [Lus10, Chapter 4], the completion $\mathcal{D}_q(\mathfrak{g}) \hat{\otimes} \mathcal{D}_q(\mathfrak{g})$ of $\mathcal{D}_q(\mathfrak{g}) \otimes \mathcal{D}_q(\mathfrak{g})$ with respect to the descending sequence of spaces

$$(\mathcal{D}_q(\mathfrak{g}) \otimes \mathcal{D}_q(\mathfrak{g}))_N = \mathcal{D}_q(\mathfrak{g}) \otimes \sum_{\text{ht}(\mu) \geq N} \mathcal{D}_q(\mathfrak{g})^{>0} \mathcal{D}_q(\mathfrak{g})^0 \mathcal{D}_q(\mathfrak{g})_{-\mu}^{<0} + \sum_{\text{ht}(\mu) \geq N} \mathcal{D}_q(\mathfrak{g})^{<0} \mathcal{D}_q(\mathfrak{g})^0 \mathcal{D}_q(\mathfrak{g})_\mu^{>0} \otimes \mathcal{D}_q(\mathfrak{g}).$$

The morphism Ψ extends by continuity to $\mathcal{D}_q(\mathfrak{g}) \hat{\otimes} \mathcal{D}_q(\mathfrak{g})$ and we consider the following element of $\mathcal{D}_q(\mathfrak{g}) \hat{\otimes} \mathcal{D}_q(\mathfrak{g})$

$$\Theta = \sum_{\substack{\mu \in Q \\ \mu \geq 0}} \Theta_\mu.$$

Then we can rewrite Proposition 1.14 as

$$\Theta \Delta(u) = (\Psi \circ \Delta^{\text{op}})(u) \Theta$$

for any $u \in \mathcal{D}_q(\mathfrak{g})$.

We also have the analogue of [Jan96, Lemma 7.4].

Lemma 1.15. — For $\mu \in Q$, $\mu \geq 0$, we have

$$(\Delta \otimes \text{id})(\Theta_\mu) = \sum_{0 \leq \nu \leq \mu} (\Theta_\nu)_{13} (1 \otimes K_\nu \otimes 1) (\Theta_{\mu-\nu})_{23},$$

and

$$(\text{id} \otimes \Delta)(\Theta_\mu) = \sum_{0 \leq \nu \leq \mu} (\Theta_\nu)_{13} (1 \otimes L_\nu^{-1} \otimes 1) (\Theta_{\mu-\nu})_{12}.$$

Proof. — First, note that for any $x \in \mathcal{D}_q(\mathfrak{g})_\mu^{>0}$

$$\Delta(x) = \sum_{\substack{0 \leq \nu \leq \mu \\ i,j}} (x, v_i^\nu v_j^{\mu-\nu}) u_i^\nu \otimes K_\nu u_j^{\mu-\nu},$$

and for any $y \in \mathcal{D}_q(\mathfrak{g})_{-\mu}^{<0}$

$$\Delta(y) = \sum_{\substack{0 \leq \nu \leq \mu \\ i,j}} (u_i^\nu u_j^{\mu-\nu}, y) L_\nu^{-1} v_j^{\mu-\nu} \otimes v_i^\nu.$$

Therefore

$$\begin{aligned}
(\Delta \otimes \text{id})(\Theta_\mu) &= \sum_k \Delta(u_k^\mu) \otimes v_k^\mu \\
&= \sum_{\substack{0 \leq \nu \leq \mu \\ i, j, k}} (u_k^\mu, v_i^\nu v_j^{\mu-\nu}) u_i^\nu \otimes K_\nu u_j^{\mu-\nu} \otimes v_k^\mu \\
&= \sum_{\substack{0 \leq \nu \leq \mu \\ i, j}} u_i^\nu \otimes K_\nu u_j^{\mu-\nu} \otimes v_i^\nu v_j^{\mu-\nu} \\
&= \sum_{0 \leq \nu \leq \mu} (\Theta_\nu)_{13} (1 \otimes K_\nu \otimes 1) (\Theta_{\mu-\nu})_{23}.
\end{aligned}$$

The proof of the other formula is similar. \square

We can translate this last lemma as equalities in $\mathcal{D}_q(\mathfrak{g}) \hat{\otimes} \mathcal{D}_q(\mathfrak{g})$. Note that for any $x \in \mathcal{D}_q(\mathfrak{g})_\mu^{>0}$ and $y \in \mathcal{D}_q(\mathfrak{g})_{-\mu}^{\leq 0}$, we have

$$\begin{aligned}
\Psi(x \otimes 1) &= x \otimes L_\mu^{-1}, & \Psi(1 \otimes x) &= K_\mu^{-1} \otimes x, \\
\Psi(y \otimes 1) &= y \otimes L_\mu, & \Psi(1 \otimes y) &= K_\mu \otimes y.
\end{aligned}$$

Therefore, we have

$$\begin{aligned}
(\Delta \otimes \text{id})(\Theta) &= \sum_{\mu \geq 0} \sum_{\nu+\eta=\mu} (\Theta_\eta)_{13} (1 \otimes K_\eta \otimes 1) (\Theta_\nu)_{23} \\
&= \sum_{\mu \geq 0} \sum_{\nu+\eta=\mu} \Psi_{23}((\Theta_\eta)_{13}) (\Theta_\nu)_{23} \\
&= \Psi_{23}(\Theta_{13}) \Theta_{23}.
\end{aligned}$$

Similarly, we have

$$(\text{id} \otimes \Delta)(\Theta) = \Psi_{12}(\Theta_{13}) \Theta_{12}.$$

We now show that the element Θ is invertible. Set $\Gamma_\mu = (S \otimes \text{id})(\Theta_\mu)(K_\mu \otimes 1)$ and $\Gamma = \sum_{\mu \geq 0} \Gamma_\mu$.

Lemma 1.16. — *We have $\Gamma\Theta = 1 = \Theta\Gamma$ in $\mathcal{D}_q(\mathfrak{g}) \hat{\otimes} \mathcal{D}_q(\mathfrak{g})$.*

Proof. — We show that for all $\mu \geq 0$,

$$\sum_{\lambda+\nu=\mu} \Gamma_\lambda \Theta_\nu = \delta_{\mu,0},$$

and

$$\sum_{\lambda+\nu=\mu} \Theta_\lambda \Gamma_\nu = \delta_{\mu,0}.$$

We start with the first formula. We may and will suppose that $\mu > 0$. As

$$\sum_{\lambda+\nu=\mu} \Gamma_\lambda \Theta_\nu = \sum_{\substack{\lambda+\nu=\mu \\ i, j}} S(u_i^\lambda) K_\lambda u_j^\nu \otimes v_i^\lambda v_j^\nu$$

is in $\mathcal{D}_q(\mathfrak{g}) \otimes \mathcal{D}_q(\mathfrak{g})_{-\mu}^{\leq 0}$, it suffices to show that applying $\text{id} \otimes (x, \cdot)$ gives zero, for all $x \in \mathcal{D}_q(\mathfrak{g})_\mu^{>0}$. But

$$\Delta(x) = \sum_{\substack{\lambda+\nu=\mu \\ i, j}} (x, v_i^\lambda v_j^\nu) u_i^\lambda \otimes K_\lambda u_j^\nu,$$

so the antipode axiom gives

$$0 = \varepsilon(x) = \sum_{\substack{\lambda+\nu=\mu \\ i,j}} \left(x, v_i^\lambda v_j^\nu \right) S(u_i^\lambda) K_\lambda u_j^\nu$$

as desired.

For the second equality, we again apply $\text{id} \otimes (x, \cdot)$ to show that

$$\sum_{\lambda+\nu=\mu} \Theta_\lambda \Gamma_\nu = \sum_{\substack{\lambda+\nu=\mu \\ i,j}} u_i^\lambda S(u_j^\nu) K_\nu \otimes v_i^\lambda v_j^\nu = 0.$$

The antipode axiom also gives

$$0 = \varepsilon(x) = \sum_{\substack{\lambda+\nu=\mu \\ i,j}} \left(x, v_i^\lambda v_j^\nu \right) u_i^\lambda S(u_j^\nu) K_{-\lambda}$$

which is, after multiplication by K_μ , the result expected. \square

We therefore have proven:

Proposition 1.17. — *The element $\Theta \in \mathcal{D}_q(\mathfrak{g}) \hat{\otimes} \mathcal{D}_q(\mathfrak{g})$ is invertible and*

- *for all $u \in \mathcal{U}_q(\mathfrak{g})$ we have $\Theta \Delta(u) = (\Psi \circ \Delta^{\text{op}})(u) \Theta$,*
- *$(\Delta \otimes \text{id})(\Theta) = \Psi_{23}(\Theta_{13}) \Theta_{23}$,*
- *$(\text{id} \otimes \Delta)(\Theta) = \Psi_{12}(\Theta_{13}) \Theta_{12}$.*

Finally, we give an explicit form of Θ (compare with [CP94, 10.1.D])

$$\Theta = \prod_{\alpha \in \Phi^+} \left(\sum_{n=0}^{+\infty} q_\alpha^{\frac{n(n-1)}{2}} [n]_{q_\alpha}! (q_\alpha - q_\alpha^{-1})^n E_\alpha^{(n)} \otimes F_\alpha^{(n)} \right).$$

2. Representation theory at a generic parameter q

We now turn to the representation theory of the quantum double $\mathcal{D}_q(\mathfrak{g})$, which is quite similar to the one of $\mathcal{U}_q(\mathfrak{g})$.

2.1. Representations of $\mathcal{D}_q(\mathfrak{g})$. — We will consider the category \mathcal{C}_q of finite dimensional $\mathcal{D}_q(\mathfrak{g})$ -modules M such that

$$M = \bigoplus_{(\lambda, \mu) \in P \times P} M_{(\lambda, \mu)},$$

where $M_{\lambda, \mu}$ denote the weight space of M associated to $(\lambda, \mu) \in P \times P$:

$$M_{\lambda, \mu} = \left\{ m \in M \mid K_i \cdot m = q_i^{\langle \lambda, \alpha_i^\vee \rangle} m, L_i \cdot m = q_i^{\langle \mu, \alpha_i^\vee \rangle} m, \forall 1 \leq i \leq n \right\}.$$

As $\mathcal{D}_q(\mathfrak{g})$ is a Hopf algebra, \mathcal{C}_q is a monoidal rigid abelian category. For $(\lambda, \mu) \in P \times P$ and $(\lambda', \mu') \in P \times P$, we write $(\lambda', \mu') \leq (\lambda, \mu)$ if $(\lambda' - \lambda, \mu' - \mu) = \sum_{\alpha \in \Delta} n_\alpha (\alpha, \alpha)$ with $n_\alpha \in \mathbb{N}$.

Proposition 2.1. — *A simple module M in \mathcal{C}_q is a highest weight module.*

Proof. — Let M be a simple module in \mathcal{C}_q . Then, there exists a weight (λ, μ) of M such that for all other weight (λ', μ') of M , we have $(\lambda, \mu) \not\leq (\lambda', \mu')$. Such a weight exists because M is finite dimensional. Pick a non-zero $v \in M_{(\lambda, \mu)}$. As, by choice of (λ, μ) , $(\lambda + \alpha_i, \mu + \alpha_i)$ is not a weight of M for any i , we have $E_i v = 0$ for any i . Hence v is a highest weight vector.

Consider now the $\mathcal{D}_q(\mathfrak{g})$ -module of M generated by v . As M is simple and $v \neq 0$, it is the entire module M : M is a highest weight module. \square

Proposition 2.2. — *Any highest weight (λ, μ) of a module in \mathcal{C}_q satisfy $\lambda + \mu \in 2P^+$.*

Proof. — Let M be a module in \mathcal{C}_q and $m \in M$ a highest weight vector of weight (λ, μ) . Let $1 \leq i \leq n$ and consider the family $(F_i^{(k)} m)_{k \in \mathbb{N}}$ of vectors in M . The vector $F_i^{(k)} m$ being of weight $(\lambda, \mu) - k(\alpha_i, \alpha_i)$ and M being finite dimensional, there exists a unique $k \in \mathbb{N}$ such that $F_i^{(k)} m \neq 0$ and $F_i^{(k+1)} m = 0$. As

$$[E_i, F_i^{(k+1)}] = F_i^{(k)} \frac{q_i^{-k} K_i - q_i^k L_i^{-1}}{q_i - q_i^{-1}},$$

we have $0 = [E_i, F_i^{(k+1)}] m = \frac{q_i^{-k+\langle \lambda, \alpha_i^\vee \rangle} - q_i^{k-\langle \mu, \alpha_i^\vee \rangle}}{q_i - q_i^{-1}} F_i^{(k)} m$. Therefore $\langle \lambda + \mu, \alpha_i^\vee \rangle = 2k$. \square

For any $\lambda \in P$, there exists a one-dimensional $\mathcal{D}_q(\mathfrak{g})$ -module with unique (hence highest) weight $(\lambda, -\lambda)$, which we denote by $L(\lambda, -\lambda)$. The elements E_i and F_i necessarily acts by 0 and the action of K_i (resp. L_i) is multiplication by $q^{\langle \lambda, \alpha_i \rangle}$ (resp. $q^{-\langle \lambda, \alpha_i \rangle}$).

We denote by \tilde{P} the subset of $P \times P$ such that the sum of the two components is in $2P$. These are precisely the weights which appear as weights of objects of the category \mathcal{C}_q :

Proposition 2.3. — *For any $(\lambda, \mu) \in P \times P$ such that $\lambda + \mu \in 2P^+$, there exists a simple module in \mathcal{C}_q of highest weight (λ, μ) .*

Proof. — Using the classification of $\mathcal{U}_q(\mathfrak{g})$ -modules (see [CP94, Proposition 10.1.1]), there exists a simple $\mathcal{D}_q(\mathfrak{g})$ -module of highest weight $(\frac{\lambda+\mu}{2}, \frac{\lambda+\mu}{2})$. Tensoring this module with $L(\frac{\lambda-\mu}{2}, -\frac{\lambda-\mu}{2})$, we obtain a simple module in \mathcal{C}_q of highest weight (λ, μ) . \square

Therefore, simple objects in \mathcal{C}_q are classified by $(\lambda, \mu) \in P \times P$ such that $\lambda + \mu \in 2P^+$. We denote \tilde{P}^+ this set of weights and by $L(\lambda, \mu)$ the simple module of highest weight (λ, μ) . Note that we also have a construction of simple modules “à la Verma”.

Proposition 2.4. — *The category \mathcal{C}_q is semisimple.*

Proof. — It suffices to show that there are no extensions $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ of objects in \mathcal{C}_q with M indecomposable and L, N non-trivial. If such an extension exists, for $1 \leq i \leq n$ the action of z_i on M has a unique eigenvalue. Therefore, for any weight (λ, μ) of M , the value $\lambda - \mu$ does not depend on the weight and is in $2P$. We tensorize the exact sequence by the invertible object $X = L(\frac{\mu-\lambda}{2}, -\frac{\mu-\lambda}{2})$ in order to obtain an exact sequence of $\mathcal{U}_q(\mathfrak{g})$ -modules. As the category of finite dimensional $\mathcal{U}_q(\mathfrak{g})$ -modules is semisimple, $M \otimes X \simeq (L \otimes X) \oplus (N \otimes X)$. Therefore $M \simeq L \oplus N$, contrary to our assumption. \square

There exists a faithful P -grading on the category \mathcal{C}_q given by the action of the central elements $(z_i)_{1 \leq i \leq n}$

$$\mathcal{C}_q = \bigoplus_{\nu \in P} \mathcal{C}_{q, \nu},$$

where $\mathcal{C}_{q, \nu}$ is additively generated by simple objects $L(\lambda, \mu)$ with $\lambda - \mu = 2\nu$, i.e. the simple objects on which the central elements $(z_i)_{1 \leq i \leq n}$ act by 2ν . Each component $\mathcal{C}_{q, \nu}$ is equivalent to the category of finite dimensional $\mathcal{U}_q(\mathfrak{g})$ -modules: it is clear for the trivial component $\mathcal{C}_{q, 0}$ and tensoring by $L(\nu, -\nu)$ gives an equivalence between $\mathcal{C}_{q, 0}$ and $\mathcal{C}_{q, \nu}$.

2.2. Character formula. — We now write a character formula, which we obtain easily using Weyl character formula (see [Hum78, 24.3]). We define an action of the Weyl group W on \tilde{P} as follows

$$s_i(\lambda, \mu) = (\lambda, \mu) - \frac{1}{2} \langle \lambda + \mu, \alpha_i^\vee \rangle (\alpha_i, \alpha_i),$$

for any $1 \leq i \leq n$. It is easily shown, by induction on the length of $w \in W$ that

$$w(\lambda, \mu) = \left(w \left(\frac{\lambda + \mu}{2} \right) + \frac{\lambda - \mu}{2}, w \left(\frac{\lambda + \mu}{2} \right) - \frac{\lambda - \mu}{2} \right).$$

We denote by $e^{(\lambda, \mu)} \in \mathbb{Z}[P \times P]$ the element of the group ring of $P \times P$ over \mathbb{Z} corresponding to (λ, μ) . The character of a module M in \mathcal{C}_q is defined to be

$$\chi_M = \sum_{(\lambda, \mu) \in P \times P} \dim(M_{(\lambda, \mu)}) e^{(\lambda, \mu)}.$$

Almost every term of the sum is zero and the support of χ_M is contained in \tilde{P} .

The usual Weyl character formula gives the character of all simple modules in \mathcal{C}_q of the form $L(\lambda, \lambda)$ with $\lambda \in P^+$:

$$\chi(\lambda, \lambda) = \frac{\sum_{w \in W} (-1)^{l(w)} e^{(w(\lambda + \rho), w(\lambda + \rho))}}{\sum_{w \in W} (-1)^{l(w)} e^{(w(\rho), w(\rho))}}.$$

As, for $(\lambda, \mu) \in \tilde{P}$, we have an isomorphism in \mathcal{C}_q

$$L(\lambda, \mu) \simeq L \left(\frac{\lambda + \mu}{2}, \frac{\lambda + \mu}{2} \right) \otimes L \left(\frac{\lambda - \mu}{2}, -\frac{\lambda - \mu}{2} \right),$$

the character of $L(\lambda, \mu)$ is

$$\chi(\lambda, \mu) = \frac{\sum_{w \in W} (-1)^{l(w)} e^{(w(\frac{\lambda + \mu}{2} + \rho), w(\frac{\lambda + \mu}{2} + \rho))}}{\sum_{w \in W} (-1)^{l(w)} e^{(w(\rho), w(\rho))}} e^{(\frac{\lambda - \mu}{2}, -\frac{\lambda - \mu}{2})},$$

which is therefore equal to

$$\chi(\lambda, \mu) = \frac{\sum_{w \in W} (-1)^{l(w)} e^{(w(\lambda + \rho, \mu + \rho))}}{\sum_{w \in W} (-1)^{l(w)} e^{(w(\rho, \rho))}}.$$

We rewrite this formula by introducing the *dot action* $w \bullet (\lambda, \mu) = w(\lambda + \rho, \mu + \rho) - (\rho, \rho)$ which stabilizes $P \times P$:

$$\chi(\lambda, \mu) = \frac{\sum_{w \in W} (-1)^{l(w)} e^{(w \bullet (\lambda, \mu))}}{\sum_{w \in W} (-1)^{l(w)} e^{(w \bullet (0, 0))}}.$$

2.3. Braiding. — Using the quasi- R -matrix of section 1.7, we endow the tensor category \mathcal{C}_q with a braiding. We moreover have to work with an extension of $\mathbb{Q}(q)$. Let L be the smallest integer such that $L \langle \lambda, \mu \rangle \in \mathbb{Z}$ for any $\lambda, \mu \in P$. Let s an indeterminate such that $s^L = q$ and we now work over the field $\mathbb{Q}(s)$.

First, as any element of \mathcal{C}_q has a finite number of weights, we see that for N sufficiently large and any $\mu \in Q$, $\mu \geq 0$ and $\text{ht}(\mu) \geq N$, the element Θ_μ of section 1.7 acts by 0 on any tensor product of two elements of \mathcal{C}_q . Therefore, for any M and M' in \mathcal{C}_q , the action of Θ defines a linear map

$$\Theta_{M, M'}: M \otimes M' \rightarrow M \otimes M'.$$

Note that this map is not $\mathcal{D}_q(\mathfrak{g})$ -linear but satisfies for any $u \in \mathcal{D}_q(\mathfrak{g})$

$$\Theta_{M, M'} \circ \Delta(u) = (\Psi \circ \Delta^{\text{op}})(u) \circ \Theta_{M, M'},$$

as linear endomorphisms of $M \otimes M'$. This follows immediately from Proposition 1.14.

To construct the braiding, we need one more ingredient. For M and M' two objects of \mathcal{C}_q , we introduce a linear map $f_{M,M'}: M \otimes M' \rightarrow M \otimes M'$ defined on weight vectors $m \in M_{\lambda,\mu}$ and $m' \in M'_{\lambda',\mu'}$ by

$$f_{M,M'}(m \otimes m') = q^{\langle \lambda, \mu' \rangle} m \otimes m',$$

where we write $q^r = s^{Lr}$ for any $r \in \frac{1}{L}\mathbb{Z}$.

Lemma 2.5. — *For any $u \in \mathcal{D}_q(\mathfrak{g}) \otimes \mathcal{D}_q(\mathfrak{g})$, we have the following equality as linear endomorphisms of $M \otimes M'$*

$$u \circ f_{M,M'} = f_{M,M'} \circ \Psi(u).$$

Proof. — It suffices to show it on the generators of $\mathcal{D}_q(\mathfrak{g}) \otimes \mathcal{D}_q(\mathfrak{g})$. It is trivial for $K_i \otimes 1$, $L_i \otimes 1$, $1 \otimes K_i$ and $1 \otimes L_i$. We now verify it for $E_i \otimes 1$. On the one hand,

$$(f_{M,M'} \circ (E_i \otimes L_i^{-1}))|_{M_{\lambda,\mu} \otimes M'_{\lambda',\mu'}} = q^{\langle \lambda + \alpha_i, \mu' \rangle - \langle \alpha_i, \mu' \rangle} (E_i \otimes 1)|_{M_{\lambda,\mu} \otimes M'_{\lambda',\mu'}}.$$

On the other hand,

$$((E_i \otimes 1) \circ f_{M,M'})|_{M_{\lambda,\mu} \otimes M'_{\lambda',\mu'}} = q^{\langle \lambda, \mu' \rangle} (E_i \otimes 1)|_{M_{\lambda,\mu} \otimes M'_{\lambda',\mu'}},$$

which concludes the proof for $E_i \otimes 1$. The other cases are similar. \square

Denote by $\tau_{V,W}: V \otimes W \rightarrow W \otimes V$ the twist of vector spaces sending $v \otimes w$ to $w \otimes v$ and define

$$c_{M,M'} = \tau \circ f_{M,M'} \circ \Theta_{M,M'}.$$

It is then a morphism in \mathcal{C}_q between $M \otimes M'$ and $M' \otimes M$. Indeed, for any $u \in \mathcal{D}_q(\mathfrak{g})$ we have

$$\begin{aligned} c_{M,M'} \circ u|_{M \otimes M'} &= \tau \circ f_{M,M'} \circ \Theta_{M,M'} \circ \Delta(u)|_{M \otimes M'} \\ &= \tau \circ f_{M,M'} \circ (\Psi \circ \Delta^{\text{op}})(u)|_{M \otimes M'} \circ \Theta_{M,M'} \\ &= \tau \circ \Delta^{\text{op}}(u)|_{M \otimes M'} \circ f_{M,M'} \circ \Theta_{M,M'} \\ &= \Delta(u)|_{M' \otimes M} \circ c_{M,M'} \\ &= u|_{M' \otimes M} \circ c_{M,M'}. \end{aligned}$$

Proposition 2.6. — *The morphisms c defined above endow \mathcal{C}_q with a braiding: the hexagon axioms [EGNO15, Definition 8.1.1] are satisfied.*

Proof. — We check that $c_{M,M' \otimes M''} = (\text{id}_{M'} \otimes c_{M,M''}) \circ (c_{M,M'} \otimes \text{id}_{M''})$. It follows from the following calculations, where we denote by $\tau_{1,23}: M \otimes (M' \otimes M'') \rightarrow (M' \otimes M'') \otimes M$ the twist, which is equal to $\tau_{23} \circ \tau_{12}$:

$$\begin{aligned} c_{M,M' \otimes M''} &= \tau_{1,23} \circ f_{M,M' \otimes M''} \circ \Theta_{M,M' \otimes M''} \\ &= \tau_{23} \circ \tau_{12} \circ (f_{M,M''})_{13} \circ (f_{M,M'})_{12} \circ (\text{id} \otimes \Delta)(\Theta)|_{M \otimes M' \otimes M''} \\ &= \tau_{23} \circ \tau_{12} \circ (f_{M,M''})_{13} \circ (f_{M,M'})_{12} \circ \Psi_{12}(\Theta_{13})|_{M \otimes M' \otimes M''} \circ (\Theta_{12})|_{M \otimes M' \otimes M''} \\ &= \tau_{23} \circ \tau_{12} \circ (f_{M,M''})_{13} \circ (\Theta_{13})|_{M \otimes M' \otimes M''} \circ (f_{M,M'})_{12}(\Theta_{M,M'} \otimes \text{id}_{M''}) \\ &= \tau_{23} \circ (f_{M,M''})_{23} \circ (\text{id}_{M'} \otimes \Theta_{M,M''}) \circ \tau_{12} \circ (f_{M,M'})_{12}(\Theta_{M,M'} \otimes \text{id}_{M''}) \\ &= (\text{id}_{M'} \otimes c_{M,M''}) \circ (c_{M,M'} \otimes \text{id}_{M''}). \end{aligned}$$

The other hexagon axiom is shown similarly. \square

2.4. Duality and pivotal structure. — The algebra $\mathcal{D}_q(\mathfrak{g})$ being a Hopf algebra, for any $\mathcal{D}_q(\mathfrak{g})$ -module M in \mathcal{C}_q , the space of linear forms has naturally a structure of left (resp. right) dual denoted by M^* (resp. *M), where the action of $u \in \mathcal{D}_q(\mathfrak{g})$ on $\varphi \in M^*$ (resp. $\varphi \in {}^*M$) is given by

$$(u \cdot \varphi)(m) = \varphi(S(u) \cdot m) \quad (\text{resp. } (u \cdot \varphi)(m) = \varphi(S^{-1}(u) \cdot m)).$$

Let $(\lambda, \mu) \in \tilde{P}^+$. The simple module $L(\lambda, \mu)$ is isomorphic to $L\left(\frac{\lambda+\mu}{2}, \frac{\lambda+\mu}{2}\right) \otimes L\left(\frac{\lambda-\mu}{2}, -\frac{\lambda-\mu}{2}\right)$ and

$$L(\lambda, \mu)^* \simeq L\left(\frac{\lambda-\mu}{2}, -\frac{\lambda-\mu}{2}\right)^* \otimes L\left(\frac{\lambda+\mu}{2}, \frac{\lambda+\mu}{2}\right)^*.$$

But the left dual of a simple $\mathcal{U}_q(\mathfrak{g})$ -module $L(\kappa, \kappa)$ is isomorphic to $L(-w_0(\kappa), -w_0(\kappa))$ [Jan96, Proposition 5.16] and the left dual of the invertible module $L(\kappa, -\kappa)$ is clearly isomorphic to $L(-\kappa, \kappa)$. Consequently,

$$L(\lambda, \mu)^* \simeq L\left(\frac{\mu-\lambda}{2}, -\frac{\mu-\lambda}{2}\right) \otimes L\left(-w_0\left(\frac{\lambda+\mu}{2}\right), -w_0\left(\frac{\lambda+\mu}{2}\right)\right) \simeq L(-w_0(\lambda, \mu)).$$

Similarly ${}^*L(\lambda, \mu) \simeq L(-w_0(\lambda, \mu))$, which also follows from the pivotal structure given below.

As for any $u \in \mathcal{D}_q(\mathfrak{g})$ we check that $S^2(u) = K_{2\rho} u K_{2\rho}^{-1}$, we have for any $\lambda \in Q$ a pivotal structure [EGNO15, Definition 4.7.7] given by

$$a_{\lambda, M}: \begin{cases} M & \longrightarrow & M^{**} \\ m & \longmapsto & \varphi \mapsto \varphi((L_\lambda K_{2\rho-\lambda}) \cdot m) \end{cases}.$$

We will consider only the pivotal structure given by $\lambda = 2\rho$, *i.e.* we choose $L_{2\rho}$ as pivotal element. This pivotal structure allows us to compute left and right quantum dimensions of modules [EGNO15, Definition 4.7.1]. It follows from the Weyl character formula that

$$\dim^+(L(\lambda, \mu)) = \frac{\sum_{w \in W} (-1)^{l(w)} q^{\langle 2\rho, (w \bullet (\lambda, \mu))_2 \rangle}}{\sum_{w \in W} (-1)^{l(w)} q^{\langle 2\rho, w \bullet (0) \rangle}}$$

and

$$\dim^-(L(\lambda, \mu)) = \frac{\sum_{w \in W} (-1)^{l(w)} q^{-\langle 2\rho, (w \bullet (\lambda, \mu))_2 \rangle}}{\sum_{w \in W} (-1)^{l(w)} q^{-\langle 2\rho, w \bullet (0) \rangle}},$$

where we denote by $(\lambda', \mu')_i$ the i th component of (λ', μ') for $i \in \{1, 2\}$.

As the actions of $L_{2\rho}$ and $K_{2\rho}$ coincide on any $\mathcal{U}_q(\mathfrak{g})$ -module, the quantum dimensions of $L(\kappa, \kappa)$ viewed as a $\mathcal{D}_q(\mathfrak{g})$ -module coincide with the quantum dimensions of $L(\kappa, \kappa)$ viewed as a $\mathcal{U}_q(\mathfrak{g})$ -module, for any $\kappa \in P^+$.

3. Specialization at a root of unity and tilting modules

The goal to this section is to construct a fusion category using the representations of $\mathcal{D}_q(\mathfrak{g})$. But the category \mathcal{C}_q has an infinite number of simple objects. We adapt the construction of the fusion categories attached to quantum groups at roots of unity to $\mathcal{D}_q(\mathfrak{g})$. Recall the notation $\mathcal{A} = \mathbb{Z}[q, q^{-1}]$.

3.1. Lusztig's integral form. — We introduce an \mathcal{A} -subalgebra of $\mathcal{D}_q(\mathfrak{g})$ which we will use to specialize $\mathcal{D}_q(\mathfrak{g})$ at a root of unity. Consider the following elements of $\mathcal{D}_q(\mathfrak{g})$:

$$\begin{aligned} \begin{bmatrix} K_i; c \\ t \end{bmatrix} &= \prod_{r=1}^t \frac{q_i^{c-r+1} K_i - q_i^{-c+r-1} K_i^{-1}}{q_i^r - q_i^{-r}}, & \begin{bmatrix} L_i; c \\ t \end{bmatrix} &= \prod_{r=1}^t \frac{q_i^{c-r+1} L_i - q_i^{-c+r-1} L_i^{-1}}{q_i^r - q_i^{-r}}, \\ \begin{bmatrix} K_i; c; L_i \\ t \end{bmatrix} &= \prod_{r=1}^t \frac{q_i^{c-r+1} K_i - q_i^{-c+r-1} L_i^{-1}}{q_i^r - q_i^{-r}}, & \begin{bmatrix} z_i; c \\ t \end{bmatrix} &= \prod_{r=1}^t \frac{q_i^{c-r+1} z_i - q_i^{-c+r-1} z_i^{-1}}{q_i^r - q_i^{-r}} \end{aligned}$$

for $1 \leq i \leq n$, $c \in \mathbb{Z}$ and $t \in \mathbb{N}$.

We define $\mathcal{D}_q^{\text{res}}(\mathfrak{g})$ as the \mathcal{A} -subalgebra of $\mathcal{D}_q(\mathfrak{g})$ generated by $E_i^{(r)}$, $F_i^{(r)}$, K_i , L_i , $\begin{bmatrix} K_i; c \\ t \end{bmatrix}$, $\begin{bmatrix} L_i; c \\ t \end{bmatrix}$, $\begin{bmatrix} z_i; c \\ t \end{bmatrix}$ and $\begin{bmatrix} K_i; c; L_i \\ t \end{bmatrix}$, for $1 \leq i \leq n$, $c \in \mathbb{Z}$ and $t \in \mathbb{N}$. The coproduct, the counit and the antipode of $\mathcal{D}_q(\mathfrak{g})$ restrict to $\mathcal{D}_q^{\text{res}}(\mathfrak{g})$ and endow $\mathcal{D}_q^{\text{res}}(\mathfrak{g})$ with a structure of a Hopf algebra.

The quotient of $\mathcal{D}_q^{\text{res}}(\mathfrak{g})$ by the Hopf ideal generated by $z_i - 1$, $\begin{bmatrix} z_i; c \\ t \end{bmatrix} - \begin{bmatrix} c \\ t \end{bmatrix}_{q_i}$, $\begin{bmatrix} K_i; c; L_i \\ t \end{bmatrix} - \begin{bmatrix} L_i; c \\ t \end{bmatrix}$ and $\begin{bmatrix} K_i; c; L_i \\ t \end{bmatrix} - \begin{bmatrix} K_i; c \\ t \end{bmatrix}$, for $1 \leq i \leq n$ is the usual Lusztig's integral form of $\mathcal{U}_q(\mathfrak{g})$ (see [CP94, Section 9.3]).

The elements of the form $\begin{bmatrix} K_i; c; L_i \\ t \end{bmatrix}$ appear naturally in the following identity, proved by induction

$$E_i^{(p)} F_i^{(r)} = \sum_{t=0}^{\min(p,r)} F_i^{(r-t)} \begin{bmatrix} K_i; 2t-r-p; L_i \\ t \end{bmatrix} E_i^{(p-t)},$$

for any $1 \leq i \leq n$, $p, r \in \mathbb{N}$.

The following formulas for the coproduct will be useful.

Proposition 3.1. — *Let $1 \leq i \leq n$ and $t \in \mathbb{N}$. Then*

1. $\Delta \left(\begin{bmatrix} K_i; 0 \\ t \end{bmatrix} \right) = \sum_{r=0}^t \begin{bmatrix} K_i; 0 \\ t-r \end{bmatrix} K_i^{-r} \otimes \begin{bmatrix} K_i; 0 \\ r \end{bmatrix} K_i^{t-r},$
2. $\Delta \left(\begin{bmatrix} L_i; 0 \\ t \end{bmatrix} \right) = \sum_{r=0}^t \begin{bmatrix} L_i; 0 \\ t-r \end{bmatrix} L_i^{-r} \otimes \begin{bmatrix} L_i; 0 \\ r \end{bmatrix} L_i^{t-r},$
3. $\Delta \left(\begin{bmatrix} K_i; 0; L_i \\ t \end{bmatrix} \right) = \sum_{r=0}^t \begin{bmatrix} K_i; 0; L_i \\ t-r \end{bmatrix} L_i^{-r} \otimes \begin{bmatrix} K_i; 0; L_i \\ r \end{bmatrix} K_i^{t-r},$
4. $\Delta \left(\begin{bmatrix} z_i; 0 \\ t \end{bmatrix} \right) = \sum_{r=0}^t \begin{bmatrix} z_i; 0 \\ t-r \end{bmatrix} z_i^{-r} \otimes \begin{bmatrix} z_i; 0 \\ r \end{bmatrix} z_i^{t-r}.$

Proof. — We only prove the first formula, the proof for the other formulas are similar. We proceed by induction on t . For $t = 0$, there is nothing to prove. For $t = 1$, we have:

$$\frac{K_i - K_i^{-1}}{q_i - q_i^{-1}} \otimes K_i + K_i^{-1} \otimes \frac{K_i - K_i^{-1}}{q_i - q_i^{-1}} = \frac{K_i \otimes K_i - K_i^{-1} \otimes K_i^{-1}}{q_i - q_i^{-1}}.$$

Now, suppose that the result is true for $t \in \mathbb{N}$. We have:

$$\begin{aligned}
 \Delta\left(\begin{bmatrix} K_i; 0 \\ t+1 \end{bmatrix}\right) &= \Delta\left(\begin{bmatrix} K_i; 0 \\ t \end{bmatrix}\right) \frac{q_i^{-t} K_i \otimes K_i - q_i^t K_i^{-1} \otimes K_i^{-1}}{q_i^{t+1} - q_i^{-t-1}} \\
 &= \sum_{r=0}^t \left(\begin{bmatrix} K_i; 0 \\ t-r \end{bmatrix} K_i^{-r} \otimes \begin{bmatrix} K_i; 0 \\ r \end{bmatrix} K_i^{t-r} \frac{q_i^{-r} (q_i^{-t+r} K_i - q_i^{t-r} K_i^{-1}) \otimes K_i}{q_i^{t+1} - q_i^{-t-1}} \right) \\
 &\quad + \sum_{r=0}^t \left(\begin{bmatrix} K_i; 0 \\ t-r \end{bmatrix} K_i^{-r} \otimes \begin{bmatrix} K_i; 0 \\ r \end{bmatrix} K_i^{t-r} \frac{q_i^{t-r} K_i^{-1} \otimes (q_i^{-r} K_i - q_i^r K_i^{-1})}{q_i^{t+1} - q_i^{-t-1}} \right) \\
 &= \sum_{r=0}^t \left(\begin{bmatrix} K_i; 0 \\ t-r+1 \end{bmatrix} K_i^{-r} \otimes \begin{bmatrix} K_i; 0 \\ r \end{bmatrix} K_i^{t-r+1} \frac{q_i^{-r} (q_i^{t-r+1} - q_i^{-t+r-1})}{q_i^{t+1} - q_i^{-t-1}} \right) \\
 &\quad + \sum_{r=0}^t \left(\begin{bmatrix} K_i; 0 \\ t-r \end{bmatrix} K_i^{-r-1} \otimes \begin{bmatrix} K_i; 0 \\ r+1 \end{bmatrix} K_i^{t-r} \frac{q_i^{t-r} (q_i^{r+1} - q_i^{-r-1})}{q_i^{t+1} - q_i^{-t-1}} \right).
 \end{aligned}$$

We just change the indes in the second sum and obtain:

$$\begin{aligned}
 \Delta\left(\begin{bmatrix} K_i; 0 \\ t+1 \end{bmatrix}\right) &= \sum_{r=0}^t \left(\begin{bmatrix} K_i; 0 \\ t-r+1 \end{bmatrix} K_i^{-r} \otimes \begin{bmatrix} K_i; 0 \\ r \end{bmatrix} K_i^{t-r+1} \frac{q_i^{-r} (q_i^{t-r+1} - q_i^{-t+r-1})}{q_i^{t+1} - q_i^{-t-1}} \right) \\
 &\quad + \sum_{r=1}^{t+1} \left(\begin{bmatrix} K_i; 0 \\ t-r+1 \end{bmatrix} K_i^{-r} \otimes \begin{bmatrix} K_i; 0 \\ r \end{bmatrix} K_i^{t-r+1} \frac{q_i^{t+1-r} (q_i^r - q_i^{-r})}{q_i^{t+1} - q_i^{-t-1}} \right).
 \end{aligned}$$

Noticing that

$$\frac{q_i^{-r} (q_i^{t-r+1} - q_i^{-t+r-1})}{q_i^{t+1} - q_i^{-t-1}} + \frac{q_i^{t+1-r} (q_i^r - q_i^{-r})}{q_i^{t+1} - q_i^{-t-1}} = \frac{q_i^{t-2r+1} - q_i^{-t-1} + q_i^{t+1} - q_i^{t-2r+1}}{q_i^{t+1} - q_i^{-t-1}} = 1,$$

leads to the expected conclusion. \square

We denote by $\mathcal{D}_q^{\text{res}}(\mathfrak{g})^{>0}$ (resp. $\mathcal{D}_q^{\text{res}}(\mathfrak{g})^0$, resp. $\mathcal{D}_q^{\text{res}}(\mathfrak{g})^{<0}$) the intersection of $\mathcal{D}_q^{\text{res}}(\mathfrak{g})$ with $\mathcal{D}_q(\mathfrak{g})^{>0}$ (resp. $\mathcal{D}_q(\mathfrak{g})^0$, resp. $\mathcal{D}_q(\mathfrak{g})^{<0}$).

It is not hard to show that for any $1 \leq i \leq n$, the automorphism T_i restricts to $\mathcal{D}_q^{\text{res}}(\mathfrak{g})$. As in Section 1.6, we have:

Proposition 3.2 ([Lus90, Theorem 6.7]). — *The elements*

$$\prod_{\alpha \in \Phi^+} E_\alpha^{(n_\alpha)}, \quad n_\alpha \in \mathbb{N}$$

form an \mathcal{A} -basis of $\mathcal{D}_q^{\text{res}}(\mathfrak{g})^{>0}$.

The elements

$$\prod_{\alpha \in \Phi^+} F_\alpha^{(n_\alpha)}, \quad n_\alpha \in \mathbb{N}$$

form an \mathcal{A} -basis of $\mathcal{D}_q^{\text{res}}(\mathfrak{g})^{<0}$.

Note that the quasi- R -matrix at the end of Section 1.7 lies in fact in $\mathcal{D}_q^{\text{res}}(\mathfrak{g})$ and the formula for its inverse shows that it also lies in $\mathcal{D}_q^{\text{res}}(\mathfrak{g})$. We still denote by Θ this element. Moreover, the algebra endomorphism Ψ of $\mathcal{D}_q(\mathfrak{g}) \otimes \mathcal{D}_q(\mathfrak{g})$ restricts to an algebra endomorphism of $\mathcal{D}_q^{\text{res}}(\mathfrak{g}) \otimes \mathcal{D}_q^{\text{res}}(\mathfrak{g})$, which we still denote by Ψ . Therefore, we still have the relations

$$\Theta \Delta(u) = (\Psi \circ \Delta^{\text{op}})(u) \Theta, \quad (\text{id} \otimes \Delta)(\Theta) = \Psi_{12}(\Theta_{13}) \Theta_{12}, \quad \text{and} \quad (\Delta \otimes \text{id})(\Theta) = \Psi_{23}(\Theta_{13}) \Theta_{23}.$$

3.2. Specialization at a root of unity. — Let $\xi \in \mathbb{C}$ be a root of unity of order l . Set $l' = l$ if l is odd and $l' = \frac{l}{2}$ if l is even so that l' is the order of ξ^2 .

$$\begin{aligned} \mathcal{A} &\longrightarrow \mathbb{C}, \\ q &\longmapsto \xi. \end{aligned}$$

The specialization of $\mathcal{D}_q(\mathfrak{g})$ at ξ is by definition the algebra $\mathcal{D}_\xi(\mathfrak{g}) = \mathcal{D}_q^{\text{res}}(\mathfrak{g}) \otimes_{\mathcal{A}} \mathbb{C}$. Introduce the integers l_i as the order of $\xi_i = \xi^{\frac{\langle \alpha_i, \alpha_i \rangle}{2}}$ and set $l'_i = l_i$ if l_i is odd and $l'_i = \frac{l_i}{2}$ if l_i is even so that l'_i is the order of ξ_i^2 .

The algebra $\mathcal{D}_\xi(\mathfrak{g})$ is generated by elements of the form

$$E_i^{(r)}, F_i^{(r)}, K_i^{\pm 1}, L_i^{\pm 1}, \begin{bmatrix} K_i; c \\ t \end{bmatrix}, \begin{bmatrix} L_i; c \\ t \end{bmatrix}, \begin{bmatrix} K_i; c; L_i \\ t \end{bmatrix} \text{ and } \begin{bmatrix} z_i; c \\ t \end{bmatrix},$$

for $1 \leq i \leq n$, $r, t \in \mathbb{N}$ and $c \in \mathbb{Z}$ given by the images in $\mathcal{D}_\xi(\mathfrak{g})$ of the corresponding elements of $\mathcal{D}_q^{\text{res}}(\mathfrak{g})$. There exist some relations in $\mathcal{D}_\xi(\mathfrak{g})$ related to the order of the ξ_i 's as

$$E_i^{l'_i} = 0, F_i^{l'_i} = 0, K_i^{2l'_i} = 1, L_i^{2l'_i} = 1, K_i^{l'_i} = L_i^{l'_i}.$$

We denote by Θ_ξ the specialization of Θ . It is still an invertible element of some completion of $\mathcal{D}_\xi(\mathfrak{g}) \otimes \mathcal{D}_\xi(\mathfrak{g})$, and we have an algebra endomorphism Ψ_ξ which is the specialization of Ψ . They satisfy the usual relations

$$\begin{aligned} \Theta_\xi \Delta(u) &= (\Psi_\xi \circ \Delta^{\text{op}})(u) \Theta_\xi, \quad (\text{id} \otimes \Delta)(\Theta_\xi) = (\Psi_\xi)_{12}((\Theta_\xi)_{13})(\Theta_\xi)_{12}, \\ &\text{and } (\Delta \otimes \text{id})(\Theta_\xi) = (\Psi_\xi)_{23}((\Theta_\xi)_{13})(\Theta_\xi)_{23}. \end{aligned}$$

3.3. Representations at a root of unity. — The representation theory of $\mathcal{U}_\xi(\mathfrak{g})$, as well the one of $\mathcal{D}_\xi(\mathfrak{g})$, is more involved. As explained in [CP94, 11.2.A], defining the weight space of weight (λ, μ) of a $\mathcal{D}_q^{\text{res}}(\mathfrak{g})$ -module M as

$$\{m \in M \mid K_i m = \xi^{\langle \lambda, \alpha_i \rangle} m, L_i m = \xi^{\langle \mu, \alpha_i \rangle} m, \forall 1 \leq i \leq n\}$$

is rather unsatisfactory: one can not distinguish the weight (λ, μ) from the weight $(\lambda + l_i \varpi_i, \mu)$ for example.

The weight space of weight (λ, μ) of M is then defined as

$$M_{\lambda, \mu} = \left\{ m \in M \mid K_i m = \xi_i^{\langle \lambda, \alpha_i^\vee \rangle} m, L_i m = \xi_i^{\langle \mu, \alpha_i^\vee \rangle} m, \begin{bmatrix} K_i; 0 \\ l'_i \end{bmatrix} m = \begin{bmatrix} \langle \lambda, \alpha_i^\vee \rangle \\ l'_i \end{bmatrix}_{\xi_i}, \right. \\ \left. \begin{bmatrix} L_i; 0 \\ l'_i \end{bmatrix} m = \begin{bmatrix} \langle \mu, \alpha_i^\vee \rangle \\ l'_i \end{bmatrix}_{\xi_i}, \forall 1 \leq i \leq n \right\}.$$

Proposition 3.3. — Let (λ, μ) and (λ', μ') be two weights of a $\mathcal{D}_\xi(\mathfrak{g})$ -module M . Then $M_{\lambda, \mu} = M_{\lambda', \mu'}$ if and only if $(\lambda, \mu) = (\lambda', \mu')$.

Proof. — We show that for ζ a root of unity of order d and with $d' = d$ if d is odd, $d' = \frac{d}{2}$ if d is even, and $r \in \mathbb{Z}$, we can recover r knowing only $\begin{bmatrix} r \\ d' \end{bmatrix}_\zeta$ and ζ^r .

Lemma 3.4. — Let $a, b \in \mathbb{N}$ with $a \geq b$. Let $a = a_1 d' + a_0$ and $b = b_1 d' + b_0$ with $0 \leq a_0, b_0 < d'$. Then

$$\begin{bmatrix} a \\ b \end{bmatrix}_\zeta = (-1)^{(d'+1)(a_1+1)b_1} (\zeta^{d'})^{(a_1+1)b_1 + a_1 b_0 + a_0 b_1} \begin{bmatrix} a_1 \\ b_1 \end{bmatrix}_\zeta \begin{bmatrix} a_0 \\ b_0 \end{bmatrix}_\zeta.$$

Proof. — We start with the following equality, valid for any $m \in \mathbb{N}$ and X an indeterminate

$$\prod_{k=0}^{m-1} (1 + \zeta^{2k} X) = \sum_{k=0}^m \zeta^{k(m-1)} \begin{bmatrix} m \\ k \end{bmatrix}_{\zeta} X^k.$$

As $\{\zeta^{-2k} \mid 0 \leq k < d'\}$ is the set of d' th-roots of 1 and

$$\prod_{k=0}^{d'-1} (1 + \zeta^{2k} X) = \zeta^{d'(d'-1)} \prod_{k=0}^{d'-1} (X + \zeta^{-2k})$$

we have

$$\prod_{k=0}^{d'-1} (1 + \zeta^{2k} X) = \zeta^{d'(d'-1)} (X^{d'} - (-1)^{d'}) = \zeta^{d'(d'+1)} (X^{d'} + (-1)^{d'+1}).$$

On the one hand we have

$$\prod_{k=0}^{a-1} (1 + \zeta^{2k} X) = \sum_{k=0}^a \zeta^{k(r-1)} \begin{bmatrix} a \\ k \end{bmatrix}_{\zeta} X^k,$$

and on the other hand

$$\begin{aligned} \prod_{k=0}^{a-1} (1 + \zeta^{2k} X) &= \left(\zeta^{d'(d'+1)} (X^{d'} + (-1)^{d'+1}) \right)^{a_1} \left(\prod_{k=0}^{a_0-1} (1 + \zeta^{2k} X) \right) \\ &= \zeta^{d'(d'+1)a_1} \left(\sum_{k=0}^{a_1} \begin{bmatrix} a_1 \\ k \end{bmatrix} (-1)^{(d'+1)(a_1-k)} X^{kd'} \right) \left(\sum_{k=0}^{a_0} \zeta^{k(a_0-1)} \begin{bmatrix} a_0 \\ k \end{bmatrix}_{\zeta} X^k \right). \end{aligned}$$

Comparing the coefficient of X^b gives us

$$\zeta^{b(a-1)} \begin{bmatrix} a \\ b \end{bmatrix}_{\zeta} = \zeta^{d'(d'+1)a_1 + b_0(a_0-1)} (-1)^{(d'+1)(a_1-b_1)} \begin{bmatrix} a_1 \\ b_1 \end{bmatrix} \begin{bmatrix} a_0 \\ b_0 \end{bmatrix}_{\zeta},$$

which, using the fact that $\zeta^{d'(d'+1)} = (-1)^{d'+1}$, leads to the desired formula. \square

Therefore, for $r \in \mathbb{N}$, we have

$$\begin{bmatrix} d' \\ r \end{bmatrix}_{\zeta} = \begin{cases} r_1 & \text{if } d' = d, \\ (-1)^{r_0+r_1+1} r_1 & \text{if } d' = \frac{d}{2} \text{ and } d' \text{ is odd,} \\ (-1)^{r_0} r_1 & \text{if } d' = \frac{d}{2} \text{ and } d' \text{ is even.} \end{cases}$$

Using $\begin{bmatrix} -r \\ d' \end{bmatrix}_{\zeta} = (-1)^{d'} \begin{bmatrix} r + d' - 1 \\ d' \end{bmatrix}_{\zeta}$, we check that this is still valid for $r \in \mathbb{Z}$.

We write $r = r_1 d' + r_0$ with $0 \leq r_0 < d'$. Let $0 \leq r' < d$ be the unique integer such that $\zeta^r = \zeta^{r'}$.

First, we suppose $d' = d$, hence d' is odd. Lemma 3.4 gives $\begin{bmatrix} r \\ d' \end{bmatrix}_{\zeta} = r_1$ and we have

$$r = \begin{bmatrix} r \\ d' \end{bmatrix}_{\zeta} d' + r'.$$

Now, we suppose that d is even. Let $r'' = r'$ if $0 \leq r' < d'$ and $r'' = r' - d'$ otherwise.

We suppose that $d' = \frac{d}{2}$ is odd. Lemma 3.4 gives $\begin{bmatrix} r \\ d' \end{bmatrix}_{\zeta} = (-1)^{r_0+r_1+1} r_1$. If $r'' = r'$ then

$$r = (-1)^{r''+1} \begin{bmatrix} r \\ d' \end{bmatrix}_{\zeta} + r'', \text{ and if } r'' \neq r' \text{ then } r = (-1)^{r''} \begin{bmatrix} r \\ d' \end{bmatrix}_{\zeta} + r''.$$

Finally, if we suppose that $d' = \frac{d}{2}$ is even, Lemma 3.4 gives $\begin{bmatrix} r \\ d' \end{bmatrix}_\zeta = (-1)^{r_0} r_1$ and $r = (-1)^{r''} \begin{bmatrix} r \\ d' \end{bmatrix}_\zeta + r''$.

Therefore, if for all $1 \leq i \leq n$, $\xi_i^{\langle \lambda, \alpha_i^\vee \rangle} = \xi_i^{\langle \lambda', \alpha_i^\vee \rangle}$ and $\begin{bmatrix} \langle \lambda, \alpha_i^\vee \rangle \\ l'_i \end{bmatrix}_{\xi_i} = \begin{bmatrix} \langle \lambda', \alpha_i^\vee \rangle \\ l'_i \end{bmatrix}_{\xi_i}$, we have $\langle \lambda, \alpha_i^\vee \rangle = \langle \lambda', \alpha_i^\vee \rangle$ for all $1 \leq i \leq n$ and hence $\lambda = \lambda'$. Similarly $\mu = \mu'$. \square

The formula for the coproduct given in Proposition 3.1 shows that the tensor product of two weight vectors of weight (λ, μ) and (λ', μ') is again a weight vector, of weight $(\lambda + \lambda', \mu + \mu')$.

We consider the category \mathcal{C}_ξ of finite dimensional $\mathcal{D}_\xi(\mathfrak{g})$ -modules M such that

$$M = \bigoplus_{(\lambda, \mu) \in P \times P} M_{\lambda, \mu}.$$

As in the generic case we have $E_i^{(r)} \cdot M_{\lambda, \mu} \subseteq M_{\lambda + r\alpha_i, \mu + r\alpha_i}$ and $F_i^{(r)} \cdot M_{\lambda, \mu} \subseteq M_{\lambda - r\alpha_i, \mu - r\alpha_i}$.

Contrary to the generic case, it may happen that $\lambda + \mu \notin 2P$, for example if one of the l_i 's is odd.

One way to construct $\mathcal{D}_\xi(\mathfrak{g})$ -modules is specialization. Let $(\lambda, \mu) \in \tilde{P}^+$. We have defined a simple highest weight $\mathcal{D}_q(\mathfrak{g})$ -module $L(\lambda, \mu)$ and we consider a sub- $\mathcal{D}_q^{\text{res}}(\mathfrak{g})$ -module $L^{\text{res}}(\lambda, \mu)$ generated by a chosen highest weight vector. Define the *Weyl module* by

$$W_\xi(\lambda, \mu) = L^{\text{res}}(\lambda, \mu) \otimes_{\mathcal{A}} \mathbb{C}.$$

This is a highest weight $\mathcal{D}_\xi(\mathfrak{g})$ -module with highest weight (λ, μ) , but it is not always a simple module. As specialization does not change the weight spaces, it is clear that the character of $W_\xi(\lambda, \mu)$ is still given by the Weyl character formula given in section 2.2. The Weyl module is a quotient of the Verma module and it has simple head which we denote $L_\xi(\lambda, \mu)$.

Similarly to the case of a generic parameter q , the choice of an L -th root $\xi^{1/L}$ of ξ turns the category of representations into a braided category, as in Section 2.3

3.4. Tilting modules. — As for $\mathcal{U}_\xi(\mathfrak{g})$, we consider the tilting modules, which have been studied for quantum groups at roots of unity by Andersen in [And92].

Definition 3.5. — *A $\mathcal{D}_\xi(\mathfrak{g})$ -module M is tilting if both M and M^* have a filtration with successive quotients being Weyl modules.*

We refer to [Saw06] for more details in the case of $\mathcal{U}_\xi(\mathfrak{g})$. We remark that weights (λ, μ) of Weyl modules always satisfy $\lambda + \mu \in 2P$; hence only these weights can appear in a tilting module.

Proposition 3.6. — *For any $(\lambda, \mu) \in \tilde{P}^+$, there exists an indecomposable tilting module $T(\lambda, \mu)$ such that $T(\lambda, \mu)_{\lambda', \mu'} = 0$ unless $(\lambda', \mu') \leq (\lambda, \mu)$ and $T(\lambda, \mu)_{\lambda, \mu}$ is of dimension one.*

Moreover, any indecomposable tilting module is isomorphic to some $T(\lambda, \mu)$.

Proof. — The existence of such tilting modules follows easily from the existence for $\mathcal{U}_\xi(\mathfrak{g})$. Indeed, there exists an indecomposable tilting module for $\mathcal{U}_\xi(\mathfrak{g})$ with a maximal vector of weight $\frac{\lambda + \mu}{2}$. Tensoring it with $L_\xi\left(\frac{\lambda - \mu}{2}, -\frac{\lambda - \mu}{2}\right)$ gives us an indecomposable tilting module with a maximal vector of weight (λ, μ) .

Let T be an indecomposable tilting $\mathcal{U}_\xi(\mathfrak{g})$ -module. Using the action of the central elements $z_i, \begin{bmatrix} z_i & 0 \\ l'_i & \end{bmatrix}$ for $1 \leq i \leq n$, one can show that $\lambda - \mu$ does not depend on the weight (λ, μ) of T . Therefore, tensoring T by $L_\xi\left(\frac{\mu-\lambda}{2}, -\frac{\mu-\lambda}{2}\right)$ gives an indecomposable tilting module for $\mathcal{U}_\xi(\mathfrak{g})$, which is isomorphic to $T(\kappa, \kappa)$ for some $\kappa \in P^+$. \square

As a direct summand of a tilting module is again a tilting module, every tilting module is a direct sum of indecomposable tilting modules. Using the fact that the tensor product of two tilting modules for $\mathcal{U}_\xi(\mathfrak{g})$ is again a tilting module, we easily show that the tensor product of two tilting modules for $\mathcal{D}_\xi(\mathfrak{g})$ is again a tilting module.

The full subcategory of \mathcal{C}_ξ with objects the tilting modules is far from being a fusion category: it is not abelian, nor semisimple. We want to semisimplify this category; hence we must understand which indecomposable tilting modules are of non-zero quantum dimension. As any indecomposable tilting module is isomorphic to some $T(\lambda, \mu) \simeq T\left(\frac{\lambda+\mu}{2}, \frac{\lambda+\mu}{2}\right) \otimes L_\xi\left(\frac{\lambda-\mu}{2}, -\frac{\lambda-\mu}{2}\right)$, we can use the results for $\mathcal{U}_\xi(\mathfrak{g})$. The next theorem follows immediately from [Saw06, Theorem 2]. Let $D = \max_{\alpha, \beta \in \Phi} \frac{\langle \alpha, \alpha \rangle}{\langle \beta, \beta \rangle}$. We have $D = 1$ in type A , D or E , $D = 2$ in type B, C or F_4 and $D = 3$ in type G_2 . Let θ_0 be the highest root of Φ if $D \mid l'$ and be the highest short root of Φ otherwise. Let C be the following set of dominant weights

$$C = \{\lambda \in P^+ \mid \langle \lambda + \rho, \theta_0 \rangle < l'\}.$$

Finally, denote by h the Coxeter number of \mathfrak{g} and by h^\vee the dual Coxeter number of \mathfrak{g} .

Theorem 3.7. — *We suppose that $l' \geq Dh^\vee$ if $D \mid l'$ and $l' > h$ otherwise. Then $T(\lambda, \mu)$ is of non-zero positive and negative quantum dimension if and only if $\frac{\lambda+\mu}{2} \in C$. Moreover, $T(\lambda, \mu) \simeq W_\xi(\lambda, \mu) \simeq L_\xi(\lambda, \mu)$.*

3.5. The semisimple category of non-negligible tilting modules. — From now on, we keep the assumption on l of Theorem 3.7. We construct a fusion category using a semisimplification of pivotal categories. As the category of tilting modules is neither abelian, nor spherical, we use the version of semisimplification of pivotal Karoubian categories as in [EO18, 2.3]. The hypothesis of [EO18, Theorem 2.6] are satisfied since

1. the category of tilting modules for the algebra $\mathcal{U}_\xi(\mathfrak{g})$ is a subcategory of an abelian category,
2. up to an invertible element, the positive and negative quantum dimensions of an indecomposable tilting module for the algebra $\mathcal{U}_\xi(\mathfrak{g})$ are equal.

We then denote by \mathcal{T}_ξ the semisimplification of the category of tilting $\mathcal{D}_\xi(\mathfrak{g})$ -modules. This is a semisimple braided pivotal tensor category and the simple objects are the indecomposable tilting modules $T(\lambda, \mu)$ such that $\frac{\lambda+\mu}{2} \in C$. This category admits a faithful P -grading

$$\mathcal{T}_\xi = \bigoplus_{\nu \in P} \mathcal{T}_{\xi, \nu}$$

where $\mathcal{T}_{\xi, \nu}$ is additively generated by simple objects $L_\xi(\lambda, \mu)$ with $\lambda - \mu = 2\nu$. As in the case of a generic parameter q , each component $\mathcal{T}_{\xi, \nu}$ is equivalent to the category of tilting modules for $\mathcal{U}_\xi(\mathfrak{g})$.

We now compute explicitly the S -matrix and the twist for \mathcal{T}_ξ . The S -matrix is the matrix indexed by $\text{Irr}(\mathcal{T}_\xi)$ given by

$$S_{(\lambda, \mu), (\lambda', \mu')} = \text{Tr}_{L_\xi(\lambda, \mu) \otimes L_\xi(\lambda', \mu')}^+ (c_{L_\xi(\lambda', \mu'), L_\xi(\lambda, \mu)} \circ c_{L_\xi(\lambda, \mu), L_\xi(\lambda', \mu')}).$$

The twist θ and the pivotal structure a are related using the Drinfeld morphism u (see [EGNO15, 8.35]): $a = u\theta$. The latter is given by the composition

$$X \xrightarrow{\text{id}_X \otimes \text{coev}_{X^*}} X \otimes X^* \otimes X^{**} \xrightarrow{c_{X,X^*} \otimes \text{id}_{X^{**}}} X^* \otimes X \otimes X^{**} \xrightarrow{\text{ev}_X \otimes \text{id}_{X^{**}}} X^{**}.$$

We recall the expression of the quasi- R -matrix Θ_ξ of $\mathcal{D}_\xi(\mathfrak{g})$

$$\Theta_\xi = \prod_{\alpha \in \Phi^+} \left(\sum_{n=0}^{+\infty} q_\alpha^{\frac{n(n-1)}{2}} [n]_{q_\alpha}! (q_\alpha - q_\alpha^{-1})^n E_\alpha^{(n)} \otimes F_\alpha^{(n)} \right).$$

Recall also that we have chosen the pivotal structure given by the element $L_{2\rho}$ so that positive and negative quantum traces of any linear map $f: M \rightarrow M$ are given by

$$\text{Tr}_M^+(f) = \text{Tr}(L_{2\rho} f | M) \quad \text{and} \quad \text{Tr}_M^-(f) = \text{Tr}(f L_{2\rho}^{-1} | M).$$

Proposition 3.8. — *Let (λ, μ) and (λ', μ') in \tilde{P}^+ . Then the S -matrix of \mathcal{T}_ξ is given by*

$$S_{(\lambda, \mu), (\lambda', \mu')} = \frac{\sum_{w \in W} (-1)^{l(w)} \xi^{\langle 2\rho + \lambda, (w \bullet (\lambda', \mu'))_2 \rangle + \langle \mu, (w \bullet (\lambda', \mu'))_1 + 2\rho \rangle}}{\sum_{w \in W} (-1)^{l(w)} \xi^{\langle 2\rho, w \bullet 0 \rangle}},$$

and the twist is

$$\theta_{L_\xi(\lambda, \mu)} = \xi^{\langle \lambda + 2\rho, \mu \rangle} \text{id}_{L_\xi(\lambda, \mu)}.$$

Proof. — Let $L_\xi(\lambda, \mu)$ be a simple object in \mathcal{T}_ξ . For any other simple object M in \mathcal{T}_ξ the map

$$(\text{id}_{L_\xi(\lambda, \mu)} \otimes \text{Tr}_M^+)(c_{M, L_\xi(\lambda, \mu)} \circ c_{L_\xi(\lambda, \mu), M})$$

is an endomorphism of the simple object $L_\xi(\lambda, \mu)$ hence is a scalar. We compute it on the highest weight vector $v_{\lambda, \mu}$ of $L_\xi(\lambda, \mu)$. Let $m \in M$ be a vector of weight (λ', μ') . We have

$$c_{M, L_\xi(\lambda, \mu)} \circ c_{L_\xi(\lambda, \mu), M}(v_{\lambda, \mu} \otimes m) = \xi^{\langle \lambda, \mu' \rangle} c_{M, L_\xi(\lambda, \mu)}(m \otimes v_{\lambda, \mu})$$

as $v_{\lambda, \mu}$ is a vector of highest weight. Now, we take the partial trace of this expression and are interested only in the component on $v_{\lambda, \mu}$. As $F_\alpha^{(n)} v_{\lambda, \mu}$ is not of weight (λ, μ) for $n > 0$, this shows that only the term $1 \otimes 1$ in Θ_ξ contributes on this component. Therefore

$$(\text{id}_{L_\xi(\lambda, \mu)} \otimes \text{Tr}_M^+)(c_{M, L_\xi(\lambda, \mu)} \circ c_{L_\xi(\lambda, \mu), M}) = \text{Tr}_M^+(\varphi_{\lambda, \mu}^M) \text{id}_{L_\xi(\lambda, \mu)}$$

where $\varphi_{\lambda, \mu}^M(m) = \xi^{\langle \lambda, \mu' \rangle + \langle \lambda', \mu \rangle} m$ for m a vector of weight (λ', μ') .

Finally, the S -matrix is given by

$$S_{(\lambda, \mu), (\lambda', \mu')} = \dim^+(L_\xi(\lambda, \mu)) \text{Tr}^+\left(\varphi_{\lambda, \mu}^{L_\xi(\lambda', \mu')}\right).$$

Using the Weyl character formula, we have

$$S_{(\lambda, \mu), (\lambda', \mu')} = \frac{\sum_{w \in W} (-1)^{l(w)} \xi^{\langle 2\rho, (w \bullet (\lambda, \mu))_2 \rangle}}{\sum_{w \in W} (-1)^{l(w)} \xi^{\langle 2\rho, w \bullet 0 \rangle}} \frac{\sum_{w \in W} (-1)^{l(w)} \xi^{\langle 2\rho + \lambda, (w \bullet (\lambda', \mu'))_2 \rangle + \langle \mu, (w \bullet (\lambda', \mu'))_1 \rangle}}{\sum_{w \in W} (-1)^{l(w)} \xi^{\langle 2\rho + \lambda + \mu, w \bullet 0 \rangle}}.$$

Note that the assumption on l ensures that the denominators are non-zero. As we have $\langle 2\rho + \lambda + \mu, w \bullet 0 \rangle = \langle 2\rho, (w^{-1} \bullet (\lambda, \mu))_2 \rangle - \langle \mu, 2\rho \rangle$ the formula for the S -matrix becomes

$$S_{(\lambda, \mu), (\lambda', \mu')} = \frac{\sum_{w \in W} (-1)^{l(w)} \xi^{\langle 2\rho + \lambda, (w \bullet (\lambda', \mu'))_2 \rangle + \langle \mu, (w \bullet (\lambda', \mu'))_1 + 2\rho \rangle}}{\sum_{w \in W} (-1)^{l(w)} \xi^{\langle 2\rho, w \bullet 0 \rangle}}.$$

We now turn to the twist. Unrolling the definition of the Drinfeld morphism, one can show that for $v_{\lambda, \mu}$ a highest weight vector of $L_\xi(\lambda, \mu)$

$$u_{L_\xi(\lambda, \mu)}(v_{\lambda, \mu}) = (\varphi \in L_\xi(\lambda, \mu)^* \mapsto \xi^{-\langle \lambda, \mu \rangle} \varphi(v_{\lambda, \mu})),$$

whereas

$$a_{L_\xi(\lambda, \mu)}(v_{\lambda, \mu}) = (\varphi \in L_\xi(\lambda, \mu)^* \mapsto \xi^{(2\rho, \mu)} \varphi(v_{\lambda, \mu})).$$

Therefore we have $\theta_{L_\xi(\lambda, \mu)} = \xi^{(\lambda+2\rho, \mu)} \text{id}_{L_\xi(\lambda, \mu)}$. \square

3.6. A partial modularization. — Contrary to the category of tilting modules for $\mathcal{U}_\xi(\mathfrak{g})$, \mathcal{T}_ξ has an infinite number of non-isomorphic simple objects. Following M\"uger [M\"ug00], we first find some objects in the symmetric center of \mathcal{T}_ξ , and we add isomorphism between these objects and the unit object.

Proposition 3.9. — *Let $v \in (l'Q^\vee) \cap P$. Then the invertible simple object $L_\xi(v, -v)$ lies in the symmetric center of \mathcal{T}_ξ . Its positive and negative quantum dimension is 1 and its twist is 1 or -1 .*

Proof. — For any simple object $L_\xi(\lambda, \mu)$ in \mathcal{T}_ξ , we compute $c_{L_\xi(\lambda, \mu), L_\xi(v, -v)} \circ c_{L_\xi(v, -v), L_\xi(\lambda, \mu)}$. As $L_\xi(v, -v) \otimes L_\xi(\lambda, \mu)$ is a simple object, the double braiding is the multiplication by a scalar. By a computation on a highest weight vector, the double braiding is the multiplication by $\xi^{(v, \mu - \lambda)}$. But as v is in $l'Q^\vee$ and $\mu - \lambda \in 2P$, we have $\langle v, \mu - \lambda \rangle \in 2l'\mathbb{Z}$, so that the double braiding is the identity.

The positive quantum dimension of $L_\xi(v, -v)$ is given by the action of $L_{2\rho}$, so is equal to $\xi^{-(2\rho, v)} = 1$ because $\rho \in P$.

The twist is given by $\xi^{(v+2\rho, -v)}$ which is obviously equal to ± 1 since $v \in l'Q^\vee$. \square

Let \mathcal{S} be the tensor subcategory of \mathcal{T}_ξ generated by the $L_\xi(v, -v)$ (which we will denote by $I(v)$), with $v \in (l'Q^\vee) \cap P$ and of twist 1. We recall the construction of the category $\mathcal{T}_\xi \rtimes \mathcal{S}$ of [M\"ug00, Definition 3.12], which is simpler in our case since the objects in \mathcal{S} are all of dimension 1. We choose for any $v, v' \in (l'Q^\vee) \cap P$ an isomorphism $\varphi_{v, v'}$ in the one dimensional space $\text{Hom}_{\mathcal{S}}(I(v) \otimes I(v'), I(v+v'))$ such that the following diagram commutes

$$\begin{array}{ccc} I(v) \otimes I(v') \otimes I(v'') & \xrightarrow{\varphi_{v, v'} \otimes \text{id}} & I(v+v') \otimes I(v'') \\ \downarrow \text{id} \otimes \varphi_{v', v''} & & \downarrow \varphi_{v+v', v''} \\ I(v) \otimes I(v'+v'') & \xrightarrow{\varphi_{v, v'+v''}} & I(v+v'+v'') \end{array} .$$

To do so, choose in any $I(v)$ a non-zero vector v_v . Then $\varphi_{v, v'}$ sends $v_v \otimes v_{v'}$ to $v_{v+v'}$.

We consider the category $\mathcal{T}_\xi \rtimes_0 \mathcal{S}$ with the same objects as \mathcal{T}_ξ and with space of morphisms between two objects X and Y

$$\text{Hom}_{\mathcal{T}_\xi \rtimes_0 \mathcal{S}}(X, Y) = \bigoplus_{v \in P \cap Q^\vee} \text{Hom}_{\mathcal{T}_\xi}(X, I(v) \otimes Y).$$

The composition of $f \in \text{Hom}_{\mathcal{T}_\xi}(X, I(v) \otimes Y)$ and of $g \in \text{Hom}(Y, I(v') \otimes Z)$ in $\mathcal{T}_\xi \rtimes \mathcal{S}$ is given by

$$X \xrightarrow{f} I(v) \otimes Y \xrightarrow{\text{id} \otimes g} I(v) \otimes I(v') \otimes Z \xrightarrow{\varphi_{v, v'} \otimes \text{id}} I(v+v') \otimes Z .$$

Due to the compatibility of the maps φ , it is easy to check that this defines an associative composition.

This category has tensor products: on objects the tensor product is the same as the one in \mathcal{T}_ξ and if $f \in \text{Hom}_{\mathcal{T}_\xi}(X, I(v) \otimes Y)$ and $g \in \text{Hom}_{\mathcal{T}_\xi}(X', I(v') \otimes Y')$, their tensor product is

defined as the composition

$$\begin{aligned} X \otimes X' &\xrightarrow{f \otimes g} I(\nu) \otimes Y \otimes I(\nu') \otimes Y' \xrightarrow{\text{id} \otimes c_{Y, I(\nu')} \otimes \text{id}} I(\nu) \otimes I(\nu') \otimes Y \otimes Y' \\ &\xrightarrow{\varphi_{\nu, \nu'} \otimes \text{id}} I(\nu + \nu') \otimes Y \otimes Y'. \end{aligned}$$

Again, the compatibility of φ ensures that this tensor product endows $\mathcal{T}_\xi \rtimes_0 \mathcal{S}$ with a structure of a semisimple tensor category, see [Müg00, Section 3.2] for further details.

The duality on \mathcal{T}_ξ extends to a duality on $\mathcal{T}_\xi \rtimes_0 \mathcal{S}$. One may check that the pivotal structure in \mathcal{T}_ξ induces a pivotal structure on $\mathcal{T}_\xi \rtimes_0 \mathcal{S}$. For this, it is crucial that the objects in \mathcal{S} are of twist 1.

Finally, the category $\mathcal{T}_\xi \rtimes_0 \mathcal{S}$ is braided because \mathcal{S} is a subcategory of the symmetric center of \mathcal{T}_ξ (see [Müg00, Lemma 3.10]).

Now, in the general case, it can happen that the category constructed above is not idempotent complete. This happens exactly when tensoring by a non-trivial simple object of \mathcal{S} has a fixed point on the set of simple objects of \mathcal{T}_ξ . But the object $I(\nu)$ is in the component $\mathcal{T}_{\xi, \nu}$ of the grading, thus tensoring by this object has no fixed points on the set of simples, provided that $\nu \neq 0$. Therefore the idempotent completion $\mathcal{T}_\xi \rtimes \mathcal{S}$ of $\mathcal{T}_\xi \rtimes_0 \mathcal{S}$ is $\mathcal{T}_\xi \rtimes_0 \mathcal{S}$ itself.

Proposition 3.10. — *The category $\mathcal{T}_\xi \rtimes \mathcal{S}$ is finite in the sense of [EGNO15, Definition 1.8.6].*

Proof. — Indeed, denoting by G the quotient of P by $\{\nu \in (l'Q^\vee) \cap P \mid \theta_{I(\nu)} = 1\}$, $\mathcal{T}_\xi \rtimes \mathcal{S}$ is a G -graded category, with each homogeneous component equivalent to the category of tilting modules for $\mathcal{W}_\xi(\mathfrak{g})$

$$\mathcal{T}_\xi \rtimes \mathcal{S} = \bigoplus_{\nu \in G} (\mathcal{T}_\xi \rtimes \mathcal{S})_\nu.$$

Therefore, $\mathcal{T}_\xi \rtimes \mathcal{S}$ has $|C||G|$ simple objects. Note that G is indeed finite since $P/((l'Q^\vee) \cap P)$ surjects on G . \square

3.7. An integral subcategory. — We consider the full subcategory of \mathcal{T}_ξ additively generated by the $L_\xi(\lambda, \mu)$ with $\lambda + \mu \in 2C$ and $\mu \in Q$. This subcategory is stable by tensor product: we can easily see this at the level of $\mathcal{D}_\xi(\mathfrak{g})$ -modules. We denote this category by $\mathbb{Z}(\mathcal{T}_\xi)$. In the case of the tilting category for $\mathcal{W}_\xi(\mathfrak{sl}_{n+1})$, Masbaum and Wenzl [MW98] show that an analogue of this subcategory is a modular category provided that l is even and l' and $n+1$ are relatively prime.

As for the category \mathcal{T}_ξ , there is an infinite number of simple objects. But the objects of the form $I(\nu)$ with $\nu \in (l'Q^\vee) \cap Q$ are in the symmetric center of $\mathbb{Z}(\mathcal{T}_\xi)$.

Lemma 3.11. — *The twist of $I(\nu)$ is 1 for any $\nu \in (l'Q^\vee) \cap Q$.*

Proof. — We want to show that for any $\nu \in (l'Q^\vee) \cap Q$, we have $\langle \nu, \nu \rangle \equiv 0 \pmod{l}$. For any value of the integer l , we have $\langle \nu, \nu \rangle \in l'^2\mathbb{Z}$. Therefore we may, and will, suppose that $l = 2l'$ with l' odd.

Suppose first that \mathfrak{g} is not of type G_2 . Then, $(l'Q^\vee) \cap Q = l'Q$ and we easily conclude because for any $\lambda \in Q$, $\langle \lambda, \lambda \rangle$ is even (recall that $\langle \cdot, \cdot \rangle$ is normalized such that $\langle \alpha, \alpha \rangle = 2$ for short simple roots).

Now suppose that \mathfrak{g} is of type G_2 . If moreover l' is not a multiple of 3, we have $(l'Q^\vee) \cap Q = l'Q$ and the conclusion follows. We finally suppose that $l = 2l'$ with l' a multiple of

3. Let α be the short simple root and β be the long simple root, so that $\alpha^\vee = \alpha$ and $\beta^\vee = \frac{\beta}{3}$. We therefore have $(l'Q^\vee) \cap Q = l'Q^\vee$ and for any $\nu = l' \nu_\alpha \alpha + \frac{l' \nu_\beta}{3} \beta \in l'Q^\vee$, we have

$$\langle \nu, \nu \rangle = l'^2 \left(2\nu_\alpha^2 - 2\nu_\alpha \nu_\beta + \frac{2}{3} \nu_\beta^2 \right) = 2l' \frac{l'}{3} (3\nu_\alpha^2 - 3\nu_\alpha \nu_\beta + \nu_\beta^2).$$

Hence $\langle \nu, \nu \rangle \equiv 0 \pmod{l}$ since l' is a multiple of 3. \square

Hence, we have shown that for any $\nu \in (l'Q^\vee) \cap Q$, $I(\nu)$ is of quantum dimension 1 and of twist 1. We again construct the category $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ and we obtain a $P/((l'Q^\vee) \cap Q)$ -graded category

$$\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S} = \bigoplus_{\nu \in P/((l'Q^\vee) \cap Q)} \mathbb{Z}(\mathcal{T}_\xi)_{\nu}.$$

Note that we do not have an equivalence of categories between $\mathbb{Z}(\mathcal{T}_\xi)_\nu$ and $\mathbb{Z}(\mathcal{T}_\xi)_0$ if $\nu \notin Q$. But we have an equivalence between $\mathbb{Z}(\mathcal{T}_\xi)_\nu$ and $\mathbb{Z}(\mathcal{T}_\xi)_{\nu'}$ if $\nu = \nu'$ in P/Q .

Proposition 3.12. — *The category $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ is finite and has $|C||Q/((l'Q^\vee) \cap Q)|$ simple objects.*

Proof. — If we choose some representatives ν_1, \dots, ν_k of P/Q in $P/((l'Q^\vee) \cap Q)$, we have $|C|$ simple objects in $\bigoplus_{i=1}^k \mathbb{Z}(\mathcal{T}_\xi)_{\nu_i}$. Therefore $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ has $|C||Q/((l'Q^\vee) \cap Q)|$ simple objects. \square

We end this part with some notation. Denote by \tilde{C} the following set of weights

$$\tilde{C} = \{(\lambda, \mu) \in P \times Q \mid \lambda + \mu \in 2C\}.$$

This set parametrizes the simple objects of $\mathbb{Z}(\mathcal{T}_\xi)$ and the group $l'Q^\vee \cap Q$ acts on it by

$$\nu \cdot (\lambda, \mu) = (\lambda + \nu, \mu - \nu)$$

for $(\lambda, \mu) \in \tilde{C}$ and $\nu \in (l'Q^\vee) \cap Q$. The set $\overline{C} = \tilde{C}/((l'Q^\vee) \cap Q)$ parametrizes the simple objects of $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$.

3.8. Non-degeneracy of $\mathcal{T}_\xi \rtimes \mathcal{S}$ and degeneracy of $\mathbb{Z}(\mathcal{T})_\xi \rtimes \mathcal{S}$. — In this subsection, we suppose that $l = 2Dd$ for $d \geq h^\vee$. Then $l' = Dd$ and $DdQ^\vee \subseteq Q$ and by Lemma 3.11 for any $\nu \in DdQ^\vee$, we have $\theta_{I(\nu)} = 1$.

Proposition 3.13. — *If moreover $\xi = \exp(\frac{i\pi}{d})$, the category $\mathcal{T}_\xi \rtimes \mathcal{S}$ is non-degenerate: if a simple object X is such that for all object Y in \mathcal{C} we have*

$$c_{X,Y} \circ c_{Y,X} = \text{id}_{Y \otimes X}$$

then $X \simeq \mathbf{1}$.

Proof. — Using the proof of [BK01, Theorem 3.3.20], we show that the square of the S-matrix of $\mathcal{T}_\xi \rtimes \mathcal{S}$ is invertible. As for any $w \in W$ and $(\lambda, \mu) \in \tilde{P}$ we have

$$w \bullet (\lambda, \mu) = \left(w \bullet \frac{\lambda + \mu}{2} + \frac{\lambda - \mu}{2}, w \bullet \frac{\lambda + \mu}{2} - \frac{\lambda - \mu}{2} \right),$$

we can rewrite the S-matrix of $\mathcal{T}_\xi \rtimes \mathcal{S}$ as

$$S_{(\lambda, \mu), (\lambda', \mu')} = \xi^{(2\rho, \mu - \eta + \mu' - \eta') - 2(\mu - \eta, \mu' - \eta')} \tilde{s}_{\eta, \eta'}$$

where $\lambda + \mu = 2\eta$, $\lambda' + \mu' = 2\eta'$ and \tilde{s} is the S-matrix of the modular category of tilting modules for $\mathcal{U}_\xi(\mathfrak{g})$ (see proof of [BK01, Theorem 3.3.20]). The simple objects of $\mathcal{T}_\xi \rtimes \mathcal{S}$

are indexed by $\{(\lambda, \mu) \in P \times P \mid \lambda + \mu \in 2C\} / DdQ^\vee$ which is in bijection with $C \times P / DdQ^\vee$ sending (λ, μ) to (η, μ) . Therefore

$$\begin{aligned}
(S^2)_{(\lambda, \mu), (\lambda'', \mu'')} &= \xi^{(2\rho, \mu - \eta + \mu'' - \eta'')} \sum_{\eta' \in C} \tilde{s}_{\eta, \eta'} \tilde{s}_{\eta', \eta''} \xi^{-2\langle \eta', 2\rho + \eta - \mu + \eta'' - \mu'' \rangle} \underbrace{\sum_{\mu' \in P / DdQ^\vee} \xi^{2\langle \mu', 2\rho + \eta - \mu + \eta'' - \mu'' \rangle}}_{= |P / DdQ^\vee| \delta_{2\rho + \eta - \mu + \eta'' - \mu'' \in DdQ^\vee}} \\
&= |P / DdQ^\vee| \delta_{2\rho + \eta - \mu + \eta'' - \mu'' \in DdQ^\vee} \xi^{(2\rho, 2\rho)} \sum_{\eta' \in C} \tilde{s}_{\eta, \eta'} \tilde{s}_{\eta', \eta''} \\
&= \delta_{2\rho + \eta - \mu + \eta'' - \mu'' \in DdQ^\vee} \delta_{\eta'', -w_0(\eta)} \kappa \\
&= \delta_{\eta'', -w_0(\eta)} \delta_{\mu'' \in 2\rho + \eta - \mu - w_0(\eta) + DdQ^\vee} \kappa,
\end{aligned}$$

where κ is a non-zero constant. As $\mu \in P / DdQ^\vee$, the matrix S^2 is, up to a non-zero constant, the invertible permutation matrix $(\delta_{(\lambda'', \mu''), (-2\rho, 2\rho) - w_0(\lambda, \mu)})_{(\lambda, \mu), (\lambda'', \mu'')}$. \square

The square of the S -matrix is not given by the duality: this is due to the fact that $\mathcal{T}_\xi \rtimes \mathcal{S}$ is not spherical, but only pivotal.

Now we turn to the category $\mathbb{Z}(\mathcal{T})_\xi \rtimes \mathcal{S}$ which is degenerate in general.

Theorem 3.14. — *The symmetric center of $\mathbb{Z}(\mathcal{T})_\xi \rtimes \mathcal{S}$ contains $|P/Q|$ simple objects. Therefore, the category $\mathbb{Z}(\mathcal{T})_\xi \rtimes \mathcal{S}$ is non-degenerate if and only if \mathfrak{g} is of type E_8, F_4 or G_2 .*

Proof. — Using [EGNO15, Lemma 8.20.9], the simple object $L_\xi(\lambda, \mu)$ is in the symmetric center of $\mathbb{Z}(\mathcal{T})_\xi \rtimes \mathcal{S}$ if and only if

$$(h_{\lambda, \mu}, h_{0,0})_{\mathbb{Z}(\mathcal{T})_\xi \rtimes \mathcal{S}} = \sum_{(\lambda', \mu') \in \bar{C}} \frac{\dim^-(L_\xi(\lambda', \mu'))}{\dim^+(L_\xi(\lambda', \mu'))} S_{(\lambda, \mu), (\lambda', \mu')} S_{(\lambda', \mu'), (0,0)} \neq 0.$$

As in the proof of Proposition 3.13, we let $2\eta = \lambda + \mu$ and $2\eta' = \lambda' + \mu'$, which are elements in C . We then have

$$\begin{aligned}
(h_{\lambda, \mu}, h_{0,0})_{\mathbb{Z}(\mathcal{T})_\xi \rtimes \mathcal{S}} &= \sum_{\eta' \in C} \sum_{\mu' \in Q / DdQ^\vee} \xi^{2\langle 2\rho, \eta' - \mu' \rangle} \xi^{(2\rho, \mu - \eta + \mu' - \eta') - 2\langle \mu - \eta, \mu' - \eta' \rangle} \tilde{s}_{\eta, \eta'} \xi^{(2\rho, \mu' - \eta')} \tilde{s}_{\eta', 0} \\
&= \xi^{(2\rho, \mu - \eta)} \sum_{\eta' \in C} \xi^{-2\langle \eta', \eta - \mu \rangle} \tilde{s}_{\eta, \eta'} \tilde{s}_{\eta', 0} \underbrace{\sum_{\mu' \in Q / DdQ^\vee} \xi^{2\langle \mu', \eta - \mu \rangle}}_{|Q / DdQ^\vee| \delta_{\mu - \eta \in DdP^\vee}} \\
&= |Q / DdQ^\vee| \delta_{\mu - \eta \in DdP^\vee} \sum_{\eta' \in C} \tilde{s}_{\eta, \eta'} \tilde{s}_{\eta', 0} \xi^{2\langle \rho + \eta', \mu - \eta \rangle}.
\end{aligned}$$

Lemma 3.15. — *Let $\gamma \in DdP^\vee$. For any $w \in W$, we have $\gamma - w(\gamma) \in DdQ^\vee$.*

Proof. — See [Bou68, VI.1.10, Proposition 27]. \square

Now, fix $\gamma \in DdP^\vee$. Following [BK01, Section 3.3], we have an action of the affine Weyl group $W^a = W \ltimes DdQ^\vee$ on P such that C contains exactly one element for every orbit with trivial stabilizer for the dot action (note that in [BK01, Section 3.3], the translation subgroup of W^a is generated by dQ^\vee , but Q^\vee is embedded in P using $D^{-1}\langle \cdot, \cdot \rangle$, so that it coincides with our notations). Using the fact that $\tilde{s}_{\eta, w \bullet \eta'} = (-1)^{l(w)} \tilde{s}_{\eta, \eta'}$ and Lemma 3.15, we see that $\tilde{s}_{\eta, \eta'} \tilde{s}_{\eta', 0} \xi^{2\langle \rho + \eta', \mu - \eta \rangle}$ is invariant by the action of W^a . We therefore can replace

the summation on C by a summation on P/DdQ^\vee and we obtain, using the formula for \tilde{s} in the proof of [BK01, Theorem 3.3.20]:

$$\begin{aligned} \sum_{\eta' \in C} \tilde{s}_{\eta, \eta'} \tilde{s}_{\eta', 0} \xi^{2\langle \rho + \eta', \gamma \rangle} &= \frac{1}{|W| \kappa} \sum_{w, w' \in W} (-1)^{l(w) + l(w')} \sum_{\eta' \in P/DdQ^\vee} \xi^{2\langle \eta' + \rho, w(\eta + \rho) + w'(\rho) + \gamma \rangle} \\ &= \frac{|P/DdQ^\vee|}{|W| \kappa} \sum_{w, w' \in W} (-1)^{l(w) + l(w')} \delta_{w(\eta + \rho) + w'(\rho) + \gamma \in DdQ^\vee}, \end{aligned}$$

where κ is a non-zero constant.

As $\gamma \in DdP^\vee$ the stabilizer of γ for the dot action is trivial, and there exist a unique $\tilde{\gamma} \in C$ and $\tilde{w} \in W$ such that $\gamma + \rho \in \tilde{w}(\tilde{\gamma} + \rho) + DdQ^\vee$. Now $w(\eta + \rho) + w'(\rho) + \gamma \in DdQ^\vee$ if and only if $w(\eta + \rho) \in -\gamma - w'(\rho) + DdQ^\vee$. But $w'(\rho) \in -w'(\gamma) + w'\tilde{w}(\tilde{\gamma} + \rho) + DdQ^\vee$ and therefore, using Lemma 3.15, $w(\eta + \rho) + w'(\rho) + \gamma \in DdQ^\vee$ if and only if $\eta + \rho \in w^{-1}w'\tilde{w}w_0(-w_0(\tilde{\gamma}) + \rho) + DdQ^\vee$. But this is possible if and only if $\eta = -w_0(\tilde{\gamma})$ and $w = w'\tilde{w}w_0$ and therefore

$$\sum_{\eta' \in C} \tilde{s}_{\eta, \eta'} \tilde{s}_{\eta', 0} \xi^{2\langle \rho + \eta', \gamma \rangle} = \kappa (-1)^{l(\tilde{w}) + l(w_0)} \delta_{\eta, -w_0(\tilde{\gamma})}.$$

Therefore, the simple objects in the symmetric center of $\mathbb{Z}(\mathcal{T})_\xi \rtimes \mathcal{S}$ are indexed by $(\lambda, \mu) \in \overline{C}$ where $\frac{\mu - \lambda}{2} = \gamma \in DdP^\vee / DdQ^\vee$ and $\frac{\mu + \lambda}{2} = -w_0(\tilde{\gamma})$, where $\tilde{\gamma}$ is the only element in C in the orbit of γ under the dot action of W^a . \square

4. The type A

In this section, we investigate in details the category $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ for \mathfrak{sl}_{n+1} at an even root of unity.

Notations. In this section $\mathfrak{g} = \mathfrak{sl}_{n+1}$ and ξ is a primitive $2d$ -th root of unity, where $d \geq n + 1$. To be consistent with the notation of Section 3 we set $l = 2d$ and $l' = d$. We use the conventions of [Bou68, Planche I] for the labelling of roots.

4.1. The category $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$. — In type A, we have $Q^\vee = Q$ therefore this category has $|C||Q/dQ| = d^n |C|$ simple objects. From the description of C given in Section 3.4, we have

$$C = \left\{ \sum_{i=1}^n \eta_i \varpi_i \in P^+ \mid \sum_{i=1}^n \eta_i \leq d - (n + 1) \right\}$$

so that $|C| = \binom{d-1}{n}$. Recall the notations \tilde{C} and \overline{C} at the end of Section 3.7.

The category $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ is not modular by Theorem 3.14. To compute the $n + 1$ simple objects in the symmetric center of $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$, we follow the proof of Theorem 3.14: for every $\gamma \in dP^\vee / dQ^\vee$, we find a representative $\tilde{\gamma} \in C$ of γ for the dot action of the affine Weyl group W^a . The group dP^\vee / dQ^\vee is generated by the image of $d\varpi_n$ and we have

$$s_n s_{n-1} \cdots s_1 (d\varpi_n + \rho) - \rho \equiv d\varpi_n - \sum_{i=1}^n s_n s_{n-1} \cdots s_{n+2-i} (\alpha_{n+1-i}) \pmod{dQ^\vee}.$$

But $s_n s_{n-1} \cdots s_{n+2-i} (\alpha_{n+1-i}) = \sum_{j=n+1-i}^n \alpha_j$ and therefore

$$s_n s_{n-1} \cdots s_1 \bullet (d\varpi_n) \equiv d\varpi_n - (n + 1)\varpi_n \pmod{dQ^\vee},$$

which is indeed an element of C . Therefore the element

$$I = L_\xi((d - (n + 1))\varpi_1 - d\varpi_n, (d - (n + 1))\varpi_1 + d\varpi_n)$$

is in the symmetric center of $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$. As $\varpi_1 + \varpi_n \in Q^\vee$, we have an isomorphism $I \simeq L_\xi((2d - (n + 1))\varpi_1, -(n + 1)\varpi_1)$ in $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$.

Proposition 4.1. — *The symmetric center of $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ is generated by I as a tensor category. The object I is of positive and negative quantum dimension $(-1)^n$ and of twist 1. Moreover, tensorisation by I has no fixed points on the set of simple objects of $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$.*

Proof. — First, the object I is invertible because the object $L_\xi((d - (n + 1))\varpi_1, (d - (n + 1))\varpi_1)$ in \mathcal{T}_ξ is invertible and the tensor product $L_\xi((d - (n + 1))\varpi_1, (d - (n + 1))\varpi_1) \otimes L_\xi(\eta, \eta)$, $\eta = \sum_{i=1}^n \eta_i \varpi_i \in C$, is given by

$$L_\xi((d - (n + 1))\varpi_1, (d - (n + 1))\varpi_1) \otimes L_\xi(\eta, \eta) \simeq L_\xi\left(\sum_{i=1}^n \eta_{i-1} \varpi_i, \sum_{i=1}^n \eta_{i-1} \varpi_i\right),$$

where we set $\eta_0 = d - (n + 1) - \sum_{i=1}^n \eta_i \geq 0$ (see the proof of [Bru00, Lemme 5.1]). Let $(\lambda, \mu) \in \tilde{C}$. Write this weight as

$$\mu = \sum_{i=1}^n \mu_i \alpha_i \quad \text{and} \quad \lambda = -\mu + 2 \sum_{i=1}^n \eta_i \varpi_i$$

with $\mu_i \in \mathbb{Z}$, $\eta_i \in \mathbb{N}$ and $\sum_{i=1}^n \eta_i \leq d - (n + 1)$. Therefore, using the braiding, we have

$$I \otimes L_\xi(\lambda, \mu) \simeq L_\xi\left(\lambda + \sum_{i=1}^n (\eta_{i-1} - \eta_i) \varpi_i + d\varpi_1, \mu + \sum_{i=1}^n (\eta_{i-1} - \eta_i) \varpi_i - d\varpi_1\right).$$

From this, we see that the objects $I^{\otimes k}$ for $0 \leq k \leq n$ are non-isomorphic. Since I is in the symmetric center of $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$, which contains $|P/Q| = n + 1$ simple objects, all the simple objects in this symmetric center are given by the powers of I .

We can compute the quantum dimension directly in \mathcal{C}_ξ . As a $\mathcal{U}_\xi(\mathfrak{sl}_{n+1})$ -module, the quantum dimension of $L_\xi((d - (n + 1))\varpi_1, (d - (n + 1))\varpi_1)$ is 1: this object is invertible and its quantum dimension is positive [BK01, Theorem 3.3.9]. Therefore, the positive quantum dimension of I is the one of $L_\xi(d\varpi_1, -d\varpi_1)$ which is $\xi^{-d\langle 2\rho, \varpi_1 \rangle} = (-1)^n$.

The twist is given by $\xi^{\langle (2d - (n + 1))\varpi_1 + 2\rho, -(n + 1)\varpi_1 \rangle} = \xi^{-2dn} = 1$. The last assertion is easy, since the grading of I is $d\varpi_1 \notin dQ$. \square

4.2. Dimension of $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ and renormalization. — Thanks to the decomposition $L_\xi(\lambda, \mu) \simeq L_\xi\left(\frac{\lambda + \mu}{2}, \frac{\lambda + \mu}{2}\right) \otimes L_\xi\left(\frac{\lambda - \mu}{2}, -\frac{\lambda - \mu}{2}\right)$ the squared norm of $L_\xi(\lambda, \mu)$ is the same as the one of $L_\xi\left(\frac{\lambda + \mu}{2}, \frac{\lambda + \mu}{2}\right)$. Therefore

$$\dim(\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}) = d^n N,$$

where N is the dimension of the fusion category of tilting modules for $\mathcal{U}_\xi(\mathfrak{sl}_{n+1})$, which is well known (see [BK01, Theorem 3.3.20], except our N is their D^2)

$$N = (n + 1)d^n (-1)^{|\Phi^+|} \prod_{\alpha \in \Phi^+} \frac{1}{(\xi^{\langle \alpha, \rho \rangle} - \xi^{-\langle \alpha, \rho \rangle})^2}.$$

The Weyl character formula gives

$$\prod_{\alpha \in \Phi^+} \xi^{\langle \alpha, \rho \rangle} - \xi^{-\langle \alpha, \rho \rangle} = \xi^{-2\langle \rho, \rho \rangle} \sum_{w \in \mathfrak{S}_{n+1}} (-1)^{l(w)} \xi^{\langle 2\rho, w \bullet 0 \rangle}.$$

Now, as the category $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ is degenerate, we construct a non-degenerate category from it. If n is even, all elements in the symmetric center are of quantum dimension 1 and of twist 1: adding an isomorphism between I and $\mathbf{1}$ as in Sections 3.6 and 3.7 yields a non-degenerate category of dimension

$$\frac{\dim(\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S})}{n+1} = d^{2n} (-1)^{|\Phi^+|} \xi^{-\langle 2\rho, 2\rho \rangle} \left(\sum_{w \in \mathfrak{S}_{n+1}} (-1)^{l(w)} \xi^{\langle 2\rho, w \bullet 0 \rangle} \right)^{-2}.$$

If n is odd, one half of the simple objects of the symmetric center are of quantum dimension 1 and of twist 1 and the other half is of quantum dimension -1 and of twist 1. We first add an isomorphism between $I \otimes I$ and $\mathbf{1}$ and we obtain a slightly degenerate category of superdimension

$$\frac{\dim(\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S})}{n+1} = d^{2n} (-1)^{|\Phi^+|} \xi^{-\langle 2\rho, 2\rho \rangle} \left(\sum_{w \in \mathfrak{S}_{n+1}} (-1)^{l(w)} \xi^{\langle 2\rho, w \bullet 0 \rangle} \right)^{-2},$$

see [Lac18] for more details. By adding an isomorphism of odd degree between I and $\mathbf{1}$, we obtain a braided pivotal superfusion category, as in [Lac18, Section 3].

In both cases, the category obtained is not spherical and we renormalize the S -matrix by a factor involving the dimension on a certain object $\bar{\mathbf{1}}$ introduced in [Lac18]: it is an object X such that the character induced by X on $\text{Gr}(\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S})$ is the negative quantum dimension. If such an object exists, it is not unique as tensorization by the symmetric center leaves invariant the character induced at the level of the Grothendieck group. Therefore, if n is even, $\bar{\mathbf{1}}$ is well defined in the modularization of $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$, and if n is odd, it is well defined in the super category associated to $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$.

Proposition 4.2. — *The object $\bar{\mathbf{1}}$ belongs to the orbit of $L_\xi(-2\rho, 2\rho)$ under tensorisation by the symmetric center.*

Proof. — We show that the character of $\text{Gr}(\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S})$ induced by $L_\xi(-2\rho, 2\rho)$ is the negative quantum dimension. Denote by χ the character of $\text{Gr}(\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S})$ defined by

$$\chi(X) = \frac{S_{L_\xi(-2\rho, 2\rho), X}}{\dim^+(L_\xi(-2\rho, 2\rho))}.$$

The formula for the S -matrix, together with $\dim^+(L_\xi(-2\rho, 2\rho)) = \xi^{\langle 2\rho, 2\rho \rangle}$ gives

$$\chi(L_\xi(\lambda, \mu)) = \frac{\sum_{w \in \mathfrak{S}_{n+1}} (-1)^{l(w)} \xi^{\langle \lambda + 2\rho, (w \bullet (-2\rho, 2\rho))_2 \rangle + \langle (w \bullet (-2\rho, 2\rho))_1 + 2\rho, \mu \rangle}}{\xi^{\langle 2\rho, 2\rho \rangle} \sum_{w \in \mathfrak{S}_{n+1}} (-1)^{l(w)} \xi^{\langle 2\rho, w \bullet 0 \rangle}}.$$

It is easy to check that $w \bullet (-2\rho, 2\rho) = w \bullet (0, 0) + (-2\rho, 2\rho)$ and therefore

$$\begin{aligned} & \langle \lambda + 2\rho, (w \bullet (-2\rho, 2\rho))_2 \rangle + \langle (w \bullet (-2\rho, 2\rho))_1 + 2\rho, \mu \rangle - \langle 2\rho, 2\rho \rangle \\ &= \langle \lambda + \mu + 2\rho, w(\rho) \rangle + \left\langle \frac{\lambda - \mu}{2} - \rho, 2\rho \right\rangle \\ &= \left\langle w^{-1} \bullet \frac{\lambda + \mu}{2}, 2\rho \right\rangle + \left\langle \frac{\lambda - \mu}{2}, 2\rho \right\rangle. \end{aligned}$$

Then, the value of χ at $L_\xi(\lambda, \mu)$ is given by

$$\begin{aligned}\chi(L_\xi(\lambda, \mu)) &= \xi^{\langle \frac{\lambda-\mu}{2}, 2\rho \rangle} \frac{\sum_{w \in \mathfrak{S}_{n+1}} (-1)^{l(w)} \xi^{\langle w \bullet \frac{\lambda+\mu}{2}, 2\rho \rangle}}{\sum_{w \in \mathfrak{S}_{n+1}} (-1)^{l(w)} \xi^{\langle 2\rho, w \bullet 0 \rangle}} \\ &= \dim^- \left(L_\xi \left(\frac{\lambda-\mu}{2}, -\frac{\lambda-\mu}{2} \right) \right) \dim^\pm \left(L_\xi \left(\frac{\lambda+\mu}{2}, \frac{\lambda+\mu}{2} \right) \right) \\ &= \dim^-(L_\xi(\lambda, \mu)),\end{aligned}$$

as stated. \square

Finally, we renormalize S with a square root of

$$\frac{\dim(\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S})}{n+1} \dim^+(\bar{\mathbf{I}}) = d^{2n} (-1)^{|\Phi^+|+n} \left(\sum_{w \in \mathfrak{S}_{n+1}} (-1)^{l(w)} \xi^{\langle 2\rho, w \bullet 0 \rangle} \right)^{-2},$$

which is

$$d^n i^{|\Phi^+|+n} \left(\sum_{w \in \mathfrak{S}_{n+1}} (-1)^{l(w)} \xi^{\langle 2\rho, w \bullet 0 \rangle} \right)^{-1}$$

up to a sign, where i is a square root of -1 in \mathbb{C} .

We choose for each orbit of simple objects of $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ a representative of the orbit under tensorisation by the symmetric center, such that $\mathbf{1}$ is the representative of the orbit of $\mathbf{1}$ and $I \otimes L_\xi(-2\rho, 2\rho)$ is the representative of the orbit of $L_\xi(-2\rho, 2\rho)$. This choice will be explained in Section 5.4.

Theorem 4.3. — *If n is even, the category $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ gives rise to a non-degenerate braided pivotal fusion category. If n is odd, the category $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ gives rise to a non-degenerate braided pivotal superfusion category. In both cases the renormalized S -matrix is given by*

$$\tilde{S}_{(\lambda, \mu), (\lambda', \mu')} = i^{-n-|\Phi^+|} \frac{\sum_{w \in W} (-1)^{l(w)} \xi^{\langle 2\rho + \lambda, (w \bullet (\lambda', \mu'))_2 \rangle + \langle \mu, (w \bullet (\lambda', \mu'))_1 + 2\rho \rangle}}{d^n}$$

and the twist by

$$\theta_{\lambda, \mu} = \xi^{\langle \lambda + 2\rho, \mu \rangle}.$$

5. Malle's \mathbb{Z} -modular datum

We refer to [Mal95] and [Cun07] for most of the material of this section. Let d and n be positive integers. Let ζ be a primitive d -th of unity.

5.1. Set up. — Let $Y = \{1, 2, \dots, nd+1\}$ of cardinal $nd+1$ and $\pi: Y \rightarrow \mathbb{N}$ be the map defined by

$$\pi(k) = \begin{cases} n & \text{if } 1 \leq k \leq n+1 \\ \lfloor \frac{k-n-2}{d-1} \rfloor & \text{if } n+2 \leq k \leq nd+1 \end{cases}.$$

Let $\Psi(Y, \pi)$ be the set of maps $f: Y \rightarrow \{0, 1, \dots, d-1\}$ such that f is strictly increasing on each $\pi^{-1}(i)$, $0 \leq i \leq n$. Since for $0 \leq i \leq n-1$ the set $\pi^{-1}(i)$ is of cardinal $d-1$, there exists a unique element $k_i(f) \in \{0, \dots, d-1\}$ such that $\{0, \dots, d-1\} = f(\pi^{-1}(i)) \cup \{k_i(f)\}$. Note that f is then determined by the values of $f(1) < \dots < f(n+1)$ and of $k_0(f), \dots, k_{n-1}(f)$. For $f \in \Psi(Y, \pi)$, we set

$$\varepsilon(f) = (-1)^{|\{(y, y') \in Y \times Y \mid y < y' \text{ and } f(y) < f(y')\}|}.$$

Let V be a \mathbb{C} -vector space of dimension d with basis $(v_i)_{0 \leq i \leq d-1}$. We denote by \mathcal{S} the square matrix $(\zeta^{ij})_{0 \leq i, j \leq d-1}$ and we will view it as an endomorphism of V . We set $\tau(d) = (-1)^{d(d-1)/2} \det(\mathcal{S}) = \prod_{0 \leq i < j \leq d-1} (\zeta^i - \zeta^j)$. We consider the automorphism of the vector space $(\bigwedge^{n+1} V) \otimes (\bigwedge^{d-1} V)^{\otimes n}$ given by $(\bigwedge^{n+1} \mathcal{S}) \otimes (\bigwedge^{d-1} \mathcal{S})^{\otimes n}$. This space has a basis given by

$$\mathbf{v}_f = (v_{f(1)} \wedge \cdots \wedge v_{f(n+1)}) \otimes (v_{f(n+2)} \wedge \cdots \wedge v_{f(n+d)}) \otimes \cdots \otimes (v_{f(n+2+(n-1)(d-1))} \wedge \cdots \wedge v_{f(n+d+1)})$$

for $f \in \Psi(Y, \pi)$. Denote by \mathbf{S} the matrix of $(\bigwedge^{n+1} \mathcal{S}) \otimes (\bigwedge^{d-1} \mathcal{S})^{\otimes n}$ in the basis $(\mathbf{v}_f)_{f \in \Psi(Y, \pi)}$

$$\left((\bigwedge^{n+1} \mathcal{S}) \otimes (\bigwedge^{d-1} \mathcal{S})^{\otimes n} \right) (\mathbf{v}_f) = \sum_{g \in \Psi(Y, \pi)} \mathbf{S}_{g, f} \mathbf{v}_g.$$

The following lemma follows immediately from [BR17, Lemma 6.2].

Lemma 5.1. — *Let $f, g \in \Psi(Y, \pi)$. Then we have*

$$\mathbf{S}_{f, g} = (-1)^{\sum_{i=0}^{n-1} (k_i(f) + k_i(g))} \frac{(-1)^{nd(d-1)/2} \tau(d)^n}{d^n} \prod_{i=0}^{n-1} \zeta^{-k_i(f)k_i(g)} \sum_{\sigma \in \mathfrak{S}_{n+1}} (-1)^{l(\sigma)} \prod_{i=1}^{n+1} \zeta^{f(i)g(\sigma(i))}.$$

5.2. Malle's \mathbb{Z} -modular datum. — Following [Mal95], we consider the family \mathcal{F} of unipotent characters of $G\left(d, 1, \frac{n(n+1)}{2}\right)$ parametrized by reduced d -symbols with values in the multiset

$$\{0^{d-1}, 1^{d-1}, \dots, (n-1)^{d-1}, n^{n+1}\}.$$

These d -symbols are in bijection with the set

$$\Psi^\#(Y, \pi) = \left\{ f \in \Psi(Y, \pi) \mid \sum_{y \in Y} f(y) \equiv n \binom{d}{2} \pmod{d} \right\}.$$

Indeed to $f \in \Psi^\#(Y, \pi)$ we associate the reduced d -symbol $S = (S_0, \dots, S_{d-1})$ with entries $i \in S_j$ for all $j \in f(\pi^{-1}(i))$. We define for $f \in \Psi^\#(Y, \pi)$

$$\text{Fr}(f) = \zeta_*^{nd(1-d^2)} \prod_{y \in Y} \zeta_*^{-6(f(y)^2 + df(y))},$$

where ζ_* is a primitive $12d$ -th root of unity such that $\zeta_*^{12} = \zeta$. Denote by \mathbb{T} the diagonal matrix with entries $(\text{Fr}(f))_{f \in \Psi^\#(Y, \pi)}$.

Following [Cun07, Section 5], we denote by $\mathbb{S} = (\mathbb{S}_{f, g})_{f, g \in \Psi^\#(Y, \pi)}$ the square matrix defined by

$$\mathbb{S}_{f, g} = \frac{(-1)^{n(d-1)}}{\tau(d)^n} \mathbf{S}_{f, g}.$$

Define $f_{\text{sp}} \in \Psi^\#(Y, \pi)$ by $f_{\text{sp}}(i) = i - 1$ for $1 \leq i \leq n + 1$ and $k_j(f_{\text{sp}}) = j + 1$ for $0 \leq j \leq n - 1$. Note that $\mathbf{S}_{f_{\text{sp}}, g} \neq 0$ for all $g \in \Psi(Y, \pi)$ as it is, up to a non-zero constant, the value of a Vandermonde determinant. The following is due to Malle [Mal95, 4.15].

Proposition 5.2. — *We have*

1. $\mathbb{S}^4 = (\mathbb{S}\mathbb{T})^3 = [\mathbb{S}^2, \mathbb{T}] = 1$.
2. \mathbb{S} is symmetric and unitary.

3. For all $f, g, h \in \Psi^\#(Y, \pi)$

$$N_{f,g}^h = \sum_{k \in \Psi^\#(Y, \pi)} \frac{\mathbb{S}_{f,k} \mathbb{S}_{g,k} \overline{\mathbb{S}_{h,k}}}{\mathbb{S}_{f_{\text{sp}},k}} \in \mathbb{Z}.$$

The special symbol f_{sp} also parametrizes a complex representation of $G\left(d, 1, \frac{n(n+1)}{2}\right)$ and the complex conjugate of this representation is parametrized by the cospecial symbol, see [Mal95, 2.A, 2.D] for more details. Explicitly, the cospecial symbol f_{cosp} is given by

$$f_{\text{cosp}}(i) = \begin{cases} 0 & \text{if } i = 1 \\ d - n + i - 2 & \text{otherwise} \end{cases} \quad \text{and} \quad k_i(f_{\text{cosp}}) = d - i - 1.$$

Given a matrix \mathbb{S} as above, we can associate a fusion algebra A : it is a free \mathbb{Z} -algebra with a base $(b_f)_{f \in \Psi^\#(Y, \pi)}$ such that the multiplication on this base is given by

$$b_f \cdot b_g = \sum_{k \in \Psi^\#(Y, \pi)} N_{f,g}^k b_k.$$

This multiplication is associative, and Cuntz made a remarkable conjecture about the algebra A :

Conjecture 5.3 ([Cun05, Vermutung 5.1.6]). — *Let A^{abs} be a free \mathbb{Z} -module equipped with a base $(b_f^{\text{abs}})_{f \in \Psi^\#(Y, \pi)}$. We define a multiplication on the base by*

$$b_f^{\text{abs}} \cdot b_g^{\text{abs}} = \sum_{k \in \Psi^\#(Y, \pi)} |N_{f,g}^k| b_k^{\text{abs}}.$$

Then this multiplication is associative.

Remark 5.4. — If there exists $(d_f)_{f \in \Psi^\#(Y, \pi)}$ with $d_f \in \{\pm 1\}$ such that the structure constants of the ring A relative to the base $(d_f b_f)_{f \in \Psi^\#(Y, \pi)}$ are positive, then this conjecture is clearly true: one have $|N_{f,g}^k| = d_f d_g d_k N_{f,g}^k$ and the associativity of A^{abs} follows from the associativity of A .

If d is even and n is odd, it seems impossible to find these $(d_f)_{f \in \Psi^\#(Y, \pi)}$, see [Lac18, Section 4] when $n = 1$.

We will compare this modular datum to the one constructed in Theorem 4.3, which will give a proof of the Conjecture 5.3. For this, we rewrite the expressions for \mathbb{S} and \mathbb{T} . First, note that $f \in \Psi(Y, \pi)$ is in $\Psi^\#(Y, \pi)$ if and only if

$$\sum_{i=1}^{n+1} f(i) \equiv \sum_{i=0}^{n-1} k_i(f) \pmod{d}.$$

Therefore, we get rid of $f(n+1)$ in the expressions of \mathbb{S} and \mathbb{T} .

Proposition 5.5. — *For $f, g \in \Psi^\#(Y, \pi)$ we have*

$$\text{Fr}(f) = \zeta^{\sum_{i=1}^n (k_{i-1}(f) - f(i))} \left(\sum_{j=1}^i f(j) - \sum_{j=1}^{i-1} k_{j-1}(f) \right)$$

and

$$\begin{aligned} \mathbb{S}_{f,g} &= \frac{1}{d^n} \varepsilon(f) \varepsilon(g) (-1)^{\sum_{i=1}^n (k_{i-1}(f) + k_{i-1}(g))} \\ &\quad \sum_{\sigma \in \mathfrak{S}_{n+1}} (-1)^{l(\sigma)} \zeta^{\sum_{i=1}^n [(k_{i-1}(f) - f(i)) (\sum_{j=1}^i (\sigma \cdot g)(j) - \sum_{j=1}^{i-1} k_{j-1}(\sigma \cdot g)) + (k_{i-1}(\sigma \cdot g) - (\sigma \cdot g)(i)) (\sum_{j=1}^i f(j) - \sum_{j=1}^{i-1} k_{j-1}(f))]} \end{aligned}$$

Proof. — Let us begin by the value of $\text{Fr}(f)$. By definition, $\text{Fr}(f) = \zeta_*^\alpha$ where

$$\alpha = nd(1-d^2) - 6 \sum_{y \in Y} (f(y)^2 + df(y)),$$

which we consider as an element of $\mathbb{Z}/12d\mathbb{Z}$. First, as $\{0, \dots, d-1\} = f(\pi^{-1}(i)) \cup \{k_i(f)\}$ for $0 \leq i \leq n-1$, we have

$$\sum_{y \in Y} (f(y)^2 + df(y)) = \sum_{i=1}^{n+1} (f(i)^2 + df(i)) + n \sum_{i=0}^{d-1} (i^2 + di) - \sum_{i=1}^n (k_{i-1}(f)^2 + dk_{i-1}(f)).$$

But $6 \sum_{i=0}^{d-1} (i^2 + di) = (d-1)d(2d-1) + 3d^2(d-1) = d(d-1)(5d-1)$ and therefore

$$d(1-d^2) - 6 \sum_{i=0}^{d-1} (i^2 + di) = 6d^2(1-d) \equiv 0 \pmod{12d},$$

so that

$$\alpha = -6 \left(\sum_{i=1}^{n+1} (f(i)^2 + df(i)) - \sum_{i=1}^n (k_{i-1}(f)^2 + dk_{i-1}(f)) \right).$$

Fix $\eta \in \mathbb{Z}$ such that $f(n+1) = \sum_{i=1}^n (k_{i-1}(f) - f(i)) + \eta d$, so that we have

$$\begin{aligned} f(n+1)^2 + df(n+1) &= \left(\sum_{i=1}^n (k_{i-1}(f) - f(i)) \right)^2 + d \sum_{i=1}^n (k_{i-1}(f) - f(i)) \\ &\quad + 2d\eta \sum_{i=1}^n (k_{i-1}(f) - f(i)) + \underbrace{d^2 \eta(1+\eta)}_{\equiv 0 \pmod{2}} \\ &\equiv \left(\sum_{i=1}^n (k_{i-1}(f) - f(i)) \right)^2 + d \sum_{i=1}^n (k_{i-1}(f) - f(i)) \pmod{2d}. \end{aligned}$$

Finally

$$\begin{aligned} \alpha &= -6 \left(\left(\sum_{i=1}^n (k_{i-1}(f) - f(i)) \right)^2 + \sum_{i=1}^n (k_{i-1}(f)^2 - f(i)^2) \right) \\ &= 12 \left(\sum_{i=1}^n f(i)(k_{i-1}(f) - f(i)) - 2 \sum_{1 \leq j < i \leq n} (k_{i-1}(f) - f(i))(k_{j-1}(f) - f(j)) \right) \\ &= 12 \sum_{i=1}^n (k_{i-1}(f) - f(i)) \left(\sum_{j=1}^i f(j) - \sum_{j=1}^{i-1} k_{j-1}(f) \right) \end{aligned}$$

which gives the expected formula for $\text{Fr}(f)$.

We now turn to the formula for \mathbb{S} . Since $\overline{\tau(d)} = (-1)^{(d-1)(d-2)/2} \tau(d)$ we have

$$\mathbb{S}_{f,g} = \frac{(-1)^{\sum_{i=0}^{n-1} (k_i(f) + k_i(g))}}{d^n} \prod_{i=0}^{n-1} \zeta^{k_i(f)k_i(\sigma \cdot g)} \sum_{\sigma \in \mathfrak{S}_{n+1}} \varepsilon(\sigma) \prod_{i=1}^{n+1} \zeta^{-f(i)(\sigma \cdot g)(i)},$$

where $\varepsilon(\sigma)$ is the parity of the permutation σ . Fix $\sigma \in \mathfrak{S}_{n+1}$ and let $h = \sigma \cdot g$. Then

$$\begin{aligned}
& \sum_{i=1}^n k_{i-1}(f)k_{i-1}(h) - \sum_{i=1}^{n+1} f(i)h(i) \\
&= \sum_{i=1}^n (k_{i-1}(f)k_{i-1}(h) - f(i)h(i)) - \sum_{1 \leq i, j \leq n} (k_{i-1}(f) - f(i))(k_{j-1}(h) - h(j)) \\
&= \sum_{i=1}^n (k_{i-1}(f) - f(i))h(i) + \sum_{i=1}^n (k_{i-1}(h) - h(i))f(i) - \sum_{1 \leq i \neq j \leq n} (k_{i-1}(f) - f(i))(k_{j-1}(h) - h(j)) \\
&= \sum_{i=1}^n \left[(k_{i-1}(f) - f(i)) \left(\sum_{j=1}^i h(j) - \sum_{j=1}^{i-1} k_{j-1}(h) \right) + (k_{i-1}(h) - h(i)) \left(\sum_{j=1}^i f(j) - \sum_{j=1}^{i-1} k_{j-1}(f) \right) \right].
\end{aligned}$$

Hence, taking the sum over $\sigma \in \mathfrak{S}_{n+1}$, we obtain the second formula. \square

5.3. Ennola d -ality. — For each d -symbol S in the family \mathcal{F} , Malle defined a polynomial $\gamma_S(q)$, which has similar properties to the degree of a unipotent character for a finite group of Lie type. These polynomials satisfy the Ennola property: there exists a bijection $\mathcal{E}: \mathcal{F} \rightarrow \mathcal{F}$ such that for every symbol S in the family \mathcal{F} , $\gamma_S(\zeta q) = \pm \gamma_{\mathcal{E}(S)}$. The symbol $\mathcal{E}(S)$ is explicitly given in the proof of [Mal95, Folgerung 3.11]. We describe it at the level of functions in $\Psi^\#(Y, \pi)$. Let $f \in \Psi^\#(Y, \pi)$, its Ennola dual $\mathcal{E}(f)$ is given by the unique function g such that

$$\begin{aligned}
k_i(g) &= \left(k_i(f) + i - \frac{n(n+3)}{2} \right)^{\text{res}} \quad \text{and} \\
\{g(i) \mid 1 \leq i \leq n+1\} &= \left\{ \left(f(i) - \frac{n(n+1)}{2} \right)^{\text{res}} \mid 1 \leq i \leq n+1 \right\},
\end{aligned}$$

$(k)^{\text{res}}$ being the remainder in the Euclidean division of k by d .

5.4. Comparison with the modular datum of the category $\mathbb{Z}(\mathcal{T}_{\bar{\zeta}}) \rtimes \mathcal{S}$. — We now compare the modular datum of Malle with the modular datum of Theorem 4.3, from which we use the notation. To each function $f \in \Psi^\#(Y, \pi)$ we associate $\mu = \sum_{i=1}^n \mu_i \alpha_i \in Q$ and $\lambda = -\mu + 2 \sum_{i=1}^n \eta_i \varpi_i \in P$ with

$$\mu_i = \sum_{j=1}^{i-1} k_{j-1}(f) - \sum_{j=1}^i f(j) \quad \text{and} \quad \eta_i = f(i+1) - f(i) - 1.$$

As f takes its values in $\{0, \dots, d-1\}$ and is strictly increasing on $\{1, \dots, n+1\}$ we have $\sum_{i=1}^n \eta_i \leq d-1-n$ and therefore $\lambda + \mu \in 2C$. We then obtain a map

$$\iota: \begin{cases} \Psi^\#(Y, \pi) & \longrightarrow \bar{C} \\ f & \longmapsto (\lambda_f, \mu_f) \end{cases}.$$

Note that the special symbol f_0 is sent to $(0, 0)$ and the cospecial symbol is sent to $(-2\rho + (2d - (n+1))\varpi_1, 2\rho - (n+1)\varpi_1)$. We define a left inverse to ι as follows. Let $(\lambda, \mu) \in \bar{C}$ and define $f_{\lambda, \mu}$ as the unique function $f \in \Psi(Y, \pi)$ such that

$$\{f(1), \dots, f(n+1)\} = \left\{ \left(-\langle \mu, \varpi_1 \rangle + \sum_{j=1}^{i-1} \left\langle \frac{\lambda + \mu}{2} + \rho, \alpha_j \right\rangle \right)^{\text{res}} \mid 1 \leq i \leq n+1 \right\}$$

and

$$k_{i-1}(f) = \left(\sum_{j=1}^i \left\langle \frac{\lambda - \mu}{2} + \rho, \alpha_j \right\rangle \right)^{\text{res}}.$$

An easy computation shows that $f_{\lambda, \mu}$ belongs to $\Psi^\#(Y, \pi)$ and that $f_{\iota(f)} = f$ for any $f \in \Psi^\#(Y, \pi)$. A straightforward computation shows the following lemma.

Lemma 5.6. — *Let $f, g \in \Psi^\#(Y, \pi)$. With the previous notation we have*

$$\begin{aligned} \langle \lambda_f + 2\rho, \mu_g \rangle + \langle \lambda_g + 2\rho, \mu_f \rangle = & - \sum_{i=1}^n (k_{i-1}(f) - f(i)) \left(\sum_{j=1}^i g(j) - \sum_{j=1}^{i-1} k_{j-1}(g) \right) \\ & - \sum_{i=1}^n (k_{i-1}(g) - g(i)) \left(\sum_{j=1}^i f(j) - \sum_{j=1}^{i-1} k_{j-1}(f) \right). \end{aligned}$$

Now, let $\zeta = \xi^{-2}$. It follows immediately that

$$\mathbb{T}_f = \theta_{L_\xi(\lambda_f, \mu_f)}$$

and noticing that $\iota(\sigma \cdot f) = s \bullet \iota(f)$ (where we extend ι to all functions $f: Y \rightarrow \{0, \dots, d-1\}$ such that f is injective on each $\pi^{-1}(i)$, the action of \mathfrak{S}_{n+1} being induced by the action on $\{1, \dots, n+1\} \subset Y$) we have

$$\mathbb{S}_{f,g} = \frac{\sum_{w \in \mathfrak{S}_{n+1}} (-1)^{l(w)} \xi^{(2\rho, w \bullet 0)}}{d^n} \varepsilon(f) \varepsilon(g) (-1)^{\sum_{i=1}^n (k_{i-1}(f) + k_{i-1}(g))} S_{(\lambda_f, \mu_f), (\lambda_g, \mu_g)}.$$

Let D be the diagonal matrix with entries $(\varepsilon(f) (-1)^{\sum_{i=1}^n k_{i-1}(f)})_{f \in \Psi^\#(Y, \pi)}$. We can now state the main theorem of this paper.

Theorem 5.7. — *If n is even (resp. odd) the braided pivotal fusion category (resp. braided pivotal superfusion category) of 4.3 is a categorification of the Malle \mathbb{Z} -modular datum:*

$$\mathbb{S}_{f,g} = i^{-|\Phi^+| - n} D \tilde{S}_{\iota(f), \iota(g)} D^{-1} \quad \text{and} \quad \mathbb{T}_f = \theta_{L_\xi(\iota(f))}.$$

Remark 5.8. — The image of ι contains exactly one representative of each orbit of simple objects of $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ under tensorisation by the symmetric center. As $I \otimes L_\xi(-2\rho, 2\rho)$ is in the image of ι , this justifies our choice of $\tilde{\mathbf{I}}$ made in Section 4.1.

Corollary 5.9. — *The Conjecture 5.3 is true: the multiplication on A^{abs} is associative.*

Proof. — If n is even, this follows from the Remark 5.4. If n is odd, then the Malle's \mathbb{Z} -modular datum is given by a slightly degenerate category, or similarly by a supercategory \mathcal{D} [Lac18, Section 3]. We have at our disposal the super-Grothendieck group of \mathcal{D} , which is a $\mathbb{Z}[\varepsilon]/(\varepsilon^2 - 1)$ -module. We then have ring isomorphisms

$$\text{sGr}(\mathcal{D})/(\varepsilon + 1) \simeq A \quad \text{and} \quad \text{sGr}(\mathcal{D})/(\varepsilon - 1) \simeq A^{\text{abs}}.$$

□

Remark 5.10. — If d and n are odd, we can find a modular subcategory of $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ which is a categorification of the Malle's \mathbb{Z} -modular datum: we simply take the full subcategory of $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ with objects of degree in Q/dQ . Therefore, there exists $(d_f)_{f \in \Psi^\#(Y, \pi)}$ with $d_f = \pm 1$ such that $d_f d_g d_h N_{f,g}^h$ is positive for any $f, g, h \in \Psi^\#(Y, \pi)$. This was also conjectured by Cuntz.

5.5. Categorification of the Ennola property. — The Ennola d -ality gives a bijection \mathcal{E} on $\Psi^\#(Y, \pi)$ and satisfies $\mathcal{E}^d = \text{id}$. Consider $(-\gamma, \gamma)$ in \overline{C} given by $\gamma = \frac{(n+1)(n+2)}{2} \varpi_1 - \rho$.

Proposition 5.11. — *Tensoring by $L_\xi(-\gamma, \gamma)$ is a categorification of the Ennola d -ality. For any $f \in \Psi^\#(Y, \pi)$ we have*

$$L_\xi(\iota(f)) \otimes L_\xi(-\gamma, \gamma) \simeq L_\xi(\iota(\mathcal{E}(f))),$$

the isomorphism being understood in the non-degenerate (super)fusion category associated to $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$.

As γ belongs to Q^\vee , it is clear that $L_\xi(-\gamma, \gamma)^{\otimes d} \simeq \mathbf{1}$.

Proof. — We write γ on the basis of simple roots

$$\gamma = \frac{1}{2} \sum_{i=1}^n (n-i+1)(n-i+2) \alpha_i.$$

Let $f \in \Psi^\#(Y, \pi)$ and let $(\lambda, \mu) = \iota(f) + (-\gamma, \gamma)$. Writing as usual $\mu = \sum_{i=1}^n \mu_i \alpha_i$ and $\lambda = -\mu + 2 \sum_{i=1}^n \eta_i \varpi_i$ we have

$$\mu_i = \sum_{j=1}^{i-1} k_{j-1}(f) - \sum_{j=1}^i f(j) + \frac{(n-i+1)(n-i+2)}{2} \quad \text{and} \quad \eta_i = f(i+1) - f(i) - 1.$$

By definition, $f_{\lambda, \mu}$ is the unique function $g \in \Psi^\#(Y, \pi)$ such that

$$\{g(1), \dots, g(n+1)\} = \left\{ \left(-\langle \mu, \varpi_1 \rangle + \sum_{j=1}^{i-1} \left\langle \frac{\lambda + \mu}{2} + \rho, \alpha_j \right\rangle \right)^{\text{res}} \mid 1 \leq i \leq n+1 \right\}$$

and

$$k_{i-1}(g) = \left(\sum_{j=1}^i \left\langle \frac{\lambda - \mu}{2} + \rho, \alpha_j \right\rangle \right)^{\text{res}}.$$

But as

$$-\langle \mu, \varpi_1 \rangle + \sum_{j=1}^{i-1} \left\langle \frac{\lambda + \mu}{2} + \rho, \alpha_j \right\rangle = f(i) - \frac{n(n+1)}{2}$$

and

$$\begin{aligned} \sum_{j=1}^i \left\langle \frac{\lambda - \mu}{2} + \rho, \alpha_j \right\rangle &= f(i+1) - \frac{n(n+1)}{2} + \mu_{i+1} - \mu_i \\ &= k_{i-1}(f) - \frac{n(n+1)}{2} + \frac{(n-i)(n-i+1)}{2} - \frac{(n-i+1)(n-i+2)}{2} \\ &= k_{i-1}(f) + i - 1 - \frac{n(n+3)}{2} \end{aligned}$$

we have $f_{\lambda, \mu} = f_{\iota(\mathcal{E}(f))}$. As the fibres of the map $(\lambda, \mu) \mapsto f_{\lambda, \mu}$ are exactly the orbits under tensorisation by the symmetric center on the set of simple objects of $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$, there exists $k \in \mathbb{Z}$ such that $L_\xi(\iota(f)) \otimes L_\xi(-\gamma, \gamma) \simeq L_\xi(\iota(\mathcal{E}(f))) \otimes I^{\otimes k}$. \square

6. \mathbb{Z} -modular data associated to exceptional complex reflection groups

The notion of “unipotent characters” has been defined for some exceptional complex reflection group by Broué, Malle and Michel [BMM99], [BMM14], which are called spetsial. The notion of families of such characters, as well as of Fourier transform and eigenvalues of the Frobenius exists, and are available in the package CHEVIE of GAP [GHL⁺96], [Mic15]. In this section, we categorify some of these \mathbb{Z} -modular data using subcategories of $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$, for a well suited simple complex Lie algebra \mathfrak{g} and ξ a well chosen root of unity. The unipotent characters of a complex reflection group G are obtained in CHEVIE and displayed with the following command (we do the example of G_4):

```
gap> G:=ComplexReflectionGroup(4);;
gap> U:=UnipotentCharacters(G);;
gap> Display(U,rec(byFamily:=true));
Unipotent characters for G4
```

Name	Degree	FakeDegree	Eigenvalue	Label
*phi{1,0}	1	1	1	
*phi{2,1}	$(3-ER(-3))/6qP'3P4P^6$	qP4	1	1.E3
#phi{2,3}	$(3+ER(-3))/6qP^3P4P'6$	q^3P4	1	1.E3^2
Z3:2	$-ER(-3)/3qP1P2P4$	0	E3^2	E3.E3^2
*phi{3,2}	q^2P3P6	q^2P3P6	1	
*phi{1,4}	$-ER(-3)/6q^4P^3P4P^6$	q^4	1	1.-E3^2
phi{2,5}	$1/2q^4P2^2P6$	q^5P4	1	1.E3^2
G4	$-1/2q^4P1^2P3$	0	-1	-E3^2.-1
Z3:11	$-ER(-3)/3q^4P1P2P4$	0	E3^2	E3.-E3
#phi{1,8}	$ER(-3)/6q^4P'3P4P'6$	q^8	1	-1.E3^2

The unipotents characters are ordered by families. In the first column, one can read the name of the unipotent characters; In the second column, one can read the degree of the unipotent character; in the third column, one can read the fake degree, for the principal series unipotent characters; in the fourth column, one can read the eigenvalues of the Frobenius; finally, in the last column, one can read some labelling compatible with a categorification of the modular datum associated with the family.

Cyclotomic polynomials are denoted by P_k, P_k', P_k'' and $E_3 = E(3)$ is a third root of unity.

One can access a family, its Fourier matrix and the eigenvalues of the Frobenius as follows (continuing the same example):

```
gap> f:=U.families[2];
Family("RZ/3^2", [6,5,8])
gap> f.fourierMat;
[ [ -2/3*E(3)-1/3*E(3)^2, -1/3*E(3)-2/3*E(3)^2, -1/3*E(3)+1/3*E(3)^2 ],
  [ -1/3*E(3)-2/3*E(3)^2, -2/3*E(3)-1/3*E(3)^2, 1/3*E(3)-1/3*E(3)^2 ],
  [ -1/3*E(3)+1/3*E(3)^2, 1/3*E(3)-1/3*E(3)^2, 1/3*E(3)-1/3*E(3)^2 ] ]
gap> f.eigenvalues
[ 1, 1, E(3)^2 ]
```

6.1. Two families attached to G_{27} . — We consider the complex reflection group denoted by G_{27} in the classification of Shephard-Todd [ST54]. Two families of unipotent characters are of size 18, the second one and the last but one. The \mathbb{Z} -modular datum they define are complex conjugate to each other, hence we will only consider the second family of

unipotent characters of G_{27} . The \mathbb{Z} -modular datum is in fact the tensor product of a \mathbb{Z} -modular datum of size 3 and of an \mathbb{N} -modular datum of size 6.

6.1.1. *The \mathbb{Z} -modular datum of size 3.* — The Fourier matrix and the eigenvalues of the Frobenius are

$$S = \frac{1}{3} \begin{pmatrix} 1 - \zeta_3 & 1 - \zeta_3^2 & \zeta_3 - \zeta_3^3 \\ 1 - \zeta_3^2 & 1 - \zeta_3 & \zeta_3^2 - \zeta_3 \\ \zeta_3 - \zeta_3^2 & \zeta_3^2 - \zeta_3 & \zeta_3 - \zeta_3^2 \end{pmatrix} \quad \text{and} \quad T = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \zeta_3^2 \end{pmatrix}$$

where ζ_3 is a primitive third root of unity. It is easily checked that it coincides with the \mathbb{Z} -modular datum associated to the non-trivial family of the cyclic complex reflection group $G(3, 1, 1)$. By Theorem 5.7, this \mathbb{Z} -modular datum is categorified by the braided pivotal slightly degenerate fusion category $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ for $\mathfrak{g} = \mathfrak{sl}_2$ and ξ a sixth root of unity such that $\xi^2 = \zeta_3^{-1}$.

Remark 6.1. — Up to conjugation by a diagonal matrix with coefficients in $\{\pm 1\}$, we recover the modular datum associated to the second family of unipotent characters of G_4 , which we gave in the beginning of this Section.

6.1.2. *The \mathbb{N} -modular datum of size 6.* — The Fourier matrix and the eigenvalues of the Frobenius are

$$S = \frac{1}{10} \begin{pmatrix} -\zeta_5^4 + \zeta_5^3 + \zeta_5^2 - \zeta_5 & -\zeta_5^4 + \zeta_5^3 + \zeta_5^2 - \zeta_5 & 2(-\zeta_5^4 + \zeta_5^3 + \zeta_5^2 - \zeta_5) & 2(-\zeta_5^4 + \zeta_5^3 + \zeta_5^2 - \zeta_5) & -5 & -5 \\ -\zeta_5^4 + \zeta_5^3 + \zeta_5^2 - \zeta_5 & -\zeta_5^4 + \zeta_5^3 + \zeta_5^2 - \zeta_5 & 2(-\zeta_5^4 + \zeta_5^3 + \zeta_5^2 - \zeta_5) & 2(-\zeta_5^4 + \zeta_5^3 + \zeta_5^2 - \zeta_5) & 5 & 5 \\ 2(-\zeta_5^4 + \zeta_5^3 + \zeta_5^2 - \zeta_5) & 2(-\zeta_5^4 + \zeta_5^3 + \zeta_5^2 - \zeta_5) & 2(3\zeta_5^4 + 2\zeta_5^3 + 2\zeta_5^2 + 3\zeta_5) & 2(-2\zeta_5^4 - 3\zeta_5^3 - 3\zeta_5^2 - 2\zeta_5) & 0 & 0 \\ 2(-\zeta_5^4 + \zeta_5^3 + \zeta_5^2 - \zeta_5) & 2(-\zeta_5^4 + \zeta_5^3 + \zeta_5^2 - \zeta_5) & 2(-2\zeta_5^4 - 3\zeta_5^3 - 3\zeta_5^2 - 2\zeta_5) & 2(3\zeta_5^4 + 2\zeta_5^3 + 2\zeta_5^2 + 3\zeta_5) & 0 & 0 \\ -5 & -5 & 0 & 0 & 5 & -5 \\ -5 & -5 & 0 & 0 & 5 & -5 \end{pmatrix}$$

and

$$T = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & \zeta_5^3 & 0 & 0 & 0 \\ 0 & 0 & 0 & \zeta_5^2 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

where ζ_5 is a primitive fifth root of unity.

Notation. In this section, \mathfrak{g} is of type B_2 with Cartan matrix

$$\begin{pmatrix} 2 & -2 \\ -1 & 2 \end{pmatrix}.$$

The short simple root is denoted by α_1 and the long simple root is denoted by α_2 . The fundamental weights are $\varpi_1 = \alpha_1 + \frac{1}{2}\alpha_2$ and $\varpi_2 = \alpha_1 + \alpha_2$.

Let ξ be a primitive twentieth root of unity such that $\xi^4 = \zeta_5$ and consider the full subcategory \mathcal{C} of $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ for \mathfrak{g} of type B_2 generated by the objects of grading 0 and $5\varpi_1$. As $10\varpi_1 \in 10Q^\vee \cap Q$, $\{0, 5\varpi_1\}$ is a subgroup of $P/(10Q^\vee \cap Q)$. Hence \mathcal{C} is stable by tensor product. The fundamental chamber C is given by

$$C = \{\lambda_1\varpi_1 + \lambda_2\varpi_2 \in P^+ \mid \lambda_1 + \lambda_2 < 3\} = \{0, \varpi_1, \varpi_2, \varpi_1 + \varpi_2, 2\varpi_1, 2\varpi_2\}.$$

Therefore, there are 6 simple objects in \mathcal{C} which are labelled by:

- in degree 0: $(0, 0), (\varpi_2, \varpi_2), (2\varpi_1, 2\varpi_1)$ and $(2\varpi_2, 2\varpi_2)$,
- in degree $5\varpi_1$: $(6\varpi_1, -4\varpi_1)$ and $(6\varpi_1 + \varpi_2, -4\varpi_1 + \varpi_2)$.

We choose the following set of representatives of the fusion category \mathcal{C} :

$$\{L_\xi(0,0), L_\xi(2\varpi_2, 2\varpi_2), L_\xi(2\varpi_1, 2\varpi_1), L_\xi(\varpi_2, \varpi_2), L_\xi(6\varpi_1 + \varpi_2, -4\varpi_1 + \varpi_2), L_\xi(6\varpi_1, -4\varpi_2)\}$$

With this order, the S -matrix and the T -matrix of the twist of \mathcal{C} are

$$S_{\mathcal{C}} = \begin{pmatrix} 1 & 1 & 2 & 2 & \zeta_5^4 - \zeta_5^3 - \zeta_5^2 + \zeta_5 & \zeta_5^4 - \zeta_5^3 - \zeta_5^2 + \zeta_5 \\ 1 & 1 & 2 & 2 & -\zeta_5^4 + \zeta_5^3 + \zeta_5^2 - \zeta_5 & -\zeta_5^4 + \zeta_5^3 + \zeta_5^2 - \zeta_5 \\ 2 & 2 & 2\zeta_5^4 + 2\zeta_5 & 2\zeta_5^3 + 2\zeta_5^2 & 0 & 0 \\ 2 & 2 & 2\zeta_5^3 + 2\zeta_5^2 & 2\zeta_5^4 + 2\zeta_5 & 0 & 0 \\ \zeta_5^4 - \zeta_5^3 - \zeta_5^2 + \zeta_5 & -\zeta_5^4 + \zeta_5^3 + \zeta_5^2 - \zeta_5 & 0 & 0 & -\zeta_5^4 + \zeta_5^3 + \zeta_5^2 - \zeta_5 & \zeta_5^4 - \zeta_5^3 - \zeta_5^2 + \zeta_5 \\ \zeta_5^4 - \zeta_5^3 - \zeta_5^2 + \zeta_5 & -\zeta_5^4 + \zeta_5^3 + \zeta_5^2 - \zeta_5 & 0 & 0 & \zeta_5^4 - \zeta_5^3 - \zeta_5^2 + \zeta_5 & -\zeta_5^4 + \zeta_5^3 + \zeta_5^2 - \zeta_5 \end{pmatrix},$$

and

$$T_{\mathcal{C}} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & \zeta_5^3 & 0 & 0 & 0 \\ 0 & 0 & 0 & \zeta_5^2 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

Renormalizing $S_{\mathcal{C}}$ with the square root of 20, which is the dimension of \mathcal{C} , gives us the negative of the matrix S .

Theorem 6.2. — *The S -matrix and the T -matrix of the modular category \mathcal{C} satisfy*

$$\frac{S_{\mathcal{C}}}{\sqrt{20}} = -S \quad T_{\mathcal{C}} = T.$$

Remark 6.3. — The category \mathcal{C} is indeed ribbon since every object is self dual.

6.2. Two families attached to G_{24} . — We consider the complex reflection group denoted by G_{24} in the classification of Shephard-Todd [ST54]. Two families of unipotent characters are of size 7, the second one and the last but one. The \mathbb{Z} -modular datum they define are complex conjugate to each other, hence we will only consider the last but one family of unipotent characters of G_{24} .

The Fourier matrix and the eigenvalues of the Frobenius are

$$S = \frac{1}{14} \begin{pmatrix} -\sqrt{-7} & \sqrt{-7} & 7 & 7 & -2\sqrt{-7} & -2\sqrt{-7} & -2\sqrt{-7} \\ \sqrt{-7} & -\sqrt{-7} & 7 & 7 & 2\sqrt{-7} & 2\sqrt{-7} & 2\sqrt{-7} \\ \frac{7}{7} & \frac{7}{7} & \frac{7}{7} & -\frac{7}{7} & 0 & 0 & 0 \\ -2\sqrt{-7} & 2\sqrt{-7} & 0 & 0 & 2\zeta_7^6 + 4\zeta_7^4 - 4\zeta_7^3 - 2\zeta_7 & -4\zeta_7^6 + 2\zeta_7^5 - 2\zeta_7^2 + 4\zeta_7 & -4\zeta_7^5 - 2\zeta_7^4 + 2\zeta_7^3 + 4\zeta_7^2 \\ -2\sqrt{-7} & 2\sqrt{-7} & 0 & 0 & -4\zeta_7^6 + 2\zeta_7^5 - 2\zeta_7^2 + 4\zeta_7 & -4\zeta_7^5 - 2\zeta_7^4 + 2\zeta_7^3 + 4\zeta_7^2 & 2\zeta_7^6 + 4\zeta_7^4 - 4\zeta_7^3 - 2\zeta_7 \\ -2\sqrt{-7} & 2\sqrt{-7} & 0 & 0 & -4\zeta_7^5 - 2\zeta_7^4 + 2\zeta_7^3 + 4\zeta_7^2 & 2\zeta_7^6 + 4\zeta_7^4 - 4\zeta_7^3 - 2\zeta_7 & -4\zeta_7^6 + 2\zeta_7^5 - 2\zeta_7^2 + 4\zeta_7 \end{pmatrix}$$

and

$$T = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \zeta_7^3 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \zeta_7^5 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \zeta_7^6 \end{pmatrix},$$

where ζ_7 is a primitive seventh root of unity and $\sqrt{-7} = \zeta_7 + \zeta_7^2 - \zeta_7^3 + \zeta_7^4 - \zeta_7^5 - \zeta_7^6$.

In [Cun05, Section 8.2.1 Beispiel 15], Cuntz showed that the fusion algebra associated to this modular datum is related to the Verlinde algebra of type B_3 at a twenty-eighth root of unity.

Let ξ be a primitive twenty-eighth root of unity such that $\xi^4 = \zeta_7$ and consider the category $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ for \mathfrak{g} of type B_3 .

Notations. In this section, \mathfrak{g} is of type B_3 with Cartan matrix

$$\begin{pmatrix} 2 & -2 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}.$$

The short simple root is denoted by α_1 and the two long simple roots are denoted by α_2 and α_3 . The fundamental weights are $\varpi_1 = \frac{3}{2}\alpha_1 + \alpha_2 + \frac{1}{2}\alpha_3$, $\varpi_2 = 2\alpha_1 + 2\alpha_2 + \alpha_3$ and $\varpi_3 = \alpha_1 + \alpha_2 + \alpha_3$.

We have $\varpi_1 = \frac{3}{2}\alpha_1^\vee + 2\alpha_2^\vee + \alpha_3^\vee$ and therefore the subgroup of $P/(14Q^\vee \cap Q)$ generated by $7\varpi_1$ is $\{0, 7\varpi_1, 14\varpi_1, 21\varpi_1\}$. Let \mathcal{C} be the full subcategory of $\mathbb{Z}(\mathcal{T}_\xi) \rtimes \mathcal{S}$ generated by the objects of grading in $\{0, 7\varpi_1, 14\varpi_1, 21\varpi_1\}$. This category is then stable by tensor product. The fundamental chamber C is

$$C = \{\lambda_1\varpi_1 + \lambda_2\varpi_2 + \lambda_3\varpi_3 \in P^+ \mid \lambda_1 + 2\lambda_2 + \lambda_3 < 3\} = \{0, \varpi_1, \varpi_2, \varpi_3, 2\varpi_1, 2\varpi_3, \varpi_1 + \varpi_3\}.$$

Therefore, there are 14 simple objects in \mathcal{C} which are labelled by:

- in degree 0: $(0, 0), (\varpi_2, \varpi_2), (\varpi_3, \varpi_3), (2\varpi_1, 2\varpi_1)$ and $(2\varpi_3, 2\varpi_3)$,
- in degree $7\varpi_1$: $(8\varpi_1, -6\varpi_1)$ and $(8\varpi_1 + \varpi_3, -6\varpi_1 + \varpi_3)$,
- in degree $14\varpi_1$: $(14\varpi_1, -14\varpi_1), (14\varpi_1 + \varpi_2, -14\varpi_1 + \varpi_2), (14\varpi_1 + \varpi_3, -14\varpi_1 + \varpi_3), (16\varpi_1, -12\varpi_1)$ and $(14\varpi_1 + 2\varpi_3, -14\varpi_1 + 2\varpi_3)$
- in degree $21\varpi_1$: $(22\varpi_1, -20\varpi_1)$ and $(22\varpi_1 + \varpi_3, -20\varpi_1 + \varpi_1)$.

Proposition 6.4. — The category \mathcal{C} is slightly degenerate. The non-unit object of its symmetric center is $L_\xi(14\varpi_1 + 2\varpi_3, -14\varpi_1 + 2\varpi_3)$ which is of dimension -1 and of twist 1.

We choose the following set of representatives of the superfusion category $\widehat{\mathcal{C}}$:

$$\{L_\xi(0, 0), L_\xi(14\varpi_1, -14\varpi_1), L_\xi(22\varpi_1, -20\varpi_1), L_\xi(22\varpi_1 + \varpi_1, -20\varpi_1 + \varpi_3), \\ L_\xi(\varpi_3, \varpi_3), L_\xi(\varpi_2, \varpi_2), L_\xi(2\varpi_1, 2\varpi_1)\}.$$

With this order, the S -matrix and the T -matrix of the twist of $\widehat{\mathcal{C}}$ are

$$S_{\widehat{\mathcal{C}}} = \begin{pmatrix} 1 & -1 & \sqrt{-7} & \sqrt{-7} & 2 & 2 & 2 \\ -1 & 1 & \sqrt{-7} & \sqrt{-7} & -2 & -2 & -2 \\ \sqrt{-7} & \sqrt{-7} & \sqrt{-7} & -\sqrt{-7} & 0 & 0 & 0 \\ \sqrt{-7} & \sqrt{-7} & -\sqrt{-7} & \sqrt{-7} & 0 & 0 & 0 \\ 2 & -2 & 0 & 0 & 2\zeta_7^6 + 2\zeta_7 & 2\zeta_7^5 + 2\zeta_7^2 & 2\zeta_7^4 + 2\zeta_7^3 \\ 2 & -2 & 0 & 0 & 2\zeta_7^5 + 2\zeta_7^2 & 2\zeta_7^4 + 2\zeta_7^3 & 2\zeta_7^6 + 2\zeta_7 \\ 2 & -2 & 0 & 0 & 2\zeta_7^4 + 2\zeta_7^3 & 2\zeta_7^6 + 2\zeta_7 & 2\zeta_7^5 + 2\zeta_7^2 \end{pmatrix}$$

and

$$T_{\widehat{\mathcal{C}}} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \zeta_7^3 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \zeta_7^5 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \zeta_7^6 \end{pmatrix}$$

This category is not spherical and the object $\bar{\mathbf{1}}$ is $L_{\xi}(14\varpi_1, -14\varpi_1)$, which is of dimension -1 . Therefore following [Lac18, Theorem 2.7], we renormalise $S_{\widehat{\mathcal{C}}}$ with the square root of -28 , which is $\text{sdim}(\mathcal{C})\dim^+(\bar{\mathbf{1}})$.

Theorem 6.5. — *The S -matrix and the T -matrix of the superfusion category $\widehat{\mathcal{C}}$ satisfy*

$$\frac{S_{\widehat{\mathcal{C}}}}{i\sqrt{28}} = S \quad \text{and} \quad T_{\widehat{\mathcal{C}}} = T,$$

where $i = \xi^7$.

References

- [And92] H. H. ANDERSEN – “Tensor products of quantized tilting modules”, *Comm. Math. Phys.* **149** (1992), no. 1, p. 149–159.
- [BK01] B. BAKALOV & A. KIRILLOV, JR. – *Lectures on tensor categories and modular functors*, University Lecture Series, vol. 21, American Mathematical Society, Providence, RI, 2001.
- [BMM99] M. BROUÉ, G. MALLE & J. MICHEL – “Towards spetses. I”, *Transform. Groups* **4** (1999), no. 2-3, p. 157–218.
- [BMM14] M. BROUÉ, G. MALLE & J. MICHEL – “Split spetses for primitive reflection groups”, *Astérisque* (2014), no. 359, p. vi+146.
- [Bou68] N. BOURBAKI – *Éléments de mathématique. Fasc. XXXIV. Groupes et algèbres de Lie. Chapitre IV: Groupes de Coxeter et systèmes de Tits. Chapitre V: Groupes engendrés par des réflexions. Chapitre VI: systèmes de racines*, Actualités Scientifiques et Industrielles, No. 1337, Hermann, Paris, 1968.
- [BR17] C. BONNAFÉ & R. ROQUIER – “An asymptotic cell category for cyclic groups”, *ArXiv e-prints* (2017), 1708.09730.
- [Bru00] A. BRUGUIÈRES – “Catégories prémodulaires, modularisations et invariants des variétés de dimension 3”, *Math. Ann.* **316** (2000), no. 2, p. 215–236.
- [CP94] V. CHARI & A. PRESSLEY – *A guide to quantum groups*, Cambridge University Press, Cambridge, 1994.
- [Cun05] M. CUNTZ – “Fourier-Matrizen und Ringe mit Basis”, Ph.D. Thesis, Universität Kassel, 2005.
- [Cun07] ———, “Fusion algebras for imprimitive complex reflection groups”, *J. Algebra* **311** (2007), no. 1, p. 251–267.
- [Dri87] V. G. DRINFELD – “Quantum groups”, in *Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Berkeley, Calif., 1986)*, Amer. Math. Soc., Providence, RI, 1987, p. 798–820.
- [EGNO15] P. ETINGOF, S. GELAKI, D. NIKSHYCH & V. OSTRIK – *Tensor categories*, Mathematical Surveys and Monographs, vol. 205, American Mathematical Society, Providence, RI, 2015.
- [EO18] P. ETINGOF & V. OSTRIK – “On semisimplification of tensor categories”, *ArXiv e-prints* (2018), 1801.04409.
- [GHL+96] M. GECK, G. HISS, F. LÜBECK, G. MALLE & G. PFEIFFER – “CHEVIE—a system for computing and processing generic character tables”, *Appl. Algebra Engrg. Comm. Comput.* **7** (1996), no. 3, p. 175–210.
- [Hum78] J. E. HUMPHREYS – *Introduction to Lie algebras and representation theory*, Graduate Texts in Mathematics, vol. 9, Springer-Verlag, New York-Berlin, 1978, Second printing, revised.
- [Jan96] J. C. JANTZEN – *Lectures on quantum groups*, Graduate Studies in Mathematics, vol. 6, American Mathematical Society, Providence, RI, 1996.

- [KS97] A. KLIMYK & K. SCHMÜDGEN – *Quantum groups and their representations*, Texts and Monographs in Physics, Springer-Verlag, Berlin, 1997.
- [Lac18] A. LACABANNE – “Slightly degenerate categories and \mathbb{Z} -modular data”, *ArXiv e-prints* (2018), 1807.00766.
- [Lus90] G. LUSZTIG – “Quantum groups at roots of 1”, *Geom. Dedicata* **35** (1990), no. 1-3, p. 89–113.
- [Lus10] ———, *Introduction to quantum groups*, Modern Birkhäuser Classics, Birkhäuser/Springer, New York, 2010, Reprint of the 1994 edition.
- [Mal95] G. MALLE – “Unipotente Grade imprimitiver komplexer Spiegelungsgruppen”, *J. Algebra* **177** (1995), no. 3, p. 768–826.
- [Mic15] J. MICHEL – “The development version of the `chevie` package of `gap3`”, *J. Algebra* **435** (2015), p. 308–336.
- [Müg00] M. MÜGER – “Galois theory for braided tensor categories and the modular closure”, *Adv. Math.* **150** (2000), no. 2, p. 151–201.
- [MW98] G. MASBAUM & H. WENZL – “Integral modular categories and integrality of quantum invariants at roots of unity of prime order”, *J. Reine Angew. Math.* **505** (1998), p. 209–235.
- [Ros90] M. ROSSO – “Analogues de la forme de Killing et du théorème d’Harish-Chandra pour les groupes quantiques”, *Ann. Sci. École Norm. Sup. (4)* **23** (1990), no. 3, p. 445–467.
- [Saw06] S. F. SAWIN – “Quantum groups at roots of unity and modularity”, *J. Knot Theory Ramifications* **15** (2006), no. 10, p. 1245–1277.
- [ST54] G. C. SHEPHARD & J. A. TODD – “Finite unitary reflection groups”, *Canadian J. Math.* **6** (1954), p. 274–304.
- [Tan92] T. TANISAKI – “Killing forms, Harish-Chandra isomorphisms, and universal R -matrices for quantum algebras”, in *Infinite analysis, Part A, B (Kyoto, 1991)*, Adv. Ser. Math. Phys., vol. 16, World Sci. Publ., River Edge, NJ, 1992, p. 941–961.

October 11, 2018

ABEL LACABANNE, Institut Montpellierain Alexander Grothendieck (CNRS: UMR 5149), Université de Montpellier, Case Courrier 051, Place Eugène Bataillon, 34095 MONTPELLIER Cedex, FRANCE
E-mail: `abel.lacabanne@umontpellier.fr`