



HAL
open science

Familial Catecholamine-Induced QT Prolongation in Unexplained Sudden Cardiac Death

Francois Huchet, Florence Kyndt, Julien Barc, Aurelie Thollet, Flavien Charpentier, Richard Redon, Jean Jacques Schott, Herve Le Marec, Vincent Probst, Jean-Baptiste Gourraud

► **To cite this version:**

Francois Huchet, Florence Kyndt, Julien Barc, Aurelie Thollet, Flavien Charpentier, et al.. Familial Catecholamine-Induced QT Prolongation in Unexplained Sudden Cardiac Death. Journal of the American College of Cardiology, 2017, Equipe I Equipe Iia, 69 (12), pp.1642–1643. 10.1016/j.jacc.2017.01.030 . hal-01832147

HAL Id: hal-01832147

<https://hal.science/hal-01832147>

Submitted on 12 Jul 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Letters

Familial Catecholamine-Induced QT Prolongation in Unexplained Sudden Cardiac Death



Despite major progress in molecular and phenotypic characterization of primary electrical disorders, many (aborted) sudden cardiac deaths (SCDs) occur in young victims without identifiable abnormalities (1). This study aimed to describe the use of mental stress test (MST) to identify catecholamine-induced QT prolongation (CIQTP) in SCD familial screening.

MST was performed in the screening of 65 consecutive families affected by unexplained SCD referred to the National Referral Centre for Inherited Cardiac Arrhythmias of Nantes. Conventional screening included echocardiography, exercise test, and epinephrine and ajmaline tests (1). MST was performed following a standardized protocol mostly based on mental arithmetic in stressful conditions (2). Two physicians blinded to patient clinical and genetic status reviewed all the electrocardiograms. Patients were considered affected for CIQTP when presenting prolonged QTc interval >480 ms or if >30 ms QT prolongation with the QTc interval >460 ms (1) during the different tests.

We fortuitously unmasked a significant prolongation of QTc duration in a young woman when mentioning the unexplained SCD of her sister at summer 2013. According to previous description of stress-induced repolarization abnormalities (2,3), we additionally performed MST during familial screening in 65 families with a familial history of SCD in young patients. Conventional screening identified 7 Brugada syndrome, 6 long QT syndrome (LQTS), and 3 familial cardiomyopathies. MST unmasked CIQTP in 4 families (mean age of SCD 18 years) (Figure 1A). In these families, 31 of the 57 tested family members were affected (QTc during MST 501 ± 36 ms vs. 430 ± 19 ms) with a flattening or double-hump T-wave aspect in 13 of 31 versus 1 of 26 in unaffected patients ($p < 0.001$) (Figure 1B).

In 9 patients (including 2 with previous reported syncope) MST was the only positive test. Eight of these patients did not receive an epinephrine test because of


contraindication (<15 years old; $n = 7$) or refusal ($n = 1$). Epinephrine tests were positive in 20 of 21 affected patients (95%), whereas the exercise test failed to identify the QT prolongation in 18 of 30 affected patients (60%). After the diagnosis of CIQTP, 14 family members were treated by beta-blocker therapy and 2 were implanted by an implantable cardioverter-defibrillator. No symptom occurred during a median follow-up of 22 months, in contrast to 6 SCDs and 7 previous syncope. Genetic screening failed to identify any mutation in the 45 susceptibility genes tested (4).

We describe MST as a diagnostic tool, able to unmask CIQTP in a context of unexplained SCD even in patients previously negative after conventional exhaustive screening. Considering the family histories and the presence of CIQTP in symptomatic family members, it is highly probable that CIQTP was responsible of the familial SCD (2,3). CIQTP is distinguished from the classical form of LQTS by the normal baseline electrocardiogram without any QT prolongation, the saddleback T-wave occurring during MST, and the absence of mutations in the LQTS genes.

Even if CIQTP may be identified by both MST and epinephrine test, MST has the advantage of being safer, easier, and quicker to perform. On the other hand, the complex use of epinephrine test can lead to contraindication or misinterpretation. The use of MST in the context of unexplained SCD and LQTS deserves further investigations and comparative studies.

Our preliminary findings have identified CIQTP as a new specific familial phenotype characterized by normal QT duration at rest but major QT lengthening during mental stress. CIQTP seems to play an important role in unexplained ventricular fibrillation and MST appears to be a simple and efficient test to identify this pathology.

Francois Huchet, MD
Florence Kyndt, MD
Julien Barc, PhD
Aurelie Thollet, PharmD, PhD
Flavien Charpentier, PhD
Richard Redon, PhD
Jean Jacques Schott, PhD
Herve le Marec, MD, PhD
Vincent Probst, MD, PhD
*Jean Baptiste Gourraud, MD, PhD



*Service de Cardiologie
 CHU de Nantes-Nord Laennec
 Boulevard du Professeur Jacques Monod
 44800 Saint-Herblain
 France
 E-mail: jeanbaptiste.gourraud@chu-nantes.fr
<http://dx.doi.org/10.1016/j.jacc.2017.01.030>

Please note: The authors have reported that they have no relationships relevant to the contents of this paper to disclose. Drs. Huchet, Kyndt, and Barc contributed equally to this work.

REFERENCES

1. Priori SG, Blomström-Lundqvist C, Mazzanti A, et al. 2015 ESC Guidelines for the management of patients with ventricular arrhythmias and the

prevention of sudden cardiac death: the Task Force for the Management of Patients with Ventricular Arrhythmias and the Prevention of Sudden Cardiac Death of the European Society of Cardiology (ESC) Endorsed by Association for European Paediatric and Congenital Cardiology (AEPC). *Eur Heart J* 2015; 36:2793-867.

2. Paavonen KJ, Swan H, Piippo K, et al. Response of the QT interval to mental and physical stress in types LQT1 and LQT2 of the long QT syndrome. *Heart Br Card Soc* 2001;86:39-44.

3. Finlay MC, Lambiase PD, Ben-Simon R, Taggart P. Effect of mental stress on dynamic electrophysiological properties of the endocardium and epicardium in humans. *Heart Rhythm* 2016;13: 175-82.

4. Le Scouarnec S, Karakachoff M, Gourraud J-B, et al. Testing the burden of rare variation in arrhythmia-susceptibility genes provides new insights into molecular diagnosis for Brugada syndrome. *Hum Mol Genet* 2015;24: 2757-63.