



HAL
open science

Embolism and mechanical resistances play a key-role in dehydration tolerance of a perennial grass *Dactylis glomerata* L

Florence Volaire, Frederic Lens, Hervé Cochard, Hueng Xu, Larissa Chacon Doria, Pauline Bristiel, Jennifer A. Balachowski, Nick P Rowe, Cyrille Violle, Catherine Picon-Cochard

► **To cite this version:**

Florence Volaire, Frederic Lens, Hervé Cochard, Hueng Xu, Larissa Chacon Doria, et al.. Embolism and mechanical resistances play a key-role in dehydration tolerance of a perennial grass *Dactylis glomerata* L. *Annals of Botany*, 2018, 122 (2), pp.325-336. 10.1093/aob/mcy073 . hal-01832287

HAL Id: hal-01832287

<https://hal.science/hal-01832287>

Submitted on 8 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Embolism and mechanical resistances play a key-role in dehydration tolerance of a perennial grass *Dactylis glomerata* L.

Thank you for agreeing to review this paper for Annals of Botany. The Annals of Botany aims to be among the very top of plant science journals and as we receive over 1000 submissions every year we need to be very selective in deciding which papers we can publish. In making your assessment of the manuscript's suitability for publication in the journal please consider the following points.

Scientific Scope

Annals of Botany welcomes papers in all areas of plant science. Papers may address questions at any level of biological organization ranging from molecular through cells and organs, to whole organisms, species, communities and ecosystems. Its scope extends to all flowering and non-flowering taxa, and to evolutionary and pathology research. Many questions are addressed using comparative studies, genetics, genomics, molecular tools, and modeling.

To merit publication in Annals of Botany, contributions should be substantial, concise, written in clear English and combine originality of content with potential general interest.

- We want to publish papers where our reviewers are enthusiastic about the science: is this a paper that you would keep for reference, or pass on to your colleagues? If the answer is "no" then please enter a low priority score when you submit your report.
- We want to publish papers with novel and original content that move the subject forward, not papers that report incremental advances or findings that are already well known in other species. Please consider this when you enter a score for originality when you submit your report.

Notes on categories of papers:

All review-type articles should be **novel, rigorous, substantial and "make a difference" to plant science**. The purpose is to summarise, clearly and succinctly, the "cutting edge" of the subject and how future research would best be directed. Reviews should be relevant to a broad audience and all should have a **strong conclusion and illustrations** including diagrams.

- *Primary Research* articles should report on original research relevant to the scope of the journal, demonstrating an important advance in the subject area, and the results should be clearly presented, novel and supported by appropriate experimental approaches. The Introduction should clearly set the context for the work and the Discussion should demonstrate the importance of the results within that context. Concise speculation, models and hypotheses are encouraged, but must be informed by the results and by the authors' expert knowledge of the subject.
- *Reviews* should place the subject in context, add significantly to previous reviews in the subject area and moving forward research in the subject area. Reviews should be selective, including the most important and best, up-to-date, references, not a blow-by-blow and exhaustive listing.
- *Research in Context* should combine a review/overview of a subject area with original research, often leading to new ideas or models; they present a hybrid of review and research. Typically a Research in Context article contains an extended Introduction that provides a general overview of the topic before incorporating new research results with a Discussion proposing general models and the impact of the research.
- *Viewpoints* are shorter reviews, presenting clear, concise and logical arguments supporting the authors' opinions, and in doing so help to stimulate discussions within the topic.
- *Botanical Briefings* are concise, perhaps more specialised reviews and usually cover topical issues, maybe involving some controversy.

1 **Embolism and mechanical resistances play a key-role in dehydration tolerance of a**
2 **perennial grass *Dactylis glomerata* L.**

3
4
5 Florence Volaire^{1*}, Frederic Lens², Hervé Cochard³, Hueng Xu², Larissa Chacon Doria², Pauline
6 Bristiel⁴, Jennifer Balachowski^{4,5}, Nick Rowe⁶, Cyrille Violle⁴, Catherine Picon-Cochard⁷

7
8 ¹INRA, USC 1338, UMR 5175, Centre d'Ecologie Fonctionnelle et Evolutive, Université de
9 Montpellier, Université Paul Valéry, EPHE, 1919 route de Mende, F-34293 Montpellier France,
10 Florence.volaire@cefe.cnrs.fr; corresponding author Tel: 0033 4 67 61 33 30

11 ²Naturalis Biodiversity Center, Leiden University, PO Box 9517, 2300RA Leiden, The
12 Netherlands, frederic.lens@naturalis.nl, larissa.chacondoria@naturalis.nl; ellie.xu@naturalis.nl

13 ³Université Clermont-Auvergne, INRA, PIAF, F-63000 Clermont-Ferrand,
14 France, herve.cochard@inra.fr

15 ⁴UMR 5175, Centre d'Ecologie Fonctionnelle et Evolutive, Université de Montpellier, Université
16 Paul Valéry, EPHE, 1919 route de Mende, F-34293 Montpellier France,
17 Pauline.bristiel@cefe.cnrs.fr; cyrille.violle@cefe.cnrs.fr

18 ⁵Department of Evolution and Ecology, University of California, Davis, CA 95616,
19 USA, jbalachowski@ucdavis.edu

20 ⁶ Univ Montpellier, UMR - AMAP, Montpellier, F-34000 France; CNRS 5120, UMR AMAP,
21 Montpellier, F-34000 France. nrowe@cirad.fr

22 ⁷INRA, VetAgroSup, UREP, F-63000 Clermont-Ferrand, France, [catherine.picon-](mailto:catherine.picon-cochard@inra.fr)
23 cochard@inra.fr

24

25 **Running title:** Embolism resistance and dehydration tolerance in a perennial grass

26

27 Original Article

28

29

30

31

32

33 **Abstract**

34

35 • **Background and Aims** More intense droughts under climate change threaten species
36 resilience. Hydraulic strategies determine drought survival in woody plants but have been hardly
37 studied in herbaceous species. We explored the intra-specific variability of hydraulic and
38 morphological traits as indicators of dehydration tolerance in a perennial grass, cocksfoot
39 (*Dactylis glomerata*) which has a large biogeographical distribution in Europe.

40 • **Methods** Twelve populations of cocksfoot originating from Mediterranean, Temperate and
41 Northern European areas were grown in a controlled environment in pots. Dehydration tolerance,
42 leaf and stem anatomical traits and xylem pressure associated with 88% or 50% loss of xylem
43 conductance (P_{88} , P_{50}) were measured.

44 • **Key Results** Across the twelve populations of cocksfoot, P_{50} ranged from -3.06 to -6.36 MPa,
45 while P_{88} ranged from -5.06 to -11.6 MPa. This large intraspecific variability of embolism
46 thresholds corresponded with the biogeographical distribution and some key traits of the
47 populations. In particular, P_{88} was correlated with dehydration tolerance ($r=-0.79$). The
48 dehydration-sensitive temperate populations exhibited the highest P_{88} (-6.1 MPa). The most
49 dehydration-tolerant Mediterranean populations had the greatest leaf dry matter content and leaf
50 fracture toughness, and the lowest P_{88} (-10.4 MPa). The Northern populations displayed
51 intermediate trait values, potentially attributable to frost resistance. The thickness of metaxylem
52 vessel walls in stems was highly correlated with P_{50} ($r=-0.92$) but no trade-off with stem
53 lignification was observed. The relevance of the linkage between hydraulic and stomatal traits is
54 discussed for drought survival in perennial grasses.

55 • **Conclusions** Compared to woody species, the large intra-specific variability in dehydration
56 tolerance and embolism resistance within cocksfoot has consequences for its sensitivity to
57 climate change. To better understand adaptive strategies of herbaceous species to increasing
58 drought and frost requires exploring further the role of hydraulic and mechanical traits using a
59 larger inter- and intra- specific range of species.

60

61 **Key-words:** cavitation resistance, *Dactylis glomerata* L., dehydration tolerance, drought survival,
62 embolism threshold, plant mortality, leaf fracture toughness, vessel wall thickness, stem anatomy,
63 intra-specific variability, perennial grass

64 INTRODUCTION

65 Higher temperatures due to climate change have led to more frequent extreme climate events
66 around the world (I.P.C.C., 2014, Orlowsky and Seneviratne, 2012, Seneviratne et al., 2014).
67 Extreme events, including intense droughts and heat waves, will become important drivers of
68 future ecosystem dynamics and function (Smith, 2011). Resilience, defined as the capacity of a
69 population or community to recover following disturbance, may therefore be the key to the
70 persistence of ecosystems (Hodgson et al., 2015; Nimmo et al., 2015).

71 Forests and grasslands are major terrestrial ecosystems, covering one-quarter and one-
72 third of the total land area, respectively. They provide many important ecosystem services
73 including timber and forage supply, carbon storage, and biodiversity preservation, but their
74 persistence is threatened under more frequent extreme climate events (Anderegg et al., 2013,
75 Pilgrim et al., 2010). Climate-induced forest dieback is an increasing global concern (Allen et al.,
76 2010) and major research efforts have been carried out to identify strategies for drought survival
77 among woody plant species (Pivovarov et al., 2015). In parallel, declines in grassland
78 productivity (Brookshire and Weaver, 2015) and long-term degradation from drought may
79 gradually become more common in the future (Ciais et al., 2005). Drought reduces plant
80 productivity and induces plant mortality in grasslands under extreme events (Griffin and
81 Hoffmann, 2012, Hodgkinson and Muller, 2005, Moran et al., 2014, Poirier et al., 2012, Van Peer
82 et al., 2001). Although the ability of woody species to survive intense drought was strongly
83 related to their embolism resistance i.e. the vulnerability to embolism (McDowell et al., 2008),
84 the traits and processes involved in the survival and recovery of herbaceous species after extreme
85 stress are still poorly characterized (Craine et al., 2013, Hoover et al., 2014).

86 Recent methodological developments have enabled **inter-specific** study of the vulnerability to
87 embolism in stems of herbaceous plants (Lens et al., 2013, Tixier et al., 2013), opening the door
88 for explorations of the role of hydraulic traits in stress survival of non-woody species. Building
89 on these recent results showing large interspecific variability in embolism resistance in grasses
90 distributed along an aridity gradient (Lens et al., 2016), the present study aimed to explore the
91 extent of intraspecific variability in the perennial grass *Dactylis glomerata* L. **This model species**
92 **was chosen since it has a large biogeographical distribution over Europe and Asia (Borrill, 1991)**
93 **and a high genetic variation within both populations and geographical origins (Lumaret, 1988,**
94 **Xie et al., 2010).**

95 Intraspecific variation with respect to embolism resistance is still poorly understood
96 (Martinez-Vilalta et al., 2009). In woody trees, phenotypic plasticity in resistance to embolism
97 was found to be low and buffered against environmental variation in a large range of *Pinus*
98 *pinaster* (Lamy et al., 2011, Lamy et al., 2014) but was not different across populations of *Fagus*
99 *sylvatica* (Herbette et al., 2010, Wortemann et al., 2011); however, not a single study has
100 assessed the intraspecific variation within an herbaceous grass species. Perennial species exhibit
101 various functional strategies to cope with moderate to severe drought (Pérez-Ramos et al., 2013,
102 Volaire, 2008, Zwicke et al., 2015), since they undergo many successive periods of stress and
103 recovery during their life cycle. Under severe drought, the basal aerial meristems and roots are
104 the key organs for plant survival and recovery, while most leaves of perennial grasses senesce to
105 reduce water loss (Volaire et al., 2009). As a result, no significant correlation was found between
106 vulnerability of leaf hydraulic conductance and whole-plant survival of drought in grasses
107 (Ocheltree et al., 2016). The role of hydraulic traits in herbaceous species should therefore be
108 analysed through a whole-plant perspective.

109 Survival in increasingly drought prone environments depends on a range of traits which include
110 sufficient mechanical strengthening to avoid irreversible effects of tissue dehydration. Embolism
111 resistant tree species are often characterized by a high wood density (Hacke et al., 2001, Lens et
112 al., 2013, Ogasa et al., 2013) although no strong evidence was that denser woods conferred
113 higher survival, or that the risk of embolism caused by wide conduits increased mortality (Russo
114 et al., 2010). In addition, increased tolerance to hydraulic dysfunction implies increased carbon
115 costs for leaf construction and water use (Nardini et al., 2012). Low embolism threshold values
116 were associated with narrower and denser vein conduits, increased thickness of conduit wall and
117 vein density and also with reduced leaf area (Nardini et al., 2012). In herbaceous species too,
118 *Arabidopsis* plants with increased wood formation in their stems were shown to be significantly
119 more embolism-resistant (Lens et al., 2013, Tixier et al., 2013) and grass species from the most
120 arid environments had more lignified stems compared to the vulnerable species native to wetter
121 habitats (Lens et al., 2016).

122 The main objective of this study is to identify relationships between embolism resistance, whole
123 plant dehydration survival, leaf traits, and stem anatomy amongst twelve populations of *Dactylis*
124 *glomerata* L originating from Scandinavia to Morocco. Given the contrasted biogeographic
125 origins of the studied populations, we expected that they differ strongly for hydraulic and

126 anatomical traits which would subsequently contribute to their dehydration tolerance. To
127 determine ecological responses of grassland species under climate extremes (Smith, 2011a), this
128 study therefore aims to (a) identify the role of embolism thresholds and associated traits for
129 drought survival and mechanical resistance in a perennial herbaceous species and (b) determine
130 the intra-specific variability of these traits as indicators of potential drought survival.

131

132 MATERIAL AND METHODS

133

134 *Plant Material*

135 Twelve populations of *Dactylis glomerata* L. (cocksfoot), including native populations and
136 cultivars, were selected from germplasm banks to be representative from three climatic regions:
137 four Mediterranean, four Temperate, and four Northern populations (Table 1). Cocksfoot (*D.*
138 *glomerata* L.) is a tetraploid ($2n=4x=28$) species from the genus *Dactylis* whose taxonomic
139 constitution is debated, but broadly consists of a complex of diploids and tetraploids (Borrill,
140 1991). The twelve populations tested were all verified to be tetraploid (Barre, pers. comm).

141

142 *Experimental sites and design*

143 Three pot-experiments were carried out to determine dehydration tolerance, embolism resistance
144 as well as anatomy and leaf traits, at the CEFÉ-CNRS research center in Montpellier, France
145 (43°6'N, 3°8'E). Seeds from each *D. glomerata* population were sown in seedling trays in
146 February 2013 and the adult plants (over one year old) were used for all subsequent experiment.

147 - Experiment 1 was carried out to determine plant dehydration tolerance measured as plant
148 survival rate in plants growing in short pots, after rewatering following an intense drought
149 (Volaire, 2018). Plants were transplanted into 4L, 18cm-diameter pots in May 2013. As
150 roots were equally limited in depth for all the populations in pots, plant drought survival
151 mirrored plants dehydration tolerance under a similar soil water availability (Volaire,
152 2008). It allowed us to discount the effect of rooting depth associated mainly with
153 dehydration avoidance (Pérez-Ramos et al., 2013, Volaire and Lelievre, 2001). For this
154 experiment, the quantity of dry substrate was measured to be identical for each of the 4L
155 pot so that the time dynamics of soil moisture could be accurately assessed and compared
156 simply by pot weight during the period of intense water deficit in summer 2014.

157 - Experiment 2 was carried out to measure embolism threshold and stem anatomy.
158 Individuals started from tillers taken from the adult plants sown in 2013 were transplanted
159 in September 2014 into 10L (24 cm-diameter) pots in order to obtain reproductive tillers
160 from spring 2015.

161 - Experiment 3 was carried out to measure biomechanical and leaf traits. Individuals taken
162 from the initial adult plants were transplanted into new 4L pots in September 2015.

163 Each 4L and 10L pot contained four clones from a single population, to ensure soil cover close to
164 field conditions, with 10 and 4 replicates for 4L pots and 10L pots respectively per population.

165 Pots were filled with a substrate composed of 65% local loamy-clay soil and 35% compost.

166 For all experiments, the plants were grown under an open-ended hoop-house and they were kept
167 fully irrigated and fertilized (40 U N after each cut) except during the imposition of the severe
168 drought in Experiment 1 during summer 2014. Pesticides were applied to prevent the
169 development of diseases caused by insects (aphids) or fungi (rust). For all experiments, pots were
170 distributed on a bench according to a full randomization design. In addition, the position of pots
171 was modified multiple times during the experiment to minimize placement effects.

172 In addition, plants of one of the Mediterranean populations (Kasbah), that was grown in a nearby
173 field in a 1.30 m deep loamy soil since 2011 (Barkaoui et al., 2016) and without irrigation nor
174 fertilization for two years, were used for stem sampling in order to assess the plasticity of
175 embolism resistance in one population. Twenty stems of this field-grown 'Kasbah' population
176 were collected in spring 2015 and processed for the measurement of stem embolism (see below).
177 No other measurement was carried out on these plants.

178 Mean temperatures were 9.5°C in winter, 16.6°C in spring and 24.2°C in summer, with 63%
179 mean air relative humidity and mean radiation of $800 \mu\text{mol m}^{-2} \text{s}^{-1}$ from spring to summer.

180

181 *Measurements*

182

183 *Dehydration tolerance under severe summer drought (Experiment 1)*

184 Plants of Experiment 1 were kept fully irrigated and fertilized until June 2014 when the severe
185 summer drought started to be imposed as follows. In June 2014, the soil water content was raised
186 to field capacity (60% soil water content for the substrate used) and irrigation was ceased
187 thereafter. Pots were weighed at frequent intervals (every 2-5 days) throughout the experiment to

188 monitor the decrease in soil water content. Once full leaf senescence was reached for all
189 populations, and when soil water content in the pots decreased to near 12% (27 to 35 days after
190 irrigation cessation), plants were harvested at 3cm from the soil surface and the pot was put back
191 under full irrigation (field capacity), thus ending the severe drought. All pots were re-watered
192 after they reached the same soil water content to ensure that plant survival corresponded to an
193 identical soil final dehydration for all populations irrespective of their growth potential. The
194 plants were irrigated for 15 days after which dehydration survival was measured. Plants that did
195 not produce new shoots after 15 days were counted as dead. The survival rate was measured as
196 the number of living plants divided by the total number of plants (plant survival; SURV %).

197
198 *Embolism Resistance* (Experiment 2)
199 In late spring 2015, three pots for each population were maintained under irrigated conditions and
200 allowed to flower. Stems were then collected and processed using the static centrifuge technique
201 (Alder et al., 1997). A negative pressure was applied to stem pieces in a standard centrifuge with
202 a custom-built, 26 cm rotor. Each stem segment was only spun once in the centrifuge. After
203 centrifugation, a 2 cm long segment containing a node was excised under water from the middle
204 of the stem segment. The segment was connected to the XYL'EM apparatus (Bronkhorst,
205 Montigny-les-Cormeilles, France) to measure its hydraulic conductance using a solution of 10
206 mM KCl and 1mM CaCl₂ in deionized ultrapure water. The sample was then resaturated with
207 pressurised water to determine the percentage of loss of conductance caused by embolised
208 conduits. For each pressure point in the vulnerability curve (VC), one to seven grass stems were
209 used with 14 to 36 pieces per curve (min-max), and one S-shaped curve per population was fitted
210 (Fig. S1) according to a sigmoid function (Pammenter and Vander Willigen, 1998). Embolism
211 resistance was expressed by the 50% or 88% percentage loss of hydraulic conductance (P₅₀, P₈₈).
212 P₅₀ is the metrics most used to compare hydraulic thresholds between species and populations
213 while P₈₈ was shown to be the embolism threshold leading to irreversible drought damage in
214 angiosperm trees (Urli et al., 2013)

215
216 *Stem anatomy*

217 To evaluate which stem characters contribute to the variation in embolism resistance observed
218 among the *D. glomerata* populations, light microscopy observations were performed on cross

219 sections of stems – previously used for the measurement of embolism resistance – at the level of
 220 internodes (from three stems per population). After vulnerability curve determination, samples
 221 were frozen until processed for anatomy. Samples were defrost in water, stored in 50% ethanol
 222 and then were embedded with LR White (hard grade, London Resin, UK) following (Hamann et
 223 al., 2011), sectioned with a rotary microtome (Leica RM 2265) equipped with disposable blades.
 224 Transverse sections of 4 μm were made, heat-attached to the slides, stained with toluidine blue,
 225 and mounted in Entellan®. The slides were observed with a Leica DM2500 light microscope
 226 equipped with a Leica DFC-425C digital camera (Leica Microscopes, Wetzlar, Germany). A
 227 range of stem anatomical characters (Lens et al., 2016), such as total stem area (A_S), area of
 228 lignified outer stem tissue (A_{LIG}), surface area of fibre wall (A_{FW}) and fibre lumen (A_{FL}),
 229 thickness of the metaxylem vessel wall (T_{VW}), hydraulically weighted diameter of vessels (D_{HV}),
 230 and traits derived from these measurements, such as the proportion of lignified tissue per total
 231 stem area (P_{LIG}), the proportion of cell wall per fiber (P_{CWF}), the proportion of total fiber wall
 232 area in lignified area ($P_{FWF} * A_{LIG}$), the proportion of total fiber wall in lignified area per stem
 233 area ($(P_{FWF} * A_{LIG}) : A_S$). Observations were made based on three individuals per population and at
 234 least 30 measurements per feature using ImageJ v 1.43 (National Institutes of Health, Bethesda,
 235 Maryland, USA).

236 The diameter of the vessels (D_V) was calculated based on the lumen area that was considered to
 237 be a circle following the equation:

238

$$239 \quad D_V = \sqrt{\frac{4A}{\pi}} \quad \text{Eqn (1)}$$

240

241 where D is the vessel diameter and A is the area of the vessel. The hydraulically weighted vessel
 242 diameter (D_{HV} ; weights diameters of vessels according to their hydraulic conductance) was
 243 calculated following the equation (Sperry et al., 1994):

244

$$245 \quad Dhv = \frac{\sum D^5}{\sum D^4} \quad \text{Eqn (2)}$$

246

247 where D is the vessel diameter as measured in eqn. (1).

248

249 *Leaf traits under full optimum conditions (Experiment 3)*

250 In grasses, leaf tissue density can be assessed both by leaf dry matter content (Cornelissen et al.,
251 2003) **which is a reasonable indirect measure of dry matter concentration (Shipley and Vu, 2002)**
252 and resistance of the leaf to mechanical fracture, that can be measured in a variety of ways (Ang
253 et al., 2008, Aranwela et al., 1999, Onoda et al., 2011). For experiment 3, in early spring (2016),
254 at the time before reproductive stage, one fully extended leaf was collected from one plant per pot
255 (four replicates). Fracture toughness was measured with a portable mechanical measuring device
256 (Instron® Inspec 2200) adapted for cutting with a single razor cut, following Ang et al (Ang et
257 al., 2008). Leaf blades were placed on the cutting platform across a gap 2 mm wide permitting
258 the passage of the diagonally angled blade in the middle of each leaf blade (single edged blade,
259 "Personna GEM, 0.23 mm thickness, Cat.# 71962). Leaf toughness, the energy required to shear
260 the leaf cross section in J m^{-2} was calculated after subtracting initial elastic loading and final
261 elastic unloading, as well as friction energy from the force distance curves. As in Ang et al.
262 (2008), the initial cut was followed by a second pass that measured just the friction force
263 generated by the blade against the cut surfaces of the leaf. Leaves were cut with a razor angle of
264 20° and at a crosshead speed of 0.1 mm s^{-1} . Leaf thickness and width were measured with a
265 binocular at the central vein of the leaf and at the widest part of the leaf respectively. The leaf
266 was then immediately placed into a tube with the cut end submerged in deionized water and
267 stored in the dark at 4°C for 24 h. The fresh and dry leaf biomass (g) were weighed to obtain leaf
268 dry matter content (LDMC; Leaf dry matter content is the oven-dry mass (mg) of a leaf divided
269 by its water-saturated fresh mass (g), expressed in mg g^{-1}) following (Perez-Harguindeguy et al.,
270 2013). This trait is used as a proxy for leaf density which is a trait more difficult to measure
271 (Shipley and Vu, 2002).

272

273 **Statistical analysis**

274 All statistical analyses were performed with Rstudio (version 0.98.501). The effect of
275 biogeographical origin was tested with anova ('lmtest' package) followed with Tukey's post hoc
276 tests. A principal component analysis (PCA; 'ade4' package) was performed on the twelve
277 variables showing significant differences between origins, scaled and centered, to visualize their
278 covariations. The four bioclimatic variables (www.worldclim.org) chosen to characterize the
279 drought and frost occurrence in the sites of origins of the populations (annual precipitations,

280 precipitations of the driest quarter, aridity index (Trabucco et al., 2008) and minimum
281 temperature of the coldest months) were used as supplementary variables in the PCA. Pairwise
282 correlations between all measured traits for each season were evaluated using Pearson's method
283 ('Hmisc' package). We tested the relationships between key-variables using simple linear
284 regressions.

285

286

287 **RESULTS**

288 Across the twelve populations of *D. glomerata* studied, P_{50} ranged from -3.06 to -6.36 MPa (Fig.
289 1a), while P_{88} ranged from -5.06 to -11.6 MPa (Fig. 1b). The Mediterranean populations had
290 significantly lower P_{50} (-5.8 and -10.6 MPa respectively) than the Temperate populations (-3.3
291 and -6.1 MPa respectively), while the Northern populations showed intermediate embolism
292 thresholds (Table 2). In addition, P_{50} of the Mediterranean population Kasbah significantly
293 differed according to the growing environment (Fig. 2), either in pots ($P_{50} = -6.05 \pm 0.31$) or in the
294 field ($P_{50} = -5.21 \pm 0.29$).

295 P_{50} and P_{88} measured on potted irrigated plants were both highly and more correlated ($r=0.88$ and
296 0.89 , $P<0.001$, Table 3) with the mean precipitation of the driest quarter of the year than with the
297 annual precipitation of the origin site. Both embolism thresholds were, however, less strongly
298 correlated with the aridity index of the site of origin ($r=0.60$, $P=0.039$). The mean aridity index
299 was 1.20 for Northern populations (less arid) compared to 0.82 (more arid) for Temperate
300 populations despite of the greater rainfall in Temperate areas. In addition, no correlation was
301 detected between cavitation resistance and minimum temperatures of the coldest months of the
302 sites studied.

303 Plant dehydration survival in pots was significantly greater (Table 2) for the Mediterranean
304 populations (58%) than for the group of Northern and Temperate ones (7 to 13%).

305 Although leaf thickness did not discriminate the populations according to their origin, leaf width
306 almost doubled with increasing latitude from on average 4.3 mm in Mediterranean populations to
307 8 mm in Northern ones. Mediterranean populations also had a higher LDMC (20% greater) and a
308 very significantly greater leaf fracture toughness (2.5 fold greater) than the Northern and
309 Temperate populations.

310 The Mediterranean populations were distinguished from the others by having thinner stems with
 311 a higher fibre wall proportion in lignified area per stem (Table 2). The traits for Northern and
 312 Temperate populations were mostly not significantly different although the proportion of total
 313 fibre wall area in lignified area was greater in Northern populations. The thickness of metaxylem
 314 vessel wall highly discriminated all origins and spanned from 1.4, 1.2 and 1.1 μm in
 315 Mediterranean Northern and Temperate populations respectively (Table 2).

316 P_{88} was highly and better correlated to dehydration survival ($r=-0.79$, Table 3, Fig. 4a) and leaf
 317 traits such as fracture toughness ($r=-0.78$, Table 3, Fig. 4b) and LDMC ($r=-0.82$, Table 3, Fig.
 318 3c), than P_{50} . A highly significant correlation was found between both embolism thresholds
 319 (Table 3, Fig. 5, 6) and the thickness of metaxylem vessel wall in stems (T_{VW} : $r=-0.92$ for P_{50}).
 320 Notably no other stem anatomical trait was correlated with either embolism thresholds. In
 321 contrast, plant dehydration survival was correlated to the total stem area (A_S : $r=-0.66$) and
 322 therefore with the area of lignified outer stem tissue (A_{LIG} : $r=-0.61$) as well as with the proportion
 323 of fibre wall in lignified area over the entire stem area ($(P_{FW} \cdot A_{LIG}) : A_S$ $r=0.59$).

324 Overall, the covariation between LDMC, leaf fracture toughness, embolism thresholds and
 325 dehydration tolerance was very significant (Table 3). Unexpectedly, leaf toughness was not
 326 correlated with any trait associated with stem lignification over the entire data set. However, a
 327 PCA (Fig. 7a, b) clearly illustrates that high embolism resistance and dehydration survival
 328 associated with small and more lignified lamina and lignified stems discriminate the group of
 329 Mediterranean populations. In contrast, Temperate and Northern populations were characterised
 330 by low dehydration survival and embolism resistance, bigger leaves and stems with thicker-
 331 walled fibres especially for Northern populations.

332

333 **DISCUSSION**

334 This study is the first to explore intraspecific variability of embolism resistance, whole plant
 335 dehydration tolerance and associated traits in an herbaceous grass species. Our results show a
 336 huge range of embolism thresholds amongst the twelve *D. glomerata* populations studied,
 337 reflecting its biogeographical distribution and corresponding with some key traits.

338 *Large intraspecific variability in cocksfoot*

339 If we compare the interspecific range for P_{50} (Lens et al., 2016) observed within 18 Poaceae
340 species native to similar mesic and xeric environments (-3.2 to -7.5 MPa) with the intraspecific
341 variability in *D. glomerata* found here (-3.06 to -6.36 MPa), the overlap is striking. Only *Stipa*
342 *pennata*, a grass native to arid rangelands exhibited a lower P_{50} (-7.5 MPa) compared to the
343 Mediterranean populations of *D. glomerata*. In addition, this intra-specific variation is much
344 greater than that reported for a dicot herbaceous perennial *Solidago Canadensis* sampled from
345 dry to humid sites and showing P_{50} ranging from -2.37 to -3.08 MPa (Nolf et al., 2014). The
346 intraspecific range in P_{50} variation within cocksfoot matches with most of the woody species
347 studied in angiosperms (-2 to -4 MPa) and gymnosperms (-4 to -8 MPa), although far from the
348 extreme embolism threshold (-18.8 MPa) found so far in the gymnosperm *Callitris tuberculata*
349 (Larter et al., 2015). This intraspecific variability is strikingly large in comparison with former
350 results on woody species. For instance, across 17 populations of *Fagus sylvatica* originating from
351 sites in a large European latitudinal gradient with annual precipitations spanning from 563 to
352 1362 mm, P_{50} was relatively constant from -2.8 to -3.2 MPa (Wortemann et al., 2011). No
353 significant differences in embolism resistance (-3.9 to -4.1 MPa) was found either across 24
354 populations of *Pinus pinaster* from Northern African and French sites (Lamy et al., 2011).
355 Nevertheless, plasticity in vulnerability of embolism (from -3 to -6 MPa) was shown in the very
356 dry limits of *Pinus canariensis* under a strong aridity gradient (Lopez et al., 2016). **It could be**
357 **interesting to explore whether** the high embolism resistance of *D. glomerata* from the most arid
358 sites ($P_{50} < -5$ MPa) could be ascribed to a shallower rooting system than woody species by
359 analogy with shrubs that were found to be more embolism tolerant than trees in the same habitat,
360 probably reflecting differences in rooting depth and drought tolerance (Brendel & Cochard,
361 2011). The remarkable range in embolism resistance could be added to the list of factors
362 contributing to explain why cocksfoot covers a greater span of biomes than that of most woody
363 species.

364 Our intraspecific comparison is based on plants grown in similar optimum pot conditions for two
365 years, underlining that the large differences within populations are genetically determined. **Yet,**
366 **our results also show a high plasticity of embolism threshold since for the tested Kasbah**
367 **population, P_{50} significantly varied between potted and field-grown plants of the same**
368 **population. Many factors differ such as plant age at sampling date as well as many biotic and**
369 **abiotic factors (soil depth, plant density...) between these two growing conditions for the same**

370 population. However, it may be hypothesized that a less negative P_{50} of field-growing plants may
371 be ascribed in particular to a deeper rooting system (130 cm in the field versus 20 cm in pots)
372 which enhances dehydration avoidance versus dehydration tolerance and may have impacted
373 hydraulic adaptations (Scholz et al., 2012). It could be interesting to measure embolism resistance
374 in the same populations of cocksfoot growing all in a similar deep soil in order to analyse the
375 impact of the expression of full rooting depth on embolism thresholds. However, it can therefore
376 be assumed that if sampling was done using populations growing in their natural environment,
377 our results would have probably generated stronger genotype x environments interactions, and
378 even further increased the range of intra-specific variability in embolism thresholds. Significant
379 site x population interactions were for instance found in beech (Herbette et al., 2010, Wortemann
380 et al., 2011) underlining the high phenotypic plasticity of embolism resistance. Further work is
381 necessary to assess intra-specific P_{50} and P_{88} plasticity as a response to environments (e.g. soil
382 depth, soil nutrients and water availability).

383 *Relationships between embolism resistance and environmental limiting factors*

384 Our results show a correlation between the annual and driest quarter precipitations of the sites
385 studied with embolism thresholds measured on plants grown in a common garden under optimum
386 irrigation. As a general pattern, dehydration survival in pots was greater for the populations
387 coming from the driest sites with P_{50} and P_{88} more negative than -4.50 and -9.3 MPa respectively.
388 Plant mortality under severe drought when the resource is limited (shallow soils) can be then
389 predicted by the hydraulic failure model assuming that a significant loss of conductivity can
390 trigger plant mortality (McDowell et al., 2013). Our results are in line with a meta-analysis of tree
391 species showing that hydraulic traits capture key mechanisms determining tree death (Anderegg
392 et al., 2016). However, embolism resistance was less strongly correlated with the aridity index of
393 the site of origin in contradiction to a former study dealing with a range of Temperate to semi-
394 arid grass species that showed a strong correlation between P_{50} and aridity (Lens et al., 2016). It
395 would be interesting to test whether this interspecific correlation would hold when including
396 herbaceous species from the highest latitudes. In *D. glomerata*, Northern populations showed a
397 trend to more negative P_{50} values than Temperate populations and are more frost tolerant,
398 suggesting that the Northern populations could have also developed greater embolism resistance
399 as a mechanism of frost tolerance (Lens et al., 2013). Both drought and freezing-thawing of stems

400 induce a loss of hydraulic conductivity in woody plants (Feng et al., 2015). Frost ‘drought’ had a
401 large effect on plant water transport, adaptations in hydraulic safety and related anatomical
402 parameters (Mayr et al., 2006) and the propensity for freeze induced embolism increases as
403 conduit diameter increases (Zanne et al., 2014). The relationships between recovery from winter
404 embolism and anatomical vessel traits also varied according to functional groups of tree species
405 (Niu et al., 2017). The role of hydraulic dysfunction under frost remains to be explored in
406 herbaceous species.

407 *Tissue lignification and embolism resistance*

408 In pseudostems, partly constituted with leaf sheaths, we found a strong positive correlation
409 between the thickness of the metaxylem vessel wall and embolism resistance while lignification
410 characters are not correlated with P_{50} nor P_{88} . Despite a significant positive correlation (Lens et
411 al., 2016) between embolism resistance and stem lignification amongst 19 grass species (in terms
412 of the thickness of the fibre walls and the relative proportion of lignified tissue per stem area),
413 this trade-off is lacking within populations of *D. glomerata*. This lack of correlation is
414 noteworthy given the huge range in P_{50} variation within cocksfoot. There are slight indications
415 that stem lignification may contribute to differences in embolism resistance. For instance, Figure
416 6ab shows that the proportion of lignified area per stem area clusters with other key traits for the
417 embolism resistant Mediterranean populations. Furthermore, the two most resistant
418 Mediterranean populations have more lignified fibres inside and surrounding the vascular bundles
419 (Fig. S2) compared to the other cocksfoot populations studied. However, this character is difficult
420 to quantify and seemingly shows at random variation amongst the remaining populations. Since
421 we have sectioned internodes, we had to restrict ourselves to measurements of solitary vessels. It
422 is plausible that our observations also reflect a correlation between more negative P_{50}/P_{88} values
423 and increased inter-vessel wall thickness compared to vessel diameter (so called thickness-to-
424 span ratio) in case of vessel multiples that are much more present in the nodal parts of the stem.
425 Vessel wall reinforcement was suggested to be required for embolism resistance in order to
426 prevent wall implosion when xylem pressure is very negative (Hacke et al., 2001). However,
427 vessels have never been observed to implode in stems, probably because embolism events occur
428 before the critical vessel implosion threshold is reached. Whatever the exact mechanism(s)
429 operating during xylem embolism dysfunction in *D. glomerata*, it appears that plants with more

430 fracture resistant leaves have stems that are more resistant to embolism than plants with less
431 resistant leaves. In addition to possibly reducing the probability of lethal embolism (Hacke and
432 Sperry, 2001), mechanically robust leaf architecture will also play a role in mechanically
433 supporting water-stressed leaves and protecting leaf meristems as discussed below. This perhaps
434 at least partly explains the tendency towards narrower, more compact, denser, tougher leaves in
435 the Mediterranean ecotypes that are more likely exposed to more drought conditions. Of course
436 leaf lamina traits are anatomically and functionally grade into the more proximal leaf sheath
437 (Bell, 1991) which is also important for mechanical robustness and resistance to dehydration of
438 the self-supporting meristems of *D. glomerata*.

439 *Drought survival in perennial herbaceous species: hydraulic leaf traits may not be crucial*

440 A general tenet of ecophysiology of drought adaptation is based on the assumption of a strong
441 linkage between hydraulic and stomatal traits especially in woody species (Skelton et al., 2015,
442 Bartlett et al., 2016) and also in herbaceous species (Holloway-Phillips and Brodribb, 2011, Lens
443 et al., 2016, Nardini et al., 2012). Regarding the extreme case of drought plant mortality in both
444 annual and perennial grasses, basal leaf meristems are the tissues surviving longer and able to
445 regenerate when the mature blades are dead (Van Peer et al., 2004). These leaf meristems
446 enclosed in mature leaf sheaths, are protected from intense evaporation and can survive more
447 intense water deficits than older tissues (Barlow et al., 1980, Munns et al., 1979). As they
448 accumulate carbohydrates such as fructans (Schnyder and Nelson, 1989, Volaire et al., 1998),
449 apices exhibit the greatest osmotic adjustment relative to other tissues during drought (Matsuda
450 and Riazi, 1981, West et al., 1990). In addition, the strategy of summer dormancy (Gillepsie and
451 Volaire, 2017, Volaire and Norton, 2006) found in some species and populations of perennial
452 species, triggers leaf senescence and meristem dehydration to reduce plant metabolism and
453 consequently enhance drought survival. Our results show that the Mediterranean populations of
454 *D. glomerata* which are the most summer dormant, ‘sacrifice’ their lamina in summer
455 (irrespectively of the water supply) but also have the highest embolism resistance and
456 dehydration tolerance. In frost-prone areas too, perennial herbaceous species avoid freezing by
457 senescing aboveground tissues and overwintering as underground storage organs (Zanne et al.,
458 2014). For dehydration tolerance under frost or drought (Gillepsie and Volaire, 2017), these
459 model species can then raise the relevance of the paradigm of hydraulic safety margins in leaves

460 for stress survival. The leaf (lamina)-level drought tolerance (Nardini and Luglio, 2014,
461 Ocheltree et al., 2016) may not be the key-level to explore drought survival in all species.
462 However, studies linking hydraulic conductivity and lamina anatomy (Scoffoni et al., 2017) could
463 be fruitful provided they take into account the continuum between expanded blades, sheaths and
464 meristematic tissues. The co-variation between hydraulic traits and anatomical traits in meristems
465 but also in root tissues should be explored further in grasses.

466 In conclusion, this study investigated dehydration survival amongst twelve populations of *D.*
467 *glomerata* from a latitudinal gradient from Mediterranean to Scandinavian areas, representing
468 only a restricted range of its much larger biogeographical distribution. *D. glomerata* represents an
469 ‘ideal’ species for analyzing the interactions genotype x environment because of its
470 biogeographical distribution is huge and spans over Europe and Asia (Borrill, 1991) and its
471 phylogenetic relationships between populations are being investigated and show high genetic
472 variation within both populations and geographical origins (Lumaret, 1988, Xie et al., 2010).
473 Compared to woody species, the large degree of intra-specific variability in dehydration tolerance
474 and embolism resistance within cocksfoot has consequences for its sensitivity to climate shifts,
475 possible migrations and future biogeographical distribution (Anderegg, 2015). In addition, given
476 that the measurement of plant drought survival is experimentally difficult to standardize, our
477 study demonstrating the key role of embolism threshold and tissue lignification as proxy for
478 dehydration tolerance provide highly relevant knowledge to screen genetic resources and new
479 breeding material.

480 Finally, this study calls for extending the numerous studies on mortality factors and lethal
481 embolism thresholds in woody species to a large range of herbaceous species, both at inter and
482 intra-specific levels, under contrasting environmental constraints.

483 **ACKNOWLEDGMENTS**

484 Thank to Pascal Chapon and Mathieu Ogier for excellent technical assistance, the ‘Terrain
485 d’expérience platform’ and the ‘Plateforme d’Analyses Chimiques en Ecologie’ both at CEFÉ-
486 CNRS (Labex CEMEB). Thanks to Mark Norton, Jeffrey Clary, Rajae Kallida, N. Shaimi, A.
487 Palmé, S. Fourtier, B. Reid, G. Testa, and V. Copani, for providing seeds. This study was
488 supported by the European Research Council (ERC) Starting Grant Project ‘Ecophysiological and

489 biophysical constraints on domestication in crop plants' [Grant ERC-StG-2014-639706-
490 CONSTRAINTS], and by the Agence National de la Recherche [PRAISE project, ANR-13-
491 BIOADAPT-0015]. Pauline Bristiel was financially supported by INRA (EA department).

492

493

494

495 LITERATURE CITED

496

497 **Alder NN, Pockman WT, Sperry JS, Nuismer S. 1997.** Use of centrifugal force in
498 the study of xylem cavitation. *Journal of Experimental Botany*, **48**: 665-674.

499 **Anderegg WRL. 2015.** Spatial and temporal variation in plant hydraulic traits and
500 their relevance for climate change impacts on vegetation. *New Phytologist*,
501 **205**: 1008-1014.

502 **Anderegg WRL, Kane JM, Anderegg LDL. 2013.** Consequences of widespread
503 tree Mortality triggered by drought and temperature stress. *Nature Climate*
504 *Change*, **3**: 30-36.

505 **Anderegg WRL, Klein T, Bartlett M, Sack L, Pellegrini AFA, Choat B, Jansen**
506 **S. 2016.** Meta-analysis reveals that hydraulic traits explain cross-species
507 patterns of drought-induced tree mortality across the globe. *Proceedings of*
508 *the National Academy of Sciences of the United States of America*, **113**:
509 5024-5029.

510 **Ang KY, Lucas PW, Tan HTW. 2008.** Novel way of measuring the fracture
511 toughness of leaves and other thin films using a single inclined razor blade.
512 *New Phytologist*, **177**: 830-837.

513 **Aranwela N, Sanson G, Read J. 1999.** Methods of assessing leaf fracture
514 properties. *New Phytologist*, **144**: 369-383.

515 **Barkaoui K, Roumet C, Volaire F. 2016.** Mean root trait more than root trait
516 diversity determines drought resilience in native and cultivated Mediterranean
517 grass mixtures. *Agriculture Ecosystems & Environment*, **231**: 122-132.

518 **Barlow EWR, Munns RE, Brady DJ. 1980.** Drought responses of apical
519 meristems. In: Turner NC, Kramer PJ, eds. *Adaptation of Plants to Water and*
520 *High Temperature Stress*. New York, USA: John Wiley & Sons.

521 **Bartlett M, Klein T, Jansen S, Choat B, Sack L. 2016.** The correlations and
522 sequence of plant stomatal, hydraulic and wilting responses to drought.
523 *Proceedings of the National Academy of Sciences of the United States of*
524 *America*, **113**: 13098-13103.

525 **Bell A. 1991.** *Plant Form*. Portland - London: Timber Press.

526 **Borrill M. 1991.** Evolution and genetic resources in cocksfoot. Amsterdam,
527 Netherlands: Elsevier Science Publishers.

528 **Brookshire ENJ, Weaver T. 2015.** Long-term decline in grassland productivity
529 driven by increasing dryness. *Nature Communications*, **6**.

530 **Ciais P, Reichstein M, Viovy N, Granier A, Ogee J, Allard V, Aubinet M,**
531 **Buchmann N, Bernhofer C, Carrara A, Chevallier F, De Noblet N,**
532 **Friend AD, Friedlingstein P, Grunwald T, Heinesch B, Keronen P,**
533 **Knohl A, Krinner G, Loustau D, Manca G, Matteucci G, Miglietta F,**
534 **Ourcival JM, Papale D, Pilegaard K, Rambal S, Seufert G, Soussana JF,**
535 **Sanz MJ, Schulze ED, Vesala T, Valentini R. 2005.** Europe-wide reduction
536 in primary productivity caused by the heat and drought in 2003. *Nature*,
537 **437**: 529-533.

538 **Craine JM, Ocheltree TW, Nippert JB, Towne EG, Skibbe AM, Kembel SW,**
539 **Fargione JE. 2013.** Global diversity of drought tolerance and grassland
540 climate-change resilience. *Nature Climate Change*, **3**: 63-67.

- 541 **Feng F, Ding F, Tyree MT. 2015.** Investigations Concerning Cavitation and Frost
542 Fatigue in Clonal 84K Poplar Using High-Resolution Cavitron Measurements.
543 *Plant Physiology*, **168**: 144-U850.
- 544 **Gillepsie L, Volaire F. 2017.** Are winter and summer dormancy symmetrical
545 seasonal adaptive strategies? The case of temperate herbaceous perennials.
546 *Annals of Botany*, **119**: 311-323.
- 547 **Griffin PC, Hoffmann AA. 2012.** Mortality of Australian alpine grasses (*Poa* spp.)
548 after drought: species differences and ecological patterns. *Journal of Plant*
549 *Ecology*, **5**: 121-133.
- 550 **Hacke U, Sperry J. 2001.** Functional and ecological xylem anatomy. *Perspectives*
551 *in Plant Ecology, Evolution and Systematics*, **4**: 97-115.
- 552 **Hacke UG, Sperry JS, Pockman WT, Davis SD, McCulloch KA. 2001.** Trends in
553 wood density and structure are linked to prevention of xylem implosion by
554 negative pressure. *Oecologia*, **126**: 457-461.
- 555 **Hamann TD, Smets E, Lens F. 2011.** A comparison of paraffin and resin-based
556 techniques used in bark anatomy. *Taxon*, **60**: 841-851.
- 557 **Herbette S, Wortemann R, Awad H, Huc R, Cochard H, Barigah TS. 2010.**
558 Insights into xylem vulnerability to cavitation in *Fagus sylvatica* L.:
559 phenotypic and environmental sources of variability. *Tree Physiology*, **30**:
560 1448-1455.
- 561 **Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. 2005.** Very high
562 resolution interpolated climate surfaces for global land areas. *International*
563 *Journal of Climatology*, **25**: 1965-1978.
- 564 **Hodgkinson KC, Muller WJ. 2005.** Death model for tussock perennial grasses: a
565 rainfall threshold for survival and evidence for landscape control of death in
566 drought. *Rangeland Journal*, **27**: 105-115.
- 567 **Holloway-Phillips MM, Brodribb TJ. 2011.** Minimum hydraulic safety leads to
568 maximum water-use efficiency in a forage grass. *Plant Cell and Environment*,
569 **34**: 302-313.
- 570 **Hoover DL, Knapp AK, Smith MD. 2014.** Resistance and resilience of a grassland
571 ecosystem to climate extremes. *Ecology*, **95**: 2646-2656.
- 572 **I.P.C.C. 2014.** International Panel of Climatic changes. Fifth assessment report
573 (AR4).
- 574 **Lamy JB, Bouffier L, Burlett R, Plomion C, Cochard H, Delzon S. 2011.**
575 Uniform Selection as a Primary Force Reducing Population Genetic
576 Differentiation of Cavitation Resistance across a Species Range. *Plos One*, **6**.
- 577 **Lamy JB, Delzon S, Bouche PS, Alia R, Vendramin GG, Cochard H, Plomion**
578 **C. 2014.** Limited genetic variability and phenotypic plasticity detected for
579 cavitation resistance in a Mediterranean pine. *New Phytologist*, **201**: 874-
580 886.
- 581 **Larter M, Brodribb TJ, Pfautsch S, Burlett R, Cochard H, Delzon S. 2015.**
582 Extreme Aridity Pushes Trees to Their Physical Limits. *Plant Physiology*, **168**:
583 804-+.
- 584 **Lens F, Picon-Cochard C, Delmas C, Signarbieux C, Buttler A, Cochard H,**
585 **Jansen S, Chauvin T, Chacon Doria L, del Arco M, Delzon S. 2016.**
586 Herbaceous Angiosperms Are Not More Vulnerable to Drought-Induced
587 Embolism Than Angiosperm Trees. *Plant Physiology*, **172**: 661-667.

- 588 **Lens F, Tixier A, Cochard H, Sperry JS, Jansen S, Herbette S. 2013.** Embolism
589 resistance as a key mechanism to understand adaptive plant strategies.
590 *Current Opinion in Plant Biology*, **16**: 287-292.
- 591 **Lopez R, Cano FJ, Choat B, Cochard H, Gil L. 2016.** Plasticity in Vulnerability to
592 Cavitation of *Pinus canariensis* Occurs Only at the Driest End of an Aridity
593 Gradient. *Frontiers in Plant Science*, **7**.
- 594 **Lumaret R. 1988.** Cytology, genetics, and evolution in the genus *Dactylis*. *Critical*
595 *Reviews in Plant Sciences*, **7**: 55-91.
- 596 **Martinez-Vilalta J, Cochard H, Mencuccini M, Sterck F, Herrero A, Korhonen**
597 **JFJ, Llorens P, Nikinmaa E, Nole A, Poyatos R, Ripullone F, Sass-**
598 **Klaassen U, Zweifel R. 2009.** Hydraulic adjustment of Scots pine across
599 Europe. *New Phytologist*, **184**: 353-364.
- 600 **Matsuda K, Riazi A. 1981.** Stress-induced osmotic adjustment in growing regions
601 of barley leaves. *Plant Physiology*, **68**: 571-576.
- 602 **Mayr S, Hacke U, Schmid P, Schwienbacher F, Gruber A. 2006.** Frost drought
603 in conifers at the alpine timberline: Xylem dysfunction and adaptations.
604 *Ecology*, **87**: 3175-3185.
- 605 **McDowell N, Pockman WT, Allen CD, Breshears DD, Cobb N, Kolb T, Plaut J,**
606 **Sperry J, West A, Williams DG, Yezzer EA. 2008.** Mechanisms of plant
607 survival and mortality during drought: why do some plants survive while
608 others succumb to drought? *New Phytologist*, **178**: 719-739.
- 609 **McDowell NG, Fisher RA, Xu CG, Domec JC, Holtta T, Mackay DS, Sperry JS,**
610 **Boutz A, Dickman L, Gehres N, Limousin JM, Macalady A, Martinez-**
611 **Vilalta J, Mencuccini M, Plaut JA, Ogee J, Pangle RE, Rasse DP, Ryan**
612 **MG, Sevanto S, Waring RH, Williams AP, Yezzer EA, Pockman WT.**
613 **2013.** Evaluating theories of drought-induced vegetation mortality using a
614 multimodel-experiment framework. *New Phytologist*, **200**: 304-321.
- 615 **Moran MS, Ponce-Campos GE, Huete A, McClaran MP, Zhang YG,**
616 **Hamerlynck EP, Augustine DJ, Gunter SA, Kitchen SG, Peters DPC,**
617 **Starks PJ, Hernandez M. 2014.** Functional response of U.S. grasslands to
618 the early 21st-century drought. *Ecology*, **95**: 2121-2133.
- 619 **Munns R, Brady CJ, Barlow EWR. 1979.** Solute accumulation in the apex and
620 leaves of wheat during water-stress. *Australian Journal of Plant Physiology*,
621 **6**: 379-389.
- 622 **Nardini A, Luglio J. 2014.** Leaf hydraulic capacity and drought vulnerability:
623 possible trade-offs and correlations with climate across three major biomes.
624 *Functional Ecology*, **28**: 810-818.
- 625 **Nardini A, Peda G, La Rocca N. 2012.** Trade-offs between leaf hydraulic capacity
626 and drought vulnerability: morpho-anatomical bases, carbon costs and
627 ecological consequences. *New Phytologist*, **196**: 788-798.
- 628 **Niu C-Y, Meinzer FC, Hao G-Y. 2017.** Divergence in strategies for coping with
629 winter embolism among co-occurring temperate tree species: the role of
630 positive xylem pressure, wood type and tree stature. *Functional Ecology (In*
631 *press)*.
- 632 **Nolf M, Pagitz K, Mayr S. 2014.** Physiological acclimation to drought stress in
633 *Solidago canadensis*. *Physiologia Plantarum*, **150**: 529-539.
- 634 **Ocheltree TW, Nippert JB, Prasad PVV. 2016.** A safety vs efficiency trade-off
635 identified in the hydraulic pathway of grass leaves is decoupled from

- 636 photosynthesis, stomatal conductance and precipitation. *New Phytologist*,
 637 **210**: 97-107.
- 638 **Ogasa M, Miki NH, Murakami Y, Yoshikawa K. 2013.** Recovery performance in
 639 xylem hydraulic conductivity is correlated with cavitation resistance for
 640 temperate deciduous tree species. *Tree Physiology*, **33**: 335-344.
- 641 **Onoda Y, Westoby M, Adler P, Choong A, Clissold F, Cornelissen J, Díaz S,**
 642 **Dominy N, Elgart A, Enrico L. 2011.** Global patterns of leaf mechanical
 643 properties. *Ecology letters*, **14**: 301-312.
- 644 **Orlowsky B, Seneviratne SI. 2012.** Global changes in extreme events: regional
 645 and seasonal dimension. *Climatic Change*, **110**: 669-696.
- 646 **Pammenter NW, Vander Willigen C. 1998.** A mathematical and statistical
 647 analysis of the curves illustrating vulnerability of xylem to cavitation. *Tree*
 648 *Physiology*, **18**: 589-593.
- 649 **Perez-Harguindeguy N, Diaz S, Garnier E, Lavorel S, Poorter H,**
 650 **Jaureguiberry P, Bret-Harte MS, Cornwell WK, Craine JM, Gurvich DE,**
 651 **Urcelay C, Veneklaas EJ, Reich PB, Poorter L, Wright IJ, Ray P, Enrico**
 652 **L, Pausas JG, de Vos AC, Buchmann N, Funes G, Quetier F, Hodgson**
 653 **JG, Thompson K, Morgan HD, ter Steege H, van der Heijden MGA, Sack**
 654 **L, Blonder B, Poschlod P, Vaieretti MV, Conti G, Staver AC, Aquino S,**
 655 **Cornelissen JHC. 2013.** New handbook for standardised measurement of
 656 plant functional traits worldwide. *Australian Journal of Botany*, **61**: 167-234.
- 657 **Pérez-Ramos IM, Volaire F, Fattet M, Blanchard A, Roumet C. 2013.**
 658 Tradeoffs between functional strategies for resource-use and drought-survival
 659 in Mediterranean rangeland species. *Environmental and Experimental Botany*,
 660 **87**: 126-136.
- 661 **Pilgrim ES, Macleod CJA, Blackwell MSA, Bol R, Hogan DV, Chadwick DR,**
 662 **Cardenas L, Misselbrook TH, Haygarth PM, Brazier RE, Hobbs P,**
 663 **Hodgson C, Jarvis S, Dungait J, Murray PJ, Firbank LG. 2010.**
 664 Interactions among agricultural production and other ecosystem services
 665 delivered from European temperate grassland systems. In: Sparks DL, ed.
 666 *Advances in Agronomy, Vol 109*. San Diego: Elsevier Academic Press Inc.
- 667 **Pivovarov AL, Pasquini S, M.E. DG, Alstad KP, Stemke JS, Santiago LS.**
 668 **2015.** Multiple strategies for drought survival among woody plant species.
 669 *Functional Ecology*, **30**: 1-10.
- 670 **Poirier M, Durand JL, Volaire F. 2012.** Persistence and production of perennial
 671 grasses under water deficits and extreme temperatures: importance of
 672 intraspecific vs. interspecific variability. *Global Change Biology*, **18**: 3632-
 673 3646.
- 674 **Russo SE, Jenkins KL, Wisser SK, Uriarte M, Duncan RP, Coomes DA. 2010.**
 675 Interspecific relationships among growth, mortality and xylem traits of woody
 676 species from New Zealand. *Functional Ecology*, **24**: 253-262.
- 677 **Schnyder H, Nelson CJ. 1989.** Growth rates and assimilate partitioning in the
 678 elongation zone of tall fescue leaf blades at high and low irradiance. *Plant*
 679 *Physiology*, **90**: 1201-1206.
- 680 **Scholz FG, Bucci SJ, Arias N, Meinzer FC, Goldstein G. 2012.** Osmotic and
 681 elastic adjustments in cold desert shrubs differing in rooting depth: coping
 682 with drought and subzero temperatures. *Oecologia*, **170**: 885-897.

- 683 **Scoffoni C, Albuquerque C, Brodersen CR, Townes SV, John GP, Cochard H,**
684 **Buckley TN, McElrone AJ, Sack L. 2017.** Leaf vein xylem conduit diameter
685 influences susceptibility to embolism and hydraulic decline. *New Phytologist*,
686 **213**: 1076-1092.
- 687 **Seneviratne SI, Donat MG, Mueller B, Alexander LV. 2014.** No pause in the
688 increase of hot temperature extremes (vol 4, pg 161, 2014). *Nature Climate*
689 *Change*, **4**: 320-320.
- 690 **Shipley B, Vu TT. 2002.** Dry matter content as a measure of dry matter
691 concentration in plants and their parts. *New Phytologist*, **153**: 359-364.
- 692 **Skelton RP, West AG, Dawson TE. 2015.** Predicting plant vulnerability to drought
693 in biodiverse regions using functional traits. *Proceedings of the National*
694 *Academy of Sciences of the United States of America*, **112**: 5744-5749.
- 695 **Smith MD. 2011.** The ecological role of climate extremes: current understanding
696 and future prospects. *Journal of Ecology*, **99**: 651-655.
- 697 **Sperry JS, Nichols KL, Sullivan JEM, Eastlack SE. 1994.** Xylem embolism in
698 ring-porous, diffuse-porous, and coniferous trees of northern Utah and
699 interior Alaska. *Ecology* **75**: 1736-1752.
- 700 **Tixier A, Cochard H, Badel E, Dusotoit-Coucaud A, Jansen S, Herbette S.**
701 **2013.** Arabidopsis thaliana as a model species for xylem hydraulics: does
702 size matter? *Journal of Experimental Botany*, **64**: 2295-2305.
- 703 **Trabucco A, Zomer RJ, Bossio DA, van Straaten O, Verchot LV. 2008.** Climate
704 change mitigation through afforestation/reforestation: A global analysis of
705 hydrologic impacts with four case studies. *Agriculture Ecosystems &*
706 *Environment*, **126**: 81-97.
- 707 **Urli M, Porte AJ, Cochard H, Guengant Y, Burlett R, Delzon S. 2013.** Xylem
708 embolism threshold for catastrophic hydraulic failure in angiosperm trees.
709 *Tree Physiology*, **33**: 672-683.
- 710 **Van Peer L, Nijs I, Bogaert J, Verelst I, Reheul D. 2001.** Survival, gap
711 formation, and recovery dynamics, in grassland ecosystems exposed to heat
712 extremes: The role of species richness. *Ecosystems*, **4**: 797-806.
- 713 **Van Peer L, Nijs I, Reheul D, De Cauwer B. 2004.** Species richness and
714 susceptibility to heat and drought extremes in synthesized grassland
715 ecosystems: compositional vs physiological effects. *Functional Ecology*, **18**:
716 769-778.
- 717 **Voltaire F. 2008.** Plant traits and functional types to characterise drought survival
718 of pluri-specific perennial herbaceous swards in Mediterranean areas.
719 *European Journal of Agronomy*, **29**: 116-124.
- 720 **Voltaire F. 2018.** A unified framework for plant drought adaptive strategies: across
721 scales and disciplines. *Global Change Biology*: 1-10.
- 722 **Voltaire F, Lelievre F. 2001.** Drought survival in *Dactylis glomerata* and *Festuca*
723 *arundinacea* under similar rooting conditions in tubes. *Plant & Soil*, **229**: 225-
724 234.
- 725 **Voltaire F, Norton M. 2006.** Summer dormancy in perennial temperate grasses.
726 *Annals of Botany*, **98**: 927-933.
- 727 **Voltaire F, Thomas H, Bertagne N, Bourgeois E, Gautier MF, Lelievre F. 1998.**
728 Survival and recovery of perennial forage grasses under prolonged
729 Mediterranean drought: water status, solute accumulation, abscisic acid

- 730 concentration and accumulation of dehydrin transcripts in bases of immature
731 leaves. *New Phytologist*, **140**: 451-460.
- 732 **West CP, Oosterhuis DM, Wullschlegel SD. 1990.** Osmotic adjustment in
733 tissues of tall fescue in response to water deficit. *Environmental and*
734 *Experimental Botany*, **30**: 149-156.
- 735 **Wortemann R, Herbette S, Barigah TS, Fumanal B, Alia R, Ducousso A,**
736 **Gomory D, Roedel-Drevet P, Cochard H. 2011.** Genotypic variability and
737 phenotypic plasticity of cavitation resistance in *Fagus sylvatica* L. across
738 Europe. *Tree Physiology*, **31**: 1175-1182.
- 739 **Xie WG, Zhang XQ, Ma X, Cai HW, Huang LK, Pen Y, Zeng B. 2010.** Diversity
740 comparison and phylogenetic relationships of cocksfoot (*Dactylis glomerata*
741 L.) germplasm as revealed by SSR markers. *Canadian Journal of Plant*
742 *Science*, **90**: 13-21.
- 743 **Zanne AE, Tank DC, Cornwell WK, Eastman JM, Smith SA, FitzJohn RG,**
744 **McGlenn DJ, O'Meara BC, Moles AT, Reich PB, Royer DL, Soltis DE,**
745 **Stevens PF, Westoby M, Wright IJ, Aarssen L, Bertin RI, Calaminus A,**
746 **Govaerts R, Hemmings F, Leishman MR, Oleksyn J, Soltis PS, Swenson**
747 **NG, Warman L, Beaulieu JM. 2014.** Three keys to the radiation of
748 angiosperms into freezing environments. *Nature*, **506**: 89-+.
- 749 **Zwicke M, Picon-Cochard C, Morvan-Bertrand A, Prud'homme M, Volaire F.**
750 **2015.** What functional strategies drive drought survival and recovery of
751 perennial species from upland grassland? *Annals of Botany*, **116**: 1001-1015.
752

Population	Origin	Country	Latitude	Longitude	Temperature (°C)			Rainfall mean (mm)			Aridity Index
					Annual mean	Min of the coldest month	Max of the warmest month	Annual	Wettest quarter	Driest quarter	
Maroc 1(ma1)	Mediterranean	Morocco	33.10	-8.08	18.0	7.0	30.2	346	7	4	0.27
Maroc 9 (ma9)	Mediterranean	Morocco	29.71	-8.96	16.2	1.9	31.1	246	13	5	0.17
Kasbah (cv kas)	Mediterranean	Morocco	32.35	-6.16	11.7	3.9	39.6	270	215	17	0.31
Sicily (sic)	Mediterranean	Italy	37.48	14.50	15.4	4.0	27.6	572	237	40	0.47
Carnac (car)	Temperate	France	47.58	-3.07	11.7	3.5	21.6	893	179	153	0.86
Fourchette (fou)	Temperate	France	48.17	-2.75	10.8	2.3	20.8	857	174	153	0.74
St Michel (stm)	Temperate	France	48.63	-1.51	11.5	2.6	22.3	731	169	149	0.78
Ludac (cv lud)	Temperate	France	46.58	0.33	11.5	9	25.0	724	152	152	0.90
Danemark (dan)	Northern	Denmark	56.46	8.66	7.8	-3.2	19.4	809	201	134	1.48
Sweden (swe)	Northern	Sweden	60.50	24.22	4.8	-9.8	21.4	613	199	93	1.10
Loke (cv lok)	Northern	Sweden	62.98	14.80	2.6	-12.6	18.9	563	213	87	1.14
Tammisto (cv tam)	Northern	Finland	60.23	23.75	4.9	-9.3	21.3	609	196	92	1.08

753

754 **Table 1.** List of the twelve populations of *Dactylis glomerata*. Climatic data associated either with their origin sites of collection or
755 sites of breeding (for cultivars cv in Temperate and Northern areas) from the WorldClim data set (<http://www.worldclim.org>; (Hijmans
756 et al., 2005). Seeds from Northern populations were provided by the Nordic Genetic Resource Center (NordGen). Temperate
757 populations were provided by the plant genetic resources information system of the National Institute for Agronomical Research
758 (INRA, France, Lusignan). The Sicilian ecotype came from University of Catania, Italy. The Moroccan ecotypes came from INRA
759 Rabat, Morocco

760 **Table 2:** Results of variance analysis and Tukey test between means of variables for the three
 761 origins of *Dactylis glomerata* populations. ***: $p < 0.001$; **: $p < 0.01$; *: $p < 0.05$; ns: non-
 762 significant. For each variable, different subscript letters indicate significant differences between
 763 origins
 764

	Biogeographic origin of <i>D. glomerata</i> populations			
	P value	Medit.	Northern	Temperate
SURV = Plant dehydration survival (%)	<0.001***	58.2 ^a	13.1 ^b	6.9 ^b
P ₅₀ = Embolism pressure threshold for 50% embolism (MPa)	0.0016**	-5.8 ^a	-4.3 ^{ab}	-3.3 ^b
P ₈₈ = Embolism pressure threshold for 88% embolism (MPa)	<0.001***	-10.4 ^a	-7.6 ^{ab}	-6.1 ^b
Lwidth = Leaf width (mm)	<0.001***	4.28 ^a	7.97 ^c	6.22 ^b
Lthick = Leaf thickness (mm)	0.13ns	0.30	0.36	0.35
LDMC = Leaf Dry Matter content (mg g DM ⁻¹)	<0.001***	232.72 ^a	185.75 ^b	188.41 ^b
LFrac = Leaf fracture resistance (J m ⁻²)	<0.001***	478.6 ^a	188.47 ^b	225.55 ^b
A _S = Total stem area (mm ²)	0.006**	2.07 ^a	3.09 ^b	2.9 ^b
A _{LIG} = lignified area (μm ²)	0.02*	0.27 ^a	0.38 ^b	0.33 ^{ab}
P _{LIG} = Proportion of lignified area per total stem area	0.02*	0.1308 ^b	0.1258 ^{ab}	0.115 ^a
A _F = fiber cell area (μm ²)	0.34ns	70.61	77.21	77.56
A _{FL} = fiber lumen area (μm ²)	0.50ns	15.95	14.16	14.76
A _{FW} = fiber wall area (μm ²)	0.12ns	54.66	63.04	62.79
P _{CWF} = Proportion of cell wall per fiber	0.029*	0.77 ^a	0.82 ^b	0.81 ^{ab}
P _{FWF} *A _{LIG} = Proportion of total fiber wall area in lignified area	0.037*	0.30 ^{ab}	0.33 ^b	0.29 ^a
(P _{FWF} *A _{LIG}):A _S = Proportion of total fiber wall in lignified area per stem area	0.002**	0.156 ^a	0.117 ^b	0.102 ^b
D _{HV} = hydraulically weighted diameter of vessels (μm)	0.17ns	23.04	21.77	21.18
T _{VW} = Thickness of metaxylem vessel wall (μm)	<0.001***	1.426 ^c	1.226 ^b	1.109 ^a

765 **Table 3:** Pearson coefficients of correlations (lower part of the table) and P values (top part of the table) between traits measured on twelve populations of *Dactylis*
 766 *glomerata*. For abbreviations and units see table 1 and 2. Climatic variables of the origin sites (or breeding sites) of the populations: P_{DRY}= Mean precipitations of
 767 the driest three months; P_{YEAR} = Mean annual precipitations, T_{MIN} = mean minimum temperatures of the coldest months, Aridity = aridity index. P<0.05 (italics,
 768 bold), P<0.01 (bold), P<0.001 (bold underlined)

	P ₈₈	P ₅₀	SURV	Lwidth	Lthick	LDMC	LFrac	A _S	A _{LIG}	P _{LIG}	A _F	A _{FL}	A _{FW}	P _{CWF}	P _{FWF*} A _{LIG}	(P _{FWF*} A _{LIG}) :A _S	D _{HV}	T _{VW}	P _{SUM}	P _{YEAR}	T _{MIN}	Aridity
P ₈₈	1	0,00	0,00	0,10	0,19	0,00	0,00	0,13	0,18	0,89	0,45	0,67	0,38	0,37	0,88	0,14	0,09	0,00	0,00	0,00	0,76	0,04
P ₅₀	<u>0,94</u>	1	0,02	0,21	0,25	0,02	0,03	0,24	0,45	0,58	0,51	0,47	0,38	0,27	0,73	0,18	0,33	0,00	0,00	0,00	0,92	0,04
SURV	<u>-0,79</u>	<u>-0,65</u>	1	0,01	0,02	0,00	0,00	0,02	0,03	0,85	0,09	0,73	0,06	0,13	0,80	0,04	0,07	0,11	0,00	0,01	0,25	0,01
Lwidth	0,50	0,39	<u>-0,71</u>	1	0,01	0,00	0,00	0,00	0,02	0,40	0,14	0,87	0,16	0,35	0,15	0,09	0,71	0,36	0,10	0,21	0,00	0,00
Lthick	0,40	0,36	<u>-0,66</u>	<u>0,72</u>	1	0,02	0,02	0,20	0,42	0,53	0,08	0,48	0,04	0,07	0,30	0,73	0,81	0,78	0,06	0,12	0,03	0,01
LDMC	<u>-0,82</u>	<u>-0,68</u>	<u>0,87</u>	<u>-0,85</u>	<u>-0,65</u>	1	0,00	0,01	0,07	0,50	0,23	0,66	0,18	0,22	0,64	0,07	0,13	0,07	0,00	0,04	0,09	0,01
LFrac	<u>-0,78</u>	<u>-0,61</u>	<u>0,92</u>	<u>-0,84</u>	<u>-0,68</u>	<u>0,97</u>	1	0,02	0,03	0,81	0,21	0,62	0,16	0,18	0,62	0,08	0,10	0,11	0,01	0,05	0,08	0,00
A _S	0,46	0,37	<u>-0,66</u>	<u>0,83</u>	0,40	<u>-0,70</u>	<u>-0,67</u>	1	0,00	0,24	0,15	0,22	0,30	0,99	0,14	0,00	0,90	0,30	0,11	0,19	0,08	0,13
A _{LIG}	0,41	0,24	<u>-0,61</u>	<u>0,65</u>	0,26	-0,54	<u>-0,62</u>	<u>0,78</u>	1	0,36	0,24	0,24	0,42	0,88	0,06	0,08	0,71	0,34	0,30	0,49	0,17	0,13
P _{LIG}	-0,04	-0,18	0,06	-0,27	-0,20	0,22	0,08	-0,37	0,29	1	0,81	0,80	0,87	0,96	0,67	0,18	0,37	0,92	0,53	0,42	0,52	0,99
A _F	0,24	0,21	-0,52	0,45	0,53	-0,37	-0,39	0,44	0,37	-0,08	1	0,66	0,00	0,06	0,09	0,99	0,87	0,96	0,19	0,23	0,41	0,32
A _{FL}	-0,14	-0,23	0,11	0,05	-0,22	0,14	0,16	0,38	0,37	-0,08	0,14	1	0,64	0,01	0,05	0,76	0,21	0,64	0,26	0,17	0,92	0,10
A _{FW}	0,28	0,28	-0,55	0,43	<u>0,59</u>	-0,41	-0,44	0,33	0,26	-0,05	<u>0,96</u>	-0,15	1	0,00	0,28	0,92	0,60	0,93	0,09	0,10	0,43	0,13
P _{CWF}	0,28	0,35	-0,46	0,29	0,53	-0,38	-0,41	0,00	-0,05	-0,02	0,56	<u>-0,73</u>	<u>0,77</u>	1	0,60	0,90	0,21	0,65	0,06	0,03	0,49	0,02
P _{FWF*} A _{LIG}	-0,05	-0,11	-0,08	0,44	0,33	-0,15	-0,16	0,45	0,56	0,14	0,52	<u>0,59</u>	0,34	-0,17	1	0,60	0,11	0,68	0,69	0,53	0,20	0,78
(P _{FWF*} A _{LIG}):A _S	-0,45	-0,42	<u>0,59</u>	-0,50	-0,11	0,53	0,53	<u>-0,76</u>	-0,52	0,41	0,00	-0,10	0,03	0,04	0,17	1	0,59	0,16	0,08	0,08	0,47	0,24
D _{HV}	-0,51	-0,31	0,53	-0,12	-0,08	0,46	0,50	0,04	-0,12	-0,28	-0,05	0,39	-0,17	-0,39	0,49	0,17	1	0,44	0,19	0,37	0,97	0,36
T _{VW}	<u>-0,85</u>	<u>-0,92</u>	0,48	-0,29	-0,09	0,53	0,49	-0,33	-0,30	0,03	0,02	0,15	-0,03	-0,14	0,13	0,44	0,25	1	0,01	0,01	0,96	0,09
P _{SUM}	<u>0,88</u>	<u>0,89</u>	<u>-0,84</u>	0,50	0,55	<u>-0,77</u>	<u>-0,74</u>	0,48	0,32	-0,20	0,40	-0,35	0,51	0,56	-0,13	-0,53	-0,40	<u>-0,71</u>	1	0,00	0,83	0,01
P _{YEAR}	<u>0,75</u>	<u>0,86</u>	<u>-0,71</u>	0,39	0,48	<u>-0,61</u>	-0,57	0,40	0,22	-0,26	0,38	-0,42	0,50	<u>0,61</u>	-0,20	-0,52	-0,29	<u>-0,70</u>	<u>0,94</u>	1	0,83	0,01
T _{MIN}	0,10	-0,03	0,36	<u>-0,82</u>	<u>-0,62</u>	0,51	0,53	-0,53	-0,42	0,21	-0,26	-0,03	-0,25	-0,22	-0,40	0,23	0,01	-0,02	-0,07	-0,07	1	0,03
Aridity	<u>0,60</u>	<u>0,60</u>	<u>-0,71</u>	<u>0,76</u>	<u>0,73</u>	<u>-0,74</u>	<u>-0,77</u>	0,47	0,46	0,01	0,32	-0,49	0,46	<u>0,65</u>	0,09	-0,37	-0,29	-0,51	<u>0,70</u>	<u>0,70</u>	<u>-0,62</u>	1

769 **Legend of figures**

770 **Fig. 1. Thresholds of embolism resistance in cocksfoot.** Means and standard errors of water
 771 potentials at 50 % embolism (a, P_{50}) and at 88% embolism (b, P_{88}) in stems of twelve populations
 772 of *Dactylis glomerata* of contrasting origins (red = Mediterranean; blue = Northern; green =
 773 Temperate)

774

775 **Fig. 2. Embolism resistance at increasing water potentials in stems of the Mediterranean**
 776 *Dactylis glomerata* population ‘Kasbah’ growing either in pot or in the field. Mean water
 777 potentials at 50 % embolism (P_{50}) are -6.1 MPa and -5.2 MPa for the population growing in pot
 778 and in the field respectively.

779 **Fig. 3. Embolism resistance and aridity of origin site in cocksfoot.** Linear relationships
 780 between mean precipitation of the driest quarter at origin site and water potential at 50%
 781 embolism (P_{50}) in twelve populations of *Dactylis glomerata* of contrasting origins (red =
 782 Mediterranean; blue = Northern; green = Temperate). Bars correspond to standard errors.

783 **Fig. 4. Embolism resistance and (a) dehydration tolerance, (b) leaf fracture toughness and**
 784 **(c) leaf dry matter content in cocksfoot.** Linear relationships between water potential at 88%
 785 embolism (P_{88}) and (a) the proportion of surviving plants at soil wilting point in short pots
 786 (dehydration tolerance), (b) leaf fracture toughness of irrigated plants and (c) leaf dry matter
 787 content (LDMC) of irrigated plants in twelve populations of *Dactylis glomerata* of contrasting
 788 origins (red = Mediterranean; blue = Northern; green = Temperate). Bars correspond to standard
 789 errors.

790

791 **Fig. 5 Embolism resistance and stem anatomy in cocksfoot.** Linear relationships between
 792 water potential at 50% embolism (P_{50}) with thickness of metaxylem vessel walls in stem in
 793 twelve populations of *Dactylis glomerata* of contrasting origins (red= Mediterranean; blue=
 794 Northern; green= Temperate). Bars correspond to standard errors.

795 **Fig. 6 Thickness of metaxylem vessel wall in stems of cocksfoot:** three populations of *Dactylis*
 796 *glomerata* of contrasting origins: a = 'car' (Temperate); b = 'swe' (Northern); c = 'ma1'
 797 (Mediterranean). **The arrows point to wall of the metaxylem vessels.**

798 **Fig. 7 PCA of 12 plant traits and 12 populations of cocksfoot.** Principal component analysis
 799 (factorial axes 1 and 2) showing (a) the distribution of the twelve traits significantly
 800 discriminating origins of populations (only P₅₀ was included since it is highly correlated to P₈₈),
 801 see table 2 and (b) twelve populations of *Dactylis glomerata* of contrasting origins (red=
 802 Mediterranean; blue= Northern; green= Temperate). **Climatic data of the origin or breeding site:**
 803 **Annual precipitations (P_{year}), precipitations of the driest quarter (P_{dry}), aridity index (aridity)**
 804 **and minimum temperature of the coldest months (T_{min}) were included as supplementary**
 805 **variables (bold and italics characters).**

806



807

808

809 **Supplementary material**

810 **S1: S curves for determination of embolism resistance in cocksfoot.** Relationships between
 811 percentage of embolism and water potential measured for 12 populations of *Dactylis glomerata*
 812 originating from Temperate (green), Northern (blue) and Mediterranean (red) areas. Fitted curves
 813 are sigmoid functions. Vertical bars correspond to standard error, n = 1 to 7. P₅₀ and R² are given
 814 for each curve

815 **S2: Overview of the stem anatomy of the 12 populations of *Dactylis glomerata* and P₈₈ (MPa)**






Fig.3




Fig. 4




Fig. 5




Fig. 6




Fig. 7

