



**HAL**  
open science

## **KV4.3 Expression Modulates NaV1.5 Sodium Current**

Vincent Portero, Ronald Wilders, Simona Casini, Flavien Charpentier, Arie O. Verkerk, Carol Ann Remme

► **To cite this version:**

Vincent Portero, Ronald Wilders, Simona Casini, Flavien Charpentier, Arie O. Verkerk, et al.. KV4.3 Expression Modulates NaV1.5 Sodium Current. *Frontiers in Physiology*, 2018, 9, pp.178. 10.3389/fphys.2018.00178 . hal-01833332

**HAL Id: hal-01833332**

**<https://hal.science/hal-01833332>**

Submitted on 13 Jul 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# K<sub>V</sub>4.3 Expression Modulates Na<sub>V</sub>1.5 Sodium Current

Vincent Portero<sup>1</sup>, Ronald Wilders<sup>2</sup>, Simona Casini<sup>1</sup>, Flavien Charpentier<sup>3</sup>, Arie O. Verkerk<sup>1,2†</sup> and Carol Ann Remme<sup>1\*†</sup>

<sup>1</sup> Department of Experimental Cardiology, Academic Medical Center, Amsterdam, Netherlands, <sup>2</sup> Department of Medical Biology, Academic Medical Center, Amsterdam, Netherlands, <sup>3</sup> L'Institut du Thorax, INSERM, CNRS, Université de Nantes, Nantes, France

## OPEN ACCESS

### Edited by:

Marcel van der Heyden,  
University Medical Center Utrecht,  
Netherlands

### Reviewed by:

Anthony Varghese,  
University of Wisconsin–River Falls,  
United States

Ricardo Caballero,  
Complutense University of Madrid,  
Spain

Susanne Kämmerer,  
Institut für Pharmakologie und  
Toxikologie, Technische Universität  
Dresden, Germany

### \*Correspondence:

Carol Ann Remme  
c.a.remme@amc.uva.nl

<sup>†</sup>These authors have contributed  
equally to this work.

### Specialty section:

This article was submitted to  
Cardiac Electrophysiology,  
a section of the journal  
Frontiers in Physiology

Received: 06 December 2017

Accepted: 20 February 2018

Published: 12 March 2018

### Citation:

Portero V, Wilders R, Casini S,  
Charpentier F, Verkerk AO and  
Remme CA (2018) K<sub>V</sub>4.3 Expression  
Modulates Na<sub>V</sub>1.5 Sodium Current.  
Front. Physiol. 9:178.  
doi: 10.3389/fphys.2018.00178

In cardiomyocytes, the voltage-gated transient outward potassium current ( $I_{to}$ ) is responsible for the phase-1 repolarization of the action potential (AP). Gain-of-function mutations in *KCND3*, the gene encoding the  $I_{to}$  carrying K<sub>V</sub>4.3 channel, have been associated with Brugada syndrome (BrS). While the role of  $I_{to}$  in the pro-arrhythmic mechanism of BrS has been debated, recent studies have suggested that an increased  $I_{to}$  may directly affect cardiac conduction. However, the effects of an increased  $I_{to}$  on AP upstroke velocity or sodium current at the cellular level remain unknown. We here investigated the consequences of K<sub>V</sub>4.3 overexpression on Na<sub>V</sub>1.5 current and consequent sodium channel availability. We found that overexpression of K<sub>V</sub>4.3 protein in HEK293 cells stably expressing Na<sub>V</sub>1.5 (HEK293-Na<sub>V</sub>1.5 cells) significantly reduced Na<sub>V</sub>1.5 current density without affecting its kinetic properties. In addition, K<sub>V</sub>4.3 overexpression decreased AP upstroke velocity in HEK293-Na<sub>V</sub>1.5 cells, as measured with the alternating voltage/current clamp technique. These effects of K<sub>V</sub>4.3 could not be explained by alterations in total Na<sub>V</sub>1.5 protein expression. Using computer simulations employing a multicellular *in silico* model, we furthermore demonstrate that the experimentally observed increase in K<sub>V</sub>4.3 current and concurrent decrease in Na<sub>V</sub>1.5 current may result in a loss of conduction, underlining the potential functional relevance of our findings. This study gives the first proof of concept that K<sub>V</sub>4.3 directly impacts on Na<sub>V</sub>1.5 current. Future studies employing appropriate disease models should explore the potential electrophysiological implications in (patho)physiological conditions, including BrS associated with *KCND3* gain-of-function mutations.

**Keywords:** transient outward current, sodium current, channels, action potential, myocyte, arrhythmias, computer simulation

## INTRODUCTION

The cardiac sodium current ( $I_{Na}$ ), generated by the *SCN5A*-encoded voltage-gated Na<sup>+</sup> channel (Na<sub>V</sub>1.5) (Gellens et al., 1992), is responsible for the initial fast upstroke of the cardiac action potential (AP). It determines excitability of myocardial cells and ensures proper conduction of the electrical impulse within the heart. Consequently, Na<sub>V</sub>1.5 channel dysfunction may lead to conduction slowing, ventricular arrhythmias, and sudden cardiac death. In particular, *SCN5A* mutations leading to loss of sodium channel function are associated with isolated (progressive) conduction slowing or block (Schott et al., 1999), sick sinus syndrome (Benson et al., 2003), and Brugada syndrome (BrS) (Chen et al., 1998; Crotti et al., 2012; Le Scouarnec et al., 2015).

In cardiomyocytes, the voltage-gated transient outward K<sup>+</sup> current (I<sub>to</sub>) is responsible for the phase-1 repolarization of the cardiac AP and thereby contributes to the refractory period and inotropic state of the myocardium. In human, I<sub>to</sub> is generated by the K<sub>V</sub>4.3 channel, which is encoded by the gene *KCND3* (Niwa and Nerbonne, 2010). Gain-of-function mutations in *KCND3*, or its regulatory subunits, have also been associated with BrS (Delpón et al., 2008; You et al., 2015; Portero et al., 2016), giving rise to an ongoing discussion on the apparent role of I<sub>to</sub> in the pro-arrhythmic mechanism of BrS (Wilde et al., 2010). Previous studies have suggested that an increased I<sub>to</sub> may directly affect cardiac conduction due to a current-to-load mismatch during the depolarization process (Hoogendijk et al., 2010a,b). However, the effects of a gain-of-function of I<sub>to</sub> on the AP upstroke velocity at the cellular level remain unknown. The characterization of various knock-out mouse models of  $\alpha$ -subunits generating the fast component of I<sub>to</sub> (*Kcnd3*, *Kcnd2*) confirmed the involvement of I<sub>to</sub> in phase-1 repolarization, but its impact on AP upstroke velocity or I<sub>Na</sub> density were not investigated (Niwa et al., 2008; Liu et al., 2015). We recently evaluated the effects of an *in silico* I<sub>to</sub> injection on AP upstroke and repolarization velocity using the dynamic clamp technique (Verkerk et al., 2016). In that study, we observed a minimal effect ( $\approx 2\%$ ) of *in-silico* I<sub>to</sub> injection on upstroke velocity but only when the injected current was large and rapidly activated at very negative potentials (around  $-50$  mV). However, while an I<sub>to</sub>-like current does not appear to directly affect the fast depolarization, evidence is increasing that various ion channel proteins may interact and thus modulate each other's expression, function, or membrane trafficking (Balse and Boycott, 2017). For example, studies highlighted an interaction and co-regulation of the Kir2.1 and Na<sub>V</sub>1.5 proteins with a direct effect on their electrophysiological properties and thus important for cardiac excitability (Milstein et al., 2012). Moreover, a recent study demonstrated a direct interaction between the K<sub>V</sub>4.3 and hERG proteins, resulting in an increase in hERG current density upon co-expression of hERG with K<sub>V</sub>4.3 (Zhao et al., 2017). Considering these novel studies and the involvement of both I<sub>Na</sub> and I<sub>to</sub> in BrS, we hypothesized that the level of expression of K<sub>V</sub>4.3 may impact on I<sub>Na</sub>.

We addressed this hypothesis by evaluating the impact of K<sub>V</sub>4.3 protein overexpression on Na<sub>V</sub>1.5-based current and sodium channel availability in HEK293 cells stably expressing Na<sub>V</sub>1.5 (HEK293-Na<sub>V</sub>1.5). We show that an overexpression of K<sub>V</sub>4.3 channels leads to a reduction of Na<sub>V</sub>1.5 current density and lower AP upstroke velocity, as measured with the alternating voltage/current clamp (VC/CC) technique. These effects of K<sub>V</sub>4.3 overexpression could not be explained by alterations in total Na<sub>V</sub>1.5 protein expression. Computer simulations furthermore indicate that the experimentally observed decreased upstroke velocity is not directly due to an increase in the K<sub>V</sub>4.3-based I<sub>to</sub>, but instead a consequence of the K<sub>V</sub>4.3 protein itself. We also demonstrate that an increase in I<sub>to</sub> as well as a decrease in I<sub>Na</sub> can affect cardiac conduction and that a combination of both can lead to dramatic consequences, underlining the potential functional relevance of our findings.

## MATERIALS AND METHODS

### HEK293-Na<sub>V</sub>1.5 Cell Culture and Transfection

To avoid artifacts due to co-transfections, we evaluated the effect of the overexpression of K<sub>V</sub>4.3 protein on Na<sub>V</sub>1.5 currents using a genetically modified cell line constitutively overexpressing the human Na<sub>V</sub>1.5 channel (van Bemmelen et al., 2004). HEK293 cells stably expressing hNa<sub>V</sub>1.5 (HEK293-Na<sub>V</sub>1.5, kindly provided by Drs. Hugues Abriel and Jean-Sébastien Rougier) were cultured in DMEM (Gibco) supplemented with 10% FBS (Biowest), L-glutamine (Gibco), penicillin-streptomycin (Gibco), and Zeomycin (Invitrogen) in a 5% CO<sub>2</sub> incubator at 37°C. Cells were transfected at 70% confluency in 36.8 mm culture wells with 1  $\mu$ g IRES-GFP or 1  $\mu$ g *KCND3*-IRES-GFP cDNA using lipofectamine (Invitrogen, Carlsbad, USA). Both IRES-GFP and *KCND3*-IRES-GFP (human *KCND3* transcript reference: NM\_004980.4) constructs contained the same M61 vector plasmid backbone (Addgene). Gene-transfer was monitored by means of green fluorescence from the IRES-GFP or *KCND3*-IRES-GFP bicistronic vector. Patch clamp experiments were performed on single fluorescent cells 2 days after transfection.

### Ventricular Cell Preparation

Animal procedures were performed in accordance with governmental and institutional guidelines for animal use in research and were approved by the Animal Experimental Committee of the Academic Medical Center, Amsterdam, The Netherlands. Single left ventricular myocytes were isolated from 3 to 5 months old FVB/N mice by enzymatic dissociation (Remme et al., 2006). Therefore, mice were anesthetized by an intraperitoneal injection of pentobarbital prior to cervical dislocation, after which the heart was excised, cannulated, and mounted on a Langendorff perfusion set-up. The hearts were perfused for 5 min with Tyrode's solution containing (in mM): 140 NaCl, 5.4 KCl, 1.8 CaCl<sub>2</sub>, 1 MgCl<sub>2</sub>, 5.5 glucose, 5 HEPES; pH 7.4 (NaOH). Subsequently, the heart was perfused for 8 min with a similar solution in which the calcium concentration was lowered to 1  $\mu$ M, after which the enzyme Liberase Blendzyme type 4 (Roche; 0.05 mg/ml) and trypsin (Boehringer, 1  $\mu$ l/ml of 2.5% solution) were added for 10 minutes. Single myocytes were obtained by gently triturating the digested tissue in the low calcium enzyme solution supplemented with bovine serum albumin (BSA, 50 mg/ml). Myocytes were washed twice in normal calcium Tyrode's solution and quiescent, rod-shaped cross-striated cells with smooth surface were selected for measurements.

### Western Blot Experiment for Total Na<sub>V</sub>1.5 Quantification

Forty-eight hours after transfection (1  $\mu$ g IRES-GFP or 1  $\mu$ g *KCND3*-IRES-GFP plasmid), HEK293-Na<sub>V</sub>1.5 cells were washed twice with PBS and lysed in PBS containing 0.1% Triton and complete mini EDTA-free protease inhibitor mixture tablet (Roche). Cell lysates were then sonicated twice for 10 s and centrifuged for 10 min at 8,000 rpm. The supernatant was quantified using the BCA kit and 40  $\mu$ g of each separated

cell lysate was loaded on a 4–20% gradient gel (Bio-Rad) after 5 min of denaturation at 60°C in Laemmli buffer. The Rb $\alpha$ GFP antibody used for Western blotting was purchased from Santa Cruz Biotechnology, Rb $\alpha$ Calnexin from Calbiochem, and Rb $\alpha$ Na<sub>V</sub>1.5 from Sigma-Aldrich. Goat anti-rabbit horseradish peroxidase-conjugated secondary antibodies were purchased from GE Healthcare Life Science. Chemo luminescence signal was acquired with the ImageQuant LAS 4000 instrument. Band intensities were quantified using the public domain ImageJ software (USA National Institutes of Health). Total Na<sub>V</sub>1.5 protein expression protein was normalized to the calnexin signal.

## Electrophysiology

### Data Acquisition

Na<sub>V</sub>1.5 and K<sub>V</sub>4.3 currents and upstroke velocities (dV/dt) were measured in the whole-cell configuration of the patch-clamp technique using an Axopatch 200B amplifier (Molecular Devices Corporation, Sunnyvale, CA, USA) or a custom-made amplifier, capable of fast switching between voltage clamp (VC) and current clamp (CC) modes (Berecki et al., 2010). Voltage control, data acquisition, and analysis were accomplished using custom software. Potentials were corrected for the calculated liquid junction potentials (Barry and Lynch, 1991). Signals were low-pass filtered with a cut-off frequency of 5 kHz and digitized at 20, 4, and 40 kHz for Na<sub>V</sub>1.5, K<sub>V</sub>4.3, and upstroke velocities, respectively. Patch pipettes were pulled from borosilicate glass (Harvard Apparatus) and had resistances of  $\approx$ 2.0 M $\Omega$  after filling with the solutions as indicated below. Series resistance was compensated by  $\geq$ 80%. Cell membrane capacitance ( $C_m$ ) was calculated by dividing the time constant of the decay of the capacitive transient upon 5 mV hyperpolarizing voltage step from  $-40$  mV by the series resistance. The average  $C_m$  was not significantly different between the IRES-GFP and K<sub>V</sub>4.3-IRES-GFP cells [IRES-GFP:  $12.27 \pm 0.76$  pF ( $n = 36$ ); K<sub>V</sub>4.3-IRES-GFP:  $12.00 \pm 0.99$  pF ( $n = 37$ ); mean  $\pm$  standard error of the mean (SEM)].

### K<sub>V</sub>4.3 and Na<sub>V</sub>1.5 Current Measurements With Conventional VC

Na<sub>V</sub>1.5 current was measured at room temperature (21°C) with patch pipettes containing (in mM): 60 CsCl, 50 aspartic acid, 11 EGTA, 1.0 CaCl<sub>2</sub>, 1.0 MgCl<sub>2</sub>, 5.0 Na<sub>2</sub>ATP, 10 HEPES, pH 7.2 (CsOH). Bath solution for Na<sub>V</sub>1.5 current measurements contained (in mM): 20 NaCl, 130 CsCl, 2 CaCl<sub>2</sub>, 1 MgCl<sub>2</sub>, 5 glucose, 10 HEPES, pH 7.4 (CsOH). K<sub>V</sub>4.3 current was measured at 37°C in Tyrode's solution with the aforementioned composition. Patch pipettes for K<sub>V</sub>4.3 current measurements were filled with solution containing (in mM): 125 K-gluc, 20 KCl, 5 NaCl, 1 MgCl<sub>2</sub>, 10 BAPTA, 5 MgATP, 10 HEPES, pH 7.2 (NMDG-OH).

Current density and gating properties were determined by means of custom voltage-clamp protocols as indicated in figure insets. Cycle lengths were 5 and 10 s for Na<sub>V</sub>1.5 and K<sub>V</sub>4.3 current, respectively. Both Na<sub>V</sub>1.5 and K<sub>V</sub>4.3 current densities were defined as the difference between peak and steady-state current, divided by  $C_m$ . To determine the activation characteristics of Na<sub>V</sub>1.5 current, current-voltage (I-V) curves were corrected for driving force and

normalized to maximum peak current. Steady-state activation and inactivation curves were fitted using the Boltzmann equation  $I/I_{\max} = A/\{1.0 + \exp[(V_{1/2} - V)/k]\}$  to determine  $V_{1/2}$  (membrane potential for the half-maximal (in)activation) and the slope factor  $k$ .

### Upstroke Velocity Measurements Using Alternating VC/CC

The alternating VC/CC technique was used to measure Na<sub>V</sub>1.5 current-driven upstrokes at physiological temperature and Na<sup>+</sup> concentrations, as described previously (Berecki et al., 2010). Patch pipette and bath solutions were similar to those used for K<sub>V</sub>4.3 current measurements (see above). HEK293-Na<sub>V</sub>1.5 cells and freshly isolated myocytes were voltage clamped at a holding potential of  $-85$  mV, a value close to the resting membrane potential of working cardiomyocytes. Next, upstroke and repolarization were elucidated by switching for 20 ms to the CC mode of the patch clamp amplifier. Upstrokes were elicited by  $1.2 \times$  threshold current pulses through the patch pipette at 0.5 Hz and the stimulus current was present during the entire 20 ms in CC mode. Maximal upstroke velocity (dV/dt<sub>max</sub>) during VC/CC, offline corrected for the contribution of stimulus current ( $I_{\text{stim}}$ ), served as an indicator of Na<sub>V</sub>1.5 current availability.

## Computer Simulations

### Numerical Reconstruction of Na<sub>V</sub>1.5 and K<sub>V</sub>4.3 Currents in HEK293 Cells

Alternating VC/CC experiments in HEK293 cells expressing Na<sub>V</sub>1.5 and K<sub>V</sub>4.3 channels were simulated in a standard cell model with intracellular and extracellular sodium and potassium concentrations similar to those of the patch-clamp experiments. The cell model contained equations for  $I_{\text{Na}}$ ,  $I_{\text{to}}$ , and  $I_{\text{stim}}$ . The  $I_{\text{Na}}$  and  $I_{\text{to}}$  equations were taken from the human ventricular cell model by Ten Tusscher and Panfilov (2006) with the  $I_{\text{Na}}$  and  $I_{\text{to}}$  densities scaled by a factor of 1.8 and 6.0, respectively, to arrive at the experimentally observed values for the maximum upstroke velocity and repolarization velocity.  $I_{\text{stim}}$  was set to 6.5 pA/pF to ensure that the maximum upstroke velocity was reached at  $\approx$ 4 ms after the stimulus onset as in the experiments. For numerical integration of the differential equations we applied a simple and efficient Euler-type integration scheme (Rush and Larsen, 1978) with a 1- $\mu$ s time step.

### Maximum Upstroke Velocity and Repolarization Velocity in Murine Left Ventricular Myocytes

The alternating VC/CC protocol of the experiments on single left ventricular myocytes was implemented in the CellML code (Cuellar et al., 2003) of the apical version of the mouse ventricular AP model by Bondarenko et al. (2004). The amplitude of the 20 ms stimulus current was set to 10 pA/pF, which made the AP upstroke occur at  $\approx$ 4 ms after the stimulus onset. The maximum upstroke velocity and repolarization velocity were determined from the time derivative of the action potential trace. The CellML code was edited and run in version 0.9.31.1409 of the Windows based Cellular Open Resource (COR) environment (Garny et al., 2003) with its standard CVODE integrator.

## Action Potential Transfer in Human Left Ventricular Myocytes

AP transfer was studied in a one-dimensional strand of human left ventricular myocytes. The strand consisted of 80 longitudinally coupled cells, each described by the Ten Tusscher and Panfilov model of a single human ventricular myocyte (Ten Tusscher and Panfilov, 2006). The intercellular coupling conductance was set to 6 nS and the myoplasmic resistivity was set to 150 Ω·cm. The first cell of the strand was stimulated at a rate of 1 Hz with a 2-ms, ≈20% supra-threshold stimulus current with an amplitude of 3.2 nA. As in other studies (Shaw and Rudy, 1997; Thomas et al., 2003; Wilders, 2012), the entire cell length (of 74 μm) was used as the spatial discretization element, with elements connected by the lumped myoplasmic resistance (calculated from the myocyte dimensions and the myoplasmic resistivity of 150 Ω·cm) and gap junctional resistance. At the selected gap junctional conductance of 6 nS, this lumped resistance was almost completely determined by the gap junctional resistance (Wilders, 2012). The aforementioned Euler-type integration scheme was used for numerical integration.

## Statistics

Data are expressed as mean ± SEM. Mann-Whitney Rank Sum test, Student *t*-test, or Two-Way Repeated Measures ANOVA followed by pairwise comparison using the Student-Newman-Keuls test were used when appropriate. The level of statistical significance was set to  $p < 0.05$ .

## RESULTS


### Characterization of K<sub>V</sub>4.3 Current in HEK293 Cells Stably Expressing Nav1.5

First, we characterized the K<sub>V</sub>4.3-driven current resulting from transfection of the *KCND3*-IRES-GFP plasmid in HEK293 cells stably expressing Nav1.5 (HEK293-Nav1.5). **Figure 1A** shows representative currents measured between −100 to 40 mV (with 10 mV increment) during 1-s voltage clamp steps from a holding potential of −80 mV in a HEK293-Nav1.5 cell transfected with either IRES-GFP (top) or *KCND3*-IRES-GFP (bottom). A prepulse of 5 ms to −40 mV was applied to activate and inactivate the Nav1.5-based sodium current present in these HEK293-Nav1.5 cells (**Figure 1A**, inset). Similar to previous reports, cells transfected with IRES-GFP showed a small endogenous current upon depolarization (**Figure 1A**, top) (Niwa et al., 2008). In contrast, cells transfected with *KCND3*-IRES-GFP displayed a large K<sub>V</sub>4.3 current which started to activate around −40 mV and increased in amplitude upon further depolarization due to enhanced activation and increased K<sup>+</sup> driving force (**Figure 1A**, bottom) (Giles and van Ginneken, 1985). **Figure 1B** shows the average I-V relationships of the K<sub>V</sub>4.3 current, which was defined as the difference between the peak and steady-state current. Its density was defined as the current amplitude divided by the membrane capacitance ( $C_m$ ), amounting to ≈450 pA/pF at 40 mV in the *KCND3*-IRES-GFP transfected HEK293-Nav1.5 cells, and >20 times smaller in the IRES-GFP transfected cells. **Figure 1C** shows the voltage-dependency of inactivation for the K<sub>V</sub>4.3 current in the cells transfected with *KCND3*-IRES-GFP,

measured using a two-pulse protocol with a 1-s prepulse to a potential between −100 and 40 mV followed by a 500-ms test pulse to 40 mV, demonstrating that the K<sub>V</sub>4.3 current is fully available at −70 mV and more negative potentials. The  $V_{1/2}$  and  $k$  of the voltage-dependency of inactivation were  $-45.9 \pm 2.6$  and  $-5.6 \pm 0.4$  mV, respectively. **Figure 1D** shows the recovery from inactivation in *KCND3*-IRES-GFP transfected HEK293-Nav1.5 cells analyzed with 200-ms pulses to 40 mV with variable inter-pulse intervals. The time constant of recovery from inactivation, analyzed with a mono-exponential fit (**Figure 1D**, dashed line) was  $160 \pm 29$  ms, demonstrating that recovery from inactivation was completed with an inter-pulse interval of 1 s and longer. The K<sub>V</sub>4.3 current biophysical properties are summarized in Table S1. These data indicate that overexpression of *KCND3*-IRES-GFP in HEK293-Nav1.5 cells results in a large  $I_{to}$ , confirming the functional expression of K<sub>V</sub>4.3 channels at the cell membrane.

### K<sub>V</sub>4.3 Expression Reduces Nav1.5 Current in HEK293-Nav1.5 Cells

Secondly, we characterized the effects of K<sub>V</sub>4.3 overexpression on Nav1.5-based current in HEK293-Nav1.5 cells. **Figure 2A** shows representative Nav1.5 current activated by 500-ms depolarizing voltage clamp steps of 5 mV increment from a holding potential of −120 mV in a IRES-GFP or *KCND3*-IRES-GFP transfected HEK293-Nav1.5 cell. In both conditions, Nav1.5 current started to activate around −60 mV, peaked around −30 mV, and subsequently decreased in amplitude due to the reduction in Na<sup>+</sup> driving force. As shown in the average I-V relationships in **Figure 2B**, Nav1.5 current density was significantly smaller in the *KCND3*-IRES-GFP transfected cells compared to IRES-GFP transfected cells. For example, at −40 mV Nav1.5 current density was  $-609 \pm 62$  and  $-447 \pm 61$  pA/pF ( $p < 0.05$ ; pairwise comparison using a Student-Newman-Keuls test following two-way repeated measures ANOVA) in IRES-GFP and *KCND3*-IRES-GFP transfected HEK293-Nav1.5 cells, respectively, indicating a decrease in Nav1.5 current density by ≈25% due to K<sub>V</sub>4.3 overexpression. Next, we determined whether the Nav1.5 current density decrease was accompanied by gating property changes. Because the current decay could not be reliably fit to a 2-exponential function, the time course of inactivation was instead determined by analyzing the time required for 50% of current decay to occur ( $t_{50\%}$ ) at −30 mV (Remme et al., 2006).  $t_{50\%}$  did not differ significantly between IRES-GFP and *KCND3*-IRES-GFP transfected cells [ $0.88 \pm 0.06$  ms ( $n = 15$ ) vs.  $0.84 \pm 0.03$  ms ( $n = 19$ ), respectively,  $p = 0.41$ ; Student's *t*-test]. For determining the voltage-dependency of activation for IRES-GFP and *KCND3*-IRES-GFP transfected cells, I-V relationships, as shown in **Figure 2B**, were first corrected for the Na<sup>+</sup> driving force. Of note, the reversal potential of Nav1.5 current calculated using the Nernst equation is evaluated at +17.58 mV, which is in line with the sodium current recordings represented in **Figure 2A**. Next, currents were normalized to their maximal amplitude and the curves were fitted to a Boltzmann distribution curve. **Figure 2C** shows overlapping curves of voltage-dependency of inactivation. The latter curves were constructed by normalizing currents to their maximal



current during the voltage clamp step to  $-20$  mV. Similarly, **Figure 2D** shows overlapping curves of voltage-dependency of activation for IRES-GFP and *KCND3*-IRES-GFP transfected cells. These data demonstrate that neither the voltage-dependency of activation nor the voltage-dependency of inactivation of Na<sub>V</sub>1.5 current were affected by K<sub>V</sub>4.3 expression. The Na<sub>V</sub>1.5 current biophysical properties are summarized in Table S1. Western blot analysis showed no differences in Na<sub>V</sub>1.5 protein expression levels between HEK293-Nav1.5 cells with overexpression of *KCND3*-IRES-GFP or GFP (**Figure 3**; original images provided in the Supplementary Material, Figure S1). Thus, our data demonstrate that K<sub>V</sub>4.3 expression reduces Na<sub>V</sub>1.5 current without affecting its gating properties or the Na<sub>V</sub>1.5 expression level.



### K<sub>V</sub>4.3 Expression Reduces Action Potential Upstroke Velocity


Thirdly, we assessed upstroke velocities in the HEK293-Nav1.5 cells, transfected with either IRES-GFP or *KCND3*-IRES-GFP plasmids. Noteworthy, HEK293 cells expressing Na<sub>V</sub>1.5 display fast depolarizations upon switching from voltage clamp (VC) to current clamp (CC) due to Na<sup>+</sup> channel activation as previously shown (Berecki et al., 2010; Verkerk et al., 2016; Lieve et al., 2017). Thus, the alternating VC/CC technique allows for a more dynamic assessment of Na<sub>V</sub>1.5 current as compared to measurements in VC configuration, in the

setting of more physiological temperature and Na<sup>+</sup> gradients across the membrane. **Figure 4A** shows typical upstrokes (top) and their time derivatives (bottom) measured upon switching from a holding potential of  $-85$  mV in VC to CC for 20-ms. Upstrokes were evoked at 0.5 Hz, a stimulus frequency at which K<sub>V</sub>4.3 current is fully available (cf. **Figure 1D**). **Figure 4B** summarizes the maximum upstroke velocities (top) and maximum repolarization velocities (bottom). On average, the maximum upstroke velocity of *KCND3*-IRES-GFP transfected cells was 21% lower than that of IRES-GFP transfected cells ( $441 \pm 17$  vs.  $556 \pm 24$  V/s;  $p < 0.05$ ). Both cell types displayed a repolarization phase following the upstroke (**Figure 4A**) with a significantly larger maximum velocity ( $73 \pm 13$  vs.  $22 \pm 3$  V/s) in *KCND3*-IRES-GFP as compared to IRES-GFP transfected cells (**Figure 4B**, bottom). In **Figure 4C**, we plotted for each cell its maximum repolarization velocity vs. its maximum upstroke velocity, further demonstrating an inverse relationship between Na<sub>V</sub>1.5-induced depolarization and K<sub>V</sub>4.3-induced repolarization velocities: K<sub>V</sub>4.3 expression increases repolarization velocity, but reduces upstroke velocity.

### Relation Between Repolarization and Depolarization in Ventricular Myocytes of Mice

The above presented data indicate that K<sub>V</sub>4.3 (over)expression decreases Na<sub>V</sub>1.5 current (**Figure 2B**) and upstroke velocity





(Figure 4B), resulting in an inverse relationship between the maximum velocities of depolarization and repolarization (Figure 4C). Next, we determined whether a relationship between repolarization and depolarization also exists in cardiomyocytes. Therefore, we re-analyzed data of alternating VC/CC measurements in murine isolated left ventricular myocytes performed for a previous study (Remme et al., 2006).

In mouse ventricular myocytes, the maximum upstroke and repolarization velocities were  $285 \pm 19$  and  $-18 \pm 2$  V/s ( $n = 18$ ), respectively. In Figure 4D, we plotted for each cell its maximum repolarization velocity vs. its maximum upstroke velocity, demonstrating an inverse relationship, similar to our experiments on HEK293 cells (Figure 4C).




## Computer Simulations

To assess the functional relevance of the aforementioned findings, we first carried out computer simulations to explore whether the presence of a K<sub>V</sub>4.3-based  $I_{to}$  *per se* may affect the maximum upstroke velocity of the HEK293-Na<sub>V</sub>1.5 cells. To this end, we constructed a model of a HEK293 cell expressing both  $I_{Na}$  and  $I_{to}$  channels to further explore the findings of the alternating VC/CC experiment of **Figures 4A,B**. The  $I_{Na}$  and  $I_{to}$  equations were taken from the human ventricular cell model by Ten Tusscher and Panfilov (2006). The characteristics of the simulated  $I_{Na}$  and  $I_{to}$  are shown in **Figures 5A–D** (and also in Figures S2, S3). As illustrated in **Figures 5E–G**,  $I_{to}$  is effectively zero at the moment of maximum upstroke


velocity, indicating that this maximum upstroke velocity is fully determined by  $I_{Na}$ . Conversely,  $I_{Na}$  is almost zero at the moment of maximum repolarization velocity. Accordingly, there is no direct contribution of  $I_{Na}$  to the maximum repolarization velocity, although it must be noted that  $I_{Na}$  may still affect repolarization velocity through its effect on AP shape. If the voltage-dependency of the  $I_{to}$  channels is shifted by  $-30$  mV, thus letting them start to activate near  $-40$  mV,  $I_{to}$  is still effectively zero at the moment of maximum upstroke velocity (data not shown).

Next, we established to which extent the K<sub>V</sub>4.3-based  $I_{to}$  and the Na<sub>V</sub>1.5-based  $I_{Na}$  *per se* modulate the maximum upstroke velocity and repolarization velocity of murine left



ventricular myocytes. To this end, we ran computer simulations of the alternating VC/CC experiment on mouse ventricular myocytes of **Figure 4D** using the apical version of the mouse ventricular AP model by Bondarenko et al. (2004), in which we increased or decreased the density of either  $I_{to}$  or  $I_{Na}$ . The characteristics of  $I_{Na}$  and  $I_{to}$  of the Bondarenko et al. (2004) model are shown in **Figures 6A–C** (and also in Figures S2, S3). As shown in **Figure 6D**, and as expected from **Figure 5**, the maximum upstroke velocity is independent of  $I_{to}$  conductance ( $G_{to}$ ), whereas  $G_{to}$  is an important determinant of repolarization velocity. The maximum repolarization velocity increases almost linearly with  $G_{to}$  (**Figure 6D**, open squares), albeit not in a

1:1 fashion—a four times increase in  $G_{to}$  (from 40 to 160% of its control value) results in a 2.3 times increase in maximum repolarization velocity, which can be explained by accompanying changes in AP shape (and thus in channel activation and inactivation) as well as the presence of other membrane currents. As shown in **Figure 6E**, the maximum upstroke velocity is strongly dependent on  $I_{Na}$  conductance ( $G_{Na}$ ), whereas repolarization velocity is only substantially affected by  $G_{Na}$  at lower values of  $G_{Na}$ . At these lower values, the AP reaches a considerably lower peak (data not shown), which in turn results in less activation of  $I_{to}$  channels as well as a lower driving force for these channels. At control or higher values



of  $G_{Na}$ , repolarization velocity is not notably dependent on  $G_{Na}$ .

Finally, we tested whether a decrease in  $G_{Na}$  can result in a loss of conduction in the presence of a concomitant increase in  $G_{to}$ . This was studied in a one-dimensional strand of poorly coupled human left ventricular myocytes, as illustrated in **Figure 7A**. Under control conditions (100%  $G_{Na}$  and 100%  $G_{to}$ ), the stimulated leftmost cell of the strand was able to drive its neighboring cells (**Figure 7B**). This was also observed upon a 20% increase in  $G_{to}$  or a 50% decrease in  $G_{Na}$  (**Figures 7C,D**, respectively). The main effect of the decrease in  $G_{Na}$  was a slowing of conduction (cf. **Table 1**). However, the introduction of a concomitant 20% increase in  $G_{to}$  led to loss of conduction: an action potential was elicited in the first cell of the strand, but this cell was unable to transfer this action potential to its neighboring cells (**Figure 7E**). Thus, a simultaneous increase in  $I_{to}$  and decrease in  $I_{Na}$  in the setting of a compromised cellular coupling, as may occur on a microscopic scale in BrS patients, may result in severe conduction alteration.


## DISCUSSION

In the present study, we report for the first time a direct effect of K<sub>V</sub>4.3 expression on Nav1.5 current and consequent sodium channel availability. We found that overexpression of K<sub>V</sub>4.3 protein reduces Nav1.5 current density (**Figure 2**) without affecting total Nav1.5 protein expression. Moreover, K<sub>V</sub>4.3 overexpression decreased the upstroke velocity, which was not

due to a direct effect of the current generated by K<sub>V</sub>4.3. These findings suggest that the observed effects of K<sub>V</sub>4.3 on Nav1.5 are due to a functional interaction between the two channel proteins rather than an electrophysiological interference. The physiological relevance of our findings was demonstrated in a multicellular *in silico* model where we found that an  $I_{to}$  increase and concomitant  $I_{Na}$  decrease is capable of impairing conduction (**Figure 7**).

## Decreased AP Upstroke Velocity Secondary to K<sub>V</sub>4.3 Overexpression

Using the alternating VC/CC technique, we observed in HEK293 cells stably expressing Nav1.5 a decrease in upstroke velocity secondary to overexpression of K<sub>V</sub>4.3. Upstroke velocity decreased to a similar extent as Nav1.5 sodium current density ( $\approx 25\%$ ) in cells overexpressing K<sub>V</sub>4.3 compared to cells overexpressing only GFP. Although the alternating VC/CC technique is not commonly used in cardiac electrophysiology, it allows detailed  $I_{Na}$  studies in a dynamic fashion under close-to-physiological conditions as occurring during cardiac action potentials (Berecki et al., 2010). Previously it has been used to demonstrate changes in  $I_{Na}$  during heart failure (Berecki et al., 2010), changes in intracellular  $Ca^{2+}$  concentration (Casini et al., 2009), and in the setting of *SCN5A* mutations (Remme et al., 2006; Lieve et al., 2017). Using dynamic clamp, we have recently assessed the effect of an  $I_{to}$ -like current on the upstroke velocity in HEK293 cells transiently overexpressing Nav1.5 channels. That study revealed only a minor influence of  $I_{to}$ , if any, on



**TABLE 1** | Action potential parameters and propagation in a strand of human left ventricular myocytes.

Settings	APA (mV)	(dV <sub>m</sub> /dt) <sub>max</sub> (V/s)	APD <sub>90</sub> (ms)	CV (cm/s)
100% G <sub>Na</sub> , 100% G <sub>To</sub>	111.3	223.5	275.6	0.260
100% G <sub>Na</sub> , 120% G <sub>To</sub>	110.5	217.1	272.1	0.245
50% G <sub>Na</sub> , 100% G <sub>To</sub>	104.6	79.0	248.1	0.163

Data obtained from computer simulations with an intercellular coupling conductance of 6 nS. APA, action potential amplitude; (dV<sub>m</sub>/dt)<sub>max</sub>, maximum upstroke velocity; APD<sub>90</sub>, action potential duration at 90% repolarization; CV, conduction velocity in the strand. Action potential parameters for cells in the middle of the strand.

upstroke velocity (Verkerk et al., 2016). Here, we confirmed these results using computer simulations based on the current densities recorded (Figure 5). The I<sub>To</sub> current is not yet present at the time of the maximum upstroke velocity confirming that I<sub>To</sub> does not affect the upstroke velocity in single cells. Moreover, no correlation between I<sub>To</sub> density and AP upstroke velocity was observed during simulations of the electrical activity of cardiomyocytes (Figure 6). Thus, the observed K<sub>V</sub>4.3-induced decrease in upstroke velocity is not a consequence of an increased K<sub>V</sub>4.3 current.

In freshly isolated left ventricular myocytes of mice, we found an inverse relationship between the maximum upstroke velocity and repolarization velocity (Figure 4D). It is tempting to speculate that such a relationship is due to effects of K<sub>V</sub>4.3 expression, but this observation might also be due to different currents or transcription gradient of both *Kcnd3* and *Scn5a* through the myocardial wall. Nevertheless, it indicates that depolarization and repolarization are not independent factors. Future overexpression or knock-down experiments in myocytes are needed to provide further insight into the potential relevance of our observation.

## Modulation of Nav1.5 Current by K<sub>V</sub>4.3 Expression

We found that K<sub>V</sub>4.3 overexpression resulted in a significantly decreased Nav1.5 current density, while neither current kinetics nor total Nav1.5 protein expression were affected. Several studies have previously reported interactions and consequent inter-relationships between sodium and potassium channels, or their associated subunits, leading to electrophysiological modifications of either current (Hu et al., 2012; Milstein et al., 2012; Matamoros et al., 2016; Utrilla et al., 2017). For instance, Kir2.1 and Nav1.5 channels when co-expressed exert a synergic effect on current density due to a distinct trafficking process as compared to when these channels are expressed separately (Utrilla et al., 2017). Concerning the direct interaction of K<sub>V</sub>4.3 and Nav1.5, a mutation in the *SCN1B* gene encoding the cardiac sodium channel β1 subunit has been found in BrS and sudden infant death syndrome patients (Hu et al., 2012). This specific *SCN1B* mutation led to a decreased I<sub>Na</sub> together with an increased I<sub>To</sub>, suggesting that Nav1.5 may physically interact with K<sub>V</sub>4.3 via β1 to form a macromolecular complex. In neurons, both sodium and potassium channels mainly localize at the

axon initial segment (AIS), a subcellular region involved in AP genesis. Ion channel recruitment to the AIS is highly dependent on the cytoskeleton and associated proteins which densely populate this region (Brachet et al., 2010; Letierrier et al., 2011; Vacher et al., 2011). A comparable structure is observed at the intercalated discs in adult cardiomyocytes, a subcellular region also characterized by a high density of sodium channels (Agullo-Pascual et al., 2014; Marsman et al., 2014; El Refaey et al., 2017). Our findings, together with existing knowledge, may suggest a potential subcellular co-localization of K<sub>V</sub>4.3 and Nav1.5 in cardiomyocytes and a possible explanation for the co-regulation of channel function. Further studies should aim at unraveling the molecular mechanism involved in our observation. A recent work also highlighted the synergic impact of K<sub>V</sub>4.3 expression on hERG channels in HEK293 cells (Zhao et al., 2017). This study together with our findings strongly suggest that the overexpression or knock-down of K<sub>V</sub>4.3 in cardiomyocytes may lead to a complex modulation of ion channels initially seen as unrelated including Nav1.5, hERG, and Kir2.1 channels.

## Potential (Patho)Physiological Implications

Our computer simulations show that a concomitant increase in I<sub>To</sub> and decrease in I<sub>Na</sub> are capable of significantly affecting cardiac conduction. Our current observations may thus be particularly relevant in the phenotypical explanation of BrS patients presenting with mutations in the *KCND3* gene or in genes encoding K<sub>V</sub>4.3 associated subunits leading to gain-of-function through an increased protein stability or membrane trafficking of K<sub>V</sub>4.3 channels. In this context, an increased I<sub>To</sub> current could be associated with a simultaneous decrease in I<sub>Na</sub> and consequent conduction slowing (Kucera et al., 2017). Such an effect is likely even more prominent within the subepicardium, a region with relatively low I<sub>Na</sub> and high I<sub>To</sub>. This may be of particular relevance in the right ventricular subepicardium and right ventricular outflow tract, where fractionated signals indicative of slowed conduction are often observed in BrS patients (Nademanee et al., 2011). Importantly, concomitant cardiac structural abnormalities, for example age-dependent development of fibrosis (Coronel et al., 2005), may further compromise conduction and hence unmask the impact of increased I<sub>To</sub> (Hoogendijk et al., 2010a). Moreover, one may speculate that the known gender differences in I<sub>To</sub> magnitude may also contribute to the male preponderance for BrS via an indirect effect on I<sub>Na</sub> (Di Diego et al., 2002).

## Limitations

While we clearly demonstrated that K<sub>V</sub>4.3 is capable of decreasing Nav1.5 current density, extrapolation to physiological cardiomyocyte conditions must be done with some caution. The upstroke velocities measured in our experiments are in the same range of values as recorded in cardiomyocytes (Berecki et al., 2010; Veerman et al., 2017). However, *KCND3* overexpression resulted in K<sub>V</sub>4.3 currents larger than the I<sub>To</sub> previously reported in various animal species and human ventricular cardiomyocytes (Niwa and Nerbonne, 2010).

Consequently, the K<sub>V</sub>4.3-induced ≈25% reduction of Na<sub>V</sub>1.5 current density and upstroke velocity reported in this study is likely smaller in animal species and human. Moreover, our experiments in HEK293 cells were performed without β subunits of either Na<sub>V</sub>1.5 or K<sub>V</sub>4.3. We therefore cannot exclude a supplementary complexity of Na<sub>V</sub>1.5 macromolecular complexes in cardiomyocytes. Nevertheless, our myocyte data at least demonstrate that depolarization is not independent of repolarization.

## CONCLUSION

Overall, this study gives the first proof of concept that the K<sub>V</sub>4.3 protein directly impacts on Na<sub>V</sub>1.5 current. Future studies employing appropriate disease models should explore the potential electrophysiological implications in (patho)physiological conditions, including BrS associated with gain-of-function mutations in *KCNQ3*.

## AUTHOR CONTRIBUTIONS

Experimental design: VP, RW, FC, AV, and CR; Data acquisition: VP, RW, and AV; Analysis and interpretation of data: VP, RW, SC, AV, and CR; Drafting manuscript: VP, RW, AV, and CR;

## REFERENCES

- Agullo-Pascual, E., Lin, X., Leo-Macias, A., Zhang, M., Liang, F.-X., Li, Z., et al. (2014). Super-resolution imaging reveals that loss of the C-terminus of connexin43 limits microtubule plus-end capture and Na<sub>V</sub>1.5 localization at the intercalated disc. *Cardiovasc. Res.* 104, 371–381. doi: 10.1093/cvr/cvu195
- Balse, E., and Boycott, H. E. (2017). Ion channel trafficking: control of ion channel density as a target for arrhythmias? *Front. Physiol.* 8:808. doi: 10.3389/fphys.2017.00808
- Barry, P. H., and Lynch, J. W. (1991). Liquid junction potentials and small cell effects in patch-clamp analysis. *J. Membr. Biol.* 121, 101–117. doi: 10.1007/BF01870526
- Benson, D. W., Wang, D. W., Dymont, M., Knilans, T. K., Fish, F. A., Strieper, M. J., et al. (2003). Congenital sick sinus syndrome caused by recessive mutations in the cardiac sodium channel gene (*SCN5A*). *J. Clin. Invest.* 112, 1019–1028. doi: 10.1172/JCI18062
- Berecki, G., Wilders, R., de Jonge, B., van Ginneken, A. C. G., and Verkerk, A. O. (2010). Re-evaluation of the action potential upstroke velocity as a measure of the Na<sup>+</sup> current in cardiac myocytes at physiological conditions. *PLoS ONE* 5:e15772. doi: 10.1371/journal.pone.0015772
- Bondarenko, V. E., Szigeti, G. P., Bett, G. C. L., Kim, S.-J., and Rasmusson, R. L. (2004). Computer model of action potential of mouse ventricular myocytes. *Am. J. Physiol. Heart Circ. Physiol.* 287, H1378–H1403. doi: 10.1152/ajpheart.00185.2003
- Brachet, A., Leterrier, C., Irondele, M., Fache, M.-P., Racine, V., Sibarita, J.-B., et al. (2010). Ankyrin G restricts ion channel diffusion at the axonal initial segment before the establishment of the diffusion barrier. *J. Cell. Biol.* 191, 383–395. doi: 10.1083/jcb.201003042
- Casini, S., Verkerk, A. O., van Borren, M. M. G. J., van Ginneken, A. C. G., Veldkamp, M. W., de Bakker, J. M. T., et al. (2009). Intracellular calcium modulation of voltage-gated sodium channels in ventricular myocytes. *Cardiovasc. Res.* 81, 72–81. doi: 10.1093/cvr/cvn274
- Chen, Q., Kirsch, G. E., Zhang, D., Brugada, R., Brugada, J., Brugada, P., et al. (1998). Genetic basis and molecular mechanism for idiopathic ventricular fibrillation. *Nature* 392, 293–296. doi: 10.1038/32675

Editing manuscript, and approval: VP, RW, SC, FC, AV, and CR; Funding: CR.

## FUNDING

This study was supported by an Innovational Research Incentives Scheme Vidi grant from ZonMw (grant no. 91714371 to CR).

## ACKNOWLEDGMENTS

We thank Drs. Hugues Abriel and Jean-Sebastien Rougier (Institute of Biochemistry and Molecular Medicine, Bern, Switzerland) for kindly providing the HEK293-Na<sub>V</sub>1.5 cell line, Dr. Ruben Coronel (Academic Medical Center, Amsterdam, The Netherlands) for inspiring discussions, and Jan G. Zegers and Dr. Antoni C.G van Ginneken (Academic Medical Center, Amsterdam, The Netherlands) for developing and providing technical assistance with the soft- and hardware.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fphys.2018.00178/full#supplementary-material>

- Coronel, R., Casini, S., Koopmann, T. T., Wilms-Schopman, F. J. G., Verkerk, A. O., de Groot, J. R., et al. (2005). Right ventricular fibrosis and conduction delay in a patient with clinical signs of Brugada syndrome: a combined electrophysiological, genetic, histopathologic, and computational study. *Circulation* 112, 2769–2777. doi: 10.1161/CIRCULATIONAHA.105.532614
- Crotti, L., Marcou, C. A., Tester, D. J., Castelletti, S., Giudicessi, J. R., Torchio, M., et al. (2012). Spectrum and prevalence of mutations involving BrS1- through BrS12-susceptibility genes in a cohort of unrelated patients referred for Brugada syndrome genetic testing: implications for genetic testing. *J. Am. Coll. Cardiol.* 60, 1410–1418. doi: 10.1016/j.jacc.2012.04.037
- Cuellar, A. A., Lloyd, C. M., Nielsen, P. F., Bullivant, D. P., Nickerson, D. P., and Hunter, P. J. (2003). An overview of CellML 1.1, a biological model description language. *Simulation* 79, 740–747. doi: 10.1177/0037549703040939
- Delpón, E., Cordeiro, J. M., Núñez, L., Thomsen, P. E. B., Guerschicoff, A., Pollevick, G. D., et al. (2008). Functional effects of *KCNE3* mutation and its role in the development of Brugada syndrome. *Circ. Arrhythm. Electrophysiol.* 1, 209–218. doi: 10.1161/CIRCEP.107.748103
- Di Diego, J. M., Cordeiro, J. M., Goodrow, R. J., Fish, J. M., Zygmunt, A. C., Pérez, G. J., et al. (2002). Ionic and cellular basis for the predominance of the Brugada syndrome phenotype in males. *Circulation* 106, 2004–2011. doi: 10.1161/01.CIR.0000032002.22105.7A
- El Refaey, M. M., and Mohler, P. J. (2017). Ankyrins and spectrins in cardiovascular biology and disease. *Front. Physiol.* 8:852. doi: 10.3389/fphys.2017.00852
- Garny, A., Kohl, P., and Noble, D. (2003). Cellular open resource (COR): a public CellML based environment for modelling biological function. *Int. J. Bifurcat. Chaos* 13, 3579–3590. doi: 10.1142/S021812740300882X
- Gellens, M. E., George, A. L., Chen, L. Q., Chahine, M., Horn, R., Barchi, R. L., et al. (1992). Primary structure and functional expression of the human cardiac tetrodotoxin-insensitive voltage-dependent sodium channel. *Proc. Natl. Acad. Sci. U.S.A.* 89, 554–558. doi: 10.1073/pnas.89.2.554
- Giles, W. R., and van Ginneken, A. C. G. (1985). A transient outward current in isolated cells from the crista terminalis of rabbit heart. *J. Physiol.* 368, 243–264. doi: 10.1113/jphysiol.1985.sp015856
- Hoogendijk, M. G., Ophof, T., Postema, P. G., Wilde, A. A. M., de Bakker, J. M. T., and Coronel, R. (2010a). The Brugada ECG pattern: a marker of channelopathy, structural heart disease, or neither? Toward a unifying

- mechanism of the Brugada syndrome. *Circ. Arrhythm. Electrophysiol.* 3, 283–290. doi: 10.1161/CIRCEP.110.937029
- Hoogendijk, M. G., Potse, M., Linnenbank, A. C., Verkerk, A. O., den Ruijter, H. M., van Amersfoort, S. C. M., et al. (2010b). Mechanism of right precordial ST-segment elevation in structural heart disease: excitation failure by current-to-load mismatch. *Heart Rhythm* 7, 238–248. doi: 10.1016/j.hrthm.2009.10.007
- Hu, D., Barajas-Martínez, H., Medeiros-Domingo, A., Crotti, L., Veltmann, C., Schimpf, R., et al. (2012). A novel rare variant in *SCN1Bb* linked to Brugada syndrome and SIDS by combined modulation of Nav1.5 and Kv4.3 channel currents. *Heart Rhythm* 9, 760–769. doi: 10.1016/j.hrthm.2011.12.006
- Kucera, J. P., Rohr, S., and Kleber, A. G. (2017). Microstructure, cell-to-cell coupling, and ion currents as determinants of electrical propagation and arrhythmogenesis. *Circ. Arrhythm. Electrophysiol.* 10:e004665. doi: 10.1161/CIRCEP.117.004665
- Le Scouarnec, S., Karakachoff, M., Gourraud, J.-B., Lindenbaum, P., Bonnaud, S., Portero, V., et al. (2015). Testing the burden of rare variation in arrhythmia-susceptibility genes provides new insights into molecular diagnosis for Brugada syndrome. *Hum. Mol. Genet.* 24, 2757–2763. doi: 10.1093/hmg/ddv036
- Leterrier, C., Vacher, H., Fache, M.-P., d'Ortoli, S. A., Castets, F., Autillo-Touati, A., et al. (2011). End-binding proteins EB3 and EB1 link microtubules to ankyrin G in the axon initial segment. *Proc. Natl. Acad. Sci. U.S.A.* 108, 8826–8831. doi: 10.1073/pnas.1018671108
- Lieve, K. V., Verkerk, A. O., Podliesna, S., van der Werf, C., Tanck, M. W., Hofman, N., et al. (2017). Gain-of-function mutation in *SCN5A* causes ventricular arrhythmias and early onset atrial fibrillation. *Int. J. Cardiol.* 236, 187–193. doi: 10.1016/j.ijcard.2017.01.113
- Liu, J., Kim, K.-H., Morales, M. J., Heximer, S. P., Hui, C.-C., and Backx, P. H. (2015). Kv4.3-encoded fast transient outward current is presented in Kv4.2 knockout mouse cardiomyocytes. *PLoS ONE* 10:e0133274. doi: 10.1371/journal.pone.0133274
- Marsman, R. F. J., Bezzina, C. R., Freiberg, F., Verkerk, A. O., Adriaens, M. E., Podliesna, S., et al. (2014). Coxsackie and adenovirus receptor is a modifier of cardiac conduction and arrhythmia vulnerability in the setting of myocardial ischemia. *J. Am. Coll. Cardiol.* 63, 549–559. doi: 10.1016/j.jacc.2013.10.062
- Matamoros, M., Pérez-Hernández, M., Guerrero-Serna, G., Amorós, I., Barana, A., Núñez, M., et al. (2016). Nav1.5 N-terminal domain binding to  $\alpha$ 1-syntrophin increases membrane density of human Kir2.1, Kir2.2 and Nav1.5 channels. *Cardiovasc. Res.* 110, 279–290. doi: 10.1093/cvr/cvw009
- Milstein, M. L., Musa, H., Balbuena, D. P., Anumonwo, J. M. B., Auerbach, D. S., Furspan, P. B., et al. (2012). Dynamic reciprocity of sodium and potassium channel expression in a macromolecular complex controls cardiac excitability and arrhythmia. *Proc. Natl. Acad. Sci. U.S.A.* 109, E2134–E2143. doi: 10.1073/pnas.1109370109
- Nademanee, K., Veerakul, G., Chandanamattha, P., Chaothawee, L., Ariyachaipanich, A., Jirasirirojanakorn, K., et al. (2011). Prevention of ventricular fibrillation episodes in Brugada syndrome by catheter ablation over the anterior right ventricular outflow tract epicardium. *Circulation* 123, 1270–1279. doi: 10.1161/CIRCULATIONAHA.110.972612
- Niwa, N., and Nerbonne, J. M. (2010). Molecular determinants of cardiac transient outward potassium current ( $I_{to}$ ) expression and regulation. *J. Mol. Cell. Cardiol.* 48, 12–25. doi: 10.1016/j.yjmcc.2009.07.013
- Niwa, N., Wang, W., Sha, Q., Marionneau, C., and Nerbonne, J. M. (2008). Kv4.3 is not required for the generation of functional  $I_{to,f}$  channels in adult mouse ventricles. *J. Mol. Cell. Cardiol.* 44, 95–104. doi: 10.1016/j.yjmcc.2007.10.007
- Portero, V., Le Scouarnec, S., Es-Salah-Lamoureux, Z., Burel, S., Gourraud, J.-B., Bonnaud, S., et al. (2016). Dysfunction of the voltage-gated K<sup>+</sup> channel  $\beta$ 2 subunit in a familial case of Brugada syndrome. *J. Am. Heart Assoc.* 5:e003122. doi: 10.1161/JAHA.115.003122
- Remme, C. A., Verkerk, A. O., Nuyens, D., van Ginneken, A. C. G., van Brunschot, S., Belterman, C. N. W., et al. (2006). Overlap syndrome of cardiac sodium channel disease in mice carrying the equivalent mutation of human *SCN5A-1795insD*. *Circulation* 114, 2584–2594. doi: 10.1161/CIRCULATIONAHA.106.653949
- Rush, S., and Larsen, H. (1978). A practical algorithm for solving dynamic membrane equations. *IEEE Trans. Biomed. Eng.* 25, 389–392. doi: 10.1109/TBME.1978.326270
- Schott, J. J., Alshinawi, C., Kyndt, F., Probst, V., Hoorntje, T. M., Hulsbeek, M., et al. (1999). Cardiac conduction defects associate with mutations in *SCN5A*. *Nat. Genet.* 23, 20–21. doi: 10.1038/12618
- Shaw, R. M., and Rudy, Y. (1997). Ionic mechanisms of propagation in cardiac tissue: roles of the sodium and L-type calcium currents during reduced excitability and decreased gap junction coupling. *Circ. Res.* 81, 727–741. doi: 10.1161/01.RES.81.5.727
- Ten Tusscher, K. H. W. J., and Panfilov, A. V. (2006). Cell model for efficient simulation of wave propagation in human ventricular tissue under normal and pathological conditions. *Phys. Med. Biol.* 51, 6141–6156. doi: 10.1088/0031-9155/51/23/014
- Thomas, S. P., Kucera, J. P., Bircher-Lehmann, L., Rudy, Y., Saffitz, J. E., and Kléber, A. G. (2003). Impulse propagation in synthetic strands of neonatal cardiac myocytes with genetically reduced levels of connexin43. *Circ. Res.* 92, 1209–1216. doi: 10.1161/01.RES.0000074916.41221.EA
- Utrilla, R. G., Nieto-Marin, P., Alfayate, S., Tinaquero, D., Matamoros, M., Pérez-Hernández, M., et al. (2017). Kir2.1-Nav1.5 channel complexes are differently regulated than Kir2.1 and Nav1.5 channels alone. *Front. Physiol.* 8:903. doi: 10.3389/fphys.2017.00903
- Vacher, H., Yang, J.-W., Cerda, O., Autillo-Touati, A., Dargent, B., and Trimmer, J. S. (2011). Cdk-mediated phosphorylation of the Kv $\beta$ 2 auxiliary subunit regulates Kv1 channel axonal targeting. *J. Cell. Biol.* 192, 813–824. doi: 10.1083/jcb.201007113
- van Bemmelen, M. X., van, Rougier, J.-S., Gavillet, B., Apothéoz, F., Daidié, D., Tateyama, M., et al. (2004). Cardiac voltage-gated sodium channel Nav1.5 is regulated by Nedd4-2 mediated ubiquitination. *Circ. Res.* 95, 284–291. doi: 10.1161/01.RES.0000136816.05109.89
- Veerman, C. C., Podliesna, S., Tadros, R., Lodder, E. M., Mengarelli, I., de Jonge, B., et al. (2017). The Brugada syndrome susceptibility gene *HEY2* modulates cardiac transmural ion channel patterning and electrical heterogeneity. *Circ. Res.* 121, 537–548. doi: 10.1161/CIRCRESAHA.117.310959
- Verkerk, A. O., Veerman, C. C., Zegers, J. G., and Wilders, R. (2016). Effects of the transient outward potassium current on action potential upstroke velocities tested using the dynamic clamp technique. *Comput. Cardiol.* 43, 257–260. doi: 10.22489/CinC.2016.076-143
- Wilde, A. A. M., Postema, P. G., Di Diego, J. M., Viskin, S., Morita, H., Fish, J. M., et al. (2010). The pathophysiological mechanism underlying Brugada syndrome: depolarization versus repolarization. *J. Mol. Cell. Cardiol.* 49, 543–553. doi: 10.1016/j.yjmcc.2010.07.012
- Wilders, R. (2012). Arrhythmogenic right ventricular cardiomyopathy: considerations from *in silico* experiments. *Front. Physiol.* 3:168. doi: 10.3389/fphys.2012.00168
- You, T., Mao, W., Cai, B., Li, F., and Xu, H. (2015). Two novel Brugada syndrome-associated mutations increase Kv4.3 membrane expression and function. *Int. J. Mol. Med.* 36, 309–315. doi: 10.3892/ijmm.2015.2223
- Zhao, X.-J., Zhu, C., Tian, L.-Y., Fu, Y.-C., Zhang, Y., Chen, X., et al. (2017). Kv4.3 modulates the distribution of hERG. *Sci. Rep.* 7:17757. doi: 10.1038/s41598-017-17837-6

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Portero, Wilders, Casini, Charpentier, Verkerk and Remme. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.