



HAL
open science

THE COST OF CONTROLLING STRONGLY DEGENERATE PARABOLIC EQUATIONS

Piermarco Cannarsa, Patrick Martinez, Judith Vancostenoble

► **To cite this version:**

Piermarco Cannarsa, Patrick Martinez, Judith Vancostenoble. THE COST OF CONTROLLING STRONGLY DEGENERATE PARABOLIC EQUATIONS. ESAIM: Control, Optimisation and Calculus of Variations, In press. hal-01834720

HAL Id: hal-01834720

<https://hal.science/hal-01834720>

Submitted on 10 Jul 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE COST OF CONTROLLING STRONGLY DEGENERATE PARABOLIC EQUATIONS

P. CANNARSA, P. MARTINEZ, AND J. VANCOSTENOBLE

ABSTRACT. We consider the typical one-dimensional strongly degenerate parabolic operator $Pu = u_t - (x^\alpha u_x)_x$ with $0 < x < \ell$ and $\alpha \in (0, 2)$, controlled either by a boundary control acting at $x = \ell$, or by a locally distributed control. Our main goal is to study the dependence of the so-called controllability cost needed to drive an initial condition to rest with respect to the degeneracy parameter α . We prove that the control cost blows up with an explicit exponential rate, as $e^{C/((2-\alpha)^2 T)}$, when $\alpha \rightarrow 2^-$ and/or $T \rightarrow 0^+$.

Our analysis builds on earlier results and methods (based on functional analysis and complex analysis techniques) developed by several authors such as Fattorini-Russel, Seidman, Güichal, Tenenbaum-Tucsnak and Lissy for the classical heat equation. In particular, we use the moment method and related constructions of suitable biorthogonal families, as well as new fine properties of the Bessel functions J_ν of large order ν (obtained by ordinary differential equations techniques).

1. INTRODUCTION

1.1. Presentation of the problem and of the main results.

The aim of this paper is to study the null controllability cost for the typical 1D degenerate parabolic operator

$$(1.1) \quad Pu = u_t - (x^\alpha u_x)_x \quad (x \in (0, 1), t > 0)$$

under the action of a boundary control H :

$$(1.2) \quad \begin{cases} u_t - (x^\alpha u_x)_x = 0, & x \in (0, 1), t > 0, \\ (x^\alpha u_x)(0, t) = 0, & t > 0, \\ u(1, t) = H(t), & t > 0, \\ u(x, 0) = u_0(x), & x \in (0, 1), \end{cases}$$

and under the action of a locally distributed control h :

$$(1.3) \quad \begin{cases} u_t - (x^\alpha u_x)_x = h(x, t)\chi_{[a,b]}(x) & x \in (0, 1), t > 0, \\ (x^\alpha u_x)(0, t) = 0, & t > 0, \\ u(1, t) = 0, & t > 0, \\ u(x, 0) = u_0(x), & x \in (0, 1). \end{cases}$$

In [11], we established the following property:

1991 *Mathematics Subject Classification.* 35K65, 33C10, 93B05, 93B60, 35P10, 34B08.

Key words and phrases. degenerate parabolic equations, null controllability, moment problem, Bessel functions.

The first author was partly supported by the University of Roma Tor Vergata (Consolidate the Foundations 2015) and Istituto Nazionale di Alta Matematica (GNAMPA 2017 Research Projects). The second author was partly supported by Istituto Nazionale di Alta Matematica (GNAMPA 2017 Research Projects).

given $\alpha \geq 1$, $T > 0$, $0 < a < b < 1$, then, for any $u_0 \in L^2(0, 1)$, problem (1. 3) admits a control $h \in L^2((a, b) \times (0, T))$ that drives the solution to 0 in time $T > 0$ *if and only if* $\alpha < 2$.

In the same way,

given $\alpha \geq 1$, $T > 0$, then, for any $u_0 \in L^2(0, 1)$, problem (1. 2) admits a control $H \in L^2(0, T)$ that drives the solution to 0 in time $T > 0$ *if and only if* $\alpha < 2$.

The aim of this paper is to understand the behavior of the cost of control as:

- $\alpha \rightarrow 2^-$ (since $\alpha = 2$ is the threshold for null controllability),
- and/or $T \rightarrow 0^+$, an issue related to the so-called 'fast control problem'.

It is well-known that the cost of control blows up when $T \rightarrow 0^+$ (at least for nondegenerate parabolic equations, as we will recall in the following), and it is expected to blow up when $\alpha \rightarrow 2^-$. In this work we will prove precise upper and lower bounds for this blow-up: denoting by $u^{(h)}$ the solution of (1. 3) for any given $h \in L^2((a, b) \times (0, T))$, we prove that the null controllability costs, defined as

$$C_{ctr-bd}(\alpha, T) := \sup_{\|u_0\|_{L^2(0,1)}=1} \inf\{\|H\|_{H^1(0,T)}, u^{(H)}(T) = 0\},$$

and

$$C_{ctr-loc}(\alpha, T) := \sup_{\|u_0\|_{L^2(0,1)}=1} \inf\{\|h\|_{L^2((0,1)\times(0,T))}, u^{(h)}(T) = 0\},$$

blow up

- as $\alpha \rightarrow 2^-$,
- and/or as $T \rightarrow 0^+$,

at a precise speed: there exist positive constants C, C' such that

$$e^{-\frac{1}{C} \frac{1}{(2-\alpha)^{4/3}} (\ln \frac{1}{2-\alpha} + \ln \frac{1}{T})} e^{\frac{C}{T(2-\alpha)^2}} \leq C_{ctr-bd}(\alpha, T) \leq e^{\frac{C'}{T(2-\alpha)^2}},$$

and, in a similar way,

$$e^{-\frac{1}{C} \frac{1}{(2-\alpha)^{4/3}} (\ln \frac{1}{2-\alpha} + \ln \frac{1}{T})} e^{\frac{C}{T(2-\alpha)^2}} \leq C_{ctr-loc}(\alpha, T) \leq e^{\frac{C'}{T(2-\alpha)^2}}.$$

(See precise statements in Theorems 2.1, 2.2, 2.3 and 2.4.)

1.2. Relation to literature.

This question of the cost of null controllability when some parameter comes into play has been studied for several equations and in several situations:

- the 'fast control problem', that is, the cost of null controllability with respect to time T as $T \rightarrow 0^+$, has been investigated for the heat operator

$$(1. 4) \quad Pu = u_t - \Delta u$$

(with a boundary or localized control) and the Schrödinger equation by several authors, see, in particular, the works by Seidman et al [58, 59], Güichal [32], Miller [47, 48, 49, 50], Tenenbaum and Tucsnak [61, 62], and the more recent papers by Lissy [42] (for dispersive equations) and Benabdallah et al [5] (for parabolic systems);

- the 'vanishing viscosity limit', that is the cost of null controllability of a heat operator with the addition of a transport term when the diffusion coefficient goes to zero:

$$(1. 5) \quad P_\varepsilon u = u_t - \varepsilon u_{xx} + Mu_x$$

(again with a boundary or localized control) has been investigated by Coron and Guerrero [17], Guerrero and Lebeau [30], Glass [28], Glass and Guerrero [29], and Lissy [43];

- the 1D degenerate parabolic equation, controlled by a boundary control acting at the degeneracy point (and $\alpha \rightarrow 1^-$, 1 being the threshold value of well-posedness in this case, see [13]).

1.3. Description of the method and connection with the literature.

For the proof of our results we follow the classical strategy which consists in:

- the spectral analysis of the associated stationary operator (see Proposition 2.4) in order to determine the eigenvalues and eigenfunctions of our problem by typical ODE techniques,
- the use of the moment method, that was developed in the seminal papers by Fattorini and Russell [23, 24], to give, at least formally, a sequence of relations satisfied by the desired control,
- the construction and the properties of suitable biorthogonal families which are the main tool (at this point we will use two extensions of the results of Seidman-Avdonin-Ivanov [59] and Güichal [32], that we proved in [13] and [14]),
- the construction of suitable controls, mainly based on the behavior of the eigenfunctions of the spectral problem in the control region.

Hence a starting point is the study of the spectral problem. In the context of degenerate parabolic equations, it is classical that the eigenfunctions of the problem are expressed in terms of Bessel functions of order $\nu_\alpha = \frac{\alpha-1}{2-\alpha}$, and the eigenvalues in terms of the zeros of these Bessel functions, see Kamke [35]. This was a crucial observation in the work by Gueye [31], where the null controllability of the degenerate heat equation for $\alpha \in [0, 1)$ was addressed for the first time when the control acts at the degeneracy point, and in [13] where we obtained optimal bounds for the cost of control for such a problem. For strongly degenerate parabolic equations, an additional source of difficulty is that the order of the useful Bessel functions blows up as $\alpha \rightarrow 2^-$. To cope with such difficulties several classical results from Watson [64] and Qu-Wong [55] will be needed.

It turns out that there is a common phenomenon in the classical fast control problem ((1. 4) when $T \rightarrow 0^+$), the vanishing viscosity problem ((1. 5) when $\varepsilon \rightarrow 0^+$), and the null controllability of the degenerate parabolic equation (1. 1) when the degeneracy parameter approaches its critical value: *the eigenvalues concentrate when parameters go to their critical values*. Such a concentration phenomenon can be observed:

- for the vanishing viscosity problem, in [17];
- for degenerate parabolic equations, once the eigenvalues have been computed, in Lemma 5.1;
- for the classical heat equation and the fast control problem, once time has been renormalized to a fixed value, see Remark 5.1.

This common feature is the key point in understanding the behavior of the control in every context. Indeed, the construction of suitable biorthogonal families is strongly related to gap properties: the gap $\lambda_{n+1} - \lambda_n \rightarrow 0$ when the degeneracy parameter goes to its critical limit, and the speed at which it goes to zero govern the upper and lower estimates for the associated biorthogonal families (since the norm of such biorthogonal families involve large products of the inverse of such differences), hence for the null controllability cost.

In the context of degenerate parabolic equations, in order to obtain optimal bounds,

- we refine classical results providing sharp gap estimates for the zeros of Bessel functions of large order, see Lemma 5.2,

- we will combine these gap estimates with some recent results [13, 14] that complete classical results of Fattorini-Russel [24], obtaining explicit and precise (upper and lower) estimates for biorthogonal families, even in short time, under some gap conditions, namely:

$$\begin{cases} \forall n \geq 1, & \gamma_{min} \leq \sqrt{\lambda_{n+1}} - \sqrt{\lambda_n} \leq \gamma_{max} \\ \forall n \geq N^*, & \gamma_{min}^* \leq \sqrt{\lambda_{n+1}} - \sqrt{\lambda_n} \leq \gamma_{max}^* \end{cases}$$

these gap conditions are a little more general than the asymptotic development of the eigenvalues used in Tenenbaum-Tucsnak [61] and Lissy [42, 43]:

$$\lambda_n = rn^2 + O(n),$$

but, which is more important, they allow us to obtain precise estimates for the biorthogonal family when some parameter comes into play, as it happens here or in 2D problems (see, e.g., [4, 25]); (the proof of the general results obtained in [13, 14] concerning biorthogonal families is based on some complex and hilbertian analysis techniques developed by Seidman-Avdonin-Ivanov [59], Güichal [32] and the adjonction of some well-chosen parameter, inspired from Tenenbaum-Tucsnak [61]);

- we complete the analysis of the asymptotic behavior of Bessel functions of large order, see Proposition 2.5; this issue is related to the so-called 'transition zone' (see Watson [64] and Krasikov [37]) and recent results of Privat-Trélat-Zuazua [53] p. 957, even though we prove Proposition 2.5 directly by estimating the norm of the solution of a second-order differential equation depending on a large parameter.

1.4. Plan of the paper.

The plan of the paper is the following.

- In section 2, we state our main results concerning the null controllability costs (Theorems 2.1-2.4) and the spectral properties of the problem (see Propositions 2.4 and 2.5).
- In section 3, we recall the main properties of Bessel functions, and prove Propositions 2.4 and 2.2.
- In section 4, we establish useful identities by the moment method.
- In section 5, we prove Theorem 2.1; the proof is based on a recent result [14] based on hilbertian techniques developed by Güichal [32], the concentration of the eigenvalues (Lemma 5.1) and a precised form of a classical property concerning the gap of the zeros of the Bessel function of large parameter (Lemma 5.2).
- In section 6 we prove Theorem 2.2; the proof is based on the construction of some suitable biorthogonal family, based on [13] and inspired by the construction of Seidman et al [59].
- In section 7, we prove Theorem 2.4, which is a direct consequence of the biorthogonal family constructed in section 6, assuming Proposition 2.5.
- In section 8, we prove Theorem 2.3; the proof is based on energy methods and uses Theorem 2.1.
- In section 9, we study the eigenfunctions in the control region, proving Proposition 2.5; the proof is based on ODE techniques.

2. THE COST OF NULL CONTROLLABILITY: MAIN RESULTS

2.1. The controllability problems.

We study the cost of null controllability of a degenerate parabolic equation, using either a boundary control acting at the non degeneracy point or a locally distributed

control. More precisely, we fix $\ell > 0$, $\alpha \geq 1$, $T > 0$, and for any $u_0 \in L^2(0, \ell)$, we wish to find a control H such that the solution of

$$(2.1) \quad \begin{cases} u_t - (x^\alpha u_x)_x = 0, & x \in (0, \ell), t > 0, \\ (x^\alpha u_x)(0, t) = 0, & t > 0, \\ u(\ell, t) = H(t), & t > 0, \\ u(x, 0) = u_0(x), & x \in (0, \ell), \end{cases}$$

also satisfies $u(\cdot, T) = 0$.

Similarly, given $0 < a < b < \ell$ and for any $u_0 \in L^2(0, \ell)$, we wish to find a control h such that the solution of

$$(2.2) \quad \begin{cases} u_t - (x^\alpha u_x)_x = h(x, t)\chi_{[a, b]}(x) & x \in (0, \ell), t > 0, \\ (x^\alpha u_x)(0, t) = 0, & t > 0, \\ u(\ell, t) = 0, & t > 0, \\ u(x, 0) = u_0(x), & x \in (0, \ell), \end{cases}$$

also satisfies $u(\cdot, T) = 0$.

In space dimension 1, these two problems are very close. From [11], we know that such controls exist if and only if $\alpha \in [1, 2)$.

2.2. Functional setting and well-posedness.

2.2.a. Functional setting and well-posedness for a locally distributed control.

For $1 \leq \alpha < 2$, we consider the following spaces :

$$H_\alpha^1(0, \ell) := \{u \in L^2(0, \ell) \mid u \text{ locally absolutely continuous in } (0, \ell], x^{\alpha/2}u_x \in L^2(0, \ell)\},$$

$$H_{\alpha,0}^1(0, \ell) := \{u \in H_\alpha^1(0, \ell) \mid u(\ell) = 0\},$$

and

$$H_\alpha^2(0, \ell) := \{u \in H_\alpha^1(0, \ell) \mid x^\alpha u_x \in H^1(0, \ell)\}$$

Then, the operator $A : D(A) \subset L^2(0, \ell) \rightarrow L^2(0, \ell)$ will be defined by

$$\begin{cases} \forall u \in D(A), & Au := (x^\alpha u_x)_x, \\ D(A) := \{u \in H_{\alpha,0}^1(0, \ell) \mid x^\alpha u_x \in H^1(0, \ell)\} = H_\alpha^2(0, \ell) \cap H_{\alpha,0}^1(0, \ell), \\ & = \{u \in L^2(0, \ell) \mid u \text{ locally absolutely continuous in } (0, \ell], \\ & \quad x^\alpha u \in H_0^1(0, \ell), x^\alpha u_x \in H^1(0, \ell) \text{ and } (x^\alpha u_x)(0) = 0\}. \end{cases}$$

Notice that, if $u \in D(A)$, then u satisfies the Neumann boundary condition $(x^\alpha u_x)(0) = 0$ at $x = 0$ and the Dirichlet boundary condition $u(\ell) = 0$ at $x = \ell$.

Then the following results hold, (see, e.g., [7] and [10]).

Proposition 2.1. *$A : D(A) \subset L^2(0, \ell) \rightarrow L^2(0, \ell)$ is a self-adjoint negative operator with dense domain.*

Hence, A is the infinitesimal generator of an analytic semigroup of contractions e^{tA} on $L^2(0, \ell)$. Given a source term h in $L^2((0, \ell) \times (0, T))$ and an initial condition $v_0 \in L^2(0, \ell)$, consider the problem

$$(2.3) \quad \begin{cases} v_t - (x^\alpha v_x)_x = h(x, t), \\ (x^\alpha v_x)(0, t) = 0, \\ v(\ell, t) = 0, \\ v(x, 0) = v_0(x). \end{cases}$$

The function $v \in C^0([0, T]; L^2(0, \ell)) \cap L^2(0, T; H_{\alpha, 0}^1(0, \ell))$ given by the variation of constant formula

$$v(\cdot, t) = e^{tA}v_0 + \int_0^t e^{(t-s)A}h(\cdot, s) ds$$

is called the mild solution of (2. 3). We say that a function

$$v \in C^0([0, T]; H_{\alpha, 0}^1(0, \ell)) \cap H^1(0, T; L^2(0, \ell)) \cap L^2(0, T; D(A))$$

is a strict solution of (2. 3) if v satisfies $v_t - (x^\alpha v_x)_x = h(x, t)$ almost everywhere in $(0, \ell) \times (0, T)$, and the initial and boundary conditions for all $t \in [0, T]$ and all $x \in [0, \ell]$.

Proposition 2.2. *If $v_0 \in H_{\alpha, 0}^1(0, \ell)$, then the mild solution of (2. 3) is the unique strict solution of (2. 3).*

The proof of Proposition 2.2 follows from classical results, see subsection 3.4.

2.2.b. *Functional setting and well-posedness for a boundary control.*

To define the solution of the boundary value problem (2. 1), we transform it into a problem with homogeneous boundary conditions and a source term (depending on the control h): formally, if u is a solution of (2. 1), then the function v defined by

$$(2. 4) \quad v(x, t) = u(x, t) - \frac{x^{2-\alpha}}{\ell^{2-\alpha}} H(t)$$

satisfies the auxiliary problem

$$(2. 5) \quad \begin{cases} v_t - (x^\alpha v_x)_x = -\frac{x^{2-\alpha}}{\ell^{2-\alpha}} H'(t) + \frac{2-\alpha}{\ell^{2-\alpha}} H(t), \\ (x^\alpha v_x)(0, t) = 0, \\ v(\ell, t) = 0, \\ v(x, 0) = u_0(x) - \frac{x^{2-\alpha}}{\ell^{2-\alpha}} H(0). \end{cases}$$

Reciprocally, given $H \in H^1(0, T)$, consider the solution v of

$$(2. 6) \quad \begin{cases} v_t - (x^\alpha v_x)_x = -\frac{x^{2-\alpha}}{\ell^{2-\alpha}} H'(t) + \frac{2-\alpha}{\ell^{2-\alpha}} H(t), \\ (x^\alpha v_x)(0, t) = 0, \\ v(\ell, t) = 0, \\ v(x, 0) = v_0(x). \end{cases}$$

Then the function u defined by

$$(2. 7) \quad u(x, t) = v(x, t) + \frac{x^{2-\alpha}}{\ell^{2-\alpha}} H(t)$$

satisfies

$$(2. 8) \quad \begin{cases} u_t - (x^\alpha u_x)_x = 0, \\ (x^\alpha u_x)(0, t) = 0, \\ u(\ell, t) = H(t), \\ u(x, 0) = v_0(x) + \frac{x^{2-\alpha}}{\ell^{2-\alpha}} H(0). \end{cases}$$

This motivates the following definition of what is the solution of the boundary value problem (2. 1):

Definition 2.1. *a) We say that $u \in C([0, T]; L^2(0, \ell)) \cap L^2(0, T; H_\alpha^1(0, \ell))$ is the mild solution of (2. 1) if v defined by (2. 4) is the mild solution of (2. 5).*

b) We say that

$$u \in C([0, T]; H_\alpha^1(0, \ell)) \cap H^1(0, T; L^2(0, \ell)) \cap L^2(0, T; H_\alpha^2(0, \ell))$$

is the strict solution of (2. 1) if v defined by (2. 4) is the strict solution of (2. 5).

Then we immediately obtain

Proposition 2.3. *a) Given $u_0 \in L^2(0, \ell)$, $H \in H^1(0, T)$, problem (2. 1) admits a unique mild solution.*

b) Given $u_0 \in H_{\alpha,0}^1(0, \ell)$, $H \in H^1(0, T)$ such that $u_0(\ell) = H(0)$, problem (2. 1) admits a unique strict solution.

The proof of Proposition 2.3 follows immediately, noting that

$$\tilde{H}(x, t) := \frac{x^{2-\alpha}}{\ell^{2-\alpha}} H(t)$$

satisfies

$$\tilde{H} \in C([0, T]; H_{\alpha}^1(0, \ell)) \cap H^1(0, T; L^2(0, \ell)) \cap L^2(0, T; H_{\alpha}^2(0, \ell)).$$

2.3. Null controllability results for the boundary control.

Consider

$$(2. 9) \quad C_{ctr-bd}(\alpha, T, \ell) := \sup_{\|u_0\|_{L^2(0, \ell)}=1} \inf\{\|H\|_{H^1(0, T)}, u^{(H)}(T) = 0\},$$

where $u^{(H)}$ is the solution of problem (2. 1). Then we prove the following estimates:

2.3.a. *Lower bound of the null controllability cost.*

Theorem 2.1. *There exists a constant $C_u > 0$ independent of $\alpha \in [1, 2)$, of $\ell > 0$ and of $T > 0$ such that*

$$(2. 10) \quad C_{ctr-bd}(\alpha, T, \ell) \geq C_u \frac{\ell^{2-\alpha}}{\sqrt{(2-\alpha)T\ell}} e^{-\pi^2 \frac{T}{\ell^{2-\alpha}}} e^{C_u \frac{\ell^{2-\alpha}}{T(2-\alpha)^2}} e^{-\frac{1}{C_u} \left(\frac{1}{(2-\alpha)^{4/3}} + \frac{\ell^{1-\alpha/2}}{2-\alpha} \right) (\ln \frac{\ell^{1-\alpha/2}}{2-\alpha} + \ln \frac{1}{T})}.$$

Remark 2.1. This proves that the cost blows up when $T \rightarrow 0^+$, or $\alpha \rightarrow 2^-$, or $\ell \rightarrow +\infty$, and at least exponentially fast. When ℓ is fixed and $T \leq T_0$, this simplifies into

$$C_{ctr-bd}(\alpha, T, \ell) \geq C_u e^{\frac{C_u}{T(2-\alpha)^2}} e^{-\frac{1}{C_u} \frac{1}{(2-\alpha)^{4/3}} (\ln \frac{1}{2-\alpha} + \ln \frac{1}{T})}.$$

2.3.b. *Upper bound of the null controllability cost.*

Theorem 2.2. *There exists a constant $C_u > 0$ independent of $\alpha \in [1, 2)$, of $\ell > 0$ and of $T > 0$ such that*

$$(2. 11) \quad C_{ctr-bd}(\alpha, T, \ell) \leq \frac{C_u}{\sqrt{(2-\alpha)T\ell}} e^{-\frac{1}{C_u} \frac{T}{\ell^{2-\alpha}}} e^{C_u \frac{\ell^{2-\alpha}}{T(2-\alpha)^2}}.$$

Remark 2.2. This proves that the cost blows up exactly exponentially fast as $T \rightarrow 0^+$, or $\alpha \rightarrow 2^-$, or $\ell \rightarrow +\infty$. When ℓ is fixed and $T \leq T_0$, this simplifies into

$$C_{ctr-bd}(\alpha, T, \ell) \leq \frac{C_u}{\sqrt{(2-\alpha)T}} e^{\frac{C_u}{T(2-\alpha)^2}}.$$

2.4. Null controllability results for the locally distributed control.

Consider

$$(2. 12) \quad C_{ctr-loc}(\alpha, T, \ell) := \sup_{\|u_0\|_{L^2(0, \ell)}=1} \inf\{\|h\|_{L^2((a,b) \times ((0, T)))}, u^{(h)}(T) = 0\},$$

where $u^{(h)}$ is the solution of problem (2. 2). For this problem, we are mainly interested in the dependence with respect to the degeneracy ($\alpha \rightarrow 2^-$) and to time (fast controls, when $T \rightarrow 0$), see Remarks 2.3 and 2.4. And we prove the following estimates:

2.4.a. *Lower bound of the null controllability cost.*

Theorem 2.3. *Given $\ell > 0$, and $0 < a < b < \ell$, there exists a constant $\tilde{C} = \tilde{C}(a, b, \ell) > 0$ independent of $\alpha \in [1, 2)$ and of $T > 0$ such that*

$$(2.13) \quad C_{ctr-loc}(\alpha, T, \ell) \geq \tilde{C} e^{\frac{\tilde{C}}{T(2-\alpha)^2}} e^{-\frac{1}{\tilde{C}} \frac{1}{(2-\alpha)^{4/3}} (\ln \frac{1}{2-\alpha} + \ln \frac{1}{T}) - \frac{1}{\tilde{C}} T} - 1.$$

Remark 2.3. In the proof of Theorem 2.3 we obtain an explicit expression of $\tilde{C}(a, b, \ell)$. And of course Theorem 2.3 proves that the cost blows up (exponentially fast) when $T \rightarrow 0^+$, or $\alpha \rightarrow 2^-$: when $T \rightarrow 0$ and/or $\alpha \rightarrow 2^-$, this simplifies into

$$C_{ctr-loc}(\alpha, T, \ell) \geq \tilde{C} e^{\frac{\tilde{C}}{T(2-\alpha)^2}} e^{-\frac{1}{\tilde{C}} \frac{1}{(2-\alpha)^{4/3}} (\ln \frac{1}{2-\alpha} + \ln \frac{1}{T})}.$$

2.4.b. *Upper bound of the null controllability cost.*

Theorem 2.4. *There exists a constant $C_u = C_u > 0$ independent of $\alpha \in [1, 2)$, of $T > 0$ and of $0 < a < b < \ell$, and $\gamma_0^* = \gamma_0^*(a, b, \ell) > 0$ such that*

$$(2.14) \quad C_{ctr-loc}(\alpha, T, \ell) \leq \frac{C_u}{\gamma_0^*} e^{C_u \frac{\ell^{2-\alpha}}{T(2-\alpha)^2}} e^{-\frac{1}{C_u} \frac{T}{\ell^{2-\alpha}}} \max\left\{\frac{1}{\sqrt{T}(2-\alpha)}, \frac{1}{\ell^{1-\alpha/2}}\right\}.$$

Remark 2.4. This proves that the cost blows up exactly exponentially fast as $T \rightarrow 0^+$, or $\alpha \rightarrow 2^-$, or $\ell \rightarrow +\infty$. When ℓ is fixed and $T \leq T_0$, this simplifies into

$$C_{ctr-loc}(\alpha, T, \ell) \leq C_u e^{C_u \frac{1}{T(2-\alpha)^2}}.$$

2.5. The eigenvalue problem.

The knowledge of the eigenvalues and associated eigenfunctions of the degenerate diffusion operator $u \mapsto -(x^\alpha u)'$, i.e. the solutions (λ, Φ) of

$$(2.15) \quad \begin{cases} -(x^\alpha \Phi'(x))' = \lambda \Phi(x) & x \in (0, \ell), \\ (x^\alpha \Phi'(x))(0) = 0, \\ \Phi(\ell) = 0. \end{cases}$$

will be essential for our purposes.

2.5.a. Eigenvalues and eigenfunctions.

It is well-known that Bessel functions play an important role in this problem, see, e.g., Kamke [35]. For $\alpha \in [1, 2)$, let

$$\nu_\alpha := \frac{\alpha - 1}{2 - \alpha}, \quad \kappa_\alpha := \frac{2 - \alpha}{2}.$$

Given $\nu \geq 0$, we denote by J_ν the Bessel function of first kind and of order ν (see section 3.2) and denote $j_{\nu,1} < j_{\nu,2} < \dots < j_{\nu,n} < \dots$ the sequence of positive zeros of J_ν . Then we have the following:

Proposition 2.4. *The eigenvalues λ for problem (2.15) are given by*

$$(2.16) \quad \forall n \geq 1, \quad \lambda_{\alpha,n} = \ell^{\alpha-2} \kappa_\alpha^2 j_{\nu_\alpha,n}^2$$

and the corresponding normalized (in $L^2(0, \ell)$) eigenfunctions takes the form

$$(2.17) \quad \Phi_{\alpha,n}(x) = \frac{\sqrt{2\kappa_\alpha}}{\ell^{\kappa_\alpha} |J'_{\nu_\alpha}(j_{\nu_\alpha,n})|} x^{(1-\alpha)/2} J_{\nu_\alpha}(j_{\nu_\alpha,n} \frac{x}{\ell})^{\kappa_\alpha}, \quad x \in (0, \ell).$$

Moreover the family $(\Phi_{\alpha,n})_{n \geq 1}$ forms an orthonormal basis of $L^2(0, \ell)$.

Remark 2.5. Gueye [31] proved Proposition 2.4 in the case $\alpha \in [0, 1)$ and when $\ell = 1$. The case $\alpha \in [1, 2)$ and $\ell \neq 1$ is very similar.

2.5.b. *The eigenfunctions in the control region.*

We will prove the following property:

Proposition 2.5. *Given $0 < a < b < \ell$, there exists $\gamma_0^* = \gamma_0^*(a, b, \ell) > 0$ such that*

$$(2.18) \quad \forall \alpha \in [1, 2), \forall m \geq 1, \quad \int_a^b \Phi_{\alpha, m}(x)^2 dx \geq \gamma_0^*(2 - \alpha).$$

It is classical in the nondegenerate case (Lagnese [38]) that

$$\inf_m \int_a^b \Phi_{\alpha, m}^2 > 0;$$

but, in our purpose of estimating the cost of null controllability, it is necessary to have a lower bound of $\int_a^b \Phi_{\alpha, m}^2$ with respect to the degeneracy parameter α when $\alpha \rightarrow 2^-$, and the dependence is given in Proposition 2.5. This does not come easily, since $\Phi_{\alpha, m}$ is solution of a second-order differential equation depending on a large parameter. We will overcome this difficulty with ODE techniques.

3. PROOF OF PROPOSITIONS 2.4 AND 2.2

In this section, we study the spectral problem (2.15) and the properties of the eigenvalues and eigenfunctions, and as a first application we deduce the well-posedness result stated in Proposition 2.2.

Let us study the spectral problem. First, one can observe that if λ is an eigenvalue, then $\lambda > 0$: indeed, multiplying (2.15) by Φ and integrating by parts, then

$$\lambda \int_0^\ell \Phi^2 = \int_0^\ell x^\alpha \Phi_x^2,$$

which implies first $\lambda \geq 0$, and next that $\Phi = 0$ if $\lambda = 0$.

3.1. The link with the Bessel's equation.

There is a change a variables that allows one to transform the eigenvalue problem (2.15) into a differential Bessel's equation (see in particular Kamke [35, section 2.162, equation (Ia), p. 440], and Gueye [31]): assume that Φ is a solution of (2.15) associated to the eigenvalue λ ; then one easily checks that the function Ψ defined by

$$(3.1) \quad \Phi(x) =: x^{\frac{1-\alpha}{2}} \Psi\left(\frac{2}{2-\alpha} \sqrt{\lambda} x^{\frac{2-\alpha}{2}}\right)$$

is solution of the following boundary problem:

$$(3.2) \quad \begin{cases} y^2 \Psi''(y) + y \Psi'(y) + (y^2 - (\frac{\alpha-1}{2-\alpha})^2) \Psi(y) = 0, & y \in (0, \frac{2}{2-\alpha} \sqrt{\lambda} \ell^{\kappa_\alpha}), \\ (2-\alpha) y^{\frac{1}{2-\alpha}} \Psi'(y) - (\alpha-1) y^{\frac{\alpha-1}{2-\alpha}} \Psi(y) \rightarrow 0 \text{ as } y \rightarrow 0, \\ \Psi\left(\frac{2}{2-\alpha} \sqrt{\lambda} \ell^{\frac{2-\alpha}{2}}\right) = 0. \end{cases}$$

3.2. Bessel's equation and Bessel's functions of order ν . For reader convenience, we recall here the definitions concerning Bessel's equation and functions together with some useful properties of these functions and of their zeros. *Throughout this section, we assume that $\nu \in \mathbb{R}_+$.*

3.2.a. Bessel's equation and Bessel's functions of order ν .

The Bessel's functions of order ν are the solutions of the following differential equation (see [64, section 3.1, eq. (1), p. 38] or [41, eq (5.1.1), p. 98]):

$$(3.3) \quad y^2 \Psi''(y) + y \Psi'(y) + (y^2 - \nu^2) \Psi(y) = 0, \quad y \in (0, +\infty).$$

The above equation is called *Bessel's equation for functions of order ν* . Of course the fundamental theory of ordinary differential equations says that the solutions of (3.3) generate a vector space S_ν of dimension 2. In the following we recall what can be chosen as a basis of S_ν .

3.2.b. Fundamental solutions of Bessel's equation when $\nu \notin \mathbb{N}$.

Assume that $\nu \notin \mathbb{N}$. When looking for solutions of (3.3) of the form of series of ascending powers of y , one can construct two series that are solutions:

$$\sum_{m \geq 0} \frac{(-1)^m}{m! \Gamma(\nu + m + 1)} \left(\frac{y}{2}\right)^{\nu+2m} \quad \text{and} \quad \sum_{m \geq 0} \frac{(-1)^m}{m! \Gamma(-\nu + m + 1)} \left(\frac{y}{2}\right)^{-\nu+2m},$$

where Γ is the Gamma function (see [64, section 3.1, p. 40]). The first of the two series converges for all values of y and defines the so-called Bessel function of order ν and *of the first kind* which is denoted by J_ν :

$$(3.4) \quad J_\nu(y) := \sum_{m=0}^{\infty} \frac{(-1)^m}{m! \Gamma(m + \nu + 1)} \left(\frac{y}{2}\right)^{2m+\nu} = \sum_{m=0}^{\infty} c_{\nu,m}^+ y^{2m+\nu}, \quad y \geq 0,$$

(see [64, section 3.1, (8), p. 40] or [41, eq. (5.3.2), p. 102]). The second series converges for all positive values of y and is evidently $J_{-\nu}$:

$$(3.5) \quad J_{-\nu}(y) := \sum_{m=0}^{\infty} \frac{(-1)^m}{m! \Gamma(m - \nu + 1)} \left(\frac{y}{2}\right)^{2m-\nu} = \sum_{m=0}^{\infty} c_{\nu,m}^- y^{2m-\nu}, \quad y > 0.$$

When $\nu \notin \mathbb{N}$, the two functions J_ν and $J_{-\nu}$ are linearly independent and therefore the pair $(J_\nu, J_{-\nu})$ forms a fundamental system of solutions of (3.3), (see [64, section 3.12, eq. (2), p. 43]).

3.2.c. Fundamental solutions of Bessel's equation when $\nu = n \in \mathbb{N}$.

Assume that $\nu = n \in \mathbb{N}$. When looking for solutions of (3.3) of the form of series of ascending powers of y , one sees that J_n and J_{-n} are still solutions of (3.3), where J_n is still given by (3.4) and J_{-n} is given by (3.5); when $\nu = n \in \mathbb{N}$, J_{-n} can be written

$$(3.6) \quad J_{-n}(y) = \sum_{m \geq n} \frac{(-1)^m}{m! \Gamma(m - n + 1)} \left(\frac{y}{2}\right)^{-n+2m}.$$

However now $J_{-n}(y) = (-1)^n J_n(y)$, hence J_n and J_{-n} are linearly dependent, (see [64, section 3.12, p. 43] or [41, eq. (5.4.10), p. 105]). The determination of a fundamental system of solutions in this case requires further investigation. In this purpose, one introduces the Bessel's functions of order ν and *of the second kind*: among the several definitions of Bessel's functions of second order, we recall here the definition by Weber. The Bessel's functions of order ν and *of second kind* are denoted by Y_ν and defined by (see [64, section 3.54, eq. (1)-(2), p. 64] or [41, eq. (5.4.5)-(5.4.6), p. 104]):

$$\begin{cases} \forall \nu \notin \mathbb{N}, & Y_\nu(y) := \frac{J_\nu(y) \cos(\nu\pi) - J_{-\nu}(y)}{\sin(\nu\pi)}, \\ \forall n \in \mathbb{N}, & Y_n(y) := \lim_{\nu \rightarrow n} Y_\nu(y). \end{cases}$$

For any $\nu \in \mathbb{R}_+$, the two functions J_ν and Y_ν always are linearly independent, see [64, section 3.63, eq. (1), p. 76]. In particular, in the case $\nu = n \in \mathbb{N}$, the

pair (J_n, Y_n) forms a fundamental system of solutions of the Bessel's equation for functions of order n .

In the case $\nu = n \in \mathbb{N}$, it will be useful to expand Y_n under the form of a series of ascending powers. This can be done using Hankel's formula, see [64, section 3.52, eq. (3), p. 62] or [41, eq. (5.5.3), p. 107]:

$$(3.7) \quad \forall n \in \mathbb{N}^*, \quad Y_n(y) = \frac{2}{\pi} J_n(y) \log\left(\frac{y}{2}\right) - \frac{1}{\pi} \sum_{m=0}^{n-1} \frac{(n-m-1)!}{m!} \left(\frac{y}{2}\right)^{2m-n} \\ - \frac{1}{\pi} \sum_{m=0}^{+\infty} \frac{(-1)^m}{m!(n+m)!} \left(\frac{y}{2}\right)^{n+2m} \left[\frac{\Gamma'(m+1)}{\Gamma(m+1)} + \frac{\Gamma'(m+n+1)}{\Gamma(m+n+1)} \right],$$

where $\frac{\Gamma'}{\Gamma}$ is the logarithmic derivative of the Gamma function, and satisfies $\frac{\Gamma'(1)}{\Gamma(1)} = -\gamma$ (here γ denotes Euler's constant) and

$$\frac{\Gamma'(m+1)}{\Gamma(m+1)} = 1 + \frac{1}{2} + \dots + \frac{1}{m} - \gamma \quad \text{for all } m \in \mathbb{N}.$$

In the case $n = 0$, the first sum in (3.7) should be set equal to zero.

3.2.d. Zeros of Bessel functions of order ν of the first kind.

The function J_ν has an infinite number of real zeros which are simple with the possible exception of $x = 0$ ([64, section 15.21, p. 478-479 applied to $C_\nu = J_\nu$] or [41, section 5.13, Theorem 2, p. 127]). We denote by $(j_{\nu,n})_{n \geq 1}$ the strictly increasing sequence of the positive zeros of J_ν :

$$0 < j_{\nu,1} < j_{\nu,2} < \dots < j_{\nu,n} < \dots$$

and we recall that

$$j_{\nu,n} \rightarrow +\infty \text{ as } n \rightarrow +\infty.$$

We will also often use the following bounds on the zeros, proved in Lorch and Muldoon [44]:

$$(3.8) \quad \forall \nu \geq \frac{1}{2}, \forall n \geq 1, \quad \pi\left(n + \frac{\nu}{4} - \frac{1}{8}\right) \leq j_{\nu,n} \leq \pi\left(n + \frac{\nu}{2} - \frac{1}{4}\right).$$

Note also that ([44]):

$$(3.9) \quad \forall \nu \in [0, \frac{1}{2}], \forall n \geq 1, \quad \pi\left(n + \frac{\nu}{2} - \frac{1}{4}\right) \leq j_{\nu,n} \leq \pi\left(n + \frac{\nu}{4} - \frac{1}{8}\right).$$

We will also use the following asymptotic development of the first zero $j_{\nu,1}$ of J_ν with respect to ν ([64, section 15.81, p. 516]) when $\nu \rightarrow +\infty$:

$$j_{\nu,1} = \nu + 1, 855757\nu^{1/3} + O(1),$$

and a similar development for $j_{\nu,2}$, extracted from [55] where it is proved that

$$(3.10) \quad \nu - \frac{a_k}{2^{1/3}} \nu^{1/3} < j_{\nu,k} < \nu - \frac{a_k}{2^{1/3}} \nu^{1/3} + \frac{3}{20} a_k^2 \frac{2^{1/3}}{\nu^{1/3}},$$

which is valid for all $\nu > 0$, all $k \geq 1$, and where a_k is the k -th negative zero of the Airy function.

3.3. Proof of Proposition 2.4.

As noted before, $\lambda = 0$ is not an admissible eigenvalue for problem (2.15), hence $\lambda > 0$. So, using (3.1), we can transform problem (2.15) into problem (3.2). In the following, because of the difference in the construction of a fundamental system of solutions of (3.3), we treat the following cases separately: $\nu_\alpha \notin \mathbb{N}$, $\nu_\alpha = n \in \mathbb{N}^*$ and $\nu_\alpha = 0$.

3.3.a. *Case $\nu_\alpha \notin \mathbb{N}$.* Let us assume that $\nu_\alpha \notin \mathbb{N}$. Then we have

$$\Phi = C_+ \Phi_+ + C_- \Phi_-$$

where Φ_+ and Φ_- are defined by

$$\Phi_+(x) := x^{\frac{1-\alpha}{2}} J_{\nu_\alpha} \left(\frac{2}{2-\alpha} \sqrt{\lambda} x^{\frac{2-\alpha}{2}} \right), \quad \Phi_-(x) := x^{\frac{1-\alpha}{2}} J_{-\nu_\alpha} \left(\frac{2}{2-\alpha} \sqrt{\lambda} x^{\frac{2-\alpha}{2}} \right).$$

Using the series expansion of J_{ν_α} and $J_{-\nu_\alpha}$, one obtains

$$(3.11) \quad \Phi_+(x) = \sum_{m=0}^{\infty} \tilde{c}_{\nu_\alpha, m}^+ x^{(2-\alpha)m}, \quad \Phi_-(x) = \sum_{m=0}^{\infty} \tilde{c}_{\nu_\alpha, m}^- x^{1-\alpha+(2-\alpha)m},$$

where the coefficients $\tilde{c}_{\nu_\alpha, m}^+$ and $\tilde{c}_{\nu_\alpha, m}^-$ are defined by

$$(3.12) \quad \tilde{c}_{\nu_\alpha, m}^+ := c_{\nu_\alpha, m}^+ \left(\frac{2}{2-\alpha} \sqrt{\lambda} \right)^{2m+\nu_\alpha}, \quad \tilde{c}_{\nu_\alpha, m}^- := c_{\nu_\alpha, m}^- \left(\frac{2}{2-\alpha} \sqrt{\lambda} \right)^{2m-\nu_\alpha}.$$

We deduce that

$$\Phi_+(x) \sim_0 \tilde{c}_{\nu_\alpha, 0}^+, \quad x^{\alpha/2} \Phi_+'(x) \sim_0 (2-\alpha) \tilde{c}_{\nu_\alpha, 1}^+ x^{1-\alpha/2},$$

$$\Phi_-(x) \sim_0 \tilde{c}_{\nu_\alpha, 0}^-, \quad x^{\alpha/2} \Phi_-'(x) \sim_0 (1-\alpha) \tilde{c}_{\nu_\alpha, 0}^- x^{-\alpha/2},$$

hence $\Phi_+ \in H_\alpha^1(0, \ell)$, while $\Phi_- \notin H_\alpha^1(0, \ell)$. Therefore, $\Phi = C_+ \Phi_+ + C_- \Phi_- \in H_\alpha^1(0, \ell)$ implies that $C_- = 0$ and $\Phi = C_+ \Phi_+$. Moreover, $x^\alpha \Phi_+'(x) \rightarrow 0$ as $x \rightarrow 0$, hence the boundary condition in 0 is automatically satisfied. Finally, the boundary condition $\Phi(\ell) = 0$ implies that there is some C_+ and some $m \in \mathbb{N}$, $m \geq 1$ such that

$$\lambda = \kappa_\alpha^2 j_{\nu_\alpha, m}^2 \ell^{\alpha-2} \quad \text{and} \quad \Phi(x) = C_+ x^{\frac{1-\alpha}{2}} J_{\nu_\alpha} \left(j_{\nu_\alpha, m} \left(\frac{x}{\ell} \right)^{\kappa_\alpha} \right).$$

In the same way, any $\Phi(x) := C x^{\frac{1-\alpha}{2}} J_{\nu_\alpha} \left(j_{\nu_\alpha, m} \left(\frac{x}{\ell} \right)^{\kappa_\alpha} \right)$ is solution of (2.15), and the family $(\Phi_n(x) := x^{\frac{1-\alpha}{2}} J_{\nu_\alpha} \left(j_{\nu_\alpha, n} \left(\frac{x}{\ell} \right)^{\kappa_\alpha} \right))_n$ forms an orthogonal family of $L^2(0, \ell)$, which is complete since the family is composed by the eigenfunctions of the operator T_α :

$$T_\alpha : L^2(0, \ell) \rightarrow L^2(0, \ell), \quad f \mapsto T_\alpha(f) := u_f$$

where $u_f \in D(A)$ is the solution of the problem $-Au_f = f$, which is self-adjoint and compact (Appendix in [1]). Finally, it remains to norm this orthogonal family:

$$\begin{aligned} \int_0^\ell x^{1-\alpha} J_{\nu_\alpha}^2 \left(j_{\nu_\alpha, n} \left(\frac{x}{\ell} \right)^{\kappa_\alpha} \right) dx &= \ell^{2-\alpha} \int_0^1 y^{1-\alpha} J_{\nu_\alpha}^2 \left(j_{\nu_\alpha, n} y^{\kappa_\alpha} \right) dy \\ &= \frac{\ell^{2-\alpha}}{\kappa_\alpha} \int_0^1 z J_{\nu_\alpha}^2 \left(j_{\nu_\alpha, n} z \right) dz = \frac{\ell^{2-\alpha}}{\kappa_\alpha} \frac{1}{2} [J_{\nu_\alpha+1}(j_{\nu_\alpha, n})]^2 = \frac{\ell^{2-\alpha} [J_{\nu_\alpha}'(j_{\nu_\alpha, n})]^2}{2\kappa_\alpha}, \end{aligned}$$

which gives us that the family given by (2.17) forms an orthonormal basis of $L^2(0, \ell)$. This ends the proof of Proposition 2.4 when $\alpha \in [1, 2)$ is such that $\nu_\alpha \notin \mathbb{N}$.

3.3.b. *Case $\nu_\alpha = n_\alpha \in \mathbb{N}^*$.* Let us assume that $\nu_\alpha = n_\alpha \in \mathbb{N}^*$. In this case, we have recalled in subsection 3.2.c that a fundamental system of the differential equation (3.3) is given by J_{n_α} and Y_{n_α} . This gives us that Φ is a linear combination of Φ_+ and $\Phi_{+,-}$, where

$$(3.13) \quad \Phi_{+,-}(x) := x^{\frac{1-\alpha}{2}} Y_{n_\alpha} \left(\frac{2}{2-\alpha} \sqrt{\lambda} x^{\frac{2-\alpha}{2}} \right).$$

As we have done above, we now study if $\Phi_{+,-} \in H_\alpha^1(0, \ell)$. First we need its decomposition in series: it follows from (3.7) that

(3. 14)

$$\Phi_{+,-}(x) = \frac{2}{\pi} \Phi_+(x) \log \left(\frac{1}{2-\alpha} \sqrt{\lambda} x^{\frac{2-\alpha}{2}} \right) + \sum_{m=0}^{n_\alpha-1} \hat{a}_m x^{(1-\alpha)+(2-\alpha)m} + \sum_{m=0}^{+\infty} \hat{b}_m x^{(2-\alpha)m},$$

where

$$\hat{a}_m := -\frac{1}{\pi} \frac{(n_\alpha - m - 1)!}{m!} \left(\frac{\sqrt{\lambda}}{2\kappa_\alpha} \right)^{2m-n_\alpha}$$

and

$$\hat{b}_m := -\frac{1}{\pi} \frac{(-1)^m}{m!(n_\alpha + m)!} \left(\frac{\sqrt{\lambda}}{2\kappa_\alpha} \right)^{2m+n_\alpha} \left[\frac{\Gamma'(m+1)}{\Gamma(m+1)} + \frac{\Gamma'(m+n_\alpha+1)}{\Gamma(m+n_\alpha+1)} \right].$$

We study the three functions that appear in the formula of $\Phi_{+,-}$. First

$$\Phi_{+,-,1}(x) := \frac{2}{\pi} \Phi_+(x) \log \left(\frac{1}{2-\alpha} \sqrt{\lambda} x^{\frac{2-\alpha}{2}} \right)$$

satisfies

$$\Phi_{+,-,1}(x) \sim_0 \frac{2\kappa_\alpha}{\pi} \tilde{c}_{n_\alpha,0}^+ \log x, \quad x^{\alpha/2} \Phi'_{+,-,1}(x) \sim_0 \frac{2\kappa_\alpha}{\pi} \tilde{c}_{n_\alpha,0}^+ x^{-1+\alpha/2},$$

hence $\Phi_{+,-,1} \in H_\alpha^1(0, \ell)$ since $\alpha > 1$. Next

$$\Phi_{+,-,2}(x) := \sum_{m=0}^{n_\alpha-1} \hat{a}_m x^{(1-\alpha)+(2-\alpha)m}$$

satisfies

$$\Phi_{+,-,2}(x) \sim_0 \hat{a}_0 x^{1-\alpha}, \quad x^{\alpha/2} \Phi'_{+,-,2}(x) \sim_0 (1-\alpha) \hat{a}_0 x^{-\alpha/2},$$

hence $\Phi_{+,-,2} \notin H_\alpha^1(0, \ell)$, since $\hat{a}_0 \neq 0$. Finally,

$$\Phi_{+,-,3}(x) := \sum_{m=0}^{+\infty} \hat{b}_m x^{(2-\alpha)m}$$

satisfies

$$\Phi_{+,-,3}(x) \sim_0 \hat{b}_0, \quad x^{\alpha/2} \Phi'_{+,-,3}(x) \sim_0 (2-\alpha) \hat{b}_1 x^{1-\alpha/2},$$

hence $\Phi_{+,-,3} \in H_\alpha^1(0, \ell)$. Thus $\Phi_{+,-} = \Phi_{+,-,1} + \Phi_{+,-,2} + \Phi_{+,-,3} \notin H_\alpha^1(0, \ell)$, and if $\Phi = C_+ \Phi_+ + C_{+,-} \Phi_{+,-} \in H_\alpha^1(0, \ell)$ then necessarily $C_{+,-} = 0$, and $\Phi = C_+ \Phi_+$. Then we are in the same position as in the previous case and the conclusion is the same.

3.3.c. *Case $\nu_\alpha = 0$ (hence $\alpha = 1$).* In this case, the first sum in the decomposition of Y_0 is equal to zero, hence we have $\Phi_{+,-} = \Phi_{+,-,1} + \Phi_{+,-,3}$. Moreover,

$$\Phi_{+,-,1}(x) \sim_0 \frac{2\kappa_1}{\pi} \tilde{c}_{0,0}^+ \ln x, \quad x^{\alpha/2} \Phi'_{+,-,1}(x) \sim_0 \frac{2\kappa_1}{\pi} \tilde{c}_{0,0}^+ x^{-1/2},$$

hence $\Phi_{+,-,1} \notin H_\alpha^1(0, \ell)$. On the contrary $\Phi_{+,-,3} \in H_\alpha^1(0, \ell)$, which implies that, once again, $\Phi_{+,-} = \Phi_{+,-,1} + \Phi_{+,-,3} \notin H_\alpha^1(0, \ell)$, and the conclusion is the same. \square

3.4. Proof of Proposition 2.2.

Since $\{\Phi_{\alpha,n}, n \geq 1\}$ is an orthonormal basis of $L^2(0, \ell)$, it suffices to observe that

$$(3. 15) \quad H_{\alpha,0}^1(0, \ell) = \left\{ u \in L^2(0, \ell), \sum_{n=1}^{\infty} \lambda_{\alpha,n}(u, \Phi_{\alpha,n})_{L^2(0,\ell)}^2 < \infty \right\} (= D((-A)^{1/2})).$$

Indeed, since A generates an analytic semigroup of negative type on $X = L^2(0, \ell)$, the conclusion follows from the variation of constant formula

$$u(t) = e^{tA}u_0 + \int_0^t e^{(t-s)A}h(\cdot, s) ds$$

and well-known maximal regularity results which ensure that both maps

$$t \mapsto e^{tA}u_0 \quad \text{and} \quad t \mapsto \int_0^t e^{(t-s)A}h(\cdot, s) ds$$

belong to $H^1(0, T; X) \cap L^2(0, T; D(A)) \cap C^0([0, T]; D((-A)^{1/2}))$ whenever $u_0 \in D((-A)^{1/2})$ and $h \in L^2(0, T; X)$ (see, e.g., [6]). Finally, in order to check (3. 15), it suffices to observe that for any $u \in D(A)$, given by

$$u = \sum_{n=1}^{\infty} (u, \Phi_{\alpha,n})_{L^2(0,\ell)} \Phi_{\alpha,n},$$

we have that

$$\int_0^\ell a(x)u_x^2 dx = -(Au, u)_{L^2(0,\ell)} = \sum_{n=1}^{\infty} \lambda_{\alpha,n}(u, \Phi_{\alpha,n})_{L^2(0,\ell)}^2. \quad \square$$

4. PRELIMINARIES: THE MOMENT METHOD

We follow the strategy initiated by Fattorini and Russell [23, 24]. The precise estimates given in Theorems 2.1-2.4 are based on identities given by the moment method. We separate the boundary case from the locally distributed case.

4.1. The boundary control problem (2. 1).

4.1.a. *The moment problem satisfied by a control $H \in L^2(0, T)$.*

In this part, we analyze the problem with formal computations. First, we expand the initial condition $u_0 \in L^2(0, \ell)$: there exists $(\mu_{\alpha,n}^0)_{n \geq 1} \in \ell^2(\mathbb{N}^*)$ such that

$$u_0(x) = \sum_{n \geq 1} \mu_{\alpha,n}^0 \Phi_{\alpha,n}(x).$$

Next we expand the solution u of (2. 1):

$$u(x, t) = \sum_{n \geq 1} \beta_{\alpha,n}(t) \Phi_{\alpha,n}(x), \quad x \in (0, \ell), \quad t \geq 0$$

with

$$\sum_{n \geq 1} \beta_{\alpha,n}(t)^2 < +\infty.$$

Therefore the controllability condition $u(\cdot, T) = 0$ becomes

$$\forall n \geq 1, \quad \beta_{\alpha,n}(T) = 0.$$

On the other hand, we observe that $w_{\alpha,n}(x, t) := \Phi_{\alpha,n}(x)e^{\lambda_{\alpha,n}(t-T)}$ is solution of the adjoint problem:

$$(4. 1) \quad \begin{cases} (w_{\alpha,n})_t + (x^\alpha(w_{\alpha,n})_x)_x = 0 & x \in (0, \ell), \quad t > 0, \\ (x^\alpha(w_{\alpha,n})_x)(0, t) = 0, \\ w_{\alpha,n}(\ell, t) = 0 & t > 0. \end{cases}$$

Multiplying (2. 1) by $w_{\alpha,n}$ and (4. 1) by u , we obtain

$$\begin{aligned}
0 &= \int_0^T \int_0^\ell w_{\alpha,n}(u_t - (x^\alpha u_x)_x) + u((w_{\alpha,n})_t + (x^\alpha(w_{\alpha,n})_x)_x) \\
&= \int_0^\ell [w_{\alpha,n}u]_0^T dx - \int_0^T [w_{\alpha,n}x^\alpha u_x]_0^\ell dt + \int_0^T [ux^\alpha(w_{\alpha,n})_x]_0^\ell dt \\
&= \int_0^\ell u(x, T)\Phi_{\alpha,n}(x)dx - \int_0^\ell u(x, 0)\Phi_{\alpha,n}(x)e^{-\lambda_{\alpha,n}T}dx + \int_0^T u(\ell, t)(x^\alpha(w_{\alpha,n})_x)(\ell, t)dt \\
&= \beta_{\alpha,n}(T) - e^{-\lambda_{\alpha,n}T}\mu_{\alpha,n}^0 + \int_0^T H(t)e^{\lambda_{\alpha,n}(t-T)}(x^\alpha\Phi'_{\alpha,n})(x=\ell)dt.
\end{aligned}$$

It follows that, if the control H drives the solution to 0 at time T , then

$$r_{\alpha,n} \int_0^T H(t)e^{-\lambda_{\alpha,n}(T-t)}dt = e^{-\lambda_{\alpha,n}T}\mu_{\alpha,n}^0,$$

where we have set

$$(4. 2) \quad r_{\alpha,n} = (x^\alpha\Phi'_{\alpha,n})(x=\ell).$$

Hence, the controllability condition $u(\cdot, T) = 0$ implies that

$$(4. 3) \quad \forall n \geq 1, \quad r_{\alpha,n} \int_0^T H(t)e^{\lambda_{\alpha,n}t}dt = \mu_{\alpha,n}^0.$$

4.1.b. *The moment problem satisfied by a control $H \in H^1(0, T)$.*

Moreover, since we want a solution of the moment problem that belongs to $H^1(0, T)$, it will be more interesting to see what its derivative has to satisfy. Integrating by parts, we have

$$\int_0^T H(t)e^{\lambda_{\alpha,n}t}dt = \left[\frac{1}{\lambda_{\alpha,n}}H(t)e^{\lambda_{\alpha,n}t} \right]_0^T - \int_0^T \frac{1}{\lambda_{\alpha,n}}H'(t)e^{\lambda_{\alpha,n}t}dt.$$

Hence the derivative H' has to satisfy

$$(4. 4) \quad -\frac{r_{\alpha,n}}{\lambda_{\alpha,n}} \int_0^T H'(t)e^{\lambda_{\alpha,n}t}dt = \mu_{\alpha,n}^0 - \frac{r_{\alpha,n}}{\lambda_{\alpha,n}} \left[H(T)e^{\lambda_{\alpha,n}T} - H(0) \right].$$

We will provide a solution of this problem that satisfies $H(0) = 0 = H(T)$.

4.1.c. *A formal solution to the moment problem, using a biorthogonal family.*

Assume that there is a family $(\sigma_{\alpha,m}^+)_{m \geq 1}$ of functions $\sigma_{\alpha,m}^+ \in L^2(0, T)$, which is biorthogonal to the family $(e^{\lambda_{\alpha,n}t})_{n \geq 1}$, which means that:

$$(4. 5) \quad \forall m, n \geq 1, \quad \int_0^T \sigma_{\alpha,m}^+(t)e^{\lambda_{\alpha,n}t}dt = \delta_{mn} = \begin{cases} 1 & \text{if } m = n, \\ 0 & \text{if } m \neq n. \end{cases}$$

Then, at least formally, the function

$$H(t) := \sum_{m=1}^{\infty} \frac{\mu_{\alpha,m}^0}{r_{\alpha,m}} \sigma_{\alpha,m}^+(t)$$

satisfies the moment problem (4. 3). To enter into our functional setting, we would need to verify that this gives a function belonging to $H^1(0, T)$, then at least to $L^2(0, T)$. For this, we will need suitable bounds on $\|\sigma_{\alpha,m}^+\|_{L^2(0, T)}$, first with respect to m (to ensure the convergence of the series that defines H), then with respect to α , to measure the null controllability cost.

Since our functional setting demands the control to belong to $H^1(0, T)$, we are going to repeat the same arguments, but with the moment problem (4. 4): set

$$\lambda_{\alpha,0} := 0,$$

and assume that we are able to construct a family $(\sigma_{\alpha,m}^+)_{m \geq 0}$ of functions $\sigma_{\alpha,m}^+ \in L^2(0, T)$, which is biorthogonal to the family $(e^{\lambda_{\alpha,n}t})_{n \geq 0}$, which means that:

$$(4. 6) \quad \forall m, n \geq 0, \quad \int_0^T \sigma_{\alpha,m}^+(t) e^{\lambda_{\alpha,n}t} dt = \delta_{mn} = \begin{cases} 1 & \text{if } m = n, \\ 0 & \text{if } m \neq n. \end{cases}$$

Then consider

$$K(t) := - \sum_{m=1}^{\infty} \frac{\lambda_{\alpha,m} \mu_{\alpha,m}^0}{r_{\alpha,m}} \sigma_{\alpha,m}^+(t), \quad \text{and} \quad H(t) := \int_0^t K(\tau) d\tau.$$

Then at least formally K solves the following moment problem

$$\forall n \geq 1, \quad - \frac{r_{\alpha,n}}{\lambda_{\alpha,n}} \int_0^T K(t) e^{\lambda_{\alpha,n}t} dt = \mu_{\alpha,n}^0;$$

moreover, if $K \in L^2(0, T)$, then $H \in H^1(0, T)$, clearly $H' = K$, and $H(0) = 0$, and moreover $H(T) = 0$ thanks to the additional property that the family $(\sigma_{\alpha,m}^+)_{m \geq 1}$ is orthogonal to $e^{\lambda_{\alpha,0}t} = 1$; hence H will be in $H^1(0, T)$ and will satisfy the moment problem (4. 4). It remains to check that all this makes sense, in particular that $K \in L^2(0, T)$. Clearly, we will need suitable L^2 bounds on the biorthogonal sequence $(\sigma_{\alpha,m}^+)_{m \geq 1}$, that will come from the study of the eigenvalues $\lambda_{\alpha,n}$, and from the behavior of the real sequence $(r_{\alpha,m}^2)_m$.

4.2. The locally distributed control problem (2. 2).

4.2.a. *The moment problem satisfied by a control $h \in L^2((a, b) \times (0, T))$.*

First we expand the initial condition $u_0 \in L^2(0, \ell)$: there exists $(\mu_{\alpha,n}^0)_{n \geq 1} \in \ell^2(\mathbb{N}^*)$ such that

$$u_0(x) = \sum_{n \geq 1} \mu_{\alpha,n}^0 \Phi_{\alpha,n}(x), \quad x \in (0, \ell).$$

Next we expand the solution u of (2. 2):

$$u(x, t) = \sum_{n \geq 1} \beta_{\alpha,n}(t) \Phi_{\alpha,n}(x), \quad x \in (0, \ell), \quad t \in (0, T), \quad \text{with} \quad \sum_{n \geq 1} \beta_{\alpha,n}(t)^2 < +\infty.$$

Once again multiplying (2. 2) by $w_{\alpha,n}(x, t) := \Phi_{\alpha,n}(x) e^{\lambda_{\alpha,n}(t-T)}$, which is solution of the adjoint problem (4. 1), one gets:

$$\begin{aligned} & \int_0^T \int_0^\ell h(x, t) \chi_{[a,b]}(x) w_{\alpha,n}(x, t) dx dt = \int_0^T \int_0^\ell w_{\alpha,n}(x, t) (u_t - (x^\alpha u_x)_x) \\ & = \int_0^\ell [w_{\alpha,n} u]_0^T - \int_0^T \int_0^\ell (w_{\alpha,n})_t u - \int_0^T [w_{\alpha,n} (x^\alpha u_x)]_0^\ell + \int_0^T \int_0^\ell (w_{\alpha,n})_x x^\alpha u_x \\ & = \int_0^\ell \Phi_{\alpha,n} u(T) - e^{-\lambda_{\alpha,n}T} \int_0^\ell \Phi_{\alpha,n} u_0 - \int_0^T [w_{\alpha,n} (x^\alpha u_x)]_0^\ell \\ & + \int_0^T [x^\alpha (w_{\alpha,n})_x u]_0^\ell - \int_0^T \int_0^\ell \left((w_{\alpha,n})_t + (x^\alpha (w_{\alpha,n})_x)_x \right) u \\ & = \int_0^\ell \Phi_{\alpha,n} u(T) - e^{-\lambda_{\alpha,n}T} \int_0^\ell \Phi_{\alpha,n} u_0. \end{aligned}$$

Hence, if h drives the solution u to 0 in time T , we obtain the following moment problem:

$$(4.7) \quad \forall n \geq 1, \quad \int_0^T \int_0^\ell h(x, t) \chi_{[a, b]}(x) \Phi_{\alpha, n}(x) e^{\lambda_{\alpha, n} t} dx dt = -\mu_{\alpha, n}^0.$$

4.2.b. *A formal solution to the moment problem, using a biorthogonal family.*

Assume for a moment that there exists a family $(\sigma_{\alpha, m}^+)_{m \geq 1}$ in $L^2(0, T)$ that satisfies (4.5). Then let us define

$$(4.8) \quad h(x, t) := \sum_{m \geq 1} -\mu_{\alpha, m}^0 \sigma_{\alpha, m}^+(t) \frac{\Phi_{\alpha, m}(x)}{\int_a^b \Phi_{\alpha, m}^2}.$$

Let us prove that, formally, h is solution of the moment problem (4.7):

$$\begin{aligned} & \int_0^T \int_0^\ell h(x, t) \chi_{[a, b]}(x) \Phi_{\alpha, n}(x) e^{\lambda_{\alpha, n} t} dx dt \\ &= \int_a^b \int_0^T \left(\sum_{m \geq 1} -\mu_{\alpha, m}^0 \sigma_{\alpha, m}^+(t) \frac{\Phi_{\alpha, m}(x)}{\int_a^b \Phi_{\alpha, m}^2} \right) \Phi_{\alpha, n}(x) e^{\lambda_{\alpha, n} t} dt dx \\ &= \int_a^b \sum_{m \geq 1} -\mu_{\alpha, m}^0 \frac{\Phi_{\alpha, m}(x) \Phi_{\alpha, n}(x)}{\int_a^b \Phi_{\alpha, m}^2} \left(\int_0^T \sigma_{\alpha, m}^+(t) e^{\lambda_{\alpha, n} t} dt \right) dx \\ &= \sum_{m \geq 1} -\mu_{\alpha, m}^0 \delta_{mn} \frac{\int_a^b \Phi_{\alpha, m}(x) \Phi_{\alpha, n}(x) dx}{\int_a^b \Phi_{\alpha, m}^2} = -\mu_{\alpha, n}^0. \end{aligned}$$

Hence, formally, h defined by (4.8) solves the moment problem. It remains to check that all this makes sense, in particular that $h \in L^2((0, \ell) \times (0, T))$. Clearly, we will need suitable L^2 bounds on the biorthogonal sequence $(\sigma_{\alpha, m}^+)_{m \geq 1}$, that will come from the study of the eigenvalues $\lambda_{\alpha, n}$, and from the behavior of the real sequence $(\int_a^b \Phi_{\alpha, m}^2)_m$ (given in Proposition 2.5).

5. PROOF OF THEOREM 2.1

In this section, we are going to work on the moment problem (4.3) given by the moment method to obtain the desired lower bound on the null controllability cost for (2.1). The proof will use in particular ideas of Güichal [32].

5.1. The contribution of the eigenfunctions to the blow-up of the null controllability cost.

Assume that $H \in H^1(0, T)$ drives the solution u of (2.1) to 0 in time T . Then H satisfies (4.3). Let us compute the coefficient that appears:

$$\begin{aligned} (5.1) \quad |r_{\alpha, n}| &= |(x^\alpha \Phi'_{\alpha, n})(x = \ell)| = \ell^\alpha \frac{\sqrt{2\kappa_\alpha}}{\ell^{\kappa_\alpha} |J'_{\nu_\alpha}(j_{\nu_\alpha, n})|} \ell^{(1-\alpha)/2} j_{\nu_\alpha, n}^{\kappa_\alpha} \frac{\kappa_\alpha}{\ell} |J'_{\nu_\alpha}(j_{\nu_\alpha, n})| \\ &= \ell^{\alpha - \kappa_\alpha + (1-\alpha)/2 - 1} \sqrt{2\kappa_\alpha}^{3/2} j_{\nu_\alpha, n}^{3/2} = \sqrt{2} \ell^{(2\alpha-3)/2} \kappa_\alpha^{3/2} j_{\nu_\alpha, n} \\ &= \sqrt{2} \ell^{(\alpha-1)/2} \sqrt{\kappa_\alpha} \sqrt{\lambda_{\alpha, n}}. \end{aligned}$$

This implies that the null controllability cost blows up, at least at a rational rate: indeed, we deduce from (4.3) that

$$\forall n \geq 1, \quad \|H\|_{L^2(0, T)} \|e^{\lambda_{\alpha, n} t}\|_{L^2(0, T)} \geq \frac{|\mu_{\alpha, n}^0|}{|r_{\alpha, n}|},$$

hence

$$\forall n \geq 1, \quad \|H\|_{L^2(0,T)} \geq \frac{|\mu_{\alpha,n}^0|}{\sqrt{2} \ell^{(\alpha-1)/2} \sqrt{\kappa_\alpha} \sqrt{\lambda_{\alpha,n}}} \sqrt{\frac{2\lambda_{\alpha,n}}{e^{2\lambda_{\alpha,n}T} - 1}}.$$

Fix $u_0 = \Phi_{\alpha,1}$. Then any control that drives $\Phi_{\alpha,1}$ to 0 in time T satisfies

$$\|H\|_{L^2(0,T)} \geq \frac{1}{\ell^{(\alpha-1)/2} \sqrt{\kappa_\alpha} \sqrt{e^{2\lambda_{\alpha,1}T} - 1}}.$$

This implies a first bound from below for the null controllability cost:

$$C_{ctr-bd} \geq \frac{1}{\ell^{(\alpha-1)/2} \sqrt{\kappa_\alpha} \sqrt{e^{2\lambda_{\alpha,1}T} - 1}}.$$

In particular, just looking the behavior with respect to $\alpha \in [1, 2)$, we see that there exists $C_{T,\ell}$ independent of $\alpha \in [1, 2)$ such that

$$C_{ctr-bd} \geq \frac{C_{T,\ell}}{\sqrt{2-\alpha}}.$$

This gives a first estimate of blow-up (that we will improve in the following).

5.2. A connection between null controllability and the existence of biorthogonal sequences.

We notice the following fact: fix $m \geq 1$ and consider the initial condition $u_0 = \Phi_{\alpha,m}$; let $H_{\alpha,m}$ be a control that drives the solution of (2. 1) to 0 in time T ; then the sequence $(r_{\alpha,m} H_{\alpha,m})_{m \geq 1}$ is biorthogonal to $(e^{\lambda_{\alpha,n}t})_{n \geq 1}$ in $L^2(0, T)$. Indeed, $H_{\alpha,m}$ satisfies (4. 3):

$$\forall n \geq 1, \quad r_{\alpha,n} \int_0^T H_{\alpha,m}(t) e^{\lambda_{\alpha,n}t} dt = \mu_{\alpha,n}^0 = \delta_{mn},$$

hence

$$\forall n \geq 1, \quad \int_0^T (r_{\alpha,m} H_{\alpha,m}(t)) e^{\lambda_{\alpha,n}t} dt = r_{\alpha,m} \frac{\delta_{mn}}{r_{\alpha,n}} = \begin{cases} 1 & \text{if } m = n, \\ 0 & \text{if } m \neq n, \end{cases}$$

hence

$$\forall m, n \geq 1, \quad \int_0^T (r_{\alpha,m} H_{\alpha,m}(t)) e^{\lambda_{\alpha,n}t} dt = \delta_{mn},$$

which means that the sequence $(r_{\alpha,m} H_{\alpha,m})_{m \geq 1}$ is biorthogonal to $(e^{\lambda_{\alpha,n}t})_{n \geq 1}$ in $L^2(0, T)$.

In the literature, there exists several bounds from below for the biorthogonal families, we refer in particular to Güichal [32] and Hansen [33]. In the following we will use two extensions of the one of Güichal [32], obtained in [13] and in [14].

5.3. The concentration of the eigenvalues.

The following observation is fundamental in the understanding of the blow-up of the null controllability cost:

Lemma 5.1. *The eigenvalues concentrate when $\alpha \rightarrow 2^-$:*

$$\forall n \geq 1, \quad \lambda_{\alpha,n+1} - \lambda_{\alpha,n} \rightarrow 0 \quad \text{as } \alpha \rightarrow 2^-.$$

Before proving Lemma 5.1, let us explain why this property is clearly important in the understanding of the blow-up of the null controllability cost: as noted before, if null controllability holds, and if $H_{\alpha,m}$ is a control that drives the solution of (2. 1) with $u_0 = \Phi_{\alpha,m}$ to 0 in time T , then $(r_{\alpha,m} H_{\alpha,m})_{m \geq 1}$ is biorthogonal to

$(e^{\lambda_{\alpha,n}t})_{n \geq 1}$ in $L^2(0, T)$; now, if additionally some eigenvalues concentrate, for example $\lambda_{\alpha,2} - \lambda_{\alpha,1} \rightarrow 0$ as $\alpha \rightarrow 2^-$, then $r_{\alpha,1}H_{\alpha,1}$ will have to satisfy

$$\int_0^T (r_{\alpha,1}H_{\alpha,1}(t))e^{\lambda_{\alpha,1}t} dt = 1, \quad \text{and} \quad \int_0^T (r_{\alpha,1}H_{\alpha,1}(t))e^{\lambda_{\alpha,2}t} dt = 0,$$

hence

$$\int_0^T (r_{\alpha,1}H_{\alpha,1}(t))(e^{\lambda_{\alpha,1}t} - e^{\lambda_{\alpha,2}t}) dt = 1;$$

but this will only be possible if $\|r_{\alpha,1}H_{\alpha,1}\|$ is sufficiently large, since $\|e^{\lambda_{\alpha,1}t} - e^{\lambda_{\alpha,2}t}\|_{L^2(0,T)}$ will be small. We will come back on this later.

Proof of Lemma 5.1. We note that

$$\lambda_{\alpha,n+1} - \lambda_{\alpha,n} = \kappa_{\alpha}^2(j_{\nu_{\alpha,n+1}}^2 - j_{\nu_{\alpha,n}}^2) = \kappa_{\alpha}^2(j_{\nu_{\alpha,n+1}} - j_{\nu_{\alpha,n}})(j_{\nu_{\alpha,n+1}} + j_{\nu_{\alpha,n}}).$$

It is classical ([36] p. 135) that

- if $\nu \in [0, \frac{1}{2}]$, the sequence $(j_{\nu,n+1} - j_{\nu,n})_n$ is nondecreasing and converges to π ,
- if $\nu \geq \frac{1}{2}$, the sequence $(j_{\nu,n+1} - j_{\nu,n})_n$ is nonincreasing and converges to π .

Then, when $\nu_{\alpha} \geq \frac{1}{2}$ (i.e. when $\alpha \in [\frac{4}{3}, 2)$), the sequence $(j_{\nu_{\alpha,n+1}} - j_{\nu_{\alpha,n}})_n$ is nonincreasing, hence

$$\lambda_{\alpha,n+1} - \lambda_{\alpha,n} \leq \kappa_{\alpha}^2(j_{\nu_{\alpha,2}} - j_{\nu_{\alpha,1}})(j_{\nu_{\alpha,n+1}} + j_{\nu_{\alpha,n}}),$$

and using (3. 8),

$$\lambda_{\alpha,n+1} - \lambda_{\alpha,n} \leq \kappa_{\alpha}^2(j_{\nu_{\alpha,2}} - j_{\nu_{\alpha,1}})\left(\pi(n+1 + \frac{\nu_{\alpha}}{2} - \frac{1}{4}) + \pi(n + \frac{\nu_{\alpha}}{2} - \frac{1}{4})\right).$$

Using (3. 10), we obtain

$$\begin{aligned} j_{\nu_{\alpha,2}} - j_{\nu_{\alpha,1}} &\leq \left(\nu_{\alpha} - \frac{a_2}{2^{1/3}}\nu_{\alpha}^{1/3} + \frac{3}{20}a_2^2\frac{2^{1/3}}{\nu_{\alpha}^{1/3}}\right) - \left(\nu_{\alpha} - \frac{a_1}{2^{1/3}}\nu_{\alpha}^{1/3}\right) \\ &= \frac{a_1 - a_2}{2^{1/3}}\nu_{\alpha}^{1/3} + \frac{3}{20}a_2^2\frac{2^{1/3}}{\nu_{\alpha}^{1/3}}. \end{aligned}$$

Hence there is some C independent of $\alpha \in [\frac{4}{3}, 2)$ such that

$$(5. 2) \quad j_{\nu_{\alpha,2}} - j_{\nu_{\alpha,1}} \leq C\nu_{\alpha}^{1/3},$$

and

$$\lambda_{\alpha,n+1} - \lambda_{\alpha,n} \leq C\nu_{\alpha}^{1/3}\kappa_{\alpha}^2(n + \nu_{\alpha}) \leq C(\kappa_{\alpha}^{2/3} + \kappa_{\alpha}^{5/3}n). \quad \square$$

Remark 5.1. A similar concentration phenomenon can be pointed out in the fast control problem for the classical heat equation

$$(5. 3) \quad \begin{cases} u_t - u_{xx} = h(x, t)\chi_{[a,b]}(x) & x \in (0, 1), \quad 0 < t < T, \\ u(0, t) = 0 = u(1, t), & 0 < t < T, \\ u(x, 0) = u_0(x), & x \in (0, 1), \\ u(x, T) = 0, & x \in (0, 1). \end{cases}$$

Indeed, as is well-known, the eigenvalues of the stationary operator associated with (5. 3) are $\lambda_n = \pi^2 n^2$ for all $n > 0$. On the other hand, if we are interested in studying the behaviour of the above system for controls yielding $u(\cdot, T) = 0$ as $T \rightarrow 0^+$, then it might be useful to normalize the time, hence to look at the normalized solution

$$v(x, \tau) = u(x, \tau T).$$

This function v is solution of the problem

$$\begin{cases} v_\tau - Tv_{xx} = Th(x, \tau T)\chi_{[a,b]}(x) & x \in (0, 1), \quad 0 < \tau < 1, \\ v(0, \tau) = 0 = v(1, \tau), & 0 < \tau < 1, \\ v(x, 0) = u_0(x), & x \in (0, 1), \\ v(x, 1) = 0, & x \in (0, 1). \end{cases}$$

Clearly, the eigenvalues of the stationary operator associated with this last problem are given by the sequence $\{T\pi^2 n^2\}_{n \geq 1}$, which concentrates as $T \rightarrow 0^+$.

5.4. An additional property of the eigenvalues.

As we recalled, it is classical ([36] p. 135) that

- if $\nu \in [0, \frac{1}{2}]$, the sequence $(j_{\nu, n+1} - j_{\nu, n})_n$ is nondecreasing and converges to π ,
- if $\nu \geq \frac{1}{2}$, the sequence $(j_{\nu, n+1} - j_{\nu, n})_n$ is nonincreasing and converges to π .

Hence there exists a rank N_ν such that

$$i \geq N_\nu \implies j_{\nu, i+1} - j_{\nu, i} \leq 2\pi.$$

However, the asymptotic development (3. 10) tells us that

$$(5. 4) \quad j_{\nu, 2} - j_{\nu, 1} \sim_{\nu \rightarrow \infty} \frac{a_1 - a_2}{2^{1/3}} \nu^{1/3}.$$

Hence this rank N_ν probably satisfies $N_\nu \rightarrow +\infty$ as $\nu \rightarrow \infty$. In the following, we estimate this N_ν (using the classical theory of Sturm concerning second order differential equations); we will need this estimate later.

Lemma 5.2. *Given $\nu \geq \frac{1}{2}$, then*

$$(5. 5) \quad \forall n > \nu, \quad j_{\nu, n+1} - j_{\nu, n} \leq 2\pi.$$

Proof of Lemma 5.2. We follow and use the proofs of section 7.3 in [36]: first we note that

$$y_\nu(x) := \sqrt{x} J_\nu(x)$$

satisfies the second-order differential equation

$$y_\nu''(x) + h_\nu(x)y_\nu(x) = 0,$$

with

$$h_\nu(x) = 1 - \frac{\nu^2 - \frac{1}{4}}{x^2}.$$

Of course, y_ν and J_ν have the same positive zeros. We are going to use the following classical property of Sturm type (see Proposition 7.6 in [36]): assume that

- $f, g : [a, b] \rightarrow \mathbb{R}$ are continuous and satisfy

$$\forall x \in [a, b], \quad f(x) < g(x),$$

- u, v are functions of class C^2 satisfying

$$\forall x \in [a, b], \quad u'' + fu = 0, \quad v'' + gv = 0,$$

- a, b are two consecutive zeros of u ,

then v has at least one zero in (a, b) .

We recall that, in a classical way ([36]), this implies that J_ν has an infinite number of positive zeros: indeed:

$$\forall x > \nu, \quad h_\nu(x) > \frac{1}{4\nu^2},$$

hence choosing

$$\begin{aligned} k &\geq 1, \quad a := 2k\nu\pi, \quad b := 2(k+1)\nu\pi, \\ f(x) &:= \frac{1}{4\nu^2}, \quad u(x) := \sin \frac{x}{2\nu}, \\ g(x) &:= h_\nu(x), \quad v(x) := y_\nu(x), \end{aligned}$$

we can apply the Sturm property, and we derive that y_ν (hence J_ν) has at least one zero on $(2k\nu\pi, 2(k+1)\nu\pi)$. From (3. 8) we also have

$$\forall k > \nu, \quad j_{\nu,k} > \pi\left(\nu + \frac{1}{4}\left(\nu - \frac{1}{2}\right)\right) =: \gamma_\nu,$$

and then we can apply the Sturm property with

$$\begin{aligned} k > \nu, \quad a &:= j_{\nu,k}, \quad b := j_{\nu,k} + \frac{\pi}{\sqrt{h_\nu(\gamma_\nu)}}, \\ f(x) &:= h_\nu(\gamma_\nu), \quad u(x) := \sin\left(\sqrt{h_\nu(\gamma_\nu)}(x - j_{\nu,k})\right), \\ g(x) &:= h_\nu(x), \quad v(x) := y_\nu(x), \end{aligned}$$

and we deduce that y_ν has at least one zero inside $(j_{\nu,k}, j_{\nu,k} + \frac{\pi}{\sqrt{h_\nu(\gamma_\nu)}})$, hence

$$j_{\nu,k+1} < j_{\nu,k} + \frac{\pi}{\sqrt{h_\nu(\gamma_\nu)}}.$$

Hence

$$\forall k > \nu, \quad j_{\nu,k+1} - j_{\nu,k} < \frac{\pi}{\sqrt{h_\nu(\gamma_\nu)}} = \frac{\pi}{\sqrt{1 - \frac{\nu^2 - \frac{1}{4}}{\gamma_\nu^2}}}.$$

It can be easily checked that

$$\forall \nu \geq \frac{1}{2}, \quad \frac{\pi}{\sqrt{1 - \frac{\nu^2 - \frac{1}{4}}{\gamma_\nu^2}}} \leq 2\pi :$$

indeed, if $\nu \geq \frac{1}{2}$, then

$$\left(1 - \frac{\nu^2 - \frac{1}{4}}{\gamma_\nu^2}\right) - \frac{1}{4} = \frac{3}{4} - \frac{\nu^2 - \frac{1}{4}}{\gamma_\nu^2} = \frac{3\gamma_\nu^2 - 4(\nu^2 - \frac{1}{4})}{4\gamma_\nu^2},$$

and

$$3\gamma_\nu^2 - 4(\nu^2 - \frac{1}{4}) = 3\left(\pi\left(\nu + \frac{1}{4}\left(\nu - \frac{1}{2}\right)\right)\right)^2 - 4\left(\nu^2 - \frac{1}{4}\right)$$

and the discriminant of this quantity is negative, hence the quantity remains positive. This implies (5. 5). \square

5.5. A lower bound of the norm of any sequence biorthogonal to $(e^{\lambda_{\alpha,n}t})_n$ when $\nu_\alpha \in [0, \frac{1}{2}]$.

If $\nu_\alpha \in [0, \frac{1}{2}]$, the gap $(j_{\nu_\alpha, n+1} - j_{\nu_\alpha, n})_n$ is nondecreasing and converges to π , hence

$$\forall n \geq 1, \quad j_{\nu_\alpha, n+1} - j_{\nu_\alpha, n} \leq \pi,$$

hence

$$\forall n \geq 1, \quad \sqrt{\lambda_{\alpha, n+1}} - \sqrt{\lambda_{\alpha, n}} = \ell^{\frac{\alpha}{2}-1} \kappa_\alpha (j_{\nu_\alpha, n+1} - j_{\nu_\alpha, n}) \leq \ell^{\frac{\alpha}{2}-1} \kappa_\alpha \pi,$$

hence

$$\forall n \geq 1, \quad \sqrt{\lambda_{\alpha, n+1}} - \sqrt{\lambda_{\alpha, n}} \leq \gamma_{max} \quad \text{with } \gamma_{max} = \ell^{\frac{\alpha}{2}-1} \kappa_\alpha \pi.$$

Let us apply the following extension of Güichal [32]:

Theorem 5.1. (Theorem 2.5 in [13]) Assume that

$$\forall n \geq 0, \quad \lambda_n \geq 0,$$

and that there is some $0 < \gamma_{\min} \leq \gamma_{\max}$ such that

$$(5.6) \quad \forall n \geq 0, \quad \gamma_{\min} \leq \sqrt{\lambda_{n+1}} - \sqrt{\lambda_n} \leq \gamma_{\max}.$$

Then there exists $c_u > 0$ independent of T , and m such that: any family $(\sigma_m^+)_{m \geq 0}$ which is biorthogonal to the family $(e^{\lambda_n t})_{n \geq 0}$ in $L^2(0, T)$ satisfies:

$$(5.7) \quad \|\sigma_m^+\|_{L^2(0, T)}^2 \geq e^{-2\lambda_m T} e^{\frac{1}{2\gamma_{\max}^2 T}} b(T, \gamma_{\max}, m),$$

with

$$(5.8) \quad b(T, \gamma_{\max}, m) = \frac{c_u^2}{C(m, \gamma_{\max}, \lambda_0)^2 T} \left(\frac{1}{2\gamma_{\max}^2 T}\right)^{2m} \frac{1}{(4\gamma_{\max}^2 T + 1)^2}.$$

and

$$(5.9) \quad C(m, \gamma_{\max}, \lambda_0) = m! 2^{m + [\frac{2\sqrt{\lambda_0}}{\gamma_{\max}}] + 1} \left(m + \left[\frac{2\sqrt{\lambda_0}}{\gamma_{\max}}\right] + 1\right).$$

Using Theorem 5.1 with $\gamma_{\max} = \ell^{\frac{\alpha}{2}-1} \kappa_\alpha \pi$, one obtains that any family $(\sigma_m^+)_{m \geq 1}$ which is biorthogonal to the family $(e^{\lambda_{\alpha, n} t})_{n \geq 1}$ in $L^2(0, T)$ satisfies a lower bound with the classical dominant exponential factor of the type $e^{C/T}$:

$$\|\sigma_m^+\|_{L^2(0, T)}^2 \geq e^{-2\lambda_{\alpha, m} T} e^{C_u \frac{\ell^{2-\alpha}}{T \kappa_\alpha^2}} \frac{1}{T} \left(\frac{\ell^{2-\alpha}}{T \kappa_\alpha^2}\right)^{2m} \left(\frac{\ell^{2-\alpha}}{1 + \frac{\ell^{2-\alpha}}{T \kappa_\alpha^2}}\right)^2 \frac{C_u^{2m}}{m!^2} 2^{-2\frac{\ell^{1-\alpha/2}}{\kappa_\alpha}} \frac{1}{(m+1)^2 + \frac{\lambda_{\alpha, 1} \ell^{1-\frac{\alpha}{2}}}{\kappa_\alpha^2 \pi^2}}.$$

This will immediately give an exponential blow-up of the cost as $T \rightarrow 0^+$, as explained in subsection 5.7, but the interesting behavior is when $\alpha \rightarrow 2^-$, and we study it in the following.

5.6. A lower bound of the norm of any sequence biorthogonal to $(e^{\lambda_{\alpha, n} t})_n$ when $\nu_\alpha \geq \frac{1}{2}$.

This is the interesting case, where $\alpha \rightarrow 2^-$. In this case, the gap $(j_{\nu_\alpha, n+1} - j_{\nu_\alpha, n})_n$ is nonincreasing and converges to π , hence

$$\forall n \geq 1, \quad \pi \leq j_{\nu_\alpha, n+1} - j_{\nu_\alpha, n} \leq j_{\nu_\alpha, 2} - j_{\nu_\alpha, 1},$$

hence

$$\forall n \geq 1, \quad \sqrt{\lambda_{\alpha, n+1}} - \sqrt{\lambda_{\alpha, n}} \leq \gamma_{\max} \quad \text{with } \gamma_{\max} = \ell^{\frac{\alpha}{2}-1} \kappa_\alpha (j_{\nu_\alpha, 2} - j_{\nu_\alpha, 1});$$

but this time, we already noted that $j_{\nu_\alpha, 2} - j_{\nu_\alpha, 1}$ behaves as $\nu_\alpha^{1/3}$ (see (5.4)), hence

$$\gamma_{\max} = c_\alpha \ell^{\frac{\alpha}{2}-1} \kappa_\alpha^{2/3},$$

with some uniformly bounded c_α .

On the other hand, we proved in Lemma 5.2 that

$$\forall n \geq \nu_\alpha + 1, \quad j_{\nu_\alpha, n+1} - j_{\nu_\alpha, n} \leq 2\pi,$$

hence

$$\forall n \geq N_*, \quad \sqrt{\lambda_{\alpha, n+1}} - \sqrt{\lambda_{\alpha, n}} \leq \gamma_{\max}^*, \quad \text{with } N_* = [\nu_\alpha] + 1, \quad \text{and } \gamma_{\max}^* = 2\pi \ell^{\frac{\alpha}{2}-1} \kappa_\alpha.$$

Note that

$$\frac{\gamma_{\max}}{\gamma_{\max}^*} = \frac{c_\alpha}{2\pi \kappa_\alpha^{1/3}} \rightarrow \infty \quad \text{as } \alpha \rightarrow 2^-.$$

In that context, when there is a 'bad' global gap γ_{\max} , and a 'good' (much smaller) asymptotic gap γ_{\max}^* , it is interesting to use the following extension of Theorem 5.1:

Theorem 5.2. (Theorem 2.2 in [14]) Assume that

$$\forall n \geq 1, \quad \lambda_n \geq 0,$$

and that there are $0 < \gamma_{min} \leq \gamma_{max}^* \leq \gamma_{max}$ such that

$$(5.10) \quad \forall n \geq 1, \quad \gamma_{min} \leq \sqrt{\lambda_{n+1}} - \sqrt{\lambda_n} \leq \gamma_{max},$$

and

$$(5.11) \quad \forall n \geq N_*, \quad \sqrt{\lambda_{n+1}} - \sqrt{\lambda_n} \leq \gamma_{max}^*.$$

Then any family $(\sigma_m^+)_{m \geq 1}$ which is biorthogonal to the family $(e^{\lambda_n t})_{n \geq 1}$ in $L^2(0, T)$ satisfies:

$$(5.12) \quad \|\sigma_m^+\|_{L^2(0, T)}^2 \geq e^{-2\lambda_m T} e^{\frac{2}{T(\gamma_{max}^*)^2}} b^*(T, \gamma_{max}, \gamma_{max}^*, N_*, \lambda_1, m)^2,$$

where b^* is rational in T (and explicitly given in Lemma 4.4 of [14]).

Applying Theorem 5.2, we obtain that any family $(\sigma_m^+)_{m \geq 1}$ which is biorthogonal to the family $(e^{\lambda_{\alpha, n} t})_{n \geq 1}$ in $L^2(0, T)$ satisfies

$$\|\sigma_m^+\|_{L^2(0, T)}^2 \geq e^{-2\lambda_{\alpha, m} T} e^{\frac{2}{4\pi^2 \kappa_\alpha^2 T \ell^{\alpha-2}}} b^*(T, \gamma_{max}, \gamma_{max}^*, N_*, \lambda_{\alpha, 1}, m)^2$$

with an explicit value of b^* (see Lemma 4.4 in [14]): when $m \leq N_*$, we have

$$(5.13) \quad b^*(T, \gamma_{max}, \gamma_{max}^*, N_*, \lambda_1, m) = C^* \frac{\sqrt{1+T\lambda_1}}{\sqrt{T}} \frac{(T(\gamma_{max}^*)^2)^{K_*+K'_*+2}}{(1+(T(\gamma_{max}^*)^2))^{N_*+K_*+K'_*+3}},$$

where

$$K_* = \left[\frac{2\sqrt{\lambda_1} + (N_* + m)\gamma_{max}}{\gamma_{max}^*} \right] - N_* + 2,$$

$$K'_* = \left[\frac{\gamma_{max}}{\gamma_{max}^*} (N_* - m) \right] - N_* + 2,$$

$$C^* = \frac{1}{(N_* + K_* + K'_* + 3)!} \frac{c_u (\gamma_{max}^*)^{2(N_*-1)}}{C^{(+)} C^{(-)},}$$

where

$$C^{(+)} = \left(\frac{\gamma_{max}}{\gamma_{max}^*} \right)^{N_*-1} \frac{(N_* + m + \left[\frac{2\sqrt{\lambda_1}}{\gamma_{max}^*} \right] + 1)!}{(m + \left[\frac{2\sqrt{\lambda_1}}{\gamma_{max}^*} \right] + 1)! \left(\left[\frac{2\sqrt{\lambda_1} + (N_* + m)\gamma_{max}}{\gamma_{max}^*} \right] + 1 \right)! (2m + \left[\frac{2\sqrt{\lambda_1}}{\gamma_{max}^*} \right] + 1)!},$$

and

$$C^{(-)} = \left(\frac{\gamma_{max}}{\gamma_{max}^*} \right)^{N_*-1} \frac{(m-1)! (N_* - m)!}{(1 + \left[\frac{\gamma_{max}}{\gamma_{max}^*} (N_* - m) \right])!}.$$

These expressions seem to be a little frightening, but we are looking for the behavior as $\alpha \rightarrow 2^-$, and this is not difficult to study: one immediately sees that

$$K_* + K'_* = c\nu_\alpha^{4/3} + c'\ell^{1-\alpha/2}\nu_\alpha + c''\nu_\alpha^{1/3}m,$$

$$(\gamma_{max}^*)^{2(N_*-1)} = e^{-c\nu_\alpha \ln \nu_\alpha - c'\nu_\alpha \ln \ell},$$

$$\frac{1}{(N_* + K_* + K'_* + 3)!} \geq e^{-C(\nu_\alpha^{4/3} + \ell^{1-\alpha/2}\nu_\alpha + \nu_\alpha^{1/3}m)(\ln \nu_\alpha + (1-\frac{\alpha}{2}) \ln \ell + \ln m)},$$

and finally

$$\frac{1}{C^{(+)} C^{(-)}} \geq e^{-c(\nu_\alpha + m)(\ln \nu_\alpha + \ln m)} \frac{1}{(m-1)!},$$

hence we obtain that

$$C^* \geq e^{-C(\nu_\alpha^{4/3} + \ell^{1-\alpha/2}\nu_\alpha + \nu_\alpha^{1/3}m)(\ln \nu_\alpha + (1-\frac{\alpha}{2}) \ln \ell + \ln m)} \frac{1}{(m-1)!}.$$

This gives that

$$b^* \geq e^{-C(\nu_\alpha^{4/3} + \ell^{1-\alpha/2} \nu_\alpha + \nu_\alpha^{1/3} m)(\ln \nu_\alpha + (1-\frac{\alpha}{2}) \ln \ell + \ln m + \ln \frac{1}{T})} \frac{1}{(m-1)!} \frac{\sqrt{1+T}}{\sqrt{T}},$$

hence

$$\|\sigma_m^+\|_{L^2(0,T)}^2 \geq \underline{b}(T, \alpha, m)^2,$$

with

$$(5.14) \quad \underline{b}(T, \alpha, m) := e^{-\lambda_{\alpha,m} T} e^{C_u \frac{\ell^{2-\alpha}}{T \kappa_\alpha^2}} \frac{1}{(m-1)!} \frac{\sqrt{1+T}}{\sqrt{T}} e^{-C(\nu_\alpha^{4/3} + \ell^{1-\alpha/2} \nu_\alpha + \nu_\alpha^{1/3} m)(\ln \nu_\alpha + (1-\frac{\alpha}{2}) \ln \ell + \ln m + \ln \frac{1}{T})}.$$

This will give the expected blow-up of the cost, as $\alpha \rightarrow 2^-$ and/or as $T \rightarrow 0^+$.

5.7. The exponential blow-up of the cost.

In the previous subsection, we obtained a bound from below for any biorthogonal sequence. But we already noted that if $u_0 = \phi_{\alpha,m}$, and if $H_{\alpha,m}$ is any control that drives u_0 to rest in time T , then $(r_{\alpha,m} H_{\alpha,m})_{m \geq 1}$ is biorthogonal to $(e^{\lambda_{\alpha,n} t})_{n \geq 1}$ in $L^2(0, T)$. Hence

$$\|r_{\alpha,m} H_{\alpha,m}\|_{L^2(0,T)} \geq \underline{b}(T, \alpha, m),$$

where $\underline{b}(T, \alpha, m)$ is given in (5.14). By definition of the cost, we obtain that

$$\forall m \geq 1, \quad C_{ctr-bd} \geq \frac{1}{|r_{\alpha,m}|} \underline{b}(T, \alpha, m),$$

which gives the expected exponential blow-up of the cost: choosing $m = 1$, and using the fact that $|r_{\alpha,1}| = \sqrt{2\kappa_\alpha \lambda_{\alpha,1}} \ell^{(\alpha-1)/2}$ and that $\lambda_{\alpha,1} = \kappa_\alpha j_{\nu_\alpha,1} \ell^{(\alpha-2)/2} \geq \frac{\pi}{4} \kappa_\alpha \nu_\alpha \ell^{(\alpha-2)/2}$, we obtain that there exists some C_u independent of all the other parameters such that

$$\begin{aligned} C_{ctr-bd} &\geq C_u \frac{\ell^{2-\alpha}}{\sqrt{\kappa_\alpha T \ell}} e^{-\lambda_{\alpha,1} T} e^{C_u \frac{\ell^{2-\alpha}}{T \kappa_\alpha^2}} e^{-C(\nu_\alpha^{4/3} + \ell^{1-\alpha/2} \nu_\alpha)(\ln \nu_\alpha + \ln \ell^{1-\alpha/2} + \ln \frac{1}{T})} \\ &\geq C_u \frac{\ell^{2-\alpha}}{\sqrt{\kappa_\alpha T \ell}} e^{-\pi^2 \frac{T}{\ell^{2-\alpha}}} e^{C_u \frac{\ell^{2-\alpha}}{T \kappa_\alpha^2}} e^{-C(\nu_\alpha^{4/3} + \ell^{1-\alpha/2} \nu_\alpha)(\ln \nu_\alpha + \ln \ell^{1-\alpha/2} + \ln \frac{1}{T})}, \end{aligned}$$

and this is (2.1) and concludes the proof of Theorem 2.1. \square

6. PROOF OF THEOREM 2.2

We will use the following result (Theorem 2.4 in [13]):

Theorem 6.1. (*Existence of a suitable biorthogonal family and upper bounds*) Assume that

$$\forall n \geq 0, \quad \lambda_n \geq 0,$$

and that there is some $\gamma_{min} > 0$ such that

$$(6.1) \quad \forall n \geq 0, \quad \sqrt{\lambda_{n+1}} - \sqrt{\lambda_n} \geq \gamma_{min}.$$

Then there exists a family $(\sigma_m^+)_{m \geq 0}$ which is biorthogonal to the family $(e^{\lambda_n t})_{n \geq 0}$ in $L^2(0, T)$:

$$(6.2) \quad \forall m, n \geq 0, \quad \int_0^T \sigma_m^+(t) e^{\lambda_n t} dt = \delta_{mn}.$$

Moreover, it satisfies: there is some universal constant C_u independent of T , γ_{min} and m such that, for all $m \geq 0$, we have

$$(6.3) \quad \|\sigma_m^+\|_{L^2(0,T)}^2 \leq C_u e^{-2\lambda_m T} e^{C_u \frac{\sqrt{\lambda_m}}{\gamma_{min}}} B(T, \gamma_{min}),$$

with

$$(6.4) \quad B(T, \gamma_{min}) = \begin{cases} \left(\frac{1}{T} + \frac{1}{T^2 \gamma_{min}^2}\right) e^{\frac{C_u}{\gamma_{min}^2 T}} & \text{if } T \leq \frac{1}{\gamma_{min}^2}, \\ C_u \gamma_{min}^2 & \text{if } T \geq \frac{1}{\gamma_{min}^2}. \end{cases}$$

Note that (6.3) and (6.4) imply that there is some universal constant C_u independent of T , γ_{min} and m such that, for all $m \geq 0$, we have

$$(6.5) \quad \|\sigma_m^+\|_{L^2(0,T)}^2 \leq C_u e^{-2\lambda_m T} e^{C_u \frac{\sqrt{\lambda_m}}{\gamma_{min}}} e^{\frac{C_u}{\gamma_{min}^2 T}} B^*(T, \gamma_{min}),$$

with

$$(6.6) \quad B^*(T, \gamma_{min}) = \frac{C_u}{T} \max\{T \gamma_{min}^2, \frac{1}{T \gamma_{min}^2}\}.$$

Now, as we have already noted, the eigenvalues of the problem satisfy

$$\forall n \geq 1, \quad \sqrt{\lambda_{\alpha, n+1}} - \sqrt{\lambda_{\alpha, n}} = \ell^{\frac{\alpha}{2}-1} \kappa_{\alpha} (j_{\nu_{\alpha, n+1}} - j_{\nu_{\alpha, n}}) \geq \begin{cases} \ell^{\frac{\alpha}{2}-1} \kappa_{\alpha} (j_{\nu_{\alpha, 2}} - j_{\nu_{\alpha, 1}}) & \text{if } \nu_{\alpha} \in [0, \frac{1}{2}], \\ \ell^{\frac{\alpha}{2}-1} \kappa_{\alpha} \pi & \text{if } \nu_{\alpha} \geq \frac{1}{2} \end{cases}.$$

Define artificially

$$\lambda_{\alpha, 0} := 0.$$

Then

$$\sqrt{\lambda_{\alpha, 1}} - \sqrt{\lambda_{\alpha, 0}} = \ell^{\frac{\alpha}{2}-1} \kappa_{\alpha} j_{\nu_{\alpha, 1}}.$$

Then consider

$$c_{\alpha} := \begin{cases} \min\{j_{\nu_{\alpha, 2}} - j_{\nu_{\alpha, 1}}, j_{\nu_{\alpha, 1}}\} & \text{if } \nu_{\alpha} \in [0, \frac{1}{2}], \\ \min\{\pi, j_{\nu_{\alpha, 1}}\} & \text{if } \nu_{\alpha} \geq \frac{1}{2} \end{cases},$$

and

$$\underline{c} := \inf_{\alpha \in [0, 2)} c_{\alpha}.$$

It is clear from (3.8)-(3.9) that $\underline{c} > 0$, and by construction we have

$$\forall n \geq 0, \quad \sqrt{\lambda_{\alpha, n+1}} - \sqrt{\lambda_{\alpha, n}} \geq \gamma_{min} \quad \text{with} \quad \gamma_{min} := \ell^{\frac{\alpha}{2}-1} \kappa_{\alpha} \underline{c}.$$

Then, applying Theorem 6.1 with $\gamma_{min} = \ell^{\frac{\alpha}{2}-1} \kappa_{\alpha} \underline{c}$, we obtain that there exists a family $(\sigma_{\alpha, m}^+)_{m \geq 0}$ biorthogonal to $(e^{\lambda_{\alpha, n} t})_{n \geq 0}$ in $L^2(0, T)$, and such that

$$\begin{aligned} \|\sigma_{\alpha, m}^+\|_{L^2(0,T)}^2 &\leq C_u e^{-2\lambda_{\alpha, m} T} e^{C_u \frac{\sqrt{\lambda_{\alpha, m}}}{\gamma_{min}}} B(T, \gamma_{min}) \\ &= C_u e^{-2\lambda_{\alpha, m} T} e^{C_u j_{\nu_{\alpha, m}}} B(T, \gamma_{min}). \end{aligned}$$

Then define

$$(6.7) \quad K(t) := - \sum_{m=1}^{\infty} \frac{\lambda_{\alpha, m} \mu_{\alpha, m}^0}{r_{\alpha, m}} \sigma_{\alpha, m}^+(t), \quad \text{and} \quad H(t) := \int_0^t K(\tau) d\tau,$$

and let us check that H is an admissible control that drives the solution of (2.1) to 0 in time T :

- first we check that $K \in L^2(0, T)$: using (5.1) and (6.3), we have

$$\sum_{m=1}^{\infty} \frac{|\lambda_{\alpha, m} \mu_{\alpha, m}^0|}{|r_{\alpha, m}|} \|\sigma_{\alpha, m}^+\|_{L^2(0,T)} \leq \left(\sum_{m=1}^{\infty} |\mu_{\alpha, m}^0|^2 \right)^{1/2} \left(\sum_{m=1}^{\infty} \frac{|\lambda_{\alpha, m}|^2}{|r_{\alpha, m}|^2} \|\sigma_{\alpha, m}^+\|_{L^2(0,T)}^2 \right)^{1/2}$$

which is finite (we will come back on this in the following); this implies that $H \in H^1(0, T)$, and of course $H(0) = 0$, and also $H(T) = 0$ using (6.2) with $n = 0$;

- next, we check that H satisfies the moment problem (4. 4):

$$\begin{aligned} \forall n \geq 1, \quad & -\frac{r_{\alpha,n}}{\lambda_{\alpha,n}} \int_0^T H'(t) e^{\lambda_{\alpha,n} t} dt + \frac{r_{\alpha,n}}{\lambda_{\alpha,n}} \left[H(T) e^{\lambda_{\alpha,n} T} - H(0) \right] \\ & = -\frac{r_{\alpha,n}}{\lambda_{\alpha,n}} \int_0^T K(t) e^{\lambda_{\alpha,n} t} dt = \mu_{\alpha,n}^0; \end{aligned}$$

- finally we check that the solution of (2. 1) satisfies $u(T) = 0$: multiplying the first equation of (2. 1) by $w_{\alpha,n}(x, t) := \Phi_{\alpha,n}(x) e^{\lambda_{\alpha,n}(t-T)}$ and integrating by parts, we obtain that

$$\forall n \geq 1, \quad \int_0^\ell u(x, T) \Phi_{\alpha,n}(x) dx = 0,$$

hence $u(T) = 0$.

Hence H is an admissible control, and therefore

$$C_{ctr-bd} \leq \frac{\|H\|_{H^1(0,T)}}{\|u_0\|_{L^2(0,\ell)}} \leq C \frac{\|K\|_{L^2(0,T)}}{\|u_0\|_{L^2(0,\ell)}},$$

hence

$$C_{ctr-bd} \leq C \left(\sum_{m=1}^{\infty} \frac{|\lambda_{\alpha,m}|^2}{|r_{\alpha,m}|^2} \|\sigma_{\alpha,m}^+\|_{L^2(0,T)}^2 \right)^{1/2}.$$

Since $r_{\alpha,m}^2 = 2\kappa_\alpha \ell^{\alpha-1} \lambda_{\alpha,m}$, we have

$$\frac{|\lambda_{\alpha,m}|^2}{|r_{\alpha,m}|^2} = \frac{\kappa_\alpha j_{\nu_\alpha,m}^2}{2\ell},$$

hence

$$\begin{aligned} C_{ctr-bd} & \leq \frac{C}{\sqrt{\ell}} \sqrt{\kappa_\alpha} \left(\sum_{m=1}^{\infty} j_{\nu_\alpha,m}^2 \|\sigma_{\alpha,m}^+\|_{L^2(0,T)}^2 \right)^{1/2} \\ & \leq \frac{C}{\sqrt{\ell}} \sqrt{\kappa_\alpha} \left(\sum_{m=1}^{\infty} j_{\nu_\alpha,m}^2 e^{-2\lambda_{\alpha,m} T} e^{C_u \frac{\sqrt{\lambda_{\alpha,m}}}{\gamma_{min}}} B(T, \gamma_{min}) \right)^{1/2} \\ & = \frac{C}{\sqrt{\ell}} \sqrt{\kappa_\alpha} \sqrt{B(T, \gamma_{min})} \left(\sum_{m=1}^{\infty} j_{\nu_\alpha,m}^2 e^{-2\lambda_{\alpha,m} T} e^{C_u \frac{\sqrt{\lambda_{\alpha,m}}}{\gamma_{min}}} \right)^{1/2} \end{aligned}$$

But

$$C_u \frac{\sqrt{\lambda_{\alpha,m}}}{\gamma_{min}} \leq \lambda_{\alpha,m} T + \frac{C_u^2}{T \gamma_{min}^2},$$

hence

$$\begin{aligned} (6. 8) \quad C_{ctr-bd} & \leq \frac{C}{\sqrt{\ell}} \sqrt{\kappa_\alpha} \sqrt{B(T, \gamma_{min})} e^{\frac{C_u^2}{2T \gamma_{min}^2}} \left(\sum_{m=1}^{\infty} j_{\nu_\alpha,m}^2 e^{-\lambda_{\alpha,m} T} \right)^{1/2} \\ & = \frac{C}{\sqrt{\ell}} \sqrt{\kappa_\alpha} \sqrt{B(T, \gamma_{min})} e^{\frac{C_u^2}{2\ell^2} \frac{\ell^2 - \alpha}{T \kappa_\alpha}} \left(\sum_{m=1}^{\infty} j_{\nu_\alpha,m}^2 e^{-j_{\nu_\alpha,m}^2 \frac{\kappa_\alpha^2 T}{\ell^2 - \alpha}} \right)^{1/2} \end{aligned}$$

It remains to estimate the last sum. We distinguish the cases $\nu_\alpha \leq \frac{1}{2}$ and $\nu_\alpha \geq \frac{1}{2}$.

Take $Y > 0$. When $\nu_\alpha \leq \frac{1}{2}$, using (3. 9) we see that

$$j_{\nu_\alpha,m}^2 e^{-j_{\nu_\alpha,m}^2 Y} \leq \pi^2 \left(m + \frac{\nu_\alpha - \frac{1}{2}}{4} \right)^2 e^{-Y \pi^2 \left(m + \frac{\nu_\alpha - \frac{1}{2}}{2} \right)^2}.$$

When $m \geq 1$, we have

$$\left(m + \frac{\nu_\alpha - \frac{1}{2}}{2} \right)^2 \geq \left(m - \frac{1}{4} \right)^2 \geq \frac{1}{2} m^2 + \frac{1}{16},$$

hence

$$j_{\nu_\alpha, m}^2 e^{-j_{\nu_\alpha, m}^2 Y} \leq \pi^2 m^2 e^{-Y\pi^2 m^2/2} e^{-Y\pi^2/16}.$$

The function $V : x \mapsto \pi^2 x^2 e^{-Y\pi^2 x^2/2}$ attains its maximum at $x_Y := \sqrt{\frac{2}{\pi^2 Y}}$, is increasing on $[0, x_Y]$, decreasing on $[x_Y, +\infty)$, and its maximum is $\frac{2}{eY}$. If $x_Y \leq 1$, then

$$\begin{aligned} \sum_{m=1}^{\infty} \pi^2 m^2 e^{-Y\pi^2 m^2/2} &= \sum_{m=1}^{\infty} V(m) = V(1) + \sum_{m=1}^{\infty} V(m+1) \\ &\leq V(1) + \sum_{m=1}^{\infty} \int_m^{m+1} V(x) dx = V(1) + \int_1^{\infty} V(x) dx. \end{aligned}$$

If $x_Y \geq 1$, then

$$\begin{aligned} \sum_{m=1}^{\infty} \pi^2 m^2 e^{-Y\pi^2 m^2/2} &= \sum_{m=0}^{\infty} V(m) \\ &= \sum_{m=0}^{[x_Y]-1} V(m) + V([x_Y]) + V([x_Y] + 1) + \sum_{m=[x_Y]+2}^{\infty} V(m) \\ &\leq \sum_{m=0}^{[x_Y]-1} \int_m^{m+1} V(x) dx + V([x_Y]) + V([x_Y] + 1) + \sum_{m=[x_Y]+2}^{\infty} \int_{m-1}^m V(x) dx \\ &\leq V([x_Y]) + V([x_Y] + 1) + \int_0^{\infty} V(x) dx \leq \frac{4}{eY} + \int_0^{\infty} V(x) dx. \end{aligned}$$

Hence in any case,

$$\begin{aligned} (6.9) \quad \sum_{m=1}^{\infty} \pi^2 m^2 e^{-Y\pi^2 m^2/2} &\leq \frac{4}{eY} + \int_0^{\infty} \pi^2 x^2 e^{-Y\pi^2 x^2/2} dx \\ &= \frac{4}{eY} + \frac{2^{3/2}}{\pi Y^{3/2}} \int_0^{\infty} s^2 e^{-s^2} ds, \end{aligned}$$

and

$$\sum_{m=1}^{\infty} j_{\nu_\alpha, m}^2 e^{-j_{\nu_\alpha, m}^2 Y} \leq \left(\frac{4}{eY} + \frac{2^{3/2}}{\pi Y^{3/2}} \int_0^{\infty} s^2 e^{-s^2} ds \right) e^{-Y\pi^2/16}.$$

hence there exists some C_u such that, when $\nu_\alpha \leq \frac{1}{2}$,

$$(6.10) \quad \forall Y > 0, \quad \sum_{m=1}^{\infty} j_{\nu_\alpha, m}^2 e^{-j_{\nu_\alpha, m}^2 Y} \leq \frac{1}{C_u Y^{3/2}} e^{-C_u Y}.$$

When $\nu_\alpha \geq \frac{1}{2}$, we proceed in the same way, using (3.8): we see that

$$j_{\nu_\alpha, m}^2 e^{-j_{\nu_\alpha, m}^2 Y} \leq \pi^2 \left(m + \frac{\nu_\alpha - \frac{1}{2}}{2} \right)^2 e^{-Y\pi^2 \left(m + \frac{\nu_\alpha - \frac{1}{2}}{4} \right)^2}.$$

But

$$\left(m + \frac{\nu_\alpha - \frac{1}{2}}{4} \right)^2 \geq \frac{1}{2} m^2 + \left(\frac{\nu_\alpha - \frac{1}{2}}{4} \right)^2,$$

hence

$$\begin{aligned} j_{\nu_\alpha, m}^2 e^{-j_{\nu_\alpha, m}^2 Y} &\leq \pi^2 \left(m + \frac{\nu_\alpha - \frac{1}{2}}{2}\right)^2 e^{-Y\pi^2 m^2/2} e^{-Y\pi^2 \left(\frac{\nu_\alpha - \frac{1}{2}}{4}\right)^2} \\ &\leq 2\pi^2 \left(m^2 + \left(\frac{\nu_\alpha - \frac{1}{2}}{2}\right)^2\right) e^{-Y\pi^2 m^2/2} e^{-Y\pi^2 \left(\frac{\nu_\alpha - \frac{1}{2}}{4}\right)^2} \\ &\leq 2 \left(1 + \left(\frac{\nu_\alpha - \frac{1}{2}}{2}\right)^2\right) e^{-Y\pi^2 \left(\frac{\nu_\alpha - \frac{1}{2}}{4}\right)^2} \pi^2 m^2 e^{-Y\pi^2 m^2/2}. \end{aligned}$$

Hence, using (6. 9), we obtain

$$\sum_{m=1}^{\infty} j_{\nu_\alpha, m}^2 e^{-j_{\nu_\alpha, m}^2 Y} \leq 2 \left(1 + \left(\frac{\nu_\alpha - \frac{1}{2}}{2}\right)^2\right) e^{-Y\pi^2 \left(\frac{\nu_\alpha - \frac{1}{2}}{4}\right)^2} \left(\frac{4}{eY} + \frac{2^{3/2}}{\pi Y^{3/2}} \int_0^\infty s^2 e^{-s^2} ds\right),$$

hence there exists some C_u such that, when $\nu_\alpha \geq \frac{1}{2}$,

$$(6. 11) \quad \forall Y > 0, \quad \sum_{m=1}^{\infty} j_{\nu_\alpha, m}^2 e^{-j_{\nu_\alpha, m}^2 Y} \leq \frac{\nu_\alpha^2}{C_u Y^{3/2}} e^{-C_u \nu_\alpha^2 Y}.$$

And then, there is some C_u independent of $\alpha \in [1, 2)$ of $T > 0$ and of $\ell > 0$ such that

$$(6. 12) \quad \sum_{m=1}^{\infty} j_{\nu_\alpha, m}^2 e^{-j_{\nu_\alpha, m}^2 \left(\frac{\kappa_\alpha^2 T}{\ell^2 - \alpha}\right)} \leq \frac{1}{C_u \kappa_\alpha^2 \left(\frac{\kappa_\alpha^2 T}{\ell^2 - \alpha}\right)^{3/2}} e^{-\frac{C_u}{\kappa_\alpha^2} \left(\frac{\kappa_\alpha^2 T}{\ell^2 - \alpha}\right)}.$$

That allows us to complete the estimate from above of the null controllability cost: we deduce from (6. 8), (6. 12), (6. 4) that

$$C_{ctr-bd} \leq C'_u \frac{1}{\sqrt{\kappa_\alpha T \ell}} e^{C'_u \frac{\ell^2 - \alpha}{\kappa_\alpha^2 T}} e^{-\frac{C_u}{2} \frac{T}{\ell^2 - \alpha}},$$

which is (2. 11). This completes the proof of Theorem 2.2. \square

7. PROOF OF THEOREM 2.4

In section 4.2, we constructed, at least formally, a control that drives the initial condition u_0 to 0 in time T . This control is given by (4. 8), and depends of a suitable biorthogonal family $\sigma_{\alpha, m}^+$ satisfying (4. 5), and of the norm of the eigenfunctions in the control region. Theorem 6.1 (in fact (6. 5) and (6. 6)) gives the existence and bounds for a biorthogonal family $(\sigma_{\alpha, m}^+)_{m \geq 1}$ satisfying (4. 5). Proposition 2.5 gives an estimate of the norm of the eigenfunctions in the control region (and will be proved in section 9). Here we use these results to prove Theorem 2.4: using Theorem 6.1 and Proposition 2.5, we have

$$\begin{aligned} \|\sigma_{\alpha, m}^+\|_{L^2(0, T)}^2 \frac{1}{\left(\int_a^b \Phi_{\alpha, m}^2\right)^2} &\leq C_u B^*(T, \gamma_{min}) e^{C_u \frac{\ell^2 - \alpha}{\kappa_\alpha^2 T}} e^{-\lambda_{\alpha, m} T} \frac{1}{\left(\gamma_0^*(2 - \alpha)\right)^2} \\ &\leq \frac{C_u B^*(T, \gamma_{min})}{|\gamma_0^*|^2 (2 - \alpha)^2} e^{C_u \frac{\ell^2 - \alpha}{\kappa_\alpha^2 T}} e^{-\kappa_\alpha^2 j_{\nu_\alpha, 1}^2 \frac{T}{\ell^2 - \alpha}}. \end{aligned}$$

Hence, there is some C_u independent of $T > 0$, $\ell > 0$, $\alpha \in [1, 2)$, $m \geq 1$ such that

$$\|\sigma_{\alpha, m}^+\|_{L^2(0, T)}^2 \frac{1}{\left(\int_a^b \Phi_{\alpha, m}^2\right)^2} \leq \frac{C_u B^*(T, \gamma_{min})}{|\gamma_0^*|^2 (2 - \alpha)^2} e^{C_u \frac{\ell^2 - \alpha}{\kappa_\alpha^2 T}} e^{-\frac{1}{C_u} \frac{T}{\ell^2 - \alpha}}.$$

Of course, if $(\mu_{\alpha,m}^0)_m \in \ell^2(\mathbb{N})$, then the series

$$\sum_{m \geq 1} |\mu_{\alpha,m}^0|^2 \|\sigma_{\alpha,m}^+\|_{L^2(0,T)}^2 \frac{1}{\left(\int_a^b \Phi_{\alpha,m}^2\right)^2}$$

is convergent. Hence the control given by the formula (4. 8) is in $L^2((0, \ell) \times (0, T))$, and

$$\begin{aligned} \|h\|_{L^2((0,\ell) \times (0,T))}^2 &= \sum_{m \geq 1} |\mu_{\alpha,m}^0|^2 \|\sigma_{\alpha,m}^+\|_{L^2(0,T)}^2 \frac{1}{\left(\int_a^b \Phi_{\alpha,m}^2\right)^2} \\ &\leq \frac{C_u B^*(T, \gamma_{min})}{|\gamma_0^*|^2 (2-\alpha)^2} e^{C_u \frac{\ell^2-\alpha}{\kappa_\alpha^2 T}} e^{-\frac{1}{C_u} \frac{T}{\ell^2-\alpha}} \sum_{m \geq 1} |\mu_{\alpha,m}^0|^2. \end{aligned}$$

Hence

$$C_{ctr-loc}^2 \leq \frac{C_u B^*(T, \gamma_{min})}{|\gamma_0^*|^2 (2-\alpha)^2} e^{C_u \frac{\ell^2-\alpha}{\kappa_\alpha^2 T}} e^{-\frac{1}{C_u} \frac{T}{\ell^2-\alpha}}.$$

This gives (2. 14) (with another constant C_u). In particular, note that the dependence in the control region appears only in γ_0^* . \square

8. PROOF OF THEOREM 2.3

Given $u_0 \in L^2(0, \ell)$, assume that $h \in L^2((a, b) \times (0, T))$ is a control that drives the solution of (2. 2) to 0 in time T . Denote

$$H(t) := u(a, t).$$

Then the function u satisfies

$$(8. 1) \quad \begin{cases} u_t - (x^\alpha u_x)_x = 0 & x \in (0, a), t > 0, \\ (x^\alpha u_x)(0, t) = 0, & t > 0, \\ u(a, t) = H(t) & t > 0, \\ u(x, 0) = u_0(x), & x \in (0, a) \end{cases}$$

and

$$u(x, T) = 0, \quad x \in (0, a),$$

hence H is a boundary control that drives the solution of (8. 1) to 0 in time T . Let us choose $m \geq 1$ and

$$u_0(x) := \begin{cases} \frac{\sqrt{2\kappa_\alpha}}{a^{\kappa_\alpha} |J'_{\nu_\alpha}(j_{\nu_\alpha, m})|} x^{(1-\alpha)/2} J_{\nu_\alpha}(j_{\nu_\alpha, m} (\frac{x}{a})^{\kappa_\alpha}), & x \in (0, a), \\ 0, & x \in (a, \ell), \end{cases}$$

in such a way that the initial condition of (8. 1) is exactly an eigenfunction of the associated Sturm-Liouville problem. Then we know from subsection 5.7 that

$$(8. 2) \quad \|r_{\alpha, m}\| \|u(a, \cdot)\|_{L^2(0, T)} = \|r_{\alpha, m} H_m\|_{L^2(0, T)} \geq \underline{b}(T, \alpha, m),$$

where \underline{b} is defined in (5. 14), but where a replaces ℓ in the expressions of $r_{\alpha, m}$ and $\underline{b}(T, \alpha, m)$.

On the other hand, energy methods tell us that the control h and the initial condition dominate the solution of (2. 2): indeed, first we have

$$\forall y \geq a, \quad -u(y, t) = \int_y^\ell u_x(x, t) dx,$$

hence

$$u(y, t)^2 = \left(\int_y^\ell u_x(x, t) dx\right)^2 \leq \left(\int_y^\ell x^\alpha u_x^2(x, t) dx\right) \left(\int_y^\ell x^{-\alpha} dx\right),$$

hence

$$\forall y \in [a, \ell), \quad u(y, t)^2 \leq C(\alpha, a, \ell) \int_0^\ell x^\alpha u_x^2(x, t) dx$$

with

$$C(\alpha, a, \ell) = \begin{cases} \frac{1}{(\alpha-1)a^{\alpha-1}} & \text{if } \alpha \in (1, 2), \\ \ln \frac{\ell}{a} & \text{if } \alpha = 1 \end{cases}.$$

Then, multiplying the first equation of (2. 2) by u , we have

$$\int_0^T \int_0^\ell u h \chi_{(a,b)} = \int_0^T \int_0^\ell u(u_t - (x^\alpha u_x)_x) = -\frac{1}{2} \int_0^\ell u_0^2 + \int_0^T \int_0^\ell x^\alpha u_x^2,$$

hence

$$\begin{aligned} \int_0^T \int_0^\ell x^\alpha u_x^2 &= \frac{1}{2} \int_0^\ell u_0^2 + \int_0^T \int_a^b u h \\ &\leq \frac{1}{2} \int_0^\ell u_0^2 + \int_0^T \int_a^b \left(C(\alpha, a, \ell) \int_0^\ell x^\alpha u_x^2(x, t) dx \right)^{1/2} |h| \\ &\leq \frac{1}{2} \int_0^\ell u_0^2 + \frac{1}{2} \int_0^T \int_0^\ell x^\alpha u_x^2(x, t) dx dt + \frac{(b-a)C(\alpha, a, \ell)}{2} \int_0^T \int_a^b h(x, t)^2 dx dt. \end{aligned}$$

We obtain that

$$\int_0^T \int_0^\ell x^\alpha u_x^2 \leq \int_0^\ell u_0^2 + (b-a)C(\alpha, a, \ell) \int_0^T \int_a^b h(x, t)^2 dx dt,$$

hence

$$\begin{aligned} \int_0^T u(a, t)^2 dt &\leq C(\alpha, a, \ell) \int_0^T \int_0^\ell x^\alpha u_x^2 \\ &\leq C(\alpha, a, \ell) \int_0^\ell u_0^2 + (b-a)C(\alpha, a, \ell)^2 \int_0^T \int_a^b h(x, t)^2 dx dt. \end{aligned}$$

The initial condition u_0 that we have chosen has an L^2 -norm equal to 1, hence

$$\int_0^T \int_a^b h(x, t)^2 dx dt \geq \frac{1}{(b-a)C(\alpha, a, \ell)^2} \int_0^T u(a, t)^2 dt - \frac{1}{(b-a)C(\alpha, a, \ell)},$$

and the lower bound (8. 2) of $\|u(a, \cdot)\|_{L^2(0, T)}$ implies that

$$\int_0^T \int_a^b h(x, t)^2 dx dt \geq \frac{1}{(b-a)C(\alpha, a, \ell)^2} \frac{b(T, \alpha, m)^2}{r_{\alpha, m}^2} - \frac{1}{(b-a)C(\alpha, a, \ell)}.$$

As we did in subsection 5.7, choosing $m = 1$, this implies that

$$\begin{aligned} \int_0^T \int_a^b h(x, t)^2 dx dt &\geq \frac{1}{(b-a)C(\alpha, a, \ell)^2} \left[C_u \frac{a^{2(2-\alpha)}}{\kappa_\alpha T a} e^{-2\pi^2 \frac{T}{a^{2-\alpha}}} e^{2C_u \frac{a^{2-\alpha}}{T(2-\alpha)^2}} \right. \\ &\quad \left. e^{-\frac{2}{C_u} \left(\frac{1}{(2-\alpha)^{4/3}} + \frac{a^{1-\alpha/2}}{2-\alpha} \right) (\ln \frac{a^{1-\alpha/2}}{2-\alpha} + \ln \frac{1}{T})} \right] - \frac{1}{(b-a)C(\alpha, a, \ell)}. \end{aligned}$$

Then the null controllability cost for (2. 2) blows up at least exponentially fast when $\alpha \rightarrow 2^-$, as stated in Theorem 2.3. One can note that the bound from below is very poor when ℓ is large. But this is due to the method: indeed, we concentrate the initial condition on the zone at the left of the control region. \square

9. THE EIGENFUNCTIONS IN THE CONTROL REGION (PROPOSITION 2.5)

The goal of this section is to prove Proposition 2.5, that was be useful to prove Theorem 2.4.

9.1. The reduction to an ordinary differential equation question.

Using Proposition 2.4, we note that

$$\begin{aligned}
\int_a^b \Phi_{\alpha,m}(x)^2 dx &= \int_a^b \frac{2\kappa_\alpha}{\ell^{2\kappa_\alpha} |J'_{\nu_\alpha}(j_{\nu_\alpha,m})|^2} x^{1-\alpha} J_{\nu_\alpha}(j_{\nu_\alpha,m}(\frac{x}{\ell})^{\kappa_\alpha})^2 dx \\
&= \frac{2\kappa_\alpha}{\ell^{2\kappa_\alpha} |J'_{\nu_\alpha}(j_{\nu_\alpha,m})|^2} \int_{j_{\nu_\alpha,m}(\frac{a}{\ell})^{\kappa_\alpha}}^{j_{\nu_\alpha,m}(\frac{b}{\ell})^{\kappa_\alpha}} \left(\ell \left(\frac{y}{j_{\nu_\alpha,m}} \right)^{1/\kappa_\alpha} \right)^{1-\alpha} J_{\nu_\alpha}(y)^2 \frac{\ell}{j_{\nu_\alpha,m}^{1/\kappa_\alpha}} \frac{1}{\kappa_\alpha} y^{\frac{1}{\kappa_\alpha}-1} dy \\
&= \frac{2\kappa_\alpha \ell^{2-\alpha}}{\kappa_\alpha \ell^{2\kappa_\alpha} j_{\nu_\alpha,m}^{\frac{1-\alpha}{\kappa_\alpha} + \frac{1}{\kappa_\alpha}} |J'_{\nu_\alpha}(j_{\nu_\alpha,m})|^2} \int_{j_{\nu_\alpha,m}(\frac{a}{\ell})^{\kappa_\alpha}}^{j_{\nu_\alpha,m}(\frac{b}{\ell})^{\kappa_\alpha}} y^{\frac{1-\alpha}{\kappa_\alpha} + \frac{1}{\kappa_\alpha} - 1} J_{\nu_\alpha}(y)^2 dy \\
&= \frac{2}{j_{\nu_\alpha,m}^2 |J'_{\nu_\alpha}(j_{\nu_\alpha,m})|^2} \int_{j_{\nu_\alpha,m}(\frac{a}{\ell})^{\kappa_\alpha}}^{j_{\nu_\alpha,m}(\frac{b}{\ell})^{\kappa_\alpha}} y J_{\nu_\alpha}(y)^2 dy,
\end{aligned}$$

where we used the change of variables

$$y = j_{\nu_\alpha,m} \left(\frac{x}{\ell} \right)^{\kappa_\alpha}, \quad x = \ell \left(\frac{y}{j_{\nu_\alpha,m}} \right)^{1/\kappa_\alpha}, \quad dx = \frac{\ell}{j_{\nu_\alpha,m}^{1/\kappa_\alpha}} \frac{1}{\kappa_\alpha} y^{\frac{1}{\kappa_\alpha}-1} dy.$$

Now introduce the function

$$K_{\nu_\alpha,m}(y) := \sqrt{y} \frac{J_{\nu_\alpha}(y)}{J'_{\nu_\alpha}(j_{\nu_\alpha,m})}.$$

With the help of $K_{\nu_\alpha,m}$, we have

$$(9.1) \quad \int_a^b \Phi_{\alpha,m}(x)^2 dx = \frac{2}{j_{\nu_\alpha,m}^2} \int_{j_{\nu_\alpha,m}(\frac{a}{\ell})^{\kappa_\alpha}}^{j_{\nu_\alpha,m}(\frac{b}{\ell})^{\kappa_\alpha}} K_{\nu_\alpha,m}(y)^2 dy.$$

Moreover, it is well known that $K_{\nu_\alpha,m}$ is solution of the second order ordinary differential equation

$$K_{\nu_\alpha,m}'' + h_{\nu_\alpha}(y) K_{\nu_\alpha,m} = 0,$$

where

$$h_{\nu_\alpha}(y) = 1 - \frac{\nu_\alpha^2 - \frac{1}{4}}{y^2}.$$

(We already recalled this in the proof of Lemma 5.2.) Hence in fact $K_{\nu_\alpha,m}$ solves the Cauchy problem

$$(9.2) \quad \begin{cases} K_{\nu_\alpha,m}'' + h_{\nu_\alpha}(y) K_{\nu_\alpha,m} = 0, \\ K_{\nu_\alpha,m}(j_{\nu_\alpha,m}) = 0, \\ K'_{\nu_\alpha,m}(j_{\nu_\alpha,m}) = \sqrt{j_{\nu_\alpha,m}}, \end{cases}$$

and we want to estimate it on the zone $[j_{\nu_\alpha,m}(\frac{a}{\ell})^{\kappa_\alpha}, j_{\nu_\alpha,m}(\frac{b}{\ell})^{\kappa_\alpha}]$. Let us normalize the localization of the Cauchy conditions and the localization of the integration interval, using a suitable change of variables: consider

$$(9.3) \quad L_{\nu_\alpha,m}(z) = \frac{-1}{\sqrt{j_{\nu_\alpha,m}}} K_{\nu_\alpha,m}(j_{\nu_\alpha,m} - z j_{\nu_\alpha,m}).$$

Then $L_{\nu_\alpha,m}$ solves the Cauchy problem

$$(9.4) \quad \begin{cases} L_{\nu_\alpha,m}'' + k_{\nu_\alpha,m}(z) L_{\nu_\alpha,m} = 0, & \text{with } k_{\nu_\alpha,m}(z) = j_{\nu_\alpha,m}^2 - \frac{\nu_\alpha^2 - \frac{1}{4}}{(1-z)^2}, \\ L_{\nu_\alpha,m}(0) = 0, \\ L'_{\nu_\alpha,m}(0) = j_{\nu_\alpha,m}, \end{cases}$$

and

$$(9.5) \quad \int_a^b \Phi_{\alpha,m}(x)^2 dx = 2 \int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} L_{\nu_\alpha,m}(z)^2 dz.$$

Once again, the term we are interested in is the norm of the solution of a Cauchy problem, but now with Cauchy conditions at the point 0, and we have to estimate its norm on some fixed interval (that does not contain 0). To do this, we are going to study the Cauchy problem (9.4).

9.2. The study of the Cauchy problem: a uniform bound on $L_{\nu_\alpha,m}$.

We begin by the following observation: when $\alpha \rightarrow 2^-$, then

$$1 - \left(\frac{b}{\ell}\right)^{\kappa_\alpha} = 1 - e^{(\frac{1}{2} \ln \frac{b}{\ell})(2-\alpha)} = -\left(\frac{1}{2} \ln \frac{b}{\ell}\right)(2-\alpha) + O((2-\alpha)^2),$$

hence

$$1 - \left(\frac{b}{\ell}\right)^{\kappa_\alpha} \sim \kappa_\alpha \ln \frac{\ell}{b} \quad \text{as } \alpha \rightarrow 2^-,$$

and in the same way

$$1 - \left(\frac{a}{\ell}\right)^{\kappa_\alpha} \sim \kappa_\alpha \ln \frac{\ell}{a} \quad \text{as } \alpha \rightarrow 2^-;$$

hence the integration interval shrinks to 0, and its length satisfies

$$\left(1 - \left(\frac{a}{\ell}\right)^{\kappa_\alpha}\right) - \left(1 - \left(\frac{b}{\ell}\right)^{\kappa_\alpha}\right) \sim \kappa_\alpha \ln \frac{b}{a} \quad \text{as } \alpha \rightarrow 2^-.$$

In particular, there exists some $0 < \gamma_*(a, b, \ell) \leq \gamma^*(a, b, \ell)$ and $\underline{\gamma}(a, b, \ell) > 0$ such that, for all $\alpha \in [1, 2)$,

$$(9.6) \quad \gamma_*(a, b, \ell) \kappa_\alpha \leq 1 - \left(\frac{b}{\ell}\right)^{\kappa_\alpha} < 1 - \left(\frac{a}{\ell}\right)^{\kappa_\alpha} \leq \gamma^*(a, b, \ell) \kappa_\alpha,$$

$$(9.7) \quad \left(1 - \left(\frac{a}{\ell}\right)^{\kappa_\alpha}\right) - \left(1 - \left(\frac{b}{\ell}\right)^{\kappa_\alpha}\right) \geq \underline{\gamma}(a, b, \ell) \kappa_\alpha.$$

Let us prove the following uniform bound:

Lemma 9.1. *There exists C_u independent of $\alpha \in [1, 2)$ and of $m \geq 1$ such that*

$$(9.8) \quad \forall \alpha \in [1, 2[, \forall m \geq 1, \forall z \in [0, 1 - \left(\frac{a}{\ell}\right)^{\kappa_\alpha}], \quad |L_{\nu_\alpha,m}(z)| \leq C_u.$$

Proof of Lemma 9.1. To obtain an integral equation satisfied by $L_{\nu_\alpha,m}$, we write the Cauchy problem (9.4) under the form

$$(9.9) \quad \begin{cases} L''_{\nu_\alpha,m} + j_{\nu_\alpha,m}^2 L_{\nu_\alpha,m} = \frac{\nu_\alpha^2 - \frac{1}{4}}{(1-z)^2} L_{\nu_\alpha,m}, \\ L_{\nu_\alpha,m}(0) = 0, \\ L'_{\nu_\alpha,m}(0) = j_{\nu_\alpha,m}. \end{cases}$$

Since the solution of the Cauchy problem

$$\begin{cases} Y'' + \omega^2 Y = g(z), \\ Y(0) = 0, \\ Y'(0) = \omega \end{cases}$$

is

$$Y(z) = \sin(\omega z) + \frac{1}{\omega} \int_0^z g(s) \sin(\omega(z-s)) ds,$$

we deduce that $L_{\nu_\alpha,m}$ satisfies

$$(9.10) \quad L_{\nu_\alpha,m}(z) = \sin(j_{\nu_\alpha,m} z) + \frac{1}{j_{\nu_\alpha,m}} \int_0^z \frac{\nu_\alpha^2 - \frac{1}{4}}{(1-s)^2} L_{\nu_\alpha,m}(s) \sin(j_{\nu_\alpha,m}(z-s)) ds.$$

Hence

$$|L_{\nu_\alpha, m}(z)| \leq 1 + \frac{1}{j_{\nu_\alpha, m}} \int_0^z \frac{|\nu_\alpha^2 - \frac{1}{4}|}{(1-s)^2} |L_{\nu_\alpha, m}(s)| ds,$$

and the classical Gronwall inequality gives that

$$|L_{\nu_\alpha, m}(z)| \leq e^{\frac{|\nu_\alpha^2 - \frac{1}{4}|}{j_{\nu_\alpha, m}} \int_0^z \frac{1}{(1-s)^2} ds} = e^{\frac{|\nu_\alpha^2 - \frac{1}{4}|}{j_{\nu_\alpha, m}} \frac{z}{1-z}}.$$

But then, since we know from (3. 10) that $j_{\nu_\alpha, m} \geq \nu_\alpha$, we have

$$\forall \alpha \in [1, 2], \forall m \geq 1, \forall z \in [0, 1 - (\frac{a}{\ell})^{\kappa_\alpha}], \quad |L_{\nu_\alpha, m}(z)| \leq e^{\frac{|\nu_\alpha^2 - \frac{1}{4}|}{\nu_\alpha} \frac{1 - (\frac{a}{\ell})^{\kappa_\alpha}}{(\frac{a}{\ell})^{\kappa_\alpha}}}.$$

Using (9. 6), we see that this is uniformly bounded with respect to α , hence we obtain (9. 8). \square

9.3. The L^2 norm of $L_{\nu_\alpha, m}$ for fixed values of ν_α .

The integral expression (9. 10) and the uniform bound (9. 8) allow us to prove the following

Lemma 9.2. *There exists $\bar{\gamma} = \bar{\gamma}(a, b, \ell)$ such that*

$$(9. 11) \quad \forall \alpha \in [1, 2], \forall m \geq 1, \quad \int_{1 - (\frac{b}{\ell})^{\kappa_\alpha}}^{1 - (\frac{a}{\ell})^{\kappa_\alpha}} L_{\nu_\alpha, m}(z)^2 dz \geq \frac{1}{2} \underline{\gamma} \kappa_\alpha - \frac{\bar{\gamma}}{j_{\nu_\alpha, m}}.$$

Proof of Lemma 9.2. From (9. 10) we have

$$\begin{aligned} \int_{1 - (\frac{b}{\ell})^{\kappa_\alpha}}^{1 - (\frac{a}{\ell})^{\kappa_\alpha}} L_{\nu_\alpha, m}(z)^2 dz &= \int_{1 - (\frac{b}{\ell})^{\kappa_\alpha}}^{1 - (\frac{a}{\ell})^{\kappa_\alpha}} \sin^2(j_{\nu_\alpha, m} z) dz \\ &\quad + \int_{1 - (\frac{b}{\ell})^{\kappa_\alpha}}^{1 - (\frac{a}{\ell})^{\kappa_\alpha}} \left(\frac{1}{j_{\nu_\alpha, m}} \int_0^z \frac{\nu_\alpha^2 - \frac{1}{4}}{(1-s)^2} L_{\nu_\alpha, m}(s) \sin(j_{\nu_\alpha, m}(z-s)) ds \right)^2 dz \\ &\quad + \int_{1 - (\frac{b}{\ell})^{\kappa_\alpha}}^{1 - (\frac{a}{\ell})^{\kappa_\alpha}} 2 \sin(j_{\nu_\alpha, m} z) \left(\frac{1}{j_{\nu_\alpha, m}} \int_0^z \frac{\nu_\alpha^2 - \frac{1}{4}}{(1-s)^2} L_{\nu_\alpha, m}(s) \sin(j_{\nu_\alpha, m}(z-s)) ds \right) dz. \end{aligned}$$

Of course

$$\int_{1 - (\frac{b}{\ell})^{\kappa_\alpha}}^{1 - (\frac{a}{\ell})^{\kappa_\alpha}} \left(\frac{1}{j_{\nu_\alpha, m}} \int_0^z \frac{\nu_\alpha^2 - \frac{1}{4}}{(1-s)^2} L_{\nu_\alpha, m}(s) \sin(j_{\nu_\alpha, m}(z-s)) ds \right)^2 dz \geq 0,$$

and

$$\begin{aligned} \int_{1 - (\frac{b}{\ell})^{\kappa_\alpha}}^{1 - (\frac{a}{\ell})^{\kappa_\alpha}} \sin^2(j_{\nu_\alpha, m} z) dz &= \int_{1 - (\frac{b}{\ell})^{\kappa_\alpha}}^{1 - (\frac{a}{\ell})^{\kappa_\alpha}} \frac{1 - \cos(2j_{\nu_\alpha, m} z)}{2} dz \\ &= \frac{1}{2} \left(\left(1 - \left(\frac{a}{\ell}\right)^{\kappa_\alpha}\right) - \left(1 - \left(\frac{b}{\ell}\right)^{\kappa_\alpha}\right) \right) - \frac{1}{2} \left[\frac{\sin 2j_{\nu_\alpha, m} z}{2j_{\nu_\alpha, m}} \right]_{1 - (\frac{b}{\ell})^{\kappa_\alpha}}^{1 - (\frac{a}{\ell})^{\kappa_\alpha}} \\ &\geq \frac{1}{2} \underline{\gamma} \kappa_\alpha - \frac{1}{2j_{\nu_\alpha, m}}. \end{aligned}$$

And using (9. 8) and (9. 6), we have

$$\begin{aligned}
& \left| \int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} 2 \sin(j_{\nu_\alpha, m} z) \left(\frac{1}{j_{\nu_\alpha, m}} \int_0^z \frac{\nu_\alpha^2 - \frac{1}{4}}{(1-s)^2} L_{\nu_\alpha, m}(s) \sin(j_{\nu_\alpha, m}(z-s)) ds \right) dz \right| \\
& \leq \frac{2}{j_{\nu_\alpha, m}} \int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} \left(\int_0^z \frac{|\nu_\alpha^2 - \frac{1}{4}|}{(1-s)^2} C_u ds \right) dz = 2C_u \frac{|\nu_\alpha^2 - \frac{1}{4}|}{j_{\nu_\alpha, m}} \int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} \frac{z}{1-z} dz \\
& \leq \frac{2C_u}{(\frac{a}{\ell})^{\kappa_\alpha}} \frac{|\nu_\alpha^2 - \frac{1}{4}|}{j_{\nu_\alpha, m}} \left[\frac{z^2}{2} \right]_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} \\
& = \frac{C_u}{(\frac{a}{\ell})^{\kappa_\alpha}} \frac{|\nu_\alpha^2 - \frac{1}{4}|}{j_{\nu_\alpha, m}} \left(\left(1 - \left(\frac{a}{\ell}\right)^{\kappa_\alpha}\right) - \left(1 - \left(\frac{b}{\ell}\right)^{\kappa_\alpha}\right) \right) \left(\left(1 - \left(\frac{a}{\ell}\right)^{\kappa_\alpha}\right) + \left(1 - \left(\frac{b}{\ell}\right)^{\kappa_\alpha}\right) \right) \\
& \leq \frac{C_u}{(\frac{a}{\ell})^{\kappa_\alpha}} \frac{|\nu_\alpha^2 - \frac{1}{4}|}{j_{\nu_\alpha, m}} 2\gamma^* (\gamma^* - \gamma_*) \kappa_\alpha^2.
\end{aligned}$$

Hence

$$\int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} L_{\nu_\alpha, m}(z)^2 dz \geq \frac{1}{2} \underline{\gamma} \kappa_\alpha - \frac{1}{2j_{\nu_\alpha, m}} - \frac{C_u}{(\frac{a}{\ell})^{\kappa_\alpha}} \frac{|\nu_\alpha^2 - \frac{1}{4}|}{j_{\nu_\alpha, m}} 2\gamma^* (\gamma^* - \gamma_*) \kappa_\alpha^2.$$

Hence there exists $\bar{\gamma} = \bar{\gamma}(a, b, \ell)$ such that

$$\int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} L_{\nu_\alpha, m}(z)^2 dz \geq \frac{1}{2} \underline{\gamma} \kappa_\alpha - \frac{\bar{\gamma}}{j_{\nu_\alpha, m}}.$$

This implies (9. 11) . □

9.4. A first consequence for the eigenfunctions on the control region.

The consequence of (9. 11) is immediate: combining (9. 5) and (9. 11), we obtain

$$(9. 12) \quad \int_a^b \Phi_{\alpha, m}(x)^2 dx \geq \underline{\gamma} \kappa_\alpha - \frac{2\bar{\gamma}}{j_{\nu_\alpha, m}},$$

hence we find again what is well-known at least in the nondegenerate case ([38]):

$$\lim_{m \rightarrow \infty} \int_a^b \Phi_{\alpha, m}(x)^2 dx > 0,$$

hence the sequence of positive terms $(\int_a^b \Phi_{\alpha, m}^2)_{m \geq 1}$ is bounded from below by a positive constant. But this constant may depend on α . At least, we obtain the following uniform result: given $\bar{\alpha}^* \in [1, 2)$, there exists $\bar{m}^* \geq 1$ such that

$$\forall \alpha \in [1, \bar{\alpha}^*], \forall m \geq \bar{m}^*, \quad \int_a^b \Phi_{\alpha, m}(x)^2 dx \geq \frac{1}{2} \underline{\gamma} \kappa_{\bar{\alpha}^*}.$$

Hence, since $\Phi_{\alpha, m}$ depends continuously on the parameter α , we obtain that, given $\bar{\alpha}^* \in [1, 2)$, the sequences $(\int_a^b \Phi_{\alpha, m}^2)_{m \geq 1}$ are bounded from below by a positive constant, uniformly with respect to $\alpha \in [1, \bar{\alpha}^*]$:

$$(9. 13) \quad \forall \bar{\alpha}^* \in [1, 2), \exists \underline{\gamma}^* > 0, \forall \alpha \in [1, \bar{\alpha}^*], \forall m \geq 1, \quad \int_a^b \Phi_{\alpha, m}(x)^2 dx \geq \underline{\gamma}^* \kappa_{\bar{\alpha}^*}.$$

This is not sufficient to conclude, since we want a lower estimate valid for all $\alpha \in [1, 2)$, but this is a first step, and we will use this partial result later.

9.5. Another integral equation for $L_{\nu_\alpha, m}$ when ν_α is large.

Now, we would like to obtain bounds from below when ν_α is large. In this case, we have to integrate $L_{\nu_\alpha, m}^2$ in an interval close to 0. So, since we are interested in looking what happens near 0, it is more interesting to write

$$\begin{aligned} k_{\nu_\alpha, m}(z) &= j_{\nu_\alpha, m}^2 - \frac{\nu_\alpha^2 - \frac{1}{4}}{(1-z)^2} = j_{\nu_\alpha, m}^2 - (\nu_\alpha^2 - \frac{1}{4}) \left(1 + \frac{1}{(1-z)^2} - 1\right) \\ &= \left(j_{\nu_\alpha, m}^2 - \nu_\alpha^2 + \frac{1}{4}\right) - (\nu_\alpha^2 - \frac{1}{4}) \frac{2z - z^2}{(1-z)^2}. \end{aligned}$$

Then we can write the Cauchy problem (9. 4) under the form

$$(9. 14) \quad \begin{cases} L''_{\nu_\alpha, m} + \left(j_{\nu_\alpha, m}^2 - \nu_\alpha^2 + \frac{1}{4}\right) L_{\nu_\alpha, m} = (\nu_\alpha^2 - \frac{1}{4}) \frac{2z - z^2}{(1-z)^2} L_{\nu_\alpha, m}, \\ L_{\nu_\alpha, m}(0) = 0, \\ L'_{\nu_\alpha, m}(0) = j_{\nu_\alpha, m}. \end{cases}$$

Since the solution of the Cauchy problem

$$\begin{cases} Y'' + \omega^2 Y = g(z), \\ Y(0) = 0, \\ Y'(0) = \rho \end{cases}$$

is

$$Y(z) = \frac{\rho}{\omega} \sin \omega z + \frac{1}{\omega} \int_0^z g(s) \sin \omega(z-s) ds,$$

we obtain a new integral equation satisfied by $L_{\nu_\alpha, m}$: denote

$$\omega_{\nu_\alpha, m} := \sqrt{j_{\nu_\alpha, m}^2 - \nu_\alpha^2 + \frac{1}{4}};$$

then we have

$$\begin{aligned} L_{\nu_\alpha, m}(z) &= \frac{j_{\nu_\alpha, m}}{\omega_{\nu_\alpha, m}} \sin(\omega_{\nu_\alpha, m} z) \\ &\quad + \frac{1}{\omega_{\nu_\alpha, m}} \int_0^z (\nu_\alpha^2 - \frac{1}{4}) \frac{2s - s^2}{(1-s)^2} L_{\nu_\alpha, m}(s) \sin(\omega_{\nu_\alpha, m}(z-s)) ds. \end{aligned}$$

Hence

$$\begin{aligned} (9. 15) \quad &\int_{1 - (\frac{b}{\ell})^{\kappa_\alpha}}^{1 - (\frac{a}{\ell})^{\kappa_\alpha}} L_{\nu_\alpha, m}(z)^2 dz = \int_{1 - (\frac{b}{\ell})^{\kappa_\alpha}}^{1 - (\frac{a}{\ell})^{\kappa_\alpha}} \left(\frac{j_{\nu_\alpha, m}}{\omega_{\nu_\alpha, m}} \sin(\omega_{\nu_\alpha, m} z) \right. \\ &\quad \left. + \frac{1}{\omega_{\nu_\alpha, m}} \int_0^z (\nu_\alpha^2 - \frac{1}{4}) \frac{2s - s^2}{(1-s)^2} L_{\nu_\alpha, m}(s) \sin(\omega_{\nu_\alpha, m}(z-s)) ds \right)^2 dz \\ &= \frac{j_{\nu_\alpha, m}^2}{\omega_{\nu_\alpha, m}^2} \int_{1 - (\frac{b}{\ell})^{\kappa_\alpha}}^{1 - (\frac{a}{\ell})^{\kappa_\alpha}} \sin^2(\omega_{\nu_\alpha, m} z) dz \\ &\quad + \frac{1}{\omega_{\nu_\alpha, m}^2} \int_{1 - (\frac{b}{\ell})^{\kappa_\alpha}}^{1 - (\frac{a}{\ell})^{\kappa_\alpha}} \left(\int_0^z (\nu_\alpha^2 - \frac{1}{4}) \frac{2s - s^2}{(1-s)^2} L_{\nu_\alpha, m}(s) \sin(\omega_{\nu_\alpha, m}(z-s)) ds \right)^2 dz \\ &+ 2 \frac{j_{\nu_\alpha, m}}{\omega_{\nu_\alpha, m}^2} \int_{1 - (\frac{b}{\ell})^{\kappa_\alpha}}^{1 - (\frac{a}{\ell})^{\kappa_\alpha}} \sin(\omega_{\nu_\alpha, m} z) \left(\int_0^z (\nu_\alpha^2 - \frac{1}{4}) \frac{2s - s^2}{(1-s)^2} L_{\nu_\alpha, m}(s) \sin(\omega_{\nu_\alpha, m}(z-s)) ds \right) dz. \end{aligned}$$

We are going to study the behavior of the first and third term of the right hand side of (9. 15), the second one being nonnegative.

9.6. The L^2 norm of $L_{\nu_\alpha, m}$ for large values of ν_α .

9.6.a. *The first term of (9. 15).*

We study

$$(9. 16) \quad \int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} \sin^2 \omega_{\nu_\alpha, m} z \, dz.$$

It appears that we need to distinguish the cases $\omega_{\nu_\alpha, m} \kappa_\alpha$ small and $\omega_{\nu_\alpha, m} \kappa_\alpha$ not small: indeed,

$$\int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} \sin^2(\omega_{\nu_\alpha, m} z) \, dz = \frac{1}{\omega_{\nu_\alpha, m}} \int_{\omega_{\nu_\alpha, m}[1-(\frac{b}{\ell})^{\kappa_\alpha}]}^{\omega_{\nu_\alpha, m}[1-(\frac{a}{\ell})^{\kappa_\alpha}]} \sin^2 x \, dx,$$

and we derive from the elementary convexity inequalities:

$$(9. 17) \quad \forall \mu \geq 0, \forall u \in [0, 1], \quad (1 - e^{-\mu})u \leq 1 - e^{-\mu u} \leq \mu u,$$

that

- first, using (9. 17) with $u = \kappa_\alpha$ and $\mu = \ln \frac{\ell}{a}$, we have

$$(9. 18) \quad [1 - (\frac{a}{\ell})^{\kappa_\alpha}] = [1 - e^{-\kappa_\alpha \ln \frac{\ell}{a}}] \leq \ln \frac{\ell}{a} \kappa_\alpha,$$

- next, using (9. 17) with $u = \kappa_\alpha$ and $\mu = \ln \frac{\ell}{b}$, we have

$$(9. 19) \quad [1 - (\frac{b}{\ell})^{\kappa_\alpha}] = [1 - e^{-\kappa_\alpha \ln \frac{\ell}{b}}] \geq (1 - e^{-\ln \frac{\ell}{b}}) \kappa_\alpha = (1 - \frac{b}{\ell}) \kappa_\alpha,$$

and (9. 18)-(9. 19) give that

$$(1 - \frac{b}{\ell}) \omega_{\nu_\alpha, m} \kappa_\alpha \leq \omega_{\nu_\alpha, m} [1 - (\frac{b}{\ell})^{\kappa_\alpha}] \leq \omega_{\nu_\alpha, m} [1 - (\frac{a}{\ell})^{\kappa_\alpha}] \leq (\ln \frac{\ell}{a}) \omega_{\nu_\alpha, m} \kappa_\alpha.$$

Hence the bounds of the integral appearing in (9. 16) are both small or both non small, depending on the value of $\omega_{\nu_\alpha, m} \kappa_\alpha$.

We prove the following

Lemma 9.3. *There exists $\eta_0 = \eta_0(a, b, \ell) > 0$, and $\gamma_0 = \gamma_0(a, b, \ell) > 0$ both independent of $\alpha \in [1, 2)$ and of $m \geq 1$ such that*

- if $\omega_{\nu_\alpha, m} \kappa_\alpha \leq \eta_0$, then

$$(9. 20) \quad \int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} \sin^2(\omega_{\nu_\alpha, m} z) \, dz \geq \gamma_0 \omega_{\nu_\alpha, m}^2 \kappa_\alpha^3.$$

- if $\omega_{\nu_\alpha, m} \kappa_\alpha \geq \eta_0$, then

$$(9. 21) \quad \int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} \sin^2(\omega_{\nu_\alpha, m} z) \, dz \geq \gamma_0 \kappa_\alpha.$$

Remark 9.1. A similar property of the function sinus appears in Haraux [34] (Lemma 1.3.2) and [52] (Theorem 1). In our case, we have to bound from below the integrals of $z \mapsto \sin^2(\omega_{\nu_\alpha, m} z)$ with respect to the size of the integration zone (and this size is small, of the order κ_α), and the coefficients $\omega_{\nu_\alpha, m}$ that appear are non integer and possibly small.

Proof of Lemma 9.3. It comes from the following observations:

- First,

$$(9. 22) \quad 0 < A < B \leq \frac{\pi}{2} \quad \implies \quad \int_A^B \sin^2 x \, dx \geq \frac{4}{\pi^2} A^2 (B - A).$$

Indeed, if $0 < A < B \leq \frac{\pi}{2}$, then

$$\forall x \in [A, B], \quad \sin x \geq \frac{2}{\pi} x,$$

hence

$$\int_A^B \sin^2 x \, dx \geq \frac{4}{\pi^2} A^2 (B - A).$$

- On the other hand, consider η_1 , A and B such that

$$0 < \eta_1 \leq A < B, \quad \text{and} \quad B - A \geq 2\eta_1.$$

Then, first the function

$$s \mapsto \int_s^{s+\eta_1} \sin^2 x \, dx$$

is continuous, 2π -periodic, and positive, hence is bounded from below by a positive constant, denoted γ_1 . Next, there exists one and only one integer k such that

$$k\eta_1 \leq B - A < (k + 1)\eta_1,$$

then

$$\begin{aligned} \int_A^B \sin^2 x \, dx &\geq \int_A^{A+k\eta_1} \sin^2 x \, dx = \sum_{j=0}^{k-1} \int_{A+j\eta_1}^{A+(j+1)\eta_1} \sin^2 x \, dx \\ &\geq \sum_{j=0}^{k-1} \gamma_1 = k\gamma_1 \geq \gamma_1 \left(\frac{B-A}{\eta_1} - 1 \right) = \gamma_1 \left(\frac{B-A}{2\eta_1} + \frac{B-A}{2\eta_1} - 1 \right) \geq \frac{\gamma_1}{2\eta_1} (B-A). \end{aligned}$$

Hence

$$(9.23) \quad \left(0 < \eta_1 \leq A < B, \quad \text{and} \quad B - A \geq 2\eta_1 \right) \\ \implies \int_A^B \sin^2 x \, dx \geq \frac{\gamma_1}{2\eta_1} (B-A) \quad \text{with} \quad \gamma_1 = \gamma_1(\eta_1).$$

Now we are in position to conclude the proof of Lemma 9.3. This is based on the observation that we are in one of the two situations studied previously: using (9.6), we see that there is some η_0 such that

$$\omega_{\nu_\alpha, m} \kappa_\alpha \leq \eta_0 \implies 0 < \omega_{\nu_\alpha, m} \left[1 - \left(\frac{b}{\ell} \right)^{\kappa_\alpha} \right] < \omega_{\nu_\alpha, m} \left[1 - \left(\frac{a}{\ell} \right)^{\kappa_\alpha} \right] \leq \frac{\pi}{2};$$

in this case, (9.22) gives that

$$\begin{aligned} &\int_{\omega_{\nu_\alpha, m} \left[1 - \left(\frac{b}{\ell} \right)^{\kappa_\alpha} \right]}^{\omega_{\nu_\alpha, m} \left[1 - \left(\frac{a}{\ell} \right)^{\kappa_\alpha} \right]} \sin^2 x \, dx \\ &\geq \frac{4}{\pi^2} \left(\omega_{\nu_\alpha, m} \left[1 - \left(\frac{b}{\ell} \right)^{\kappa_\alpha} \right] \right)^2 \left(\omega_{\nu_\alpha, m} \left[1 - \left(\frac{a}{\ell} \right)^{\kappa_\alpha} \right] - \omega_{\nu_\alpha, m} \left[1 - \left(\frac{b}{\ell} \right)^{\kappa_\alpha} \right] \right) \\ &= \frac{4}{\pi^2} \omega_{\nu_\alpha, m}^3 \left[1 - \left(\frac{b}{\ell} \right)^{\kappa_\alpha} \right]^2 \left(\left[1 - \left(\frac{a}{\ell} \right)^{\kappa_\alpha} \right] - \left[1 - \left(\frac{b}{\ell} \right)^{\kappa_\alpha} \right] \right), \end{aligned}$$

hence, thanks to (9.6) and (9.7), there exists some $\gamma(a, b, \ell) > 0$ such that

$$\omega_{\nu_\alpha, m} \kappa_\alpha \leq \eta_0 \implies \int_{\omega_{\nu_\alpha, m} \left[1 - \left(\frac{b}{\ell} \right)^{\kappa_\alpha} \right]}^{\omega_{\nu_\alpha, m} \left[1 - \left(\frac{a}{\ell} \right)^{\kappa_\alpha} \right]} \sin^2 x \, dx \geq \gamma(a, b, \ell) \omega_{\nu_\alpha, m}^3 \kappa_\alpha^3,$$

which gives (9.20).

Now if $\omega_{\nu_\alpha, m} \kappa_\alpha \geq \eta_0$: then, thanks to (9.6) and (9.7), there is some $\eta_1 = \eta_1(a, b, \ell)$ such that

$$\eta_1 < \omega_{\nu_\alpha, m} \left[1 - \left(\frac{b}{\ell} \right)^{\kappa_\alpha} \right] < \omega_{\nu_\alpha, m} \left[1 - \left(\frac{a}{\ell} \right)^{\kappa_\alpha} \right],$$

and

$$\omega_{\nu_\alpha, m} \left[1 - \left(\frac{a}{\ell} \right)^{\kappa_\alpha} \right] - \omega_{\nu_\alpha, m} \left[1 - \left(\frac{b}{\ell} \right)^{\kappa_\alpha} \right] \geq 2\eta_1.$$

Then (9. 23) gives that there exists $\gamma_1 = \gamma_1(a, b, \ell)$ such that

$$\int_{\omega_{\nu_\alpha, m}[1-(\frac{b}{\ell})^{\kappa_\alpha}]}^{\omega_{\nu_\alpha, m}[1-(\frac{a}{\ell})^{\kappa_\alpha}]} \sin^2 x \, dx \geq \frac{\gamma_1}{2\eta_1} \left(\omega_{\nu_\alpha, m}[1-(\frac{a}{\ell})^{\kappa_\alpha}] - \omega_{\nu_\alpha, m}[1-(\frac{b}{\ell})^{\kappa_\alpha}] \right),$$

hence, once again with (9. 7), we obtain that

$$\omega_{\nu_\alpha, m} \kappa_\alpha \geq \eta_0 \quad \implies \quad \int_{\omega_{\nu_\alpha, m}[1-(\frac{b}{\ell})^{\kappa_\alpha}]}^{\omega_{\nu_\alpha, m}[1-(\frac{a}{\ell})^{\kappa_\alpha}]} \sin^2 x \, dx \geq \frac{\gamma_1}{2\eta_1} \omega_{\nu_\alpha, m} \underline{\gamma} \kappa_\alpha,$$

which gives (9. 21). \square

9.6.b. *The third term of (9. 15).*

We study

$$M_3 := \int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} \sin(\omega_{\nu_\alpha, m} z) \left(\int_0^z (\nu_\alpha^2 - \frac{1}{4}) \frac{2s - s^2}{(1-s)^2} L_{\nu_\alpha, m}(s) \sin(\omega_{\nu_\alpha, m}(z-s)) \, ds \right) dz,$$

and we prove the following

Lemma 9.4. *Choose $\eta_0 = \eta_0(a, b, \ell) > 0$ given in Lemma 9.3. Then there exists $\gamma'_0 = \gamma'_0(a, b, \ell) > 0$ independent of $\alpha \in [1, 2)$ and of $m \geq 1$ such that*

- if $\omega_{\nu_\alpha, m} \kappa_\alpha \leq \eta_0$, then

$$(9. 24) \quad |M_3| \leq \gamma'_0 \omega_{\nu_\alpha, m} \kappa_\alpha^2,$$

- if $\omega_{\nu_\alpha, m} \kappa_\alpha \geq \eta_0$, then

$$(9. 25) \quad |M_3| \leq \gamma'_0 \kappa_\alpha,$$

hence in any case

$$(9. 26) \quad |M_3| \leq \gamma'_0 \kappa_\alpha.$$

Proof of Lemma 9.4. First we prove (9. 24). Assume that $\omega_{\nu_\alpha, m} \kappa_\alpha \leq \eta_0$. Then, since we have proved in Lemma 9.1 that $L_{\nu_\alpha, m}$ is uniformly bounded in $[0, 1-(\frac{a}{\ell})^{\kappa_\alpha}]$, we have

$$\begin{aligned} |M_3| &\leq \int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} |\sin \omega_{\nu_\alpha, m} z| \left(\int_0^z |\nu_\alpha^2 - \frac{1}{4}| \frac{|2s - s^2|}{(1-s)^2} |L_{\nu_\alpha, m}(s)| |\sin \omega_{\nu_\alpha, m}(z-s)| \, ds \right) dz \\ &\leq \int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} \omega_{\nu_\alpha, m} z \left(\int_0^z |\nu_\alpha^2 - \frac{1}{4}| \frac{2s}{(1-s)^2} C_u \, ds \right) dz \\ &\leq \omega_{\nu_\alpha, m} |\nu_\alpha^2 - \frac{1}{4}| C'_u \int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} z^3 \, dz \\ &\leq \omega_{\nu_\alpha, m} |\nu_\alpha^2 - \frac{1}{4}| C'_u (1 - (\frac{a}{\ell})^{\kappa_\alpha})^3 \left((1 - (\frac{a}{\ell})^{\kappa_\alpha}) - 1 - (\frac{b}{\ell})^{\kappa_\alpha} \right) \\ &\leq C''_u \omega_{\nu_\alpha, m} |\nu_\alpha^2 - \frac{1}{4}| \kappa_\alpha^4 \leq C''_u \omega_{\nu_\alpha, m} \kappa_\alpha^{-2} \kappa_\alpha^4 = C''_u \omega_{\nu_\alpha, m} \kappa_\alpha^2, \end{aligned}$$

which is (9. 24) (which implies (9. 26)).

Now we prove (9. 25). Assume that $\omega_{\nu_\alpha, m} \kappa_\alpha \geq \eta_0$. Then, once again using the fact that $L_{\nu_\alpha, m}$ is uniformly bounded in $[0, 1-(\frac{a}{\ell})^{\kappa_\alpha}]$ (Lemma 9.1), we have

$$\begin{aligned} |M_3| &\leq \int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} |\sin \omega_{\nu_\alpha, m} z| \left(\int_0^z |\nu_\alpha^2 - \frac{1}{4}| \frac{|2s - s^2|}{(1-s)^2} |L_{\nu_\alpha, m}(s)| |\sin \omega_{\nu_\alpha, m}(z-s)| \, ds \right) dz \\ &\leq \int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} \left(\int_0^z |\nu_\alpha^2 - \frac{1}{4}| C'_u s \, ds \right) dz \leq C'_u |\nu_\alpha^2 - \frac{1}{4}| \left[\frac{z^3}{6} \right]_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} \\ &\leq C'''_u |\nu_\alpha^2 - \frac{1}{4}| (1 - (\frac{a}{\ell})^{\kappa_\alpha})^2 \left((1 - (\frac{a}{\ell})^{\kappa_\alpha}) - 1 - (\frac{b}{\ell})^{\kappa_\alpha} \right) \leq C'''_u \kappa_\alpha^{-2} \kappa_\alpha^3 = C'''_u \kappa_\alpha \end{aligned}$$

which is (9. 25). \square

9.6.c. *The L^2 norm of $L_{\nu_\alpha, m}$ for large values of ν_α .*

We prove the following

Lemma 9.5. *Choose $\eta_0 = \eta_0(a, b, \ell) > 0$ and $\gamma_0 = \gamma_0(a, b, \ell) > 0$ given in Lemma 9.3. Then there exists $\bar{\alpha} \in [1, 2)$ such that*

$$(9. 27) \quad \forall \alpha \in [\bar{\alpha}, 2), \forall m \geq 1, \quad \int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} L_{\nu_\alpha, m}(z)^2 dz \geq \frac{\gamma_0}{2} \frac{j_{\nu_\alpha, m}^2 \kappa_\alpha^3}{1 + \omega_{\nu_\alpha, m}^2 \kappa_\alpha^2}.$$

Proof of Lemma 9.5. We start from (9. 15), that gives

$$(9. 28) \quad \int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} L_{\nu_\alpha, m}(z)^2 dz \geq \frac{j_{\nu_\alpha, m}^2}{\omega_{\nu_\alpha, m}^2} \int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} \sin^2(\omega_{\nu_\alpha, m} z) dz \\ + 2 \frac{j_{\nu_\alpha, m}}{\omega_{\nu_\alpha, m}^2} \int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} \sin(\omega_{\nu_\alpha, m} z) \left(\int_0^z (\nu_\alpha^2 - \frac{1}{4}) \frac{2s - s^2}{(1-s)^2} L_{\nu_\alpha, m}(s) \sin(\omega_{\nu_\alpha, m}(z-s)) ds \right) dz.$$

First we prove (9. 27) when $\omega_{\nu_\alpha, m} \kappa_\alpha \leq \eta_0$. Using (9. 20) and (9. 24) in (9. 28), we have

$$\int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} L_{\nu_\alpha, m}(z)^2 dz \geq \frac{j_{\nu_\alpha, m}^2}{\omega_{\nu_\alpha, m}^2} \gamma_0 \omega_{\nu_\alpha, m}^2 \kappa_\alpha^3 - 2 \frac{j_{\nu_\alpha, m}}{\omega_{\nu_\alpha, m}^2} \gamma_0' \omega_{\nu_\alpha, m} \kappa_\alpha^2 \\ = \gamma_0 j_{\nu_\alpha, m}^2 \kappa_\alpha^3 - 2 \gamma_0' \frac{j_{\nu_\alpha, m}}{\omega_{\nu_\alpha, m}} \kappa_\alpha^2 = j_{\nu_\alpha, m}^2 \kappa_\alpha^3 \left(\gamma_0 - \frac{2 \gamma_0'}{j_{\nu_\alpha, m} \omega_{\nu_\alpha, m} \kappa_\alpha} \right).$$

Now remember that (3. 10) ([55]) says that

$$j_{\nu, k} \geq \nu - \frac{a_1}{2^{1/3}} \nu^{1/3},$$

where $a_1 < 0$. Hence

$$j_{\nu_\alpha, m} - \nu_\alpha \geq -\frac{a_1}{2^{1/3}} \nu_\alpha^{1/3},$$

and

$$j_{\nu_\alpha, m} + \nu_\alpha \geq 2\nu_\alpha,$$

therefore

$$\omega_{\nu_\alpha, m} = \sqrt{j_{\nu_\alpha, m}^2 - \nu_\alpha^2 + \frac{1}{4}} \geq \sqrt{-\frac{2a_1}{2^{1/3}} \nu_\alpha^{4/3}}.$$

Since $j_{\nu_\alpha, m} \geq \nu_\alpha$, this implies that

$$j_{\nu_\alpha, m} \omega_{\nu_\alpha, m} \kappa_\alpha \geq \left(-\frac{2a_1}{2^{1/3}} \right)^{1/2} \nu_\alpha^{5/3} \kappa_\alpha.$$

That last quantity goes to infinity when $\alpha \rightarrow 2^-$, hence there exists $\alpha_0 \in [1, 2)$ such that

$$\forall \alpha \in [\alpha_0, 2), \forall m \geq 1, \quad \int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} L_{\nu_\alpha, m}(z)^2 dz \geq \frac{\gamma_0}{2} j_{\nu_\alpha, m}^2 \kappa_\alpha^3,$$

which implies (9. 27) when $\omega_{\nu_\alpha, m} \kappa_\alpha \leq \eta_0$.

Next we prove (9. 27) when $\omega_{\nu_\alpha, m} \kappa_\alpha \geq \eta_0$. Using (9. 21) and (9. 25) in (9. 28), we have

$$\int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} L_{\nu_\alpha, m}(z)^2 dz \geq \frac{j_{\nu_\alpha, m}^2}{\omega_{\nu_\alpha, m}^2} \gamma_0 \kappa_\alpha - 2 \frac{j_{\nu_\alpha, m}}{\omega_{\nu_\alpha, m}^2} \gamma_0' \kappa_\alpha = \frac{j_{\nu_\alpha, m}^2 \kappa_\alpha}{\omega_{\nu_\alpha, m}^2} \left(\gamma_0 - 2 \frac{\gamma_0'}{j_{\nu_\alpha, m}} \right).$$

Hence, once again, there exists $\alpha_1 \in [1, 2)$ such that

$$\forall \alpha \in [\alpha_1, 2), \forall m \geq 1, \quad \int_{1-(\frac{b}{\ell})^{\kappa_\alpha}}^{1-(\frac{a}{\ell})^{\kappa_\alpha}} L_{\nu_\alpha, m}(z)^2 dz \geq \frac{\gamma_0}{2} \frac{j_{\nu_\alpha, m}^2 \kappa_\alpha}{\omega_{\nu_\alpha, m}^2}.$$

Finally, since

$$\inf\left\{a, \frac{1}{b}\right\} \geq \frac{a}{1+ab},$$

we obtain that

$$\begin{aligned} \int_{1-(\frac{b}{2})^{\kappa_\alpha}}^{1-(\frac{a}{2})^{\kappa_\alpha}} L_{\nu_\alpha, m}(z)^2 dz &\geq \inf\left\{\frac{\gamma_0}{2} j_{\nu_\alpha, m}^2 \kappa_\alpha^3, \frac{\gamma_0}{2} \frac{j_{\nu_\alpha, m}^2 \kappa_\alpha}{\omega_{\nu_\alpha, m}^2}\right\} \\ &= \frac{\gamma_0}{2} j_{\nu_\alpha, m}^2 \kappa_\alpha^3 \inf\left\{1, \frac{1}{\omega_{\nu_\alpha, m}^2 \kappa_\alpha^2}\right\} \geq \frac{\gamma_0}{2} j_{\nu_\alpha, m}^2 \kappa_\alpha^3 \frac{1}{1 + \omega_{\nu_\alpha, m}^2 \kappa_\alpha^2}, \end{aligned}$$

which is (9. 27). \square

9.7. Proof of Proposition 2.5.

We deduce from (9. 5) and (9. 27) that, for $\alpha \geq \bar{\alpha}$,

$$\begin{aligned} \forall m \geq 1, \quad \int_a^b \Phi_{\alpha, m}(x)^2 dx &\geq \gamma_0 \frac{j_{\nu_\alpha, m}^2 \kappa_\alpha^3}{1 + \omega_{\nu_\alpha, m}^2 \kappa_\alpha^2} = \gamma_0 \kappa_\alpha \frac{j_{\nu_\alpha, m}^2 \kappa_\alpha^2}{1 + \omega_{\nu_\alpha, m}^2 \kappa_\alpha^2} \\ &= \gamma_0 \kappa_\alpha \frac{\omega_{\nu_\alpha, m}^2 \kappa_\alpha^2 + (\nu_\alpha^2 - \frac{1}{4}) \kappa_\alpha^2}{1 + \omega_{\nu_\alpha, m}^2 \kappa_\alpha^2}. \end{aligned}$$

Since $\nu_\alpha \kappa_\alpha \rightarrow \frac{1}{2}$ as $\alpha \rightarrow 2^-$, there exists $\bar{\alpha}^* \in [\bar{\alpha}, 2)$ such that $(\nu_\alpha^2 - \frac{1}{4}) \kappa_\alpha^2 \geq \frac{1}{8}$ for all $\alpha \in [\bar{\alpha}^*, 2)$. Then, for all $\alpha \in [\bar{\alpha}^*, 2)$, we have

$$\frac{\omega_{\nu_\alpha, m}^2 \kappa_\alpha^2 + (\nu_\alpha^2 - \frac{1}{4}) \kappa_\alpha^2}{1 + \omega_{\nu_\alpha, m}^2 \kappa_\alpha^2} \geq \frac{\omega_{\nu_\alpha, m}^2 \kappa_\alpha^2 + \frac{1}{8}}{1 + \omega_{\nu_\alpha, m}^2 \kappa_\alpha^2} \geq \frac{1}{8}.$$

Hence

$$(9. 29) \quad \forall \alpha \in [\bar{\alpha}^*, 2), \forall m \geq 1, \quad \int_a^b \Phi_{\alpha, m}(x)^2 dx \geq \frac{\gamma_0}{8} \kappa_\alpha.$$

But we already proved in (9. 13) that $(\int_a^b \Phi_{\alpha, m}^2)_{m \geq 1}$ is uniformly bounded from below when $\alpha \in [1, \bar{\alpha}^*]$. Hence (9. 13) and (9. 29) give (2. 18) and the proof of Proposition 2.5 is completed. \square

Acknowledgements. The authors wish to thank the referees for their constructive comments.

REFERENCES

- [1] F. Alabau-Boussouira, P. Cannarsa and G. Fragnelli, *Carleman estimates for degenerate parabolic operators with applications to null controllability*, J. evol. equ. 6 (2006).
- [2] F. Ammar Khodja, A. Benabdallah, M. González-Burgos, L. de Teresa, *The Kalman condition for the boundary controllability of coupled parabolic systems. Bounds on biorthogonal families to complex matrix exponentials*, J. Math. Pures Appl. 96 (2011), p. 555-590.
- [3] F. Ammar Khodja, A. Benabdallah, M. González-Burgos, L. de Teresa, *Minimal time for the null controllability of parabolic systems: The effect of the condensation index of complex sequences*, Journal of Functional Analysis 267 (2014), p. 2077-2151.
- [4] K. Beauchard, P. Cannarsa, R. Guglielmi, *Null controllability of Grushin-type operators in dimension two*, J. Eur. Math. Soc. (JEMS) 16 (2014), No. 1, 67-101.
- [5] A. Benabdallah, F. Boyer, M. González-Burgos, G. Olive, *Sharp estimates of the one-dimensional boundary control cost for parabolic systems and applications to the N-dimensional boundary null controllability in cylindrical domains*, SIAM J. Control Optim., Vol. 52, No. 5, pp. 2970-3001.
- [6] A. Bensoussan, G. Da Prato, M. C. Delfour, S. K. Mitter, *Representation and control of infinite-dimensional systems. Vol. 1*, Systems and Control: Foundations and Applications, Birkhauser Boston, 1992.
- [7] M. Campiti, G. Metafuno, D. Pallara, *Degenerate self-adjoint evolution equations on the unit interval*, Semigroup Forum, 57, (1998), 1-36.
- [8] P. Cannarsa, G. Fragnelli, D. Rocchetti, *Null controllability of degenerate parabolic operators with drift*, Netw. Heterog. Media 2 (2007), no. 4, 695-715.

- [9] P. Cannarsa, G. Fragnelli, D. Rocchetti, *Controllability results for a class of one-dimensional degenerate parabolic problems in nondivergence form*, J. evol. equ. 8 (2008), 583–616.
- [10] P. Cannarsa, P. Martinez, J. Vancostenoble, *Null Controllability of degenerate heat equations*, Adv. Differential Equations 10 (2005), 2, 153-190.
- [11] P. Cannarsa, P. Martinez, J. Vancostenoble, *Carleman estimates for a class of degenerate parabolic operators*, SIAM J. Control Optim. 47, (2008), no. 1, 1–19.
- [12] P. Cannarsa, P. Martinez, J. Vancostenoble, *Global Carleman estimates for degenerate parabolic operators with applications*, Memoirs of the American Mathematical Society (2016), Vol. 239.
- [13] P. Cannarsa, P. Martinez, J. Vancostenoble, *The cost of controlling weakly degenerate parabolic equations by boundary controls*, Math. Control Relat. Fields (2017), Vol 7, No 2, p. 171-211.
- [14] P. Cannarsa, P. Martinez, J. Vancostenoble, *Precise estimates for biorthogonal families under asymptotic gap conditions*, submitted, and arXiv:1706.02435
- [15] P. Cannarsa, D. Rocchetti, J. Vancostenoble, *Generation of analytic semi-groups in L^2 for a class of second order degenerate elliptic operators*, Control Control and Cybernetics vol. 37 (2008) No. 4, p. 831-878.
- [16] P. Cannarsa, J. Tort, M. Yamamoto, *Unique continuation and approximate controllability for a degenerate parabolic equation*, Appl. Anal. 91, No. 8, 1409-1425 (2012).
- [17] J.M. Coron, S. Guerrero, *Singular optimal control: A linear 1-D parabolic-hyperbolic example*, Asymp. Anal. 44, No 3-4, (2005), p. 237-257.
- [18] S. Ervedoza, E. Zuazua, *Sharp observability estimates for heat equations*, Arch. Ration. Mech. Anal. 202, No 3, 975-1017.
- [19] S. Ervedoza, E. Zuazua, *Observability of heat processes by transmutation without geometric restrictions*, Math. Control Relat. Fields 1, No 2, 177-187 (2011).
- [20] L. Escauriaza, G. Seregin, V. Šverák, *Backward uniqueness for the heat operator in half-space*, St. Petersburg Math. J. 15 (2004), no. 1, 139–148.
- [21] W. N. Everitt, A. Zettl, *On a class of integral inequalities*, J. London Math. Soc. (2) 17 (1978) 291-303.
- [22] W.N. Everitt, *A catalogue of Sturm-Liouville differential equations*, Sturm-Liouville Theory, 271-331, Birkhäuser, Basel (2005).
- [23] H. O. Fattorini, D. L. Russel, *Exact Controllability Theorems for Linear Parabolic Equations in One Space Dimension*, Arch. Rat. Mech. Anal. 4, 272-292 (1971).
- [24] H. O. Fattorini, D. L. Russel, *Uniform bounds on biorthogonal functions for real exponentials with an application to the control theory of parabolic equations*, Quart. Appl. Math. 32 (1974/75), 45-69.
- [25] H. O. Fattorini, *Boundary control of temperature distributions in a parallelepipedon*, SIAM J. Control, Vol 13, No 1 (1975).
- [26] E. Fernandez-Cara, E. Zuazua, *The cost of approximate controllability for heat equations: the linear case*, Adv. Differential equations 5 (2000), No 4-6, 465-514.
- [27] A. V. Fursikov, O. Yu. Imanuvilov, *Controllability of evolution equations*. Lecture Notes Ser. 34, Seoul National University, Seoul, Korea, 1996.
- [28] O. Glass, *A complex-analytic approach to the problem of uniform controllability of transport equation in the vanishing viscosity limit*, J. Funct. Anal. 258 (2010), No 3, p. 852-868.
- [29] O. Glass, S. Guerrero, *Uniform controllability of a transport equation in zero diffusion-dispersion limit*, Mathematical Models and Methods in Applied Sciences, Vol 19, No. 9 (2009) 1567-1601.
- [30] S. Guerrero, G. Lebeau, *Singular optimal control for a transport-diffusion equation*, Communications in Partial Differential Equations 32 (2007), no. 10-12, 18, 1813–1836.
- [31] M. Gueye, *Exact boundary controllability of 1-D parabolic and hyperbolic degenerate equations*, SIAM J. Control Optim. 52 (2014), No 4, p. 2037-2054.
- [32] E.N. Güichal, *A lower bound of the norm of the control operator for the heat equation*, Journal of Mathematical Analysis and Applications 110 (1985), p. 519-527.
- [33] S. Hansen, *Bounds on functions biorthogonal to sets of complex exponentials; control of damped elastic systems*, Journal of Math. Anal. and Appl., 158 (1991), 487-508.
- [34] A. Haraux, *On a completion problem in the theory of distributed control of wave equations*, Nonlinear partial differential equations and their applications. Collège de France Seminar, Vol. X (Paris, 1987-1988), Pitman Res. Notes Math. Ser., Longman Sci. Tech., Harlow, 220 (1991), 241-271.
- [35] E. Kamke, *Differentialgleichungen: Lösungsmethoden und Lösungen. Band 1: Gewöhnliche Differentialgleichungen*. 3rd edition, Chelsea Publishing Company, New York, 1948.
- [36] V. Komornik, P. Loret, *Fourier Series in Control Theory*, Springer, Berlin, 2005.

- [37] I. Krasikov, *On the Bessel function $J_\nu(x)$ in the transition zone*, LMS Journal of Computation and Mathematics 17(01), January 2014.
- [38] J. Lagnese, *Control of wave processes with distributed controls supported on a subregion*, SIAM J. Control Optim. 21 (1983), no. 1, 68-85.
- [39] L.J. Landau, *Bessel functions: monotonicity and bounds*, Journal of the London Mathematical Society 61 (2000) 197-215.
- [40] G. Lebeau, L. Robbiano, *Contrôle exact de l'équation de la chaleur*, Comm. Partial Differential Equations 20 (1995), p. 335-356.
- [41] N.N. Lebedev, *Special Functions and their Applications*, Dover Publications, New York, 1972
- [42] P. Lissy, *On the cost of fast controls for some families of dispersive or parabolic equations in one space dimension*, SIAM J. Control Optim. 52 (2014), no. 4, 26512676.
- [43] P. Lissy, *Explicit lower bounds for the cost of fast controls for some 1-D parabolic or dispersive equations, and a new lower bound concerning the uniform controllability of the 1-D transport-diffusion equation*, J. Differential Equations 259 (2015), no. 10, 53315352.
- [44] L. Lorch, M.E. Muldoon, *Monotonic sequences related to zeros of Bessel functions*, Numer. Algor (2008) 49, p. 221-233.
- [45] W.A.J. Luxemburg, J. Korevaar, *Entire functions and Müntz-Szász type approximation*, Transactions of the American Math. Society 157 (1971), p. 23-37.
- [46] P. Martinez, J. Vancostenoble, *Carleman estimates for one-dimensional degenerate heat equations*. J. Evol. Eq 6 (2006), no. 2, 325-362.
- [47] L. Miller, *Geometric bounds on the growth rate of null-controllability cost for the heat equation in small time*, J. Differential Equations 204 (2004), No 1, p. 202-226.
- [48] L. Miller, *How violent are fast controls for Schrödinger and plate vibrations?*, Arch. Ration. Mech. Anal. 172 (2004) 429456.
- [49] L. Miller, *Controllability cost of conservative systems: Resolvent condition and transmutation*, J. Funct. Anal. 218 (2005) 425444.
- [50] L. Miller, *The control transmutation method and the cost of fast controls*, SIAM J. Control Optim. 45 (2006) 762772.
- [51] F. W. Olver, *Asymptotics and special functions*, New York, Academic press, 1974.
- [52] Y. Privat, E. Trélat and E. Zuazua, *Optimal observation of the one-dimensional wave equation*, J. Fourier Anal. Appl., 19(2013), 514-544.
- [53] Y. Privat, E. Trélat and E. Zuazua, *Optimal Shape and Location of Sensors for Parabolic Equations with Random Initial Data*, Arch. Rational Mech. Anal. 216 (2015) p. 921-981.
- [54] Y. Privat, E. Trélat and E. Zuazua, *Actuator design for parabolic distributed parameter systems with the moment method*, SIAM J. Control Optim. Vol. 55 (No 2), p. 1128-1152.
- [55] C. K. Qu, R. Wong, *"Best possible" upper and lower bounds for the zeros of the Bessel function $J_\nu(x)$* , Trans. Amer. Math. Soc. 351 (1999), no. 7, 2833-2859.
- [56] R.M. Redheffer, *Elementary remarks on completeness*, Duke Math. Journal 35 (1968), p. 103-116.
- [57] L. Schwartz, *Étude des sommes d'exponentielles*, deuxième édition. Paris, Hermann 1959.
- [58] T.I. Seidman, *Two results on exact boundary control of parabolic equations*, Appl. Math. Optim. 11 (1984), No 2, p. 145-152.
- [59] T.I. Seidman, S.A. Avdonin, S.A. Ivanov, *The "window problem" for series of complex exponentials*, J. Fourier Anal. Appl. 6 (2000), No 3, p. 233-254.
- [60] T.I. Seidman, *How violent are fast controls?*, Math. Control Signals Systems 1 (1988) 8995.
- [61] G. Tenenbaum, M. Tucsnak, *New blow-up rates for fast controls of Schrödinger and heat equations*, J. Differential Equations 243 (2007), p. 70-100.
- [62] G. Tenenbaum, M. Tucsnak, *On the null controllability of diffusion equations*, ESAIM: COCV 17 (2011) 10881100.
- [63] E.C. Titchmarsh, *Eigenfunction expansions associated with second-order differential equations, part II*, Oxford University Press, 1958.
- [64] G. N. Watson, *A treatise on the theory of Bessel functions*, second edition, Cambridge University Press, Cambridge, England, 1944.
- [65] R.M. Young, *An Introduction to Nonharmonic Fourier Series*, Academic Press, 1980.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI ROMA "TOR VERGATA", VIA DELLA RICERCA SCIENTIFICA, 00133 ROMA, ITALY

E-mail address: `cannarsa@mat.uniroma2.it`

INSTITUT DE MATHÉMATIQUES DE TOULOUSE; UMR 5219, UNIVERSITÉ DE TOULOUSE; CNRS, UPS IMT F-31062 TOULOUSE CEDEX 9, FRANCE

E-mail address: `patrick.martinez@math.univ-toulouse.fr`

INSTITUT DE MATHÉMATIQUES DE TOULOUSE; UMR 5219, UNIVERSITÉ DE TOULOUSE; CNRS, UPS IMT F-31062 TOULOUSE CEDEX 9, FRANCE

E-mail address: `Judith.Vancostenoble@math.univ-toulouse.fr`