



Immersion of transitive tournaments in digraphs with large minimum outdegree

William Lochet

► To cite this version:

William Lochet. Immersion of transitive tournaments in digraphs with large minimum outdegree. Journal of Combinatorial Theory, Series B, 2019, 134, pp.350-353. 10.1016/j.jctb.2018.05.004 . hal-01835124

HAL Id: hal-01835124

<https://hal.science/hal-01835124>

Submitted on 16 Jul 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Immersion of transitive tournaments in digraphs with large minimum outdegree*

William Lochet^{a,b}

^a Université Côte d’Azur, CNRS, I3S, UMR 7271,
Sophia Antipolis, France

^bLaboratoire d’Informatique du Parallélisme
UMR 5668 ENS Lyon - CNRS - UCBL - INRIA
Université de Lyon, France

Abstract

We prove the existence of a function $h(k)$ such that every simple digraph with minimum outdegree greater than $h(k)$ contains an immersion of the transitive tournament on k vertices. This solves a conjecture of Devos, McDonald, Mohar and Scheide.

In this note, all digraphs are without loops. Let D be a digraph. We denote by $V(D)$ its vertex set and $A(D)$ its arc set. A digraph D is *simple* if there is at most one arc from x to y for any $x, y \in V(D)$. Note that arcs in opposite directions are allowed. The *multiplicity* of a digraph D is the maximum number of parallel arcs in the same direction in D . For an arc $a = (u, v)$ of a digraph D , we say that u is the *tail* of a and v its *head*. The *outdegree* (resp. *indegree*) of a vertex v , denoted by $d^+(v)$ (resp. $d^-(v)$), is equal to the number of arcs a of D such that v is the tail (resp. head) of a . The *outneighbourhood* (resp. *inneighbourhood*) of a vertex v , denoted by $N^+(v)$ (resp. $N^-(v)$), is the set of vertices y such that (v, y) (resp. (y, v)) is an arc of D . A *directed path* P in a digraph D is a set of vertices x_1, \dots, x_k such that $(x_i, x_{i+1}) \in A(D)$ for all $1 \leq i \leq k-1$. A *directed cycle* C in a digraph D is a set of vertices x_1, \dots, x_k such that $(x_i, x_{i+1}) \in A(D)$ for all $1 \leq i \leq k-1$ and $(x_k, x_1) \in A(D)$. A digraph D is a *tournament* if exactly one of (u, v) and (v, u) is an arc of D for all distinct $u, v \in V(D)$. The *transitive tournament* on k vertices, denoted by TT_k , is the unique tournament on k vertices without any directed cycle. The *complete digraph* on k vertices is the simple digraph on k vertices with every possible arc.

We say that a digraph D contains an *immersion* of a digraph H if there exists a mapping such that vertices of H are mapped to distinct vertices of D , and the arcs of H are mapped to directed paths joining the corresponding pairs of vertices of D , in such a way that these paths are pairwise arc-disjoint. If the directed paths are pairwise internally vertex-disjoint, we say that D contains a *subdivision* of H .

*This work was supported by ANR under contract STINT ANR-13-BS02-0007.

Understanding the necessary conditions for undirected graphs to contain a subdivision of a clique is a very natural and well-studied question. One of the most important examples is the following result by Mader [6]:

Theorem 1 ([6]). *For every $k \geq 1$, there exists an integer $f(k)$ such that every undirected graph with minimum degree greater than $f(k)$ contains a subdivision of K_k .*

Bollobás and Thomason [1] as well as Komlós and Szemerédi [4] proved that $f(k) = O(k^2)$. In the case of digraphs, there exist examples of digraphs with large out- and indegree without a subdivision of the complete digraph on three vertices, as shown by Thomassen [7] (see also [3] for a simpler construction). However Mader [5] conjectured that an analogue should hold for transitive tournaments TT_k in digraphs with large minimum outdegree.

Conjecture 2 ([5]). *For every $k \geq 1$, there exists an integer $g(k)$ such that every simple digraph with minimum outdegree greater than $g(k)$ contains a subdivision of TT_k .*

The question turned out to be much more difficult than the undirected case, as the existence of $g(5)$ remains unknown. Weakening the statement, Devos, McDonald, Mohar and Scheide [3] made the following conjecture replacing subdivision with immersion and proved it for the case of Eulerian digraphs.

Conjecture 3 ([3]). *For every $k \geq 1$, there exists an integer $h(k)$ such that every simple digraph with minimum outdegree greater than $h(k)$ contains an immersion of TT_k .*

As with subdivisions, Devos et al. showed in [3] the existence of digraphs with large out- and indegree without an immersion of the complete digraph on three vertices. Finding the right value for $h(k)$ in the case of undirected graphs is an interesting question on its own (see [2] for more details).

The goal of this note is to present a proof of this conjecture. Let $F(k, l)$ be the digraph consisting of k vertices x_1, \dots, x_k and l arcs from x_i to x_{i+1} for every $1 \leq i \leq k-1$. It is clear that $F(k, \binom{k}{2})$ contains an immersion of TT_k , so the following theorem implies Conjecture 3.

Theorem 4. *For every $k \geq 1$ and l , there exists a function $f(k, l)$ such that every digraph with minimum outdegree greater than $f(k, l)$ and multiplicity at most kl contains an immersion of $F(k, l)$.*

Proof. We prove the result for $f(k, l) = 2k^3l^2$ and $l \geq 2$. We proceed by induction on k . For $k = 1$ this is trivial because $F(1, l)$ is one vertex. Suppose now that the result holds for k and assume for a contradiction that it does not hold for $k + 1$. Let D be the digraph with the smallest number of arcs and vertices such that D has multiplicity at most $(k + 1)l$, all but at most $c_1 = k + (k + 1)l$ vertices have outdegree at least $f(k + 1, l)$ and without an immersion of $F(k + 1, l)$. By minimality of D , every vertex has outdegree exactly $f(k + 1, l)$, except c_1 of them with outdegree 0. Call T the set of vertices of outdegree 0. Suppose we want to remove arcs from D such that the multiplicity of the remaining digraph is at most kl , while keeping the minimum outdegree as large as possible. For a vertex v , the worst case is when, for every vertex $y \in N^+(v)$, the multiplicity of (v, y) is equal to $(k + 1)l$. In this case we have to remove at most l arcs for each of the $\frac{f(k+1, l)}{(k+1)l}$ vertices of $N^+(v)$. Therefore, removing T and some of the parallel arcs, we obtain a digraph of outdegree greater than $d' = f(k+1, l) - c_1(k+1)l - \frac{f(k+1, l)}{k+1}$ with multiplicity kl . Because $f(k+1, l) - f(k, l) = 2(3k^2 + 3k + 1)l^2$ and $c_1(k + 1)l + \frac{f(k+1, l)}{(k+1)} = k(k + 1)l + 3(k + 1)^2l^2$, we get that $d' \geq f(k, l)$ and by induction there

exists an immersion of $F(k, l)$ in $D - T$. Call $X = \{x_1, \dots, x_k\}$ the set of vertices of the immersion and, numbering the paths arbitrarily, $P_{i,j}$ the j th directed path of this immersion from x_i to x_{i+1} . We can assume this immersion is of minimum size, so that every vertex in $P_{i,j}$ has exactly one outgoing arc in $P_{i,j}$. Let D' be the digraph obtained from D by removing all the arcs of the $P_{i,j}$ and the vertices x_1, \dots, x_{k-1} . By the previous remark, the outdegree of each vertex in D' is either 0 if this vertex belongs to T or at least $f(k+1, l) - (k-1)l - (k-1)(k+1)l$.

For every vertex $y \in D' - x_k$, there do not exist l arc-disjoint directed paths from x_k to y in D' , for otherwise there would be an immersion of $F(k+1, l)$ in D . Hence, by Menger's Theorem there exists a set E_y of less than l arcs such that there is no directed path from x_k to y in $D' \setminus E_y$. Define C_y for every vertex $y \in D' - x_k$ as the set of vertices which can reach y in $D' \setminus E_y$. Now take Y a minimal set such that $\cup_{y \in Y} C_y$ covers $D' - x_k$. We claim that Y consists of at least $c_2 \geq \frac{f(k+1, l) - (k-1)l - (k-1)(k+1)l}{l} \geq 2c_1$ elements, as $\cup_{y \in Y} E_y$ must contain all the arcs of D' with x_k as tail.

For each $y \in Y$, define S_y as the set of vertices which belong to C_y and no other $C_{y'}$ for $y' \in Y$. Since Y is minimal, every S_y is non-empty. Note that for $u \in S_y$, if there exists $y' \in Y \setminus y$ and $v \in C_{y'}$ such that $uv \in A(D)$, then $uv \in E_{y'}$. Note that $T \subset Y$ as vertices in T have outdegree 0 and if $y \in Y \setminus T$ then S_y consists only of vertices of outdegree $f(k+1, l)$ in D .

Let R be the digraph with vertex set Y and arcs from y to y' if there is an arc from S_y to $C_{y'}$. As noted before, $d_R^-(y) \leq |E_y| \leq l$. The average outdegree of the vertices of $Y \setminus T$ in R is then at most $\frac{c_1 l + (c_2 - c_1)l}{c_2 - c_1} \leq 2l$. Let y be a vertex of $R \setminus T$ with outdegree at most this average. Let H be the digraph induced on D' by the vertices in S_y to which we add X , all the arcs that existed in D (with multiplicity) from vertices of S_y to vertices of X and the following arcs: For each $P_{i,j}$, let $z_1, z_2, \dots, z_l = P_{i,j} \cap S_y$, where z_i appears before z_{i+1} on $P_{i,j}$ and add all the arcs (z_i, z_{i+1}) to H . Note that, if (x, y) is an arc of D' , then by minimality of the immersion of $F(k, l)$, every time x appears before y on some $P_{i,j}$, then $P_{i,j}$ uses one of the arcs (x, y) . Thus for each pair of vertices x and y in H , either $(x, y) \in A(D)$ and the number of (x, y) arcs in H is equal to the one in D , or $(x, y) \notin A(D)$ and the number of (x, y) arcs in H is bounded by $(k-1)l$. This implies that H has multiplicity at most $(k+1)l$.

Claim 4.1. *H is a digraph with multiplicity at most $(k+1)l$, such that all but at most c_1 vertices have outdegree greater than $f(k+1, l)$ and H does not contain an immersion of $F(k+1, l)$.*

Proof of the claim. Suppose H contains an immersion of $F(k+1, l)$, then by replacing the new arcs by the corresponding directed paths along the $P_{i,j}$ we get an immersion of $F(k+1, l)$ in D . Moreover, we claim that the number of vertices in H with outdegree smaller than $f(k+1, l)$ is at most $k + 2l + (k-1)l = c_1$. Indeed, the vertices of H that can have outdegree smaller in H than in D are the x_i , or the vertices with outgoing arcs in $E_{y'}$ for some $y' \in Y \setminus y$, or the vertices along the $P_{i,j}$. But with the additions of the new arcs, we know that there is at most one vertex per path $P_{i,j}$ that loses some outdegree in H . \diamond

However, since H is strictly smaller than D , we reach a contradiction. \square

Acknowledgements

The author wishes to thank Frédéric Havet and Stéphan Thomassé for their useful comments on the manuscript.

References

- [1] B. Bollobás and A. Thomason, Proof of a conjecture of Mader, Erdős and Hajnal on topological complete subgraphs, *European Journal of Combinatorics* **19** (1998), 883–887.
- [2] M. Devos, Z. Dvořák, J. Fox, J. McDonald, B. Mohar, and D. Scheide, Minimum degree condition forcing complete graph immersion, *Combinatorica* **34** (2014), 279–298.
- [3] M. DeVos, J. McDonald, B. Mohar, and D. Scheide. Immersing complete digraphs. *European Journal of Combinatorics*, **33** (2012), 1294–1302.
- [4] J. Komlós and E. Szemerédi, Topological Cliques in Graphs II, *Combinatorics, Probability and Computing* **5** (1996), 70–90.
- [5] W. Mader, Degree and Local Connectivity in Digraphs, *Combinatorica* **5** (1985), 161–165
- [6] W. Mader, Homomorphieeigenschaften und mittlere Kantendichte von Graphen. *Math. Annalen* **174** (1967), 265–268.
- [7] C. Thomassen, Even Cycles in Directed Graphs, *European Journal of Combinatorics* **6** (1985), 85–89.