



**HAL**  
open science

## Stress responses, outer membrane permeability control and antimicrobial resistance in Enterobacteriaceae 2 3

Sushovan Dam, Jean-Marie Pages, Muriel Masi

### ► To cite this version:

Sushovan Dam, Jean-Marie Pages, Muriel Masi. Stress responses, outer membrane permeability control and antimicrobial resistance in Enterobacteriaceae 2 3. Microbiology, 2018, 10.1099/mic.0.000613 . hal-01840466

HAL Id: hal-01840466

<https://hal.science/hal-01840466v1>

Submitted on 16 Jul 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1 **Stress responses, outer membrane permeability control and antimicrobial resistance in**  
2 **Enterobacteriaceae**

3

4 Sushovan Dam, Jean-Marie Pagès\* and Muriel Masi

5

6

7

8 UMR\_MD-1, Aix-Marseille Univ. & IRBA, 27 Boulevard Jean Moulin, 13005 Marseille,  
9 France.

10

11 \*Corresponding author:

12 [jean-marie.pages@univ-amu.fr](mailto:jean-marie.pages@univ-amu.fr)

13 +33 (0)4 91 32 46 97

14

15 Key words: Enterobacteriaceae, envelope stress responses, outer membrane permeability,  
16 porins, drug translocation, multidrug resistance.

17

18 Category: Regulation

19

20 Word count (from introduction to conclusion):

21

22 Abbreviations: outer membrane (OM), inner membrane (IMI), peptidoglycan (PG),  
23 lipopolysaccharide (LPS), antimicrobial resistance (AMR), multidrug resistance (MDR),  
24 envelope stress response (ESR), two-component system (TCS), small regulatory RNA  
25 (sRNA).

26

27 **Abstract**

28 Bacteria have evolved several strategies to survive a myriad of harmful conditions in the  
29 environment and in hosts. In Gram-negative bacteria, responses to nutrient limitation,  
30 oxidative or nitrosative stress, envelope stress, exposure to antimicrobials and other growth-  
31 limiting stresses have been linked to the development of antimicrobial resistance. This results  
32 from the activation of protective changes to cell physiology (decreased outer membrane  
33 permeability), resistance transporters (drug efflux pumps), resistant lifestyles (biofilms,  
34 persistence) and/or resistance mutations (target mutations, production of antibiotic  
35 modification/degradation enzymes). In targeting and interfering with essential physiological  
36 mechanisms, antimicrobials themselves are considered as stresses to which protective  
37 responses have also evolved. In this review, we focus on envelope stress responses that affect  
38 the expression of outer membrane porins and their impact on antimicrobial resistance. We  
39 also discuss evidences that indicate the role of antimicrobials as signaling molecules in  
40 activating envelope stress responses.

## 41 **Introduction**

42 Antimicrobial resistance (AMR) is broadly recognized as a growing threat for human health  
43 [1, 2, 3]. As such, increasing antibiotic treatment failures due to multidrug resistant (MDR)  
44 bacteria have stirred the urgent need to better understand the underlying molecular  
45 mechanisms and promote innovation with the development of new antibiotics and alternative  
46 therapies [4, 5]. The efficacy of antibacterial compounds depends on their capacity to reach  
47 inhibitory concentrations at the vicinity of their target. This is particularly challenging for  
48 drugs directed against Gram-negative bacteria, which exhibit a complex envelope comprising  
49 two membranes and transmembrane efflux pumps [6, 7]. The Gram-negative envelope  
50 comprises an inner membrane (IM), which is a symmetric phospholipid bilayer; a thin  
51 peptidoglycan (PG) layer ensuring the cell shape; and an outer membrane (OM) that is an  
52 asymmetric bilayer, composed of an inner leaflet of phospholipids and an outer leaflet of  
53 lipopolysaccharide (LPS) [8]. The OM is a barrier to both hydrophobic and hydrophilic  
54 compounds, including necessary nutrients, metabolic substrates and antimicrobials, but  
55 access is provided by the water filled  $\beta$ -barrel channels called porins [6, 9, 10, 11, 12]. In  
56 *Escherichia coli*, the channels of the general porins OmpF and OmpC are size restricted and  
57 show a preference for passage of hydrophilic charged compounds, including antibiotics such  
58 as  $\beta$ -lactams and fluoroquinolones. These porins are conserved throughout the phylum of  $\gamma$ -  
59 proteobacteria [13]. Additionally, tripartite RND (Resistance-Nodulation-cell Division)  
60 efflux pumps, such as AcrAB-TolC in *E. coli*, play a major role in removing antibiotics from  
61 the periplasm [7, 12]. Not surprisingly, MDR clinical isolates of *Enterobacteriaceae*  
62 generally exhibit porin loss and/or increased efflux, which act in synergy to reduce the  
63 intracellular accumulation of antibiotics below the threshold that would be efficient for  
64 activity [10].

65 Given the importance of the OM in controlling the uptake of beneficial as well as toxic  
66 compounds, one can expect that the expression of porins depends on environmental stresses  
67 and is well-coordinated at the transcriptional and post-transcriptional levels [10, 14-17]. In  
68 this review, we will address the porin-mediated influx of antibiotics and give a perspective on  
69 the factors, including major regulatory pathways and antibiotic stresses, which control porin  
70 expression in *E. coli* and closely relative *Enterobacteriaceae*. Additionally, we will discuss  
71 the recent clinical data that illustrate the bacterial strategies using porins modifications to  
72 limit antibiotic entry.

73

## 74 **Antibiotic stresses**

75 Bacteria are present in a wide range of environments in which they are exposed to diverse  
76 toxic compounds or growth-limiting conditions. These include antibiotics used in the medical  
77 environment and agricultural settings. The last few decades have been marked by the constant  
78 increase of (multi)drug resistant clinical isolates to which we responded by increasing  
79 antibiotic dosing. Therefore, antibiotics are present almost everywhere at different  
80 concentrations [18]. Although MDR still emerges from bacterial exposure to antibiotic  
81 concentrations that are higher than the minimal inhibitory concentrations (MIC, defined as  
82 the lowest concentration of a drug that inhibit bacterial growth in defined laboratory  
83 conditions), the effects of subinhibitory concentrations on bacterial physiology and AMR was  
84 mostly disregarded. Importantly, studies in this field have shown that low antibiotic  
85 concentrations affect bacteria at least at four different levels: as selectors of resistance (by  
86 enriching resistant bacteria within populations and selecting for *de novo* resistance mutations)  
87 [19]; (ii) as contributors of genetic and phenotypic heterogeneity [20]; (iii) as intercellular  
88 signals [21]; (iv) as inducers of persistence [22]. In this regard, Viveiros and colleagues have  
89 demonstrated the induction of high-level resistance to tetracycline (TET) in susceptible *E.*  
90 *coli* K12 obtained by gradual, step-wise increase exposure to subinhibitory concentrations of  
91 the antibiotic [23]. Increased expression of the AcrAB efflux pump was found responsible for  
92 resistance to TET, which could also be reversed by the use of the efflux pump inhibitor  
93 phenylalanine-arginine- $\beta$ -naphthylamide (PA $\beta$ N). Interestingly, the TET-resistant strain also  
94 exhibited MDR due to repression of OmpF and OmpC expression [24]. Important questions  
95 arise from this and other related studies. First is whether the target for signaling resistance is  
96 the same as the target that is inhibited by the antibiotic. In case the antibiotic itself but not a  
97 secondary metabolite is the signaling molecule, this could be determined by examining  
98 whether the response is alleviated by a target mutation that prevents drug binding. Second is  
99 whether and how the antibiotic (or a secondary metabolite) interferes with the ESRs  
100 described above. Here, comparative transcriptomics between susceptible and resistant strains  
101 would be a valuable tool to answer this question.

102

## 103 **Global regulators**

104 In *Enterobacteriaceae*, the development of MDR is under positive regulation by global  
105 transcriptional activators that include members of Ara/XylS superfamily such as MarA,  
106 RamA (absent in *E. coli*) and Rob as well as the oxidative stress regulon SoxSR [10, 25-29].  
107 Mutations in the corresponding genes are well-documented and induce the overproduction of

108 efflux pumps with concomitant repression of porin expression both directly and indirectly  
109 [10]. These mechanisms are reviewed in details in Davin *et al.* [10]. Negative regulation by  
110 repressors of porins also plays a major role. OmpX is a small OM channel [30], of which  
111 overexpression is associated with a decreased expression of Omp36 (the OmpC ortholog of  
112 *Enterobacter aerogenes*) and a decreased susceptibility to  $\beta$ -lactams [31, 32]. Studies have  
113 indicated that expression OmpX itself is controlled by a number of environmental factors  
114 including salicylate via MarA and paraquat via SoxS [33] A very rapid MarA-dependent  
115 response pathway for upregulation of *ompX* has been shown to occur within 60-120 min upon  
116 cell exposure to salicylate [32]. This work by Dupont *et al.* identified dramatic decrease in  
117 OmpF levels, as a first line of defense, with simultaneous development of resistance to  $\beta$ -  
118 lactams and fluoroquinolones by altering OM permeability.

119

### 120 **Envelope stress responses**

121 All living organisms have stress responses that allow them to sense and respond to  
122 environmental damaging conditions by remodeling gene expression. As such, Gram-negative  
123 bacteria possess stress responses that are uniquely targeted to the cell envelope, including  
124 membranes and cell wall. These envelope stress responses (ESRs) are the EnvZ/OmpR,  
125 CpxAR (Cpx), BaeRS, and Rcs phosphorelays, the stress responsive alternative sigma factor  
126  $\sigma^E$ , and the phage shock response [34-37] in *E. coli* and closely related *Enterobacteriaceae*.  
127 Each of these ESRs is activated following the perturbation of particular components of the  
128 envelope or exposure to particular environmental stresses. Although ESRs are important for  
129 reacting to damaging conditions, stress proteins also play important roles in the maintenance  
130 of basic cellular physiology [38, 39]. This is particularly true for the  $\sigma^E$ -dependent stress  
131 response in *E. coli*, as the *rpoE* gene, which encodes  $\sigma^E$ , is essential for viability [40]. Here,  
132 we will essentially focus on ESRs that impact on AMR by regulating porin expression  
133 together with many other targets (regulons) — namely EnvZ/OmpR, Cpx and  $\sigma^E$  (see below  
134 and key figure). Additionally, with the recent highlights and advances in RNA-based  
135 techniques [41], the repertoire of small regulatory RNAs (sRNAs) has vastly increased so as  
136 to and their impact on the OM is continuously emerging [15, 17]. sRNAs alter gene  
137 expression, allowing fast adjustment to different growth conditions [42]. Noteworthy, ESRs  
138 are often interconnected, regulate and are regulated by sRNAs in order control target genes  
139 both at the transcriptional and post-transcriptional levels [15-17, 43, 44] (see below and key  
140 figure).

141 Osmolarity was one of the earliest stresses described to influence OmpF and OmpC  
142 expression via the EnvZ/OmpR two-component system (TCS) [45, 46]. EnvZ is a membrane-  
143 bound sensor kinase, and OmpR is a cytosolic response regulator, which binds to the  
144 promoter region of the porin genes. Upon activation, EnvZ autophosphorylates and the high  
145 energy phosphoryl group from EnvZ is subsequently transferred to a conserved Asp residue  
146 on OmpR. Phosphorylated OmpR (OmpR~P) serves as a transcription factor that  
147 differentially modulates the expression of the *ompF* and *ompC* porin genes [45]. At low  
148 osmolarity, high levels of OmpR~P activates *ompF* transcription, whereas at high osmolarity,  
149 low levels of OmpR~P represses *ompF* transcription and activates *ompC* transcription [47].  
150 This differential production of OmpF and OmpC is consistent with that in high osmolarity  
151 environments, such as in the hosts where nutrients are abundant the least permeable pore  
152 channel OmpC is predominant, thus limiting the uptake of toxic bile salts; whereas in low  
153 osmolarity environments where nutrients are scarce, the most permeable pore channel OmpF  
154 is expressed [6]. OmpF and OmpC transcriptional regulation by EnvZ/OmpR is also triggered  
155 by local anesthetics, pH, and nutrient limitation [46].

156 Accumulation of misfolded OM proteins in the periplasm, presumably reflecting problems in  
157 protein assembly or transport across the IM, can be detected by regulatory sensors that  
158 activate either the Cpx TCS or the alternative sigma factor  $\sigma^E$ .  $\sigma^E$  and Cpx are the two major  
159 regulation pathways that control the envelope integrity with overlapping regulon members  
160 [48-51] but respond to different inducing cues [35]. It is possible that these poorly defined  
161 signals (see below) act by causing accumulation of misfolded proteins. However, misfolded  
162 proteins are not the inducing signal *per se*, as some induce  $\sigma^E$  but not Cpx and *vice versa*.  
163 Recent studies rather suggest that Cpx responds to IM perturbations, while  $\sigma^E$  is activated by  
164 signals at the OM. The Cpx system comprises the CpxA sensor kinase and response regulator  
165 CpxR. Envelope stresses including alkaline pH, periplasmic protein misfolding, IM  
166 abnormalities such as misfolded transporters or accumulation of the lipid II precursor, induce  
167 the dissociation of the accessory protein CpxP from CpxA, trigger CpxA-mediated  
168 phosphorylation of CpxR, and altered expression of protein foldases and proteases,  
169 respiratory complexes, IM transporters, and cell wall biogenesis enzymes [37, 48, 49], all of  
170 which affect resistance to a number of antibiotics, particularly aminoglycosides and  $\beta$ -  
171 lactams [37, 49, 52-54]. The Cpx-mediated regulation of porins occurs at several levels. At  
172 the transcriptional level, CpxR~P has been shown to bind directly the *ompF* and *ompC*  
173 promoters [55]. More recently, it has been found that the small IM protein MzrA connects

174 Cpx and EnvZ/OmpR [56]. In this pathway and upon the activation of Cpx, MzrA interacts  
175 directly with EnvZ, which in turn, stabilizes OmpR~P [57]. In sensing different signals, the  
176 interconnection between Cpx and EnvZ/OmpR allows cells to adapt to diverse environmental  
177 stresses. Finally, although Cpx contributes to AMR by regulating a number of genes [37, 49,  
178 52-54], its precise role and that of other TCSs in the development of MDR in clinical isolates  
179 are still poorly documented [58]. On the other hand, the stress responsive sigma factor  $\sigma^E$  is  
180 induced by stresses that disturb the OM and its regulon members comprise genes that  
181 facilitate the biogenesis of OM components, including proteins, lipoproteins and LPS [59-  
182 67]. In the absence of inducing signals,  $\sigma^E$  is held at the cytoplasmic side of the IM by the  
183 anti-sigma factor RseA. At the periplasmic side of the IM, RseB binds to RseA, thus  
184 enhancing the inhibition of  $\sigma^E$ . Upon activation,  $\sigma^E$  is released from RseA by a proteolytic  
185 cascade that starts with the sequential degradation of the periplasmic and transmembrane  
186 domains of RseA by DegS and RseP, respectively, followed by the degradation of the  
187 cytoplasmic domain of RseA by ClpXP [68]. Interestingly, proteolysis of RseA is triggered  
188 by the binding of a conserved peptide found at the C-terminus of OM proteins, which is  
189 normally buried in folded porin trimers, to DegS in conjunction with the release of RseB  
190 from RseA upon binding of LPS intermediates [69, 70]. Of note, the  $\sigma^E$ -dependent repression  
191 of porin synthesis only occurs at the post-transcriptional level, wherein base-pairing sRNAs  
192 inhibits translation of *omp* mRNAs (see below) in order to maintain the envelope homeostasis  
193 under stress conditions, as porins are major abundant proteins under normal growth  
194 conditions [6].

195 The post-transcriptional repression of OmpF by the sRNA MicF has been discovered in 1984  
196 [71-73]. This 93-nucleotide (nt) RNA is transcribed in the opposite direction to the *ompC*  
197 gene and acts by direct base-pairing to a region that encompasses the ribosome binding site  
198 (RBS) and the start codon of the *ompF* mRNA, thus preventing translation initiation [74].  
199 The expression of the MicF sRNA itself is subjected to multiple signals and regulatory  
200 pathways [75]. Positive regulation includes EnvZ/OmpR in high osmolarity conditions [76],  
201 SoxS in response to oxidative stress [77] and MarA in response to antibiotic stress [25]. The  
202 109-nt MicC sRNA has been identified more recently and shown to repress OmpC by direct  
203 base-pairing to a 5' untranslated region of the *ompC* mRNA [78]. Interestingly, *micC* is  
204 transcribed in the opposite direction to the *ompN* gene that encodes a quiescent porin  
205 homologous to OmpF and OmpC [79]. We have recently shown that *ompN* and *micC* are  
206 subjected to dual regulation upon exposure to certain antimicrobials such as  $\beta$ -lactams in a



207  $\sigma^E$ -dependent manner [80]. This is consistent with that *ompN-micC* and *ompC-micF* share  
208 similar genetic organization and that *ompC* and *micF* are co-induced under specific  
209 conditions (i. e. high osmolarity via EnvZ/OmpR). The last decade has been marked by the  
210 identification and characterization of several sRNAs. These are differentially expressed and  
211 have been assigned to various important regulatory pathways in *E. coli* and *Salmonella*.  
212 Interestingly, many pathways regulate and are regulated by sRNAs [43, 44]. A prime  
213 example is EnvZ/OmpR, which activates the expression of MicF (that target *ompF*), OmrA  
214 and OmrB (that target *ompT* and mRNA of OM channels for iron-siderophore complexes)  
215 [81]; OmrA and OmrB, in turn, repress the translation of the *ompR* mRNA, creating a  
216 negative feedback loop [82]. Others examples include the well-conserved  $\sigma^E$ -regulated  
217 sRNAs RybB (that target *ompC* and *lamB* in *E. coli*; *ompN* and *ompW* in *Salmonella*), MicA  
218 (*ompA*), RseX (*ompC* and *ompA*), CyaR (*ompX*) and MicL (that represses translation of the  
219 major OM lipoprotein Lpp) [43, 66, 83-90] (key figure). Of note all these sRNAs are *trans*-  
220 acting, function by imperfect base pairing with multiple mRNA targets and require the help  
221 of the RNA chaperone Hfq [15-17].

222

### 223 **Porin alterations in clinical isolates**

224 Combined regulations contributed by different stressors leads to hampering of the drug  
225 accumulation inside cells under the threshold for bacterial death. In one such study in *K.*  
226 *pneumoniae*, preferential expression of OmpK37 was detected in porin-deficient strains [92].  
227 Amino acid sequencing showed that OmpK37 is highly homologous to quiescent porins  
228 OmpS2 from *Salmonella enterica* serovar Typhimurium and OmpN from *E. coli*. Liposome  
229 swelling assay with purified porins determined that OmpK37 also has a narrower pore, which  
230 was responsible for higher MICs of cefotaxime and ceftazidime antibiotics because of lower  
231 drug diffusion. A very recent study identified mutation in the *pho* regulon of an extensively  
232 drug resistant strain of *K. pneumoniae* demonstrating downregulation of *phoE* gene by  
233 mutations in *phoR* and *phoB*. Here the PhoE porin, which is normally involved in phosphate  
234 transport, promotes restoration of ceftazidime and carbapenem resistance [93]. This is an  
235 interesting example of a regulatory mutation that effects porin expression, and clinically  
236 favors AMR under antibiotic therapy.

237 A wide array of chemicals including disinfectants and antibiotics has been shown to modulate  
238 the OM permeability including expression of porins [94]. In addition, several studies have

239 described the effect of imipenem on porin loss or loss of function mutations in clinical  
240 isolates of *Enterobacteriaceae* [58, 95-100].

241 Porins are trimers of 16-stranded  $\beta$ -barrels, each monomer formed of a central channel  
242 constricted by loop 3 that folds inward, thereby restricting the size of the channel. The  
243 presence of acidic residues in loop 3 facing a cluster of basic residues on the opposite side of  
244 the pore creates a strong transversal electric field [6, 101, 102]. This so-called eyelet or  
245 constriction region determines the channel size and ion selectivity, with OmpF being more  
246 permeable than OmpC. This latter observation was first attributed to the OmpC pore being  
247 slightly more constricted in this porin compared to OmpF [101, 102]. Although the two  
248 porins share high sequence similarity, the pore interior is more negative in OmpC than in  
249 OmpF [102]. This can also account for the low permeability of OmpC for anionic  $\beta$ -lactams  
250 [103, 104]. Moreover, the replacement of all ten titratable residues that differ between OmpC  
251 and OmpF in the pore-lining region leads to the exchange of antibiotic permeation properties  
252 [105]. Together, these structural and functional data clearly demonstrate that the charge  
253 distribution at pore linings, but not pore size, is a critical parameter that physiologically  
254 distinguishes OmpC from OmpF.

255 Functional mutations in porin genes leading to reduced permeability are another strategy  
256 found in MDR bacteria. In two documented cases,  $\beta$ -lactam-resistant clinical isolates of *E.*  
257 *aerogenes* contained Omp36 (an OmpC homologue) that carried the mutation G112D in L3  
258 [106, 97]. The homologous mutation G119D in OmpF of *E. coli* narrows the size of the  
259 channel as the large side chain of Asp protrudes into the channel lumen and confers a drastic  
260 reduction in  $\beta$ -lactam susceptibility [107]. Consistently, the Omp36 G112D mutant of *E.*  
261 *aerogenes* was characterized by a 3-fold decrease in ion conductance and a significant  
262 decrease in cephalosporin sensitivity (*e. g.* MICs of cefotaxime, cefpirome, cefepime and  
263 ceftazidime were 7 to 9 fold higher in the clinical isolate as compared to that in a sensitive  
264 reference strain) and a cross resistance to carbapenems [106, 97]. Recent studies also found a  
265 series of OmpC mutants that were isolated from a patient with chronic *E. coli* infection and  
266 additive mutations that conferred increased resistance to a variety of antibiotics, including  
267 cefotaxime, ceftazidime, imipenem, meropenem and ciprofloxacin [108, 109]. Low *et al.*  
268 demonstrated that subtle changes in OmpC in clinical isolates of *E. coli* altered antibiotic  
269 permeability and thus cell viability [108]. Seven isolates collected over a two year clinical  
270 treatment exhibited increased levels of antibiotic resistance. These isolates exhibited the same  
271 two mutations (D18E and S274F) in the OmpC porin with increased levels of antibiotic

272 resistance, thus pointing towards the possible functional role of these mutations in antibiotic  
273 influx.

274 It is worthwhile to note that from our knowledge, porin mutations causing reduced  
275 permeability have only been described in OmpC-type porins in *E coli* and *E aerogenes*.  
276 Interestingly, this type of porin is expressed under high osmolarity, the same environment the  
277 bacteria encounters the hosts. This gives an essential outlook on the host induced  
278 modifications that possibly occurs in these pathogens during infection. Heeding to this sort of  
279 information can be highly beneficial for designing drugs with an improved diffusion across  
280 the bacterial outer membrane.

281

## 282 **Conclusion**

283 It is noteworthy that the sRNA-mediated stress response mechanism has multiple benefits for  
284 bacteria as compared to regulation by protein. Since sRNAs are produced during  
285 transcription, the later stages of translation and post translational modification processes in  
286 the cell is completely skipped proving to be time and energy efficient for the cell. Not to  
287 forget the energy saved in porin assembly and discarding of misfolded proteins, which in  
288 itself can induce another stress response mechanism.

289 Decreased porin expression has been observed as a rapid response to toxic molecules and  
290 antibiotics within less than 60 minutes. Many sRNAs act at the post-transcriptional level,  
291 which ensures a rapid response to stressful conditions. In addition, the versatility of sRNAs  
292 ensures another level of gene regulation along with protein transcriptional regulators, thus  
293 contributing to an additional layer of tighter regulation. Taking into account the major role of  
294 the CpxAR and EnvZ/OmpR regulators in response to stressors such as antibiotics, it will be  
295 interesting to develop some assays allowing the detection of these kinds of mutations inside  
296 clinical isolate. This original diagnostic maybe used for determining the prevalence of these  
297 regulation events in clinical strain that have undergone antibiotic stress.

298 Targeting the early transcriptional step of antibiotic stress response regulatory mechanism is  
299 much more logical, especially when we have reports of OMP expression being regulated  
300 (both up and downregulation) within 60 minutes of stress appearance [32]. This will  
301 especially promote bypassing of aforementioned mutations in porins in clinical strains that  
302 are selected during antibiotic treatment. Targeting of sRNA or sRNA regulators such as  
303 MicF or Hfq can rejuvenate failing antimicrobial therapies in regards with membrane  
304 impermeability. They can be original targets for increasing the efficiency of existing drugs by  
305 providing fitness reduction in bacteria. As of now, a cyclic peptide RI20 has been identified

306 to inhibit Hfq-mediated repression of gene, by binding with proximal binding site of Hfq  
307 [110]. Another approach will be to inhibit sRNA interfering with porin expression that is  
308 involved in drug translocation. Recently, a small molecule was used to target human  
309 microRNA (miR)-525 precursors as an anti-cancer strategy [111]. This promising discovery  
310 can be repeated in bacteria for manipulating sRNA levels, which may save the failing  
311 antibiotic therapies.

312 Predictability of an efficient drug based on the SICAR (Structure Intracellular Concentration  
313 Activity Relationship) concept, is a step up to efficient drug designing. Briefly, SICAR  
314 connects the physicochemical drug properties to the efficacy of translocation through the  
315 bacterial membrane and the resulting intracellular accumulation. To achieve this goal, an  
316 extensive knowledge of the OM permeability control, including the contribution of sRNAs, is  
317 required.

318 **Funding information:** The research leading to the discussions presented here was conducted  
319 as part of the Marie Curie Initial Training Network TRANSLOCATION consortium and has  
320 received support from the ITN-2013-607694-Translocation (SD). This work was also  
321 supported by Aix-Marseille Univ. and Service de Santé des Armées.

322

323 **Acknowledgments:** We thank all the members of the UMR\_MD1, especially Estelle Dumont  
324 and Julia Vergalli, for helpful discussions throughout this work.

325

326 **Conflicts of interest:** The authors declare no conflict of interest. The founding sponsors had  
327 no role in the design of the study; in the collection, analyses, or interpretation of data; in the  
328 writing of the manuscript, and in the decision to publish the results.

329 **References**

- 330 1. **O'Neill, J.** [http://amr-review.org/sites/default/files/Tackling%20drug-resistant%](http://amr-review.org/sites/default/files/Tackling%20drug-resistant%20infections%20%20An%20overview%20of%20our%20work_IncVaccineLR_NOCROPS.pdf)  
331 [20infections%20%20An%20overview%20of%20our%20work\\_IncVaccineLR\\_NOCROPS.p](http://amr-review.org/sites/default/files/Tackling%20drug-resistant%20infections%20%20An%20overview%20of%20our%20work_IncVaccineLR_NOCROPS.pdf)  
332 [df](http://amr-review.org/sites/default/files/Tackling%20drug-resistant%20infections%20%20An%20overview%20of%20our%20work_IncVaccineLR_NOCROPS.pdf) (accessed March 3, 2016).
- 333 2. **WHO.** [http://apps.who.int/iris/bitstream/10665/112642/1/9789241564748\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/112642/1/9789241564748_eng.pdf)  
334 (accessed March 3, 2016).
- 335 3. **NIAID.** <http://www.niaid.nih.gov/topics/antimicro>  
336 [topics/antimicrobialresistance/documents/arstrategicplan2014.pdf](http://www.niaid.nih.gov/topics/antimicrobialresistance/documents/arstrategicplan2014.pdf) (accessed March 3, 2016).
- 337 4. **Stavenger RA, Winterhalter M.** TRANSLOCATION project: how to get good  
338 drugs into bad bugs. *Sci Transl Med* 2014;6:228ed7.
- 339 5. **Laxminarayan R, Matsoso P, Pant S, Brower C, Røttingen, JA et al.** Access to  
340 effective antimicrobials: a worldwide challenge. *The Lancet* 2016;384:168-175.
- 341 6. **Nikaido H.** Molecular basis of bacterial outer membrane permeability revisited.  
342 *Microbiol Mol Biol Rev* 2003;67:593-656.
- 343 7. **Nikaido H, Pagès JM.** Broad-specificity efflux pumps and their role in multidrug  
344 resistance of Gram-negative bacteria. *FEMS Microbiol Rev* 2012;36:340-63.
- 345 8. **Silhavy TJ, Kahne D, Walker S.** The bacterial cell envelope. *Cold Spring Harb*  
346 *Perspect Bio.* 2010;2:a000414.
- 347 9. **Pagès JM, James, CE, Winterhalter M.** The porin and the permeating antibiotic: a  
348 selective diffusion barrier in Gram-negative bacteria. *Nat Rev Microbiol* 2008;6:893-903.
- 349 10. **Davin-Régli A, Bolla JM, James CE, Lavigne JP, Chevalier J et al.** Membrane  
350 permeability and regulation of drug "influx and efflux" in enterobacterial pathogens. *Curr*  
351 *Drug Targets* 2008;9:750-9.
- 352 11. **Delcour AH.** Outer membrane permeability and antibiotic resistance. *Biochim*  
353 *Biophys Acta* 2009;1794:808-16.
- 354 12. **Masi M, Réfrégiers M, Pos KM, Pagès JM.** Mechanisms of envelope permeability  
355 and antibiotic influx and efflux in Gram-negative bacteria. *Nat Microbiol* 2017;2:17001.
- 356 13. **Nguyen TX, Alegre ER, Kelley ST.** Phylogenetic analysis of general bacterial  
357 porins: a phylogenomic case study. *J Mol Microbiol Biotechnol* 2006;11:291-301.
- 358 14. **Masi M, Pagès JM.** Structure, function and regulation of outer membrane proteins  
359 involved in drug transport in *Enterobacteriaceae*: the OmpF/C-TolC Case. *Open Microbiol J*  
360 2013;7:22–33.
- 361 15. **Guillier M, Gottesman S, Storz G.** Modulating the outer membrane with small  
362 RNAs. *Genes Dev* 2006;20(17):2338-48.

- 363 16. **Vogel J, Papenfort K.** Small non-coding RNAs and the bacterial outer membrane.  
364 *Curr Opin Microbiol* 2006;9:605-611.
- 365 17. **Valentin-Hansen P, Johansen J, Rasmussen AA.** Small RNAs controlling outer  
366 membrane porins. *Curr Opin Microbiol* 2007;10(2):152-155.
- 367 18. **Andersson DI, Hughes D.** Microbiological effects of sublethal levels of antibiotics.  
368 *Nat Rev Microbiol* 2014;12:465-478.
- 369 19. **Gullberg E, Cao S, Berg OG, Ilbäck C, Sandegren L et al.** Selection of resistant  
370 bacteria at very low antibiotic concentrations. *PLoS Pathog* 2011;7:e1002158.
- 371 20. **Gutierrez A, Laureti L, Crussard S, Abida H, Rodríguez-Rojas A et al.**  $\beta$ -lactam  
372 antibiotics promote bacterial mutagenesis via an RpoS-mediated reduction in replication  
373 fidelity. *Nat Commun* 2013;4:1610.
- 374 21. **Lee HH, Molla MN, Cantor CR, Collins JJ.** Bacterial charity work leads to  
375 population-wide resistance. *Nature* 2010;467:82-85.
- 376 22. **Pu Y, Zhao Z, Li Y, Zou J, Ma Q et al.** Enhanced efflux activity facilitates drug  
377 tolerance in dormant bacterial cells. *Mol Cell* 2016;62:284-294.
- 378 23. **Viveiros M, Jesus A, Brito M, Leandro C, Martins M et al.** Inducement and  
379 reversal of tetracycline resistance in *Escherichia coli* K-12 and expression of proton gradient-  
380 dependent multidrug efflux pump genes. *Antimicrob Agents Chemother* 2005;49:3578-3582.
- 381 24. **Viveiros M, Dupont M, Rodrigues L, Couto I, Davin-Regli A et al.** Antibiotic  
382 stress, genetic response and altered permeability of *E. coli*. *PLoS One* 2007;2(4):e365.
- 383 25. **Cohen SP, McMurry LM, Levy SB.** MarA locus causes decreased expression of  
384 OmpF porin in multiple-antibiotic-resistant (Mar) mutants of *Escherichia coli*. *J Bacteriol*  
385 1988;170:5416-5422.
- 386 26. **George AM, Hall RM, Stokes HW.** Multidrug resistance in *Klebsiella pneumoniae*:  
387 a novel gene, *ramA*, confers a multidrug resistance phenotype in *Escherichia coli*.  
388 *Microbiology* 1995;141:1909-1920
- 389 27. **Chubiz LM, Rao CV.** Role of the *mar-sox-rob* regulon in regulating outer membrane  
390 porin expression. *J Bacteriol.* 2011;193:2252-2260.
- 391 28. **Chollet R, Bollet C, Chevalier J, Mallaéa M, Pagès JM et al.** *mar* operon involved  
392 in multidrug resistance of *Enterobacter aerogenes*. *Antimicrob Agents Chemother*  
393 2002;46:1093-1097.
- 394 29. **Chollet R, Chevalier J, Bollet C, Pagès JM, Davin-Régli A.** RamA is an alternate  
395 activator of the multidrug resistance cascade in *Enterobacter aerogenes*. *Antimicrob Agents*  
396 *Chemother* 2004;48:2518-2523.

- 397 30. **Dupont M, Dé E, Chollet R, Chevalier J, Pagès JM.** *Enterobacter aerogenes*  
398 OmpX, a cation-selective channel mar- and osmo-regulated. *FEBS Lett* 2004;569:27-30.
- 399 31. **Gayet S, Chollet R, Molle G, Pagès JM, Chevalier J.** Modification of outer  
400 membrane protein profile and evidence suggesting an active drug pump in *Enterobacter*  
401 *aerogenes* clinical strains. *Antimicrob Agents Chemother.* 2003;47:1555-1559.
- 402 32. **Dupont M, James CE, Chevalier J, Pagès JM.** An early response to environmental  
403 stress involves regulation of OmpX and OmpF, two enterobacterial outer membrane pore-  
404 forming proteins. *Antimicrob Agents Chemother* 2007;51:3190-3198.
- 405 33. **Meccas J, Welch R, Erickson JW, Gross CA.** Identification and characterization of  
406 an outer membrane protein, OmpX, in *Escherichia coli* that is homologous to a family of  
407 outer membrane proteins including Ail of *Yersinia enterocolitica*. *J Bacteriol* 1995;177:799-  
408 804.
- 409 34. **Majdalani N, Gottesman S.** The Rcs phosphorelay: a complex signal transduction  
410 system. *Annu Rev Microbiol* 2005;59:379–405.
- 411 35. **Ruiz N, Silhavy TJ.** Sensing external stress: watchdogs of the *Escherichia coli* cell  
412 envelope. *Curr Opin Microbiol* 2005;8:122-126.
- 413 36. **Rowley G, Spector M, Kormanec J, Roberts M.** Pushing the envelope:  
414 extracytoplasmic stress responses in bacterial pathogens. *Nat Rev Microbiol* 2006;4:383-394.
- 415 37. **Raivio TL.** Everything old is new again: an update on current research on the Cpx  
416 envelope stress response. *Biochim Biophys Acta* 2014;1843:1529-1541.
- 417 38. **Hayden JD, Ades SE.** The extracytoplasmic stress factor, sigmaE, is required to  
418 maintain cell envelope integrity in *Escherichia coli*. *PLoS One* 2008;3(2):e1573.
- 419 39. **Delhaye A, Collet JF, Laloux G.** Fine-Tuning of the Cpx envelope stress response is  
420 required for cell wall homeostasis in *Escherichia coli*. *MBio* 2016;7(1):e00047-16.
- 421 40. **De Las Penas A, Connolly L, Gross CA.** SigmaE is an essential sigma factor in  
422 *Escherichia coli*. *J Bacteriol* 1997;179:6862-6864.
- 423 41. **Wassarman KM, Kiley PJ.** Global approaches for finding small RNA and small  
424 open reading frame functions. *J Bacteriol* 2010;192:26-28.
- 425 42. **Gottesman S.** The small RNA regulators of *Escherichia coli*: roles and mechanisms.  
426 *Annu Rev Microbiol* 2004;58:303-328.
- 427 43. **Thompson KM, Rhodius VA, Gottesman S.** SigmaE regulates and is regulated by a  
428 small RNA in *Escherichia coli*. *J Bacteriol* 2007;189:4243-4256.
- 429 44. **Vogt SL, Evans AD, Guest RL, Raivio TL.** The Cpx envelope stress response  
430 regulates and is regulated by small noncoding RNAs. *J Bacteriol* 2014;196:4229-4238.



- 431 45. **Mizuno T, Mizushima S.** Signal transduction and gene regulation through the  
432 phosphorylation of two regulatory components: the molecular basis for the osmotic  
433 regulation of the porin genes. *Mol Microbiol* 1990;4:1077-1082.
- 434 46. **Pratt LA, Hsing W, Gibson KE, Silhavy TJ.** From acids to *osmZ*: multiple factors  
435 influence synthesis of the OmpF and OmpC porins in *Escherichia coli*. *Mol Microbiol*  
436 1996;20:911-7.
- 437 47. **Cai SJ, Inouye M.** EnvZ-OmpR interaction and osmoregulation in *Escherichia coli*. *J*  
438 *Biol Chem* 2002;277:24155-61.
- 439 48. **Price NL, Raivio TL.** Characterization of the Cpx regulon in *Escherichia coli* strain  
440 MC4100. *J Bacteriol* 2009;191:1798-1815.
- 441 49. **Raivio TL, Leblanc SK, Price NL.** The *Escherichia coli* Cpx envelope stress  
442 response regulates genes of diverse function that impact antibiotic resistance and membrane  
443 integrity. *J Bacteriol* 2013;195:2755-2767.
- 444 50. **Guest RL, Wang J, Wong JL, Raivio TL.** A bacterial stress response regulates  
445 respiratory protein complexes to control envelope stress adaptation. *J Bacteriol* 2017;199:pii:  
446 e00153-17.
- 447 51. **Dartigalongue C, Missiakas D, Raina S.** Characterization of the *Escherichia coli*  
448 sigma E regulon. *J Biol Chem* 2000;276:20866-20875.
- 449 52. **Guest RL, Raivio TL.** Role of the Gram-negative envelope stress response in the  
450 presence of antimicrobial agents. *Trends Microbiol* 2016;24:377-390.
- 451 53. **Mahoney TF, Silhavy TJ.** The Cpx stress response confers resistance to some, but  
452 not all, bactericidal antibiotics. *J Bacteriol* 2013;195(9):1869-74.
- 453 54. **Moreau PL.** Protective role of the RpoE ( $\sigma$ E) and Cpx envelope stress responses  
454 against gentamicin killing of nongrowing *Escherichia coli* incubated under aerobic,  
455 phosphate starvation conditions. *FEMS Microbiol Lett* 2014;357:151-156.
- 456 55. **Batchelor E, Walthers D, Kenney LJ, Goulian M.** The *Escherichia coli* CpxA-  
457 CpxR envelope stress response system regulates expression of the porins OmpF and OmpC. *J*  
458 *Bacteriol* 2005;187:5723-5731.
- 459 56. **Gerken H, Charlson ES, Cicirelli EM, Kenney LJ, Misra R.** MzrA: a novel  
460 modulator of the EnvZ/OmpR two-component regulon. *Mol Microbiol.* 2009;72:1408-1422.
- 461 57. **Gerken H, Misra R.** MzrA-EnvZ interactions in the periplasm influence the  
462 EnvZ/OmpR two-component regulon. *J Bacteriol* 2010;192:6271-8.

- 463 58. **Philippe N, Maigre L, Santini S, Pinet E, Claverie JM et al.** In vivo evolution of  
464 bacterial resistance in two cases of *Enterobacter aerogenes* infections during treatment with  
465 imipenem. *PLoS One* 2015;10:e0138828.
- 466 59. **Mecsas J, Rouviere PE, Erickson JW, Donohue TJ, Gross CA.** The activity of  
467 sigma E, an *Escherichia coli* heat-inducible sigma-factor, is modulated by expression of outer  
468 membrane proteins. *Genes Dev* 1993;7:2618-2628.
- 469 60. **Missiakas D, Mayer MP, Lemaire M, Georgopoulos C, Raina S.** Modulation of  
470 the *Escherichia coli* sigmaE (RpoE) heat-shock transcription-factor activity by the RseA,  
471 RseB and RseC proteins. *Mol Microbiol.* 1997;24(2):355-371.
- 472 61. **Rhodium VA, Suh WC, Nonaka G, West J, Gross CA.** Conserved and variable  
473 functions of the sigmaE stress response in related genomes. *PLoS Biol.* 2006;4:e2.
- 474 62. **Noor R, Murata M, Nagamitsu H, Klein G, Raina S, Yamada M.** Dissection of  
475 sigma(E)-dependent cell lysis in *Escherichia coli*: roles of RpoE regulators RseA, RseB and  
476 periplasmic folding catalyst PpiD. *Genes Cells* 2009;14:885-899.
- 477 63. **Gogol EB, Rhodium VA, Papenfort K, Vogel J, Gross CA.** Small RNAs endow a  
478 transcriptional activator with essential repressor functions for single-tier control of a global  
479 stress regulon. *Proc Natl Acad Sci U S A* 2011;108:12875-12880.
- 480 64. **Klein G, Lindner B, Brade H, Raina S.** Molecular basis of lipopolysaccharide  
481 heterogeneity in *Escherichia coli*: envelope stress-responsive regulators control the  
482 incorporation of glycoforms with a third 3-deoxy- $\alpha$ -D-manno-oct-2-ulosonic acid and  
483 rhamnose. *J Biol Chem* 2011;286:42787-42807.
- 484 65. **Klein G, Kobylak N, Lindner B, Stupak A, Raina S.** Assembly of  
485 lipopolysaccharide in *Escherichia coli* requires the essential LapB heat shock protein. *J Biol*  
486 *Chem* 2014;289:14829-14853.
- 487 66. **Guo MS, Updegrave TB, Gogol EB, Shabalina SA, Gross CA, Storz G.** MicL, a  
488 new  $\sigma$ E-dependent sRNA, combats envelope stress by repressing synthesis of Lpp, the major  
489 outer membrane lipoprotein. *Genes Dev* 2014;28:1620-1634.
- 490 67. **Klein G, Stupak A, Biernacka D, Wojtkiewicz P, Lindner B et al.** Multiple  
491 transcriptional factors regulate transcription of the *rpoE* gene in *Escherichia coli* under  
492 different growth conditions and when the lipopolysaccharide biosynthesis is defective. *J Biol*  
493 *Chem* 2016;291:22999-23019.
- 494 68. **Ades SE.** Regulation by destruction: design of the sigmaE envelope stress response.  
495 *Curr Opin Microbiol* 2008;11:535-540.

- 496 69. **Walsh NP, Alba BM, Bose B, Gross CA, Sauer RT.** OMP peptide signals initiate  
497 the envelope-stress response by activating DegS protease via relief of inhibition mediated by  
498 its PDZ domain. *Cell* 2003;113:61-71.
- 499 70. **Lima S, Guo MS, Chaba R, Gross CA, Sauer RT.** Dual molecular signals mediate  
500 the bacterial response to outer-membrane stress. *Science* 2013;340(6134):837-841.
- 501 71. **Mizuno T, Chou MY, Inouye M.** A unique mechanism regulating gene expression:  
502 translational inhibition by a complementary RNA transcript (micRNA). *Proc Natl Acad Sci U*  
503 *SA* 1984;81:1966-1970.
- 504 72. **Delihias N.** Discovery and characterization of the first non-coding RNA that regulates  
505 gene expression, *micF* RNA: A historical perspective. *World J Biol Chem* 2015;6:272-280.
- 506 73. **Inouye M.** The first demonstration of RNA interference to inhibit mRNA function.  
507 *Gene* 2016;592(2):332-333.
- 508 74. **Andersen J, Delihias N.** *micF* RNA binds to the 5' end of *ompF* mRNA and to a  
509 protein from *Escherichia coli*. *Biochemistry* 1990;29:9249-9256.
- 510 75. **Delihias N, Forst S.** MicF: an antisense RNA gene involved in response of  
511 *Escherichia coli* to global stress factors. *J Mol Biol* 2001;313:1-12.
- 512 76. **Ramani N, Hedeshian M, Freundlich M.** *micF* antisense RNA has a major role in  
513 osmoregulation of OmpF in *Escherichia coli*. *J Bacteriol* 1994;176:5005-5010.
- 514 77. **Chou JH, Greenberg JT, Demple B.** Post-transcriptional repression of *Escherichia*  
515 *coli* OmpF protein in response to redox stress: positive control of the *micF* antisense RNA by  
516 the *soxRS* locus. *J Bacteriol* 1993;175:1026-1031.
- 517 78. **Chen S, Zhang A, Blyn LB, Storz G.** MicC, a second small-RNA regulator of Omp  
518 protein expression in *Escherichia coli*. *J Bacteriol* 2004;186:6689-6697.
- 519 79. **Prilipov A, Phale PS, Koebnik R, Widmer C, Rosenbusch JP.** Identification and  
520 characterization of two quiescent porin genes, *nmpC* and *ompN*, in *Escherichia coli* B<sup>E</sup>. *J*  
521 *Bacteriol* 1998;180:3388-3392.
- 522 80. **Dam S, Pagès JM, Masi M.** Dual regulation of the small RNA MicC and the  
523 quiescent porin OmpN in response to antibiotic stress in *Escherichia coli*. *Antibiotics* (Basel)  
524 2017;6:pii: E33.
- 525 81. **Guillier M, Gottesman S.** Remodelling of the *Escherichia coli* outer membrane by  
526 two small regulatory RNAs. *Mol Microbiol* 2006;59:231-47.
- 527 82. **Guillier M, Gottesman S.** The 5' end of two redundant sRNAs is involved in the  
528 regulation of multiple targets, including their own regulator. *Nucleic Acids Res*  
529 2008;36:6781-6794.


- 530 83. **Rasmussen AA, Eriksen M, Gilany K, Udesen C, Franch T *et al.*** Regulation of  
531 *ompA* mRNA stability: the role of a small regulatory RNA in growth phase-dependent  
532 control. *Mol Microbiol* 2005;58:1421-1429.
- 533 84. **Johansen J, Rasmussen AA, Overgaard M, Valentin-Hansen P.** Conserved small  
534 non-coding RNAs that belong to the sigmaE regulon: role in down-regulation of outer  
535 membrane proteins. *J Mol Biol* 2006;364:1-8.
- 536 85. **Skovierova H, Rowley G, Rezuchova B, Homerova D, Lewis *et al.*** Identification  
537 of the sigmaE regulon of *Salmonella enterica* serovar Typhimurium. *Microbiology*  
538 2006;152:1347-1359.
- 539 86. **Douchin V, Bohn C, Bouloc P.** Down-regulation of porins by a small RNA bypasses  
540 the essentiality of the regulated intramembrane proteolysis protease RseP in *Escherichia coli*.  
541 *J Biol Chem* 2006;281:12253-12259.
- 542 87. **Papenfort K, Pfeiffer V, Mika F, Lucchini S, Hinton JC *et al.*** SigmaE-dependent  
543 small RNAs of *Salmonella* respond to membrane stress by accelerating global *omp* mRNA  
544 decay. *Mol Microbiol* 2006 Dec;62(6):1674-88.
- 545 88. **Bossi L, Figueroa-Bossi N.** A small RNA downregulates LamB maltoporin in  
546 *Salmonella*. *Mol Microbiol* 2007;65(3):799-810.
- 547 89. **Udekwu KI, Wagner EG.** Sigma E controls biogenesis of the antisense RNA MicA.  
548 *Nucleic Acids Res* 2007;35:1279-1288.
- 549 90. **Johansen J, Eriksen M, Kallipolitis B, Valentin-Hansen P.** Down-regulation of  
550 outer membrane proteins by noncoding RNAs: unraveling the cAMP-CRP- and sigmaE-  
551 dependent CyaR-OmpX regulatory case. *J Mol Biol.* 2008;383:1-9.
- 552 91. **Castillo-Keller M, Vuong P, Misra R.** Novel mechanism of *Escherichia coli* porin  
553 regulation. *J Bacteriol* 2006;188:576-586.
- 554 92. **Doménech-Sánchez A, Hernández-Allés S, Martínez-Martínez L, Benedí VJ,  
555 Albertí S.** Identification and characterization of a new porin gene of *Klebsiella pneumoniae*:  
556 its role in beta-lactam antibiotic resistance. *J Bacteriol* 1999;181:2726-2732.
- 557 93. **Knopp M, Andersson DI.** Amelioration of the fitness costs of antibiotic resistance  
558 due to reduced outer membrane permeability by upregulation of alternative porins. *Mol Biol*  
559 *Evol* 2015;32:3252-3263.
- 560 94. **Pomposiello PJ, Bennik MH, Demple B.** Genome-wide transcriptional profiling of  
561 the *Escherichia coli* responses to superoxide stress and sodium salicylate. *J Bacteriol*  
562 2001;183:3890-902.

- 563 95. **Bornet C, Davin-Régli A, Bosi C, Pagès JM, Bollet C.** Imipenem resistance of  
564 *Enterobacter aerogenes* mediated by outer membrane permeability. *J Clin Microbiol*  
565 2000;38:1048-1052.
- 566 96. **Bornet C, Chollet R, Malléa M, Chevalier J, Davin-Régli A et al.** Imipenem and  
567 expression of multidrug efflux pump in *Enterobacter aerogenes*. *Biochem Biophys Res*  
568 *Commun* 2003;301:985-990.
- 569 97. **Thiolas A, Bornet C, Davin-Régli A, Pagès JM, Bollet C.** Resistance to imipenem,  
570 cefepime, and cefpirome associated with mutation in Omp36 osmoporin of *Enterobacter*  
571 *aerogenes*. *Biochem Biophys Res Commun* 2004;317(3):851-856.
- 572 98. **Thiolas A, Bollet C, La Scola B, Raoult D, Pagès JM.** Successive emergence of  
573 *Enterobacter aerogenes* strains resistant to imipenem and colistin in a patient. *Antimicrob*  
574 *Agents Chemother* 2005;49:1354-1358.
- 575 99. **Lavigne JP, Sotto A, Nicolas-Chanoine MH, Bouziges N, Pagès JM et al.** An  
576 adaptive response of *Enterobacter aerogenes* to imipenem: regulation of porin balance in  
577 clinical isolates. *Int J Antimicrob Agents* 2013;41:130-136.
- 578 100. **Pavez M, Vieira C, de Araujo MR, Cerda A, de Almeida LM et al.** Molecular  
579 mechanisms of membrane impermeability in clinical isolates of *Enterobacteriaceae* exposed  
580 to imipenem selective pressure. *Int J Antimicrob Agents* 2016;48:78-85.
- 581 101. **Cowan SW, Schirmer T, Rummel G, Steiert M, Ghosh R et al.** Crystal structures explain  
582 functional properties of two *E. coli* porins. *Nature* 1992;358:727-733.
- 583 102. **Baslé A, Rummel G, Storici P, Rosenbusch JP, Schirmer T.** Crystal structure of  
584 osmoporin OmpC from *E. coli* at 2.0 Å. *J Mol Biol* 2006;362:933-942.
- 585 103. **Nikaido H, Rosenberg EY.** Porin channels in *Escherichia coli*: studies with  
586 liposomes reconstituted from purified proteins. *J Bacteriol* 1983;153:241-252.
- 587 104. **Nikaido H, Rosenberg EY, Foulds, J.** Porin channels in *Escherichia coli*: studies  
588 with beta-lactams in intact cells. *J Bacteriol* 1983;153:232-240.
- 589 105. **Kojima S, Nikaido H.** High salt concentrations increase permeability through OmpC  
590 channels of *Escherichia coli*. *J Biol Chem* 2014;289:26464-26473.
- 591 106. **Dé E, Baslé A, Jaquinod M, Saint N, Malléa M et al.** A new mechanism of  
592 antibiotic resistance in *Enterobacteriaceae* induced by a structural modification of the major  
593 porin. *Mol Microbiol* 2001;41:189-198.
- 594 107. **Simonet V, Malléa M, Pagès JM.** Substitutions in the eyelet region disrupt cefepime  
595 diffusion through the *Escherichia coli* OmpF channel. *Antimicrob Agents Chemother*  
596 2000;44:311-315.

- 597 108. **Low AS, MacKenzie FM, Gould IM, Booth IR.** Protected environments allow  
598 parallel evolution of a bacterial pathogen in a patient subjected to long-term antibiotic  
599 therapy. *Mol Microbiol* 2001;42:619-630.
- 600 109. **Lou H, Chen M, Black SS, Bushell SR, Ceccarelli M. et al.** Altered antibiotic  
601 transport in OmpC mutants isolated from a series of clinical strains of multi-drug resistant *E.*  
602 *coli*. *PLoS One* 2011;6(10):e25825.
- 603 110. **El-Mowafi SA, Alumasa JN, Ades SE, Keiler KC.** Cell-based assay to identify  
604 inhibitors of the Hfq-sRNA regulatory pathway. *Antimicrob Agents Chemother*  
605 2014;58:5500-5509.
- 606 111. **Childs-Disney JL, Disney MD.** Small molecule targeting of a microRNA associated  
607 with hepatocellular carcinoma. *ACS Chem Biol* 2016;11:375-380.

608 **Key figure: Major regulatory pathways of porin regulation in *E. coli*:** EnvZ/OmpR [46],  
609 CpxAR and sigma E ( $\sigma^E$ ) [35] stress response systems are shown, along with known inducing  
610 cues and targets relevant to porin regulation. The upregulation is shown with thick green  
611 arrows, while the downregulation is shown with red lines. In the EnvZ/OmpR TCS,  
612 activation of the response regulator OmpR results in phosphorylated and OmpR~P  
613 downregulates the expression of OmpF both at the transcriptional and post-transcriptional  
614 levels, the latter via the MicF sRNA. The *mar-sox-rob* regulons also downregulate OmpF  
615 expression via MicF. Both the CpxAR and  $\sigma^E$  responses are activated by a variety of  
616 envelope stresses. For clarity, only periplasmic misfolded OMPs are represented here. On one  
617 hand, CpxR~P alters expression of multiple genes, including that of *micF*. On the other hand,  
618 the anti-sigma factor RseA is degraded by the successive action of two proteases, DegS and  
619 RseP at the periplasmic and the cytoplasmic site. Another protease, ClpXP specifically  
620 degrades the cytoplasmic RseA portion bound to  $\sigma^E$ , leading to its release. A number of  $\sigma^E$ -  
621 regulated sRNAs are indicated: MicC [78] downregulates OmpC and is coupled with *ompN*  
622 upregulation [80]; sRNA regulation of porins via CyaR [90], IpeX [91], RseX [86] and RybB  
623 [84, 88] are shown accompanied by their activators and porin targets; CyaR negatively  
624 regulates the expression of single channeled porin OmpX [30], which in turn negatively  
625 regulates the major porin OmpC. The details of all these interconnected pathways are  
626 discussed thoroughly in the text.

627



628