

Across the Gap: Geochronological and Sedimentological Analyses from the Late Pleistocene-Holocene Sequence of Goda Buticha, Southeastern Ethiopia

Chantal Tribolo, Asfawossen Asrat, Jean-Jacques Bahain, Cécile Chapon, Eric Douville, Carole Fragnol, Marion Hernandez, Erella Hovers, Alice Leplongeon,

Loïc Martin, et al.

▶ To cite this version:

Chantal Tribolo, Asfawossen Asrat, Jean-Jacques Bahain, Cécile Chapon, Eric Douville, et al.. Across the Gap: Geochronological and Sedimentological Analyses from the Late Pleistocene-Holocene Sequence of Goda Buticha, Southeastern Ethiopia. PLoS ONE, 2017, 12 (1), pp.e0169418. 10.1371/journal.pone.0169418 . hal-01841520

HAL Id: hal-01841520 https://hal.science/hal-01841520

Submitted on 28 Oct 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Citation: Tribolo C, Asrat A, Bahain J-J, Chapon C, Douville E, Fragnol C, et al. (2017) Across the Gap: Geochronological and Sedimentological Analyses from the Late Pleistocene-Holocene Sequence of Goda Buticha, Southeastern Ethiopia. PLoS ONE 12 (1): e0169418. doi:10.1371/journal.pone.0169418

Editor: Nuno Bicho, Universidade do Algarve, PORTUGAL

Received: July 26, 2016

Accepted: December 17, 2016

Published: January 26, 2017

Copyright: This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the <u>Creative</u> Commons CC0 public domain dedication.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This research benefited from funding of the National Geographic Society (<u>http://</u> <u>nationalgeographic.org/</u>), grants # 8110-06 and 8510-08 (ZA), of the Wenner-Gren Foundation (<u>http://www.wennergren.org/</u>), grant # ICRG e 102, (ZA and DP), of the Hebrew University of Jerusalem (<u>http://new.huji.ac.il/en</u>) (EH), of the French National Research Agency (<u>http://www.</u> **RESEARCH ARTICLE**

Across the Gap: Geochronological and Sedimentological Analyses from the Late Pleistocene-Holocene Sequence of Goda Buticha, Southeastern Ethiopia

Chantal Tribolo¹*, Asfawossen Asrat², Jean-Jacques Bahain³, Cécile Chapon³, Eric Douville⁴, Carole Fragnol¹, Marion Hernandez¹, Erella Hovers^{5,6}, Alice Leplongeon^{3,7}, Loïc Martin¹, David Pleurdeau³, Osbjorn Pearson⁸, Simon Puaud³, Zelalem Assefa⁹

1 IRAMAT-CRP2A, UMR 5060, CNRS, Université de Bordeaux, Pessac, France, 2 School of Earth Sciences, Addis Ababa University, Addis Ababa, Ethiopia, 3 Laboratoire « Histoire naturelle de l'Homme préhistorique », UMR 7194, CNRS, Département de Préhistoire du Muséum National d'Histoire Naturelle, Paris, France, 4 Laboratoire des Sciences du Climat et de l'Environnement, UMR 8212, CEA-CNRS, Gif-sur-Yvette, France, 5 Institute of Archaeology, Hebrew University, Jerusalem, Israel, 6 Institute of Human Origins, Arizona State University, Phoenix, United States of America, 7 McDonald Institute for Archaeological Research, Cambridge University, Cambridge, United Kingdom, 8 Department of Anthropology, University of New Mexico, Albuquerque, United States of America, 9 Department of Anthropology, Archaeobiology Program & Human Origins Program, National Museum of Natural History, Washington, D.C., United States of America

* ctribolo@u-bordeaux-montaigne.fr

Abstract

Goda Buticha is a cave site near Dire Dawa in southeastern Ethiopia that contains an archaeological sequence sampling the late Pleistocene and Holocene of the region. The sedimentary sequence displays complex cultural, chronological and sedimentological histories that seem incongruent with one another. A first set of radiocarbon ages suggested a long sedimentological gap from the end of Marine Isotopic Stage (MIS) 3 to the mid-Holocene. Macroscopic observations suggest that the main sedimentological change does not coincide with the chronostratigraphic hiatus. The cultural sequence shows technological continuity with a late persistence of artifacts that are usually attributed to the Middle Stone Age into the younger parts of the stratigraphic sequence, yet become increasingly associated with lithic artifacts typically related to the Later Stone Age. While not a unique case, this combination of features is unusual in the Horn of Africa. In order to evaluate the possible implications of these observations, sedimentological analyses combined with optically stimulated luminescence (OSL) were conducted. The OSL data now extend the radiocarbon chronology up to 63 ± 7 ka; they also confirm the existence of the chronological gap between 24.8 ± 2.6 ka and 7.5 ± 0.3 ka. The sedimentological analyses suggest that the origin and mode of deposition were largely similar throughout the whole sequence, although the anthropic and faunal activities increased in the younger levels. Regional climatic records are used to support the sedimentological observations and interpretations. We discuss the implications of the sedimentological and dating analyses for understanding cultural processes in the region.



agence-nationale-recherche.fr), projects # ANR-09-JCJC-0123-01 (CT), # ANR-14-CE31-0023-03, and # ANR-10-LABX-52, of the Regional Priority Program « Heritage, Resources, Governance » (HEREGO, coordinated by the Institut de Recherche pour le Developpement) (DP), the Doctoral School of the Museum National d'Histoire Naturelle in Paris (France) and UMR CNRS 7194 (AL), the Smithsonian NMNH small grants, and the European Union's Horizon 2020 research and innovation program under the Marie Sklodowska-Curie grant (https://ec.europa.eu/programmes/ horizon2020/en/h2020-section/marie-sklodowskacurie-actions), agreement #655459. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

1-Introduction

A large body of recent and ongoing archaeological researches in sub-Saharan Africa has focused on the transition between the Middle Stone Age (MSA) and the Later Stone Age (LSA), e.g. at Border cave [1], Mumba Rock Shelter [2], and Enkapune Ya Muto [3]. These studies have extended the date of the earliest LSA occurrences back to ca. 50 ka BP. Typically the start of LSA is marked by the appearance of more complex hunting and fishing technologies (hafted microliths, bows and arrows, polished bone tools) or symbolic items (e.g., ostrich eggshell or marine shell beads, engraved ocher, eggshells, bones and wood [4–6]), even though some of these items appeared earlier during the MSA (e.g., during the Stillbay and Howiesons Poort in southern Africa [7–9]). These studies indicate that the emergence of the LSA may involve variable technical, symbolic or social aspects in different geographic regions and different points in time. The mosaic pattern might be partly related to human dispersals and demographic shifts within and out of Africa, assumed to occur from ca. 70 ka onward, e.g. [1,10–13].

In the Horn of Africa, the evidence and processes of the MSA-LSA transition are poorly understood. Indeed, despite a history of extensive archaeological surveys [14–19], comparatively few stratified sites document this period. In Ethiopia, such sites include K'one [20], Porc-Epic [21–23], Aladi Springs [16, 24, 25], Ziway-Shala Basin sites [26], and Mochena Borago [27]. In Somalia, the site of Midhishi 2 also falls within this time range [28, 29], (Fig 1). In most of these sites, the archaeological sequences are short, discontinuous, and/or not well dated. At Mochena Borago and the Ziway-Shala Basin sites, which record a relatively long time span from Marine Isotope Stage (MIS) 4–3 to the Holocene, the sequences seem to share a major chronostratigraphic gap broadly corresponding to MIS 2 [26, 27].

A similar gap has been observed at Goda Buticha in southeastern Ethiopia [30]. The archaeological stratigraphic sequence was first dated with radiocarbon from ~ 46 ka cal BP for the lower layers, to ~ 1.5 ka cal BP near the present-day surface level. The dates revealed a chronological gap, from ~ 33 ka cal BP (end of MIS 3) to ~ 7.8 ka cal BP (mid-Holocene) in the sedimentary sequence [30] (see §2.2 for details). The cultural assemblages of Goda Buticha do not show a clear technological rupture before and after the gap. The lowest layers show MSA technical characteristics with few LSA components, while the upper layers contain typical MSA components, together with artefacts with clear LSA characteristics (see §2.2). The consistency of the radiocarbon ages, which display no significant chronostratigraphic inversion, and the absence of clear evidence for large-scale disturbances of the stratigraphic layers such as redeposition by cut and fill, argue against post-depositional mixing as an explanatory mechanism for this pattern. The late persistence of MSA components is unusual (though not unknown) in the region [30, 31], and required corroboration through sedimentological and dating analyses that allow better control for temporal and depositional processes.

Recently, genetic studies have shown the important role of the Horn of Africa in Upper Pleistocene and Holocene human dispersals from and into Africa [32–36]. There have also been suggestions that certain areas in the Horn of Africa, specifically the Southwest High Plateau, served as biogeographic refugia to Upper Pleistocene populations during times of climatic instability or deterioration [27, 37]. Such scenarios are important when we evaluate the sequence from Goda Buticha and its place within the larger picture of human history on the African continent.

The main goals of this study were i) to extend the age model of the stratigraphic sequence by dating the lowermost levels, where no charcoal could be dated; ii) to test the robustness of the ¹⁴C chronology by using an additional dating technique, and in particular to confirm or refute the existence of the chronological gap; iii) if confirmed, to investigate how the gap is manifested in the sedimentological and geochemical site formation processes. In this paper,



Fig 1. The Horn of Africa with location of Goda Buticha and sites mentioned in the text.

we present detailed sedimentological analyses combined with a new chronological framework based on optically stimulated luminescence ages (OSL) on sediments from the various stratigraphic units of Goda Buticha.

2-The Goda Buticha (Buticha Cave) sequence

2.1 Site setting

In 2007, geological and archaeological survey of cave formations in eastern Ethiopia led to the discovery of several previously unknown or poorly known caves bearing anthropogenic deposits [38]. Among them, Goda Buticha ("Cave of the Ghost"), revealed a deep archaeological sequence, comprising multiple Paleolithic occupation horizons [30]. The cave, situated ~15 km west of the city of Dire Dawa at an elevation of 1382 m above mean sea level (asl), is located in the escarpment of the Southeastern Ethiopian Highlands to the south, just outside the Main Ethiopian Rift to the north (Fig 2). It is carved into Jurassic crystalline limestone beds in a cliff side and comprises two main galleries oriented NNW and NE, connected by a short perpendicular passage. The sedimentological sequence is deposited in the NNW gallery of the branching cave system [38].

We thank the Ethiopian Authority for Research and Conservation of Cultural Heritage (ARCCH) for authorization to excavate in Goda Buticha. The excavations at Goda Buticha took place in 2008 and 2011. At a distance of four meters from the cave entrance two square-meters were dug inside the cave, revealing a > 230 cm thick sequence (Fig 2). More than 5 m³ of sediments have been removed and sieved, and several thousand archaeological finds (lithics, faunal remains, and pottery) have been plotted in 3D or recovered from a 2 mm-mesh sieve. Some of the archaeological materials and their general chronostratigraphic context have been studied and published elsewhere [30, 31, 38, 39, 40]. In the following section, we summarize the main features and results.

2.2 Stratigraphy, ¹⁴C chronology and cultural sequence at Goda Buticha

The excavated stratigraphic sequence of Goda Buticha site is 230 cm thick, but the bedrock has not been reached. The top is partially capped by deposits of modern bat guano. The sequence is divided into two distinct complexes [30] (Fig 3 and S1 Fig): Complex II (also named "Lower Complex" in previous publications) at the bottom (at elevation of 0 to 180 cm above datum),





Fig 2. Map of the Goda Buticha area (a), view of the escarpment where the cave is located (b) and map of the cave showing the excavated area (c).

and Complex I (also "Upper Complex" in earlier publications) at the top (180 to 230 cm above datum). The sediments are dry and powdery in both complexes. Two sedimentological subsets were identified in Complex II: (i) from Layer IIf to Layer IId, the sediment is silty and contains limestone clasts. In Layer IIf, the sediment is partially impregnated by calcium carbonate; (ii) from Layer IIc to Layer IIa, the sediment also has a silty matrix but the limestone fragments are more numerous than in the lower sub-unit and of more diverse sizes, ranging from gravels to blocks. The contact between the two subsets is only weakly distinguished in the field by this difference in limestone concentration and size, but there is no marked discontinuity.

Complex I is formed by a succession of ashy layers (Layers Ih to Ia1). The sediments are dark in color, sandy to silty, with a high calcium carbonate content and abundant micromammals remains. They are devoid of a coarse fraction.

Fourteen AMS radiocarbon analyses on charcoal samples [30], have provided ages for Complex II (Table 1): (i) the lower sub-unit of Complex II is of Upper Pleistocene age, dating from at least 46273 ± 2084 cal BP (42500 ± 1000 BP, top of Layer IIf) to 33843 ± 405 cal BP (29680 ± 230 BP, Layer IId), thus falling within MIS 3; (ii) the upper sub-unit of Complex II is of a Holocene age with dates from 7791 ± 126 cal BP (6920 ± 50 BP, Layer IIc) to 4676 ± 144 cal BP (4130 ± 30 BP, Layer IIb). Two additional charcoal samples have provided consistent late Holocene ages for Complex I, of 4060 ± 90 cal BP (3725 ± 30 BP, Layer If) and 1485 ± 75 cal BP (1610 ± 30 BP, Layer Ic). Note that there is some scatter in the ¹⁴C ages and thus there are a few chronostratigraphic reversals (within Layer IIf and within Layer IIc), but there is no sign of significant mixing between the samples of Pleistocene and Holocene age, or between the samples of mid and late Holocene.

The dates suggest a chronological and stratigraphic gap within Complex II. This gap does not coincide with the main visible change in the sediments (which is between Complex II and



Fig 3. Stratigraphic section of the Goda Buticha sequence. The locations of the OSL samples, ¹⁴C samples, and samples for sedimentological analyses are shown. Note that the nomenclature has changed for the layers of Complex I compared to previous publications [30].

Complex I) or with any abrupt cultural change [30]. The lower sub-unit of Complex II (Layers IId-IIf) shows clear MSA technical features, with few LSA components (Fig 4). The assemblage is oriented towards flake production together with relatively high frequencies of items related to blade production and with typical MSA unifacial and bifacial points. Both flakes and blades were obtained through the use of a planimetric conception of debitage, including Levallois methods, with typical MSA unifacial and bifacial points. The upper sub-unit (Layers IIa-IIc) shows a clear tendency for the production of blades, bladelets, and non-geometric microliths in tandem with typical MSA components (e.g. unifacial or bifacial points and Levallois products [31, 40]). Complex I, despite a very low density of artifacts, is characterized by very different technical features, such as the abundant use of obsidian to produce small flakes and blades that were then transformed into geometric microliths [30, 39].

A small sample (n = 340) of mammalian faunal remains were piece-plotted during the excavation. Several hundreds of small fragments of medium to large-sized mammals, as yet unstudied, were collected from the sieved sediments. Of the piece-plotted faunal remains, 10% are

Complex	Elevation above datum (cm)	Layer	Grid square	Lab référence	Number in Fig 3	Uncalibrated age	Calibrated Age BP
Complex I	200–210	lc	A1a/A1b	SacA#42484	12	1610 ± 30	1487 ± 47
	190–180	lf	B1a/B1b	SacA#42483	11	3725 ± 30	4067 ± 55
Complex II upper sub-	180–170						
unit	170–160	llb	B1c/B1d	SacA#27480	1	4130 ± 30	4680 ± 81
	160–150						
	150–140	llc	A1c/A1d	SacA#27479	3	6410 ± 30	7351 ± 43
	140–130	llc	B1a/B1b	SacA#35205	4**	6200 ± 30	7094 ± 57
	140–130	llc	B1c/B1d	beta #255124	4*	5590 ± 50	6372 ± 47
	130–120	llc	A1a/A1b	beta#255123	5*	6920 ± 50	7757 ± 56
	130–120	llc	B1a/B1b	beta#269596	5**	6940 ± 40	7771 ± 51
	130–120	llc	B1c/B1d	beta#269597	5***	6430 ± 40	7357 ± 42
	120–110	llc	A1a/A1b	beta#269594	6*	6470 ± 40	7376 ± 40
	120–110	llc	B1a/B1b	beta#269595	6**	6370 ± 50	7314 ± 60
	120–110	llc	B1c/B1d	beta#255122	6***	6440 ± 50	7360 ± 46
Complex II lower sub-	110–100						
unit	100–90	lld	A1c/A1d	beta#269593	8*	29680 ± 230	33839 ± 198
	100–90	lld	A1c/A1d	beta#255121	8**	31590 ± 270	35486 ± 306
	90–80						
	80–70						
	70–60	llf		SacA#35207	9	42500 ± 1000	46089 ± 1042
	60–50	llf		SacA#35206	10	38870 ± 670	42907 ± 525

Table 1. Radiocarbon ages for charcoal samples from Goda Buticha.

PLOS ONE

All the ¹⁴C samples are charcoals. The 95% confidence interval is indicated for all data. The ¹⁴C ages have been calibrated using Oxcal and the Intcal13 calibration curve [41, 42].

doi:10.1371/journal.pone.0169418.t001

teeth or jaws, 10% are long bone shaft fragments, and the rest (80%) are unidentifiable bone fragments. Many of the remains are in friable condition, with poor surface preservation, which discouraged initial efforts to identify any anthropogenic or carnivore modifications. Thus, the preliminary analysis focused on taxonomic identifications [30].

A fairly diverse set of taxa has been identified, including Procaviidae, Lagomorpha, *Equus*, *Phacochoerus*, a small carnivore, and several types of bovids including bovini, antilopini, *Redunca*, and *Tragelaphus*. No domesticated fauna were recovered from any of the stratigraphic levels at the site. Nearly 100 fragments of ostrich eggshell (some measuring up to 4–5 cm in diameter) occurred throughout the sediment column, with frequencies increasing from the bottom to the top of the sequence. Likewise, micro-faunal remains, sometimes heavily burned, occur throughout the sequence, but their relative frequency and the proportion of burned specimens both decrease with depth. A detailed analysis of the microfauna and its paleoenvironmental implication will be published elsewhere, so the data will not be discussed here.

In addition to the rich archaeological assemblages, the excavation yielded ten hominin specimens [30]. A cranial vault fragment was found in Complex II (Layer IId), at an elevation of 40 cm above datum and is potentially older than ~42–46 ka cal BP (Table 1). The human remains of Goda Buticha, recovered from the MIS 3 layers (i.e. Layer IId, close to the 47–42 ka ¹⁴C cal BP samples) and from the Holocene deposits (i.e., Layer IIc dated from ca. 7 to 4 ka cal BP) are of importance given the paucity of human fossils from the time range of MIS 3 to 1 [43] and given the genetic and archaeological arguments for out of Africa and back to Africa movements during the Upper Pleistocene and Holocene [33, 35, 36].



Fig 4. Lithic artifacts from Complex I and Layers IIc and IId-IIf of Goda Buticha. Complex I: a, b: cores; c-e: geometric microliths. Layer IIc: f: Levallois preferential core; g: blade core; h: crested blade; i, j: backed pieces, i is microlithic; k, l: elongated backed points; m, n: bifacial retouched points. Layers IId-IIf: o,p: cores; p is Levallois-like preferential core; q: Levallois blade; r,s,t: retouched points.

3- Materials and Methods

To obtain the goals of this research regarding the chronology of the site and its depositional dynamics, the entire stratigraphic sequence was sampled at high resolution (Fig 3). Nine samples were analyzed for OSL dating at the Institut de Recherche sur les ArchéoMATériaux (IRAMAT), University of Bordeaux-Montaigne. Each of the nine OSL samples were divided into two sub-samples, one of which was added to the separate batch of sixteen sediment samples exported to the Museum National d'Histoire Naturelle (MNHN) in Paris for sedimento-logical and geochemical analyses.

PLOS ONE

3.1 Sedimentological analyses

After dry sieving through a 2 mm mesh in the lab, each sample was split into several aliquots. The first one was used for quantification of carbonates using a calcimeter and organic material by calcination at 375 °C for 16 hours. The second aliquot was used for measurement of pH and grain size analyses. After wet sieving and drying, the 50–200 μ m fraction was observed under an optical microscope for the determination of heavy and light minerals and for the analysis of the quartz grain morphoscopy. Magnetic susceptibility was measured at low frequency (0.47 kHz) for each bulk sample. Another portion of each bulk sample was powdered and used for X Ray Diffraction (XRD) analyses. Mineralogical identifications and semi-quantitative estimates were then performed. Further details about the analytical procedures are given in Appendix 1.

3.2 Optically stimulated luminescence (OSL) dating analyses

Luminescence is a paleodosimetric dating method. In order to calculate ages, two main parameters must be determined: the total dose accumulated by the samples since the dated event (equivalent dose, De) and the dose delivered by the environment per unit of time (dose rate, Dr). The dated event is the last exposure to sunlight (or of heating to high temperature), which resets the OSL signal. The methods for estimating the De rely on the measurements of luminescence signals, the intensity of which is related to the amount of dose received since the last reset [44, 45]. We describe here the main procedures of the analyses that were conducted for the determination of the De and Dr. Details about the preparation of the samples, measurement units and tests of the protocols for the De estimate can be found in Appendix 2. Comments specific to the beta Dr estimation are provided in Appendix 3.

Collection of the nine sediment samples, labeled GDB1 to 9 (Fig 3), was conducted at night, using subdued orange light, and the samples were immediately packed into black plastic bags. In the laboratory, quartz grains in the size range of $200-250 \mu m$ were extracted from each sample following standard procedures (Appendix 2). Tests for the protocols of the De determinations were conducted on both multi-grains and single-grain aliquots, while for the final De measurements single-grain aliquots were preferred since they avoid any averaging effects. The single aliquot and regenerative dose protocol was applied (SAR [46, 47]; Appendix 2).

The total Dr can be decomposed into cosmic, gamma, alpha and beta Dr. The cosmic Dr were calculated by applying the equation of Prescott and Hutton [48], taking into account the geometry of the cave ceiling. The gamma Dr were estimated from field gamma spectrometry (using a LaBr₃(Ce) 1.5x1.5 probe, Canberra), conducted using the threshold technique [49, 50]. The alpha Dr were assumed to be negligible since the external part of the grains had been etched. The beta Dr was estimated from the U, Th and K contents of each sample following Martin [51, 52]. This protocol intends to investigate the effect of heterogeneity at the millimeter scale on the Dr estimate (Appendix 3). After drying, fine grinding and sealing of the samples, U, Th and K contents of the fine (< 2 mm) and coarse (2 mm-1 cm) fractions were analyzed separately with high-resolution gamma spectrometry (HRGS), except for the coarse fraction of samples GDB3, 4, 7 and 8. For these sub-samples, Inductively Coupled Plasma Spectrometry (ICP-MS and ICP-AES) was used instead, because the prepared amount of each sub-sample was too small for standard HGRS. The beta Dr was then calculated using the attenuation factors for the grain size of Guérin et al. [53] and specific dose rates of Guérin et al. [50]. Water content (see §4.2.1) was incorporated into dose rate calculations to account for attenuation.





4- Results

4.1 Sedimentology

The results of the geological analyses are displayed in Fig 5 (grain size analyses, quartz grain morphoscopy, carbonate and organic quantification, pH values, and magnetic susceptibility) and in Fig 6 and S1 Table (XRD analyses).

Twenty mineralogical species corresponding to 6 chemical groups were identified by XRD: magmatic minerals (quartz, feldspars, muscovite, chlorite, hornblend, rutile), carbonates (calcite, dolomite), evaporites (gypsum, anhydrite, halite, sylvite), phosphates (carbonate hydroxyl-apatite, dittmarite, whitlockite), calcium oxalates (weddellite, whewellite) and clays (kaolinite, montmorillonite). The minerals (determined on sieved and decarbonated sediment) are dominated by magmatic minerals (25 to 55%) and are present in all samples. Calcite (determined from the raw sediment) is abundant (5 to 70%) and present in the entire sequence; it dominates in the lower part of Complex I (Layers Ig/If), and the middle part of Complex II (Layer IId).

Complex II. Complex II is characterized by brown, well-sorted silts and fine sands (Fig 5A). The clay concentrations are low (< 5%) but omnipresent (Fig 5A). The morphoscopy analyses show that on average, 80% of the quartz grains are sub-rounded and lusterless while 20% are angular or sub-rounded and shiny (Fig 5B). The variation of the morphoscopic classes through Complex II is not statistically significant. In addition to quartz and feldspar (plagioclase and orthoclase) grains, apatite and a limited variety of heavy minerals including







green hornblend, zircon and sphene are observed. Calcium carbonates are more abundant in Layer IId (~22% in volume) than in the other layers of Complex II (~5 to 10%) (Fig 5C), but the limestone clasts are more frequent in the upper part (IIc). Halite and sylvite occur in low quantities (~2%) in all layers. Calcium phosphates (carbonate hydroxyl-apatite, whitlockite and dittmarite) are present at the bottom (Layers IIe, IIf) and at the top (Layer IIc) of Complex II. The alternating presence of gypsum and calcium oxalates (weddellite and whewellite) divides the infilling into two parts below and above Z~150 cm, i.e. from IIf to middle of IIc and from middle of IIc to IIa (Fig 6, S1 Table).

Organic matter is present in all layers (though it was macroscopically observed only in some of them) (Fig 5D) and its percentage (~10% in weight) remains constant through Complex II up to Z~150 cm. In the upper part of Complex II (above Z~150 cm, i.e. top of Layer IIc to Layer IIa), the organic content is significantly higher, up to 34%. This is likely because microfaunal remains and seeds are more frequent than in the lower part (Fig 5D). The top of Layer IIc (sample S10) in particular corresponds essentially to a dark brown accumulation of biogenic elements including fragments of insect carapaces and tiny balls that can be attributed to bat metabolic wastes. Throughout the stratigraphy, the pH is neutral or slightly basic (pH between 7 and 7.5) but becomes slightly acidic (pH 6.2) at the top of Complex II, above Z~150 cm (Fig 5E). The magnetic susceptibility values (Fig 5F) are nearly constant through Complex II, slightly decreasing in its upper layers, above Z~150 cm (top of IIc and IIb).

Complex I. Layer Ih differs markedly from the other layers of Complex I and is described separately. The other Complex I layers are characterized by moderately well sorted silts and fine sands (Fig 5A). The sediments consist mainly of ashes and become richer in microfaunal bones in the upper levels. The mineralogical content (quartz, feldspar and heavy minerals) and the quartz grain morphoscopy are close to, but more variable than, those of Complex II. Calcite is present in all these layers and is particularly abundant in Layers Ig to If (31 to 76%, S1 Table). Evaporites are present in the form of gypsum (only in Ie), anhydrite (in Ie, Id, Ia3), halite and sylvite (in all these layers). Calcium phosphates are found over the whole Complex I except dittmarite, which is present only in If and Ih.

The dark grey Layer Ih (sample S8) shows significantly high organic matter content (~75% in weight) associated with abundant charcoal fragments (< 1 mm), burned insect carapaces, and microfaunal remains (Fig 5D). The sorting of silts and fine sand in Ih is better than that of other layers in Complex I. In this layer, the quartz grains are sub-rounded and lusterless (Fig 5B). Moreover, Ih differs by its paucity in calcite, and the presence of dittmarite and Ca-oxalates, both nearly absent in the other layers of Complex I.

The pH is neutral but slightly acidic in Layer Ih (6.7) and slightly basic in Layer Ig (7.25) and If (7.8). The magnetic susceptibility values are more variable in Complex I than in Complex II (Fig 5F), ranging from 37 to $320 \ 10^{-8} \ m^3/kg$.

4.2 Optically stimulated luminescence dating analyses

4.2.1 Dose rate stability over time and beta dose rates. In this section we first discuss the reliability of the Dr (i.e. whether the current Dr can be representative of the past mean Dr), based on the examination of the U, Th and K contents. Once we have established that the current Dr is meaningful for dating purposes, we discuss the accuracy of the beta Dr estimate, bearing in mind that the heterogeneity of the sediments at a millimetric scale must be considered (Appendix 3).

U, Th and K contents are presented in Table 2A, 2B and 2C for the fine (< 2 mm), coarse (2 mm - 1 cm) and total (< 1 cm) fractions, respectively. For the fine fraction, the most striking pattern is the high K content, ranging between 1.7 and 7.6% and reaching its highest values in the upper layers of Complex I. The K content can be related to the presence of orthoclase, sylvite and illite. Its contribution to the total Dr can be roughly estimated to range between 75 and 92%. It is therefore important to ensure that the K content has been stable over the burial time. We argue here that this was likely the case in Complex II. Firstly, because of the presence of soluble minerals (gypsum and sylvite), which suggests negligible percolation: water dripping, leading to the crystallization of gypsum, was quickly followed by evaporation. Sylvite crystallizes directly from bat urine and cannot be associated with waterlogged conditions. As the water content is currently low (0-6% in Table 3), it can be safely assumed that it was also the case during the deposition of the sediments coeval to the formation of the evaporites. Moreover, there is no strong evidence for diagenetic processes that would be expected if significant percolation had occurred in such an environment (e.g. [54]). A second argument for the stability of the K content over time is the nearly constant ratio of $K/^{232}$ Th and 226 Ra/ 232 Th, at elevation from 7 cm to 163 cm above datum (Fig 7): ²³²Th is not soluble while ⁴⁰K and part of the U chain are. The consistency of the ratio is likely not fortuitous and suggests that conditions were relatively similar during the period of sediment deposition into the cave. The change of this ratio above 163 cm is likely attributable either to a change in the nature of the sediments (Complex I versus Complex II) and/or to additional input of K and U. This is further discussed in section 4.2.3 and section 5.

It can also be noted that the U chain is slightly in disequilibrium (e.g., pre-²²⁶Ra/post-²²⁶Ra activities ratio from -39 to +31% of unity) (S2 Table). However, the impact of this disequilibrium

Table 2. U, Th and K contents.

PLOS ONE

а	sample GDB			²³⁸ U	(ppm)			²³² Th	(ppm)	K	(%)
		(23	⁴ Th)	(226	³ Ra)	(21	⁰ Pb)				
	9	1.28	±0.10	1.64	±0.03	1.17	±0.16	2.23	±0.09	7.595	±0.056
	8	0.68	±0.04	0.99	±0.01	1.09	±0.10	1.84	±0.06	2.523	±0.023
	7	1.91	±0.14	1.99	±0.04	1.72	±0.13	4.14	±0.20	3.527	±0.029
	6	2.26	±0.22	1.93	±0.05	1.66	±0.16	4.47	±0.27	3.065	±0.031
	5	2.25	±0.16	1.61	±0.03	1.27	±0.12	4.03	±0.19	2.747	±0.024
	4	2.06	±0.16	1.68	±0.03	1.17	±0.12	4.13	±0.20	2.913	±0.026
	3	1.40	±0.08	1.04	±0.01	0.89	±0.10	2.94	±0.11	2.011	±0.020
	2	1.30	±0.07	1.06	±0.01	0.81	±0.09	2.58	±0.08	1.700	±0.016
	1	1.07	±0.07	1.11	±0.02	1.02	±0.11	2.56	±0.10	1.756	±0.020
b	sample GDB			²³⁸ U	(ppm)			²³² Th	(ppm)	K	(%)
		(23	⁴ Th)	(226	³ Ra)	(21	⁰ Pb)				
	9	1.57	±0.11	1.36	±0.02	0.85	±0.22	1.49	±0.06	5.247	±0.067
	8	1.25	±0.04	nd		nd		2.05	±0.04	1.975	±0.101
	7	1.72	±0.09	nd		nd		1.86	±0.07	3.497	±0.178
	6	2.02	±0.12	1.47	±0.03	1.46	±0.22	1.65	±0.07	1.143	±0.021
	5	1.31	±0.15	0.80	±0.03	0.80	±0.29	1.14	±0.08	0.846	±0.022
	4	0.83	±0.02	nd		nd		0.50	±0.00	0.840	±0.043
	3	0.69	±0.01	nd		nd		1.52	±0.01	1.040	±0.053
	2	2.17	±0.14	1.25	±0.03	1.02	±0.28	1.32	±0.08	0.881	±0.022
	1	0.88	±0.10	0.63	±0.02	0.45	±0.20	1.66	±0.06	1.103	±0.022
С	sample GDB	mass pro	oportion of		²³⁸ U	(ppm)		²³² Th	(ppm)	K	(%)
		coarse	fraction	(23	⁴ Th)	(226	³ Ra)				
	9	2	2%	1.46	±0.11	1.57	±0.03	2.02	±0.07	6.944	±0.047
	8	1	6%	0.77	±0.03			1.87	±0.05	2.436	±0.025
	7	6	5%	1.88	±0.12			3.71	±0.16	3.521	±0.041
	6	1	3%	2.18	±0.19	1.84	±0.04	4.07	±0.24	2.750	±0.027
	5	1	3%	1.88	±0.13	1.29	±0.02	2.89	±0.12	1.994	±0.017
	4	2	1%	2.00	±0.15			3.97	±0.19	2.821	±0.025
	3	7	7%	1.35	±0.08			2.84	±0.10	1.942	±0.019
	2	1	2%	1.41	±0.07	1.08	±0.01	2.42	±0.07	1.598	±0.014
	1	2	0%	1.01	±0.05	0.96	±0.01	2.28	±0.08	1.547	±0.015

a) U, Th and K contents for the < 2 mm fraction deduced from high resolution gamma spectrometry. The U contents are deduced from the activity of different daughters in the decay chain: mainly ²³⁴Th, ²¹⁴ Pb and ²¹⁴Bi in equilibrium with ²²⁶Ra, and ²¹⁰Pb. b) U, Th and K contents for the 2 mm-1 cm fraction deduced from high resolution gamma spectrometry (GDB1, 2, 5, 6, 9) or ICP-MS or AES (GDB3, 4, 7, 8). c) Mean U, Th and K contents calculated from a and b, weighted by the mass proportion of the < 2 mm and 2 mm-1 cm fractions. The proportion of the 2 mm– 1 cm fraction per sample is indicated (i.e. mass of 2 mm-1 cm fraction on total dry mass of material < 1 cm) These content values represent the contents that would have been obtained if the whole < 1 cm fraction had been crushed, homogenized and analyzed.

doi:10.1371/journal.pone.0169418.t002

might have on the total Dr over time is almost negligible (U contribution 4-13% compared to the contribution of the K (75-92%) to the total Dr).

The U, Th and K contents in the coarse fraction are significantly different from those of the fine fraction (e.g., K content for the coarse fraction of GDB5 is 69% lower than for the fine fraction—Table 2B). The beta Dr were calculated separately from the mean U, Th and K contents of the fine material (contents in Table 2A and dose rate beta-1 in Table 3) and from the whole material < 1 cm (contents in Table 2C and dose rates beta-2 in Table 3). It is expected



Table 3. Dose rates for each sample.

sample	Complex	current						c	lose rat	te (Gy/ka	I)					
GDB		water content	ga	mma	cos	mics	be	ta-1	tot	al -1	be	ta-2	tot	al -2	total-	mean
9	Complex I	6%	0.814	±0.087	0.088	±0.011	5.350	±0.691	6.252	±0.697	5.000	±0.589	5.902	±0.594	6.077	±0.639
8		0%	1.072	±0.123	0.088	±0.011	1.962	±0.249	3.122	±0.278	1.913	±0.216	3.073	±0.243	3.098	±0.259
7		3%	0.704	±0.078	0.088	±0.011	2.783	±0.340	3.575	±0.349	2.778	±0.305	3.570	±0.314	3.572	±0.330
6	Complex II	2%	0.748	±0.083	0.088	±0.011	2.492	±0.300	3.328	±0.312	2.299	±0.245	3.135	±0.257	3.231	±0.280
5	upper sub- unit	5%	0.527	±0.057	0.088	±0.011	2.162	±0.258	2.777	±0.265	1.979	±0.211	2.594	±0.218	2.686	±0.238
4	Complex II	4%	0.527	±0.057	0.088	±0.011	2.301	±0.277	2.916	±0.283	2.228	±0.241	2.843	±0.247	2.879	±0.263
3	lower sub-	4%	0.588	±0.149	0.088	±0.011	1.593	±0.192	2.269	±0.243	1.538	±0.167	2.214	±0.222	2.241	±0.232
2	unit	4%	0.507	±0.055	0.088	±0.011	1.356	±0.162	1.951	±0.171	1.289	±0.136	1.884	±0.146	1.918	±0.157
1		6%	0.535	±0.057	0.088	±0.011	1.354	±0.162	1.977	±0.172	1.255	±0.135	1.878	±0.145	1.927	±0.157

The uncertainties include both statistic and systematic errors. Beta-1 and total-1 refer to the beta and total dose rates calculated from the U, Th and K contents of the < 2 mm material, and -2 refers to the beta and total dose rates calculated from the whole sediment U, Th, K contents (excluding > 1 cm material). Their mean has been used for the final age calculations.

doi:10.1371/journal.pone.0169418.t003

on theoretical grounds that the true beta Dr falls between these two estimates. In order to confirm this and evaluate which beta Dr was closest to the true beta Dr, simulations were carried out (Appendix 3). It was observed that the true beta Dr always ranges between the beta-1 Dr and the beta-2 Dr in the different cases that were investigated. Consequently, a mean value should be the best option to estimate the beta Dr for the samples. However, the calculations





doi:10.1371/journal.pone.0169418.g007

show that for these samples, the fine (beta-1 and total-1) and fine plus coarse (beta-2 and total-2) Dr are relatively close to each other (ratio of total-1 to total-2 < 1.07) and statistically indistinguishable. Therefore, the correction calculated by this approach remains small for the Goda Buticha samples: the large differences in radio-isotope content is counterbalanced by the low relative weight of the coarse fraction (4 to 22% of the total, Table 2C) and the contribution of other components of the Dr (gamma and cosmics).

The total Dr ranges between 1.92 ± 0.16 Gy/ka to 3.57 ± 0.33 Gy/ka for GBD1 to 8 and up to 6.08 ± 0.64 Gy/ka for GDB9 (Table 3).

4.2.2. Equivalent dose distributions. The reliability of the SAR protocol for the GDB samples in this study was tested on multi-grain aliquots with dose recovery tests and preheat plateau tests. Dose recovery tests were also conducted on single grains, prior to final single-grain De measurements that were used for age determination. The tests and their results are described in Appendix 2.

Single-grain D_e distributions are plotted in Fig.8. The central D_e (after the central age model—CAM—of [55]) increases significantly from top to bottom, from 3.8 ± 0.2 Gy to 112.9 ± 5.3 Gy (Table 4), and the over-dispersions (OD—i.e. dispersion of the De beyond the statistical dispersion expected in a lognormal distribution) increase with De, from $20 \pm 2\%$ to $41 \pm 4\%$ except for sample GDB9 ($30 \pm 3\%$) and 5 ($49 \pm 3\%$) (Fig.9).

The interpretation of the De distributions, which reflect the pre- and post-deposition history of the quartz grains, is not straightforward, the main question being whether the distribution is consistent with a single "mean" age or presents multiple components, due to poor bleaching or bioturbation. In Appendix 2 we suggest that–based on the geological data, on the comparison between single and multi-grain De_s , and on the comparison of the natural and artificial (beta or gamma dose given in the laboratory) OD–the De distributions in this study are consistent with single dose components.

For the final calculation of the relevant "mean" De from each De distribution, several models are currently discussed in the literature: the widely used central age model (CAM) of Galbraith et al. [55], the recently developed Bayesian model of Combès et al. [57] and the simple, non-weighted arithmetic mean advocated by Guérin et al. [56]. In Table 4, the three estimates are indicated for each sample. Presentation of a detailed comparison of the three models is beyond the scope of this paper. It can simply be noted that they are statistically indistinguishable for the youngest samples (GDB 9, 8, 7 and 6). For the older samples, the variations between the estimates increase slightly, but the Bayesian estimate is in most cases consistent at one or two sigma with the CAM or arithmetic mean (except for GDB5 when compared to the unweighted mean, and for GDB4 when compared to the CAM, though in this last case, the discrepancy is low- <6%). Since the Bayesian SAR model is statistically more robust and allows a better treatment of the error estimates [57], it has been used for the age calculations and comparison with the ¹⁴C data. The Bayesian De range between 4.0 ± 0.2 Gy and 121.9 ± 5.1 Gy.

4.2.3. OSL ages and comparison with the ¹⁴C **chronology.** The OSL age estimates (Table 5) are shown in Fig 10 together with the ¹⁴C calibrated ages (Table 1). The OSL ages extend the chronostratigraphy of the sequence to the start of MIS 3 or end of MIS 4 (63 ± 7 ka). The age of 43 ± 5 ka for sample GDB3 is consistent with the two ¹⁴C ages, though those were in apparent chronostratigraphic reversal. The chronological gap within Complex II at ~105 cm above datum that was suggested by the ¹⁴C dates is confirmed by the OSL results, albeit it is now of a shorter duration (between 24.8 ± 2.6 ka–OSL—and 7.5 ka cal BP - ¹⁴C). The age for sample GDB5 at the base of Layer IIa of Complex II (6.9 ± 0.7 ka) is also in agreement with the ¹⁴C date. The general agreement between the ¹⁴C and OSL data for Complex II, from Layer IIf up to the top of Layer IIc, argues both against any post-depositional mixing and against any Dr instability.



Fig 8. Radial plots for the De distributions of each sample. The number of measured and accepted grains, central De (CAM) and over-dispersion can be found in Table 4. The over-dispersion due to the measurement, estimated to be at least 19% has been added quadratically to the relative uncertainty. The shaded region emphases the central De ± 2 sigma.

doi:10.1371/journal.pone.0169418.g008

For the top of Complex II (sample GDB6) and all of Complex I, the two chronologies differ. The OSL ages of 3.3 ± 0.4 and 0.7 ± 0.1 ka are younger than the ¹⁴C ages of 4680 ± 81 cal BP and 1485 ± 75 cal BP, respectively. A possible explanation would be a late input of potassium in bat urine from the current surface. In such a case, the current Dr would be overestimated compared to the past mean Dr, leading to underestimate of the calculated OSL ages. For GDB6, though the K/²³²Th ratio is similar to those for Complex II, suggesting beta Dr stability, the proximity (< 30 cm) to the upper most layers, receiving a late input of K, would imply increased gamma Dr.

5-Discussion

5.1 Interpretations of the sedimentological data

Analyses of the sediments provide information about the mode of deposition and post-depositional evolution of sediments in Goda Buticha. There is no clear indication of an erosional surface where the chronostratigraphic hiatus appears, suggesting an absence of deposition rather than post-depositional erosion. The consistency of the magnetic susceptibility from the bottom of Complex II up to its upper part suggests the absence of pedogenesis, as do the absence of pedofeatures such as CaCO₃ redistribution, clay illuviation, organic matter concentration or pH variations.

The consistency of the magnetic susceptibility from the bottom of Complex II up to its upper part also suggests a constant main source for the sediments, and a deposition under highly similar conditions, irrespective of the significant chronological hiatus in the sediment sequence. This is further supported by the other analyses. Over all, the sedimentological analyses show the association of four modes of sedimentation:

(i) A ubiquitous allochtonous geogenic sedimentation. Well sorted silts and the dominance of sub-rounded and lusterless quartz grains are attributed to aeolian dusts. The low but omnipresent clay concentrations could indicate the persistence of run-off inputs into the cave, though they could also be due to *in situ* alteration of feldspars. The mineralogical associations

sample GDB	preheat param	neters	n	0\	/er			De	(Gy)		
	natural and regenerative dose	test dose		dispe (?	ersion %)	CA	М	unwe arithme	ighted tic mean	bayesiar	central
9	260°C 10s	160 cut	55/200	30	±3	3.8	±0.2	4.0	±0.2	4.0	±0.2
8	260°C 10s	240 cut	100/400	20	±2	5.9	±0.1	6.0	±0.1	5.9	±0.1
7	260°C 10s	220 10s	47/400	21	±3	8.5	±0.3	8.7	±0.3	8.7	±0.3
6	260°C 10s	220 10s	38/300	26	±4	10.7	±0.5	11.4	±0.6	10.8	±0.5
5	260°C 10s	220 10s	188/1100	49	±3	19.2	±0.7	22.7	±1.0	18.4	±0.6
4	260°C 10s	220 10s	168/800	32	±3	67.0	±2.0	77.5	±2.6	71.3	±1.7
3	260°C 10s	160 cut	100/700	39	±4	88.4	±4.1	104	±4.7	96.6	±4.1
1	260°C 10s	160 cut	106/700	41	±4	112.9	±5.3	134	±7.3	121.9	±5.1

Table 4. Results of the De analyses.

The uncertainties include the counting uncertainty and the systematic uncertainty due to the reproducibility of the illumination. The number "n" of accepted/ measured single grains is indicated. The De are calculated using the CAM [55], or unweighted mean [56] or Bayesian central [57] models.

doi:10.1371/journal.pone.0169418.t004



Fig 9. Over-dispersion in function of the dose. Black dots: beta dose given inside the reader (dose recovery test); white dot: gamma dose given outside the reader; grey dots: natural dose. A longer explanation of the OD of the beta and gamma dosed samples is given in Appendix 2.

doi:10.1371/journal.pone.0169418.g009

of this geogenic fraction of the sediments are characteristic of granitic or clastic sedimentary rocks: the magmatic minerals (silicates and oxides, including heavy minerals) could derive from erosion of Precambrian gneissic, granitic and migmatitic basement rocks that form the ridges above the tectonically downfaulted sedimentary sequence of the Dire Dawa area [58]. Another possible origin is the Triassic and Cretaceous sandstones, which are present in the Harrar plateau and exposed in the immediate vicinity of the site [59, 60]. These rocks have been continuously eroded by streams flowing from the plateau into the rift, accumulating thick alluvial fans and sheetwash deposits in the extended rift basin in front of the cave opening.

(ii) An autochthonous geogenic sedimentation, characterized by limestone clasts within Complex II layers and calcite matrix: the limestone in which the cave formed [61] is likely the local source for those.

(iii) An ubiquitous biogenic sediment composed of numerous micro-vertebrates, seeds and insect remains. In particular in Complex I, the anthropic and biologic activities could be the main reason for the moderate sorting of the sediments, due to trampling effects.

•									
sample GDB	Complex	Layer	Dose ra	te (Gy/ka)	(Gy/ka) De (C		Age	Age (ka)	
9	Complex I	le	6.077	±0.639	4.0	±0.2	0.7	±0.1	
8	_	la	3.098	±0.259	5.9	±0.1	1.9	±0.2	
7		lla	3.572	±0.330	8.7	±0.3	2.4	±0.3	
6	Complex II upper sub-unit	llb	3.231	±0.280	10.8	±0.5	3.3	±0.4	
5	_	llc	2.686	±0.238	18.4	±0.6	6.9	±0.7	
4	Complex II lower sub-unit	llb/llc	2.879	±0.263	71.3	±1.7	24.8	±2.6	
3		lle	2.241	±0.232	96.6	±4.1	43	±5	
2	_	lle/llf	1.918	±0.157	nd		nd		
1		llf	1.927	±0.157	121.9	±5.1	63	±7	

The ages are calculated using the Bayesian approach developed by Combès et al. (2015) (Table 4) and the dose rates presented on Table 3.

doi:10.1371/journal.pone.0169418.t005

Table 5. OSL ages.



Fig 10. Plots of age estimates as a function of the elevation within the stratigraphy. The intervals represent the confidence interval at 95%. Black: ¹⁴C data; grey: OSL data. The inset zooms in the < 10 ka data.

PLOS

ONE

(iv) A secondary chemical formation during and after deposition as attested by mineralogical associations through XRD analyses. The presence of calcium phosphates, evaporites (halite, sylvite and gypsum) and calcium oxalate minerals could be linked to the accumulation of bat guano and urine, which leached and reacted with the calcium carbonates (limestone of the cave) at relatively high temperatures and thus at high evaporation rates [62–64]. In particular, Ca-oxalates, which appear in Complex I, can be related to bat guano and to human-made fires that burned wood or other plants. It is likely that the chemically unstable Ca-oxalates formed in vegetal ashes (see for example [65]) produced calcite by CO_2 absorption (e.g. in Layers Ia). Evaporitic halite and sylvite are observed in all layers while gypsum, which is produced by the reaction of urine on limestone, is present in the lower part of Complex II up to $Z\sim150$ cm. In the upper layers of Complex I, gypsum is replaced by anhydrite, another likely consequence of fires in the cave: heat leads to dehydration of gypsum or to evaporation of saline solutions such as bat urines and thus to the formation of anhydrite [63].

In summary, the sedimentation in Goda Buticha cave was mainly induced by aeolian action in both Complex II and Complex I, with a small contribution of run-off. This is consistent with the absence of typical features or morphologies that could be linked to sedimentary dynamics of fluvial origin, like cross beddings, positive grading, laminations, channels and erosive contact. The geological analyses show that the origin and mode of deposition before and after the chronological gap are identical but, in Complex I, anthropic and faunal activities are higher.

It is possible to interpret the results of the geological analyses in terms of environmental conditions in the cave's environments during the time of deposition. The conservation of very soluble minerals (such as gypsum, halite, and sylvite) in the lower part of the sequence (Layer

IId-IIf) is a good indicator of negligible water circulation (at least vertically) into the sediments, and indicate a local low-moisture regime during and after the sedimentation of the lower part of the sequence. Moreover, the small sample of identifiable fauna contains some open woodland taxa suggesting ecotone conditions including a source of water that could support grasslands.

Regional climate records display a complex picture, though there are few relevant records for the MIS 3 and MIS 2 periods. The period 50–15 ka is poorly represented in the paleoclimate literature of the region. Those few paleoclimatic sequences that are available are mostly from the Northwest Ethiopian Highlands, which belong to a different climate regime (different moisture source), making comparison with the Goda Buticha site complicated. Better records are available for the African Humid Period. Though some earlier ¹⁴C dated records from Lake Abhe (e.g., [66, 67]) indicate relatively moist conditions during the period 50–31 ka, recent, well-constrained and higher resolution data from the Ethiopian rift suggest intermediate moisture conditions interrupted by drier periods between ~50–35 ka, more pronounced aridity between 35–19 ka, and increasingly moist conditions since then, especially during the African Humid Period (~ 15–5 ka BP), and finally arid conditions during the Late Holocene [68, 69]. Stable isotope records on speleothems from the Mechara area, some 150 km to the southwest of Goda Buticha, show warm and wet conditions in the SE Ethiopian plateau during the Early to Middle Holocene followed by dry conditions during the Late Holocene [70–72].

Therefore, the chronostratigraphic hiatus at Goda Buticha coincides temporally first with the main part of the dry period that started at around 35 ka, then with the African Humid Period and the Early Holocene wet phase. The aeolian activity, which was the main transport agent throughout the sequence, apparently occurred during both wet and dry phases. It is therefore likely that the sedimentological parameters at Goda Buticha did not linearly or immediately respond to the regional climatic shifts. In particular, the regional climate record suggests that the aeolian deposition does not necessarily correspond to the driest phases.

5.2 Implications for human occupation

The association in Goda Buticha of late MSA technological characteristics with LSA components during the mid-Holocene is unusual in the Horn of Africa [30]. However, it has already been described for Gobedra [73] and for the lower units of Laas Geel shelter 7 [74], dated to the Terminal Pleistocene. The new chronological and sedimentological data argue against post-depositional mixing as an explanation for the presence of MSA characteristics in the upper levels, where LSA technological characteristics become more frequent. The data confirm that in Goda Buticha the Holocene MSA characteristics post-date a gap of some 17,500 years (from ~25 to ~ 7.5 ka) in the depositional sequence.

The earlier part of the temporal gap in the Goda Buticha sequence coincides with MIS 2, which was arid [68, 69]. Continuity of human occupation may not have been possible in this area during that time. As no archaeological data are currently available in the Horn of Africa for the period corresponding to MIS 2, local persistence of technological traditions currently seems an unlikely explanation for the Holocene MSA traits in the upper part of the sequence [31]. Reintroduction of cultural traits or technological convergence are possible alternative explanations for the MSA-like technical features of Layer IIc, but they cannot be tested regionally [75].

Only a few sites in the Horn of Africa have yielded contemporaneous Late Pleistocene (Mochena Borago, Porc-Epic or Ziway-Shala Basin) and/or Holocene occupations (Mochena Borago, Ziway-Shala Basin). The emerging regional variability of the lithic assemblages, with shared technical components as well as specificities, may derive from differences in their specific geographic locations and local environmental conditions, influences of resource

availability, subsistence strategies, and mobility patterns. Regional-scale population movements within Africa (from possible refugia areas) also may have acted as a trigger for converging and/or recurring technological behaviours leading to mosaic patterns of cultural variability in the region [31, 69, 76].

6- Conclusion

The OSL dating of the whole sequence of Goda Buticha confirms and refines the previous chronostratigraphy that was based on C14 dating [30]. In particular, the results highlight a chronological and sedimentological hiatus from ca 25 to 7.5 ka cal BP. However, the depositional hiatus is hardly visible through the sedimentological analyses, except for an increase in limestone fragments after the gap. Aeolian transport has remain the main transport agent, irrespective of regional climate changes, while the sedimentary hiatus, was coeval first to very arid conditions, then to wet phases. The relationship between the sedimentary sequence and regional / local climate variability remains to be studied in detail through on-site proxies for climate change, for example micromorphological work and a systematic study of the ubiquitous micromammals in the sequence.

Our current results validate previous observations that an assemblage with both MSA and LSA technological features post-dates the hiatus. These results bear on the high technical variability of the Late Pleistocene and Holocene in the Horn of Africa and emphasize the complexity of the shift from industries labelled "Middle Stone Age" to those labelled "Late Stone Age".

Appendix 1: protocols for geological analyses

Each sediment sample was split into several aliquots. After dry sieving at 2 mm, two $\sim 8 \text{ cm}^3$ aliquots were retained for subsequent analyses. The first aliquot was decarbonated with 10% diluted hydrochloric acid. At the end of the reaction, the sample was washed and centrifuged several times to eliminate the acidic solution. It was then mixed in distilled water by magnetic stirring for 4 hours. In the final stage of the stirring, and without stopping, a small volume of sediment was sampled using a needle with a 2 mm opening and analyzed with a magnetometer. The magnetic susceptibility of each sample was determined at low frequency (0.47 kHz) using the Bartington Susceptibility Meter model (MS2B). Air measurements were made before and after each sediment sample analysis. Results presented in Fig 5F are the average of 3 successive measurements.

The grain-size distribution was determined on portions of the first aliquot before and after the decarbonation. It was measured with a laser Mastersizer 2000 granulometer, which estimates the mean diameters of the particles within a size range of 0.02 to 2000 μ m.

Another portion of the decarbonated aliquot was used to measure organic matter. It was weighted, carbonized continuously in a muffle furnace at 375°C for 16 hours and weighted again. The loss of weight is attributed to the burning of the organic matter.

The pH value was determined on the second aliquot. Twenty ml of sediment were mixed with 20 ml of distilled water and stirred for four hours. Then the pH of the sediment was measured by plunging a pH-meter probe (Hanna Instruments 211) into the mixture. Subsequently, the second aliquot was decarbonated and rinsed several times. Wet sieving was then performed in order to collect the 50–200 μ m grain size fraction. This fraction was dried in the oven at 40°C and weighed. Heavy and light minerals were separated using bromoform at 2.89 g/cm³ following the Duplaix method [77]. Each fraction was weighed. The heavy minerals were put into Canada balsam and the mineralogy as well as number of grains (>100 grains) were determined with a polarized microscope. The light fraction was used for the analysis of the quartz grain morphoscopy. The morphoscopic class was determined from the degree of rounding and the surface aspect of at least 100 grains.

Mineralogical analyses were performed by XRD on powders of bulk samples. Analysis was conducted at the Department of "Histoire de la Terre" of the MNHN (UMR 7207 CR2P) on a D2 Phaser Bruker diffractometer equipped with a copper anode source (Cu K α radiation). After the mineralogical identification, semi-quantitative estimates of mineral percentages were calculated with Diffrac.Eva software.

Appendix 2: OSL, protocols and comments

Equivalent Doses

Sample preparation. Samples were dried and sieved with a 2 mm mesh. The < 2 mm fraction was then split into two parts. One was used for the determination of the U, Th, and K content, the second was used for the De determination. For this last one, extraction of 200–250 µm quartz grains was performed following a standard procedure: wet sieving, HCl 10% then H₂O₂ 30% for removal of carbonates and organic materials respectively, high density (2.58 and 2.72) solution of sodium heteropolytungstate for separation of quartz, feldspars and heavy minerals, and HF 40% etching for 60 min (followed by HCl) to remove the matter irradiated by alpha particles. Finally, an IRSL test [78] was performed to confirm the absence of any feldspar contamination. The quartz grains were mounted either on stainless steel discs with silicon oil (1 mm mask) or on single grain discs with 100 holes of 300 µm diameter and 300 µm depth each.

Measurements. All luminescence measurements were performed with a Risø TL/OSL-DA 15 reader equipped with a 10 mW Nd:YVO₄ diode-pumped laser (532 nm) for single grain excitation [79–81] and with clusters of blue (470 ± 20 nm) or infra-red (870 ± 30 nm) LEDs for multi-grain excitation. The reproducibility of the green laser excitation was estimated to be 1.9 to 2.7% and the reproducibility of the blue LED was 1.5%. Detection in the UV range (280–380 nm) was performed with a Q9235 Photomultiplier tube preceded by 7.5 mm of Hoya U340 filter. Irradiations were performed with a 90 Sr/ 90 Y beta source delivering 0.16 Gy/s to the 200–250 µm quartz grains.

Protocol and analyses. The single-grain SAR protocol is presented in Fig 11. Analyses were performed with Analyst V.3.24 [82]. Exponential plus linear functions were used for curve fitting. For multi-grain measurements, the blue LEDs were stimulated for 40s; the first 0.8 s and last 4.16 s were used for signal and background respectively. Similarly the first 0.06 s and last 0.12 s (of 1 s of stimulation) were used for signal and background in single grain measurements, respectively.

The quartz samples from Buticha are remarkably bright and the single grain output is rather good (> 10% of the measured grains can be selected for the De determination). An example of glow and growth curve is presented in S2 Fig. It could be verified that the multi-grain signal is dominated by the fast component (Fig 12).

Testing the protocol. The appropriateness of the parameters for the SAR protocol were checked: dose recovery tests and preheat plateau tests were conducted. For the dose recovery test, three multi-grain aliquots of each sample were bleached (1 hour in a solar simulator), given a known dose close to the grossly estimated equivalent dose (De) and the SAR protocol was applied. The unweighted arithmetic means are compared to the expected dose (S3 Fig). The dose recovery ratios were found to be consistent with unity at two sigma or within $\pm 10\%$ of unity for a test dose preheat of 220°C for 10 s for GDB 4, 5, 6 and 7, 160°C cut-heat for GDB 1, 3 and 9 and 240°C cut-heat for GDB8 (Table 6A). GDB2 was not tested any further.

A second set of dose recovery tests was then performed with single grain discs using the parameters previously selected. Rejection criteria, based on the recycling ratio (within 0.9–1.1), signal to noise ratio (> 3), precision of the natural test dose signal (RSD > 10%), and



Fig 11. SAR protocol used for the single-grain De determination. The preheat parameters were fixed for each sample according to the results of the dose recovery tests (Table 6). For the multi-grain De determination, the protocol is the same, but for the excitation, performed with blue LED for 40 s.

doi:10.1371/journal.pone.0169418.g011

recuperation (sensitivity corrected signal for a 0 dose > 5% of the sensitivity corrected natural signal) were used. The grains were also rejected when there was no intercept of the natural signal with the growth curve, or in case of intercept with infinite error estimate (saturation). Then, the dose was estimated either using an unweighted mean [56], or the central dose model (CAM, [55]), or the Bayesian SAR model [57]. The efficiency of the SAR parameters was confirmed (Table 6A).

Preheat plateau tests were performed using multi-grain discs for the samples whose De distributions were the least dispersed, i.e. GDB 4, 6, 7, 8 and 9. The preheat of the natural or regenerative dose was set at 220, 240, 260 or 280°C for 10 s (4 to 6 discs for each temperature) while the preheat of the test dose was chosen according to the results of the dose recovery tests. No significant dependence of the De on the preheat temperature was observable (S4 Fig). Therefore, we used a 260°C preheat for De determination for each of the GBD samples in this study (Table 4).

Are the De distributions consistent with a single dose component?. The sedimentological analyses suggest that the fine sands and silts, especially those of Complex II, have an aeolian origin (see §5.1). Moreover, the samples were collected close to the cave entrance where the sediments are exposed to sunlight. The quartz grains are clean, i.e. there is no embedded crust that would prevent the light from going through. Therefore, poor bleaching seems unlikely. As for the likelihood of bioturbation, burrows were observed in the section but the OSL sampling was cautiously conducted away from those. Burrowing by insects, however, might be more difficult to detect.

Some researchers have also suggested that the consistency between the multi-grain De and the single-grain De could be used as an evidence for single component (e.g. [83]). The multi-grain CAM De (mostly from the preheat plateau tests) (<u>S3 Table</u>) and the single-grain CAM



Fig 12. LM-OSL curves. For each sample a regenerative dose was given to a multi-grain aliquot. After a 260°C for 10 s preheat, the intensity of the blue LED was increased linearly from 0 to 50 mW/cm² for 500 s. The curves are compared to the LM-OSL glow curve of the Risoe calibration quartz (in red), which is known to be dominated by a fast component.

PLOS ONE

De (Table 4) are consistent at one or two sigma for all samples except for GDB9 (3 sigma) and GDB1. For GDB1 and GDB3 however, the number of multi-grain measurements is low (n = 6 and 5 respectively) and the calculated CAM De_s might not be significant because of the scatter in these samples.

Besides these observations, the OD is considered as a key parameter for interpreting the De distribution, as it should reflect the story of the sample: it depends on intrinsic factors (e.g., how well each grain responds to the protocol) as well as extrinsic factors (bleaching before and bioturbation after deposition, as well as heterogeneity of the Dr at the millimeter/micrometer scale). However, there is no typical value of OD for well-bleached undisturbed samples (e.g., [84, 85]). More particularly, it has been shown from laboratory-dosed samples that the OD 1) increases with the central (or mean) De, 2) is higher for samples dosed outside the OSL reader than for those directly dosed inside it [85, 86]. It was also shown with modeling and simulations that 3) the OD increases when–at constant mean radioactive content–the proportion of radioactive hotspots (e.g., coarse potassium feldspar grains) increases. High heterogeneity (i.e. high proportion of hotspots) generally occurs with low radioactive contents (e.g. a few feldspar

в																
irradiation type	sample GDB	preheat paramete	ers for	given		multigra	Ë					single grai	L			
		natural and regenerative dose	test dose	dose (Gy)	c	Ta	ţi	c	ove disper (%	er sion)	ratio-	CAM	ratio- unv me	veighted an	ratio -b <i>e</i> estim	iyesian ates
beta	6	260 10s	160 cut	9	e	1.01	±0.02	91/200	1	Ŧ	1.04	±0.01	1.04	±0.02	1.07	±0.01
	8	260 10s	240 cut	9	e	1.00	±0.02	pu	pu		pu		pu		pu	
	7	260 10s	220 10s	6	e	0.98	±0.02	26/200	8	±2	1.04	±0.02	1.07	±0.03	1.08	±0.03
	9	260 10s	220 10s	6	ю	0.99	±0.03	60/300	6	±2	1.00	±0.02	1.01	±0.02	1.06	±0.02
	ъ	260 10s	220 10s	18	ო	0.99	±0.02	64/300	14	±2	0.99	±0.02	1.01	±0.02	1.05	±0.02
	4	260 10s	220 10s	55	ო	0.98	±0.03	54/200	16	±3	0.95	±0.03	1.00	±0.03	0.98	±0.03
	ო	260 10s	160 cut	74	ო	0.96	±0.04	44/200	20	±3	0.95	±0.04	1.03	±0.06	1.03	±0.04
	-	260 10s	160 cut	111	ო	0.96	±0.05	25/300	22	±5	1.01	±0.06	1.22	±0.16	1.04	±0.07
q																
gamma	8	260 10s	240 cut	6				228/700	19	Ŧ						
a) dose recovery	r tests: the samp	oles were bleached in	a solar simu	lator and give	en a kn	own bet	a dose in	the OSL re	ader; b)	Minim	al intrinsi	c OD esti	imate: the	e sample	was blea	ched in
a solar simulator	and given a kno	own gamma dose out	side the OSL	- reader. The	numbe	er "n" of i	nulti-gra	in aliquots or	of sele	cted o	er meas	ured sing	gle grain i	s indicate	d, as we	ll as the

ven doses.
oratory gi
ts for labo
of the tes
Results
Table 6.

ratio of estimated to given beta dose. The ratio are given following the central De model (CAM) of [55], the Bayesian model of [57], or the arithmetic mean advocated by [56]. ົກ doi:10.1371/journal.pone.0169418.t006 . D

grains in a quartz sand dune) [84, 87]. The OD for the artificially dosed grains and most of the natural samples of Goda Buticha follow these patterns (Tables 4 and 6, and Fig 9). This statement, detailed in the following sentences (1-2-3), is based on the assumption that the OD of the different samples from Goda Buticha can be compared to each other. The samples have indeed similar geological origin, and similar LM-OSL glow curves. This assumption might have some limitations since, for example, different preheat parameters had be used (after dose recovery tests) and different artificial or natural doses are concerned. The patterns seem to be followed for three reasons: (1) The OD of the bleached and beta-dosed samples (dose recovery tests) increases from 8 ± 2 to $22 \pm 5\%$ when the given dose increases from 6 to 111 Gy. (2) One sample has been bleached and gamma dosed (GDB8). Its OD is $19 \pm 1\%$ for 9 Gy, to be compared to 10% for about 10 beta-derived Gy for other samples (Table 6B). This value gives the minimum OD to be expected for the natural samples with doses over 10 Gy. (3) The natural OD for GDB 6, 7 and 8 are only slightly higher (20 ± 2 to $26 \pm 4\%$), suggesting that the supplementary OD due to external factors is weak. This can be expected from the very high Dr (i.e., K content) for these samples (Table 3). It is therefore reasonable to consider that these De distributions are consistent with a single De component. The increase of OD for GDB 4, 3 and 1 could be related to the increase of the CAM De, though it is difficult to ascertain that no other factors interfere. Meanwhile, we have also used a single-dose model for these samples. GDB5 and 9 have higher OD compared to what could be expected from the mean or CAM De and Dr for other samples. Whether this is due to the fact that these samples are not fully comparable to the others (e.g., different preheat parameters were used) or due to beta Dr heterogeneity is not conclusive. The Dr modeling that has been performed on GDB5 (Appendix 3) cannot help to answer this question because the model used is not of sufficient accuracy for that purpose (for example, the possible heterogeneities within the fine matrix were not taken into account). Only the comparison with the ¹⁴C data suggests *a posteriori* that the hypothesis of a single component distribution is in fact reasonable, at least for GDB5.

Appendix 3: Further comments on the beta Dr

The calculation of the beta Dr is complex because all the sediment samples show heterogeneities at the millimeter scale, with a fine fraction (mainly < 1 mm) embedded in coarser material (bones, charcoals, calcareous crusts, different types of lithic fragments, etc.). Since the travel range for the beta particles is about 2 mm in rocks and sediments, the possible influence of this coarse material on the mean Dr received by the quartz grains requires special attention [51]. While the > 1 cm particles can be assumed to have no impact on the beta Dr because of their strong auto-absorption [87], particles in the thickness range between 2 mm and 1 cm require a separate analysis.

In the following discussion we must distinguish:

 D_{fine} : theoretical dose rate (i.e. energy received per unit of mass and time) in the quartz grains calculated from the radioisotope content of the fine (< 2 mm) fraction, following the infinite matrix assumption.

 D_{total} : theoretical dose rate in the quartz grains calculated from the radioisotope content of the fine and coarse fraction (< 1 cm) (which will be referred as total fraction), following the infinite matrix assumption.

 D_{grains} : the true dose rate in the quartz grains.

*Em*_{fine}: the energy emitted per unit of time and per unit of mass by the fine fraction.

 Em_{total} : the energy emitted per unit of time and per unit of mass by the total fraction, equivalent to the infinite matrix dose rate [88] of these mixed fractions.

 D_{fine} and D_{total} are calculated respectively using the equations:

$$D_{fine} = Em_{fine} * F_{size} * WF_{fine}$$

 $D_{total} = Em_{total} * F_{size} * WF_{total}$

where F_{size} is an attenuation factor that depends on the grain size and has been estimated by Guérin et al. (48) and WF_{fine} and WF_{total} are the attenuation factors due to the dose rate part absorbed by the water content in the fine and total fractions respectively, (as defined by [89]).

As discussed in the main text, the coarse fraction (2 mm– 1 cm) generally has lower radioisotope content than the fine matrix (< 2 mm) (Table 2A and 2B). Therefore, the infinite matrix assumption should not apply. How far from the beta D_{fine} or D_{total} is the true beta D_{grains} ? Two effects have to be considered, i.e., auto-absorption (or size of the emitter) and dilution (or distance between emitter and dosimeter):

- -1). the amount of energy absorbed in the coarse fraction is more important than the energy emitted within it [90]; this implies that the quartz grains in the matrix receive a lower amount of energy than if the fine matrix only was present, i.e. $D_{fine} > D_{grains}$
- -2). only a part of the quartz grains are affected by the presence of the coarse material as many are actually far from them. Using the D_{total} instead of D_{grains} leads to dilution of the radioisotope content of the fine fraction and affects (excessively) all the quartz grains; this implies $D_{total} < D_{grains}$.

Therefore, D_{grains} is bracketed by D_{fine} and D_{total} .

The calculation of D_{fine} and D_{total} for the samples shows that they are actually close to each other, despite the large difference in radioisotope contents. We have tested the predominance of one or the other of the above mentioned effects on the beta dose rate with a Geant4 simulation [51, 53, 91-93]. Sample GDB5 was chosen since it presents the largest difference between D_{fine} and D_{total} . The fine fraction was modelled after its grain size histogram, chemical content and radioisotope contents; the coarse material was modelled considering 6.5% in weight of bones and 93.5% of stones, following weighting and sorting after dry sieving. The chemical composition of the bones was assumed and the chemical composition of the rocks was estimated using EDX-SEM analyses. Materials > 1 cm were not modelled since auto-absorption was assumed to dominate [90]. The different objects are represented by spheres of different sizes and chemical compositions, randomly located in a homogeneous medium, that represents the major part of the fine fraction (i.e. the matrix): the coarse fraction is represented by spheres of rocks of 5 mm diameter and spheres of bone material of 1.7 mm diameter (presenting the same beta auto-absorption than a 2 cm long bone) (Fig 13). Ten percent of the grains of the fine fraction between 100 and 300 µm diameter were also set as quartz spheres in the matrix. The remaining 90% of grains were not represented, as they have been taken into account in the matrix material composition. The quartz grains in the fine fraction are small enough in regards to the beta particle range that the effect on their Dr of the neighboring quartz grains can be ignored. The current water content of GDB5 is 6% (water/dry sediment). We have considered 3 values of moisture: 0% for a dry sediment, 6% in mass if the water is equally distributed in the fine and coarse fractions, 7.3% in mass if the water is in the fine fraction only.

The emission of beta particles with spectra of 40 K, U series and Th series from the coarse fraction or from fine fraction were simulated based on the measurement of U, Th, K contents in these fractions. The dose received in the quartz grains was recorded. Fig 14 presents the results for the 209 μ m grain size. A comparison is made between the effective beta dose rates in



Fig 13. representation of the modelled sediment. The 5 mm diameter green spheres represent rock fragments, the 1.7 mm yellow spheres represent bones (with an identical auto-absorption of 2 cm long bones), the 100 to 300 μ m red spheres represent a fraction of the quartz grains that are used as dosimeters, and the stacking box is a cube with a side length of 19.92 mm.

the grains (D_{grains}) in different conditions (with respect to radioactive element and moisture) and the dose rate calculated using the standard method applied to the fine or total fraction $(D_{fine} \text{ and } D_{total})$. The dose rates are normalized to D_{grains} in order to simplify the reading of the graph and to be independent of the simulation time.

Note that neither D_{fine} nor D_{total} seems to predominate in the effective D_{grains} , whatever the moisture and the radioactive element considered. Note also that the merit of this modelling is to allow a relative comparison of the various estimates of the beta dose rate, although their simulated values cannot be directly used for the age calculation due to the rather rough representation of the model.



Fig 14. Comparison of dose rates from the simulation and from the infinite matrix assumption. White bars: D_{fine}/D_{grains} ; grey bars: D_{tota}/D_{grain} ; black bars: D_{grains}/D_{grains} (i.e. unity).

doi:10.1371/journal.pone.0169418.g014

Supporting Information

S1 Table. Semi-quantitative results of XRD analyses for the Goda Buticha sediment sam-

ples. –: absent; ++: $1\% < x \le 5\%$; +: $\le 1\%$. The percentages are relative to the peaks heights in the diffractogram.

(PDF)

S2 Table. Activities for the < 2 mm fraction. The ratios allowing comparison of the activities of the top, middle and bottom of the ²³⁸U decay chain are indicated. (PDF)

S3 Table. Equivalents doses for the multi-grain aliquots. The test dose preheats for GDB8 are 20°C lower than the corresponding regenerative dose preheat. N corresponds to the number of measured and accepted aliquots. (PDF)

S1 Fig. Photo of the section of the Goda Buticha sedimentary sequence in 2008. (JPG)

S2 Fig. Example of glow (a) and growth curve (b) for one sample (GDB4, multi-grain aliquot) of Goda Buticha. (PDF)

S3 Fig. Results of various dose recovery tests for multi-grain aliquots. The test dose preheats were either 160°C cut (orange dots), or 220°C for 10 s (blue dots), or 240°C cut (grey dots). Each point is the arithmetic mean of three estimates. (PDF)

S4 Fig. Preheat plateaus. Each dot represents the De for one measured multi-grain aliquot. Preheat parameters, number of aliquots and central (CAM) De_s can be found in <u>S3 Table</u>. (JPG)

Acknowledgments

We thank the Ethiopian Authority for Research and Conservation of Cultural Heritage (ARCCH) for authorization to excavate in Goda Buticha. We would like to acknowledge the efforts of Habtewold H/Michael, Tilahun G/Selassie, Workalemahu Bekele, Hadis, Dejene Dendana Gulti, as well as the local people for their help and assistance during fieldwork. The authors thank Mrs. M.-M. Blanc-Valleron (UMR 7207 CR2P CNRS/MNHN) for her availability, and ease of access and use to her laboratory and the diffractometer, and Sébastien Nomade (UMR 8212 LSCE CNRS/CEA) for providing some radiocarbon analysis facilities. We are grateful to the French Center for Ethiopian Studies for logistical support. We thank the Tourism and Culture Offices of the Eastern and Western Harerghe, the Dire Dawa Administration, and the Harari National Regional State for fieldwork administrative support.

Author Contributions

Conceptualization: CT AA JJB EH AL DP OP ZA.

Formal analysis: LM.

Funding acquisition: CT EH AL DP ZA.

Investigation: CT AA JJB CC ED CF MH LM SP.

Methodology: CT JJB CC ED MH LM SP.

Project administration: DP ZA EH.

Resources: CT AA JJB MH EH AL DP OP ZA.

Supervision: DP EH ZA.

Validation: CT CF MH CC SP.

Visualization: CT AA JJB CC EH AL LM DP.

Writing – original draft: CT AA JJB CC MH EH AL LM DP OP SP.

Writing - review & editing: CT AA JJB CC MH EH AL LM DP OP SP.

References

- Villa P, Soriano S, Tsanova T, Degano I, Higham TFG, d'Errico F, et al. Border Cave and the beginning of the Later Stone Age in South Africa. Proceedings of the National Academy of Sciences. 2012; 109 (33): 13208–13.
- Gliganic LA, Jacobs Z, Roberts RG, Dominguez-Rodrigo M, Mabulla AZ. New ages for Middle and Later Stone Age deposits at Mumba rockshelter, Tanzania: Optically stimulated luminescence dating of quartz and feldspar grains. Journal of Human Evolution. Elsevier; 2012; 62(4): 533–47. doi: 10.1016/j. jhevol.2012.02.004 PMID: 22425240
- Ambrose SH. Chronology of the Later Stone Age and food production in East Africa. Journal of Archaeological Science. 1998; 25: 377–92.
- Wadley L. The Pleistocene Later Stone Age south of the Limpopo River. Journal of World Prehistory. 1993; 7(3): 243–96.
- 5. Barham L, Mitchell P. The first Africans: African archaeology from the earliest toolmakers to most recent foragers. Cambridge University Press; 2008.
- 6. Deacon J. The Later Stone Age of southernmost Africa. Oxford BAR. 1984.
- McBrearty S, Brooks AS. The revolution that wasn't: a new interpretation of the origin of modern human behavior. Journal of Human Evolution. 2000; 39(5): 453–563. doi: 10.1006/jhev.2000.0435 PMID: 11102266
- Texier P-JJ, Porraz G, Parkington JE, Rigaud J-P, Poggenpoel C, Tribolo C. The context, form and significance of the MSA engraved ostrich eggshell collection from Diepkloof Rock Shelter, Western Cape, South Africa. Journal of Archaeological Science. 2013; 40: 3412–31.
- Henshilwood CS, Marean CW, Milo RG, Yates R. An early bone tool industry from the Middle Stone Age at Blombos Cave, South Africa: implications for the origins of modern human behaviour, symbolism and language. Journal of Human Evolution. 2001; 41: 631–78. doi: <u>10.1006/jhev.2001.0515</u> PMID: <u>11782112</u>
- Mellars P, Boyle K, Bar-Yosef O, Stringer C. Rethinking the human revolution. Radiation Measurements. Cambridge: Cambridge: McDonald Institute for Archaeological Research; 2007.
- d'Errico F, Backwell L, Villa P, Degano I, Lucejko JJ, Higham TFG, et al. Early evidence of San material culture represented by organic artifacts from Border Cave, South Africa. Proceedings of the National Academy of Sciences. 2012; 109: 13214–9.
- 12. Ambrose SH. Small things remembered: Origins of early microlithic industries in sub-Saharan Africa. Archeological Papers of the American Anthropological Association. 2002; 12: 9–29.
- Hoffecker JF. The spread of modern humans in Europe. Proceedings of the National Academy of Sciences. 2009; 106(38): 16040–5.
- Teilhard de Chardin P. Le Paléolithique en Somalie française et en Abyssinie. L'Anthropologie. 1930; 40: 331–4.
- 15. Bailloud G. La préhistoire de l'Ethiopie. Cahiers de l'Afrique et de l'Asie. 1958; 5: 15–43.
- Clark JD, Williams MAJ. Recent archaeological research in southeastern Ethiopia. 1974–1975. Annales d'Ethiopie. 1978; 19–44.
- Joussaume R. L'art rupestre de l'Ethiopie. In: Roubet C, Hugot HJ, Souville G, éditeurs. Préhistoire Africaine: Mélanges offerts au doyen Lionel Balout. A.D.P.F; 1981. p. 159–74.

- Breuil H, Teilhard de Chardin P, Wernet P. Le paléolithique du Harrar. L'Anthropologie. 1951; 55: 219– 28.
- Cervicek P. Rock paintings of Laga Oda (Ethiopia). Paideuma, Paideuma: Mitteilungen zur Kulturkunde. 1971; 17: 121–36.
- **20.** Kurashina H. An examination of prehistoric lithic technology in east-central Ethiopia. Berkeley: University of California; 1978.
- Clark JD, Williamson KD, Michels JW, Marean CW. A Middle Stone Age occupation site at Porc Epic Cave, Dire Dawa (east-central Ethiopia). African Archaeological Review. 1984; 2: 37–71.
- 22. Assefa Z. Faunal remains from Porc-Epic: paleoecological and zooarchaeological investigations from a Middle Stone Age site in southeastern Ethiopia. Journal of Human Evolution. 2006; 51: 50–75. doi: <u>10.</u> 1016/j.jhevol.2006.01.004 PMID: 16545861
- Pleurdeau D. Human technical behavior in the African Middle Stone Age: the lithic assemblage of Porc-Epic cave (Dire Dawa, Ethiopia). African Archaeological Review. 2005; (22): 177–97.
- Brandt SA. The Upper Pleistocene and early Holocene prehistory of the Horn of Africa. African Archaeological Review. 1986; 4(1): 41–82.
- Gossa T, Sahle Y, Negash A. A reassessment of the Middle and Later Stone Age lithic assemblages from Aladi Springs, southern Afar Rift, Ethiopia. Azania: Archaeological Research in Africa. 2012; 47 (2): 210–22.
- Ménard C, Bon F, Dessie A, Bruxelles L, Douze K, Fauvelle F-X, et al. Late Stone Age variability in the Main Ethiopian Rift: New data from the Bulbula River, Ziway–Shala basin. Quaternary International. 2014; 343: 53–68.
- Brandt SA, Fisher EC, Hildebrand EA, Vogelsang R, Ambrose SH, Lesur J, et al. Early MIS 3 occupation of Mochena Borago Rockshelter, Southwest Ethiopian Highlands: implications for Late Pleistocene archaeology, paleoenvironments and modern human dispersals. Quaternary International. 2012; 274: 38–54.
- **28.** Gresham TH. An investigation of an upper Pleistocene archaeological site in Northern Somalia. M.A. Thesis. [Athens]: University of Georgia; 1984.
- 29. Brandt SA, Brook GA. Archaeological and palaeoenvironmental research in Northern Somalia. Current Anthropology. 1984; 25(1): 119–21.
- Pleurdeau D, Hovers E, Assefa Z, Asrat A, Pearson O, Bahain J-J, et al. Cultural change or continuity in the late MSA/Early LSA of southeastern Ethiopia? The site of Goda Buticha, Dire Dawa area. Quaternary International. 2014; 343: 117–35.
- Leplongeon A. La transition Middle Stone Age/Later Stone Age en Afrique de l'Est (Éthiopie). Muséum National d'Histoire Naturelle. Ph. D. thesis [Paris]; 2013. <u>https://drive.google.com/open?id=0B-J7agZgpknpcGlfMTN4VUhTRkk</u>
- Lopez S, van Dorp L, Hellenthal G. Human Dispersal Out of Africa: A Lasting Debate. Evolutionary Bioinformatics. 2016; 11(2): 57–68.
- Olivieri A, Achilli A, Pala M, Battaglia V, Fornarino S, Al-Zahery N, et al. The mtDNA Legacy of the Levantine Early Upper Palaeolithic in Africa. Science. 2006; 314(5806): 1767–70. doi: 10.1126/ science.1135566 PMID: 17170302
- Luis JR, Rowold DJ, Regueiro M, Caeiro B, Cinnioğlu C, Roseman C, et al. The Levant versus the Horn of Africa: Evidence for Bidirectional Corridors of Human Migrations. The American Journal of Human Genetics. 2004; 74(3): 532–44. doi: 10.1086/382286 PMID: 14973781
- 35. Hodgson JA, Mulligan CJ, Al-Meeri A, Raaum RL. Early back-to-Africa migration into the Horn of Africa. PLoS Genet. 2014.
- Gallego Llorente M, Jones ER, Eriksson A, Siska V, Arthur KW, Arthur JW, et al. Ancient Ethiopian genome reveals extensive Eurasian admixture in Eastern Africa. Science. American Association for the Advancement of Science; 2015; 350(6262): 820–2. doi: 10.1126/science.aad2879 PMID: 26449472
- Stewart JR, Stringer CB. Human Evolution Out of Africa: The role of refugia and climate change. Science. 2012; 335(6074): 1317–21. doi: 10.1126/science.1215627 PMID: 22422974
- Assefa Z, Pleurdeau D, Duquesnoy F, Hovers E, Pearson O, Asrat A, et al. Survey and explorations of caves in southeastern Ethiopia: Middle Stone Age and Later Stone Age archaeology and Holocene rock art. Quaternary International. 2014; 343: 136–47.
- Leplongeon A. Microliths in the Middle and Later Stone Age of eastern Africa: new data from Porc-Epic and Goda Buticha cave sites, Ethiopia. Quaternary International. 2014; 343: 100–16.
- **40.** Leplongeon A, Pleurdeau D, Hovers E. Late Pleistocene and Holocene lithic variability in Southeastern Ethiopia: implications for the understanding of the Middle and Late Stone Age of the Horn of Africa. Journal of African Archaeology. Accepted.

- **41.** Bronk Ramsey C, Lee S. Recent and Planned Developments of the Program OxCal. Radiocarbon. 2013; 55(3–4): 720.
- 42. Reimer PJ, Bard E, Bayliss A, Beck JW, Blackwell PG, Bronk Ramsey C, et al. IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years cal BP. Radiocarbon. 2013; 55(4): 1869.
- Crevecœur I. Étude anthropologique du squelette du Paléolithique supérieur de Nazlet Khater 2 (Égypte). Apport à la compréhension de la variabilité passée des hommes modernes. Leuven University Press; 2008.
- 44. Huntley DJ. Optical dating of sediments. Nature. 1985; 313(5998): 105-7.
- Aitken MJ. An introduction to optical dating: the dating of Quaternary sediments by the use of photonstimulated luminescence. Oxford University Press; 1998.
- Murray AS, Wintle AG. Luminescence dating of quartz using an improved single-aliquot regenerativedose protocol. Radiation Measurements. 2000; 32: 57–73.
- Murray AS, Wintle AG. The single aliquot regenerative dose protocol: potential for improvements in reliability. Radiation Measurements. 2003; 37(4–5): 377–81.
- Prescott JR, Hutton JT. Cosmic ray contributions to dose rates for luminescence and ESR dating: large depths and long-term time variations. Radiation Measurements. 1994; 23(2): 497–500.
- Mercier N, Falguères C. Field gamma dose-rate measurement with a Nal(TI) detector: re-evaluation of the "threshold" technique. Ancient TL. 2007; 25(1): 1–4.
- Guérin G, Mercier N. Determining gamma dose rates by field gamma spectroscopy in sedimentary media: results of Monte Carlo simulations. Radiation Measurements. 2011; 46(2): 190–5.
- Martin L. Caractérisation et modélisation d'objets archéologiques en vue de leur datation par des méthodes paléo-dosimétriques. Simulation des paramètres dosimétriques sous Geant4. 2015. <u>http://www.theses.fr/s77802</u>.
- Martin L, Incerti S, Mercier N. DosiVox: Implementing Geant 4-based software for dosimetry simulations relevant to luminescence and ESR dating techniques. Ancient TL. 2015; 33(1): 1–10.
- 53. Guérin G, Mercier N, Nathan R, Adamiec G, Lefrais Y. On the use of the infinite matrix assumption and associated concepts: a critical review. Radiation Measurements. 2012; 47: 778–85.
- Mercier N, Valladas H, Schiegl S, Yosef OB, Weiner S. Thermoluminescence dating and the problem of geochemical evolution of sediments—a case study: the Mousterian levels at Hayonim. Israel Journal of Chemistry. 1995; 35(2): 137–41.
- Galbraith RF, Roberts RG, Laslett GM, Yoshida H, Olley JM. Optical dating of single and multiple grains of quartz from Jinmium rock shelter, northern Australia: Part I, experimental design and statistical models. Archaeometry. 1999; 41(2): 339–64.
- 56. Guérin G, Frouin M, Talamo S, Aldeias V, Bruxelles L, Chiotti L, et al. A multi-method luminescence dating of the Palaeolithic sequence of La Ferrassie based on new excavations adjacent to the La Ferrassie 1 and 2 skeletons. Journal of Archaeological Science. 2015; 58: 147–66.
- Combès B, Philippe A, Lanos P, Mercier N, Tribolo C, Guibert P, et al. A Bayesian central equivalent dose model for optically stimulated luminescence dating. Quaternary Geochronology. 2015; 28: 62–70.
- 58. Asrat A, Barbey P, Gleizes G. The Precambrian geology of Ethiopia: a review. 2001; 8: 271–88.
- Bosellini A, Russo A, Assefa G. The Mesozoic succession of Dire Dawa, Harar Province, Ethiopia. Journal of African Earth Sciences. 2001; 32: 403–17.
- Getaneh W. Geochemistry provenance and depositional tectonic setting of the Adigrat Sandstone northern Ethiopia. Journal of African Earth Sciences. 2002; 35: 185–98.
- Asrat A. The Rock-Hewn churches of Tigrai, northern Ethiopia: a geological perspective. geoarchaeology: an international journal. 2002; 17(7): 649–63.
- Martini J, Kavalieris I. Mineralogy of the Transvaal caves. Transactions of the geological society of South Africa. 1978; 81: 47–54.
- 63. Hill C, Forti P. Cave Minerals of the World: National Speleological Society (NSS). Huntsville, Alabama.
- Kašpar J. Genése guanových minerálů z jeskyně Domice. Věstník SGÚ RCS. Praha; 1934; 10: 104– 11.
- Brochier JÉ. Les sédiments anthropiques. Géologie de la Préhistoire: Méthodes, Techniques et Applications. Méthodes d'étude et perspectives Géologie de la Préhistoire. 2002. p. 453–73.
- Gasse E, Street FA. Late Quaternary lake-level fluctuations and environments of the northern Rift Valley and Afar region (Ethiopia and Djibouti). Palaeogeography, Palaeoclimatology, Palaeoecology. 1978; 24(4): 279–325.
- 67. Gasse F, Rognon P, Street FA. Quaternary history of the Afar and Ethiopian Rift lakes. In: Williams MA, Faure H, éditeurs. The Sahara and the Nile. Rotterdam: A.A. Balkema; 1980. p. 361–400.

- Foerster V, Junginger A, Langkamp O, Gebru T, Asrat A, Umer M, et al. Climatic change recorded in the sediments of the Chew Bahir basin, southern Ethiopia, during the last 45,000 years. Quaternary International. 2012; 274: 25–37.
- Foerster V, Vogelsang R, Junginger A, Asrat A, Lamb HF, Schaebitz F, et al. Environmental change and human occupation of southern Ethiopia and northern Kenya during the last 20,000 years. Quaternary Science Reviews. 2015; 129: 333–40.
- Asrat A, Baker A, Leng M, Gunn J, Umer M. Environmental monitoring in the Mechara caves, Southeastern Ethiopia: implications for speleothem palaeoclimate studies. International Journal of Speleology. 2008; 37: 207–20.
- Asrat A, Baker A, Leng MJ, Van Calsteren P, Smith C, Kohsiek P. A high-resolution multi-proxy stalagmite record from Mechara, Southeastern Ethiopia: palaeohydrological implications for speleothem palaeoclimate reconstruction. Journal of Quaternary Science. 2007; 22(1): 53–63.
- Baker A, Asrat A, Leng M, Thomas L, Fairchild IJ. Decadal-scale holocene climate variability recorded in an Ethiopian stalagmite. The Holocene. 2010:20; 827–36.
- Phillipson DW. The Excavation of Gobedra Rock-Shelter, Axum: an early occurrence of cultivated finger millet in northern Ethiopia. Azania: Archaeological Research in Africa. 1977; 12(1): 53–82.
- 74. Gutherz X, Diaz A, Ménard C, Bon F, Douze K, Léa V, et al. The Hargeisan revisited: Lithic industries from shelter 7 of Laas Geel, Somaliland and the transition between the Middle and Late Stone Age in the Horn of Africa. Quaternary International. 2014; 343: 69–84.
- 75. Ménard C, Bon F. Hiatus et lacune. Occupation du Rift éthiopien à la fin du Pleistocene et au debut de l'Holocene. In: Naudinot N, Meignen L, Binder D, Querre G, éditeurs. Antibes: Editions APDCA; 2015.
- Pleurdeau D. Gestion des matières premières et comportements techniques dans le Middle Stone Age africain. British Archaeological Reports Limited; 2004.
- 77. Duplaix S. Détermination microscopique des minéraux des sables. Beranger C, éditeur. 1958.
- Duller GA. Distinguishing quartz and feldspar in single grain luminescence measurements. Radiation Measurements. 2003; 37: 161–5.
- Duller GA, Bøtter-Jensen L, Kohsiek P, Murray AS. A high-sensitivity optically stimulated luminescence scanning system for measurement of single sand-sized grains. Radiation Protection Dosimetry. 1999; 84: 325–30.
- Bøtter-Jensen L, Andersen CE, Duller GA, Murray AS. Developments in radiation, stimulation and observation facilities in luminescence measurements. Radiation Measurements. 2003; 37: 535–41.
- Bøtter-Jensen L, Bulur E, Duller GA. Advances in luminescence instrument systems. Radiation Measurements. 2000; 32: 523–8.
- Duller GA. The Analyst software package for luminescence data: overview and recent improvements. Ancient TL. 2015; 33(1): 35–42.
- Bateman MD, Boulter CH, Carr AS, Frederick CD, Peter D, Wilder M. Detecting post-depositional sediment disturbance in sandy deposits using optical luminescence. Quaternary Geochronology. 2007; 2 (1–4): 57–64.
- 84. Guérin G, Jain M, Mercier N. Modelling dose rate to single grains of quartz in well-sorted sand samples: The dispersion arising from the presence of potassium feldspars and implications for single grain OSL dating. Quaternary Geochronology. 2015; 27: 52–65.
- Thomsen KJ, Murray A, Jain M. The dose dependency of the over-dispersion of quartz OSL single grain dose distributions. Radiation Measurements. 2012; 47(9): 732–9.
- Thomsen KJ, Murray AS, Bøtter-Jensen L. Sources of variability in OSL dose measurements using single grains of quartz. Radiation Measurements. 2005; 39(1): 47–61.
- Mayya YS, Morthekai P, Murari MK, Singhvi AK. Towards quantifying beta microdosimetric effects in single-grain quartz dose distribution. Radiation Measurements. 2006; 41(7–8): 1032–9.
- 88. Aitken MJ. Thermoluminescence dating. London: Academic Press; 1985
- **89.** Zimmerman DW. Thermoluminescent dating using fine grains from pottery. Archaeometry. 1971; 13 (1): 29–52.
- Nathan R.P., Thomas P.J., Jain M., Murray A.S., Rhodes E.J. Environmental dose rate heterogeneity of beta radiation and its implications for luminescence dating: Monte Carlo modelling and experimental validation. Radiation Measurements. 2003; 37, 305–313.
- **91.** Martin L, Mercier N, Incerti S. Geant4 simulations for sedimentary grains in infinite matrix conditions: The case of alpha dosimetry. Radiation Measurements. 2014; 70: 39–47.
- 92. Guérin G. Modélisation et simulations numériques des effets dosimétriques dans les sédiments quaternaires: application aux méthodes de datation par luminescence. 2011. <u>http://www.theses.fr/</u> 163762791.

93. Mercier N, Incerti S, Lefrais Y, Pecheyran C, Guérin G, Jarry M, et al. Dosimetric study of sediments at the beta dose rate scale: Characterization and modelization with the DosiVox software. Radiation Measurements. 2015; 81: 134–41.