



**HAL**  
open science

## Introduction to the versatility of biomimetic membranes to characterize modulators of $A\beta_{1-42}$ fibrillization with potential interest in Alzheimer's disease treatment

Willy Smeralda, Marc Since, Michael Boisserie, Sophie Corvaisier, Julien Cardin, Aurélie Malzert-Fréon

### ► To cite this version:

Willy Smeralda, Marc Since, Michael Boisserie, Sophie Corvaisier, Julien Cardin, et al.. Introduction to the versatility of biomimetic membranes to characterize modulators of  $A\beta_{1-42}$  fibrillization with potential interest in Alzheimer's disease treatment. 32ème Journées Franco-Belges de Chimie Thérapeutique et 26ème Rencontres Internationales des Pharmacochimistes de l'Arc Atlantique, , Jun 2018, Asnelles sur Mer, France. 2018. hal-01844495

**HAL Id: hal-01844495**

**<https://hal.science/hal-01844495>**

Submitted on 19 Jul 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Introduction to the versatility of biomimetic membranes to characterize modulators of Aβ<sub>1-42</sub> fibrillization with potential interest in Alzheimer's disease treatment

Willy Smeralda<sup>1</sup>, Marc Since<sup>1</sup>, Michael Boisserie<sup>2</sup>, Sophie Corvaisier<sup>1</sup>, Julien Cardin<sup>2</sup>, Aurélie Malzert-Fréon<sup>1</sup>  
<sup>1</sup>Centre d'Etudes et de Recherche sur le Médicament de Normandie (CERMN), Normandie Univ, UNICAEN, 14000 Caen, France  
<sup>2</sup>NIMPH Team, CIMAP CNRS UMR6252, EnsiCaen-UNICAEN-CEA, 14050 Caen Cedex, France  
 E-mail : [willy.smeralda@unicaen.fr](mailto:willy.smeralda@unicaen.fr)




## INTRODUCTION

**Alzheimer's disease (AD)** is the most widespread form of senile dementia worldwide and represents the leading socioeconomic problem in healthcare. The onset and the progression of this neurodegenerative disease is associated with the aggregation of the amyloid-β peptide (Aβ) through a fibrillization process. Hence, an attractive therapeutic strategy against this disorder is the development of molecules able to interfere at specific steps of this fibrillization pathway<sup>1</sup>, *i.e.* able to inhibit the formation of the soluble toxic oligomeric Aβ aggregates.

To identify such promising therapeutic molecules, experimental methods are required *in vitro* to precisely follow and characterize the state (monomer/oligomers/fibrils) and the conformation of Aβ peptide during its fibrillization process. In the same time, these methods must be simple enough to remain compatible with the High Throughput Screening (HTS) of new compounds with potential interest in AD treatment.

In the present study, we propose to combine experimental methods to allow a multiparameter characterization of potential modulators of Aβ<sub>1-42</sub> fibrillization. These kinetic studies are carried out in presence of liposomes as biomimetic neuronal membranes. Indeed, it has been established that lipids of the neuronal membrane are an important factor in the cascade of events that leads to the formation of the amyloid fibers.


Adapalene Bexarotene Indometacin O-Vanillin Chicago sky blue 6B



We used ThT to monitor<sup>3</sup> the effect of these molecules on the aggregation of the peptide in their presence at different concentrations.

## Kinetic study of Aβ<sub>1-42</sub> aggregation

In order to be able to characterize new compounds we have selected 5 molecules whose anti-aggregative properties are already described in the scientific literature.<sup>1,2</sup>




Aβ<sub>1-42</sub> kinetic in presence of o-vanillin

Quantitative effect observed in the presence of o-vanillin. In addition to inhibiting the formation of oligomers, it also inhibits the formation of fibers.



## Conformational study of Aβ<sub>1-42</sub> aggregation

In parallel, considering that the fibers formed by Aβ are mainly organized into β-sheets, assays were correlated with the secondary structure analysis of the peptide using ATR-FTIR spectroscopy.<sup>5</sup>




ATR-FTIR spectra of aggregated Aβ<sub>1-42</sub>

Structure	Amide I frequency (cm <sup>-1</sup> )
α-helical	1655-1665
α-helical/β-sheet	1650-1660
β-sheet	1625-1640
Aggregated strands	1610-1620


ATR-FTIR spectra of aggregated Aβ<sub>1-42</sub> at 10μM. The amide band I position is characteristic of the β sheets.

Using a reference (peptide alone), it will be possible to make relative comparisons and to study possible effects of the molecules of interest on the peptide structuration.



Aβ<sub>1-42</sub> kinetic in presence of indometacin

Kinetic effect observed in the presence of indometacin. Inhibiting the formation of oligomers (critical step of aggregation) results in an acceleration of the aggregation.



It is therefore possible to discriminate different types of molecule actions thanks to ThT monitoring, which is a simple and effective method that is perfectly adapted to screening.

However, experiments will be performed on variants of Aβ<sub>1-42</sub><sup>4</sup> to confirm our results and support our conclusions.

## Molecule/membrane interactions

In order to take into account the role of the membrane in the peptide aggregation but also because the molecules tested will be potential drug candidates, it is essential to study the interactions between the molecules and the membranes.

First, using a complex model of liposomes (SPC/CHOL/PG<sub>60</sub>:20:20) to mimic neuronal membranes, we have transposed a methodology of the scientific literature<sup>6</sup> to study of interactions between modulators of Aβ<sub>1-42</sub> aggregation and membranes.


Only one of the compounds demonstrated interaction with the liposome membranes.

These results were correlated with toxicity studies where adapalene is also the only compound to have exerted a membrane destabilization.

In order to study the toxicity, we once again used liposomes using the liposome leakage assay<sup>7</sup> method.

	Adapalene	Bexarotene	Chicago SB6B	Indometacin	o-Vanillin
Structure					
Log D7,4 (MarvinSketch)	3,9	3,82	-4,95	0,27	1,86
Log Kp	3,6	<1	<1	<1	<1

Partition coefficient determination of aggregation modulators



Liposome leakage by aggregation modulators

These are preliminary results and confirmation remains necessary. The combination of these two experiments allows a more complete prediction of the interaction of molecules with membranes.

## CONCLUSION

From results obtained to date, it appears that the tested modulators present a range of effects that can affect defined stages of the aggregation of the Aβ<sub>1-42</sub> peptide. Conformational studies with ATR-FTIR spectroscopy will give us the possibility to analyse if any molecules have an effect on the secondary structure. Especially, on anti-parallel β-sheets structure that characterizes the presence of toxic oligomeric species<sup>8</sup>.

In the long term the results will make it possible to characterize the kinetic effects of the reference molecules and may be applied to original molecules synthesized at the CERMN.

This study highlights also some advantages of the use of liposomes to characterize and discriminate newly Aβ aggregation modulators synthesized by chemists and could even be expanded to other amyloid diseases.

**ACKNOWLEDGEMENTS :** W. Smeralda was recipient of a doctoral fellowship from the French State (Research Ministry). We thank Lipoid GmbH (Ludwigshafen, Germany) for providing the phospholipids for this study. We also thanks the Normandy Doctoral School of Chemistry for financial support and all our collaborators.

**References :**  
 (1) Habchi, J.; Chia, S.; Limbocker, R.; Mannini, B.; Ahn, M.; Perni, M.; Hansson, O.; Arosio, P.; Kumita, J. R.; Challa, P. K.; et al. Systematic Development of Small Molecules to Inhibit Specific Microscopic Steps of Aβ<sub>42</sub> Aggregation in Alzheimer's Disease. *Proc. Natl. Acad. Sci.* 2017, 114 (2), E200-E208.  
 (2) Necula, M.; Kaye, R.; Milton, S.; Glabe, C. G. Small Molecule Inhibitors of Aggregation Indicate That Amyloid β Oligomerization and Fibrillization Pathways Are Independent and Distinct. *J. Biol. Chem.* 2007, 282 (14), 10311-10324.  
 (3) Voropai, E. S.; Samstov, M. P.; Kaplevskii, K. N.; Maskevich, A. A.; Stepiro, V. I.; Povarova, O. I.; Kuznetsova, I. M.; Turoverov, K. K.; Fink, A. L.; Uverskii, V. N. Spectral Properties of Thioflavin T and Its Complexes with Amyloid Fibrils. *J. Appl. Spectrosc.* 2003, 70 (6), 868-874.  
 (4) Henry, S.; Vignaud, H.; Bobo, C.; Decossas, M.; Lambert, O.; Harte, E.; Alves, I. D.; Cullin, C.; Lecomte, S. Interaction of Aβ<sub>1-42</sub> Amyloids with Lipids Promotes "Off-Pathway" Oligomerization and Membrane Damage. *Biomacromolecules* 2015, 16 (3), 944-950.  
 (5) Sarroukh, R.; Goormaghtigh, E.; Ruyschaert, J.-M.; Raussens, V. ATR-FTIR: A "rejuvenated" Tool to Investigate Amyloid Proteins. *Biochim. Biophys. Acta BBA - Biomembr.* 2013, 1828 (10), 2328-2338.  
 (6) Magalhães, L. M.; Nunes, C.; Lúcio, M.; Segundo, M. A.; Reis, S.; Lima, J. L. F. C. High-Throughput Microplate Assay for the Determination of Drug Partition Coefficients. *Nat. Protoc.* 2010, 5 (11), 1823-1830.  
 (7) Jimah, J.; Schlesinger, P.; Tolia, N. Liposome Disruption Assay to Examine Lytic Properties of Biomolecules. *BIO-Protoc.* 2017, 7 (15).  
 (8) Vignaud, H.; Bobo, C.; Lascu, I.; Sörgjerd, K. M.; Zako, T.; Maeda, M.; Salin, B.; Lecomte, S.; Cullin, C. A Structure-Toxicity Study of Aβ<sub>42</sub> Reveals a New Anti-Parallel Aggregation Pathway. *PLoS ONE* 2013, 8 (11), e80262.