



Sign changes of Kloosterman sums and exceptional characters

Sary Drappeau, James Maynard

► To cite this version:

Sary Drappeau, James Maynard. Sign changes of Kloosterman sums and exceptional characters. Proceedings of the American Mathematical Society, 2019, 147, pp.77-90. 10.1090/proc/14239 . hal-01847957

HAL Id: hal-01847957

<https://hal.science/hal-01847957>

Submitted on 18 Jun 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SIGN CHANGES OF KLOOSTERMAN SUMS AND EXCEPTIONAL CHARACTERS

SARY DRAPPEAU AND JAMES MAYNARD

ABSTRACT. We prove that the existence of exceptional real zeroes of Dirichlet L -functions would lead to cancellations in the sum $\sum_{p \leq x} \text{Kl}(1, p)$ of Kloosterman sums over primes, and also to sign changes of $\text{Kl}(1, n)$, where n runs over integers with exactly two prime factors. Our arguments involve a variant of Bombieri's sieve, bounds for twisted sums of Kloosterman sums, and work of Fouvry and Michel on sums of $|\text{Kl}(1, n)|$.

1. INTRODUCTION

Kloosterman sums. For $n \in \mathbb{N}_{>0}$ and a residue class $a \pmod{n}$, define the normalized Kloosterman sum as

$$\text{Kl}(a, n) = \frac{1}{\sqrt{n}} \sum_{\substack{\nu \pmod{n} \\ (\nu, n)=1}} e\left(\frac{\nu + a\bar{\nu}}{n}\right),$$

where we write $e(z) = e^{2\pi iz}$ and $\bar{\nu}\nu \equiv 1 \pmod{n}$. These sums have a long history [Poi11, Klo27], at the intersection of algebraic geometry and automorphic forms. The Weil bound [Wei48, Mat11] yields $|\text{Kl}(a, n)| \leq 2^{\omega(n)}$ if $32 \nmid n$ and $|\text{Kl}(a, n)| \leq 2^{\omega(n)+1/2}$ in general, where $\omega(n)$ is the number of distinct prime factors of n . In particular, for a prime p

$$|\text{Kl}(1, p)| \leq 2.$$

Let $\theta_{a,p} \in [0, \pi]$ be such that $\text{Kl}(a, p) = 2 \cos(\theta_{a,p})$. The “vertical” Sato-Tate law, due to Katz [Kat88], asserts that the numbers

$$\{\theta_{a,p} \mid 1 \leq a < p\}$$

become equidistributed, as $p \rightarrow \infty$, with respect to the Sato-Tate measure $\frac{2}{\pi} \sin(\theta)^2 d\theta$. The “horizontal” Sato-Tate conjecture is the claim that the numbers

$$\{\theta_{1,p} \mid p \leq x\}$$

become equidistributed with respect to the same measure, as $x \rightarrow \infty$. This would of course imply that

$$\sum_{p \leq x} \text{Kl}(1, p) = o(\pi(x)) \quad (x \rightarrow \infty).$$

Unfortunately the horizontal Sato-Tate conjecture is still open, and very little is known about this sum. Fouvry and Michel [FM07] have obtained significant partial progress on replacing primes by almost-primes: they show that

$$\sum_{\substack{n < x \\ p|n \Rightarrow p > x^{1/23.9}}} (|\text{Kl}(1, n)| \pm \text{Kl}(1, n)) \gg \frac{x}{\log x}.$$

In particular, it follows that there are infinitely many sign changes in the set $\{\text{Kl}(1, n), \omega(n) \leq 23\}$. After further work by many authors [FM03a, SF07, SF09, Mat11, Xi15a], the best known current result is due to Xi [Xi15b] and shows that there are infinitely many sign changes in the set $\{\text{Kl}(1, n), \omega(n) \leq 7\}$. We refer to the recent preprint [Xi18] for more references and related questions.

Date: March 14, 2018.

2010 *Mathematics Subject Classification.* 11L05, 11N36 (Primary); 11N75, 11L20, 11M20 (Secondary).

We thank É. Fouvry for helpful remarks concerning this work, and an anonymous referee for comments which helped improve the manuscript. Part of this work was done during a visit of JM to Aix-Marseille university, supported by the French-Austrian joint project MuDeRa (FWF I-1751-N26, ANR-14-CE34-0009).

Landau-Siegel zeroes. In this work we study the implications of the existence of Landau-Siegel zeroes on this question. For $D \geq 3$ an integer and $\chi \pmod{D}$ a real primitive character, define the value

$$\eta_\chi := L(1, \chi) \log D.$$

It would follow from GRH that $\eta_\chi \gg (\log D)/\log \log D$ and so $\eta_\chi \rightarrow \infty$ as $D \rightarrow \infty$. Unconditionally, however, Siegel's non-effective lower-bound [Sie35] $\eta_\chi \gg_\varepsilon D^{-\varepsilon}$ remains unsurpassed. It is as yet not even known if η_χ is bounded away from 0; this is equivalent to the non-existence of real zeroes of $L(s, \chi)$ close to 1. On the other hand, if there were a sequence of characters $\chi_1 \pmod{D_1}, \chi_2 \pmod{D_2}, \dots$ such that $\eta_{\chi_i} \rightarrow 0$, then many desirable consequences would follow: the existence of twin primes [HB83], equidistribution of primes in arithmetic progressions to large moduli [FI03], primes in very short intervals [FI04], or prime values of discriminants of elliptic curves [FI05, FI13], for example.

We are interested in obtaining cancellations in the sum

$$\sum_{p \leq x} \text{Kl}(1, p)$$

which go beyond the bound $2\pi(x)$ implied by the Weil bound should such exceptional characters exist.

Theorem 1.1. *Let $\varepsilon > 0$. Then there are constants $A, B > 0$, depending only on ε , such that for $D \geq 3$, $x \geq D^A$ and any primitive real character $\chi \pmod{D}$, we have*

$$\left| \sum_{p < x} \text{Kl}(1, p) \right| \leq \pi(x) \left(\varepsilon + BL(1, \chi) \log x \right).$$

This statement is unconditional, but is only non-trivial if the value η_χ is suitably small. We note that if there is a sequence of characters with $\eta_{\chi_i} \rightarrow 0$, then Theorem 1.1 shows that for a suitable sequence of values of x ,

$$\sum_{p < x} \text{Kl}(1, p) = o(\pi(x)),$$

as predicted by the horizontal Sato-Tate conjecture.

Unfortunately we do not know unconditionally the expected lower bound

$$(1.1) \quad \sum_{p < x} |\text{Kl}(1, p)| \gg \pi(x).$$

In particular, even if there was a sequence of characters with $\eta_{\chi_i} \rightarrow 0$, we would not be able to conclude from Theorem 1.1 that there are even infinitely many sign changes in the sequence $\text{Kl}(1, p)$. If instead of considering primes we consider products of exactly two primes, then the equivalent lower bound to (1.1) is known thanks to work of Fouvry-Kowalski-Michel [FKM14]. For technical reasons, when working with products of two primes we consider the variant

$$S(x) = \sum_{p, q \text{ prime}} \phi\left(\frac{pq}{x}\right) \log(pq)(\log p)(\log q) \text{Kl}(1, pq)$$

where $\phi : \mathbb{R}_+ \rightarrow \mathbb{C}$ is a smooth function compactly supported inside \mathbb{R}_+^* . We note that the Weil bound implies unconditionally that $S(x) \ll x \log^2 x$, whilst a variant of the horizontal Sato-Tate conjecture would suggest that we should have $S(x) = o_\phi(x \log^2 x)$.

Theorem 1.2. *Let $\varepsilon > 0$. Then there exist $A, B > 0$, depending at most on ε , such that for any $D \geq 3$, $x \geq D^A$ and any primitive real character $\chi \pmod{D}$, we have*

$$(1.2) \quad |S(x)| \ll_\phi x (\log x)^2 \{ \varepsilon + BL(1, \chi) \log x \}.$$

The implied constant depends only on the function ϕ .

As with Theorem 1.1, this is unconditional but non-trivial only if a sequence of exceptional characters exist. Thus, in the presence of Siegel zeros, we are able to establish infinitely many sign changes of $\text{Kl}(1, pq)$.

Corollary 1.3. *For some absolute constants $A, c > 0$, if $\eta_\chi \leq c$, then every interval $[x, 2x] \subset [D^A, D^{100A}]$ contains two numbers (n_1, n_2) with $\omega(n_1) = \omega(n_2) = 2$, and $\text{Kl}(1, n_1) \text{Kl}(1, n_2) < 0$.*

Proof. Choose ϕ to be real-valued with $\phi \geq \mathbf{1}_{[1, 2]}$, and consider the unsigned sum

$$A(x) = \sum_{\substack{p, q \\ x < pq \leq 2x}} (\log pq)(\log p)(\log q) |\text{Kl}(1, pq)|.$$

We have the following lower bound, due to Fouvry-Kowalski-Michel [FKM14, Proposition 5.1]:

$$(1.3) \quad A(x) \gg x(\log x)^2 \quad (x \geq 2).$$

Comparing (1.2) and (1.3) yields the claimed statement. \square

Notations. We denote by $P^-(n)$ (resp. $P^+(n)$) the smallest (resp. largest) prime factor of n , with the conventions $P^-(1) = \infty$ and $P^+(1) = 1$. The letter p denotes a prime.

2. OUTLINE

If there is a character $\chi \pmod{D}$ such that $L(1, \chi)$ is very small, then ‘most’ primes $p \in [D^A, D^{100A}]$ have $\chi(p) = -1 = \mu(p)$, for some suitable constant A . By multiplicativity, this means that the (poorly understood) Moebius function μ can be well-approximated by the character χ (which is periodic \pmod{D}), and so better understood) on integers $n < D^{100A}$ with no small prime factors. In particular, for $n \leq D^{100A}$

$$(2.1) \quad \Lambda(n) = (\mu * \log)(n) \approx (\chi * \log)(n),$$

and so we can approximate Λ by the convolution of two simpler sequences. By applying the hyperbola method, this would allow us to estimate a sum $\sum_{n < x} \Lambda(n) a_n$ provided we could suitably estimate

$$(2.2) \quad \sum_{d \leq Y_1} \chi(d) \sum_{\substack{n \leq x \\ d|n}} a_n \log \frac{n}{d}, \quad \text{and} \quad \sum_{d \leq Y_2} \log d \sum_{\substack{n < x \\ d|n}} a_n \chi\left(\frac{n}{d}\right)$$

for some choice of Y_1, Y_2 with $Y_1 Y_2 = x = D^{100A}$. Often one can suitably estimate such sums for $Y_1 = Y_2 = x^{1/2-\varepsilon}$, which just falls short of this requirement. Much of the work on the distribution of primes under the assumption of a Siegel-Landau zero followed this strategy, and the key technical challenge is then to obtain a suitable estimate for one of the sums in (2.2) with Y_1 or Y_2 slightly beyond $x^{1/2}$.

In our situation, $a_n = \text{Kl}(1, n)$, and estimates for the two sums in (2.2) with $Y_1 = Y_2 = x^{1/2-\varepsilon}$ are obtained in essentially the same way as Fouvry and Michel [FM07]. Unfortunately we do not know how to extend this work beyond $x^{1/2}$, and so this strategy fails. However, in the convolution identity $\Lambda(n) = \sum_{d|n} \mu(d) \log(n/d)$ it is only terms with $d \in [x^{1/2-\varepsilon}, x^{1/2+\varepsilon}]$ which we are unable to handle. We note the alternative identity

$$\Lambda(n) = \sum_{d|n} \mu(d) \left(\log \frac{\sqrt{n}}{d} \right).$$

The presence of the term $\log(\sqrt{n}/d)$ means we expect that terms with $d \approx \sqrt{n}$ to contribute less, and so we might hope that these central values would be negligible. This is a variant of the idea that Bombieri introduced in his asymptotic sieve [Bom76], where terms in $\Lambda_2(n) = \sum_{d|n} \mu(d) \log^2(n/d)$ from $d < n^{1-\varepsilon}$ could be handled by assumptions on equidistribution of congruence sums, and terms with $d \in [n^{1-\varepsilon}, n]$ could be bounded by virtue of the fact that $\log^2(n/d)$ was small in this range.

Unfortunately, as in Bombieri’s work, this strategy fails if we wish to count primes. To maintain the feature that the support is essentially restricted to numbers with no small prime factors one multiplies by a short sieve weight, which loses a factor ε^2 from the two variables d and n/d . This precisely cancels out the gains of a factor ε^2 coming from the range of d and from the size of $\log(\sqrt{n}/d)$. Whilst this issue might appear to be a technicality, at least in Bombieri’s work this is an expression of the fundamental parity problem of sieve methods. If instead we counted with a weight involving a higher power of $\log(\sqrt{n}/d)$ (thereby counting products of a bounded number of primes), then this strategy can succeed.

In our case, we are interested in $a_n = \text{Kl}(1, n)$. Although in general we expect the Weil bound $|\text{Kl}(1, n)| \leq 2^{\omega(n)}$ to be essentially sharp, for *most* integers n we expect $|\text{Kl}(1, n)|$ is actually much smaller than this. Indeed, the horizontal Sato-Tate conjecture would predict that for any fixed a , the average size of $|\text{Kl}(a, p)|$ is $\frac{2}{\pi} \int_0^\pi 2|\cos(t)| \sin^2(t) dt = 8/3\pi < 1$. By multiplicativity, we might then expect $|\text{Kl}(1, n)| \approx (8/3\pi)^{\omega(n)}$ on average over n . Fouvry and Michel [FM03b, FM06] combined an argument of Hooley [Hoo64] based on the identity $\text{Kl}(1, ab) = \text{Kl}(\overline{a^2}, b) \text{Kl}(\overline{b^2}, a)$ (for coprime a, b), with the *vertical* Sato-Tate law, to show unconditionally that the factor $2^{\omega(n)}$ can be indeed improved to $(8/3\pi)^{\omega(n)}$ when considering suitable averages. Since $8/3\pi < 1$, numbers with a larger number of prime factors contribute less to the problematic sums, and so there is less of a loss from being restricted to a short sieve weight. This ultimately allows us to win an additional factor of $(\varepsilon^{1-8/3\pi})^2$ for these sums

involving middle sized d , which is enough to conclude that such terms make a negligible contribution, and so we are able to bound $\sum_{p < x} \text{Kl}(1, p)$.

3. PREPARATORY LEMMAS

3.1. Level of distribution for twisted Kloosterman sums. Here we make precise the claim that the sums in (2.2) can be estimated with $Y_1 = Y_2 = x^{1/2-\varepsilon}$ by a variation of the work of Fouvry-Michel [FM07].

Proposition 3.1. *Let $\varepsilon > 0$ be fixed. There exists $\eta = \eta(\varepsilon) > 0$, such that for any real $x \geq 2$, positive integer $D \leq x^\eta$, all characters $\chi \pmod{D}$, and any smooth function $\phi : \mathbb{R}_+^* \rightarrow \mathbb{C}$ compactly supported inside \mathbb{R}_+^* , we have*

$$(3.1) \quad \sum_{q \leq x^{1/2-\varepsilon}} \left| \sum_{n \equiv 0 \pmod{q}} \phi\left(\frac{n}{x}\right) \chi(n) \text{Kl}(1, n) \right| \ll_{\varepsilon, \phi} x^{1-\eta}.$$

Remark. Note that the case $D = 1$ of the previous statement is a weaker form of Proposition 2.1 of [FM07].

Proof. We may plainly assume that the sums are restricted to $(q, D) = (n, D) = 1$. The bound we claim is a variant of Proposition 2.1 of [FM07], which is based on the Kuznetsov formula [Kuz80, DI83], the Weil bound for Kloosterman sums, and a uniform bound $\theta \leq 1/4 - \varepsilon$ towards Ramanujan-Petersson, which was first due to Luo-Rudnick-Sarnak [LRS95].

The difference in our case is the presence of the character. Recently, Blomer and Milićević [BM15] have succeeded in analysing such sums in the context of modular forms with non-trivial nebentypus; another argument was used in [Dra17], which is simpler for our purpose here. We will rely on work of Topacogullari [Top15] to estimate the spectral sums.

In our case, we will use the notations and normalization described in section 4.1.2 of [Dra17]. Our aim is to apply the Kuznetsov formula [Dra17, Lemma 4.5] for the group $\Gamma_0(qD)$, nebentypus $\bar{\chi} \pmod{D}$, with cusps ∞ and $1/q$, and parameters $m \leftarrow D$, $n \leftarrow 1$. For each q in the left-hand side of (3.1), Lemma 4.3 of [Dra17], with the choice of scaling matrices (depending only on q and D) given there, yields

$$\chi(n) \text{Kl}(1, n) = \chi(q) e(-\bar{q}/D) S_{\infty, 1/q}(D, 1; n\sqrt{D}).$$

Let $\kappa \in \{0, 1\}$ be such that $\chi(-1) = (-1)^\kappa$. The Kuznetsov formula with test function $\psi(t) = \phi(4\pi/(tx)) \sqrt{4\pi/(tx)}$ yields

$$\overline{\chi(q)} e(\bar{q}/D) \sum_{n \equiv 0 \pmod{q}} \phi\left(\frac{n}{x}\right) \chi(n) \text{Kl}(1, n) = \sqrt{x} \{ \mathcal{H} + \mathcal{E} + \mathcal{M} \},$$

where

$$\begin{aligned} \mathcal{H} &= \sum_{\substack{k > \kappa \\ k \equiv \kappa \pmod{2}}} \sum_{f \in \mathcal{B}_k(qD, \bar{\chi})} \dot{\psi}(k) \Gamma(k) \overline{\rho_{f, \infty}(D)} \rho_{f, 1/q}(1), \\ \mathcal{E} &= \sum_{\mathfrak{c} \text{ sing.}} \frac{1}{4\pi} \int_{-\infty}^{\infty} \frac{\tilde{\psi}(t)}{\cosh(\pi t)} \overline{\rho_{\mathfrak{c}, \infty}(D, t)} \rho_{\mathfrak{c}, 1/q}(1, t) dt, \\ \mathcal{M} &= \sum_{f \in \mathcal{B}(qD, \bar{\chi})} \frac{\tilde{\psi}(t_f)}{\cosh(\pi t_f)} \overline{\rho_{f, \infty}(D)} \rho_{f, 1/q}(1). \end{aligned}$$

Split $\mathcal{M} = \mathcal{M}_1 + \mathcal{M}_2$ where \mathcal{M}_1 is the contribution of those f with $t_f \in \mathbb{R}$, and \mathcal{M}_2 is the remainder contribution, which consists of f with $t_f \in [-i/4, i/4]$. Consider first \mathcal{M}_2 . Using the bound $|\tilde{\psi}(t_f)| \ll x^{2|t_f|}$, we obtain by the Cauchy-Schwarz inequality

$$\mathcal{M}_2 \ll \left(\sum_{f \in \mathcal{B}(qD, \bar{\chi})} x^{2|t_f|} |\rho_{f, \infty}(D)|^2 \right)^{1/2} \left(\sum_{f \in \mathcal{B}(qD, \bar{\chi})} x^{2|t_f|} |\rho_{f, 1/q}(1)|^2 \right)^{1/2}.$$

We have $q\sqrt{D} \leq x^{1/2-\varepsilon+\eta} \leq x^{1/2}$, so that Lemma 2.9 of [Top15] may be applied to both sums, which yields, as $x \rightarrow \infty$,

$$(3.2) \quad \mathcal{M}_2 \ll x^{o(1)} \left(\frac{\sqrt{xD}}{q} \right)^{4\theta} \left\{ 1 + \frac{\sqrt{D}}{q} \right\} \ll x^{2\eta} (x/q^2)^{2\theta}.$$

The contribution of \mathcal{E} , \mathcal{H} and \mathcal{M}_1 is handled by similar standard arguments [DI83, page 267], using instead Lemma 2.8 of [Top15]. We obtain

$$(3.3) \quad \mathcal{H} + \mathcal{E} + \mathcal{M}_1 \ll x^{o(1)} D^{1/2} \ll x^\eta$$

as $x \rightarrow \infty$. Grouping our bounds (3.3) and (3.2), we find

$$\sum_{q \leq x^{1/2-\varepsilon}} \left| \sum_{n \equiv 0 \pmod{q}} \phi\left(\frac{n}{x}\right) \chi(n) \text{Kl}(1, n) \right| \ll_{\varepsilon, \phi} x^{1+\eta-\varepsilon} + x^{1+2\eta-(1/2-2\theta)\varepsilon}.$$

By work of Kim-Sarnak [Kim03], we are ensured that $\theta \leq 7/64 < 1/4$, and therefore our claimed bound follows if $Q \leq x^{1/2-\varepsilon}$ and $0 < \eta \leq 3\varepsilon/16$. \square

From Proposition 3.1 and the convolution $\mu^2(n) = \sum_{d^2|n} \mu(d)$, we deduce the analogous bound with n restricted to square-free numbers.

Corollary 3.2. *In the setting and notations of Proposition 3.1, we have*

$$(3.4) \quad \sum_{q \leq x^{1/2-\varepsilon}} \left| \sum_{n \equiv 0 \pmod{q}} \mu^2(n) \phi\left(\frac{n}{x}\right) \chi(n) \text{Kl}(1, n) \right| \ll_{\varepsilon, \phi} x^{1-\eta}$$

for a possibly smaller value of $\eta > 0$, but depending on ε at most.

3.2. Sieve weights. To control the terms around the central point $d \in [x^{1/2-\varepsilon}, x^{1/2+\varepsilon}]$ we will introduce a short sieve weight to maintain the feature that our sums are essentially supported on integers free of small prime factors. This is also crucial for carrying out the approximation (2.1) as effectively as possible. We recall the construction of the β -sieve from [FI10], and a few of its relevant properties for our application. Let

$$\theta(n) = \sum_{d|n} \xi_d$$

be an upper-bound β -sieve of level y and dimension $\kappa \geq 0$, for the primes less than z . The parameters y , z and κ will be chosen later to be small powers of x with the condition that $z^2 \leq y$. (We will ultimately choose $z = x^{\varepsilon/(\log \varepsilon)^2}$ and $y = x^\varepsilon$.) In particular, we have $\xi_1 = 1$ and $|\xi_d| \leq 1$; (ξ_d) is supported on integers up to y free of prime factor $> z$, and we have $\theta(n) \geq 0$ for all $n \geq 1$. Moreover, for any multiplicative function f with $f(p) \in [0, \kappa]$, we have

$$(3.5) \quad \sum_d \frac{\xi_d f(d)}{d} \ll \prod_{p \leq z} \left(1 + \frac{f(p)}{p}\right)^{-1}.$$

For some technical simplifications, we will work with the smoothed version

$$\theta'(n) := \sum_{d|n} \xi_d \log\left(\frac{n}{d}\right) = \sum_{d|n} \theta\left(\frac{n}{d}\right) \Lambda(d).$$

Note that $\theta'(n) = \log n$ if $P^-(n) > z$.

The key property we use of θ' is the following estimate on averages of multiplicative functions weighted by the sieve weight.

Lemma 3.3. *Let the multiplicative function f be supported on squarefree numbers, with $0 \leq f(p) \leq \kappa$. Then*

$$(3.6) \quad \sum_{n \leq x} \theta'(n) f(n) \ll x \prod_{z < p \leq x} \left(1 + \frac{f(p)}{p}\right).$$

Proof. Denote S the sum on the left-hand side of (3.6). By definition of θ' , we have

$$S = \sum_{n \leq x} \sum_{p|n} f(n) \theta\left(\frac{n}{p}\right) \log p \leq \sum_{n \leq x} f(n) \theta(n) \sum_{p \leq x/n} f(p) \log p$$

by positivity of the summand. Therefore,

$$S \ll_\kappa x \sum_{n \leq x} \frac{\theta(n) f(n)}{n} \leq x \sum_{P^+(n) \leq x} \frac{\theta(n) f(n)}{n}.$$

Opening the convolution in $\theta(n)$, we arrive at

$$(3.7) \quad \begin{aligned} S &\ll_{\kappa} x \sum_{P^+(d) \leq x} \frac{\xi_d f(d)}{d} \sum_{\substack{P^+(n) \leq x \\ (n,d)=1}} \frac{f(n)}{n} \\ &= x \prod_{p \leq x} \left(1 + \frac{f(p)}{p}\right) \sum_{P^+(d) \leq x} \frac{\xi_d f(d)}{d} \prod_{p|d} \left(1 + \frac{f(p)}{p}\right)^{-1}. \end{aligned}$$

By (3.5), the sum over d above is $\ll_{\kappa} \prod_{p \leq x} (1 + \frac{f(p)}{p})^{-1}$, and our claimed bound follows. \square

4. ARGUMENT FOR THE UPPER BOUND

In what follows, we let $\varepsilon > 0$ be fixed, and $\eta > 0$ be a small parameter, to be chosen in terms of ε . We assume that $D \leq x^{\eta}$.

4.1. Initial setup. For a positive integer r , define the arithmetic functions $\tilde{\Lambda}_r(n)$ by

$$\tilde{\Lambda}_r(n) = \sum_{d|n} \mu(d) \left(\log \frac{\sqrt{n}}{d} \right)^r.$$

We note that $\tilde{\Lambda}_1(n) = \Lambda(n)$ and $\tilde{\Lambda}_2(n) = \Lambda_2(n) - \Lambda(n) \log n$. On squarefree n , $\tilde{\Lambda}_1$ is supported on primes, and $\tilde{\Lambda}_2$ is supported on products of exactly two primes.

Let ϕ be smooth bounded compactly supported function approximating the characteristic function $\mathbf{1}_{[0,1]}$ of the interval $[0, 1]$ from below, such that ϕ and $\mathbf{1}_{[0,1]}$ differ only on $[0, \delta]$ and $[1 - \delta, 1]$, for some small constant $\delta > 0$. If we can show that for any $\varepsilon > 0$

$$(4.1) \quad \left| \sum_n \mu^2(n) \phi\left(\frac{n}{x}\right) (\log n) \tilde{\Lambda}(n) \text{Kl}(1, n) \right| \ll_{\phi} \varepsilon x \log x + O_{\varepsilon}(x L(1, \chi) \log^2 x),$$

then, using the Weil bound to control the contribution from $n < \delta x$ or $n > (1 - \delta)x$, we see from partial summation that

$$\left| \sum_{p \leq x} \text{Kl}(1, p) \right| \ll \delta \pi(x) + O_{\delta}(\varepsilon \pi(x)) + O_{\delta, \varepsilon}(L(1, \chi) \log x).$$

Thus, by choosing δ small and then ε sufficiently small in terms of δ , we see it suffices to show (4.1) for any fixed $\varepsilon > 0$.

Since $\mu^2(n) \tilde{\Lambda}(n)$ is only supported on primes, we have

$$\sum_n \mu^2(n) \phi\left(\frac{n}{x}\right) (\log n) \tilde{\Lambda}(n) \text{Kl}(1, n) = \sum_{(n, D)=1} \mu^2(n) \phi\left(\frac{n}{x}\right) \theta'(n) \tilde{\Lambda}(n) \text{Kl}(1, n) + O_{\varepsilon}(x^{\varepsilon}).$$

For products of two primes, we note that for all $x \geq 2$,

$$(4.2) \quad \sum_{\substack{n \leq x \\ n \neq pq \text{ for all } p \neq q, \\ p, q \text{ prime}}} \tilde{\Lambda}_2(n) \ll x,$$

with the main contribution arising from pairs $n = pq^2$ (p, q primes), while

$$(4.3) \quad \sum_{\substack{n \leq x \\ (n, D) > 1}} \tilde{\Lambda}_2(n) \ll x \log \log x$$

for $2 \leq D \leq x$. Thus

$$(4.4) \quad \begin{aligned} &\sum_{\substack{p, q \\ p, q \text{ prime}}} \phi\left(\frac{pq}{x}\right) (\log pq) (\log p) (\log q) \text{Kl}(1, pq) \\ &= \sum_{\substack{p, q \\ p, q \text{ prime}}} \phi\left(\frac{pq}{x}\right) \theta'(pq) (\log p) (\log q) \text{Kl}(1, pq) + O(x (\log x) (\log z)) \\ &= \frac{1}{2} \sum_{(n, D)=1} \mu^2(n) \phi\left(\frac{n}{x}\right) \theta'(n) \tilde{\Lambda}_2(n) \text{Kl}(1, n) + O(x (\log x) (\log z)). \end{aligned}$$

We have used both (4.2) and (4.3) in the last line.

Thus, we see that to establish Theorem 1.1 and Theorem 1.2 it is sufficient to show for every $\varepsilon > 0$, every smooth compactly supported ϕ and each $r \in \{1, 2\}$ that

$$(4.5) \quad S(x) = S(x, r) = \sum_{(n, D)=1} \mu^2(n) \phi\left(\frac{n}{x}\right) \theta'(n) \tilde{\Lambda}_r(n) \text{Kl}(1, n) \ll_{\phi} \varepsilon x \log^r x + O_{\varepsilon}(x L(1, \chi) \log^{r+1} x),$$

where D and x satisfy the hypotheses of Theorem 1.1 and χ is any real primitive character (mod D). Let

$$\nu(n) = (\mu * \mu \chi)(n) = \sum_{n=uv} \mu(u) \mu(v) \chi(v).$$

We write

$$(4.6) \quad \tilde{\Lambda}_r(n) = \frac{1}{2^r} \sum_{n=abc} \nu(b) \chi(a) \left(\log \frac{ab}{c} \right)^r,$$

so that for $r \in \{1, 2\}$,

$$(4.7) \quad S(x) = \frac{1}{r 2^r} \sum_{\substack{(a, b, c) \in \mathbb{N}^3 \\ (abc, D)=1}} \mu^2(abc) \phi\left(\frac{abc}{x}\right) \theta'(abc) \nu(b) \chi(a) \left(\log \frac{ab}{c} \right)^r \text{Kl}(1, abc).$$

We split the sum over (a, b, c) into $b > 1$ or $b = 1$ and $\min\{a, c\} \leq x^{1/2}/y^3$, or $b = 1$ and $\min\{a, c\} > x^{1/2}/y^3$. This gives

$$(4.8) \quad r 2^r S(x) = S_L(x) + S_N(x) + S_C(x),$$

with

$$(4.9) \quad S_L(x) = \sum_{\substack{(a, b, c) \in \mathbb{N}^3 \\ b > 1 \\ (abc, D)=1}} \mu^2(abc) \phi\left(\frac{abc}{x}\right) \theta'(abc) \nu(b) \chi(a) \left(\log \frac{ab}{c} \right)^r \text{Kl}(1, abc),$$

$$(4.10) \quad S_N(x) = \sum_{\substack{(a, c) \in \mathbb{N}^2 \\ \min\{a, c\} \leq \sqrt{x}/y^3 \\ (ac, D)=1}} \mu^2(ac) \phi\left(\frac{ac}{x}\right) \theta'(ac) \chi(a) \left(\log \frac{a}{c} \right)^r \text{Kl}(1, ac),$$

$$(4.11) \quad S_C(x) = \sum_{\substack{(a, c) \in \mathbb{N}^2 \\ \min\{a, c\} > \sqrt{x}/y^3 \\ (ac, D)=1}} \mu^2(ac) \phi\left(\frac{ac}{x}\right) \theta'(ac) \chi(a) \left(\log \frac{a}{c} \right)^r \text{Kl}(1, ac).$$

We can control $S_L(x)$ by $L(1, \chi)$, since $\mu^2 \nu$ is supported on integers with all prime factors satisfying $\chi(p) = 1$ and θ' is concentrated on integers free of small prime factors. Thus if $L(1, \chi)$ is small, we expect the support to be a lacunary sequence and so the sum $S_L(x)$ to be small.

We can control $S_N(x)$ by the level of distribution estimates of Section 3.1, since the sum over one of a or c is a long sum.

Finally, S_C is the contribution near the central point, and we will show this is small since $|\log(a/c)| \leq 6 \log y + O_{\phi}(1)$, which is small compared with $\log x$, a is restricted to a short range on the logarithmic scale, and $\text{Kl}(1, n)$ is typically small on numbers with many prime factors.

4.2. Bounding the lacunary sum S_L . We begin by bounding $S_L(x)$ in terms of $L(1, \chi)$.

Lemma 4.1. *For all small enough $\varepsilon > 0$, and $D^8 \leq x^{\varepsilon^2} \leq z \leq y^{1/(\log \varepsilon)^2}$, we have*

$$S_L(x) \ll_{\phi} x (\log x)^r \left(\varepsilon^{-\kappa-2} L(1, \chi) \log x + \varepsilon \right),$$

Proof. We note that for squarefree n , we have $|\nu(n)| = (1 * \chi)(n)$, and we recall that $\theta'(n) = (\Lambda * \theta)(n)$. Thus, bounding the summand of $S_L(x)$ and letting $d = ac$, and then expanding the definition of θ , we

find

$$\begin{aligned} S_L(x) &\ll (\log x)^r \sum_{\substack{db \ll x \\ b > 1}} (1 * \chi)(b) \theta'(bd) \tau(bd)^2 \mu(bd)^2 \\ &= (\log x)^r \sum_{\substack{bef \ll x \\ b > 1}} (1 * \chi)(b) \theta(be) \Lambda(f) \tau(bef)^2 \mu(bef)^2. \end{aligned}$$

When $f > z$ is prime we see $\theta(be) = \theta(bef)$. Thus the contribution from $f > z$ can be bounded by

$$(4.12) \quad (\log x)^{r+1} \sum_{\substack{bg \ll x \\ b > 1}} (1 * \chi)(b) \theta(bg) \mu^2(bg) \tau(bg)^4.$$

The contribution from $f \leq z$ can be bounded by

$$(4.13) \quad (\log x)^r \sum_{f \ll z} \Lambda(f) \sum_{\substack{be \ll x/f \\ b > 1}} (1 * \chi)(b) \theta(be) \mu^2(be) \tau(be)^2.$$

These are precisely sums of the type considered in Section 24.7 of [FI10]. Indeed, there it is established [FI10, equation (24.58)] that if $D^8 \leq z \leq y \leq Y^{1/8}$ and ρ is a completely multiplicative function with $0 \leq \rho(p) \leq \kappa/6$ then

$$\sum_{\substack{ab \leq Y \\ b > 1}} (1 * \chi)(b) \theta(ab) \rho(ab) \ll \frac{Y}{\log Y} \left(L(1, \chi) \log Y + e^{-t} \right) \left(\frac{\log Y}{\log z} \right)^{\kappa/2+1},$$

where $t = \log y / \log z$. Since $\mu^2(n) \tau(n)^k \leq 2^{k\Omega(n)}$ ($k = 2, 4$), we may apply this to the bounds (4.12) and (4.13) above, and find that provided $D^8 \leq z \leq y \leq x^{1/9}$ and $\kappa > 6 \times 2^4$, we have

$$\begin{aligned} (\log x)^{r+1} \sum_{\substack{bg \ll x \\ b > 1}} (1 * \chi)(b) \theta(bg) \tau(bg)^4 &\ll x (\log x)^r \left(L(1, \chi) \log x + e^{-t} \right) \left(\frac{\log x}{\log z} \right)^{\kappa/2+1}, \\ (\log x)^r \sum_{f \ll z} \Lambda(a) \sum_{\substack{be \ll x/f \\ b > 1}} (1 * \chi)(b) \theta(be) \tau(be)^2 &\ll (\log x)^{r-1} \left(L(1, \chi) \log x + e^{-t} \right) \left(\frac{\log x}{\log z} \right)^{\kappa/2+1} \sum_{f \ll z} \frac{\Lambda(f)}{f} \\ &\ll x (\log x)^r \left(L(1, \chi) \log x + e^{-t} \right) \left(\frac{\log x}{\log z} \right)^{\kappa/2}. \end{aligned}$$

Putting this together, we find

$$S_L(x) \ll x (\log x)^r \left(L(1, \chi) \log x + e^{-t} \right) \left(\frac{\log x}{\log z} \right)^{\kappa/2+1}.$$

If $x^{\varepsilon^2} \leq z \leq y^{1/(\log \varepsilon)^2}$, then this gives

$$(4.14) \quad S_L(x) \ll x (\log x)^r \left(\varepsilon^{-\kappa-2} L(1, \chi) \log x + \varepsilon \right),$$

as required. \square

4.3. Bounding the non-central sum S_N .

Lemma 4.2. *Let $y \geq x^\varepsilon$. Then there is a constant $\eta > 0$ depending only on ε such that*

$$S_N(x) \ll_\varepsilon x^{1-\eta}.$$

Proof. Denote by $\chi_0(n)$ the principal character (mod D). Opening the summation in $\theta'(ac)$, we obtain

$$\begin{aligned} S_N(x) &\leq \sum_{d \leq y} \sum_{a < \sqrt{x}/y^3} \left| \sum_{n \equiv 0 \pmod{[a,d]}} \mu^2(n) \phi\left(\frac{n}{x}\right) \chi_0(n) \left(\log\left(\frac{a^2}{n}\right) \right)^r \left(\log \frac{n}{d} \right) \text{Kl}(1, n) \right| \\ &\quad + \sum_{d \leq y} \sum_{c < \sqrt{x}/y^3} \left| \sum_{n \equiv 0 \pmod{[c,d]}} \mu^2(n) \phi\left(\frac{n}{x}\right) \chi(n) \left(\log\left(\frac{c^2}{n}\right) \right)^r \left(\log \frac{n}{d} \right) \text{Kl}(1, n) \right| \\ (4.15) \quad &\ll \sup_{\chi_1 \pmod{D}} \sup_{0 \leq j \leq r+1} \sum_{q \leq \sqrt{x}/y} \tau(q) (\log x)^{r+1} \left| \sum_{n \equiv 0 \pmod{q}} \mu^2(n) \phi\left(\frac{n}{x}\right) \chi_1(n) (\log(n/x))^j \text{Kl}(1, n) \right|. \end{aligned}$$

Recalling that we assume $y \geq x^\varepsilon$ and applying Corollary 3.2, we obtain

$$(4.16) \quad S_N(x) \ll_\varepsilon x^{1-\eta}$$

for some η depending at most on ε . □

4.4. Bounding the central point sum S_C .

Lemma 4.3. *Let $x^{\varepsilon^2} \leq z \leq y^{1/(\log \varepsilon)^2}$ and $y \leq x^{2\varepsilon}$. Then we have*

$$S_C(x) \ll \varepsilon^{10} x (\log x)^r + \varepsilon^{3/10} x (\log x)^r \left(\frac{\log y}{\log z} \right)^2.$$

Proof. Recall that the sum $S_C(x)$ is defined in (4.11). By construction, for all (a, c) in the summation range, we have $|\log(a/c)| \ll \log y$.

Therefore, by the triangle inequality and the support condition on ϕ , our task is to bound

$$(\log y)^r \sum_{\substack{x/y^3 \leq a \ll xy^3 \\ ac \asymp x}} \mu^2(ac) \theta'(ac) |\text{Kl}(1, ac)|.$$

Following an idea of Hooley [Hoo64], we let a parameter $u \in [y, x^{1/4}]$ be given, and decompose accordingly

$$a = a_1 a_2, \quad P^+(a_1) \leq u, \quad P^-(a_2) > u,$$

and similarly let $c = c_1 c_2$. Notice that

$$\theta'(ac) = \theta(a_1 c_1) \log(a_2 c_2) + \theta'(a_1 c_1) \ll \theta(a_1 c_1) \log x + \theta'(a_1 c_1)$$

since $P^-(a_2 c_2) > u > z$. We write

$$(4.17) \quad \sum_{\substack{a_1 a_2 c_1 c_2 \ll x \\ x^{1/2}/y^3 < a_1 a_2 \ll x^{1/2} y^3 \\ P^+(a_1 c_1) \leq u < P^-(a_2 c_2)}} \mu^2(a_1 a_2 c_1 c_2) \theta'(a_1 a_2 c_1 c_2) |\text{Kl}(1, a_1 a_2 c_1 c_2)| = S_1 + S_2,$$

where S_1 is the contribution of $a_1 c_1 \leq x^{1/10}$, and S_2 is the contribution of $a_1 c_1 > x^{1/10}$.

Concerning S_1 , we have

$$S_1 \ll \sum_{\substack{a_1 c_1 \leq x^{1/10} \\ P^+(a_1 c_1) \leq u}} \mu^2(a_1 c_1) \{ \theta(a_1 c_1) \log x + \theta'(a_1 c_1) \} T(a_1, c_1),$$

with

$$T(a_1, c_1) = \sum_{\substack{a_2 c_2 \ll x/a_1 c_1 \\ P^-(a_2 c_2) > u \\ x^{1/2}/y^3 < a_1 a_2 \ll x^{1/2} y^3}} \mu^2(a_2 c_2) |\text{Kl}(1, a_1 a_2 c_1 c_2)|.$$

By a reasoning identical to page 277 of [FM03b], for $a_1 c_1 \leq x^{1/10}$, we have

$$\begin{aligned} T(a_1, c_1) &\leq \sum_{\substack{b \pmod{a_1 c_1} \\ (b, a_1 c_1) = 1}} |\text{Kl}(b^2, a_1 c_1)| \sum_{\substack{a_2 c_2 \ll x/a_1 c_1 \\ x^{1/2}/a_1 y^3 < a_2 \ll x^{1/2} y^3/a_1 \\ P^-(a_2 c_2) > u \\ b a_2 c_2 \equiv 1 \pmod{a_1 c_1}}} \mu^2(a_2 c_2) 2^{\omega(a_2 c_2)} \\ &\ll \sum_{\substack{b \pmod{a_1 c_1} \\ (b, a_1 c_1) = 1}} |\text{Kl}(b^2, a_1 c_1)| \frac{x}{a_1 c_1 \varphi(a_1 c_1)} \frac{1}{\log x} \left(\frac{\log x}{\log u} \right)^2 \sum_{\substack{x^{1/2}/a_1 y^3 < a_2 \ll x^{1/2} y^3/a_1 \\ P^-(a_2) > u}} \frac{2^{\omega(a_2)}}{a_2} \\ &\ll \frac{x}{a_1 c_1 \varphi(a_1 c_1)} \frac{\log y}{\log^2 x} \left(\frac{\log x}{\log u} \right)^4 \sum_{\substack{b \pmod{a_1 c_1} \\ (b, a_1 c_1) = 1}} |\text{Kl}(b^2, a_1 c_1)| \\ &= \frac{x \log y}{\log^2 x} \left(\frac{\log x}{\log u} \right)^4 \frac{2^{\omega(a_1 c_1)} \kappa(a_1 c_1)}{a_1 c_1} \end{aligned}$$

for a multiplicative function κ . Fouvry and Michel [FM03b, equation (3.4)] show that

$$\kappa(p) = \frac{4}{3\pi} + O(p^{-1/4}),$$

which they deduce from Katz's Sato-Tate law for Kloosterman sums [Kat88]. We insert this back into S_1 , and we relax the summation conditions $a_1 c_1 \leq x^{1/10}$. Letting

$$f(n) := \mu^2(n) 4^{\omega(n)} \kappa(n),$$

we obtain

$$(4.18) \quad S_1 \ll \frac{x \log y}{\log^2 x} \left(\frac{\log x}{\log u} \right)^4 \sum_{P^+(n) \leq u} \frac{(\theta(n) \log x + \theta'(n)) f(n)}{n}.$$

Note that

$$f(p) = \frac{16}{3\pi} + O(p^{-1/4}) \quad (p \leq u).$$

Taking Euler products, computations similar to (3.7) yield

$$(4.19) \quad \sum_{P^+(n) \leq u} \frac{\theta(n) f(n)}{n} \ll \prod_{z < p \leq u} \left(1 + \frac{f(p)}{p} \right) \ll \left(\frac{\log u}{\log z} \right)^{16/(3\pi)}.$$

On the other hand,

$$\sum_{P^+(n) \leq u} \frac{\theta'(n) f(n)}{n} = \sum_{\substack{q \leq u \\ q \text{ prime}}} \frac{f(q) \log q}{q} \sum_{\substack{P^+(n) \leq u \\ (n, q) = 1}} \frac{\theta(n) f(n)}{n} \ll (\log u) \left(\frac{\log u}{\log z} \right)^{16/(3\pi)}$$

by dropping the condition $(n, q) = 1$ and using (4.19). Inserting in (4.18), we find

$$(4.20) \quad S_1 \ll x \left(\frac{\log y}{\log x} \right) \left(\frac{\log x}{\log u} \right)^4 \left(\frac{\log u}{\log z} \right)^{16/(3\pi)}.$$

Consider now the contribution S_2 to (4.17). Using the Weil bound $|\text{Kl}(1, n)| \leq 2^{\omega(n)}$, we have

$$S_2 \leq \sum_{n \ll x} \theta'(n) g_0(n),$$

where

$$g_0(n) := \begin{cases} \mu^2(n) 4^{\omega(n)} & \text{if } \prod_{p|n, p \leq u} p > x^{1/4}, \\ 0 & \text{otherwise.} \end{cases}$$

By Rankin's inequality, we have $g_0(n) \leq x^{-1/(4 \log u)} g_1(n)$, where

$$g_1(n) := \mu^2(n) 4^{\omega(n)} \prod_{p|n, p \leq u} p^{1/\log u}.$$

Note that the function g_1 is multiplicative with

$$g(p) = \begin{cases} 4p^{1/\log u} & \text{if } p \leq u \\ 4 & \text{if } p > u. \end{cases}$$

In any case we have $g(p) \leq 4e$. Assuming $\kappa \geq 4e$, we obtain

$$S_2 \leq x^{-1/(4 \log u)} \sum_{n \ll x} \theta'(n) g_1(n) \ll x \left(\frac{\log x}{\log u} \right)^4 \left(\frac{\log u}{\log z} \right)^{4e} \exp \left\{ -\frac{\log x}{4 \log u} \right\}.$$

We now choose $u = x^{1/(\log \varepsilon)^2}$. Recalling that we have assumed $x^{\varepsilon^2} \leq z$, we see that this gives

$$(4.21) \quad S_2(x) \ll \varepsilon^{10} x.$$

Similarly, we substitute $u = x^{1/(\log \varepsilon)^2}$ into (4.20). Using the assumption $z \leq y \leq x^{2\varepsilon}$ and noting that $2 - 16/3\pi > 3/10$, we find

$$(4.22) \quad \begin{aligned} (\log y) S_1 &\ll x \log x \left(\log \frac{1}{\varepsilon} \right)^4 \left(\frac{\log y}{\log x} \right)^{2-16/3\pi} \left(\frac{\log y}{\log z} \right)^{16/3\pi} \\ &\ll \varepsilon^{3/10} x \log x \left(\frac{\log y}{\log z} \right)^2. \end{aligned}$$

Recalling that $S_C(x) \ll (S_1 + S_2)(\log y)^r$, that $r \in \{1, 2\}$, and inserting the bounds (4.22) and (4.21), we obtain

$$S_C(x) \ll \varepsilon^{10} x (\log x)^r + \varepsilon^{3/10} x (\log x)^r \left(\frac{\log y}{\log z} \right)^2,$$

as required. □

4.5. Conclusion. We choose

$$y = x^\varepsilon, \quad z = x^{\varepsilon/(\log \varepsilon)^2}.$$

With this choice, we see that y and z satisfy the assumptions of lemmas 4.1, 4.2 and 4.3. Thus, provided $D^8 \leq x^{\varepsilon^2}$ we may apply these Lemmas, giving

$$\begin{aligned} S(x) &\ll S_L(x) + S_N(x) + S_C(x) \\ &\ll \varepsilon^{-\kappa-2} x L(1, \chi) (\log x)^{r+1} + \varepsilon x (\log x)^r + O_\varepsilon(x^{1-\eta}) + \varepsilon^{3/10} x (\log x)^r \left(\log \frac{1}{\varepsilon}\right)^4 \end{aligned}$$

for some quantity $\eta > 0$ depending only on ε . Reinterpreting ε , we obtain the claimed statement (4.5), which then gives Theorems 1.1 and Theorem 1.2.

REFERENCES

- [BM15] V. Blomer and D. Milićević, *Kloosterman sums in residue classes*, J. Eur. Math. Soc. (JEMS) **17** (2015), no. 1, 51–69.
- [Bom76] E. Bombieri, *On twin almost primes*, Acta Arith. **28** (1975/76), no. 2, 177–193.
- [DI83] J.-M. Deshouillers and H. Iwaniec, *Kloosterman sums and Fourier coefficients of cusp forms*, Invent. Math. **70** (1982/83), no. 2, 219–288.
- [Dra17] S. Drappeau, *Sums of Kloosterman sums in arithmetic progressions, and the error term in the dispersion method*, Proc. London Math. Soc. **114** (2017), no. 4, 684–732.
- [FKM14] É. Fouvry, E. Kowalski, and P. Michel, *Algebraic trace functions over the primes*, Duke Math. J. **163** (2014), no. 9, 1683–1736.
- [FM03a] É. Fouvry and P. Michel, *Crible asymptotique et sommes de Kloosterman*, Proceedings of the Session in Analytic Number Theory and Diophantine Equations, Bonner Math. Schriften, vol. 360, Univ. Bonn, Bonn, 2003, p. 27.
- [FM03b] ———, *Sommes de modules de sommes d’exponentielles*, Pacific J. Math. **209** (2003), no. 2, 261–288.
- [FM06] ———, *Errata to the article: “Sums of moduli of exponential sums” (French) [Pacific J. Math. **209** (2003), no. 2, 261–288; mr1978371]*, Pacific J. Math. **225** (2006), no. 1, 199–200.
- [FM07] ———, *Sur le changement de signe des sommes de Kloosterman*, Ann. of Math. (2) **165** (2007), no. 3, 675–715.
- [FI03] J. B. Friedlander and H. Iwaniec, *Exceptional characters and prime numbers in arithmetic progressions*, Int. Math. Res. Not. (2003), no. 37, 2033–2050.
- [FI04] ———, *Exceptional characters and prime numbers in short intervals*, Selecta Math. (N.S.) **10** (2004), no. 1, 61–69.
- [FI05] ———, *The illusory sieve*, Int. J. Number Theory **1** (2005), no. 4, 459–494.
- [FI10] ———, *Opera de cribro*, American Mathematical Society Colloquium Publications, vol. 57, American Mathematical Society, Providence, RI, 2010.
- [FI13] ———, *Exceptional discriminants are the sum of a square and a prime*, Q. J. Math. **64** (2013), no. 4, 1099–1107.
- [HB83] D. R. Heath-Brown, *Prime twins and Siegel zeros*, Proc. London Math. Soc. (3) **47** (1983), no. 2, 193–224.
- [Hoo64] C. Hooley, *On the distribution of the roots of polynomial congruences*, Mathematika **11** (1964), 39–49.
- [Kat88] N. M. Katz, *Gauss sums, Kloosterman sums, and monodromy groups*, Annals of Mathematics Studies, vol. 116, Princeton University Press, Princeton, NJ, 1988.
- [Kim03] H. H. Kim, *Functoriality for the exterior square of GL_4 and the symmetric fourth of GL_2* , J. Amer. Math. Soc. **16** (2003), no. 1, 139–183, With appendix 1 by D. Ramakrishnan and appendix 2 by H. Kim and P. Sarnak.
- [Klo27] H. D. Kloosterman, *On the representation of numbers in the form $ax^2 + by^2 + cz^2 + dt^2$* , Acta Math. **49** (1927), no. 3–4, 407–464.
- [Kuz80] N. V. Kuznetsov, *The Petersson conjecture for cusp forms of weight zero and the Linnik conjecture. Sums of Kloosterman sums*, Mat. Sb. (N.S.) **111(153)** (1980), no. 3, 334–383, 479.
- [LRS95] W. Luo, Z. Rudnick, and P. Sarnak, *On Selberg’s eigenvalue conjecture*, Geom. Funct. Anal. **5** (1995), no. 2, 387–401.
- [Mat11] K. Matomäki, *A note on signs of Kloosterman sums*, Bull. Soc. Math. France **139** (2011), no. 3, 287–295.
- [Poi11] H. Poincaré, *Fonctions modulaires et fonctions fuchsienues*, Ann. Fac. Sci. Toulouse Sci. Math. Sci. Phys. (3) **3** (1911), 125–149.
- [Sie35] C. L. Siegel, *Über die Classenzahl quadratischer Zahlkörper.*, Acta Arith. **1** (1935), 83–86.
- [SF07] J. Sivak-Fischler, *Crible étrange et sommes de Kloosterman*, Acta Arith. **128** (2007), no. 1, 69–100.
- [SF09] ———, *Crible asymptotique et sommes de Kloosterman*, Bull. Soc. Math. France **137** (2009), no. 1, 1–62.
- [Top15] B. Topalogullari, *On a certain additive divisor problem*, preprint (2015), arXiv, <http://arxiv.org/abs/1512.05770v2>.
- [Wei48] A. Weil, *On some exponential sums*, Proc. Nat. Acad. Sci. U. S. A. **34** (1948), 204–207.
- [Xi15a] P. Xi, *Sign changes of Kloosterman sums with almost prime moduli*, Monatsh. Math. **177** (2015), no. 1, 141–163.
- [Xi15b] ———, *Sign changes of Kloosterman sums with almost prime moduli. II*, Int. Math. Res. Notices (2015), to appear.
- [Xi18] ———, *When Kloosterman sums meet Hecke eigenvalues*, preprint (2018), arXiv, <http://arxiv.org/abs/1801.07658.v2>.

AIX MARSEILLE UNIVERSITÉ, CNRS, CENTRALE MARSEILLE, I2M UMR 7373, 13453 MARSEILLE, FRANCE
E-mail address: `sary-aurelien.drappeau@univ-amu.fr`

MATHEMATICAL INSTITUTE, RADCLIFFE OBSERVATORY QUARTER, WOODSTOCK ROAD, OXFORD OX2 6GG, UNITED KINGDOM
E-mail address: `james.alexander.maynard@gmail.com`