



HAL
open science

Eliminating Reflection from Type Theory

Théo Winterhalter, Matthieu Sozeau, Nicolas Tabareau

► **To cite this version:**

Théo Winterhalter, Matthieu Sozeau, Nicolas Tabareau. Eliminating Reflection from Type Theory. CPP 2019 - 8th ACM SIGPLAN International Conference on Certified Programs and Proofs, Jan 2019, Lisbonne, Portugal. pp.91-103, 10.1145/3293880.3294095 . hal-01849166v3

HAL Id: hal-01849166

<https://hal.science/hal-01849166v3>

Submitted on 22 Nov 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Eliminating Reflection from Type Theory

To the Legacy of Martin Hofmann

Théo Winterhalter
Gallinette Project-Team, Inria
Nantes, France
theo.winterhalter@inria.fr

Matthieu Sozeau
Pi.R2 Project-Team, Inria and IRIF
Paris, France
matthieu.sozeau@inria.fr

Nicolas Tabareau
Gallinette Project-Team, Inria
Nantes, France
nicolas.tabareau@inria.fr

Abstract

Type theories with equality reflection, such as extensional type theory (ETT), are convenient theories in which to formalise mathematics, as they make it possible to consider provably equal terms as convertible. Although type-checking is undecidable in this context, variants of ETT have been implemented, for example in NuPRL and more recently in Andromeda. The actual objects that can be checked are not proof-terms, but derivations of proof-terms. This suggests that any derivation of ETT can be translated into a typecheckable proof term of intensional type theory (ITT). However, this result, investigated categorically by Hofmann in 1995, and 10 years later more syntactically by Oury, has never given rise to an effective translation. In this paper, we provide the first effective syntactical translation from ETT to ITT with uniqueness of identity proofs and functional extensionality. This translation has been defined and proven correct in Coq and yields an executable plugin that translates a derivation in ETT into an actual Coq typing judgment. Additionally, we show how this result is extended in the context of homotopy type theory to a two-level type theory.

Keywords dependent types, translation, formalisation

1 Introduction

Type theories with equality reflection, such as extensional type theory (ETT), are convenient theories in which to formalise mathematics, as they make it possible to consider provably equal terms as convertible, as expressed in the following typing rule:

$$\frac{\Gamma \vdash_x e : u =_A v}{\Gamma \vdash_x u \equiv v : A} \quad (1)$$

Here, the type $u =_A v$ is Martin-Löf's identity type with only one constructor $\text{refl } u : u =_A u$ which represents proofs

Permission to make digital or hard copies of part or all of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for third-party components of this work must be honored. For all other uses, contact the owner/author(s).

Conference'17, July 2017, Washington, DC, USA

© 2018 Copyright held by the owner/author(s).

ACM ISBN 978-x-xxxx-xxxx-x/YY/MM.

<https://doi.org/10.1145/nnnnnnn.nnnnnnn>

of equality inside type theory, whereas $u \equiv v : A$ means that u and v are convertible in the theory—and can thus be silently replaced by one another in any term. Several variants of ETT have been considered and implemented, for example in NuPRL [Allen et al. 2000] and more recently in Andromeda [Bauer et al. 2016]. The prototypical example of the use of equality reflection is the definition of a coercion function between two types A and B that are equal (but not convertible) by taking a term of type A and simply returning it as a term of type B :

$$\lambda A B (e : A = B) (x : A). x : \Pi A B. A = B \rightarrow A \rightarrow B.$$

In intensional type theory (ITT), this term does not type-check because x of type A can not be given the type B by conversion. In ETT, however, equality reflection can be used to turn the witness of equality into a proof of conversion and thus the type system validates the fact that x can be given the type B . This means that one needs to guess equality proofs during type-checking, because the witness of equality has been lost at the application of the reflection rule. Guessing it was not so hard in this example but is in general undecidable, as one can for instance encode the halting problem of any Turing machine as an equality in ETT. That is, the actual objects that can be checked in ETT are not terms, but instead derivations of terms. It thus seems natural to wonder whether any derivation of ETT can be translated into a typecheckable term of ITT. And indeed, it is well known that one can find a corresponding term of the same type in ITT by *explicitly* transporting the term x of type A using the elimination of internal equality on the witness of equality e , noted e_* :

$$\lambda A B (e : A = B) (x : A). e_* x : \Pi A B. A = B \rightarrow A \rightarrow B.$$

This can be seen as a way to make explicit the *silent* use of reflection. Furthermore, by making the use of transport as economic as possible, the corresponding ITT term can be seen as a compact witness of the derivation tree of the original ETT term.

This result has first been investigated categorically in the pioneering work of Hofmann [1995, 1997], by showing that the term model of ITT can be turned into a model of ETT by quotienting this model with propositional equality. However, it is not clear how to extend this categorical construction to an explicit and constructive translation from a derivation in ETT to a term of ITT. In 2005, this result has been investigated more syntactically by Oury [2005]. However, his

111 presentation does not give rise to an effective translation.
 112 By an *effective* translation we mean that it is entirely con-
 113 structive and can be used to deterministically *compute* the
 114 translation of a given ETT typing derivation. Two issues
 115 prevent deriving an *effective* translation from Oury's presen-
 116 tation, and it is the process of actual formalisation of the
 117 result in a proof assistant that led us to these discoveries.
 118 First, his handling of related contexts is not explicit enough,
 119 which we fix by framing the translation using ideas coming
 120 from the parametricity translation (section 1.2). Addition-
 121 ally, Oury's proof requires an additional axiom in ITT on
 122 top of functional extensionality and uniqueness of identity
 123 proofs, that has no clear motivation and can be avoided by
 124 considering an annotated syntax (section 2.1).

125 **Contributions.** In this paper, we present the first effective
 126 syntactical translation from ETT to ITT (assuming unique-
 127 ness of identity proofs (UIP) and functional extensionality in
 128 ITT). By syntactical translation, we mean an explicit transla-
 129 tion from a derivation $\Gamma \vdash_x t : T$ of ETT (the x index testifies
 130 that it is a derivation in ETT) to a context Γ' , term t' and
 131 type T' of ITT such that $\Gamma' \vdash t' : T'$ in ITT. This translation
 132 enjoys the additional property that if T can be typed in ITT,
 133 *i.e.*, $\Gamma \vdash T$, then $T' \equiv T$. This means in particular that a theo-
 134 rem proven in ETT but whose statement is also valid in ITT
 135 can be automatically transferred to a theorem of ITT. For
 136 instance, one could use a *local* extension of the Coq proof
 137 assistant with a reflection rule, without being forced to rely
 138 on the reflection in the entire development.
 139

140 This translation can be seen as a way to build a syntac-
 141 tical model of ETT from a model of ITT as described more
 142 generally in Boulier et al. [2017] and has been entirely pro-
 143 grammed and formalised in Coq [Coq development team
 144 2017]. For this, we rely on TemplateCoq¹ [Anand et al. 2018],
 145 which provides a reifier for Coq terms as represented in
 146 Coq's kernel as well as a formalisation of the type system of
 147 Coq. Thus, our formalisation of ETT is just given by adding
 148 the reflection rule to a subset of the original type system
 149 of Coq. This allows us to extract concrete Coq terms and
 150 types from a closed derivation of ETT, using a little trick to
 151 incorporate Inductive types and induction. We do not treat
 152 cumulativity of universes which is an orthogonal feature of
 153 Coq's type theory.

154 **Outline of the Paper.** Before going into the technical devel-
 155 opment of the translation, we explain its main ingredients
 156 and differences with previous works. Then, in Section 2,
 157 we define the extensional and intensional type theories we
 158 consider. In Section 3, we define the main ingredient of the
 159 translation, which is a relation between terms of ETT and
 160 terms in ITT. Then, the translation is given in Section 4. Sec-
 161 tion 5 describes the Coq formalisation and Sections 6 and 7

166 discuss limitations and related work. The main proofs are
 167 given in detail in Appendices B and C.

168 The Coq formalisation can be found in [https://github.com/](https://github.com/TheoWinterhalter/ett-to-itt)
 169 [TheoWinterhalter/ett-to-itt](https://github.com/TheoWinterhalter/ett-to-itt).
 170

171 1.1 On the Need for UIP and Functional 172 Extensionality.

173 Our translation targets ITT plus UIP and functional exten-
 174 sionality, which correspond to the two following axioms
 175 (where \square_i denotes the universe of types at level i):
 176

$$176 \text{ UIP} : \Pi(A : \square_i) (x y : A) (e e' : x = y). e = e'$$

$$177 \text{ FunExt} : \Pi(A : \square_i) (B : A \rightarrow \square_i) (f g : \Pi(x : A). B x).$$

$$178 (\Pi(x : A). f x = g x) \rightarrow f = g$$

179 The first axiom says that any two proofs of the same equality
 180 are equal, and the other one says that two (dependent) func-
 181 tions are equal whenever they are pointwise equal². These
 182 two axioms are perfectly valid statements of ITT and they
 183 can be proven in ETT. Indeed, UIP can be shown to be equiv-
 184 alent to the Streicher's axiom K
 185

$$186 \text{ K} : \Pi(A : \square_i). \Pi(x : A). \Pi(e : x = x). e = \text{refl}_x$$

187 using the elimination on the identity type. But K is provable
 188 in ETT by considering the type
 189

$$190 \Pi(A : \square_i). \Pi(x y : A). \Pi(e : x = y). e = \text{refl}_x$$

191 which is well typed (using the reflection rule to show that e
 192 has type $x = x$) and which can be inhabited by elimination of
 193 the identity type. In the same way, functional extensionality
 194 is provable in ETT because
 195

$$196 \Pi(x : A). f x = g x$$

$$197 \rightarrow x : A \vdash f x \equiv g x \quad \text{by reflection}$$

$$198 \rightarrow (\lambda(x : A). f x) \equiv (\lambda(x : A). g x) \quad \text{by congruence of } \equiv$$

$$199 \rightarrow f \equiv g \quad \text{by } \eta\text{-law}$$

$$200 \rightarrow f = g$$

201 Therefore, applying our translation to the proofs of those
 202 theorems in ETT gives corresponding proofs of the same
 203 theorems in ITT. However, UIP is independent from ITT, as
 204 first shown by Hofmann and Streicher using the groupoid
 205 model [Hofmann and Streicher 1998], which has recently
 206 been extended in the setting of univalent type theory using
 207 the simplicial or cubical models [Bezem et al. 2013; Kapulkin
 208 and Lumsdaine 2012]. Similarly, Boulier *et al.* have shown
 209 that functional extensionality is independent from ITT using
 210 a simple syntactical translation [Boulier et al. 2017].

211 Therefore, our translation provides proofs of axioms in-
 212 dependent from ITT, which means that the target of the
 213 translation already needs to have both UIP and functional
 214 extensionality. Part of our work is to show formally that they
 215 are the only axioms required.
 216

217 ²In Homotopy Type Theory (HoTT) [Univalent Foundations Program 2013],
 218 the functional extensionality axiom is stated in a more complete way, using
 219 the notion of adjoint equivalences, but this more complete way collapses to
 220 our simpler statement in presence of UIP.

163 ¹<https://template-coq.github.io/template-coq/>

1.2 Heterogeneous Equality and the Parametricity Translation.

The basic idea behind the translation from ETT to ITT is to interpret conversion using the internal notion of equality, *i.e.*, the identity type. But this means that two terms of two convertible types that were comparable in ETT become comparable in ITT only up-to the equality between the two types. One possible solution to this problem is to consider a native heterogeneous equality, such as *John Major equality* introduced by McBride [2000]. However, to avoid adding additional axioms to ITT as done by Oury [2005], we prefer to encode this heterogeneous equality using the following dependent sums:

$$t \text{ }_{T \cong U} \cong u := \Sigma(p : T = U). p_* t = u.$$

During the translation, the same term occurring twice can be translated in two different manners, if the corresponding typing derivations are different. Even the types of the two different translations may be different. However, we have the strong property that any two translations of the same term only differ in places where transports of proof of equality have been injected. To keep track of this property, we introduce the relation $t \sim t'$ between two terms of ITT, of possibly different types. The crux of the proof of the translation is to guarantee that for every two terms t_1 and t_2 such that $\Gamma \vdash t_1 : T_1$, $\Gamma \vdash t_2 : T_2$ and $t_1 \sim t_2$, there exists p such that $\Gamma \vdash p : t_1 \text{ }_{T_1} \cong_{T_2} t_2$. However, during the proof, variables of different but (propositionally) equal types are introduced and the context cannot be maintained to be the same for both t_1 and t_2 . Therefore, the translation needs to keep track of this duplication of variables, plus a proof that they are heterogeneously equal. This mechanism is similar to what happens in the (relational) internal parametricity translation in ITT introduced by Bernardy et al. [2012] and recently rephrased in the setting of TemplateCoq [Anand et al. 2018]. Namely, a context is not translated as a telescope of variables, but as a telescope of triples consisting of two variables plus a witness that they are in the parametric relation. In our setting, this amounts to consider telescope of triples consisting of two variables plus a witness that they are heterogeneously equal. We can express this by considering the following dependent sums:

$$\text{Pack } A_1 A_2 := \Sigma(x : A_1). \Sigma(y : A_2). x \text{ }_{A_1} \cong_{A_2} y.$$

This presentation inspired by the parametricity translation is crucial in order to get an effective translation, because it is necessary to keep track of the evolution of contexts when doing the translation on open terms. This ingredient is missing in Oury's work [Oury 2005], which prevents him from deducing an effective (*i.e.*, constructive and computable) translation from his theorem.

2 Definitions of Extensional and Intensional Type Theories

This section presents the common syntax, typing and main properties of ETT and ITT. Our type theories feature a universe hierarchy, dependent products and sums as well as Martin L of's identity types.

2.1 Syntax of ETT and ITT

The common syntax of ETT and ITT is given in Figure 1. It features: dependent products $\Pi(x : A). B$, with (annotated) λ -abstractions and (annotated) applications, negative dependent sums $\Sigma(x : A). B$ with (annotated) projections, sorts \square_i , identity types $u =_A v$ with reflection and elimination as well as terms realising UIP and functional extensionality. Annotating terms with otherwise computationally irrelevant typing information is a common practice when studying the syntax of type theory precisely (see [Streicher 1993] for a similar example). We will write $A \rightarrow B$ for $\Pi(_ : A). B$ the non-dependent product / function type.

We consider a fixed universe hierarchy without cumulativity, which ensures in particular uniqueness of typing (2.2) which is important for the translation.

About Annotations. Although it may look like a technical detail, the use of annotation is more fundamental in ETT than it is in ITT (where it is irrelevant and doesn't affect the theory). And this is actually one of the main differences between our work (and that of Martin Hofmann [1995] who has a similar presentation) and the work of Oury [2005].

Indeed, by using the standard model where types are interpreted as cardinals rather than sets, it is possible to see that the equality $\text{nat} \rightarrow \text{nat} = \text{nat} \rightarrow \text{bool}$ is independent from the theory, it is thus possible to assume it (as an axiom, or for those that would still not be convinced, simply under a λ that would introduce this equality). In that context, the identity map $\lambda(x : \text{nat}). x$ can be given the type $\text{nat} \rightarrow \text{bool}$ and we thus type $(\lambda(x : \text{nat}). x) 0 : \text{bool}$. Moreover, the β -reduction of the non-annotated system used by Oury concludes that this expression reduces to 0, but cannot be given the type bool (as we said, the equality $\text{nat} \rightarrow \text{nat} = \text{nat} \rightarrow \text{bool}$ is independent from the theory, so the context is consistent). This means we lack subject reduction in this case (or uniqueness of types, depending on how we see the issue). Our presentation has a blocked β -reduction limited to matching annotations: $(\lambda(x : A). B. t) @_{x:A. B} u = t[x \leftarrow u]$, from which subject reduction and uniqueness of types follow.

Although subtle, this difference is responsible for Oury's need for an extra axiom. Indeed, to treat the case of equality of applications in his proof, he needs to assume the congruence rule for heterogeneous equality of applications, which is not provable when formulated with John Major equality (Fig. 2). Thanks to annotations and our notion of heterogeneous equality, we can prove this congruence rule for applications.

331	s	$::= \square_i \ (i \in \mathbb{N})$	sorts (universes)	386
332	T, A, B, t, u, v	$::= x \mid \lambda(x : A).B.t \mid t @_{x:A.B} u \mid \Pi(x : A). B \mid s$	dependent λ -calculus	387
333		$\mid \langle u; v \rangle_{x:A.B} \mid \pi_1^{x:A.B} p \mid \pi_2^{x:A.B} p \mid \Sigma(x : A). B$	dependent pairs	388
334		$\mid \text{refl}_A u \mid J(A, u, x.e.P, w, v, p) \mid u =_A v$	propositional equality	389
335		$\mid \text{funext}(x : A, B, f, g, e) \mid \text{uip}(A, u, v, p, q)$	equality axioms	390
336	Γ, Δ	$::= \bullet \mid \Gamma, x : A$	contexts	391
337				392

Figure 1. Common syntax of ETT and ITT

$$\text{JMAPP} \quad \frac{f_1 \ \forall(x:U_1).V_1 \cong_{\forall(x:U_2).V_2} f_2 \quad u_1 \ U_1 \cong_{U_2} u_2}{f_1 \ u_1 \ V_1[x \leftarrow u_1] \cong_{V_2[x \leftarrow u_2]} f_2 \ u_2}$$

Figure 2. Congruence of heterogeneous equality

2.2 The Typing Systems

As usual in dependent type theory, we consider contexts which are telescopes whose declarations may depend on any variable already introduced. We note $\Gamma \vdash t : A$ to say that t has type A in context Γ . $\Gamma \vdash A$ shall stand for $\Gamma \vdash A : s$ for some sort s and similarly $\Gamma \vdash A \equiv B$ stands for $\Gamma \vdash A \equiv B : s$.

We use two relations $(s, s') \in \text{Ax}$ (written (s, s') for short) and $(s, s', s'') \in \text{R}$ (written (s, s', s'')) to constrain the sorts in the typing rules for universes, dependent products and dependent sums, as is done in any Pure Type System (PTS). In our case, because we do not have cumulativity, the rules are as follows:

$$(\square_i, \square_{i+1}) \in \text{Ax} \quad (\square_i, \square_j, \square_{\max(i,j)}) \in \text{R}$$

We give the typing rules of ITT in Figure 3. The rules are standard and we do not explain them. Let us just point out the conversion rule, which says that $u : A$ can be given the type $u : B$ when $A \equiv B$, i.e., when A and B are convertible. As the notion of conversion is central in our work—the conversion of ETT being translated to an equality in ITT—we provide an exhaustive definition of it, with computational conversion rules (including β -conversion or reduction of the elimination principle of equality over reflexivity, see Figure 4), however congruence conversion rules can be found in Appendix A (Figure 6). Note that we use Christine Paulin-Möhrling’s variant of the J rule rather than Martin-Löf’s original formulation. Although pretty straightforward, being precise here is very important, as for instance the congruence rule for λ -terms is the reason why functional extensionality is derivable in ETT. Congruence of equality terms is a standard extension of congruence to the new principles we add (UIP and functional extensionality).

ETT is thus simply an extension of ITT (we write \vdash_x for the associated typing judgment) with the reflection rule on equality, which axiomatises that propositionally equal terms are convertible (see Equation 1). Note that, as already mentioned, in the presence of reflection and J, UIP is derivable

so we could remove it from ETT, but keeping it allows us to share a common syntax which makes the statements of theorems simpler and does not affect the development.

2.3 General Properties of ITT and ETT

We now state the main properties of both ITT and ETT. We do not detail their proof as they are standard and can be found in the Coq formalisation.

First, although not explicit in the typing system, weakening is admissible in ETT and ITT.

Lemma 2.1 (Weakening). *If $\Gamma \vdash \mathcal{J}$ and Δ extends Γ (possibly interleaving variables) then $\Delta \vdash \mathcal{J}$.*

Then, as mentioned above, the use of a non-cumulative hierarchy allows us to prove that a term t can be given at most one type in a context Γ , up-to conversion.

Lemma 2.2 (Uniqueness of typing). *If $\Gamma \vdash u : T_1$ and $\Gamma \vdash u : T_2$ then $\Gamma \vdash T_1 \equiv T_2$.*

Finally, an important property of the typing system (seen as a mutual inductive definition) is the possibility to deduce hypotheses from their conclusion, thanks to inversion of typing. Note that it is important here that our syntax is annotated for applications and projections as it provides a richer inversion principle.

Lemma 2.3 (Inversion of typing).

1. *If $\Gamma \vdash x : T$ then $(x : A) \in \Gamma$ and $\Gamma \vdash A \equiv T$.*
2. *If $\Gamma \vdash \square_i : T$ then $\Gamma \vdash \square_{i+1} \equiv T$.*
3. *If $\Gamma \vdash \Pi(x : A). B : T$ then $\Gamma \vdash A : s$ and $\Gamma, x : A \vdash B : s'$ and $\Gamma \vdash s'' \equiv T$ for some (s, s', s'') .*
4. *If $\Gamma \vdash \lambda(x : A).B.t : T$ then $\Gamma \vdash A : s$ and $\Gamma, x : A \vdash B : s'$ and $\Gamma, x : A \vdash t : B$ and $\Gamma \vdash \Pi(x : A). B \equiv T$.*
5. *If $\Gamma \vdash u @_{x:A.B} v : T$ then $\Gamma \vdash A : s$ and $\Gamma, x : A \vdash B : s'$ and $\Gamma \vdash u : \Pi(x : A). B$ and $\Gamma \vdash v : A$ and $\Gamma \vdash B[x \leftarrow u] \equiv T$.*
6. *... Analogous for the remaining term and type constructors.*

Proof. Each case is proven by induction on the derivation (which corresponds to any number of applications of the conversion rule following one introduction rule). \square

3 Relating Translated Expressions

We want to define a relation on terms that equates two terms that are the same up to transport. This begs the question of

Well-formedness of contexts.

$$\frac{}{\vdash \bullet} \quad \frac{\vdash \Gamma \quad \Gamma \vdash A}{\vdash \Gamma, x : A} (x \notin \Gamma)$$

Types.

$$\frac{\vdash \Gamma}{\Gamma \vdash s : s'} (s, s') \quad \frac{\Gamma \vdash A : s \quad \Gamma, x : A \vdash B : s'}{\Gamma \vdash \Pi(x : A). B : s''} (s, s', s'')$$

$$\frac{\Gamma \vdash A : s \quad \Gamma, x : A \vdash B : s'}{\Gamma \vdash \Sigma(x : A). B : s''} (s, s', s'')$$

$$\frac{\Gamma \vdash A : s \quad \Gamma \vdash u : A \quad \Gamma \vdash v : A}{\Gamma \vdash u =_A v : s}$$

Structural rules.

$$\frac{\vdash \Gamma \quad (x : A) \in \Gamma}{\Gamma \vdash x : A} \quad \frac{\Gamma \vdash u : A \quad \Gamma \vdash A \equiv B : s}{\Gamma \vdash u : B}$$

 λ -calculus terms.

$$\frac{\Gamma \vdash A : s \quad \Gamma, x : A \vdash B : s' \quad \Gamma, x : A \vdash t : B}{\Gamma \vdash \lambda(x : A). B.t : \Pi(x : A). B}$$

$$\frac{\Gamma \vdash A : s \quad \Gamma, x : A \vdash B : s' \quad \Gamma \vdash t : \Pi(x : A). B \quad \Gamma \vdash u : A}{\Gamma \vdash t @_{x:A.B} u : B[x \leftarrow u]}$$

$$\frac{\Gamma \vdash A : s \quad \Gamma, x : A \vdash B : s' \quad \Gamma \vdash v : B[x \leftarrow u]}{\Gamma \vdash \langle u; v \rangle_{x:A.B} : \Sigma(x : A). B}$$

$$\frac{\Gamma \vdash p : \Sigma(x : A). B}{\Gamma \vdash \pi_1^{x:A.B} p : A} \quad \frac{\Gamma \vdash p : \Sigma(x : A). B}{\Gamma \vdash \pi_2^{x:A.B} p : B[x \leftarrow \pi_1^{x:A.B} p]}$$

Equality terms.

$$\frac{\Gamma \vdash A : s \quad \Gamma \vdash u : A}{\Gamma \vdash \text{refl}_A u : u =_A u} \quad \frac{\Gamma \vdash e_1, e_2 : u =_A v}{\Gamma \vdash \text{uip}(A, u, v, e_1, e_2) : e_1 = e_2}$$

$$\frac{\Gamma \vdash A : s \quad \Gamma \vdash u, v : A \quad \Gamma, x : A, e : u =_A x \vdash P : s' \quad \Gamma \vdash p : u =_A v \quad \Gamma \vdash w : P[x \leftarrow u, e \leftarrow \text{refl}_A u]}{\Gamma \vdash J(A, u, x.e.P, w, v, p) : P[x \leftarrow v, e \leftarrow p]}$$

$$\frac{\Gamma \vdash f, g : \Pi(x : A). B \quad \Gamma \vdash e : \Pi(x : A). f @_{x:A.B} x =_B g @_{x:A.B} x}{\Gamma \vdash \text{funext}(x : A, B, f, g, e) : f = g}$$

Figure 3. Typing rules

what notion of transport is going to be used. Transport can be defined from elimination of equality as follows:

Definition 3.1 (Transport). Given $\Gamma \vdash p : T_1 =_s T_2$ and $\Gamma \vdash t : T_1$ we define the transport of t along p , written $p_* t$, as $J(s, T_1, X.e. T_1 \rightarrow X, \lambda(x : T_1). T_1.x, T_2, p) @_{T_1.T_2} t$ such that $\Gamma \vdash p_* t : T_2$.

However, in order not to confuse the transports added by the translation with the transports that were already present in the source, we consider p_* as part of the syntax in the reasoning. It will be unfolded to its definition only after the complete translation is performed. This idea is not novel as Hofmann already had a Subst operator that was part of his ITT (noted TT_1 in his paper [Hofmann 1995]).

We first define the (purely syntactic) relation \sqsubset between ETT terms and ITT terms in Figure 5 stating that the ITT term is simply a decoration of the first term by transports. Its purpose is to state how close to the original term its translation is. Then, we extend this relation to a similarity relation \sim on ETT terms by taking its symmetric and transitive closure:

$$\sim := (\sqsubset \cup \sqsubset^{-1})^+$$

Lemma 3.2 (\sim is an equivalence relation). \sim is reflexive, symmetric and transitive.

Proof. For reflexivity we proceed by induction on the term. \square

The goal is to prove that two terms in this relation, that are well-typed in the target type theory, are heterogeneously equal. As for this notion, we recall the definition we previously gave: $t \vDash_U u := \Sigma(p : T = U). p_* t = u$. This definition of heterogeneous equality can be shown to be reflexive, symmetric and transitive. Because of UIP, heterogeneous equality collapses to equality when taken on the same type.

Lemma 3.3. If $\Gamma \vdash e : u \vDash_A v$ then there exists p such that $\Gamma \vdash p : u =_A v$.

Proof. This holds thanks to UIP on equality, which implies K, and so the proof of $A = A$ can be taken to be reflexivity. \square

Note. In particular, \cong on types corresponds to equality. This is not as trivial as it sounds, one might be concerned about what happens if we have $\Gamma \vdash e : A \vDash_{s'} B$ with two distinct sorts s and s' . We would thus have $s = s'$, however, for this to be well-typed, we need to give a common type to s and s' , which can only be achieved if s and s' are actually the same sort.

Before we can prove the fundamental lemma stating that two terms in relation are heterogeneously equal, we need to consider another construction. As explained in the introduction, when proving the property by induction on terms, we introduce variables in the context that are equal only up-to heterogeneous equality. This phenomenon is similar to what happens in the parametricity translation [Bernardy et al. 2012]. Our fundamental lemma on the decoration relation \sim assumes two related terms of potentially different

Computation.

$$\frac{\Gamma \vdash A : s \quad \Gamma, x : A \vdash B : s' \quad \Gamma, x : A \vdash t : B \quad \Gamma \vdash u : A}{\Gamma \vdash (\lambda(x : A).B.t) @_{x:A.B} u \equiv t[x \leftarrow u] : B[x \leftarrow u]}$$

$$\frac{\Gamma \vdash A : s \quad \Gamma \vdash u : A \quad \Gamma, x : A, e : u =_A x \vdash P : s' \quad \Gamma \vdash w : P[x \leftarrow u, e \leftarrow \text{refl}_A u]}{\Gamma \vdash J(A, u, x.e.P, w, u, \text{refl}_A u) \equiv w : P[x \leftarrow u, e \leftarrow \text{refl}_A u]}$$

$$\frac{\Gamma \vdash A : s \quad \Gamma \vdash u : A \quad \Gamma, x : A \vdash B : s' \quad \Gamma \vdash v : B[x \leftarrow u]}{\Gamma \vdash \pi_1^{x:A.B} \langle u; v \rangle_{x:A.B} \equiv u : A}$$

$$\frac{\Gamma \vdash A : s \quad \Gamma \vdash u : A \quad \Gamma, x : A \vdash B : s' \quad \Gamma \vdash v : B[x \leftarrow u]}{\Gamma \vdash \pi_2^{x:A.B} \langle u; v \rangle_{x:A.B} \equiv v : B[x \leftarrow u]}$$

Conversion.

$$\frac{\Gamma \vdash t_1 \equiv t_2 : T_1 \quad \Gamma \vdash T_1 \equiv T_2}{\Gamma \vdash t_1 \equiv t_2 : T_2}$$

Figure 4. Main conversion rules (omitting congruence rules)

types T_1 and T_2 to produce an heterogeneous equality between them. For induction to go through under binders (e.g. for dependent products and abstractions), we hence need to consider the two terms under different, but heterogeneously equal contexts. Therefore, the context we produce will not only be a telescope of variables, but rather a telescope of triples consisting of two variables of possibly different types, and a witness that they are heterogeneously equal. To make this precise, we define the following macro:

$$\text{Pack } A_1 A_2 := \Sigma(x : A_1). \Sigma(y : A_2). x \cong y$$

together with its projections

$$\text{Proj}_1 p := \pi_1 p \quad \text{Proj}_2 p := \pi_2 p \quad \text{Proj}_e p := \pi_2 \pi_2 p.$$

We can then extend this notion canonically to contexts of the same length that are well formed using the same sorts:

$$\text{Pack } (\Gamma_1, x : A_1) (\Gamma_2, x : A_2) := (\text{Pack } \Gamma_1 \Gamma_2), x : \text{Pack } (A_1[\gamma_1]) (A_2[\gamma_2])$$

$$\text{Pack } \bullet \bullet := \bullet.$$

When we pack contexts, we also need to apply the correct projections for the types in that context to still make sense. Assuming two contexts Γ_1 and Γ_2 of the same length, we can define left and right substitutions:

$$\gamma_1 := [x \leftarrow \text{Proj}_1 x \mid (x : _) \in \Gamma_1]$$

$$\gamma_2 := [x \leftarrow \text{Proj}_2 x \mid (x : _) \in \Gamma_2].$$

These substitutions implement lifting of terms to packed contexts: $\Gamma, \text{Pack } \Gamma_1 \Gamma_2 \vdash t[\gamma_1] : A[\gamma_1]$ whenever $\Gamma, \Gamma_1 \vdash t : A$ (resp. $\Gamma, \text{Pack } \Gamma_1 \Gamma_2 \vdash t[\gamma_2] : A[\gamma_2]$ whenever $\Gamma, \Gamma_2 \vdash t : A$).

For readability, when Γ_1 and Γ_2 are understood we will write Γ_p for $\text{Pack } \Gamma_1 \Gamma_2$.

Implicitly, whenever we use the notation $\text{Pack } \Gamma_1 \Gamma_2$ it means that the two contexts are of the same length and well-formed with the same sorts. We can now state the fundamental lemma.

Lemma 3.4 (Fundamental lemma). *Let t_1 and t_2 be two terms. If $\Gamma, \Gamma_1 \vdash t_1 : T_1$ and $\Gamma, \Gamma_2 \vdash t_2 : T_2$ and $t_1 \sim t_2$ then there exists p such that $\Gamma, \text{Pack } \Gamma_1 \Gamma_2 \vdash p : t_1[\gamma_1]_{T_1[\gamma_1]} \cong_{T_2[\gamma_2]} t_2[\gamma_2]$.*

Proof. The proof is by induction on the derivation of $t_1 \sim t_2$. We show the three most interesting cases:

- VAR

$$\frac{}{x \sim x}$$

If x belongs to Γ , we apply reflexivity—together with uniqueness of typing (2.2)—to conclude. Otherwise, $\text{Proj}_e x$ has the expected type (since $x[\gamma_1] \equiv \text{Proj}_1 x$ and $x[\gamma_2] \equiv \text{Proj}_2 x$).

- APPLICATION

$$\frac{t_1 \sim t_2 \quad A_1 \sim A_2 \quad B_1 \sim B_2 \quad u_1 \sim u_2}{t_1 @_{x:A_1.B_1} u_1 \sim t_2 @_{x:A_2.B_2} u_2}$$

We have $\Gamma, \Gamma_1 \vdash t_1 @_{x:A_1.B_1} u_1 : T_1$ and $\Gamma, \Gamma_2 \vdash t_2 @_{x:A_2.B_2} u_2 : T_2$ which means by inversion (2.3) that the subterms are well-typed. We apply the induction hypothesis and then conclude.

- TRANSPORTLEFT

$$\frac{t_1 \sim t_2}{p_* t_1 \sim t_2}$$

We have $\Gamma, \Gamma_1 \vdash p_* t_1 : T_1$ and $\Gamma, \Gamma_2 \vdash t_2 : T_2$. By inversion (2.3) we have $\Gamma, \Gamma_1 \vdash p : T'_1 = T_1$ and $\Gamma, \Gamma_1 \vdash t_1 : T'_1$. By induction hypothesis we have e such that $\Gamma, \Gamma_p \vdash e :$

$$\begin{array}{c}
\frac{t_1 \sqsubset t_2}{t_1 \sqsubset p_* t_2} \\
\frac{}{x \sqsubset x} \quad \frac{A_1 \sqsubset A_2 \quad B_1 \sqsubset B_2}{\Pi(x : A_1). B_1 \sqsubset \Pi(x : A_2). B_2} \\
\frac{A_1 \sqsubset A_2 \quad B_1 \sqsubset B_2}{\Sigma(x : A_1). B_1 \sqsubset \Sigma(x : A_2). B_2} \\
\frac{A_1 \sqsubset A_2 \quad u_1 \sqsubset u_2 \quad v_1 \sqsubset v_2}{u_1 =_{A_1} v_1 \sqsubset u_2 =_{A_2} v_2} \quad \frac{}{s \sqsubset s} \\
\frac{A_1 \sqsubset A_2 \quad B_1 \sqsubset B_2 \quad t_1 \sqsubset t_2}{\lambda(x : A_1). B_1. t_1 \sqsubset \lambda(x : A_2). B_2. t_2} \\
\frac{t_1 \sqsubset t_2 \quad A_1 \sqsubset A_2 \quad B_1 \sqsubset B_2 \quad u_1 \sqsubset u_2}{t_1 @_{x:A_1.B_1} u_1 \sqsubset t_2 @_{x:A_2.B_2} u_2} \\
\frac{A_1 \sqsubset A_2 \quad B_1 \sqsubset B_2 \quad t_1 \sqsubset t_2 \quad u_1 \sqsubset u_2}{\langle t_1; u_1 \rangle_{x:A_1.B_1} \sqsubset \langle t_2; u_2 \rangle_{x:A_2.B_2}} \\
\frac{A_1 \sqsubset A_2 \quad B_1 \sqsubset B_2 \quad p_1 \sqsubset p_2}{\pi_1^{x:A_1.B_1} p_1 \sqsubset \pi_1^{x:A_2.B_1} p_2} \\
\frac{A_1 \sqsubset A_2 \quad B_1 \sqsubset B_2 \quad p_1 \sqsubset p_2 \quad A_1 \sqsubset A_2 \quad u_1 \sqsubset u_2}{\pi_2^{x:A_1.B_1} p_1 \sqsubset \pi_2^{x:A_2.B_2} p_2 \quad \text{refl}_{A_1} u_1 \sqsubset \text{refl}_{A_2} u_2} \\
\frac{A_1 \sqsubset A_2 \quad B_1 \sqsubset B_2 \quad f_1 \sqsubset f_2 \quad g_1 \sqsubset g_2 \quad e_1 \sqsubset e_2}{\text{funext}(x : A_1, B_1, f_1, g_1, e_1) \sqsubset \text{funext}(x : A_2, B_2, f_2, g_2, e_2)} \\
\frac{A_1 \sqsubset A_2 \quad u_1 \sqsubset u_2 \quad v_1 \sqsubset v_2 \quad p_1 \sqsubset p_2 \quad q_1 \sqsubset q_2}{\text{uip}(A_1, u_1, v_1, p_1, q_1) \sqsubset \text{uip}(A_2, u_2, v_2, p_2, q_2)} \\
\frac{u_1 \sqsubset u_2 \quad P_1 \sqsubset P_2 \quad w_1 \sqsubset w_2 \quad v_1 \sqsubset v_2 \quad p_1 \sqsubset p_2}{J(A_1, u_1, x.e.P_1, w_1, v_1, p_1) \sqsubset J(A_2, u_2, x.e.P_2, w_2, v_2, p_2)}
\end{array}$$

Figure 5. Relation \sqsubset

$t_1[\gamma_1] \cong t_2[\gamma_2]$. From transitivity and symmetry we only need to provide a proof of $t_1[\gamma_1] \cong p[\gamma_1]_* t_1[\gamma_1]$ which is inhabited by $\langle p[\gamma_1]; \text{refl}(p[\gamma_1]_* t_1[\gamma_1]) \rangle_{_}$.

The complete proof can be found in Appendix B. \square

We can also prove that \sim preserves substitution.

Lemma 3.5. *If $t_1 \sim t_2$ and $u_1 \sim u_2$ then $t_1[x \leftarrow u_1] \sim t_2[x \leftarrow u_2]$.*

Proof. We proceed by induction on the derivation of $t_1 \sim t_2$. \square

4 Translating ETT to ITT

4.1 The Translation

We now define the translations (let us stress the plural here) of an extensional judgment. We extend \sqsubset canonically to contexts ($\Gamma \sqsubset \bar{\Gamma}$ when they bind the same variables and the types are in relation for \sqsubset).

Before defining the translation, we define a set $\llbracket \Gamma \vdash_x t : A \rrbracket$ of typing judgments in ITT associated to a typing judgment $\Gamma \vdash_x t : A$ in ETT. The idea is that this set describes all the possible translations that lead to the expected property. When $\bar{\Gamma} \vdash \bar{t} : \bar{A} \in \llbracket \Gamma \vdash_x t : A \rrbracket$, we say that $\bar{\Gamma} \vdash \bar{t} : \bar{A}$ realises $\Gamma \vdash_x t : A$. The translation will be given by showing that this set is inhabited by induction on the derivation.

Definition 4.1 (Characterisation of possible translations).

- For any $\vdash_x \Gamma$ we define $\llbracket \vdash_x \Gamma \rrbracket$ as a set of valid judgments (in ITT) such that $\vdash \bar{\Gamma} \in \llbracket \vdash_x \Gamma \rrbracket$ if and only if $\Gamma \sqsubset \bar{\Gamma}$.
- Similarly, $\bar{\Gamma} \vdash \bar{t} : \bar{A} \in \llbracket \Gamma \vdash_x t : A \rrbracket$ iff $\bar{\Gamma} \in \llbracket \vdash_x \Gamma \rrbracket$ and $A \sqsubset \bar{A}$ and $t \sqsubset \bar{t}$.

In order to better master the shape of the produced realiser, we state the following lemma which shows that it has the same head type constructor as the type it realises. This is important for instance for the case of an application, where we do not know a priori if the translated function has a dependent product type, which is required to be able to use the typing rule for application.

Lemma 4.2. *We can always choose types \bar{T} that have the same head constructor as T .*

Proof. Assume we have $\bar{\Gamma} \vdash \bar{t} : \bar{T} \in \llbracket \Gamma \vdash_x t : T \rrbracket$. By definition of \sqsubset , $T \sqsubset \bar{T}$ means that \bar{T} is shaped $p_* q_* \dots r_* \bar{T}'$ with \bar{T}' having the same head constructor as T . By inversion (2.3), the subterms are typable, including \bar{T}' . Actually, from inversion, we even get that the type of \bar{T}' is a universe. Then, using lemma 3.4 and lemma 3.3, we get $\bar{\Gamma} \vdash e : \bar{T} = \bar{T}'$. We conclude with $\bar{\Gamma} \vdash e_* \bar{t} : \bar{T}' \in \llbracket \Gamma \vdash_x t : T \rrbracket$. \square

Finally, in order for the induction to go through, we need to know that when we have a realiser of a derivation $\Gamma \vdash_x t : T$, we can pick an arbitrary other type realising $\Gamma \vdash_x T$ and still get a new derivation realising $\Gamma \vdash_x t : T$ with that type. This is important for instance for the case of an application, where the type of the domain of the translated function may differ from the type of the translated argument. So we need to be able to change it *a posteriori*.

Lemma 4.3. *When we have $\bar{\Gamma} \vdash \bar{t} : \bar{T} \in \llbracket \Gamma \vdash_x t : T \rrbracket$ and $\bar{\Gamma} \vdash \bar{t}' : \bar{T}' \in \llbracket \Gamma \vdash_x T \rrbracket$ then we also have $\bar{\Gamma} \vdash \bar{t}' : \bar{T}' \in \llbracket \Gamma \vdash_x t : T \rrbracket$ for some \bar{t}' .*

Proof. By definition we have $T \sqsubset \bar{T}$ and $T \sqsubset \bar{T}'$ and thus $T \sim \bar{T}$ and $T \sim \bar{T}'$, implying $\bar{T} \sim \bar{T}'$ by transitivity (3.2). By lemma 3.4 (in the case $\Gamma_1 \equiv \Gamma_2 \equiv \bullet$) we get $\bar{\Gamma} \vdash p : \bar{T} \cong \bar{T}'$ for some p . By lemma 3.3 (and lemma 4.2 to give universes as types to \bar{T} and \bar{T}') we can assume $\bar{\Gamma} \vdash p : \bar{T} = \bar{T}'$. Then $\bar{\Gamma} \vdash p_* \bar{t} : \bar{T}'$ is still a translation since \sqsubset ignores transports. \square

We can now define the translation. This is done by mutual induction on context well-formedness, typing and conversion derivations. Indeed, in order to be able to produce a realiser by induction, we need to show that every conversion in ETT is translated as an heterogeneous equality in ITT.

Theorem 4.4 (Translation).

- If $\vdash_x \Gamma$ then there exists $\vdash \bar{\Gamma} \in \llbracket \vdash_x \Gamma \rrbracket$,
- If $\Gamma \vdash_x t : T$ then for any $\vdash \bar{\Gamma} \in \llbracket \vdash_x \Gamma \rrbracket$ there exist \bar{t} and \bar{T} such that $\bar{\Gamma} \vdash \bar{t} : \bar{T} \in \llbracket \Gamma \vdash_x t : T \rrbracket$,
- If $\Gamma \vdash_x u \equiv v : A$ then for any $\vdash \bar{\Gamma} \in \llbracket \vdash_x \Gamma \rrbracket$ there exist $A \sqsubset \bar{A}, A \sqsubset \bar{A}', u \sqsubset \bar{u}, v \sqsubset \bar{v}$ and \bar{e} such that $\bar{\Gamma} \vdash \bar{e} : \bar{u} \bar{A} \cong_{\bar{A}} \bar{v}$.

Proof. We prove the theorem by induction on the derivation in the extensional type theory. We only show the two most interesting cases of application and conversion. The complete proof is given in Appendix C.

- APPLICATION

$$\frac{\Gamma \vdash_x A : s \quad \Gamma, x : A \vdash_x B : s' \quad \Gamma \vdash_x t : \Pi(x : A). B \quad \Gamma \vdash_x u : A}{\Gamma \vdash_x t @_{x:A.B} u : B[x \leftarrow u]}$$

Using IH together with lemmata 4.2 and 4.3 we get $\bar{\Gamma} \vdash \bar{A} : s$ and $\bar{\Gamma}, x : \bar{A} \vdash \bar{B} : s'$ and $\bar{\Gamma} \vdash \bar{t} : \Pi(x : \bar{A}). \bar{B}$ and $\bar{\Gamma} \vdash \bar{u} : \bar{A}$ meaning we can conclude $\bar{\Gamma} \vdash \bar{t} @_{x:\bar{A}. \bar{B}} \bar{u} : \bar{B}[x \leftarrow \bar{u}] \in \llbracket \Gamma \vdash_x t @_{x:A.B} u : B[x \leftarrow u] \rrbracket$.

- CONVERSION

$$\frac{\Gamma \vdash_x u : A \quad \Gamma \vdash_x A \equiv B}{\Gamma \vdash_x u : B}$$

By IH and lemma 3.3 we have $\bar{\Gamma} \vdash \bar{e} : \bar{A} = \bar{B}$ which implies $\bar{\Gamma} \vdash \bar{A} \in \llbracket \Gamma \vdash_x A \rrbracket$ by inversion (2.3), thus, from lemma 4.3 and IH we get $\bar{\Gamma} \vdash \bar{u} : \bar{A}$, yielding $\bar{\Gamma} \vdash \bar{e}_* \bar{u} : \bar{B} \in \llbracket \Gamma \vdash_x u : B \rrbracket$. \square

4.2 Meta-theoretical Consequences

We can check that all ETT theorems whose type are typable in ITT have proofs in ITT as well:

Corollary 4.5 (Preservation of ITT). *If $\vdash_x t : T$ and $\vdash T$ then there exist \bar{t} such that $\vdash \bar{t} : T \in \llbracket \vdash_x t : T \rrbracket$.*

Proof. Since $\vdash \bullet \in \llbracket \vdash_x \bullet \rrbracket$, by Theorem (4.4), there exists \bar{t} and \bar{T} such that $\vdash \bar{t} : \bar{T} \in \llbracket \vdash_x t : T \rrbracket$. But as $\vdash T$, we have

$\vdash T \in \llbracket \vdash_x T \rrbracket$, and, using Lemma 4.3, we obtain $\vdash \bar{t} : T \in \llbracket \vdash_x t : T \rrbracket$. \square

Corollary 4.6 (Relative consistency). *Assuming ITT is consistent, there is no term t such that $\vdash_x t : \Pi(A : \square_0). A$.*

Proof. Assume such a t exists. By the Corollary 4.5, because $\vdash \Pi(A : \square_0). A$, there exists \bar{t} such that $\vdash \bar{t} : \Pi(A : \square_0). A$ which contradicts the assumed consistency of ITT. \square

4.3 Optimisations

Up until now, we remained silent about one thing: the size of the translated terms. Indeed, the translated term is a decoration of the initial one by transports which appear in many locations. For example, at each application we use a transport by lemma 4.2 to ensure that the term in function position is given a function type. In most cases—in particular when translating ITT terms—this produces unnecessary transports (often by reflexivity) that we wish to avoid.

In order to limit the size explosion, in the above we use a different version of transport, namely transport' such that

$$\text{transport}'_{A_1, A_2}(p, t) = t \quad \text{when } A_1 =_{\alpha} A_2 \\ = p_* t \quad \text{otherwise.}$$

The idea is that we avoid *trivially* unnecessary transports (we do not deal with β -conversion for instance). We extend this technique to the different constructors of equality (symmetry, transitivity, ...) so that they reduce to reflexivity whenever possible. Take transitivity for instance:

$$\text{transitivity}'(\text{refl } u, q) = q \\ \text{transitivity}'(p, \text{refl } u) = p \\ \text{transitivity}'(p, q) = \text{transitivity}(p, q).$$

We show these *defined terms* enjoy the same typing rules as their counterparts and use them instead. In practice it is enough to recover the exact same term when it is typed in ITT.

5 Formalisation with Template-Coq

We have formalised the translation in the setting of TemplateCoq [Anand et al. 2018] in order to have a more precise proof, but also to evidence the fact that the translation is indeed constructive and can be used to perform computations.

TemplateCoq is a Coq library that has a representation of Coq terms as they are in Coq's kernel (in particular using de Bruijn indices for variables) and a (partial) implementation of the type checking algorithm (not checking guardedness of fixpoints or positivity of inductive types). It comes with a Coq plugin that permits to quote Coq terms into their representations, and to produce Coq terms from their representation (if they indeed denote well-typed terms). We have integrated our formalisation within that framework in order to ensure our presentations of ETT and ITT are close to Coq, but also to take advantage of the quoting mechanism

to produce terms using the interactive mode (in particular we get to use tactics). Note that we also rely on Mangin and Sozeau's Equations [Sozeau 2010] plugin to derive nice dependent induction principles.

Our formalisation takes full advantage of its easy interfacing with TemplateCoq: we define two theories, namely ETT and ITT, but ITT enjoys a lot of syntactic sugar by having things such as transport, heterogeneous equality and packing as part of the syntax. The operations regarding these constructors—in particular the tedious ones—are written in Coq and then quoted to finally be *realised* in the translation from ITT to TemplateCoq.

Interoperability with TemplateCoq. The translation we define from ITT to TemplateCoq is not proven correct, but it is not really important as it can just be seen as a feature to observe the produced terms in a nicer setting. In any case, TemplateCoq does not yet provide a complete formalisation of CIC rules, as guard checking of recursive definitions and strict positivity of inductive type declarations are not formalised yet.

We also provide a translation from TemplateCoq to ETT that we will describe more extensively with the examples (5.4).

5.1 Quick Overview of the Formalisation

The file `SAST.v` contains the definition of the (common) abstract syntax of ETT and ITT in the form of an inductive definition with de Bruijn indices for variables (like in TemplateCoq). Sorts are defined separately in `Sorts.v` and we will address them later in Section 5.3.

```

Inductive sterm : Type :=
| sRel (n : nat)
| sSort (s : sort)
| sProd (nx : name) (A B : sterm)
| sLambda (nx : name) (A B t : sterm)
| sApp (u : sterm) (nx : name) (A B v : sterm)
| sEq (A u v : sterm)
| sRefl (A u : sterm)
| (* ... *) .

```

The files `ITyping.v` and `XTyping.v` define respectively the typing judgments for ITT and ETT, using mutual inductive types. Then, most of the files are focused on the meta-theory of ITT and can be ignored by readers who don't need to see yet another proof of subject reduction.

The most interesting files are obviously those where the fundamental lemma and the translation are formalised: `FundamentalLemma.v` and `Translation.v`. For instance, here is the main theorem, as stated in our formalisation:

```

Theorem complete_translation Σ :
  type_glob Σ ->
  (forall Γ (h : XTyping.wf Σ Γ), Σ Γ', Σ |--i Γ' # [Γ]) *
  (forall Γ t A (h : Σ ;; Γ |-x t : A)
    Γ' (hΓ : Σ |--i Γ' # [Γ]),
    Σ A' t', Σ ;; Γ' |--- [t'] : A' # [Γ |--- [t] : A]) *
  (forall Γ u v A (h : Σ ;; Γ |-x u = v : A)
    Γ' (hΓ : Σ |--i Γ' # [Γ]),
    Σ A' A'' u' v' p', eqtrans Σ Γ A u v Γ' A' A'' u' v' p').

```

Herein `type_glob Σ` refers to the fact that some global context is well-typed, its purpose is detailed in Section 5.2. The

fact that the theorem holds in Coq ensures we can actually compute a translated term and type out of a derivation in ETT.

5.2 Inductive Types and Recursion

In the proof of Section 4, we didn't mention anything about inductive types, pattern-matching or recursion as it is a bit technical on paper. In the formalisation, we offer a way to still be able to use them, and we will even show how it works in practice with the examples (5.4).

The main guiding principle is that inductive types and induction are orthogonal to the translation, they should more or less be translated to themselves. To realise that easily, we just treat an inductive definition as a way to introduce new constants in the theory, one for the type, one for each constructor, one for its elimination principle, and one equality per computation rule. For instance, the natural numbers can be represented by having the following constants in the context:

```

nat      : □0
0        : nat
S        : nat → nat
natrec   : ∀P, P 0 → (∀m, P m → P (S m)) → ∀n, P n
natrec0 : ∀P Pz Ps, natrec P Pz Ps 0 = Pz
natrecS  : ∀P Pz Ps n,
           natrec P Pz Ps (S n) = Ps n (natrec P Pz Ps n)

```

Here we rely on the reflection rule to obtain the computational behavior of the eliminator `natrec`.

This means for instance that we do not consider inductive types that would only make sense in ETT, but we deem this not to be a restriction and to the best of our knowledge isn't something that is usually considered in the literature. With that in mind, our translation features a global context of typed constants with the restriction that the types of those constants should be well-formed in ITT. Those constants are thus used as black boxes inside ETT.

With this we are able to recover what we were missing from Coq, without having to deal with the trouble of proving that the translation doesn't break the guard condition of fixed points, and we are instead relying on a more type-based approach.

5.3 About Universes and Homotopy

The experienced reader might have noticed that our treatment of universes (except perhaps for the absence of cumulativity) was really superficial and the notion of sorts used is rather orthogonal to our main development. This is even more apparent in the formalisation. Indeed, we didn't fix a specific universe hierarchy, but instead specify what properties it should have, in what is reminiscent to a (functional³) PTS formulation.

³Meaning the sort of a sort, and the sort of a product are functions, necessary to the uniqueness of types (2.2).

```

991 Class Sorts.notion := {
992   sort : Type ;
993   succ : sort -> sort ;
994   prod_sort : sort -> sort -> sort ;
995   sum_sort : sort -> sort -> sort ;
996   eq_sort : sort -> sort ;
997   eq_dec : forall s z : sort, {s = z} + {s <> z} ;
998   succ_inj : forall s z, succ s = succ z -> s = z
999 }.

```

From the notion of sorts, we require functions to get the sort of a sort, the sort of a product from the sorts of its arguments, and (crucially) the sort of an identity type. We also require some measure of decidable equality and injectivity on those.

This allows us to instantiate this by a lot of different notions including the one presented earlier in the paper or even its extension with a universe Prop of propositions (like CIC [Bertot and Castéran 2004]). We present here two instances that have their own interest.

Type in Type. One of the instances we provide is one with only one universe Type, with the inconsistent typing rule Type : Type. Although inconsistent, this allows us to interface with TemplateCoq, without the—for the time being—very time-consuming universe constraint checking.

Homotopy Type System and Two-Level Type Theory. Another interesting application (or rather instance) of our formalisation is a translation from Homotopy Type System (HTS) [Voevodsky 2013] to Two-Level Type Theory (2TT) [Altenkirch et al. 2016; Annenkov et al. 2017].

HTS and 2TT arise from the incompatibility between UIP—recall it is provable in ETT—and univalence. The idea is to have two distinct notions of equality in the theory, a *strict* one satisfying UIP, and a *fibrant* one corresponding to the homotopy type theory equality, possibly satisfying univalence. This actually induces a separation in the types of the theory: some of them are called *fibrant* and the fibrant or homotopic equality can only be eliminated on those. HTS can be seen as an extension of 2TT with reflection on the strict equality just like ETT is an extension of ITT.

We can recover HTS and 2TT in our setting by taking F_i and U_i as respectively the fibrant and strict universes of those theories (for $i \in \mathbb{N}$), along with the following PTS rules:

$$\begin{array}{ll}
(F_i, F_{i+1}) & \in Ax & (U_i, U_{i+1}) & \in Ax \\
(F_i, F_j, F_{\max(i,j)}) & \in R & (F_i, U_j, U_{\max(i,j)}) & \in R \\
(U_i, F_j, U_{\max(i,j)}) & \in R & (U_i, U_j, U_{\max(i,j)}) & \in R
\end{array}$$

and the fact that the sort of the (strict) identity type on $A : s$ is the *strictified* version of s , i.e., U_i for $s = U_i$ or $s = F_i$. In order to have the fibrant equality, one simply needs to do as in Section 5.2.

In short, the translation from HTS to 2TT is basically the same as the one from ETT to ITT we presented in this paper, and this fact is factorised through our formalisation.

5.4 ETT-flavoured Coq: Examples

In this section we demonstrate how our translation can bring extensionality to the world of Coq in action. The examples can be found in plugin_demo.v.

First, a pedestrian approach. We would like to begin by showing how one can write an example step by step before we show how it can be used in practice. For this we use a self-contained example without any inductive types or recursion, illustrating a very simple case of reflection. The term we want to translate is the following:

$$\lambda A B e x. x : \Pi A B. A = B \rightarrow A \rightarrow B$$

This is, in some sense the identity, relying on the equality $e : A = B$ to convert $x : A$ to $x : B$. Of course, this definition isn't accepted in Coq because the conversion doesn't hold in ITT.

Fail **Definition** pseudoid (A B : **Type**) (e : A = B) (x : A) : B := x.

However, we still want to be able to write it *in some way*, in order to avoid manipulating de Bruijn indices directly. For this, we use a little trick by first defining a Coq axiom to represent an ill-typed term:

Axiom candidate : forall A B (t : A), B.

candidate A B t is a candidate t of type A to inhabit type B. We complete this by adding a notation that is reminiscent to Agda's [Norell 2007] hole mechanism.

Notation "'{!} t !'" := (candidate _ _ t).

We can now write the ETT function within Coq.

Definition pseudoid (A B : **Type**) (e : A = B) (x : A) : B := {! x !}.

We can then quote the term and its type to TemplateCoq thanks to the Quote **Definition** command provided by the plugin.

Quote **Definition** pseudoid_term :=
ltac:(let t := eval compute in pseudoid in exact t).

Quote **Definition** pseudoid_type :=
ltac:(let T := type of pseudoid in exact T).

The terms that we get are now TemplateCoq terms, representing Coq syntax. We need to put them in ETT, meaning adding the annotations, and also removing the candidate axiom. This is the purpose of the fullquote function that we provide in our formalisation.

Definition pretm_pseudoid :=
Eval lazy in fullquote (2^18) Σ [] pseudoid_term empty empty nomap.

Definition tm_pseudoid :=
Eval lazy in match pretm_pseudoid with
| Success t => t
| Error _ => sRel 0
end.

Definition prety_pseudoid :=
Eval lazy in fullquote (2^18) Σ [] pseudoid_type empty empty nomap.

Definition ty_pseudoid :=
Eval lazy in match prety_pseudoid with
| Success t => t
| Error _ => sRel 0
end.

tm_pseudoid and ty_pseudoid correspond respectively to the ETT representation of pseudoid and its type. We then produce, using our home-brewed Ltac type-checking tactic,

the corresponding ETT typing derivation (notice the use of reflection to typecheck).

```

1103 Lemma type_pseudoid :  $\Sigma i$  ;; [] |-x tm_pseudoid : ty_pseudoid.
1104 Proof.
1105   unfold tm_pseudoid, ty_pseudoid.
1106   ettccheck. cbn.
1107   eapply reflection with (e := sRel 1).
1108   ettccheck.
1109 Defined.

```

We can then translate this derivation, obtain the translated term and then convert it to TemplateCoq.

```

1110 Definition itt_pseudoid : sterm :=
1111   Eval lazy in
1112   let '(_ ; t ; _) :=
1113     type_translation type_pseudoid istrans_nil
1114   in t.

```

```

1114 Definition tc_pseudoid : tsl_result term :=
1115   Eval lazy in
1116   tsl_rec (2 ^ 18)  $\Sigma$  [] itt_pseudoid empty.

```

Once we have it, we *unquote* the term to obtain a Coq term (notice that the only use of reflection has been replaced by a transport).

```

1120 fun (A B : Type) (e : A = B) (x : A) => transport e x
1121   : forall A B : Type, A = B -> A -> B

```

Making a Plugin with TemplateCoq. All of this work is pretty systematic. Fortunately for us, TemplateCoq also features a monad to reify Coq commands which we can use to *program* the translation steps. As such we have written a complete procedure, relying on Coq type checkers we wrote for ITT and ETT, which can generate equality obligations.

Thanks to this, the user doesn't have to know about the details of implementation of the translation, and stay within the Coq ecosystem.

For instance, our previous example now becomes:

```

1132 Definition pseudoid (A B : Type) (e : A = B) (x : A) : B := {! x !}.

```

```

1133 Run TemplateProgram (Translate  $\epsilon$  "pseudoid").

```

This produces a Coq term `pseudoid'` corresponding to the translation. Notice how the user doesn't even have to provide any proof of equality or derivations of any sort. The derivation part is handled by our own typechecker while the obligation part is solved automatically by the Coq obligation mechanism.

About inductive types. As we promised, our translation is able to handle inductive types. For this consider the inductive type of vectors (or length-indexed lists) below, together with a simple definition (we will remain in ITT for simplicity).

```

1145 Inductive vec A : nat -> Type :=
1146   | vnil : vec A 0
1147   | vcons : A -> forall n, vec A n -> vec A (S n).

```

```

1148 Arguments vnil {_}.
1149 Arguments vcons {_} _ _ .

```

```

1150 Definition vv := vcons 1 _ vnil.

```

This time, in order to apply the translation we need to extend the translation context with `nat` and `vec`.

```

1153 Run TemplateProgram (
1154    $\Theta$  <- TranslateConstant  $\epsilon$  "nat" ;;
1155    $\Theta$  <- TranslateConstant  $\Theta$  "vec" ;;

```

```

1156   Translate  $\Theta$  "vv"
1157 ).

```

Here, ϵ is the empty translation context and the command `TranslateConstant` enriches it with the types the inductives and of their constructors. The translation context then also contains associative tables between our own representation of constants and those of Coq. Unsurprisingly, the translated Coq term is the same as the original term.

Reversal of vectors. Next, we tackle a motivating example: reversal on vectors. Indeed, if you want to implement this operation, the same way you would do it on lists, you end up having a conversion problem:

```

1169 Fail Definition vrev {A n m} (v : vec A n) (acc : vec A m)
1170   : vec A (n + m) :=
1171   vec_rect A (fun n _ => forall m, vec A m -> vec A (n + m))
1172     (fun m acc => acc)
1173     (fun a n _ rv m acc => rv _ (vcons a m acc))
1174     n v m acc.

```

The recursive call returns a vector of length $n + S m$ where the context expects one of length $S n + m$. In ITT these types are not convertible. This example is thus a perfect fit for ETT where we can use the fact that these two expressions always compute to the same thing when instantiated with concrete numbers.

```

1179 Definition vrev {A n m} (v : vec A n) (acc : vec A m)
1180   : vec A (n + m) :=
1181   vec_rect A (fun n _ => forall m, vec A m -> vec A (n + m))
1182     (fun m acc => acc)
1183     (fun a n _ rv m acc => {! rv _ (vcons a m acc) !})
1184     n v m acc.

```

```

1184 Run TemplateProgram (
1185    $\Theta$  <- TranslateConstant  $\epsilon$  "nat" ;;
1186    $\Theta$  <- TranslateConstant  $\Theta$  "vec" ;;
1187    $\Theta$  <- TranslateConstant  $\Theta$  "Nat.add" ;;
1188    $\Theta$  <- TranslateConstant  $\Theta$  "vec_rect" ;;
1189   Translate  $\Theta$  "vrev"
1190 ).

```

This generates four obligations that are all solved automatically. One of them contains a proof of $S n + m = n + S m$ while the remaining three correspond to the computation rules of addition (as mentioned before, `add` is simply a constant and does not compute in our representation, hence the need for equalities). The returned term is the following, with only one transport remaining (remember our interpretation map removes unnecessary transports).

```

1197 fun (A : Type) (n m : nat) (v : vec A n) (acc : vec A m) =>
1198   vec_rect A
1199     (fun n _ => forall m, vec A m -> vec A (n + m))
1200     (fun m acc => acc)
1201     (fun a n0 v0 rv m0 acc0 =>
1202       transport (vrev_obligation_3 A n m v acc a n0 v0 rv m0 acc0)
1203         (rv (S m0) (vcons a m0 acc0))) n v m acc
1204   : forall A n m, vec A n -> vec A m -> vec A (n + m)

```

5.5 Towards an Interfacing between Andromeda and Coq

Andromeda [Bauer et al. 2016] is a proof assistant implementing ETT in a sense that is really close to our formalisation. Aside from a concise nucleus with a basic type theory, most things happen with the declaration of constants with given

types, including equalities to define the computational behaviour of eliminators for instance. This is essentially what we do in our formalisation. Furthermore, their theory relies on $\text{Type} : \text{Type}$, meaning, our modular handling of universes can accommodate for this as well.

All in all, it should be possible in the near future to use our translation to produce Coq terms out of Andromeda developments. Note that this would not suffer from the difficulties in generating typing derivations since Andromeda does it for you.

5.6 Composition with other Translations

This translation also enables the formalisation of translations that target ETT rather than ITT and still get mechanised proofs of (relative) consistency by composition with this ETT to ITT translation. This could also be used to implement plugins based on the composition of translations. In particular, supposing we have a theory which forms a subset of ETT and whose conversion is decidable. Using this translation, we could formalise it as an embedded domain-specific type theory and provide an automatic translation of well-typed terms into witnesses in Coq. This would make it possible to extend conversion with the theory of lists for example.

This would provide a simple way to justify the consistency of CoqMT [Jouannaud and Strub 2017] for example, seeing it as an extensional type theory where reflection is restricted to equalities on a specific domain whose theory is decidable.

6 Limitations and Axioms

Currently, the representation of terms and derivations and the computational content of the proof only allow us to deal with the translation of relatively small terms but we hope to improve that in the future. As we have seen, the actual translation involves the computational content of lemmata of inversion, substitution, weakening and equational reasoning and thus cannot be presented as a simple recursive definition on derivations.

As we already mentioned, the axioms K and functional extensionality are both necessary in ITT if we want the translation to be conservative as they are provable in ETT [Hofmann 1995]. However, one might still be concerned about having axioms as they can for instance hinder canonicity of the system. In that respect, K isn't really a restriction since it preserves canonicity. The best proof of that is probably Agda itself which natively features K—in fact, one needs to explicitly deactivate it with a flag if they wish to work without.

The case of functional extensionality is trickier. It is still possible to realise the axiom by composing our translation with a setoid interpretation [Altenkirch 1999] which validates it, or by going into a system featuring it, for instance by implementing Observational Type Theory [Altenkirch et al. 2007] like EPIGRAM [McBride 2004].

7 Related Works and Conclusion

The seminal works on the precise connection between ETT and ITT go back to Streicher [1993] and Hofmann [1995, 1997]. In particular, the work of Hofmann provides a categorical answer to the question of consistency and conservativity of ETT over ITT with UIP and functional extensionality. Ten years later, Oury [2005, 2006] provided a translation from ETT to ITT with UIP and functional extensionality and other axioms (mainly due to technical difficulties). Although a first step towards a move from categorical semantics to a syntactic translation, his work does not stress any constructive aspect of the proof and shows that there merely exist translations in ITT to a typed term in ETT.

van Doorn et al. [2013] have later proposed and formalised a similar translation between a PTS with and without explicit conversion. This does not entail anything about ETT to ITT but we can find similarities in that there is a witness of conversion between any term and itself under an explicit conversion, which internalises irrelevance of explicit conversions. This morally corresponds to a Uniqueness of Conversions principle.

The Program [Sozeau 2007] extension of Coq performs a related coercion insertion algorithm, between objects in subsets on the same carrier or in different instances of the same inductive family, assuming a proof-irrelevance axiom. Inserting coercions locally is not as general as the present translation from ETT to ITT which can insert transports in any context.

In this paper we provide the first effective translation from ETT to ITT with UIP and functional extensionality. The translation has been formalised in Coq using TemplateCoq, a meta-programming plugin of Coq. This translation is also effective in the sense that we can produce in the end a Coq term using the TemplateCoq denotation machinery. With ongoing work to extend the translation to the inductive fragment of Coq, we are paving the way to an extensional version of the Coq proof assistant which could be translated back to its intensional version, allowing the user to navigate between the two modes, and in the end produce a proof term checkable in the intensional fragment.

References

- Stuart F. Allen, Robert L. Constable, Richard Eaton, Christoph Kreitz, and Lori Lorigo. 2000. The Nuprl Open Logical Environment. In *Automated Deduction - CADE-17, 17th International Conference on Automated Deduction, Pittsburgh, PA, USA, June 17-20, 2000, Proceedings (Lecture Notes in Computer Science)*, David A. McAllester (Ed.), Vol. 1831. Springer, 170–176. https://doi.org/10.1007/10721959_12
- T. Altenkirch. 1999. Extensional equality in intensional type theory. In *Proceedings. 14th Symposium on Logic in Computer Science (Cat. No. PR00158)*. 412–420. <https://doi.org/10.1109/LICS.1999.782636>
- Thorsten Altenkirch, Paolo Capriotti, and Nicolai Kraus. 2016. Extending Homotopy Type Theory with Strict Equality. *CoRR* abs/1604.03799 (2016). arXiv:1604.03799 <http://arxiv.org/abs/1604.03799>

- 1321 Thorsten Altenkirch, Conor McBride, and Wouter Swierstra. 2007. Observational equality, now!. In *Proceedings of the 2007 workshop on Programming languages meets program verification*. ACM, 57–68. 1376
- 1322 Abhishek Anand, Simon Boulrier, Cyril Cohen, Matthieu Sozeau, and Nicolas Tabareau. 2018. Towards Certified Meta-Programming with Typed Template-Coq. In *Interactive Theorem Proving - 9th International Conference, ITP 2018, Held as Part of the Federated Logic Conference, FloC 2018, Oxford, UK, July 9-12, 2018, Proceedings (Lecture Notes in Computer Science)*, Jeremy Avigad and Assia Mahboubi (Eds.), Vol. 10895. Springer, 20–39. https://doi.org/10.1007/978-3-319-94821-8_2 1377
- 1323 Danil Annenkov, Paolo Capriotti, and Nicolai Kraus. 2017. Two-Level Type Theory and Applications. *CoRR* abs/1705.03307 (2017). arXiv:1705.03307 <http://arxiv.org/abs/1705.03307> 1378
- 1324 Andrej Bauer, Gaëtan Gilbert, Philipp G. Haselwarter, Matija Pretnar, and Chris Stone. 2016. The ‘Andromeda’ prover. (2016). <http://www.andromeda-prover.org/> 1379
- 1325 Jean-philippe Bernardy, Patrik Jansson, and Ross Paterson. 2012. Proofs for free: Parametricity for dependent types. *Journal of Functional Programming* 22, 2 (2012), 107–152. 1380
- 1326 Yves Bertot and Pierre Castéran. 2004. Interactive Theorem Proving and Program Development. (2004). 1381
- 1327 Marc Bezem, Thierry Coquand, and Simon Huber. 2013. A Model of Type Theory in Cubical Sets. (December 2013). <http://www.cse.chalmers.se/~coquand/mod1.pdf> 1382
- 1328 Simon Boulrier, Pierre-Marie Pédrot, and Nicolas Tabareau. 2017. The Next 700 Syntactical Models of Type Theory. In *Certified Programs and Proofs - CPP 2017*. 182–194. 1383
- 1329 The Coq development team. 2017. *The Coq proof assistant reference manual*. LogiCal Project. <http://coq.inria.fr> Version 8.7. 1384
- 1330 Martin Hofmann. 1995. Conservativity of equality reflection over intensional type theory. In *International Workshop on Types for Proofs and Programs*. Springer, 153–164. 1385
- 1331 Martin Hofmann. 1997. *Extensional constructs in intensional type theory*. Springer. 1386
- 1332 Martin Hofmann and Thomas Streicher. 1998. The Groupoid Interpretation of Type Theory. In *Twenty-five years of constructive type theory (Venice, 1995)*. Oxford Logic Guides, Vol. 36. Oxford Univ. Press, New York, 83–111. <http://www.tcs.informatik.uni-muenchen.de/lehre/SS97/types-vl/venedig.ps> 1387
- 1333 Jean-Pierre Jouannaud and Pierre-Yves Strub. 2017. Coq without Type Casts: A Complete Proof of Coq Modulo Theory. In *LPAR-21, 21st International Conference on Logic for Programming, Artificial Intelligence and Reasoning, Maun, Botswana, May 7-12, 2017 (EPIc Series in Computing)*, Thomas Eiter and David Sands (Eds.), Vol. 46. EasyChair, 474–489. <http://www.easychair.org/publications/paper/340342> 1388
- 1334 Chris Kapulkin and Peter LeFanu Lumsdaine. 2012. The simplicial model of univalent foundations. *arXiv preprint arXiv:1211.2851* (2012). 1389
- 1335 Conor McBride. 2000. *Dependently typed functional programs and their proofs*. Ph.D. Dissertation. University of Edinburgh. 1390
- 1336 Conor McBride. 2004. Epigram: Practical programming with dependent types. In *International School on Advanced Functional Programming*. Springer, 130–170. 1391
- 1337 Ulf Norell. 2007. *Towards a practical programming language based on dependent type theory*. Vol. 32. Citeseer. 1392
- 1338 Nicolas Oury. 2005. Extensionality in the calculus of constructions. In *International Conference on Theorem Proving in Higher Order Logics*. Springer, 278–293. 1393
- 1339 Nicolas Oury. 2006. *Egalité et filtrage avec types dépendants dans le calcul des constructions inductives*. Ph.D. Dissertation. <http://www.theses.fr/2006PA112136> Thèse de doctorat dirigée par Paulin-Mohring, Christine Informatique Paris 11 2006. 1394
- 1340 Matthieu Sozeau. 2007. Program-ing Finger Trees in Coq. In *ICFP’07*. ACM Press, Freiburg, Germany, 13–24. <https://doi.org/10.1145/1291151.1291156> 1395
- 1341 Matthieu Sozeau. 2010. Equations: A Dependent Pattern-Matching Compiler. In *Interactive Theorem Proving, First International Conference, ITP 2010, Edinburgh, UK, July 11-14, 2010. Proceedings (Lecture Notes in Computer Science)*, Matt Kaufmann and Lawrence C. Paulson (Eds.), Vol. 6172. Springer, 419–434. https://doi.org/10.1007/978-3-642-14052-5_29 1396
- 1342 Thomas Streicher. 1993. *Investigations into intensional type theory*. The Univalent Foundations Program. 2013. *Homotopy Type Theory: Univalent Foundations of Mathematics*. Institute for Advanced Study. 1397
- 1343 Floris van Doorn, Herman Geuvers, and Freek Wiedijk. 2013. Explicit convertibility proofs in pure type systems. In *Proceedings of the Eighth ACM SIGPLAN international workshop on Logical frameworks & meta-languages: theory & practice*. ACM, 25–36. 1398
- 1344 Vladimir Voevodsky. 2013. A simple type system with two identity types. (2013). <https://ncatlab.org/homotopytypetheory/files/HTS.pdf> 1399

A Complementary rules

Equivalence relation.

$$\frac{\Gamma \vdash u : A}{\Gamma \vdash u \equiv u : A} \quad \frac{\Gamma \vdash u \equiv v : A}{\Gamma \vdash v \equiv u : A} \quad \frac{\Gamma \vdash u \equiv v : A \quad \Gamma \vdash v \equiv w : A}{\Gamma \vdash u \equiv w : A}$$

Congruence of type constructors.

$$\frac{\Gamma \vdash A_1 \equiv A_2 : s \quad \Gamma, x : A_1 \vdash B_1 \equiv B_2 : s'}{\Gamma \vdash \Pi(x : A_1). B_1 \equiv \Pi(x : A_2). B_2 : s''} (s, s', s'') \quad \frac{\Gamma \vdash A_1 \equiv A_2 : s \quad \Gamma, x : A_1 \vdash B_1 \equiv B_2 : s'}{\Gamma \vdash \Sigma(x : A_1). B_1 \equiv \Sigma(x : A_2). B_2 : s''} (s, s', s'')$$

$$\frac{\Gamma \vdash A_1 \equiv A_2 : s \quad \Gamma \vdash u_1 \equiv u_2 : A_1 \quad \Gamma \vdash v_1 \equiv v_2 : A_1}{\Gamma \vdash u_1 =_{A_1} v_1 \equiv u_2 =_{A_2} v_2 : s}$$

Congruence of λ -calculus terms.

$$\frac{\Gamma \vdash A_1 \equiv A_2 : s \quad \Gamma, x : A_1 \vdash B_1 \equiv B_2 : s' \quad \Gamma, x : A_1 \vdash t_1 \equiv t_2 : B_1}{\Gamma \vdash \lambda(x : A_1). B_1. t_1 \equiv \lambda(x : A_2). B_2. t_2 : \Pi(x : A_1). B_1}$$

$$\frac{\Gamma \vdash A_1 \equiv A_2 : s \quad \Gamma, x : A_1 \vdash B_1 \equiv B_2 : s' \quad \Gamma \vdash t_1 \equiv t_2 : \Pi(x : A_1). B_1 \quad \Gamma \vdash u_1 \equiv u_2 : A_1}{\Gamma \vdash t_1 @_{x:A_1.B_1} u_1 \equiv t_2 @_{x:A_1.B_1} u_2 : B_1[x \leftarrow u_1]}$$

$$\frac{\Gamma \vdash A_1 \equiv A_2 : s \quad \Gamma \vdash u_1 \equiv u_2 : A_1 \quad \Gamma, x : A_1 \vdash B_1 \equiv B_2 : s' \quad \Gamma \vdash v_1 \equiv v_2 : B_1[x \leftarrow u_1]}{\Gamma \vdash \langle u_1; v_1 \rangle_{x:A_1.B_1} \equiv \langle u_2; v_2 \rangle_{x:A_2.B_2} : \Sigma(x : A_1). B_1}$$

$$\frac{\Gamma \vdash A_1 \equiv A_2 : s \quad \Gamma, x : A_1 \vdash B_1 \equiv B_2 : s' \quad \Gamma \vdash p_1 \equiv p_2 : \Sigma(x : A_1). B_1}{\Gamma \vdash \pi_1^{x:A_1.B_1} p_1 \equiv \pi_1^{x:A_2.B_2} p_2 : A_1}$$

$$\frac{\Gamma \vdash A_1 \equiv A_2 : s \quad \Gamma, x : A_1 \vdash B_1 \equiv B_2 : s' \quad \Gamma \vdash p_1 \equiv p_2 : \Sigma(x : A_1). B_1}{\Gamma \vdash \pi_2^{x:A_1.B_1} p_1 \equiv \pi_2^{x:A_2.B_2} p_2 : B_1[x \leftarrow \pi_1^{x:A_1.B_1} p_1]}$$

Congruence of equality terms.

$$\frac{\Gamma \vdash A_1 \equiv A_2 : s \quad \Gamma \vdash u_1 \equiv u_2 : A}{\Gamma \vdash \text{refl}_{A_1} u_1 \equiv \text{refl}_{A_2} u_2 : u_1 =_{A_1} u_1}$$

$$\frac{\Gamma \vdash A_1 \equiv A_2 : s \quad \Gamma \vdash u_1 \equiv u_2 : A_1 \quad \Gamma \vdash v_1 \equiv v_2 : A_1 \quad \Gamma, x : A_1, e : u_1 =_{A_1} x \vdash P_1 \equiv P_2 : s' \quad \Gamma \vdash p_1 \equiv p_2 : u_1 =_{A_1} v_1 \quad \Gamma \vdash w_1 \equiv w_2 : P_1[x \leftarrow u_1, e \leftarrow \text{refl}_{A_1} u_1]}{\Gamma \vdash J(A_1, u_1, x.e.P_1, w_1, v_1, p_1) \equiv J(A_2, u_2, x.e.P_2, w_2, v_2, p_2) : P[x \leftarrow v_1, e \leftarrow p_1]}$$

$$\frac{\Gamma \vdash A_1 \equiv A_2 : s \quad \Gamma, x : A_1 \vdash B_1 \equiv B_2 : s' \quad \Gamma \vdash f_1 \equiv f_2 : \Pi(x : A_1). B_1 \quad \Gamma \vdash g_1 \equiv g_2 : \Pi(x : A_1). B_1 \quad \Gamma \vdash e_1 \equiv e_2 : \Pi(x : A_1). f_1 @_{x:A_1.B_1} x =_{B_1} g_1 @_{x:A_1.B_1} x}{\Gamma \vdash \text{funext}(x : A_1, B_1, f_1, g_1, e_1) \equiv \text{funext}(x : A_2, B_2, f_2, g_2, e_2) : f_1 = g_1}$$

$$\frac{\Gamma \vdash A_1 \equiv A_2 \quad \Gamma \vdash u_1 \equiv u_2 : A_1 \quad \Gamma \vdash v_1 \equiv v_2 : A_2 \quad \Gamma \vdash p_1 \equiv p_2 : u_1 =_{A_1} v_1 \quad \Gamma \vdash q_1 \equiv q_2 : u_1 =_{A_1} v_1}{\Gamma \vdash \text{uip}(A_1, u_1, v_1, p_1, q_1) \equiv \text{uip}(A_2, u_2, v_2, p_2, q_2) : p_1 = q_1}$$

Figure 6. Congruence rules

B Proof of the fundamental lemma

Lemma B.1 (Fundamental lemma). *Let t_1 and t_2 be two terms. If $\Gamma, \Gamma_1 \vdash t_1 : T_1$ and $\Gamma, \Gamma_2 \vdash t_2 : T_2$ and $t_1 \sim t_2$ then there exists p such that $\Gamma, \text{Pack } \Gamma_1 \Gamma_2 \vdash p : t_1[\gamma_1]_{T_1[\gamma_1]} \cong_{T_2[\gamma_2]} t_2[\gamma_2]$.*

For readability we will abbreviate the left and right substitutions $_{-}[\gamma_1]$ and $_{-}[\gamma_2]$ by \uparrow and \downarrow respectively.

Proof. We prove it by induction on the derivation of $t_1 \sim t_2$.

- VAR

$$\frac{}{x \sim x}$$

If x belongs to Γ , we apply reflexivity—together with uniqueness of typing (2.2)—to conclude. Otherwise, $\text{Proj}_e x$ has the expected type (since $x[\gamma_1] \equiv \text{Proj}_1 x$ and $x[\gamma_2] \equiv \text{Proj}_2 x$).

- TRANSPORTLEFT

$$\frac{t_1 \sim t_2}{p_* t_1 \sim t_2}$$

We have $\Gamma, \Gamma_1 \vdash p_* t_1 : T_1$ and $\Gamma, \Gamma_2 \vdash t_2 : T_2$. By inversion (2.3) we have $\Gamma, \Gamma_1 \vdash p : T'_1 = T_1$ and $\Gamma, \Gamma_1 \vdash t_1 : T'_1$. Then by induction hypothesis we have e such that $\Gamma, \Gamma_p \vdash e : t_1 \uparrow \cong_{T_2 \downarrow} t_2 \downarrow$. From transitivity and symmetry we only need to provide a proof of $t_1 \uparrow \cong p \downarrow_* t_1 \downarrow$ which is inhabited by $\langle p \downarrow; \text{refl } (p \downarrow_* t_1 \downarrow) \rangle_{-,-}$.

- TRANSPORTRIGHT

$$\frac{t_1 \sim t_2}{t_1 \sim p_* t_2}$$

Similarly.

- PRODUCT

$$\frac{A_1 \sim A_2 \quad B_1 \sim B_2}{\Pi(x : A_1). B_1 \sim \Pi(x : A_2). B_2}$$

We have $\Gamma, \Gamma_1 \vdash \Pi(x : A_1). B_1 : T_1$ and $\Gamma, \Gamma_2 \vdash \Pi(x : A_2). B_2 : T_2$ so by inversion (2.3) we have $\Gamma, \Gamma_1 \vdash A_1 : s_1$ and $\Gamma, \Gamma_1, x : A_1 \vdash B_1 : s'_1$ and $\Gamma, \Gamma_1 \vdash s''_1 \equiv T_1$ for $(s_1, s'_1, s''_1) \in \mathbb{R}$ (and similarly with 2s). By induction hypothesis we have $\Gamma, \Gamma_p \vdash p_A : A_1 \uparrow \cong_{A_2 \downarrow} A_2 \downarrow$ and $\Gamma, \Gamma_p, x : \text{Pack } A_1 A_2 \vdash p_B : B_1 \uparrow \cong_{B_2 \downarrow} B_2 \downarrow$ hence the result (using UIP and functional extensionality, refer to the formalisation and especially to the file `Quotes.v` for more details on how to realise this equality).

- EQUALITY

$$\frac{A_1 \sim A_2 \quad u_1 \sim u_2 \quad v_1 \sim v_2}{u_1 =_{A_1} v_1 \sim u_2 =_{A_2} v_2}$$

We have $\Gamma, \Gamma_1 \vdash u_1 =_{A_1} v_1 : T_1$ and $\Gamma, \Gamma_2 \vdash u_2 =_{A_2} v_2 : T_2$ so, by inversion (2.3), we have $\Gamma, \Gamma_1 \vdash A_1 : s_1$ and $\Gamma, \Gamma_1 \vdash u_1 : A_1$ and $\Gamma, \Gamma_1 \vdash v_1 : A_1$ as well as $\Gamma, \Gamma_1 \vdash s_1 \equiv T_1$ (and the same with 2s). By induction hypothesis we thus have $\Gamma, \Gamma_p \vdash p_A : A_1 \cong_{A_2} A_2$ and $\Gamma, \Gamma_p \vdash p_u : u_1 \cong_{u_2} u_2$ and $\Gamma, \Gamma_p \vdash p_v : v_1 \cong_{v_2} v_2$. We can thus conclude.

- REFLEXIVITY

$$\frac{}{s \sim s}$$

This one holds by reflexivity and uniqueness of typing (2.2) (indeed, $s \uparrow \equiv s$ and $s \downarrow \equiv s$).

- LAMBDA

$$\frac{A_1 \sim A_2 \quad B_1 \sim B_2 \quad t_1 \sim t_2}{\lambda(x : A_1). B_1.t_1 \sim \lambda(x : A_2). B_2.t_2}$$

We have $\Gamma, \Gamma_1 \vdash \lambda(x : A_1). B_1.t_1 : T_1$ and $\Gamma, \Gamma_2 \vdash \lambda(x : A_2). B_2.t_2 : T_2$, thus, by inversion 2.3 the subterms are well-typed and we can apply induction hypothesis. The conclusion follows similarly to the Π case.

- APPLICATION

$$\frac{t_1 \sim t_2 \quad A_1 \sim A_2 \quad B_1 \sim B_2 \quad u_1 \sim u_2}{t_1 @_{x:A_1}. B_1 u_1 \sim t_2 @_{x:A_2}. B_2 u_2}$$

We have $\Gamma, \Gamma_1 \vdash t_1 @_{x:A_1}. B_1 u_1 : T_1$ and $\Gamma, \Gamma_2 \vdash t_2 @_{x:A_2}. B_2 u_2 : T_2$ which means by inversion (2.3) that the subterms are well-typed. We apply the induction hypothesis and then conclude.

- REFLEXIVITY

$$\frac{A_1 \sim A_2 \quad u_1 \sim u_2}{\text{refl}_{A_1} u_1 \sim \text{refl}_{A_2} u_2}$$

We have $\Gamma, \Gamma_1 \vdash \text{refl}_{A_1} u_1 : T_1$ and $\Gamma, \Gamma_2 \vdash \text{refl}_{A_2} u_2 : T_2$ so by inversion (2.3) we have $\Gamma, \Gamma_1 \vdash A_1 : s_1$ and $\Gamma, \Gamma_1 \vdash u_1 : A_1$ (same with 2s). By IH we have $A_1 \upharpoonright \cong A_2 \upharpoonright$ and $u_1 \upharpoonright_{A_1} \cong_{A_2} u_2 \upharpoonright$. The proof follows easily.

- FUNEXT

$$\frac{A_1 \sim A_2 \quad B_1 \sim B_2 \quad f_1 \sim f_2 \quad g_1 \sim g_2 \quad e_1 \sim e_2}{\text{funext}(x : A_1, B_1, f_1, g_1, e_1) \sim \text{funext}(x : A_2, B_2, f_2, g_2, e_2)}$$

Similar.

- UIP

$$\frac{A_1 \sim A_2 \quad u_1 \sim u_2 \quad v_1 \sim v_2 \quad p_1 \sim p_2 \quad q_1 \sim q_2}{\text{uip}(A_1, u_1, v_1, p_1, q_1) \sim \text{uip}(A_2, u_2, v_2, p_2, q_2)}$$

Similar.

- J

$$\frac{A_1 \sim A_2 \quad u_1 \sim u_2 \quad P_1 \sim P_2 \quad w_1 \sim w_2 \quad v_1 \sim v_2 \quad p_1 \sim p_2}{J(A_1, u_1, x.e.P_1, w_1, v_1, p_1) \sim J(A_2, u_2, x.e.P_2, w_2, v_2, p_2)}$$

Similar.

□

C Correctness of the translation

Theorem C.1 (Translation).

- If $\vdash_x \Gamma$ then there exists $\bar{\Gamma} \in \llbracket \vdash_x \Gamma \rrbracket$,
- If $\Gamma \vdash_x t : T$ then for any $\bar{\Gamma} \in \llbracket \vdash_x \Gamma \rrbracket$ there exist \bar{t} and \bar{T} such that $\bar{\Gamma} \vdash \bar{t} : \bar{T} \in \llbracket \Gamma \vdash_x t : T \rrbracket$,
- If $\Gamma \vdash_x u \equiv v : A$ then for any $\bar{\Gamma} \in \llbracket \vdash_x \Gamma \rrbracket$ there exist $A \sqsubset \bar{A}, A \sqsubset \bar{A}', u \sqsubset \bar{u}, v \sqsubset \bar{v}$ and \bar{e} such that $\bar{\Gamma} \vdash \bar{e} : \bar{u}_{\bar{A}} \cong_{\bar{A}'} \bar{v}$.

Proof. We prove the theorem by induction on the derivation in the extensional type theory. In most cases we need to assume some $\bar{\Gamma}$, translation of the context, we will implicitly refer to $\bar{\Gamma}$ in such cases as the one given as hypothesis.

- EMPTY

$$\frac{}{\vdash_x \bullet}$$

We have $\vdash \bullet \in \llbracket \vdash_x \bullet \rrbracket$.

- EXTEND

$$\frac{\vdash_x \Gamma \quad \Gamma \vdash_x A}{\vdash_x \Gamma, x : A} (x \notin \Gamma)$$

By IH we have $\vdash \bar{\Gamma} \in \llbracket \vdash_x \Gamma \rrbracket$ and, using $\bar{\Gamma}$ as well as lemma 4.2, $\bar{\Gamma} \vdash \bar{A} : s \in \llbracket \Gamma \vdash_x A : s \rrbracket$. Thus $\vdash \bar{\Gamma}, x : \bar{A} \in \llbracket \vdash_x \Gamma, x : A \rrbracket$.

- SORT

$$\frac{\vdash_x \Gamma}{\Gamma \vdash_x s : s'} (s, s')$$

We have $\bar{\Gamma} \vdash s : s' \in \llbracket \Gamma \vdash_x s : s' \rrbracket$.

- PRODUCT

$$\frac{\Gamma \vdash_x A : s \quad \Gamma, x : A \vdash_x B : s'}{\Gamma \vdash_x \Pi(x : A). B : s''} (s, s', s'')$$

By IH and lemma 4.2 we have $\bar{\Gamma} \vdash \bar{A} : s$, meaning $\vdash \bar{\Gamma}, x : \bar{A} \in \llbracket \vdash_x \Gamma, x : A \rrbracket$, and then $\bar{\Gamma}, x : \bar{A} \vdash \bar{B} : s'$. We thus conclude $\bar{\Gamma} \vdash \Pi(x : \bar{A}). \bar{B} : s'' \in \llbracket \Gamma \vdash_x \Pi(x : A). B : s'' \rrbracket$.

- SIGMA

$$\frac{\Gamma \vdash_x A : s \quad \Gamma, x : A \vdash_x B : s'}{\Gamma \vdash_x \Sigma(x : A). B : s''} (s, s', s'')$$

Similar.

1761 • EQUALITY

$$\frac{\Gamma \vdash_x A : s \quad \Gamma \vdash_x u : A \quad \Gamma \vdash_x v : A}{\Gamma \vdash_x u =_A v : s}$$

1762 By IH and lemma 4.2 we have $\bar{\Gamma} \vdash \bar{A} : s$, and—using lemma 4.3—we also have $\bar{\Gamma} \vdash \bar{u} : \bar{A}$ and $\bar{\Gamma} \vdash \bar{v} : \bar{A}$. Then
1763 $\bar{\Gamma} \vdash \bar{u} =_{\bar{A}} \bar{v} : s \in \llbracket \Gamma \vdash_x u =_A v : s \rrbracket$.

1764 • VARIABLE

$$\frac{\vdash_x \Gamma \quad (x : A) \in \Gamma}{\Gamma \vdash_x x : A}$$

1765 We have $\vdash \bar{\Gamma} \in \llbracket \vdash_x \Gamma \rrbracket$ (as we assumed, this is not an instance of the induction hypothesis) and $(x : A) \in \Gamma$. By definition
1766 of $\Gamma \sqsubset \bar{\Gamma}$ we also have some $(x : \bar{A}) \in \bar{\Gamma}$ with $A \sqsubset \bar{A}$, thus $\bar{\Gamma} \vdash x : \bar{A} \in \llbracket \Gamma \vdash_x x : A \rrbracket$.

1767 • CONVERSION

$$\frac{\Gamma \vdash_x u : A \quad \Gamma \vdash_x A \equiv B}{\Gamma \vdash_x u : B}$$

1768 By IH and lemma 3.3 we have $\bar{\Gamma} \vdash \bar{e} : \bar{A} = \bar{B}$ which implies $\bar{\Gamma} \vdash \bar{A} \in \llbracket \Gamma \vdash_x A \rrbracket$ by inversion (2.3), thus, from lemma 4.3 and
1769 IH we get $\bar{\Gamma} \vdash \bar{u} : \bar{A}$, yielding $\bar{\Gamma} \vdash \bar{e}_* \bar{u} : \bar{B} \in \llbracket \Gamma \vdash_x u : B \rrbracket$.

1770 • LAMBDA

$$\frac{\Gamma \vdash_x A : s \quad \Gamma, x : A \vdash_x B : s' \quad \Gamma, x : A \vdash_x t : B}{\Gamma \vdash_x \lambda(x : A).B.t : \Pi(x : A). B}$$

1771 By IH and lemma 4.2 we have $\bar{\Gamma} \vdash \bar{A} : s$ and thus $\vdash \bar{\Gamma}, x : \bar{A} \in \llbracket \vdash_x \Gamma, x : A \rrbracket$, meaning we can apply IH and lemma 4.2 to
1772 the second hypothesis to get $\bar{\Gamma}, x : \bar{A} \vdash \bar{B} : s' \in \llbracket \Gamma, x : A \vdash_x B : s' \rrbracket$ and then IH and lemma 4.3 to get $\bar{\Gamma}, x : \bar{A} \vdash \bar{t} : \bar{B} \in$
1773 $\llbracket \Gamma, x : A \vdash_x t : B \rrbracket$. All of this yields $\bar{\Gamma} \vdash \lambda(x : \bar{A}).\bar{B}.\bar{t} : \Pi(x : \bar{A}). \bar{B} \in \llbracket \Gamma \vdash_x \lambda(x : A).B.t : \Pi(x : A). B \rrbracket$.

1774 • APPLICATION

$$\frac{\Gamma \vdash_x A : s \quad \Gamma, x : A \vdash_x B : s' \quad \Gamma \vdash_x t : \Pi(x : A). B \quad \Gamma \vdash_x u : A}{\Gamma \vdash_x t @_{x:A.B} u : B[x \leftarrow u]}$$

1775 Using IH together with lemmata 4.2 and 4.3 we get $\bar{\Gamma} \vdash \bar{A} : s$ and $\bar{\Gamma}, x : \bar{A} \vdash \bar{B} : s'$ and $\bar{\Gamma} \vdash \bar{t} : \Pi(x : \bar{A}). \bar{B}$ and $\bar{\Gamma} \vdash \bar{u} : \bar{A}$
1776 meaning we can conclude $\bar{\Gamma} \vdash \bar{t} @_{x:\bar{A}.\bar{B}} \bar{u} : \bar{B}[x \leftarrow \bar{u}] \in \llbracket \Gamma \vdash_x t @_{x:A.B} u : B[x \leftarrow u] \rrbracket$.

1777 • PAIR

$$\frac{\Gamma \vdash_x u : A \quad \Gamma \vdash_x A : s \quad \Gamma, x : A \vdash_x B : s' \quad \Gamma \vdash_x v : B[x \leftarrow u]}{\Gamma \vdash_x \langle u; v \rangle_{x:A.B} : \Sigma(x : A). B}$$

1778 Using IH with lemmata 4.2 and 4.3 we translate all the hypotheses to conclude $\bar{\Gamma} \vdash \langle \bar{u}; \bar{v} \rangle_{x:\bar{A}.\bar{B}} : \Sigma(x : \bar{A}). \bar{B} \in \llbracket \Gamma \vdash_x$
1779 $\langle u; v \rangle_{x:A.B} : \Sigma(x : A). B \rrbracket$.

1780 • PROJ₁

$$\frac{\Gamma \vdash_x p : \Sigma(x : A). B}{\Gamma \vdash_x \pi_1^{x:A.B} p : A}$$

1781 Similar.

1782 • PROJ₂

$$\frac{\Gamma \vdash_x p : \Sigma(x : A). B}{\Gamma \vdash_x \pi_2^{x:A.B} p : B[x \leftarrow \pi_1^{x:A.B} p]}$$

1783 Similar.

1784 • REFLEXIVITY

$$\frac{\Gamma \vdash_x A : s \quad \Gamma \vdash_x u : A}{\Gamma \vdash_x \text{refl}_A u : u =_A u}$$

1785 By IH we have $\bar{\Gamma} \vdash \bar{u} : \bar{A}$ and thus $\bar{\Gamma} \vdash \text{refl}_{\bar{A}} \bar{u} : \bar{u} =_{\bar{A}} \bar{u} \in \llbracket \Gamma \vdash_x \text{refl}_A u : u =_A u \rrbracket$.

1786 • J

$$\frac{\Gamma \vdash_x A : s \quad \Gamma \vdash_x u, v : A \quad \Gamma, x : A, e : u =_A x \vdash_x P : s' \quad \Gamma \vdash_x p : u =_A v \quad \Gamma \vdash_x w : P[x \leftarrow u, e \leftarrow \text{refl}_A u]}{\Gamma \vdash_x J(A, u, x.e.P, w, v, p) : P[x \leftarrow v, e \leftarrow p]}$$

1787 By IH and lemma 4.2 we have $\bar{\Gamma} \vdash \bar{A} : s$. From this and IH and lemma 4.3 we have $\bar{\Gamma} \vdash \bar{u}, \bar{v} : \bar{A}$. We can thus deduce
1788 $\vdash \bar{\Gamma}, x : \bar{A}, e : \bar{u} =_{\bar{A}} x \in \llbracket \Gamma, x : A, e : u =_A x \rrbracket$ which in turn gives us $\bar{\Gamma}, x : \bar{A}, e : \bar{u} =_{\bar{A}} x \vdash \bar{P} : s'$. Similarly we

also get $\bar{\Gamma} \vdash \bar{p} : \bar{u} =_{\bar{A}} \bar{v}$ and $\bar{\Gamma} \vdash \bar{w} : \bar{P}[x \leftarrow \bar{u}, e \leftarrow \text{refl}_{\bar{A}} \bar{u}]$. All of this allows us to conclude $\bar{\Gamma} \vdash \text{J}(\bar{A}, \bar{u}, x.e.\bar{P}, \bar{w}, \bar{v}, \bar{p}) : \bar{P}[x \leftarrow \bar{v}, e \leftarrow \bar{p}] \in \llbracket \Gamma \vdash_x \text{J}(A, u, x.e.P, w, v, p) : P[x \leftarrow v, e \leftarrow p] \rrbracket$.

- FUNEXT

$$\frac{\Gamma \vdash f, g : \Pi(x : A). B \quad \Gamma \vdash e : \Pi(x : A). f @_{x:A.B} x =_B g @_{x:A.B} x}{\Gamma \vdash \text{funext}(x : A, B, f, g, e) : f = g}$$

Similar.

- UIP

$$\frac{\Gamma \vdash_x e_1, e_2 : u =_A v}{\Gamma \vdash_x \text{uip}(A, u, v, e_1, e_2) : e_1 = e_2}$$

Similar.

- BETA

$$\frac{\Gamma \vdash_x A : s \quad \Gamma, x : A \vdash_x B : s' \quad \Gamma, x : A \vdash_x t : B \quad \Gamma \vdash_x u : A}{\Gamma \vdash_x (\lambda(x : A).B.t) @_{x:A.B} u \equiv t[x \leftarrow u] : B[x \leftarrow u]}$$

From IH and the lemmata, we even get the conversion, we conclude using reflexivity.

- PROJ₁-RED

$$\frac{\Gamma \vdash_x A : s \quad \Gamma \vdash_x u : A \quad \Gamma, x : A \vdash_x B : s' \quad \Gamma \vdash_x v : B[x \leftarrow u]}{\Gamma \vdash_x \pi_1^{x:A.B} \langle u; v \rangle_{x:A.B} \equiv u : A}$$

Likewise.

- PROJ₂-RED

$$\frac{\Gamma \vdash_x A : s \quad \Gamma \vdash_x u : A \quad \Gamma, x : A \vdash_x B : s' \quad \Gamma \vdash_x v : B[x \leftarrow u]}{\Gamma \vdash_x \pi_2^{x:A.B} \langle u; v \rangle_{x:A.B} \equiv v : B[x \leftarrow u]}$$

Likewise.

- J-RED

$$\frac{\Gamma \vdash_x A : U_i \quad \Gamma \vdash_x u : A \quad \Gamma, x : A, e : u =_A x \vdash_x P : U_j \quad \Gamma \vdash_x w : P[x \leftarrow u, e \leftarrow \text{refl}_A u]}{\Gamma \vdash_x \text{J}(A, u, x.e.P, w, u, \text{refl}_A u) \equiv w : P[x \leftarrow u, e \leftarrow \text{refl}_A u]}$$

Likewise.

- CONV-REFL

$$\frac{\Gamma \vdash_x u : A}{\Gamma \vdash_x u \equiv u : A}$$

We conclude from IH and reflexivity of \cong .

- CONV-SYM

$$\frac{\Gamma \vdash_x u \equiv v : A}{\Gamma \vdash_x v \equiv u : A}$$

We conclude from IH and symmetry of \cong .

- CONV-TRANS

$$\frac{\Gamma \vdash_x u \equiv v : A \quad \Gamma \vdash_x v \equiv w : A}{\Gamma \vdash_x u \equiv w : A}$$

We conclude from IH and transitivity of \cong .

- CONV-CONV

$$\frac{\Gamma \vdash_x t_1 \equiv t_2 : T_1 \quad \Gamma \vdash_x T_1 \equiv T_2}{\Gamma \vdash_x t_1 \equiv t_2 : T_2}$$

By IH (and lemma 3.3) we have $\bar{\Gamma} \vdash \bar{e} : \bar{t}_1 \bar{T}_1 \cong_{\bar{T}_1} \bar{t}_2$ and $\bar{\Gamma} \vdash \bar{p} : \bar{T}_1'' = \bar{T}_2$. Also from lemmata 3.4 and 3.3 we have $\bar{T}_1' = \bar{T}_1''$ and $\bar{T}_1 = \bar{T}_1''$, meaning we get $\bar{T}_1' = \bar{T}_2$ and $\bar{T}_1 = \bar{T}_2$. This allows us to conclude by transporting along the aforementioned equalities.

- CONV-PROD

$$\frac{\Gamma \vdash_x A_1 \equiv A_2 : s \quad \Gamma, x : A_1 \vdash_x B_1 \equiv B_2 : s'}{\Gamma \vdash_x \Pi(x : A_1). B_1 \equiv \Pi(x : A_2). B_2 : s''} (s, s', s'')$$

We conclude exactly like we did in the proof of lemma 3.4.

- All congruences hold like in proof of lemma 3.4.

- CONV-EQ

$$\frac{\Gamma \vdash_x e : u =_A v}{\Gamma \vdash_x u \equiv v : A}$$

By IH and lemma 4.2 we have $\bar{\Gamma} \vdash \bar{e} : \bar{u} =_{\bar{A}} \bar{v} \in \llbracket \Gamma \vdash_x e : u =_A v \rrbracket$ which yields the conclusion we wanted.

□

1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035

2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090