



**HAL**  
open science

## On extensions of symmetric operators

Namig Guliyev

► **To cite this version:**

Namig Guliyev. On extensions of symmetric operators. *Operators and Matrices*, 2020, 14 (1), pp.71-75. 10.7153/oam-2020-14-05 . hal-01852568

**HAL Id: hal-01852568**

**<https://hal.science/hal-01852568>**

Submitted on 2 Aug 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ON EXTENSIONS OF SYMMETRIC OPERATORS

NAMIG J. GULIYEV

ABSTRACT. We give an explicit description of all minimal self-adjoint extensions of a densely defined, closed symmetric operator in a Hilbert space with deficiency indices  $(1, 1)$ .

There is a widely used linearization technique in the theory of Sturm–Liouville problems with boundary and/or discontinuity conditions polynomially dependent on the eigenvalue parameter. One considers a Hilbert (or Pontryagin) space of the form  $L^2 \oplus \mathbb{C}^k$  and constructs a self-adjoint operator in this space such that the eigenvalue problem for this operator and the original boundary value problem become equivalent, in the sense that their eigenvalues coincide, the eigenfunctions of the latter problem are in one-to-one correspondence with the first components of the eigenvectors of the former problem, and so on (see, e.g., [1], [2], [8], and the references therein). Fulton [6, Remark 2.1] attributes this technique to Friedman [5, pp. 205–207]. The purpose of this short paper is to show that a straightforward generalization of this technique gives an explicit description of all minimal self-adjoint extensions of a densely defined, closed symmetric operator with deficiency indices  $(1, 1)$  (see below for definitions).

Let  $A$  be a densely defined, closed symmetric operator in a separable Hilbert space  $\mathfrak{H}$  with deficiency indices  $(1, 1)$ . Let  $\{\mathbb{C}, \Gamma_0, \Gamma_1\}$  be a *boundary triplet* for  $A^*$ . This means that  $\Gamma_0, \Gamma_1: \mathfrak{D}(A^*) \rightarrow \mathbb{C}$  are two linear mappings such that abstract Green’s identity

$$\langle A^*x, y \rangle_{\mathfrak{H}} - \langle x, A^*y \rangle_{\mathfrak{H}} = \Gamma_1x \cdot \overline{\Gamma_0y} - \Gamma_0x \cdot \overline{\Gamma_1y}, \quad x, y \in \mathfrak{D}(A^*)$$

holds and the mapping  $\Gamma: \mathfrak{D}(A^*) \rightarrow \mathbb{C}^2, x \mapsto (\Gamma_0x, \Gamma_1x)$  is surjective [4], [12, Chapter 14]. The domain of  $A$  then coincides with the kernel of  $\Gamma$ :

$$\mathfrak{D}(A) = \{x \in \mathfrak{D}(A^*) \mid \Gamma_0x = \Gamma_1x = 0\}. \quad (1)$$

If  $\tilde{A}$  is a self-adjoint operator in a Hilbert space  $\tilde{\mathfrak{H}} \supset \mathfrak{H}$  such that  $A \subset \tilde{A}$ , then  $\tilde{A}$  is called a *self-adjoint extension* of  $A$ . A self-adjoint extension  $\tilde{A}$  is called *minimal* if no nontrivial subspace of  $\tilde{\mathfrak{H}} \ominus \mathfrak{H}$  is reducing for  $\tilde{A}$  [10, Section 4], or equivalently [12, Lemma 5.17]

$$\tilde{\mathfrak{H}} = \overline{\text{span} \left\{ \left( \tilde{A} - \lambda I \right)^{-1} x \mid x \in \mathfrak{H}, \lambda \in \mathbb{C} \setminus \mathbb{R} \right\}},$$

where span denotes the linear span.

---

2010 *Mathematics Subject Classification.* 47A20, 47B25.

*Key words and phrases.* symmetric operator, self-adjoint extension, generalized resolvent.

A holomorphic operator-valued function  $R(\lambda)$  on  $\mathbb{C} \setminus \mathbb{R}$  is called a *generalized resolvent* of  $A$  if

$$R(\lambda) = P_{\mathfrak{H}} \left( \tilde{A} - \lambda I \right)^{-1} \Big|_{\mathfrak{H}}$$

for some self-adjoint extension  $\tilde{A}$ , where  $P_{\mathfrak{H}}$  is the orthogonal projection onto  $\mathfrak{H}$ . For every generalized resolvent there is a unique (up to unitary equivalence) minimal self-adjoint extension with this property (see [10, Theorem 8]). On the other hand, there is a one-to-one correspondence between generalized resolvents  $R(\lambda)$  and functions  $\omega(\lambda)$  holomorphic on the open upper half-plane  $\mathbb{C}_+$  with  $|\omega(\lambda)| \leq 1$ : for every  $x \in \mathfrak{H}$  the value  $y := R(\lambda)x$  satisfies the equation

$$A^*y - \lambda y = x$$

and the condition

$$(\omega(\lambda) - 1)\Gamma_1 y - i(\omega(\lambda) + 1)\Gamma_0 y = 0$$

(see [3]). Denoting

$$f(\lambda) := \frac{i\omega(\lambda) + i}{1 - \omega(\lambda)}$$

we obtain a one-to-one correspondence between generalized resolvents  $R(\lambda)$  and Herglotz–Nevanlinna functions  $f$ , i.e. functions holomorphic on  $\mathbb{C}_+$  with  $\text{Im } f(\lambda) \geq 0$  (cf. [13, Subsection 1.2]). The above condition then becomes

$$\Gamma_1 y + f(\lambda)\Gamma_0 y = 0. \quad (2)$$

The ‘‘Dirichlet’’ condition  $\Gamma_0 y = 0$  corresponds to  $f = \infty$ . We will denote the generalized resolvent corresponding to  $f$  by  $R_f(\lambda)$ . It should be noted that complete parameterizations of all generalized resolvents in terms of Herglotz–Nevanlinna functions were first obtained independently by Naimark [11] and Krein [9].

We are now ready to construct our minimal self-adjoint extension corresponding to a Herglotz–Nevanlinna function  $f$ . This function has a unique representation of the form [12, Appendix F]

$$f(\lambda) = h_0\lambda + h + \int_{-\infty}^{+\infty} \left( \frac{1}{t - \lambda} - \frac{t}{1 + t^2} \right) d\sigma(t),$$

where  $h_0 \geq 0$ ,  $h \in \mathbb{R}$ , and

$$\int_{-\infty}^{+\infty} \frac{d\sigma(t)}{1 + t^2} < \infty$$

(the reader may refer to [7, Appendix A] for some examples of such representations). If  $h_0 > 0$  then we consider the Hilbert space  $\tilde{\mathfrak{H}} := \mathfrak{H} \oplus L^2(\mathbb{R}; d\sigma) \oplus \mathbb{C}$  with inner product given by

$$\langle \tilde{x}, \tilde{y} \rangle_{\tilde{\mathfrak{H}}} := \langle x_0, y_0 \rangle_{\mathfrak{H}} + \int_{-\infty}^{+\infty} x_1(t)\overline{y_1(t)} d\sigma(t) + \frac{x_2\overline{y_2}}{h_0}$$

for

$$\tilde{x} = \begin{pmatrix} x_0 \\ x_1 \\ x_2 \end{pmatrix}, \quad \tilde{y} = \begin{pmatrix} y_0 \\ y_1 \\ y_2 \end{pmatrix} \in \tilde{\mathfrak{H}},$$

and define the operator

$$\tilde{A}\tilde{x} := \begin{pmatrix} A^*x_0 \\ tx_1(t) - \Gamma_0x_0 \\ \Gamma_1x_0 + h\Gamma_0x_0 + \int_{-\infty}^{+\infty} \left( x_1(t) - \frac{t}{1+t^2}\Gamma_0x_0 \right) d\sigma(t) \end{pmatrix}$$

with

$$\mathfrak{D}(\tilde{A}) := \{\tilde{x} \in \tilde{\mathfrak{H}} \mid x_0 \in \mathfrak{D}(A^*), tx_1(t) - \Gamma_0x_0 \in L^2(\mathbb{R}; d\sigma), x_2 = -h_0\Gamma_0x_0\}.$$

If  $h_0 = 0$  then we set  $\tilde{\mathfrak{H}} := \mathfrak{H} \oplus L^2(\mathbb{R}; d\sigma)$ , and define  $\tilde{A}$  by

$$\tilde{A}\tilde{x} := \begin{pmatrix} A^*x_0 \\ tx_1(t) - \Gamma_0x_0 \end{pmatrix}$$

and

$$\mathfrak{D}(\tilde{A}) := \left\{ \tilde{x} \in \tilde{\mathfrak{H}} \mid x_0 \in \mathfrak{D}(A^*), tx_1(t) - \Gamma_0x_0 \in L^2(\mathbb{R}; d\sigma), \right. \\ \left. \Gamma_1x_0 + h\Gamma_0x_0 + \int_{-\infty}^{+\infty} \left( x_1(t) - \frac{t}{1+t^2}\Gamma_0x_0 \right) d\sigma(t) = 0 \right\}.$$

The extension  $\tilde{A}$  is canonical (i.e.  $\tilde{\mathfrak{H}} = \mathfrak{H}$ ) if and only if  $f$  is a real constant (or  $\infty$ ).

**Theorem.** *For each Herglotz–Nevanlinna function  $f$  the operator  $\tilde{A}$  defined above is a minimal self-adjoint extension of the operator  $A$  with the corresponding generalized resolvent  $R_f(\lambda)$ .*

*Proof.* We will consider the case  $h_0 > 0$ ; the other case can be proved similarly. To prove the self-adjointness, let  $\tilde{y}, \tilde{z} \in \tilde{\mathfrak{H}}$  be such that

$$\langle \tilde{A}\tilde{x}, \tilde{y} \rangle_{\tilde{\mathfrak{H}}} = \langle \tilde{x}, \tilde{z} \rangle_{\tilde{\mathfrak{H}}} \quad (3)$$

for all  $\tilde{x} \in \mathfrak{D}(\tilde{A})$ . Taking into account (1) and setting  $x_1(t) \equiv 0 = x_2$  we obtain

$$\langle Ax_0, y_0 \rangle_{\mathfrak{H}} = \langle x_0, z_0 \rangle_{\mathfrak{H}}, \quad x_0 \in \mathfrak{D}(A).$$

Hence  $y_0 \in \mathfrak{D}(A^*)$  and  $z_0 = A^*y_0$ . By surjectivity of  $\Gamma$ , there exists  $x_0 \in \mathfrak{D}(A^*)$  with  $\Gamma_0x_0 = 0$  and  $\Gamma_1x_0 = 1$ . Then for this  $x_0$  and  $x_1(t) \equiv 0$  the equality (3) gives  $y_2 = -h_0\Gamma_0y_0$ . Setting  $x_0 = 0$  in (3) we get

$$\int_{-\infty}^{+\infty} x_1(t) \overline{(ty_1(t) - \Gamma_0y_0 - z_1(t))} d\sigma(t) = 0$$

for all  $x_1(t) \in L^2(\mathbb{R}; d\sigma)$  with  $tx_1(t) \in L^2(\mathbb{R}; d\sigma)$ . Since such functions are dense in  $L^2(\mathbb{R}; d\sigma)$ , we obtain  $ty_1(t) - \Gamma_0y_0 = z_1(t) \in L^2(\mathbb{R}; d\sigma)$ . Finally choosing  $x_0 \in \mathfrak{D}(A^*)$  with  $\Gamma_0x_0 = 1$ , the equality (3) yields

$$z_2 = \Gamma_1y_0 + h\Gamma_0y_0 + \int_{-\infty}^{+\infty} \left( y_1(t) - \frac{t}{1+t^2}\Gamma_0y_0 \right) d\sigma(t).$$

Thus  $\tilde{y} \in \mathfrak{D}(\tilde{A})$  and  $\tilde{A}\tilde{y} = \tilde{z}$ .

To see that the generalized resolvent corresponding to  $\tilde{A}$  is  $R_f(\lambda)$ , it suffices to check that if

$$\tilde{y} = \begin{pmatrix} y_0 \\ y_1 \\ y_2 \end{pmatrix} = (\tilde{A} - \lambda I)^{-1} \begin{pmatrix} x \\ 0 \\ 0 \end{pmatrix}$$

then  $y_0$  satisfies (2), and this is straightforward. Finally, to check the minimality, we need to verify that the linear span of all  $\tilde{y}$  of this form with all possible values of  $x \in \mathfrak{H}$  and  $\lambda \in \mathbb{C} \setminus \mathbb{R}$  is dense in  $\tilde{\mathfrak{H}}$ . To this end, let  $\tilde{z} \in \tilde{\mathfrak{H}}$  be orthogonal to all such  $\tilde{y}$ , i.e.

$$\langle \tilde{z}, \tilde{y} \rangle_{\tilde{\mathfrak{H}}} = \langle z_0, y_0 \rangle_{\mathfrak{H}} + \int_{-\infty}^{+\infty} z_1(t) \frac{\overline{\Gamma_0 y_0}}{t - \lambda} d\sigma(t) - z_2 \overline{\Gamma_0 y_0} = 0$$

for all  $y_0 \in \mathfrak{D}(A^*)$ . Surjectivity of  $\Gamma$  implies  $z_0 = 0$  and

$$\int_{-\infty}^{+\infty} \frac{z_1(t)}{t - \lambda} d\sigma(t) = z_2, \quad \lambda \in \mathbb{C} \setminus \mathbb{R}.$$

Now the Stieltjes–Perron inversion formula [12, Theorem F.2] applied to the regular complex Borel measure  $z_1 d\sigma$  yields  $z_1(t) = 0$  for  $d\sigma$ -a.e.  $t$  and consequently  $z_2 = 0$ .  $\square$

#### REFERENCES

- [1] C. Bartels, S. Currie, M. Nowaczyk, and B. A. Watson, *Sturm–Liouville problems with transfer condition Herglotz dependent on the eigenparameter: Hilbert space formulation*, Integral Equations Operator Theory **90** (2018), no. 3, Art. 34, 20 pp. [arXiv:1804.07149](https://arxiv.org/abs/1804.07149)
- [2] J. Behrndt and F. Philipp, *Finite rank perturbations in Pontryagin spaces and a Sturm–Liouville problem with  $\lambda$ -rational boundary conditions*, Indefinite inner product spaces, Schur analysis, and differential equations, Birkhäuser/Springer, Cham, 2018, pp. 163–189.
- [3] V. M. Bruk, *A certain class of boundary value problems with a spectral parameter in the boundary condition* (Russian), Mat. Sb. (N.S.) **100(142)** (1976), no. 2, 210–216.
- [4] V. Derkach, *Boundary triplets, Weyl functions, and the Krein formula*, Operator theory, Springer, Basel, 2015, pp. 183–218.
- [5] B. Friedman, *Principles and techniques of applied mathematics*, John Wiley & Sons, Inc., New York, 1956.
- [6] C. T. Fulton, *Two-point boundary value problems with eigenvalue parameter contained in the boundary conditions*, Proc. Roy. Soc. Edinburgh Sect. A **77** (1977), no. 3-4, 293–308.
- [7] F. Gesztesy and E. Tsekanovskii, *On matrix-valued Herglotz functions*, Math. Nachr. **218** (2000), 61–138. [arXiv:funct-an/9712004](https://arxiv.org/abs/funct-an/9712004)
- [8] N. J. Guliyev, *Essentially isospectral transformations and their applications*, preprint. [arXiv:1708.07497](https://arxiv.org/abs/1708.07497)
- [9] M. G. Krein, *On Hermitian operators with deficiency indices one* (Russian), Dokl. Akad. Nauk SSSR **43** (1944), 339–342.
- [10] M. A. Naimark, *Spectral functions of a symmetric operator* (Russian), Izv. Akad. Nauk SSSR. Ser. Mat. **4** (1940), no. 3, 277–318.
- [11] M. A. Naimark, *On spectral functions of a symmetric operator* (Russian), Izv. Akad. Nauk SSSR. Ser. Mat. **7** (1943), no. 6, 285–296.
- [12] K. Schmüdgen, *Unbounded self-adjoint operators on Hilbert space*, Springer, Dordrecht, 2012.
- [13] A. V. Shtraus, *On spectral functions of differential operators* (Russian), Izv. Akad. Nauk SSSR. Ser. Mat. **19** (1955), no. 4, 201–220.

INSTITUTE OF MATHEMATICS AND MECHANICS, AZERBAIJAN NATIONAL ACADEMY OF SCIENCES,  
 9 B. VAHABZADEH STR., AZ1141, BAKU, AZERBAIJAN.  
*E-mail address:* njguliyev@gmail.com