



HAL
open science

Existence of strong solutions to a fluid–structure system with a structure given by a finite number of parameters

Guillaume Delay

► **To cite this version:**

Guillaume Delay. Existence of strong solutions to a fluid–structure system with a structure given by a finite number of parameters. 2019. hal-01859849v2

HAL Id: hal-01859849

<https://hal.science/hal-01859849v2>

Preprint submitted on 11 Apr 2019 (v2), last revised 6 Mar 2020 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Existence of strong solutions to a fluid–structure system with a structure given by a finite number of parameters*

Guillaume Delay[†]

November 28, 2018

Abstract. We study the existence of strong solutions to a 2d fluid–structure system. The fluid is modelled by the incompressible Navier–Stokes equations. The structure represents a steering gear and is described by a finite number of parameters. Its equations are derived from a virtual work principle. The global domain represents a wind tunnel and imposes mixed boundary conditions to the fluid velocity. Our method reposes on the analysis of the linearized system. Under a compatibility condition on the initial data, we can guarantee local existence in time of strong solutions to the fluid–structure problem.

MSC numbers: 74F10, 74H20, 74H25, 74H30, 76D03.

1 Introduction

The goal of this study is to prove the existence of a unique solution to a fluid–structure problem. The fluid is modelled by the incompressible Navier–Stokes equations and the structure, immersed in the fluid, is governed by a finite number of parameters.

For the sake of simplicity, only two parameters θ_1 and θ_2 are considered to describe the motion of the structure. However, all results remain valid for any finite number of parameters.

1.1 Modelling of the problem

The considered structure lies inside an open bounded domain $\Omega \subset \mathbb{R}^2$ and deforms itself over time. The couple of parameters (θ_1, θ_2) lies in an admissible domain \mathbb{D}_Θ which is an open connected subset of \mathbb{R}^2 . Let S_{ref} , a smooth closed connected subset of Ω , be the reference configuration for the structure (for instance S_{ref} is the volume occupied by the structure for $\theta_1 = \theta_2 = 0$). We consider a function \mathbf{X} defined on $\mathbb{D}_\Theta \times S_{\text{ref}}$ such that $\mathbf{X}(\theta_1, \theta_2, \mathbf{y})$ is the position of the matter associated to the point \mathbf{y} in the reference configuration S_{ref} .

The volume occupied by the structure for the parameters $(\theta_1, \theta_2) \in \mathbb{D}_\Theta$ is a closed bounded connected subset of Ω denoted $S(\theta_1, \theta_2) = \mathbf{X}(\theta_1, \theta_2, S_{\text{ref}})$. We further assume that for every $(\theta_1, \theta_2) \in \mathbb{D}_\Theta$, $S(\theta_1, \theta_2) \subset \Omega$, i.e. there is no contact between the structure $S(\theta_1, \theta_2)$ and the boundary of the domain $\partial\Omega$.

1.1.1 Motivations

Structures depending only on a finite number of parameters arise in the field of aeronautics. For instance, let us consider a steering gear structure. In a first approach, we can model this structure by two rigid solids. Solid S_1 is tied to the fixed frame by a pivoting link O and solid S_2 is tied to solid S_1 by a pivoting link P . The whole model is represented in Fig. 1a. Note that S_1 can be thought of as the aerofoil of a wing and S_2 as a steering gear such as an aileron. For a given $S_{\text{ref}} \subset \Omega$, the function \mathbf{X}^a representing the motion of this structure with respect to (θ_1, θ_2) is given below

$$\mathbf{X}^a(\theta_1, \theta_2, \mathbf{y}) = \chi_{S_1}(\mathbf{y})R_{\theta_1}\mathbf{y} + \chi_{S_2}(\mathbf{y})(R_{\theta_1}\mathbf{y}_P^{\text{ref}} + R_{\theta_1+\theta_2}(\mathbf{y} - \mathbf{y}_P^{\text{ref}})), \quad \forall \mathbf{y} \in S_{\text{ref}}, \quad \forall (\theta_1, \theta_2) \in \mathbb{D}_\Theta,$$

where $R_\theta = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$ is the rotation matrix of angle θ , $\mathbf{y}_P^{\text{ref}} = (y_{P,1}, y_{P,2})$ is the coordinate of the point P in the reference configuration S_{ref} and χ_E , the characteristic function over a set $E \subset \mathbb{R}^2$, is given below

$$\forall \mathbf{y} \in \Omega, \quad \chi_E(\mathbf{y}) = \begin{cases} 1 & \text{if } \mathbf{y} \in E, \\ 0 & \text{else.} \end{cases} \quad (1.1)$$

In the previous example, the admissible domain \mathbb{D}_Θ for (θ_1, θ_2) is chosen such that no overlaps of the structure occur.

*The author is partially supported by IFSMACS ANR–15–CE40–0010

[†]Institut de Mathématiques de Toulouse, Université Paul Sabatier – 118, route de Narbonne F-31062 Toulouse Cedex 9, France (guillaume.delay@math.univ-toulouse.fr or guillaume.delay@enpc.fr)




Figure 1: Three different kinds of structure deformation.

Note that for $\theta_2 \neq 0$, the function $\mathbf{X}^a(\theta_1, \theta_2, \cdot)$ is not a diffeomorphism as it is discontinuous through the interface $\partial S_1 \cap \partial S_2$ between the two solids. In the same way, for $\theta_2 \neq 0$, the velocity field is discontinuous inside the structure (we denote $\dot{\theta}_2$ the time derivative of θ_2). In other words, if we keep S_1 at rest and rotate S_2 around P , a discontinuity of the velocity appears through the interface between the two solids. This discontinuity can reduce the regularity expected for the fluid velocity. Indeed, if we assume no-slip boundary conditions between the fluid and the structure and if at time t the trace of the velocity is discontinuous on $\partial S(\theta_1(t), \theta_2(t))$, then a Sobolev embedding argument shows that we cannot hope for a better regularity in space for the velocity of the fluid than the Sobolev space $L^2(0, T; \mathbf{H}^1(\Omega \setminus S(\theta_1(t), \theta_2(t))))^1$, while for strong solutions we usually expect the velocity in the Sobolev space $L^2(0, T; \mathbf{H}^2(\Omega \setminus S(\theta_1(t), \theta_2(t))))^1$.

This loss of regularity would harm the estimates of the nonlinear terms (see Appendix B). That is why we consider a smooth approximation \mathbf{X}^b of the deformation \mathbf{X}^a .

In the sequel, $\mathbf{y} = (y_1, y_2)$ is the Lagrangian coordinate and $\mathbf{y}^\perp = (-y_2, y_1)$ is normal to \mathbf{y} . The behaviour of the smooth structure is represented in Fig. 1b, we give \mathbf{X}^b below

$$\mathbf{X}^b(\theta_1, \theta_2, \mathbf{y}) = g_{\theta_1}(y_1)\mathbf{e}_{r_1} + g_{\theta_2}(y_1)\mathbf{e}_{r_2} + y_2 \frac{\mathbf{N}(y_1)}{\|\mathbf{N}(y_1)\|}, \quad \mathbf{y} = (y_1, y_2) \in S_{\text{ref}}, \quad (\theta_1, \theta_2) \in \mathbb{D}_\Theta, \quad (1.2)$$

where g_{θ_1} and g_{θ_2} are real-valued functions, \mathbb{D}_Θ and S_{ref} are given on page 1. We use the notations: $\mathbf{e}_{r_1} = (\cos \theta_1, \sin \theta_1)$, $\mathbf{e}_{r_2} = (\cos(\theta_1 + \theta_2), \sin(\theta_1 + \theta_2))$, $\mathbf{N}(y_1) = g'_{\theta_1}(y_1)\mathbf{e}_{\theta_1} + g'_{\theta_2}(y_1)\mathbf{e}_{\theta_2}$, where $\mathbf{e}_{\theta_1} = \mathbf{e}_{r_1}^\perp$ and $\mathbf{e}_{\theta_2} = \mathbf{e}_{r_2}^\perp$. Moreover, we have $\|\mathbf{N}(y_1)\| = ((N_1(y_1))^2 + (N_2(y_1))^2)^{1/2}$, where N_i is the i^{th} coordinate of \mathbf{N} .

The function $y_1 \mapsto g_{\theta_1}(y_1)\mathbf{e}_{r_1} + g_{\theta_2}(y_1)\mathbf{e}_{r_2}$ gives for $(\theta_1, \theta_2) \in \mathbb{D}_\Theta$ the position of a reference curve. Every fibre of matter that is normal to this curve in the reference configuration stays normal when (θ_1, θ_2) changes. The normal direction to the curve at abscissa y_1 is given by $\mathbf{N}(y_1)$. This model is inspired by the fish-like model described in [28, Section 7].

To enforce smoothness of \mathbf{X}^b , g_{θ_1} and g_{θ_2} are taken as \mathcal{C}^∞ functions which are smooth approximations of respectively $y_{P,1} + (y_1 - y_{P,1})\chi_{[0, y_{P,1}]}(y_1)$ and $(y_1 - y_{P,1})\chi_{[y_{P,1}, y_{\text{max}}]}(y_1)$, where χ_I is defined in a similar way as (1.1) for $I \subset \mathbb{R}$. For instance, let $\varepsilon > 0$ and consider μ_ε a \mathcal{C}^∞ cut-off function such that

$$\begin{cases} \mu_\varepsilon(y_1) = 1, & \text{for } y_1 < y_{P,1}, \\ \mu_\varepsilon(y_1) \in [0, 1], & \text{for } y_{P,1} \leq y_1 \leq y_{P,1} + \varepsilon, \\ \mu_\varepsilon(y_1) = 0, & \text{for } y_{P,1} + \varepsilon < y_1. \end{cases}$$

Then, we can use

$$\begin{cases} g_{\theta_1}(y_1) = y_{P,1} + \mu_\varepsilon(y_1)(y_1 - y_{P,1}), \\ g_{\theta_2}(y_1) = (1 - \mu_\varepsilon(y_1))(y_1 - y_{P,1}), \end{cases}$$

in (1.2) to get a smooth deformation as represented in Fig. 1b. The velocity field of the structure is not any more discontinuous, we can thus expect the fluid velocity to have the usual regularity of strong solutions.

Remark 1.1. When ε tends to 0, these functions become

$$\begin{cases} g_{\theta_1}(y_1) = \chi_{[a, b]}(y_1)y_1 + \chi_{[b, c]}(y_1)y_{P,1}, \\ g_{\theta_2}(y_1) = \chi_{[b, c]}(y_1)(y_1 - y_{P,2}). \end{cases} \quad (1.3)$$

In this case, we recover the behaviour of a pivoting structure with two rigid solids (see Fig. 1c), corresponding to a transformation denoted \mathbf{X}^c . However, with this definition, the two solids overlap each other, so that we will not use it neither in the sequel. Also let us remark that the limit \mathbf{X}^c of our smooth approximation \mathbf{X}^b is not the original model \mathbf{X}^a .

¹These spaces are given here in an informal manner. They will be defined more precisely later.




Figure 2: Correspondance between real and reference structure configurations.

Now, let us show that we can choose \mathbb{D}_Θ such that for every $(\theta_1, \theta_2) \in \mathbb{D}_\Theta$, $\mathbf{X}^b(\theta_1, \theta_2, \cdot)$ is a \mathcal{C}^∞ diffeomorphism. We can compute the jacobian $\mathcal{J}_{\mathbf{X}^b}(\theta_1, \theta_2, \cdot)$ of $\mathbf{X}^b(\theta_1, \theta_2, \cdot)$. It fulfils

$$\det(\mathcal{J}_{\mathbf{X}^b}(\theta_1, \theta_2, \mathbf{y})) = \|\mathbf{N}(y_1)\| + \frac{y_2}{\|\mathbf{N}(y_1)\|^2} \sin(\theta_2)(g''_{\theta_1}(y_1)g'_{\theta_2}(y_1) - g''_{\theta_2}(y_1)g'_{\theta_1}(y_1)).$$

The detail of this computation is shown in Appendix A. This implies that for a given reference configuration and for θ_2 small enough, $\det(\mathcal{J}_{\mathbf{X}^b}(\theta_1, \theta_2, \mathbf{y})) > 0$ for every $\mathbf{y} \in S_{\text{ref}}$. Hence, according to the inverse theorem, the function $\mathbf{X}^b(\theta_1, \theta_2, \cdot)$ is a bijection. It is then a \mathcal{C}^∞ diffeomorphism for θ_2 small enough.

We shall therefore keep in mind only the example of \mathbf{X}^b (see Fig. 1b), though our original motivation was to deal with \mathbf{X}^a (see Fig. 1a). More generally, our approach will be applicable to many more choices of deformations \mathbf{X} . Let us list below the assumptions used in the sequel.

Modelling Assumptions.

- For every $\mathbf{y} \in S_{\text{ref}} = S(0, 0)$, $\mathbf{X}(0, 0, \mathbf{y}) = \mathbf{y}$. (1.4)
- S_{ref} is a smooth simply connected closed subset of Ω . (1.5)
- For every $(\theta_1, \theta_2) \in \mathbb{D}_\Theta$, we have $\mathbf{X}(\theta_1, \theta_2, S_{\text{ref}}) \subset \Omega$ and $\inf_{(\theta_1, \theta_2) \in \mathbb{D}_\Theta} d(\mathbf{X}(\theta_1, \theta_2, S_{\text{ref}}), \partial\Omega) > 0$. (1.6)
- For every $(\theta_1, \theta_2) \in \mathbb{D}_\Theta$, $\mathbf{X}(\theta_1, \theta_2, \cdot)$ is a \mathcal{C}^∞ diffeomorphism from S_{ref} to its image $S(\theta_1, \theta_2)$. (1.7)
- The function \mathbf{X} is \mathcal{C}^∞ on $\mathbb{D}_\Theta \times S_{\text{ref}}$. (1.8)
- The functions $\partial_{\theta_1}\mathbf{X}(\theta_1, \theta_2, \cdot)$ and $\partial_{\theta_2}\mathbf{X}(\theta_1, \theta_2, \cdot)$ form a free family in $\mathbf{L}^2(\partial S_{\text{ref}})$ for every (θ_1, θ_2) in \mathbb{D}_Θ . (1.9)

In (1.4), we have assumed that $S_{\text{ref}} = S(0, 0)$ to ease the study. In (1.6), we assume that the structure stays away from the boundary of the wind tunnel. Assumption (1.7) enables us to use a change of variables. This is a crucial step in our approach, as we shall see in Section 3.1. Assumption (1.8) ensures continuity of the velocity field inside the structure and on its boundary. This assumption could be weakened, as \mathcal{C}^n would be sufficient for n large enough, but we keep \mathcal{C}^∞ for simplicity. In our approach, Assumption (1.9) is natural and mandatory to determine the equations of the structure, as we shall see below in Section 1.1.2.

The inverse diffeomorphism of $\mathbf{X}(\theta_1, \theta_2, \cdot)$ whose existence is guaranteed by (1.7) is denoted $\mathbf{Y}(\theta_1, \theta_2, \cdot)$, we have

$$\forall (\theta_1, \theta_2) \in \mathbb{D}_\Theta, \quad \forall \mathbf{y} \in S_{\text{ref}}, \quad \mathbf{Y}(\theta_1, \theta_2, \mathbf{X}(\theta_1, \theta_2, \mathbf{y})) = \mathbf{y}. \quad (1.10)$$

The diffeomorphisms $\mathbf{X}(\theta_1, \theta_2, \cdot)$ and $\mathbf{Y}(\theta_1, \theta_2, \cdot)$ are illustrated in Fig. 2.

Remark 1.2. The choice of \mathbf{X}^b allows changes of volume for the structure and then for the fluid. In our setting where mixed boundary conditions are considered, this is not a problem. However, in the case, for instance, of homogeneous Dirichlet boundary conditions on $\partial\Omega$, the diffeomorphism \mathbf{X}^b would have to be modified to ensure a constant volume to the structure.

1.1.2 Dynamics of the structure

In order to simplify the equations of the structure, we consider the following assumption for the dynamics of the structure.

Modelling Assumption.

$$\left| \begin{array}{l} \bullet \text{ No friction and no elastic forces are considered in the structure.} \end{array} \right. \quad (1.11)$$

The equations satisfied by the structure are obtained by a virtual work principle [2, p. 14–17]. We know that the admissible parameters of the structure are $(\theta_1, \theta_2) \in \mathbb{D}_\Theta$, and that the admissible velocities \mathbf{v}_s satisfy

$$\mathbf{v}_s \in \text{Vect}(\partial_{\theta_1}\mathbf{X}(\theta_1, \theta_2, \cdot), \partial_{\theta_2}\mathbf{X}(\theta_1, \theta_2, \cdot)).$$

Thus, the virtual work principle can be formulated for every time $t \in [0, T]$ as

$$\left\{ \begin{array}{l} \text{Find } (\theta_1(t), \theta_2(t)) \in \mathbb{D}_\Theta, \text{ such that for every } \mathbf{w} \in \text{Vect}(\partial_{\theta_1}\mathbf{X}(\theta_1(t), \theta_2(t), \cdot), \partial_{\theta_2}\mathbf{X}(\theta_1(t), \theta_2(t), \cdot)), \\ \int_{S_{\text{ref}}} \rho \left(\frac{d^2}{dt^2}(\mathbf{X}(\theta_1(t), \theta_2(t), \mathbf{y})) - \mathbf{f}_{\text{body}}(t, \mathbf{X}(\theta_1(t), \theta_2(t), \mathbf{y})) \right) \cdot \mathbf{w}(\mathbf{y}) \, d\mathbf{y} \\ - \int_{\partial S(\theta_1(t), \theta_2(t))} \mathbf{f}_{\mathcal{F} \rightarrow S}(\gamma_x) \cdot \mathbf{w}(\mathbf{Y}(\theta_1(t), \theta_2(t), \gamma_x)) \, d\gamma_x = 0, \end{array} \right. \quad (1.12)$$

where \mathbf{f}_{body} is a distributed source term in the body (modelling for instance the gravity), ρ is a positive constant that represents the mass per unit volume of the structure in the reference configuration S_{ref} and $\mathbf{f}_{\mathcal{F} \rightarrow S}$ is the force exerted by the fluid on the structure along $\partial S(\theta_1(t), \theta_2(t))$.

Note that the presence of \mathbf{f}_{body} is compatible with Assumption (1.11), as this term represents external forces. It does not depend on θ_1, θ_2 and their derivatives.

Remark 1.3. Assumption (1.11) has been used in (1.12) as no interior works have been considered.

Let us denote respectively $\dot{\theta}_j$ and $\ddot{\theta}_j$ the first and second time derivatives of the function θ_j . Then, the velocity of the structure can be written as

$$\mathbf{v}_s(t, \mathbf{y}) = \frac{d}{dt} \mathbf{X}(\theta_1(t), \theta_2(t), \mathbf{y}) = \sum_{j=1}^2 \dot{\theta}_j(t) \partial_{\theta_j} \mathbf{X}(\theta_1(t), \theta_2(t), \mathbf{y}), \quad \forall t \in [0, T], \quad \forall \mathbf{y} \in S_{\text{ref}}, \quad (1.13)$$

and its acceleration as

$$\frac{d}{dt} \mathbf{v}_s(t, \mathbf{y}) = \frac{d^2}{dt^2}(\mathbf{X}(\theta_1(t), \theta_2(t), \mathbf{y})) = \sum_{j=1}^2 \ddot{\theta}_j(t) \partial_{\theta_j} \mathbf{X}(\theta_1(t), \theta_2(t), \mathbf{y}) + \sum_{j,k=1}^2 \dot{\theta}_j(t) \dot{\theta}_k(t) \partial_{\theta_j \theta_k} \mathbf{X}(\theta_1(t), \theta_2(t), \mathbf{y}).$$

Now, problem (1.12) can be rewritten as follows

$$\left\{ \begin{array}{l} \text{Find } (\theta_1, \theta_2) \in \mathbb{D}_\Theta, \text{ such that for every } i \in \{1, 2\}, \text{ we have,} \\ \int_{S_{\text{ref}}} \rho \sum_{j=1}^2 \ddot{\theta}_j \partial_{\theta_j} \mathbf{X}(\theta_1, \theta_2, \mathbf{y}) \cdot \partial_{\theta_i} \mathbf{X}(\theta_1, \theta_2, \mathbf{y}) \, d\mathbf{y} = - \int_{S_{\text{ref}}} \rho \sum_{j,k=1}^2 \dot{\theta}_j \dot{\theta}_k \partial_{\theta_j \theta_k} \mathbf{X}(\theta_1, \theta_2, \mathbf{y}) \cdot \partial_{\theta_i} \mathbf{X}(\theta_1, \theta_2, \mathbf{y}) \, d\mathbf{y} \\ + \int_{S_{\text{ref}}} \mathbf{f}_{\text{body}}(t, \mathbf{X}(\theta_1, \theta_2, \mathbf{y})) \cdot \partial_{\theta_i} \mathbf{X}(\theta_1, \theta_2, \mathbf{y}) \, d\mathbf{y} \\ + \int_{\partial S(\theta_1, \theta_2)} \mathbf{f}_{\mathcal{F} \rightarrow S}(\gamma_x) \cdot \partial_{\theta_i} \mathbf{X}(\theta_1, \theta_2, \mathbf{Y}(\theta_1, \theta_2, \gamma_x)) \, d\gamma_x. \end{array} \right.$$

Let us denote the structure body source term

$$(f_s)_i = \int_{S_{\text{ref}}} \mathbf{f}_{\text{body}}(t, \mathbf{X}(\theta_1, \theta_2, \mathbf{y})) \cdot \partial_{\theta_i} \mathbf{X}(\theta_1, \theta_2, \mathbf{y}) \, d\mathbf{y}. \quad (1.14)$$

On a matrix form, the equations of the structure read

$$\mathcal{M}_{\theta_1, \theta_2} \begin{pmatrix} \ddot{\theta}_1 \\ \ddot{\theta}_2 \end{pmatrix} = \mathbf{M}_\mathbf{I}(\theta_1, \theta_2, \dot{\theta}_1, \dot{\theta}_2) + \mathbf{M}_\mathbf{A}(\theta_1, \theta_2, \mathbf{f}_{\mathcal{F} \rightarrow S}) + \mathbf{f}_s \quad \text{on } (0, T), \quad (1.15)$$

where $\mathbf{f}_s = ((f_s)_1, (f_s)_2)$ and

$$\mathcal{M}_{\theta_1, \theta_2} = \begin{pmatrix} (\partial_{\theta_1} \mathbf{X}(\theta_1, \theta_2, \cdot), \partial_{\theta_1} \mathbf{X}(\theta_1, \theta_2, \cdot))_S & (\partial_{\theta_2} \mathbf{X}(\theta_1, \theta_2, \cdot), \partial_{\theta_1} \mathbf{X}(\theta_1, \theta_2, \cdot))_S \\ (\partial_{\theta_1} \mathbf{X}(\theta_1, \theta_2, \cdot), \partial_{\theta_2} \mathbf{X}(\theta_1, \theta_2, \cdot))_S & (\partial_{\theta_2} \mathbf{X}(\theta_1, \theta_2, \cdot), \partial_{\theta_2} \mathbf{X}(\theta_1, \theta_2, \cdot))_S \end{pmatrix} \in \mathbb{R}^{2 \times 2}, \quad (1.16)$$

$$\mathbf{M}_\mathbf{I}(\theta_1, \theta_2, \dot{\theta}_1, \dot{\theta}_2) = \begin{pmatrix} -(\dot{\theta}_1^2 \partial_{\theta_1 \theta_1} \mathbf{X}(\theta_1, \theta_2, \cdot) + 2\dot{\theta}_1 \dot{\theta}_2 \partial_{\theta_1 \theta_2} \mathbf{X}(\theta_1, \theta_2, \cdot) + \dot{\theta}_2^2 \partial_{\theta_2 \theta_2} \mathbf{X}(\theta_1, \theta_2, \cdot), \partial_{\theta_1} \mathbf{X}(\theta_1, \theta_2, \cdot))_S \\ -(\dot{\theta}_1^2 \partial_{\theta_1 \theta_1} \mathbf{X}(\theta_1, \theta_2, \cdot) + 2\dot{\theta}_1 \dot{\theta}_2 \partial_{\theta_1 \theta_2} \mathbf{X}(\theta_1, \theta_2, \cdot) + \dot{\theta}_2^2 \partial_{\theta_2 \theta_2} \mathbf{X}(\theta_1, \theta_2, \cdot), \partial_{\theta_2} \mathbf{X}(\theta_1, \theta_2, \cdot))_S \end{pmatrix} \in \mathbb{R}^2, \quad (1.17)$$




Figure 3: The geometrical configuration.

where $(\cdot, \cdot)_S$ is the scalar product

$$(\Phi, \Psi)_S = \int_{S_{\text{ref}}} \rho \Phi(\mathbf{y}) \cdot \Psi(\mathbf{y}) \, d\mathbf{y}, \quad (1.18)$$

and

$$\mathbf{M}_A(\theta_1, \theta_2, \mathbf{f}_{\mathcal{F} \rightarrow S}) = \begin{pmatrix} \int_{\partial S(\theta_1, \theta_2)} \partial_{\theta_1} \mathbf{X}(\theta_1, \theta_2, \mathbf{Y}(\theta_1, \theta_2, \gamma_x)) \cdot \mathbf{f}_{\mathcal{F} \rightarrow S}(\gamma_x) \, d\gamma_x \\ \int_{\partial S(\theta_1, \theta_2)} \partial_{\theta_2} \mathbf{X}(\theta_1, \theta_2, \mathbf{Y}(\theta_1, \theta_2, \gamma_x)) \cdot \mathbf{f}_{\mathcal{F} \rightarrow S}(\gamma_x) \, d\gamma_x \end{pmatrix} \in \mathbb{R}^2. \quad (1.19)$$

The matrix $\mathcal{M}_{\theta_1, \theta_2}$ in (1.16) is the Gram matrix of the family $(\partial_{\theta_1} \mathbf{X}(\theta_1, \theta_2), \partial_{\theta_2} \mathbf{X}(\theta_1, \theta_2))$ with respect to the scalar product $(\cdot, \cdot)_S$. It is invertible due to Assumption (1.9) (if two \mathcal{C}^∞ functions are colinear in $\mathbf{L}^2(S_{\text{ref}})$ then they are colinear in $\mathbf{L}^2(\partial S_{\text{ref}})$).

We also consider the following initial position and velocity for the structure

$$\begin{cases} \theta_1(0) = \theta_{1,0}, & \theta_2(0) = \theta_{2,0}, \\ \dot{\theta}_1(0) = \omega_{1,0}, & \dot{\theta}_2(0) = \omega_{2,0}. \end{cases} \quad (1.20)$$

1.1.3 Equations of the fluid

In our study, the global domain $\Omega = (0, L) \times (0, 1)$ represents a wind tunnel of length $L > 0$, see Fig. 3. Hence its boundary is composed of four regions: an inflow region $\Gamma_i = \{0\} \times (0, 1)$, a bottom region $\Gamma_b = (0, L) \times \{0\}$, a top region $\Gamma_t = (0, L) \times \{1\}$ and an outflow region $\Gamma_N = \{L\} \times (0, 1)$. We denote $\Gamma_w = \Gamma_t \cup \Gamma_b$ the part of the boundary corresponding to walls and $\Gamma_D = \Gamma_i \cup \Gamma_w$ the part of the boundary where Dirichlet conditions are imposed.

At time t , the structure occupies the volume $S(\theta_1(t), \theta_2(t))$, therefore the fluid fills the domain $\mathcal{F}(\theta_1(t), \theta_2(t)) = \Omega \setminus S(\theta_1(t), \theta_2(t))$.

The velocity of the fluid is modelled by the incompressible Navier–Stokes equations

$$\begin{cases} \frac{\partial \mathbf{u}}{\partial t}(t, \mathbf{x}) + (\mathbf{u} \cdot \nabla) \mathbf{u}(t, \mathbf{x}) - \text{div} \sigma_F(\mathbf{u}(t, \mathbf{x}), p(t, \mathbf{x})) = \mathbf{f}_{\mathcal{F}}(t, \mathbf{x}), & t \in (0, T), \quad \mathbf{x} \in \mathcal{F}(\theta_1(t), \theta_2(t)), \\ \text{div} \mathbf{u}(t, \mathbf{x}) = 0, & t \in (0, T), \quad \mathbf{x} \in \mathcal{F}(\theta_1(t), \theta_2(t)), \\ \mathbf{u}(t, \mathbf{x}) = \mathbf{u}^i(t, \mathbf{x}), & t \in (0, T), \quad \mathbf{x} \in \Gamma_i, \\ \mathbf{u}(t, \mathbf{x}) = 0, & t \in (0, T), \quad \mathbf{x} \in \Gamma_w, \\ \sigma_F(\mathbf{u}(t, \mathbf{x}), p(t, \mathbf{x})) \mathbf{n}(\mathbf{x}) = 0, & t \in (0, T), \quad \mathbf{x} \in \Gamma_N, \\ \mathbf{u}(0, \mathbf{x}) = \mathbf{u}_0(\mathbf{x}), & \mathbf{x} \in \mathcal{F}(\theta_{1,0}, \theta_{2,0}), \end{cases} \quad (1.21)$$

where $\mathbf{u}(t, \mathbf{x})$ and $p(t, \mathbf{x})$ are velocity and pressure of the fluid at point \mathbf{x} and time t , and

$$\sigma_F(\mathbf{u}, p) = \nu(\nabla \mathbf{u} + (\nabla \mathbf{u})^T) - pI,$$

is the stress tensor of the fluid, where $\nu > 0$ is the kinematic viscosity of the fluid. The vector \mathbf{n} denotes the unit outward normal to Ω . The term $\mathbf{f}_{\mathcal{F}}(t, \mathbf{x})$ in (1.21)₁ is a force per unit mass exerted on the fluid. Moreover, a nonhomogeneous Dirichlet boundary condition with datum \mathbf{u}^i is imposed on the inflow region Γ_i and we consider an initial datum \mathbf{u}_0 for the fluid velocity. Of course, these equations should be completed with suitable boundary conditions on $\partial S(\theta_1(t), \theta_2(t))$ that are made precise in Section 1.1.4.

1.1.4 Interface conditions between the fluid and the structure

The velocity \mathbf{u} of the fluid fulfils an adherence condition with the boundary of the structure whose velocity is given in (1.13),

$$\mathbf{u}(t, \mathbf{X}(\theta_1(t), \theta_2(t), \mathbf{y})) = \sum_{j=1}^2 \dot{\theta}_j(t) \partial_{\theta_j} \mathbf{X}(\theta_1(t), \theta_2(t), \mathbf{y}), \quad t \in (0, T), \quad \mathbf{y} \in \partial S_{\text{ref}}.$$

Note that this no-slip boundary condition corresponds to the continuity of the velocity through the interface between the fluid and the structure and can also be rewritten as

$$\mathbf{u}(t, \mathbf{x}) = \sum_{j=1}^2 \dot{\theta}_j(t) \partial_{\theta_j} \mathbf{X}(\theta_1(t), \theta_2(t), \mathbf{Y}(\theta_1(t), \theta_2(t), \mathbf{x})), \quad t \in (0, T), \quad \mathbf{x} \in \partial S(\theta_1(t), \theta_2(t)). \quad (1.22)$$

The forces exerted by the fluid on the structure are given by the stress tensor of the fluid

$$\mathbf{f}_{\mathcal{F} \rightarrow S}(t, \mathbf{x}) = -\sigma_F(\mathbf{u}, p) \mathbf{n}_{\theta_1, \theta_2}(t, \mathbf{x}), \quad t \in (0, T), \quad \mathbf{x} \in \partial S(\theta_1(t), \theta_2(t)), \quad (1.23)$$

where $\mathbf{n}_{\theta_1, \theta_2}(\mathbf{x})$ is the outward unit normal to the fluid domain $\mathcal{F}(\theta_1(t), \theta_2(t))$ on $\partial S(\theta_1(t), \theta_2(t))$.

1.1.5 The complete set of equations

The full set of equations is given by (1.15), (1.20), (1.21), (1.22) and (1.23). Note that the coupling between the fluid and the structure appears in equations (1.21) (the fluid domain depends on θ_1 and θ_2), (1.22) and (1.23).

The considered system is given by the following set of equations

$$\left\{ \begin{array}{ll} \frac{\partial \mathbf{u}}{\partial t}(t, \mathbf{x}) + (\mathbf{u}(t, \mathbf{x}) \cdot \nabla) \mathbf{u}(t, \mathbf{x}) - \text{div } \sigma_F(\mathbf{u}(t, \mathbf{x}), p(t, \mathbf{x})) = \mathbf{f}_{\mathcal{F}}(t, \mathbf{x}), & t \in (0, T), \quad \mathbf{x} \in \mathcal{F}(\theta_1(t), \theta_2(t)), \\ \text{div } \mathbf{u}(t, \mathbf{x}) = 0, & t \in (0, T), \quad \mathbf{x} \in \mathcal{F}(\theta_1(t), \theta_2(t)), \\ \mathbf{u}(t, \mathbf{x}) = \sum_{j=1}^2 \dot{\theta}_j(t) \partial_{\theta_j} \mathbf{X}(\theta_1(t), \theta_2(t), \mathbf{Y}(\theta_1(t), \theta_2(t), \mathbf{x})), & t \in (0, T), \quad \mathbf{x} \in \partial S(\theta_1(t), \theta_2(t)), \\ \mathbf{u}(t, \mathbf{x}) = \mathbf{u}^i(t, \mathbf{x}), & t \in (0, T), \quad \mathbf{x} \in \Gamma_i, \\ \mathbf{u}(t, \mathbf{x}) = 0, & t \in (0, T), \quad \mathbf{x} \in \Gamma_w, \\ \sigma_F(\mathbf{u}(t, \mathbf{x}), p(t, \mathbf{x})) \mathbf{n}(\mathbf{x}) = 0, & t \in (0, T), \quad \mathbf{x} \in \Gamma_N, \\ \mathbf{u}(0, \mathbf{x}) = \mathbf{u}_0(\mathbf{x}), & \mathbf{x} \in \mathcal{F}(\theta_{1,0}, \theta_{2,0}), \\ \mathcal{M}_{\theta_1, \theta_2} \begin{pmatrix} \ddot{\theta}_1 \\ \ddot{\theta}_2 \end{pmatrix} = \mathbf{M}_I(\theta_1, \theta_2, \dot{\theta}_1, \dot{\theta}_2) + \mathbf{M}_A(\theta_1, \theta_2, -\sigma_F(\mathbf{u}, p) \mathbf{n}_{\theta_1, \theta_2}) + \mathbf{f}_s, & t \in (0, T), \\ \theta_1(0) = \theta_{1,0}, \quad \theta_2(0) = \theta_{2,0}, \\ \dot{\theta}_1(0) = \omega_{1,0}, \quad \dot{\theta}_2(0) = \omega_{2,0}. \end{array} \right. \quad (1.24)$$

1.2 Statement of the main result

In this section, after setting up the functional framework, we present our existence result for solutions to problem (1.24). In the sequel, $\mathcal{F}_0 = \mathcal{F}(\theta_{1,0}, \theta_{2,0})$ denotes the initial fluid domain and $S_0 = S(\theta_{1,0}, \theta_{2,0})$ the initial configuration of the structure. For the sake of simplicity, the initial parameters of the structure are taken equal to zero,

$$\theta_{1,0} = \theta_{2,0} = 0.$$

This can be done without loss of generality by the change of variables

$$(\theta_1, \theta_2) \mapsto (\theta_1 - \theta_{1,0}, \theta_2 - \theta_{2,0}).$$

Moreover, the reference configuration for the structure S_{ref} and for the fluid \mathcal{F}_{ref} are taken as the initial configuration,

$$S_{\text{ref}} = S_0 = S(0, 0), \quad \mathcal{F}_{\text{ref}} = \mathcal{F}_0 = \mathcal{F}(0, 0).$$

1.2.1 Functional spaces

Sobolev spaces. In the sequel, $H^s(\mathcal{F}_0)$ is the usual Sobolev space of order $s \geq 0$. We identify $L^2(\mathcal{F}_0)$ with $H^0(\mathcal{F}_0)$. We will denote $\mathbf{L}^2(\mathcal{F}_0) = (L^2(\mathcal{F}_0))^2$, $\mathbf{H}^s(\mathcal{F}_0) = (H^s(\mathcal{F}_0))^2$ and so on.

Corners issues. The domain considered for the fluid has four corners of angle $\pi/2$. The ones that are located between Dirichlet and Neumann boundary conditions induce singularities, we denote them $A = (L, 1)$

and $B = (L, 0)$ (see Fig. 3). We also denote $\mathcal{J}_{d,n} = \{A, B\}$ the set of these corners and we define the distance of a point \mathbf{x} from these corners

$$\text{for } j \in \mathcal{J}_{d,n}, \quad \text{for } \mathbf{x} \in \Omega, \quad r_j(\mathbf{x}) = d(\mathbf{x}, j). \quad (1.25)$$

Note that corners between two Dirichlet boundary conditions do not induce singularities as soon as suitable compatibility conditions are satisfied. We report to [22] for more details.

Weighted Sobolev spaces. The strong solution to the Stokes problem in the domain with corners A and B and with a source term in $\mathbf{L}^2(\mathcal{F}_0)$ belongs to a classical Sobolev space of lower order than what we usually have with smooth domains. In order to get the usual gain of regularity between solutions and source terms, we have to study the solution in adapted Sobolev spaces. As the loss of regularity is located around corners A and B , we can recover the usual regularity if we consider norms that are suitably weighted near these corners. The weighted Sobolev spaces are then defined for $\beta > 0$ as

$$\begin{aligned} \mathbf{H}_\beta^2(\mathcal{F}_0) &= \{\mathbf{u} \text{ with } \|\mathbf{u}\|_{\mathbf{H}_\beta^2(\mathcal{F}_0)} < +\infty\}, \\ H_\beta^1(\mathcal{F}_0) &= \{p \text{ with } \|p\|_{H_\beta^1(\mathcal{F}_0)} < +\infty\}, \end{aligned}$$

where the norms $\|\cdot\|_{\mathbf{H}_\beta^2(\mathcal{F}_0)}$ and $\|\cdot\|_{H_\beta^1(\mathcal{F}_0)}$ are given by

$$\|\mathbf{u}\|_{\mathbf{H}_\beta^2(\mathcal{F}_0)}^2 = \sum_{|\alpha|=0}^2 \sum_{i=1}^2 \int_{\mathcal{F}_0} \left(\prod_{j \in \mathcal{J}_{d,n}} r_j^{2\beta}(\mathbf{y}) \right) |\partial^\alpha u_i(\mathbf{y})|^2 d\mathbf{y}, \quad (1.26)$$

and

$$\|p\|_{H_\beta^1(\mathcal{F}_0)}^2 = \sum_{|\alpha|=0}^1 \int_{\mathcal{F}_0} \left(\prod_{j \in \mathcal{J}_{d,n}} r_j^{2\beta}(\mathbf{y}) \right) |\partial^\alpha p(\mathbf{y})|^2 d\mathbf{y}. \quad (1.27)$$

Here the sum is on all multi-index α of length $|\alpha| \leq 2$ for (1.26) and $|\alpha| \leq 1$ for (1.27) and r_j is defined in (1.25).

Steady Stokes problem with corners. The following lemma from [24] explains how and why the spaces \mathbf{H}_β^2 and H_β^1 appear in the context of corners. It gives the result expected for the steady Stokes problem in \mathcal{F}_0 with weighted Sobolev spaces and the regularity obtained in the classical Sobolev spaces.

Lemma 1.4. [24, Theorem 2.5.] *Let us assume that $\mathbf{f}_\mathcal{F} \in \mathbf{L}^2(\mathcal{F}_0)$. The unique solution (\mathbf{u}, p) to the Stokes problem*

$$\begin{cases} -\operatorname{div} \sigma_F(\mathbf{u}, p) = \mathbf{f}_\mathcal{F} & \text{in } \mathcal{F}_0, \\ \operatorname{div} \mathbf{u} = 0 & \text{in } \mathcal{F}_0, \\ \mathbf{u} = 0 & \text{on } \Gamma_D \cup \partial S_0, \\ \sigma_F(\mathbf{u}, p)\mathbf{n} = 0 & \text{on } \Gamma_N, \end{cases} \quad (1.28)$$

belongs to $\mathbf{H}_\beta^2(\mathcal{F}_0) \times H_\beta^1(\mathcal{F}_0)$ for some $\beta \in (0, 1/2)$ and to $\mathbf{H}^{3/2+\varepsilon_0}(\mathcal{F}_0) \times H^{1/2+\varepsilon_0}(\mathcal{F}_0)$ for some $\varepsilon_0 \in (0, 1/2)$. Moreover, we have the following estimate

$$\|\mathbf{u}\|_{\mathbf{H}_\beta^2(\mathcal{F}_0) \cap \mathbf{H}^{3/2+\varepsilon_0}(\mathcal{F}_0)} + \|p\|_{H_\beta^1(\mathcal{F}_0) \cap H^{1/2+\varepsilon_0}(\mathcal{F}_0)} \leq C \|\mathbf{f}_\mathcal{F}\|_{\mathbf{L}^2(\mathcal{F}_0)}. \quad (1.29)$$

Let us denote \mathbf{n}_0 the outward unit normal to \mathcal{F}_0 on ∂S_0 . The regularity proven in Lemma 1.4 gives a meaning to all integrations by parts as $p|_{\partial \mathcal{F}_0}$ and $\partial_{\mathbf{n}_0} \mathbf{u}|_{\partial \mathcal{F}_0}$ are well defined traces for $(\mathbf{u}, p) \in \mathbf{H}^{3/2+\varepsilon_0}(\mathcal{F}_0) \times H^{1/2+\varepsilon_0}(\mathcal{F}_0)$.

Also note that according to [15, Theorem 1.4.3.1], there exists a continuous extension operator from $\mathbf{H}^s(\mathcal{F}_0)$ to $\mathbf{H}^s(\mathbb{R}^2)$ for every $s > 0$. This implies that all the classical Sobolev embeddings and interpolations are valid despite the presence of corners as they can be led in \mathbb{R}^2 .

Remark 1.5. Lemma 1.4 uses the geometry of the problem. Especially, we do not consider junctions between two segments where Neumann boundary conditions are imposed. All junctions between segments are right angles. To consider other angles, the reader can report to [22]. Note that the value of ε_0 and β depends on those angles.

1.2.2 Local existence of a strong solution to the problem

The diffeomorphism Φ . A classical difficulty in fluid–structure problems is that the fluid domain changes over time. The classical way to get rid of this difficulty is to use a change of variables on \mathbf{u} and p in order to bring the study back into a fixed domain. This procedure uses a diffeomorphism that we have to define properly. When the state of the structure depends only on a finite number of parameters, it is convenient to construct this diffeomorphism as an extension of the deformations of the structure into the fluid domain.

The diffeomorphism used is defined as an extension of the diffeomorphism \mathbf{X} given for the structure. Hence, we need the extension operator defined below.

Lemma 1.6. *There exists a linear extension operator $\mathcal{E} : \mathbf{W}^{3,\infty}(S_0) \rightarrow \mathbf{W}^{3,\infty}(\Omega) \cap \mathbf{H}_0^1(\Omega)$ such that for every $\mathbf{w} \in \mathbf{W}^{3,\infty}(S_0)$,*

- (i) $\mathcal{E}(\mathbf{w}) = \mathbf{w}$ in S_0 ,
- (ii) $\mathcal{E}(\mathbf{w})$ has support within $\Omega_\varepsilon = \{\mathbf{x} \in \Omega \mid d(\mathbf{x}, \partial\Omega) > \varepsilon\}$ for some $\varepsilon > 0$ such that $d(S(\theta_1, \theta_2), \partial\Omega) > 2\varepsilon$ for all $(\theta_1, \theta_2) \in \mathbb{D}_\Theta$,
- (iii) $\|\mathbf{w}\|_{\mathbf{W}^{3,\infty}(\Omega)} \leq C\|\mathbf{w}\|_{\mathbf{W}^{3,\infty}(S_0)}$, for some $C > 0$.

Proof. Extension results are classical, we can for instance find an extension result for smooth domains in [19, Lemma 12.2]. We can get the result by multiplying the extension function of [19, Lemma 12.2] by a cut-off function in $\mathcal{D}(\Omega_\varepsilon)$. Note that the existence of $\varepsilon > 0$ fulfilling (ii) is a consequence of Assumption (1.6). \square

Then we define the following function

$$\Phi(\theta_1, \theta_2, \mathbf{y}) = \mathbf{y} + \mathcal{E}\left(\mathbf{X}(\theta_1, \theta_2, \cdot) - \text{Id}\right)(\mathbf{y}), \quad \forall (\theta_1, \theta_2) \in \mathbb{D}_\Theta, \quad \forall \mathbf{y} \in \Omega, \quad (1.30)$$

where Id denotes the identity function.

We have $\nabla\Phi(0, 0, \mathbf{y}) = I$ for every $\mathbf{y} \in \Omega$, hence $\det(\nabla\Phi(0, 0, \mathbf{y})) = 1$. Then for every $(\theta_1, \theta_2) \in \mathbb{D}_\Theta$ small enough, the function $\Phi(\theta_1, \theta_2, \cdot)$ is a diffeomorphism close to the identity function. We denote $\Psi(\theta_1, \theta_2, \cdot)$ the inverse diffeomorphism of $\Phi(\theta_1, \theta_2, \cdot)$, i.e.

$$\forall (\theta_1, \theta_2) \in \mathbb{D}_\Theta, \quad \forall \mathbf{y} \in \Omega, \quad \Psi(\theta_1, \theta_2, \Phi(\theta_1, \theta_2, \mathbf{y})) = \mathbf{y}. \quad (1.31)$$

We can prove that Φ and Ψ belong to $\mathcal{C}^\infty(\mathbb{D}_\Theta, \mathbf{W}^{3,\infty}(\Omega))$. These diffeomorphisms are represented in Fig. 2.

The properties of \mathcal{E} imply that

$$\text{for every } (\theta_1, \theta_2) \in \mathbb{D}_\Theta, \quad \Phi(\theta_1, \theta_2, S_0) = S(\theta_1, \theta_2) \quad \text{and} \quad \forall \mathbf{y} \in \Omega \setminus \Omega_\varepsilon, \quad \Phi(\theta_1, \theta_2, \mathbf{y}) = \mathbf{y}. \quad (1.32)$$

The inflow boundary conditions. We use the following space to define the admissible boundary data on the inflow part of the boundary Γ_i ,

$$\mathbf{U}^i = \left\{ \mathbf{u}^i \in \mathbf{H}^{3/2}(\Gamma_i) \text{ with } \mathbf{u}^i|_{\partial\Gamma_i} = 0, \quad \begin{array}{l} \int_0^{1/4} \frac{|\partial_{y_2} u_2^i(y_2)|^2}{y_2} dy_2 < +\infty, \\ \int_{3/4}^1 \frac{|\partial_{y_2} u_2^i(y_2)|^2}{1-y_2} dy_2 < +\infty \end{array} \right\}. \quad (1.33)$$

The conditions with integrals in the definition of \mathbf{U}^i are chosen to match the homogeneous boundary conditions on Γ_w . We now state the following existence theorem.

Theorem 1.7 (Local existence in time of a solution). *Let $T_0 > 0$, let $\mathbf{u}^i \in H^1(0, T_0; \mathbf{U}^i)$, $\mathbf{u}_0 \in \mathbf{H}^1(\mathcal{F}_0)$ and $(\omega_{1,0}, \omega_{2,0}) \in \mathbb{R}^2$ satisfying the compatibility conditions*

$$\begin{cases} \text{div } \mathbf{u}_0 = 0 & \text{in } \mathcal{F}_0, \\ \mathbf{u}_0(\cdot) = \sum_{j=1}^2 \omega_{j,0} \partial_{\theta_j} \mathbf{X}(0, 0, \cdot) & \text{on } \partial S_0, \\ \mathbf{u}_0 = \mathbf{u}^i(0, \cdot) & \text{on } \Gamma_i, \\ \mathbf{u}_0 = 0 & \text{on } \Gamma_w. \end{cases} \quad (1.34)$$

Let $\mathbf{f}_{\mathcal{F}} \in L^2(0, T_0; \mathbf{W}^{1,\infty}(\Omega))$ and $\mathbf{f}_{\mathbf{s}} \in L^2(0, T_0; \mathbb{R}^2)$. Then there exists a time $T \in (0, T_0]$ such that problem (1.24) admits a unique solution $(\mathbf{u}, p, \theta_1, \theta_2)$ with the following regularity

$$\begin{aligned} & (\theta_1, \theta_2) \in H^2(0, T; \mathbb{D}_\Theta), \\ & \mathbf{u}(t, \Phi(\theta_1(t), \theta_2(t), \mathbf{y})) \in L^2(0, T; \mathbf{H}_\beta^2(\mathcal{F}_0)) \cap \mathcal{C}^0([0, T]; \mathbf{H}^1(\mathcal{F}_0)) \cap H^1(0, T; \mathbf{L}^2(\mathcal{F}_0)), \\ & p(t, \Phi(\theta_1(t), \theta_2(t), \mathbf{y})) \in L^2(0, T; H_\beta^1(\mathcal{F}_0)). \end{aligned}$$

Moreover, we have the estimate

$$\begin{aligned} & \|\mathbf{u}(t, \Phi(\theta_1(t), \theta_2(t), \mathbf{y}))\|_{L^2(0, T; \mathbf{H}_\beta^2(\mathcal{F}_0)) \cap \mathcal{C}^0([0, T]; \mathbf{H}^1(\mathcal{F}_0)) \cap H^1(0, T; \mathbf{L}^2(\mathcal{F}_0))} \\ & + \|p(t, \Phi(\theta_1(t), \theta_2(t), \mathbf{y}))\|_{L^2(0, T; H_\beta^1(\mathcal{F}_0))} + \|(\theta_1, \theta_2)\|_{H^2(0, T; \mathbb{D}_\Theta)} \\ & \leq C(\|\mathbf{u}_0\|_{\mathbf{H}^1(\mathcal{F}_0)} + |\omega_{1,0}| + |\omega_{2,0}| + \|\mathbf{f}_{\mathcal{F}}\|_{L^2(0, T_0; \mathbf{L}^2(\mathcal{F}_0))} + \|\mathbf{u}^i\|_{H^1(0, T_0; \mathbf{H}^{3/2}(\Gamma_i))} + \|\mathbf{f}_{\mathbf{s}}\|_{L^2(0, T_0; \mathbb{R}^2)}). \end{aligned}$$

The proof of Theorem 1.7 mainly follows the one in [9] and is presented in Section 3.3.

Remark 1.8. Note that we have the regularity $\mathbf{H}^{3/2+\varepsilon_0}$ for the fluid velocity because of the embedding $\mathbf{H}_\beta^2(\mathcal{F}_0) \hookrightarrow \mathbf{H}^{3/2+\varepsilon_0}(\mathcal{F}_0)$.

Remark 1.9. The study in dimension three would require to adapt the functional framework with more intricate weighted Sobolev spaces (see [22]). An adaptation of Lemmas 1.4, 2.10 and estimate (B.32) would also be required.

1.3 Scientific context

Fluid–structure interaction problems have been considered in several works. The structure is often rigid and immersed in an incompressible fluid [10, 14, 16, 17, 29, 31, 32] or a compressible one [6, 20, 21]. More complex structures have been studied for instance in [11, 27] where a plate immersed in an incompressible fluid has been considered or in [3, 18] where the authors studied the interaction between a 1D beam and a 2D fluid.

The interaction of an elastic structure with a compressible fluid has been studied in [8, 7] and with an incompressible fluid in [5].

Deforming structures that have a given deformation have been considered to model fish–like swimmers [23, 28]. However, they do not fit our framework since, in the present study, the deformations of the structure fulfill an ODE.

The case of a deformable structure depending on a finite numbers of degrees of freedom can be found in [9]. The model of the structure approximates the linear elasticity equations and it depends only on a finite number of degrees of freedom. However, to the best of our knowledge, besides the rigid solids, we have not found any work dealing with a structure depending naturally on a finite number of degrees of freedom fulfilling an equation. In that sense, the modelling proposed in the present article is original.

Moreover, additional difficulties are induced by the corners on $\partial\Omega$, more information can be found in [22, 24].

1.4 Outline of the paper

In Section 2, we study the linearized problem in the fixed domain \mathcal{F}_0 . We prove existence of strong solutions to this linearized problem. Then, in Section 3, we prove local existence of strong solutions to the nonlinear system. The computation of $\det(\mathcal{J}_{\mathbf{x}^b})$ is reported in Appendix A. The proof of the estimates of the nonlinear terms can be found in Appendix B.

2 Existence of solution to the linearized problem

In this section we study the linearization of problem (1.24) around the null state, first with only source terms \mathbf{f} and \mathbf{s} and then with all source terms and boundary data. These equations are written in the fixed domain \mathcal{F}_0 using a change of variables explained in Section 3.1. In the sequel, $(\tilde{\mathbf{u}}, \tilde{p})$ denotes the velocity and the pressure of the fluid in the fixed domain \mathcal{F}_0 . We denote $T > 0$ the considered final time.

2.1 Linearized problem with nonhomogeneous source terms

Let us study the following problem

$$\left\{ \begin{array}{ll} \frac{\partial \tilde{\mathbf{u}}}{\partial t} - \nu \Delta \tilde{\mathbf{u}} + \nabla \tilde{p} = \mathbf{f} & \text{in } (0, T) \times \mathcal{F}_0, \\ \operatorname{div} \tilde{\mathbf{u}} = 0 & \text{in } (0, T) \times \mathcal{F}_0, \\ \tilde{\mathbf{u}} = \dot{\theta}_1 \partial_{\theta_1} \Phi(0, 0, \cdot) + \dot{\theta}_2 \partial_{\theta_2} \Phi(0, 0, \cdot) & \text{on } (0, T) \times \partial S_0, \\ \tilde{\mathbf{u}} = 0 & \text{on } (0, T) \times \Gamma_i, \\ \tilde{\mathbf{u}} = 0 & \text{on } (0, T) \times \Gamma_w, \\ \sigma_F(\tilde{\mathbf{u}}, \tilde{p}) \mathbf{n} = 0 & \text{on } (0, T) \times \Gamma_N, \\ \tilde{\mathbf{u}}(0, \cdot) = \mathbf{u}_0(\cdot) & \text{in } \mathcal{F}_0, \end{array} \right. \quad (2.1)$$

$$\mathcal{M}_{0,0} \begin{pmatrix} \ddot{\theta}_1 \\ \ddot{\theta}_2 \end{pmatrix} = \begin{pmatrix} \int_{\partial S_0} [\tilde{p} I - \nu(\nabla \tilde{\mathbf{u}} + (\nabla \tilde{\mathbf{u}})^T)] \mathbf{n}_0 \cdot \partial_{\theta_1} \Phi(0, 0, \gamma_y) \, d\gamma_y \\ \int_{\partial S_0} [\tilde{p} I - \nu(\nabla \tilde{\mathbf{u}} + (\nabla \tilde{\mathbf{u}})^T)] \mathbf{n}_0 \cdot \partial_{\theta_2} \Phi(0, 0, \gamma_y) \, d\gamma_y \end{pmatrix} + \mathbf{s} \quad \text{on } (0, T),$$

$$\begin{array}{ll} \theta_1(0) = 0, & \theta_2(0) = 0, \\ \dot{\theta}_1(0) = \omega_{1,0}, & \dot{\theta}_2(0) = \omega_{2,0}, \end{array}$$

where the unknowns are $(\tilde{\mathbf{u}}, \tilde{p}, \theta_1, \theta_2)$ and the source terms are $(\mathbf{f}, \mathbf{s}) \in L^2(0, T; \mathbf{L}^2(\mathcal{F}_0)) \times L^2(0, T; \mathbb{R}^2)$. We will show later that this system corresponds to the linearization of the nonlinear problem (1.24) transported in the fixed initial configuration \mathcal{F}_0 .

Remark 2.1. The state $(\tilde{\mathbf{u}}, \tilde{p}, \theta_1, \theta_2)$ of problem (2.1) can be reduced to $(\tilde{\mathbf{u}}, \tilde{p}, \dot{\theta}_1, \dot{\theta}_2)$. Considering the velocity of the structure instead of its position is sufficient to solve (2.1). However, we prefer to consider the full state $(\tilde{\mathbf{u}}, \tilde{p}, \theta_1, \theta_2)$, as it is useful in the sequel to deal with the nonlinear problem.

In the sequel, the following spaces are used

$$\mathbf{U}_T = L^2(0, T; \mathbf{H}_\beta^2(\mathcal{F}_0)) \cap \mathcal{C}^0([0, T]; \mathbf{H}^1(\mathcal{F}_0)) \cap H^1(0, T; \mathbf{L}^2(\mathcal{F}_0)), \quad (2.2)$$

$$\mathbf{P}_T = L^2(0, T; H_\beta^1(\mathcal{F}_0)), \quad (2.3)$$

$$\Theta_T = H^2(0, T; \mathbb{R}^2), \quad (2.4)$$

$$\mathbf{F}_T = L^2(0, T; \mathbf{L}^2(\mathcal{F}_0)), \quad (2.5)$$

$$\mathbf{G}_T = H^1(0, T; \mathbf{H}^{3/2}(\partial S_0)), \quad (2.6)$$

$$\mathbf{S}_T = L^2(0, T; \mathbb{R}^2). \quad (2.7)$$

We endow Θ_T with the following norm

$$\|(\theta_1, \theta_2)\|_{\Theta_T} = \|(\theta_1, \theta_2)\|_{H^2(0, T)} + \|(\theta_1, \theta_2)\|_{L^\infty(0, T)} + \|(\dot{\theta}_1, \dot{\theta}_2)\|_{L^\infty(0, T)},$$

the other spaces are endowed with their natural norms. The norm $\|\cdot\|_{\Theta_T}$ has been chosen so that we have the estimate $\|(\theta_1, \theta_2)\|_{L^\infty(0, T)} + \|(\dot{\theta}_1, \dot{\theta}_2)\|_{L^\infty(0, T)} \leq C\|(\theta_1, \theta_2)\|_{\Theta_T}$ where C does not depend on T . Note that with the natural norm of Θ_T , C would depend on T .

Let us fix an arbitrary time $T_0 > 0$, e.g. $T_0 = 1$. We want to prove the following result.

Proposition 2.2. *There exists a constant $C > 0$ such that for all $T \in (0, T_0)$, C does not depend on T , for all $\mathbf{u}_0 \in \mathbf{H}^1(\mathcal{F}_0)$ and $(\omega_{1,0}, \omega_{2,0}) \in \mathbb{R}^2$ satisfying the compatibility conditions (1.34) (with $\mathbf{u}^i = 0$) and every $(\mathbf{f}, \mathbf{s}) \in \mathbf{F}_T \times \mathbf{S}_T$, problem (2.1) admits a unique solution*

$$(\tilde{\mathbf{u}}, \tilde{p}, \theta_1, \theta_2) \in \mathbf{U}_T \times \mathbf{P}_T \times \Theta_T.$$

Moreover, the following estimate holds

$$\|\tilde{\mathbf{u}}\|_{\mathbf{U}_T} + \|\tilde{p}\|_{\mathbf{P}_T} + \|(\theta_1, \theta_2)\|_{\Theta_T} \leq C(\|\mathbf{u}_0\|_{\mathbf{H}^1(\mathcal{F}_0)} + |\omega_{1,0}| + |\omega_{2,0}| + \|\mathbf{f}\|_{\mathbf{F}_T} + \|\mathbf{s}\|_{\mathbf{S}_T}). \quad (2.8)$$

In order to prove Proposition 2.2, we will study problem (2.1) under its semigroup formulation. Let us define the space

$$\mathbb{H} = \left\{ (\tilde{\mathbf{u}}, \theta_1, \theta_2, \omega_1, \omega_2) \in \mathbf{L}^2(\mathcal{F}_0) \times \mathbb{R}^4, \quad \begin{array}{l} \operatorname{div} \tilde{\mathbf{u}} = 0 \text{ in } \mathcal{F}_0, \quad \tilde{\mathbf{u}} \cdot \mathbf{n} = 0 \text{ on } \Gamma_D, \\ \tilde{\mathbf{u}} \cdot \mathbf{n}_0 = \sum_j \omega_j \partial_{\theta_j} \Phi(0, 0, \cdot) \cdot \mathbf{n}_0 \text{ on } \partial S_0 \end{array} \right\}, \quad (2.9)$$

where \mathbf{n}_0 is the unit outward normal to the fluid domain \mathcal{F}_0 . The space \mathbb{H} is endowed with the scalar product $(\cdot, \cdot)_0$ of $\mathbf{L}^2(\mathcal{F}_0) \times \mathbb{R}^4$ defined by

$$((\tilde{\mathbf{u}}^a, \theta_1^a, \theta_2^a, \omega_1^a, \omega_2^a), (\tilde{\mathbf{u}}^b, \theta_1^b, \theta_2^b, \omega_1^b, \omega_2^b))_0 = \int_{\mathcal{F}_0} \tilde{\mathbf{u}}^a \cdot \tilde{\mathbf{u}}^b \, dy + (\theta_1^a \ \theta_2^a) \begin{pmatrix} \theta_1^b \\ \theta_2^b \end{pmatrix} + (\omega_1^a \ \omega_2^a) \mathcal{M}_{0,0} \begin{pmatrix} \omega_1^b \\ \omega_2^b \end{pmatrix},$$

where $\mathcal{M}_{0,0}$ is the matrix defined in (1.16) with $\theta_1 = \theta_2 = 0$. We also define

$$\mathbb{V} = \left\{ (\tilde{\mathbf{u}}, \theta_1, \theta_2, \omega_1, \omega_2) \in \mathbf{H}^1(\mathcal{F}_0) \times \mathbb{R}^4, \quad \begin{array}{l} \operatorname{div} \tilde{\mathbf{u}} = 0 \text{ in } \mathcal{F}_0, \quad \tilde{\mathbf{u}} = 0 \text{ on } \Gamma_D, \\ \tilde{\mathbf{u}} = \sum_j \omega_j \partial_{\theta_j} \Phi(0, 0, \cdot) \text{ on } \partial S_0 \end{array} \right\}, \quad (2.10)$$

endowed with the scalar product $(\cdot, \cdot)_1$ of $\mathbf{H}^1(\mathcal{F}_0) \times \mathbb{R}^4$ defined by

$$((\tilde{\mathbf{u}}^a, \theta_1^a, \theta_2^a, \omega_1^a, \omega_2^a), (\tilde{\mathbf{u}}^b, \theta_1^b, \theta_2^b, \omega_1^b, \omega_2^b))_1 = \int_{\mathcal{F}_0} (\tilde{\mathbf{u}}^a \cdot \tilde{\mathbf{u}}^b + \nabla \tilde{\mathbf{u}}^a : \nabla \tilde{\mathbf{u}}^b) \, dy + (\theta_1^a \ \theta_2^a) \begin{pmatrix} \theta_1^b \\ \theta_2^b \end{pmatrix} + (\omega_1^a \ \omega_2^a) \mathcal{M}_{0,0} \begin{pmatrix} \omega_1^b \\ \omega_2^b \end{pmatrix}.$$

In the sequel, we denote $(f_j)_{j=1,2} = \begin{pmatrix} f_1 \\ f_2 \end{pmatrix}$.

Lemma 2.3. *The orthogonal space to \mathbb{H} with respect to the scalar product $(\cdot, \cdot)_0$ is*

$$(\mathbb{H})^\perp = \left\{ \left(\nabla p, 0, 0, -\mathcal{M}_{0,0}^{-1} \left(\int_{\partial S_0} p \mathbf{n}_0 \cdot \partial_{\theta_j} \Phi(0, 0, \gamma_y) \, d\gamma_y \right)_{j=1,2} \right) \text{ with } p \in H^1(\mathcal{F}_0), p = 0 \text{ on } \Gamma_N \right\}.$$

Proof. Let $(\tilde{\mathbf{u}}^a, \theta_1^a, \theta_2^a, \omega_1^a, \omega_2^a) \in \mathbf{L}^2(\mathcal{F}_0) \times \mathbb{R}^4$ such that for every $(\tilde{\mathbf{u}}^b, \theta_1^b, \theta_2^b, \omega_1^b, \omega_2^b) \in \mathbb{H}$,

$$((\tilde{\mathbf{u}}^a, \theta_1^a, \theta_2^a, \omega_1^a, \omega_2^a), (\tilde{\mathbf{u}}^b, \theta_1^b, \theta_2^b, \omega_1^b, \omega_2^b))_0 = 0.$$

By taking $\tilde{\mathbf{u}}^b = 0$ and $\omega_1^b = \omega_2^b = 0$, we easily obtain $\theta_1^a = \theta_2^a = 0$. With $\omega_1^b = \omega_2^b = 0$, we also get

$$\int_{\mathcal{F}_0} \tilde{\mathbf{u}}^a \cdot \tilde{\mathbf{u}}^b \, d\mathbf{y} = 0, \quad \forall \tilde{\mathbf{u}}^b \in \mathbf{L}^2(\mathcal{F}_0) \text{ such that } \operatorname{div} \tilde{\mathbf{u}}^b = 0 \text{ in } \mathcal{F}_0 \text{ and } \tilde{\mathbf{u}}^b \cdot \mathbf{n}_0 = 0 \text{ on } \Gamma_D \cup \partial S_0,$$

which implies, according to [24, Lemma 2.2], $\tilde{\mathbf{u}}^a = \nabla p$, where $p \in H^1(\mathcal{F}_0)$ and $p = 0$ on Γ_N . Now,

$$\int_{\mathcal{F}_0} \nabla p \cdot \tilde{\mathbf{u}}^b \, d\mathbf{y} + \sum_{j,k} \omega_j^a \omega_k^b (\partial_{\theta_j} \mathbf{X}(0, 0, \cdot), \partial_{\theta_k} \mathbf{X}(0, 0, \cdot))_S = 0,$$

becomes with the divergence formula and the compatibility condition in (2.9)

$$\sum_j \omega_j^b \int_{\partial S_0} p \mathbf{n}_0 \cdot \partial_{\theta_j} \Phi(0, 0, \gamma_y) \, d\gamma_y + \sum_{j,k} \omega_j^a \omega_k^b (\partial_{\theta_j} \mathbf{X}(0, 0, \cdot), \partial_{\theta_k} \mathbf{X}(0, 0, \cdot))_S = 0,$$

then

$$\int_{\partial S_0} p \mathbf{n}_0 \cdot \partial_{\theta_j} \Phi(0, 0, \gamma_y) \, d\gamma_y + \sum_k \omega_k^a (\partial_{\theta_j} \mathbf{X}(0, 0, \cdot), \partial_{\theta_k} \mathbf{X}(0, 0, \cdot))_S = 0,$$

which yields a first inclusion. The converse inclusion is obtained via an integration by parts. \square

We define the operator $(A, D(A))$ on \mathbb{H} as

$$D(A) = \left\{ (\tilde{\mathbf{u}}, \theta_1, \theta_2, \omega_1, \omega_2) \in \mathbb{V}, \tilde{\mathbf{u}} \in \mathbf{H}^{3/2+\varepsilon_0}(\mathcal{F}_0), \exists \tilde{p} \in H^{1/2+\varepsilon_0}(\mathcal{F}_0) \text{ such that } \operatorname{div} \sigma_F(\tilde{\mathbf{u}}, \tilde{p}) \in \mathbf{L}^2(\mathcal{F}_0) \text{ and } \sigma_F(\tilde{\mathbf{u}}, \tilde{p}) \mathbf{n} = 0 \text{ on } \Gamma_N \right\}, \quad (2.11)$$

where ε_0 is introduced in Lemma 1.4, and

$$A \begin{pmatrix} \tilde{\mathbf{u}} \\ \theta_1 \\ \theta_2 \\ \omega_1 \\ \omega_2 \end{pmatrix} = \Pi_{\mathbb{H}} \begin{pmatrix} \operatorname{div} \sigma_F(\tilde{\mathbf{u}}, \tilde{p}) \\ \omega_1 \\ \omega_2 \\ \mathcal{M}_{0,0}^{-1} \left(\int_{\partial S_0} -\sigma_F(\tilde{\mathbf{u}}, \tilde{p}) \mathbf{n}_0 \cdot \partial_{\theta_j} \Phi(0, 0, \gamma_y) \, d\gamma_y \right)_{j=1,2} \end{pmatrix}, \quad (2.12)$$

where $\Pi_{\mathbb{H}}$ is the orthogonal projector from $\mathbf{L}^2(\mathcal{F}_0) \times \mathbb{R}^4$ onto \mathbb{H} with respect to $(\cdot, \cdot)_0$.

Remark 2.4. The use of \tilde{p} in the definition of $(A, D(A))$ is useful to guarantee that $\operatorname{div} \sigma_F(\tilde{\mathbf{u}}, \tilde{p})$ belongs to $\mathbf{L}^2(\mathcal{F}_0)$ and then that the application of $\Pi_{\mathbb{H}}$ in the right hand-side of (2.12) makes sense.

Lemma 2.5. *The operator A is uniquely defined.*

Proof. Let $(\tilde{\mathbf{u}}, \theta_1, \theta_2, \omega_1, \omega_2) \in D(A)$ and consider two functions $p, q \in H^{1/2+\varepsilon_0}(\mathcal{F}_0)$ satisfying the conditions appearing into the definition of $D(A)$. Then, $\operatorname{div} \sigma_F(0, p - q) = -\nabla(p - q) \in \mathbf{L}^2(\mathcal{F}_0)$ implies $p - q \in H^1(\mathcal{F}_0)$, and $\sigma_F(0, p - q) \mathbf{n} = 0$ on Γ_N implies $p - q = 0$ on Γ_N .

Now,

$$\begin{aligned} & \begin{pmatrix} \operatorname{div} \sigma_F(\tilde{\mathbf{u}}, p) \\ \omega_1 \\ \omega_2 \\ \mathcal{M}_{0,0}^{-1} \left(\int_{\partial S_0} -\sigma_F(\tilde{\mathbf{u}}, p) \mathbf{n}_0 \cdot \partial_{\theta_j} \Phi(0, 0, \gamma_y) \, d\gamma_y \right)_{j=1,2} \end{pmatrix} - \begin{pmatrix} \operatorname{div} \sigma_F(\tilde{\mathbf{u}}, q) \\ \omega_1 \\ \omega_2 \\ \mathcal{M}_{0,0}^{-1} \left(\int_{\partial S_0} -\sigma_F(\tilde{\mathbf{u}}, q) \mathbf{n}_0 \cdot \partial_{\theta_j} \Phi(0, 0, \gamma_y) \, d\gamma_y \right)_{j=1,2} \end{pmatrix} \\ &= \begin{pmatrix} \nabla(p - q) \\ 0 \\ 0 \\ -\mathcal{M}_{0,0}^{-1} \left(\int_{\partial S_0} (p - q) \mathbf{n}_0 \cdot \partial_{\theta_j} \Phi(0, 0, \gamma_y) \, d\gamma_y \right)_{j=1,2} \end{pmatrix}, \end{aligned}$$

which belongs to \mathbb{H}^\perp according to Lemma 2.3. Therefore A is uniquely defined. \square

Before going further, let us point out that $D(A)$ can be characterized as follows.

Lemma 2.6. *We have*

$$D(A) = \left\{ (\tilde{\mathbf{u}}, \theta_1, \theta_2, \omega_1, \omega_2) \in \mathbb{V}, \tilde{\mathbf{u}} \in \mathbf{H}_\beta^2(\mathcal{F}_0), \exists \tilde{p} \in H_\beta^1(\mathcal{F}_0) \text{ such that } \operatorname{div} \sigma_F(\tilde{\mathbf{u}}, \tilde{p}) \in \mathbf{L}^2(\mathcal{F}_0) \text{ and } \sigma_F(\tilde{\mathbf{u}}, \tilde{p}) \mathbf{n} = 0 \text{ on } \Gamma_N \right\}.$$

Proof. Assume that $(\tilde{\mathbf{u}}, \theta_1, \theta_2, \omega_1, \omega_2)$ belongs to $D(A)$ given by (2.11). Then $(\tilde{\mathbf{u}}, \tilde{p})$ satisfies

$$\begin{cases} \operatorname{div} \sigma_F(\tilde{\mathbf{u}}, \tilde{p}) & \in \mathbf{L}^2(\mathcal{F}_0), \\ \operatorname{div} \tilde{\mathbf{u}} = 0 & \text{in } \mathcal{F}_0, \\ \tilde{\mathbf{u}} = \sum_j \omega_j \partial_{\theta_j} \Phi(0, 0, \cdot) & \text{on } \partial S_0, \\ \tilde{\mathbf{u}} = 0 & \text{on } \Gamma_D, \\ \sigma_F(\tilde{\mathbf{u}}, \tilde{p}) \mathbf{n} = 0 & \text{on } \Gamma_N. \end{cases}$$

According to [24, Theorem 2.16], there exists $\mathbf{v}_s \in \mathbf{H}^2(\mathcal{F}_0)$ such that

$$\begin{cases} \operatorname{div} \sigma_F(\mathbf{v}_s, 0) = 0 & \text{in } \mathcal{F}_0, \\ \operatorname{div} \mathbf{v}_s = 0 & \text{in } \mathcal{F}_0, \\ \mathbf{v}_s = \sum_j \omega_j \partial_{\theta_j} \Phi(0, 0, \cdot) & \text{on } \partial S_0, \\ \mathbf{v}_s = 0 & \text{on } \Gamma_D, \\ \sigma_F(\mathbf{v}_s, 0) \mathbf{n} = 0 & \text{on } \Gamma_N. \end{cases}$$

Let $\mathbf{f} = -\operatorname{div} \sigma_F(\tilde{\mathbf{u}}, \tilde{p}) \in \mathbf{L}^2(\mathcal{F}_0)$. Then $(\tilde{\mathbf{u}} - \mathbf{v}_s, \tilde{p})$ satisfies

$$\begin{cases} -\operatorname{div} \sigma_F(\tilde{\mathbf{u}} - \mathbf{v}_s, \tilde{p}) = \mathbf{f} & \text{in } \mathcal{F}_0, \\ \operatorname{div} (\tilde{\mathbf{u}} - \mathbf{v}_s) = 0 & \text{in } \mathcal{F}_0, \\ \tilde{\mathbf{u}} - \mathbf{v}_s = 0 & \text{on } \Gamma_D \cup \partial S_0, \\ \sigma_F(\tilde{\mathbf{u}} - \mathbf{v}_s, \tilde{p}) \mathbf{n} = 0 & \text{on } \Gamma_N. \end{cases}$$

According to Lemma 1.4, $\tilde{\mathbf{u}} - \mathbf{v}_s \in \mathbf{H}_{\beta}^2(\mathcal{F}_0) \cap \mathbf{H}^{3/2+\varepsilon_0}(\mathcal{F}_0)$, $\tilde{p} \in H_{\beta}^1(\mathcal{F}_0) \cap \mathbf{H}^{1/2+\varepsilon_0}(\mathcal{F}_0)$. This ends the proof. \square

We define the bilinear form a_1 on $\mathbb{V} \times \mathbb{V}$ for every $(\tilde{\mathbf{u}}^a, \theta_1^a, \theta_2^a, \omega_1^a, \omega_2^a)$ and $(\tilde{\mathbf{u}}^b, \theta_1^b, \theta_2^b, \omega_1^b, \omega_2^b)$ in \mathbb{V} by

$$a_1((\tilde{\mathbf{u}}^a, \theta_1^a, \theta_2^a, \omega_1^a, \omega_2^a), (\tilde{\mathbf{u}}^b, \theta_1^b, \theta_2^b, \omega_1^b, \omega_2^b)) = \frac{\nu}{2} \int_{\mathcal{F}_0} (\nabla \tilde{\mathbf{u}}^a + (\nabla \tilde{\mathbf{u}}^a)^T) : (\nabla \tilde{\mathbf{u}}^b + (\nabla \tilde{\mathbf{u}}^b)^T) \, dy.$$

We define the operator $(A_1, D(A))$ on \mathbb{H} by

$$D(A_1) = \{ \mathbf{z} \in \mathbb{V} \text{ with } \tilde{\mathbf{z}} \mapsto a_1(\mathbf{z}, \tilde{\mathbf{z}}) \text{ is } \mathbb{H} \text{-continuous} \},$$

and

$$\forall \mathbf{z} \in D(A_1), \quad \forall \tilde{\mathbf{z}} \in \mathbb{V}, \quad (A_1 \mathbf{z}, \tilde{\mathbf{z}})_0 = -a_1(\mathbf{z}, \tilde{\mathbf{z}}).$$

Lemma 2.7. *We have*

$$D(A_1) = D(A),$$

and

$$A_1 \begin{pmatrix} \tilde{\mathbf{u}} \\ \theta_1 \\ \theta_2 \\ \omega_1 \\ \omega_2 \end{pmatrix} = \Pi_{\mathbb{H}} \begin{pmatrix} \operatorname{div} \sigma_F(\tilde{\mathbf{u}}, \tilde{p}) \\ 0 \\ 0 \\ \mathcal{M}_{0,0}^{-1} \left(\int_{\partial S_0} -\sigma_F(\tilde{\mathbf{u}}, \tilde{p}) \mathbf{n}_0 \cdot \partial_{\theta_j} \Phi(0, 0, \gamma_y) \, d\gamma_y \right)_{j=1,2} \end{pmatrix}.$$

Proof. The inclusion $D(A) \subset D(A_1)$ comes easily. Moreover, for every $\mathbf{z} = (\tilde{\mathbf{u}}, \theta_1, \theta_2, \omega_1, \omega_2) \in D(A)$, an integration by parts yields

$$\forall \tilde{\mathbf{z}} \in \mathbb{V}, \quad (A_1 \mathbf{z}, \tilde{\mathbf{z}})_0 = \left(\begin{pmatrix} \operatorname{div} \sigma_F(\tilde{\mathbf{u}}, \tilde{p}) \\ 0 \\ 0 \\ \mathcal{M}_{0,0}^{-1} \left(\int_{\partial S_0} -\sigma_F(\tilde{\mathbf{u}}, \tilde{p}) \mathbf{n}_0 \cdot \partial_{\theta_j} \Phi(0, 0, \gamma_y) \, d\gamma_y \right)_{j=1,2} \end{pmatrix}, \tilde{\mathbf{z}} \right)_0.$$

Let us now prove the reverse inclusion $D(A_1) \subset D(A)$. Let $\mathbf{z} = (\tilde{\mathbf{u}}, \theta_1, \theta_2, \omega_1, \omega_2) \in D(A_1)$. According to Riesz representation theorem, there exists $\mathbf{f} \in \mathbb{H}$ such that

$$\forall \tilde{\mathbf{z}} \in \mathbb{V}, \quad a_1(\mathbf{z}, \tilde{\mathbf{z}}) = (\mathbf{f}, \tilde{\mathbf{z}})_0.$$

We write $\mathbf{f} = (\mathbf{f}_{\mathbf{u}}, \mathbf{f}_{\theta_1}, \mathbf{f}_{\theta_2}, \mathbf{f}_{\omega_1}, \mathbf{f}_{\omega_2})$. For $\tilde{\mathbf{v}} \in \mathcal{D}_{\operatorname{div}} = \{ \mathbf{u} \in (\mathcal{C}_c^\infty(\mathcal{F}_0))^2 \text{ with } \operatorname{div} \mathbf{u} = 0 \}$, we know that $(\tilde{\mathbf{v}}, 0, 0, 0, 0)$ belongs to \mathbb{V} , and an integration by parts yields

$$a_1(\mathbf{z}, (\tilde{\mathbf{v}}, 0, 0, 0, 0)) = \int_{\mathcal{F}_0} (-\operatorname{div} \sigma_F(\tilde{\mathbf{u}}, 0)) \cdot \tilde{\mathbf{v}} \, dy = \int_{\mathcal{F}_0} \mathbf{f}_{\mathbf{u}} \cdot \tilde{\mathbf{v}} \, dy.$$

Then, according to [30, Lemma 2.2.2], there exists $\widehat{q} \in L^2(\mathcal{F}_0)$ such that

$$-\operatorname{div} \sigma_F(\tilde{\mathbf{u}}, \widehat{q}) = \mathbf{f}_u \text{ in } \mathcal{F}_0, \quad (2.13)$$

and thus $\operatorname{div} \sigma_F(\tilde{\mathbf{u}}, \widehat{q})$ belongs to $\mathbf{L}^2(\mathcal{F}_0)$, which gives a meaning to $\sigma_F(\tilde{\mathbf{u}}, \widehat{q})\mathbf{n}_0$ on $\partial\mathcal{F}_0$.

Now, let us prove that $\sigma_F(\tilde{\mathbf{u}}, \widehat{q})\mathbf{n}$ is constant along Γ_N . Let $\mathbf{g} \in (\mathcal{C}_c^\infty(\Gamma_N))^2$ fulfilling $\int_{\Gamma_N} \mathbf{g} \cdot \mathbf{n} \, d\gamma_y = 0$. According to [13, Theorem IV.1.1], there exists $\mathbf{v}_g \in \mathbf{H}^1(\mathcal{F}_0)$ satisfying

$$\begin{cases} \operatorname{div} \mathbf{v}_g = 0 \text{ in } \mathcal{F}_0, \\ \mathbf{v}_g = 0 \text{ on } \Gamma_D \cup \partial S_0, \\ \mathbf{v}_g = \mathbf{g} \text{ on } \Gamma_N. \end{cases}$$

We know that $(\mathbf{v}_g, 0, 0, 0, 0)$ belongs to \mathbb{V} . An integration by parts yields

$$a_1(\mathbf{z}, (\mathbf{v}_g, 0, 0, 0, 0)) = \int_{\mathcal{F}_0} (-\operatorname{div} \sigma_F(\tilde{\mathbf{u}}, \widehat{q})) \cdot \mathbf{v}_g \, dy + \int_{\Gamma_N} \sigma_F(\tilde{\mathbf{u}}, \widehat{q})\mathbf{n} \cdot \mathbf{g} \, d\gamma_y = \int_{\mathcal{F}_0} \mathbf{f}_u \cdot \mathbf{v}_g \, dy,$$

and with (2.13) we get

$$\int_{\Gamma_N} \sigma_F(\tilde{\mathbf{u}}, \widehat{q})\mathbf{n} \cdot \mathbf{g} \, d\gamma_y = 0.$$

The previous equality holds for every $\mathbf{g} \in (\mathcal{C}_c^\infty(\Gamma_N))^2$ fulfilling $\int_{\Gamma_N} \mathbf{g} \cdot \mathbf{n} \, d\gamma_y = 0$, then there exists a constant c such that $\sigma_F(\tilde{\mathbf{u}}, \widehat{q})\mathbf{n} = c \mathbf{n}$ on Γ_N .

Let $q = \widehat{q} - c \in L^2(\mathcal{F}_0)$, we have $\operatorname{div} \sigma_F(\tilde{\mathbf{u}}, q) = \operatorname{div} \sigma_F(\tilde{\mathbf{u}}, \widehat{q})$ and $\sigma_F(\tilde{\mathbf{u}}, q)\mathbf{n} = 0$ on Γ_N . Moreover, $(\tilde{\mathbf{u}}, q)$ satisfies

$$\begin{cases} \operatorname{div} \sigma_F(\tilde{\mathbf{u}}, q) & \in \mathbf{L}^2(\mathcal{F}_0), \\ \operatorname{div} \tilde{\mathbf{u}} = 0 & \text{in } \mathcal{F}_0, \\ \tilde{\mathbf{u}} = 0 & \text{on } \Gamma_D, \\ \tilde{\mathbf{u}} = \sum_j \omega_j \partial_{\theta_j} \Phi(0, 0, \cdot) & \text{on } \partial S_0, \\ \sigma_F(\tilde{\mathbf{u}}, q)\mathbf{n} = 0 & \text{on } \Gamma_N. \end{cases}$$

We finish this proof with a lifting of the boundary datum on ∂S_0 [24, Theorem 2.16] and Lemma 1.4. We get $D(A_1) \subset D(A)$, thus concluding the proof of Lemma 2.7. \square

The key point of this section is the following lemma.

Lemma 2.8. *The operator A generates an analytic semigroup on \mathbb{H} . Moreover, for $\lambda \in \mathbb{R}$ large enough, $\lambda I - A$ is positive and $D((\lambda I - A)^{1/2}) = \mathbb{V}$.*

Proof. We first prove the properties of Lemma 2.8 on the self-adjoint operator A_1 and then we extend it to A with a perturbation argument.

The bilinear form a_1 is symmetric and according to Korn's inequality [12, p. 110], there exists $c > 0$ such that

$$\forall (\tilde{\mathbf{u}}, \theta_1, \theta_2, \omega_1, \omega_2) \in \mathbb{V}, \quad a_1((\tilde{\mathbf{u}}, \theta_1, \theta_2, \omega_1, \omega_2), (\tilde{\mathbf{u}}, \theta_1, \theta_2, \omega_1, \omega_2)) + \frac{\nu}{2} \|\tilde{\mathbf{u}}\|_{\mathbf{L}^2(\mathcal{F}_0)}^2 \geq c \|\tilde{\mathbf{u}}\|_{\mathbf{H}^1(\mathcal{F}_0)}^2,$$

so that,

$$\forall (\tilde{\mathbf{u}}, \theta_1, \theta_2, \omega_1, \omega_2) \in \mathbb{V}, \quad a_1((\tilde{\mathbf{u}}, \theta_1, \theta_2, \omega_1, \omega_2), (\tilde{\mathbf{u}}, \theta_1, \theta_2, \omega_1, \omega_2)) + \max\left(\frac{\nu}{2}, c\right) \|(\tilde{\mathbf{u}}, \theta_1, \theta_2, \omega_1, \omega_2)\|_{\mathbb{H}}^2 \geq c \|(\tilde{\mathbf{u}}, \theta_1, \theta_2, \omega_1, \omega_2)\|_{\mathbb{V}}^2.$$

Then we can easily conclude that $D((-A_1)^{1/2}) = \mathbb{V}$. Moreover, according to [4, Theorem 2.12, p. 115], A_1 generates an analytic semigroup on \mathbb{H} .

Now, we use the fact that $A - A_1 \in \mathcal{L}(\mathbb{H})$, then according to [25, Corollary 2.2.], A generates an analytic semigroup on \mathbb{H} .

A consequence of the previous result is that there exists $\lambda > 0$ such that $\lambda I - A$ is positive. Moreover, $D(\lambda I - A) = D(A_1)$, then by interpolation, $D((\lambda I - A)^{1/2}) = D((-A_1)^{1/2}) = \mathbb{V}$. \square

We are now in position to prove Proposition 2.2.

Proof of Proposition 2.2. Let us denote $\mathbf{F} = \Pi_{\mathbb{H}}(\mathbf{f}, 0, 0, \mathcal{M}_{0,0}^{-1}\mathbf{s})$ and $\mathbf{z}_0 = (\mathbf{u}_0, 0, 0, \omega_{1,0}, \omega_{2,0})$. We have $\mathbf{F} \in L^2(0, T; \mathbb{H})$ and, according to the compatibility conditions (1.34) (with $\mathbf{u}^i = 0$), $\mathbf{z}_0 \in D(A^{1/2}) = \mathbb{V}$.

According to [4, Theorem 3.1, p. 143] and Lemma 2.8, the problem

$$\begin{cases} \mathbf{z}'(t) = A\mathbf{z}(t) + \mathbf{F}(t), & t \geq 0, \\ \mathbf{z}(0) = \mathbf{z}_0, \end{cases} \quad (2.14)$$

admits a unique solution $\mathbf{z} \in L^2(0, T; D(A)) \cap H^1(0, T; \mathbb{H})$ and there exists $C > 0$ such that

$$\|\mathbf{z}\|_{L^2(0, T; D(A)) \cap H^1(0, T; \mathbb{H})} \leq C(\|\mathbf{F}\|_{L^2(0, T; \mathbb{H})} + \|\mathbf{z}_0\|_{\mathbb{V}}). \quad (2.15)$$

With the Sobolev embedding

$$L^2(0, T; D(A)) \cap H^1(0, T; \mathbb{H}) \hookrightarrow \mathcal{C}^0([0, T]; \mathbb{V}),$$

we have

$$\|\mathbf{z}\|_{L^2(0, T; D(A)) \cap \mathcal{C}^0([0, T]; \mathbb{V}) \cap H^1(0, T; \mathbb{H})} \leq C(\|\mathbf{F}\|_{L^2(0, T; \mathbb{H})} + \|\mathbf{z}_0\|_{\mathbb{V}}). \quad (2.16)$$

Moreover, C can be taken independent from $T \in (0, T_0)$. To prove this statement, we consider

$$\forall t \in [0, T_0], \quad \tilde{\mathbf{F}}(t) = \begin{cases} \mathbf{F}(t) & \text{if } t \in [0, T], \\ 0 & \text{if } t \in]T, T_0]. \end{cases}$$

If $\tilde{\mathbf{z}}$ is the solution on $[0, T_0]$ of

$$\begin{cases} \tilde{\mathbf{z}}' = A\tilde{\mathbf{z}} + \tilde{\mathbf{F}}, \\ \tilde{\mathbf{z}}(0) = \mathbf{z}_0, \end{cases}$$

then for $t \leq T$, $\tilde{\mathbf{z}}(t) = \mathbf{z}(t)$. We have the inequality

$$\|\tilde{\mathbf{z}}\|_{L^2(0, T_0; D(A)) \cap \mathcal{C}^0([0, T_0]; \mathbb{V}) \cap H^1(0, T_0; \mathbb{H})} \leq C(\|\tilde{\mathbf{F}}\|_{L^2(0, T_0; \mathbb{H})} + \|\mathbf{z}_0\|_{\mathbb{V}}),$$

where C does not depend on T . Moreover,

$$\|\tilde{\mathbf{z}}\|_{L^2(0, T_0; D(A)) \cap \mathcal{C}^0([0, T_0]; \mathbb{V}) \cap H^1(0, T_0; \mathbb{H})} \leq \|\tilde{\mathbf{z}}\|_{L^2(0, T_0; D(A)) \cap \mathcal{C}^0([0, T_0]; \mathbb{V}) \cap H^1(0, T_0; \mathbb{H})},$$

and

$$\|\tilde{\mathbf{F}}\|_{L^2(0, T_0; \mathbb{H})} = \|\mathbf{F}\|_{L^2(0, T; \mathbb{H})}.$$

By combining these arguments, we get (2.16) with C independent from T .

Now, if we write $\mathbf{z} = (\tilde{\mathbf{u}}, \theta_1, \theta_2, \omega_1, \omega_2)$, problem (2.14) becomes

$$\frac{d}{dt} \begin{pmatrix} \tilde{\mathbf{u}} \\ \theta_1 \\ \theta_2 \\ \omega_1 \\ \omega_2 \end{pmatrix} = \Pi_{\mathbb{H}} \begin{pmatrix} \operatorname{div} \sigma_F(\tilde{\mathbf{u}}, p) + \mathbf{f} \\ \omega_1 \\ \omega_2 \\ \mathcal{M}_{0,0}^{-1} \left(\mathbf{s} + \left(\int_{\partial S_0} -\sigma_F(\tilde{\mathbf{u}}, p) \mathbf{n}_0 \cdot \partial_{\theta_j} \Phi(0, 0, \cdot) d\gamma_y \right)_{j=1..2} \right) \end{pmatrix},$$

where $p \in L^2(0, T; H_{\beta}^1(\mathcal{F}_0))$. Then, Lemma 2.3 implies that there exists $q \in L^2(0, T; H^1(\mathcal{F}_0))$ such that $(\tilde{\mathbf{u}}, p+q, \theta_1, \theta_2)$ satisfies the linear problem (2.1). Moreover, according to (2.16), we have $(\theta_1, \theta_2) \in H^2(0, T; \mathbb{R}^2)$, $\tilde{\mathbf{u}} \in H^1(0, T; \mathbf{L}^2(\mathcal{F}_0)) \cap \mathcal{C}^0([0, T]; \mathbf{H}^1(\mathcal{F}_0)) \cap L^2(0, T; \mathbf{H}_{\beta}^2(\mathcal{F}_0))$, $\tilde{p} = p + q \in L^2(0, T; H_{\beta}^1(\mathcal{F}_0))$ and

$$\begin{aligned} & \|\tilde{\mathbf{u}}\|_{L^2(0, T; \mathbf{H}_{\beta}^2(\mathcal{F}_0)) \cap \mathcal{C}^0([0, T]; \mathbf{H}^1(\mathcal{F}_0)) \cap H^1(0, T; \mathbf{L}^2(\mathcal{F}_0))} + \|\tilde{p}\|_{L^2(0, T; H_{\beta}^1(\mathcal{F}_0))} + \|(\theta_1, \theta_2)\|_{\Theta_T} \\ & \leq C(\|\mathbf{u}_0\|_{\mathbf{H}^1(\mathcal{F}_0)} + |\omega_{1,0}| + |\omega_{2,0}| + \|\mathbf{f}\|_{L^2(0, T; \mathbf{L}^2(\mathcal{F}_0))} + \|\mathbf{s}\|_{L^2(0, T; \mathbb{R}^2)}). \end{aligned}$$

This concludes the proof of Proposition 2.2. \square

2.2 Linearized problem with nonhomogeneous boundary data

Let us now consider two more nonhomogeneous data: one datum \mathbf{g} on the boundary of the structure ∂S_0 and one datum \mathbf{u}^i on the inflow boundary region Γ_i . Let $T_0 > 0$, we study

$$\left\{ \begin{array}{ll} \frac{\partial \tilde{\mathbf{u}}}{\partial t} - \nu \Delta \tilde{\mathbf{u}} + \nabla \tilde{p} = \mathbf{f} & \text{in } (0, T) \times \mathcal{F}_0, \\ \operatorname{div} \tilde{\mathbf{u}} = 0 & \text{in } (0, T) \times \mathcal{F}_0, \\ \tilde{\mathbf{u}} = \dot{\theta}_1 \partial_{\theta_1} \Phi(0, 0, \cdot) + \dot{\theta}_2 \partial_{\theta_2} \Phi(0, 0, \cdot) + \mathbf{g} & \text{on } (0, T) \times \partial S_0, \\ \tilde{\mathbf{u}} = \mathbf{u}^i & \text{on } (0, T) \times \Gamma_i, \\ \tilde{\mathbf{u}} = 0 & \text{on } (0, T) \times \Gamma_w, \\ \sigma_F(\tilde{\mathbf{u}}, \tilde{p}) \mathbf{n} = 0 & \text{on } (0, T) \times \Gamma_N, \\ \tilde{\mathbf{u}}(0, \mathbf{y}) = \mathbf{u}_0(\mathbf{y}) & \text{in } \mathcal{F}_0, \\ \mathcal{M}_{0,0} \begin{pmatrix} \ddot{\theta}_1 \\ \ddot{\theta}_2 \end{pmatrix} = \begin{pmatrix} \int_{\partial S_0} [\tilde{p} I - \nu(\nabla \tilde{\mathbf{u}} + (\nabla \tilde{\mathbf{u}})^T)] \mathbf{n}_0 \cdot \partial_{\theta_1} \Phi(0, 0, \gamma_y) d\gamma_y \\ \int_{\partial S_0} [\tilde{p} I - \nu(\nabla \tilde{\mathbf{u}} + (\nabla \tilde{\mathbf{u}})^T)] \mathbf{n}_0 \cdot \partial_{\theta_2} \Phi(0, 0, \gamma_y) d\gamma_y \end{pmatrix} + \mathbf{s} & \text{on } (0, T), \\ \theta_1(0) = 0, \quad \theta_2(0) = 0, \\ \dot{\theta}_1(0) = \omega_{1,0}, \quad \dot{\theta}_2(0) = \omega_{2,0}, \end{array} \right. \quad (2.17)$$

where the source terms and data are $\mathbf{f} \in L^2(0, T; \mathbf{L}^2(\mathcal{F}_0))$, $\mathbf{g} \in H^1(0, T; \mathbf{H}^{3/2}(\partial S_0))$, $\mathbf{u}^i \in H^1(0, T_0; \mathbf{U}^i)$ and $\mathbf{s} \in L^2(0, T; \mathbb{R}^2)$.

We prove in this section the following result.

Proposition 2.9. *There exists a constant $C > 0$ such that for all $T \in (0, T_0)$, C does not depend on T , for all $\mathbf{u}^i \in H^1(0, T_0; \mathbf{U}^i)$, $\mathbf{u}_0 \in \mathbf{H}^1(\mathcal{F}_0)$ and $(\omega_{1,0}, \omega_{2,0}) \in \mathbb{R}^2$ satisfying the compatibility conditions (1.34) and every $(\mathbf{f}, \mathbf{g}, \mathbf{s}) \in \mathbb{F}_T \times \mathbb{G}_T \times \mathbb{S}_T$ with $\mathbf{g}(0) = 0$, problem (2.17) admits a unique solution*

$$(\tilde{\mathbf{u}}, \tilde{p}, \theta_1, \theta_2) \in \mathbf{U}_T \times \mathbb{P}_T \times \Theta_T,$$

with

$$\|\tilde{\mathbf{u}}\|_{\mathbf{U}_T} + \|\tilde{p}\|_{\mathbb{P}_T} + \|(\theta_1, \theta_2)\|_{\Theta_T} \leq C(\|\mathbf{u}_0\|_{\mathbf{H}^1(\mathcal{F}_0)} + |\omega_{1,0}| + |\omega_{2,0}| + \|\mathbf{f}\|_{\mathbb{F}_T} + \|\mathbf{g}\|_{\mathbb{G}_T} + \|\mathbf{s}\|_{\mathbb{S}_T} + \|\mathbf{u}^i\|_{H^1(0, T_0; \mathbf{U}^i)}). \quad (2.18)$$

Proposition 2.9 is proven at the end of the section. The proof uses the following lifting result for the terms \mathbf{g} and \mathbf{u}^i .

Lemma 2.10. *For every $\mathbf{g} \in \mathbf{H}^{3/2}(\partial S_0)$ and every $\mathbf{u}^i \in \mathbf{U}^i$, there exists $\bar{\mathbf{u}} \in \mathbf{H}^2(\mathcal{F}_0)$ satisfying*

$$\left\{ \begin{array}{ll} \operatorname{div} \bar{\mathbf{u}} = 0 & \text{in } \mathcal{F}_0, \\ \bar{\mathbf{u}} = \mathbf{g} & \text{on } \partial S_0, \\ \bar{\mathbf{u}} = \mathbf{u}^i & \text{on } \Gamma_i, \\ \bar{\mathbf{u}} = 0 & \text{on } \Gamma_w, \\ (\nabla \bar{\mathbf{u}} + (\nabla \bar{\mathbf{u}})^T) \mathbf{n} = 0 & \text{on } \Gamma_N, \end{array} \right. \quad (2.19)$$

with

$$\|\bar{\mathbf{u}}\|_{\mathbf{H}^2(\mathcal{F}_0)} \leq C(\|\mathbf{u}^i\|_{\mathbf{U}^i} + \|\mathbf{g}\|_{\mathbf{H}^{3/2}(\partial S_0)}). \quad (2.20)$$

Note that despite the presence of corners, we recover the expected regularity of the lifting for smooth domains.

Remark 2.11. For the sake of readability, from this point onwards all terms $d\mathbf{y}$ and $d\gamma_y$ are omitted in the integrals.

Proof of Lemma 2.10. The lifting result has been established for $\mathbf{u}^i = 0$ on the inflow region in [24, Theorem 2.16]. We first lift the inflow boundary condition $\mathbf{u}^i \neq 0$ in Ω and then we use the aforementioned result.

Lifting of the inflow boundary condition. Let us look for a function \mathbf{v} defined on the entire domain Ω and satisfying

$$\left\{ \begin{array}{ll} \operatorname{div} \mathbf{v} = 0 & \text{in } \Omega, \\ \mathbf{v} = \mathbf{u}^i & \text{on } \Gamma_i, \\ \mathbf{v} = 0 & \text{on } \Gamma_w, \\ (\nabla \mathbf{v} + (\nabla \mathbf{v})^T) \mathbf{n} = 0 & \text{on } \Gamma_N. \end{array} \right. \quad (2.21)$$

As \mathbf{v} is divergence-free and Ω is simply connected, we look for it under the form $\mathbf{v} = \nabla^\perp \psi$, where ψ is a scalar-valued function. In the geometry considered, Γ_i , Γ_t , Γ_b and Γ_N are straight lines, hence $\partial_{\mathbf{n}}$ is written as $\pm \partial_{y_1}$ or $\pm \partial_{y_2}$ according on the considered part of the boundary.

We can prove that ψ has to satisfy the conditions

$$\begin{aligned} \partial_{y_2}\psi &= -u_1^i & \text{and } \partial_{y_1}\psi &= u_2^i & \text{on } \Gamma_i, \\ \partial_{y_1}\psi &= 0 & \text{and } \partial_{y_2}\psi &= 0 & \text{on } \Gamma_b, \\ \partial_{y_1}\psi &= 0 & \text{and } \partial_{y_2}\psi &= 0 & \text{on } \Gamma_t, \\ \partial_{y_1}\partial_{y_2}\psi &= 0 & \text{and } \partial_{y_1}^2\psi - \partial_{y_2}^2\psi &= 0 & \text{on } \Gamma_N. \end{aligned}$$

We choose to meet these conditions in the following way:

$$\begin{aligned} \psi(y_2) &= -\int_0^{y_2} u_1^i & \text{and } \partial_{y_1}\psi &= u_2^i & \text{on } \Gamma_i, \\ \psi &= 0 & \text{and } \partial_{y_2}\psi &= 0 & \text{on } \Gamma_b, \\ \psi &= -\int_{\Gamma_i} u_1^i & \text{and } \partial_{y_2}\psi &= 0 & \text{on } \Gamma_t, \\ \psi(y_2) &= -\eta(y_2) \int_{\Gamma_i} u_1^i, & \partial_{y_1}\psi &= 0 & \text{and } \partial_{y_1}^2\psi - \partial_{y_2}^2\psi &= -d_{y_2}^2\eta(y_2) \int_{\Gamma_i} u_1^i & \text{on } \Gamma_N, \end{aligned} \tag{2.22}$$

where η is a \mathcal{C}^∞ function on $[0, 1]$ satisfying

$$\forall y_2 \in [0, 1], \quad \eta(y_2) = \begin{cases} 0 & \text{if } y_2 \in [0, 1/4], \\ \in [0, 1] & \text{if } y_2 \in]1/4, 3/4[, \\ 1 & \text{if } y_2 \in [3/4, 1]. \end{cases} \tag{2.23}$$

The theorem [15, Theorem 1.6.1.5, p.69] with $m = 3$ and $d = 2$ gives the existence of $\psi \in H^3(\Omega)$ fulfilling (2.22) under the compatibility conditions:

$$\text{there exist } \alpha_1 \text{ and } \alpha_2 > 0 \text{ such that } \begin{cases} \int_0^{\alpha_1} \frac{|\partial_{y_2} u_2^i|^2}{y_2} < +\infty, \\ \int_{1-\alpha_2}^1 \frac{|\partial_{y_2} u_2^i|^2}{1-y_2} < +\infty. \end{cases} \tag{2.24}$$

These conditions are the ones in the definition of \mathbf{U}^i in (1.33) with $\alpha_1 = \alpha_2 = 1/4$. Moreover we have the estimate

$$\|\mathbf{v}\|_{\mathbf{H}^2(\Omega)} \leq c\|\psi\|_{H^3(\Omega)} \leq C\|\mathbf{u}^i\|_{\mathbf{H}^{3/2}(\Gamma_i)}. \tag{2.25}$$

The divergence-free field $\mathbf{v} = \nabla^\perp \psi \in \mathbf{H}^2(\Omega)$ satisfies (2.21).

Lifting of the structure velocity. Now, $\tilde{\mathbf{v}} = \bar{\mathbf{u}} - \mathbf{v}|_{\mathcal{F}_0}$ has to satisfy

$$\begin{cases} \operatorname{div} \tilde{\mathbf{v}} = 0 & \text{in } \mathcal{F}_0, \\ \tilde{\mathbf{v}} = \mathbf{g} - \mathbf{v} & \text{on } \partial S_0, \\ \tilde{\mathbf{v}} = 0 & \text{on } \Gamma_i, \\ \tilde{\mathbf{v}} = 0 & \text{on } \Gamma_w, \\ (\nabla \tilde{\mathbf{v}} + (\nabla \tilde{\mathbf{v}})^T) \mathbf{n} = 0 & \text{on } \Gamma_N. \end{cases}$$

According to [24, Theorem 2.16], such $\tilde{\mathbf{v}}$ exists in $\mathbf{H}^2(\mathcal{F}_0)$ as soon as $\mathbf{g} - \mathbf{v} \in \mathbf{H}^{3/2}(\partial S_0)$. Moreover, we have the estimate

$$\|\tilde{\mathbf{v}}\|_{\mathbf{H}^2(\mathcal{F}_0)} \leq C\|\mathbf{g} - \mathbf{v}\|_{\mathbf{H}^{3/2}(\partial S_0)} \leq C(\|\mathbf{g}\|_{\mathbf{H}^{3/2}(\partial S_0)} + \|\mathbf{v}\|_{\mathbf{H}^2(\Omega)}). \tag{2.26}$$

This yields the expected result since $\bar{\mathbf{u}} = \tilde{\mathbf{v}} + \mathbf{v}|_{\mathcal{F}_0}$, the estimate (2.20) comes from (2.25) and (2.26). \square

We can now prove Proposition 2.9 in the following way.

Proof of Proposition 2.9. Let $\mathbf{u}^i \in H^1(0, T_0; \mathbf{U}^i)$, $\mathbf{u}_0 \in \mathbf{H}^1(\mathcal{F}_0)$ and $(\omega_{1,0}, \omega_{2,0}) \in \mathbb{R}^2$ satisfying the compatibility conditions (1.34). Let $(\mathbf{f}, \mathbf{g}, \mathbf{s}) \in \mathbb{F}_T \times \mathbb{G}_T \times \mathbb{S}_T$ with $\mathbf{g}(0) = 0$.

Let $\bar{\mathbf{u}} \in H^1(0, T; \mathbf{H}^2(\mathcal{F}_0))$ be such that $\bar{\mathbf{u}}(t)$ is the solution to (2.19), it fulfils

$$\|\bar{\mathbf{u}}\|_{H^1(0, T; \mathbf{H}^2(\mathcal{F}_0))} \leq C(\|\mathbf{u}^i\|_{H^1(0, T_0; \mathbf{U}^i)} + \|\mathbf{g}\|_{H^1(0, T; \mathbf{H}^{3/2}(\partial S_0))}). \tag{2.27}$$

The lifting $\bar{\mathbf{u}}$ also belongs to $\mathcal{C}^0([0, T]; \mathbf{H}^2(\mathcal{F}_0))$, and as $\mathbf{g}(0) = 0$, we have

$$\|\bar{\mathbf{u}}\|_{\mathcal{C}^0([0, T]; \mathbf{H}^2(\mathcal{F}_0))} \leq C(\|\mathbf{u}^i\|_{H^1(0, T_0; \mathbf{U}^i)} + \|\mathbf{g}\|_{H^1(0, T; \mathbf{H}^{3/2}(\partial S_0))}), \tag{2.28}$$

where C does not depend on T .

Let $(\hat{\mathbf{u}}, \tilde{p}, \theta_1, \theta_2)$ be the solution to

$$\left\{ \begin{array}{ll} \frac{\partial \hat{\mathbf{u}}}{\partial t} - \nu \Delta \hat{\mathbf{u}} + \nabla \tilde{p} = \mathbf{f} - \frac{\partial \bar{\mathbf{u}}}{\partial t} + \nu \Delta \bar{\mathbf{u}} & \text{in } (0, T) \times \mathcal{F}_0, \\ \operatorname{div} \hat{\mathbf{u}} = 0 & \text{in } (0, T) \times \mathcal{F}_0, \\ \hat{\mathbf{u}} = \dot{\theta}_1 \partial_{\theta_1} \Phi(0, 0, \cdot) + \dot{\theta}_2 \partial_{\theta_2} \Phi(0, 0, \cdot) & \text{on } (0, T) \times \partial S_0, \\ \hat{\mathbf{u}} = 0 & \text{on } (0, T) \times \Gamma_i, \\ \hat{\mathbf{u}} = 0 & \text{on } (0, T) \times \Gamma_w, \\ \sigma_F(\hat{\mathbf{u}}, \tilde{p}) \mathbf{n} = 0 & \text{on } (0, T) \times \Gamma_N, \\ \hat{\mathbf{u}}(0, \cdot) = \mathbf{u}_0(\cdot) - \bar{\mathbf{u}}(0, \cdot) & \text{in } \mathcal{F}_0, \\ \mathcal{M}_{0,0} \begin{pmatrix} \ddot{\theta}_1 \\ \ddot{\theta}_2 \end{pmatrix} = \begin{pmatrix} \int_{\partial S_0} [\tilde{p} I - \nu(\nabla(\hat{\mathbf{u}} + \bar{\mathbf{u}}) + (\nabla(\hat{\mathbf{u}} + \bar{\mathbf{u}}))^T] \mathbf{n}_0 \cdot \partial_{\theta_1} \Phi(0, 0, \gamma_y) \\ \int_{\partial S_0} [\tilde{p} I - \nu(\nabla(\hat{\mathbf{u}} + \bar{\mathbf{u}}) + (\nabla(\hat{\mathbf{u}} + \bar{\mathbf{u}}))^T] \mathbf{n}_0 \cdot \partial_{\theta_2} \Phi(0, 0, \gamma_y) \end{pmatrix} + \mathbf{s} & \text{on } (0, T), \\ \theta_1(0) = 0, \quad \theta_2(0) = 0, \\ \dot{\theta}_1(0) = \omega_{1,0}, \quad \dot{\theta}_2(0) = \omega_{2,0}. \end{array} \right.$$

We have

$$\begin{aligned} \mathbf{f} - \frac{\partial \bar{\mathbf{u}}}{\partial t} + \nu \Delta \bar{\mathbf{u}} &\in L^2(0, T; \mathbf{L}^2(\mathcal{F}_0)), \\ \mathbf{u}_0(\cdot) - \bar{\mathbf{u}}(0, \cdot) &= 0 \text{ on } \Gamma_i, \\ s_j + \int_{\partial S_0} -\nu(\nabla \hat{\mathbf{u}} + (\nabla \hat{\mathbf{u}})^T) \mathbf{n}_0 \cdot \partial_{\theta_j} \Phi(0, 0, \gamma_y) &\in L^2(0, T). \end{aligned}$$

Then, according to Proposition 2.2, $(\hat{\mathbf{u}}, \tilde{p}, \theta_1, \theta_2) \in \mathbb{U}_T \times \mathbb{P}_T \times \Theta_T$ and we have (2.8) with $\hat{\mathbf{u}}$ instead of $\tilde{\mathbf{u}}$.

Now, we consider $\tilde{\mathbf{u}} = \hat{\mathbf{u}} + \bar{\mathbf{u}}$, then $(\tilde{\mathbf{u}}, \tilde{p}, \theta_1, \theta_2) \in \mathbb{U}_T \times \mathbb{P}_T \times \Theta_T$ and (2.18) is a consequence of (2.27)–(2.28) and (2.8). \square

Remark 2.12. Note that a larger space than $H^1(0, T_0; \mathbf{U}^i)$ could be considered for \mathbf{u}^i . Indeed, we use a lifting in space only, inducing the requirement $\mathbf{u}^i \in H^1(0, T_0; \mathbf{U}^i)$. Using a space-time lifting would be slightly more complicated (see [26]), but would allow a larger space for the inflow boundary datum \mathbf{u}^i .

3 Local existence of solution to the full problem

In this section, we study the nonlinear problem. We recall that $\theta_{1,0} = \theta_{2,0} = 0$. At first, we rewrite the equations (1.24) in the fixed domain \mathcal{F}_0 , then we prove existence of a solution to this problem.

3.1 The equations in a fixed domain

Our goal is to write the equations (1.24) in the fixed domain \mathcal{F}_0 . To do so, we use the diffeomorphism defined in (1.30). We denote \mathcal{J}_Φ its Jacobian matrix and $\operatorname{cof}(\mathcal{J}_\Phi)$ the cofactor matrix of \mathcal{J}_Φ . We use the change of variables

$$\forall t \in [0, T], \quad \forall \mathbf{y} \in \mathcal{F}_0, \quad \begin{cases} \tilde{\mathbf{u}}(t, \mathbf{y}) = \operatorname{cof}(\mathcal{J}_\Phi(\theta_1(t), \theta_2(t), \mathbf{y}))^T \mathbf{u}(t, \Phi(\theta_1(t), \theta_2(t), \mathbf{y})), \\ \tilde{p}(t, \mathbf{y}) = p(t, \Phi(\theta_1(t), \theta_2(t), \mathbf{y})). \end{cases}$$

This choice is motivated by the fact that, according to [9, Lemma 3.1], we get $\operatorname{div} \tilde{\mathbf{u}} = 0$.

In the sequel, v_i denotes the i^{th} component of the vector \mathbf{v} . We recall that $\Psi(\theta_1, \theta_2, \cdot)$ is the inverse diffeomorphism of $\Phi(\theta_1, \theta_2, \cdot)$. To compute the equations satisfied by $(\tilde{\mathbf{u}}, \tilde{p}, \theta_1, \theta_2)$, we use the following explicit formula:

$$\mathbf{u}(t, \mathbf{x}) = \operatorname{cof}(\mathcal{J}_\Psi(\theta_1(t), \theta_2(t), \mathbf{x}))^T \tilde{\mathbf{u}}(t, \Psi(\theta_1(t), \theta_2(t), \mathbf{x})),$$

we have

$$\begin{aligned} \partial_t \mathbf{u}(t, \mathbf{x}) &= \operatorname{cof} \left(\frac{d}{dt} \mathcal{J}_\Psi(\theta_1(t), \theta_2(t), \mathbf{x}) \right)^T \tilde{\mathbf{u}}(t, \Psi(\theta_1(t), \theta_2(t), \mathbf{x})) \\ &\quad + \operatorname{cof}(\mathcal{J}_\Psi(\theta_1(t), \theta_2(t), \mathbf{x}))^T \partial_t \tilde{\mathbf{u}}(t, \Psi(\theta_1(t), \theta_2(t), \mathbf{x})) \\ &\quad + \operatorname{cof}(\mathcal{J}_\Psi(\theta_1(t), \theta_2(t), \mathbf{x}))^T \nabla_{\mathbf{y}} \tilde{\mathbf{u}}(t, \Psi(\theta_1(t), \theta_2(t), \mathbf{x})) \frac{d}{dt} \Psi(\theta_1(t), \theta_2(t), \mathbf{x}), \\ \partial_{x_j} \mathbf{u}(t, \mathbf{x}) &= \operatorname{cof}(\partial_{x_j} \mathcal{J}_\Psi(\theta_1(t), \theta_2(t), \mathbf{x}))^T \tilde{\mathbf{u}}(t, \Psi(\theta_1(t), \theta_2(t), \mathbf{x})) \\ &\quad + \operatorname{cof}(\mathcal{J}_\Psi(\theta_1(t), \theta_2(t), \mathbf{x}))^T \nabla_{\mathbf{y}} \tilde{\mathbf{u}}(t, \Psi(\theta_1(t), \theta_2(t), \mathbf{x})) \partial_{x_j} \Psi(\theta_1(t), \theta_2(t), \mathbf{x}), \end{aligned}$$

and

$$\begin{aligned}
\partial_{x_j}^2 \mathbf{u}(t, \mathbf{x}) &= \text{cof}(\partial_{x_j}^2 \mathcal{J}_{\Psi}(\theta_1(t), \theta_2(t), \mathbf{x}))^T \tilde{\mathbf{u}}(t, \Psi(\theta_1(t), \theta_2(t), \mathbf{x})) \\
&\quad + 2\text{cof}(\partial_{x_j} \mathcal{J}_{\Psi}(\theta_1(t), \theta_2(t), \mathbf{x}))^T \nabla_{\mathbf{y}} \tilde{\mathbf{u}}(t, \Psi(\theta_1(t), \theta_2(t), \mathbf{x})) \partial_{x_j} \Psi(\theta_1(t), \theta_2(t), \mathbf{x}) \\
&\quad + \text{cof}(\mathcal{J}_{\Psi}(\theta_1(t), \theta_2(t), \mathbf{x}))^T \sum_k \partial_{y_k} \nabla_{\mathbf{y}} \tilde{\mathbf{u}}(t, \Psi(\theta_1(t), \theta_2(t), \mathbf{x})) \partial_{x_j} \Psi(\theta_1(t), \theta_2(t), \mathbf{x}) \partial_{x_j} \Psi_k(\theta_1(t), \theta_2(t), \mathbf{x}) \\
&\quad + \text{cof}(\mathcal{J}_{\Psi}(\theta_1(t), \theta_2(t), \mathbf{x}))^T \nabla_{\mathbf{y}} \tilde{\mathbf{u}}(t, \Psi(\theta_1(t), \theta_2(t), \mathbf{x})) \partial_{x_j}^2 \Psi(\theta_1(t), \theta_2(t), \mathbf{x}).
\end{aligned}$$

Problem (1.24) in the fixed domain reads (2.17) where $\mathbf{f}, \mathbf{g}, \mathbf{s}$ are defined by

$$\begin{cases} \mathbf{f} = \mathbf{F}(\theta_1, \theta_2, \tilde{\mathbf{u}}, \tilde{p}) + \mathbf{f}_{\mathcal{F}}(t, \Phi(\theta_1(t), \theta_2(t), \mathbf{y})), \\ \mathbf{g} = \mathbf{G}(\theta_1, \theta_2, \dot{\theta}_1, \dot{\theta}_2), \\ \mathbf{s} = \mathbf{S}(\theta_1, \theta_2, \tilde{\mathbf{u}}, \tilde{p}) + \mathbf{f}_{\mathbf{s}}, \end{cases} \quad (3.1)$$

where \mathbf{F}, \mathbf{G} and \mathbf{S} are nonlinear terms. We use the decomposition $\mathbf{F}(\theta_1, \theta_2, \tilde{\mathbf{u}}, \tilde{p}) = \mathbf{F}^1 + \mathbf{F}^2 + \mathbf{F}^3 + \mathbf{F}^4 + \mathbf{F}^5$. We write $\Phi(\theta_1, \theta_2, \cdot)$ under the simpler notation Φ . The nonlinear terms are given as follows:

$$\begin{aligned}
\mathbf{F}^1(\theta_1, \theta_2, \tilde{\mathbf{u}}) &= (I - \text{cof}(\mathcal{J}_{\Psi}(\theta_1, \theta_2, \Phi))^T) \frac{\partial \tilde{\mathbf{u}}}{\partial t_T}, \\
\mathbf{F}^2(\theta_1, \theta_2, \tilde{\mathbf{u}}) &= -\text{cof}((\partial_t \mathcal{J}_{\Psi}(\theta_1, \theta_2, \cdot)) \circ \Phi) \tilde{\mathbf{u}}(t, \mathbf{y}) - \text{cof}(\mathcal{J}_{\Psi}(\theta_1, \theta_2, \Phi))^T (\nabla_{\mathbf{y}} \tilde{\mathbf{u}}) ((\partial_t \Psi(\theta_1, \theta_2, \cdot)) \circ \Phi), \\
\mathbf{F}^3(\theta_1, \theta_2, \tilde{\mathbf{u}})_i &= \nu \sum_{j,k,\ell,m} \text{cof}(\mathcal{J}_{\Psi}(\theta_1, \theta_2, \Phi))_{ki} \frac{\partial^2 \tilde{u}_k}{\partial y_{\ell} \partial y_m} \frac{\partial \Psi_{\ell}}{\partial x_j}(\theta_1, \theta_2, \Phi) \frac{\partial \Psi_m}{\partial x_j}(\theta_1, \theta_2, \Phi) \\
&\quad + 2\nu \sum_{j,k,\ell} \text{cof}(\partial_{x_j} \mathcal{J}_{\Psi}(\theta_1, \theta_2, \Phi))_{ki} \frac{\partial \tilde{u}_k}{\partial y_{\ell}} \frac{\partial \Psi_{\ell}}{\partial x_j}(\theta_1, \theta_2, \Phi) \\
&\quad + \nu \sum_{j,k,\ell} \text{cof}(\mathcal{J}_{\Psi}(\theta_1, \theta_2, \Phi))_{ki} \frac{\partial \tilde{u}_k}{\partial y_{\ell}} \frac{\partial^2 \Psi_{\ell}}{\partial x_j^2}(\theta_1, \theta_2, \Phi) \\
&\quad + \nu \sum_{j,k} \text{cof}(\partial_{x_j}^2 \mathcal{J}_{\Psi}(\theta_1, \theta_2, \Phi))_{ki} \tilde{u}_k - \nu \Delta_{\mathbf{y}} \tilde{u}_i(t, \mathbf{y}), \\
\mathbf{F}^4(\theta_1, \theta_2, \tilde{\mathbf{u}})_i &= - \sum_{j,k,r} \text{cof}(\mathcal{J}_{\Psi}(\theta_1, \theta_2, \Phi))_{kj} \text{cof}(\partial_{x_j} \mathcal{J}_{\Psi}(\theta_1, \theta_2))_{ri} \tilde{u}_k \tilde{u}_r \\
&\quad - \sum_{k,r} \det(\mathcal{J}_{\Psi}(\theta_1, \theta_2, \Phi))^2 \frac{\partial \Phi_i}{\partial y_r} \frac{\partial \tilde{u}_r}{\partial y_k} \tilde{u}_k, \\
\mathbf{F}^5(\theta_1, \theta_2, \tilde{p}) &= (I - \mathcal{J}_{\Psi}(\theta_1, \theta_2, \Phi))^T \nabla_{\mathbf{y}} \tilde{p}, \\
\mathbf{G}(\theta_1, \theta_2, \omega_1, \omega_2) &= \sum_{j=1}^2 \omega_j \left(\text{cof}(\mathcal{J}_{\Phi}(\theta_1, \theta_2, \mathbf{y}))^T \partial_{\theta_j} \Phi(\theta_1, \theta_2, \mathbf{y}) - \partial_{\theta_j} \Phi(0, 0, \mathbf{y}) \right), \\
\mathbf{S}(\theta_1, \theta_2, \tilde{\mathbf{u}}, \tilde{p}) &= -(\mathcal{M}_{\theta_1, \theta_2} - \mathcal{M}_{0,0}) \begin{pmatrix} \ddot{\theta}_1 \\ \ddot{\theta}_2 \end{pmatrix} + \mathbf{M}_{\mathbf{I}}(\theta_1, \theta_2, \dot{\theta}_1, \dot{\theta}_2) \\
&\quad + \left(\int_{\partial S_0} |\mathcal{J}_{\Phi} \mathbf{t}_0| [\tilde{p} I - \nu(\mathcal{G}(\theta_1, \theta_2, \tilde{\mathbf{u}}) + \mathcal{G}(\theta_1, \theta_2, \tilde{\mathbf{u}})^T)] \mathbf{n}_{\theta_1, \theta_2}(\Phi) \cdot \partial_{\theta_1} \Phi(\theta_1, \theta_2, \gamma_y) \right) \\
&\quad - \left(\int_{\partial S_0} |\mathcal{J}_{\Phi} \mathbf{t}_0| [\tilde{p} I - \nu(\mathcal{G}(\theta_1, \theta_2, \tilde{\mathbf{u}}) + \mathcal{G}(\theta_1, \theta_2, \tilde{\mathbf{u}})^T)] \mathbf{n}_{\theta_1, \theta_2}(\Phi) \cdot \partial_{\theta_2} \Phi(\theta_1, \theta_2, \gamma_y) \right) \\
&\quad - \left(\int_{\partial S_0} [\tilde{p} I - \nu(\nabla \tilde{\mathbf{u}} + (\nabla \tilde{\mathbf{u}})^T)] \mathbf{n}_0 \cdot \partial_{\theta_1} \Phi(0, 0, \gamma_y) \right) \\
&\quad - \left(\int_{\partial S_0} [\tilde{p} I - \nu(\nabla \tilde{\mathbf{u}} + (\nabla \tilde{\mathbf{u}})^T)] \mathbf{n}_0 \cdot \partial_{\theta_2} \Phi(0, 0, \gamma_y) \right),
\end{aligned} \quad (3.2)$$

where \mathbf{t}_0 is a unitary tangent vector to ∂S_0 , $\mathbf{M}_{\mathbf{I}}$ and $\mathcal{M}_{\theta_1, \theta_2}$ are defined in (1.16), (1.17) and

$$\mathcal{G}(\theta_1, \theta_2, \tilde{\mathbf{u}})_{ij} = \sum_k \text{cof}[\partial_{x_j} \mathcal{J}_{\Psi}(\theta_1, \theta_2, \cdot) \circ \Phi]_{ki} \tilde{u}_k + \sum_{k,\ell} \text{cof}(\mathcal{J}_{\Psi}(\theta_1, \theta_2, \Phi))_{ki} \frac{\partial \tilde{u}_k}{\partial y_{\ell}} \frac{\partial \Psi_{\ell}}{\partial x_j}(\theta_1, \theta_2, \Phi). \quad (3.3)$$

For every $T > 0$, we define the space

$$\mathbb{N}_T = \{(\tilde{\mathbf{u}}, \tilde{p}, \theta_1, \theta_2) \in \mathbb{U}_T \times \mathbb{P}_T \times \Theta_T \text{ with } \theta_1(0) = \theta_2(0) = 0, \text{ and } \forall t \in [0, T], (\theta_1, \theta_2)(t) \in \mathbb{D}_{\Theta}\},$$

that we endow with the norm

$$\|(\tilde{\mathbf{u}}, \tilde{p}, \theta_1, \theta_2)\|_{\mathbb{N}_T} = \|\tilde{\mathbf{u}}\|_{\mathbb{U}_T} + \|\tilde{p}\|_{\mathbb{P}_T} + \|(\theta_1, \theta_2)\|_{\Theta_T}. \quad (3.4)$$

We can state the following theorem.

Theorem 3.1. *Let $T_0 > 0$. Let $\mathbf{u}^i \in H^1(0, T_0; \mathbf{U}^i)$ and $(\mathbf{f}_{\mathcal{F}}, \mathbf{f}_{\mathbf{s}}) \in L^2(0, T_0; \mathbf{W}^{1,\infty}(\Omega)) \times L^2(0, T_0; \mathbb{R}^2)$. For every $(\mathbf{u}_0, \omega_{1,0}, \omega_{2,0}) \in \mathbf{H}^1(\mathcal{F}_0) \times \mathbb{R}^2$ satisfying the compatibility conditions (1.34), there exists $T \in (0, T_0]$ such that problem (2.17) where the source terms are given by (3.1) admits a unique solution $(\tilde{\mathbf{u}}, \tilde{p}, \theta_1, \theta_2) \in \mathbf{N}_T$ satisfying the following estimate*

$$\|\tilde{\mathbf{u}}, \tilde{p}, \theta_1, \theta_2\|_{\mathbf{N}_T} \leq C(\|\mathbf{u}_0\|_{\mathbf{H}^1(\mathcal{F}_0)} + |\omega_{1,0}| + |\omega_{2,0}| + \|\mathbf{f}_{\mathcal{F}}\|_{L^2(0, T_0; \mathbf{L}^2(\mathcal{F}_0))} + \|\mathbf{u}^i\|_{H^1(0, T_0; \mathbf{U}^i)} + \|\mathbf{f}_{\mathbf{s}}\|_{L^2(0, T_0)}),$$

where C does not depend on T , $\mathbf{f}_{\mathcal{F}}$, $\mathbf{f}_{\mathbf{s}}$ and \mathbf{u}^i .

This theorem is the rewriting of Theorem 1.7 in the fixed domain \mathcal{F}_0 . To prove Theorem 3.1, we use the results of Section 2 and a fixed point argument.

3.2 Proof of Theorem 3.1

Proof. We work in the fixed fluid domain \mathcal{F}_0 . Let $T_0 > 0$.

Let $\mathbf{u}^i \in H^1(0, T_0; \mathbf{U}^i)$ and $(\mathbf{u}_0, \omega_{1,0}, \omega_{2,0}) \in \mathbf{H}^1(\mathcal{F}_0) \times \mathbb{R}^2$ satisfying the compatibility conditions (1.34).

We define an application Λ^T on \mathbf{N}_T such that for every $(\bar{\mathbf{u}}, \bar{p}, \bar{\theta}_1, \bar{\theta}_2) \in \mathbf{N}_T$, $(\tilde{\mathbf{u}}, \tilde{p}, \theta_1, \theta_2) = \Lambda^T(\bar{\mathbf{u}}, \bar{p}, \bar{\theta}_1, \bar{\theta}_2) \in \mathbf{U}_T \times \mathbf{P}_T \times \Theta_T$ is the solution to problem (2.17), where the nonhomogeneous terms are given by

$$\begin{aligned} \mathbf{f} &= \mathbf{F}(\bar{\theta}_1, \bar{\theta}_2, \bar{\mathbf{u}}, \bar{p}) + \mathbf{f}_{\mathcal{F}}(t, \Phi(\bar{\theta}_1, \bar{\theta}_2, \mathbf{y})), \\ \mathbf{g} &= \mathbf{G}(\bar{\theta}_1, \bar{\theta}_2, \dot{\bar{\theta}}_1, \dot{\bar{\theta}}_2), \\ \mathbf{s} &= \mathbf{S}(\bar{\theta}_1, \bar{\theta}_2, \bar{\mathbf{u}}, \bar{p}) + \mathbf{f}_{\mathbf{s}}, \end{aligned}$$

where \mathbf{F} , \mathbf{G} and \mathbf{S} are given by (3.2). Note that Λ^T is well defined, indeed if $(\bar{\mathbf{u}}, \bar{p}, \bar{\theta}_1, \bar{\theta}_2) \in \mathbf{N}_T$, we have $\mathbf{G}(\bar{\theta}_1, \bar{\theta}_2, \dot{\bar{\theta}}_1, \dot{\bar{\theta}}_2)(t=0) = 0$ and we prove below that $(\mathbf{f}, \mathbf{g}, \mathbf{s})$ belongs to $\mathbf{F}_T \times \mathbf{G}_T \times \mathbf{S}_T$ (see Lemma 3.2 with $(\tilde{\mathbf{u}}^a, \tilde{p}^a, \theta_1^a, \theta_2^a) = (0, 0, 0, 0)$). Then according to Proposition 2.9, $\Lambda^T(\bar{\mathbf{u}}, \bar{p}, \bar{\theta}_1, \bar{\theta}_2)$ is uniquely defined. Note that Λ^T depends on the initial data $(\mathbf{u}_0, \omega_{1,0}, \omega_{2,0})$ and on the source term \mathbf{u}^i .

We take

$$R = 2C(\|\mathbf{u}^i\|_{H^1(0, T_0; \mathbf{U}^i)} + \|\mathbf{u}_0\|_{\mathbf{H}^1(\mathcal{F}_0)} + |\omega_{1,0}| + |\omega_{2,0}| + \|\mathbf{f}_{\mathcal{F}}\|_{L^2(0, T_0; \mathbf{L}^2(\mathcal{F}_0))} + \|\mathbf{f}_{\mathbf{s}}\|_{L^2(0, T_0)}),$$

where C is the constant of Proposition 2.9, so that Proposition 2.9 gives

$$\|\Lambda^T(0, 0, 0, 0)\|_{\mathbf{N}_T} \leq C(\|\mathbf{u}^i\|_{H^1(0, T_0; \mathbf{U}^i)} + \|\mathbf{u}_0\|_{\mathbf{H}^1(\mathcal{F}_0)} + |\omega_{1,0}| + |\omega_{2,0}| + \|\mathbf{f}_{\mathcal{F}}\|_{L^2(0, T_0; \mathbf{L}^2(\mathcal{F}_0))} + \|\mathbf{f}_{\mathbf{s}}\|_{L^2(0, T_0)}) = R/2. \quad (3.5)$$

The strategy adopted is based on the existence of $T > 0$ such that Λ^T is a contraction on

$$\mathbb{B}_R(T) = \{ (\tilde{\mathbf{u}}, \tilde{p}, \theta_1, \theta_2) \in \mathbf{N}_T \quad \text{with} \quad \|(\tilde{\mathbf{u}}, \tilde{p}, \theta_1, \theta_2)\|_{\mathbf{N}_T} \leq R \}. \quad (3.6)$$

The domain \mathbb{D}_{Θ} is an open subset of \mathbb{R}^2 and $(0, 0) \in \mathbb{D}_{\Theta}$, then there exists $r > 0$ such that $B((0, 0), r) \subset \mathbb{D}_{\Theta}$. Then for $T < r/R$, if $(\cdot, \cdot, \theta_1, \theta_2) \in \mathbb{B}_R(T)$, we have $\|\dot{\theta}_j\|_{L^\infty(0, T)} \leq R$ and $\theta_j(0) = 0$, then

$$\|\theta_j\|_{L^\infty(0, T)} \leq T\|\dot{\theta}_j\|_{L^\infty(0, T)} \leq RT \leq r,$$

and we have for all $t \in (0, T)$, $(\theta_1(t), \theta_2(t)) \in \mathbb{D}_{\Theta}$. In the sequel we choose $T_0 > 0$ such that $T_0 < r/R$. Hence $\forall T \in [0, T_0]$, $\Lambda^T : \mathbb{B}_R(T) \rightarrow \mathbf{N}_T$.

The solution to the nonlinear problem will be obtained as a fixed point of the application Λ^T on $\mathbb{B}_R(T)$. We use the estimates of the following lemma.

Lemma 3.2. *For every $R' > 0$, there exists a constant $C' = C'(R') > 0$, such that for every $T \in (0, T_0)$, and every $(\tilde{\mathbf{u}}^j, \tilde{p}^j, \theta_1^j, \theta_2^j) \in \mathbb{B}_{R'}(T)$, we have*

$$\|\mathbf{F}(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a, \tilde{p}^a) - \mathbf{F}(\theta_1^b, \theta_2^b, \tilde{\mathbf{u}}^b, \tilde{p}^b)\|_{\mathbf{F}_T} \leq C'T^{1/4}(\|\tilde{\mathbf{u}}^a - \tilde{\mathbf{u}}^b\|_{\mathbf{U}_T} + \|\tilde{p}^a - \tilde{p}^b\|_{\mathbf{P}_T} + \|\theta^a - \theta^b\|_{\Theta_T}), \quad (3.7)$$

$$\|\mathbf{G}(\theta_1^a, \theta_2^a, \dot{\theta}_1^a, \dot{\theta}_2^a) - \mathbf{G}(\theta_1^b, \theta_2^b, \dot{\theta}_1^b, \dot{\theta}_2^b)\|_{\mathbf{G}_T} \leq C'T^{1/2}\|\theta^a - \theta^b\|_{\Theta_T}, \quad (3.8)$$

$$\|\mathbf{S}(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a, \tilde{p}^a) - \mathbf{S}(\theta_1^b, \theta_2^b, \tilde{\mathbf{u}}^b, \tilde{p}^b)\|_{\mathbf{S}_T} \leq C'T^{1/2}(\|\tilde{\mathbf{u}}^a - \tilde{\mathbf{u}}^b\|_{\mathbf{U}_T} + \|\tilde{p}^a - \tilde{p}^b\|_{\mathbf{P}_T} + \|\theta^a - \theta^b\|_{\Theta_T}), \quad (3.9)$$

$$\|\mathbf{f}_{\mathcal{F}}(t, \Phi(\theta_1^a, \theta_2^a, \mathbf{y})) - \mathbf{f}_{\mathcal{F}}(t, \Phi(\theta_1^b, \theta_2^b, \mathbf{y}))\|_{\mathbf{F}_T} \leq C'T\|\theta^a - \theta^b\|_{\Theta_T}. \quad (3.10)$$

These estimates are proven in Appendix B.

For $(\tilde{\mathbf{u}}^j, \tilde{p}^j, \theta_1^j, \theta_2^j) \in \mathbb{B}_R(T)$, Proposition 2.9 yields the estimate

$$\begin{aligned} &\|\Lambda^T(\tilde{\mathbf{u}}^a, \tilde{p}^a, \theta_1^a, \theta_2^a) - \Lambda^T(\tilde{\mathbf{u}}^b, \tilde{p}^b, \theta_1^b, \theta_2^b)\|_{\mathbf{N}_T} \\ &\leq C(\|\mathbf{F}(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a, \tilde{p}^a) - \mathbf{F}(\theta_1^b, \theta_2^b, \tilde{\mathbf{u}}^b, \tilde{p}^b)\|_{\mathbf{F}_T} + \|\mathbf{G}(\theta_1^a, \theta_2^a, \dot{\theta}_1^a, \dot{\theta}_2^a) - \mathbf{G}(\theta_1^b, \theta_2^b, \dot{\theta}_1^b, \dot{\theta}_2^b)\|_{\mathbf{G}_T} \\ &\quad + \|\mathbf{S}(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a, \tilde{p}^a) - \mathbf{S}(\theta_1^b, \theta_2^b, \tilde{\mathbf{u}}^b, \tilde{p}^b)\|_{\mathbf{S}_T} + \|\mathbf{f}_{\mathcal{F}}(t, \Phi(\theta_1^a, \theta_2^a, \mathbf{y})) - \mathbf{f}_{\mathcal{F}}(t, \Phi(\theta_1^b, \theta_2^b, \mathbf{y}))\|_{\mathbf{F}_T}), \end{aligned} \quad (3.11)$$

and with Lemma 3.2, we have

$$\|\Lambda^T(\tilde{\mathbf{u}}^a, \tilde{p}^a, \theta_1^a, \theta_2^a) - \Lambda^T(\tilde{\mathbf{u}}^b, \tilde{p}^b, \theta_1^b, \theta_2^b)\|_{\mathbf{N}_T} \leq KT^{1/4}(\|\tilde{\mathbf{u}}^a - \tilde{\mathbf{u}}^b\|_{\mathbf{U}_T} + \|\tilde{p}^a - \tilde{p}^b\|_{\mathbf{P}_T} + \|\theta^a - \theta^b\|_{\Theta_T}), \quad (3.12)$$

where $K = 4CC'(R)$ depends on R but not on T . For $T \in (0, T_0)$ such that

$$T < r/R \quad \text{and} \quad KT^{1/4} \leq 1/2,$$

estimates (3.12) and (3.5) yield

$$\|\Lambda^T(\tilde{\mathbf{u}}, \tilde{p}, \theta_1, \theta_2)\|_{\mathbf{N}_T} \leq \|\Lambda^T(0, 0, 0, 0)\|_{\mathbf{N}_T} + KT^{1/4}(\|\tilde{\mathbf{u}}\|_{\mathbf{U}_T} + \|\tilde{p}\|_{\mathbf{P}_T} + \|\theta\|_{\Theta_T}) \leq R/2 + KRT^{1/4} \leq R.$$

Then, $\forall T \in [0, T]$, $\Lambda^T : \mathbb{B}_R(T) \rightarrow \mathbb{B}_R(T)$ and according to (3.12), the application Λ^T is a contraction on $\mathbb{B}_R(T)$.

We have proven that $\Lambda^T : \mathbb{B}_R(T) \rightarrow \mathbb{B}_R(T)$ is a contraction. Then, according to the Picard fixed point theorem, there exists a unique fixed point to Λ^T in $\mathbb{B}_R(T)$. This fixed point is the solution to problem (2.17) where the source terms are given by (3.1). This proves Theorem 3.1. \square

3.3 Proof of the result in the moving domain, Theorem 1.7

We consider $(\tilde{\mathbf{u}}, \tilde{p}, \theta_1, \theta_2)$ in \mathbf{N}_T the solution to problem (2.17) with (3.1) given by Theorem 3.1. Let $\mathbf{u}(t, \mathbf{x}) = \text{cof}(\mathcal{J}_{\Psi}(\theta_1(t), \theta_2(t), \mathbf{x}))^T \tilde{\mathbf{u}}(t, \Psi(\theta_1(t), \theta_2(t), \mathbf{x})))$ and $p(t, \mathbf{x}) = \tilde{p}(t, \Psi(\theta_1(t), \theta_2(t), \mathbf{x}))$. Then the quadruplet $(\mathbf{u}, p, \theta_1, \theta_2)$ is solution to the problem in the moving domain, i.e. problem (1.24). This proves Theorem 1.7.

Appendix

A Computation of $\det(\mathcal{J}_{\mathbf{X}^b})$

We denote the components of the jacobian matrix as follows

$$\mathcal{J}_{\mathbf{X}^b} = \begin{pmatrix} \partial_1 X_1 & \partial_2 X_1 \\ \partial_1 X_2 & \partial_2 X_2 \end{pmatrix}.$$

Hence, we have

$$\begin{aligned} \det(\mathcal{J}_{\mathbf{X}^b}) &= \partial_1 X_1 \times \partial_2 X_2 - \partial_2 X_1 \times \partial_1 X_2 \\ &= \left(g'_{\theta_1} \mathbf{e}_{r1} + g'_{\theta_2} \mathbf{e}_{r2} + y_2 \left(\frac{\mathbf{N}'}{\|\mathbf{N}\|} - \frac{(\|\mathbf{N}\|)' \mathbf{N}}{\|\mathbf{N}\|^2} \right) \right) \times \left(\frac{\mathbf{N}}{\|\mathbf{N}\|} \right)_2 \\ &\quad - \left(g'_{\theta_1} \mathbf{e}_{r1} + g'_{\theta_2} \mathbf{e}_{r2} + y_2 \left(\frac{\mathbf{N}'}{\|\mathbf{N}\|} - \frac{(\|\mathbf{N}\|)' \mathbf{N}}{\|\mathbf{N}\|^2} \right) \right)_1 \times \left(\frac{\mathbf{N}}{\|\mathbf{N}\|} \right)_1 \\ &= \left(g'_{\theta_1} \mathbf{e}_{r1} + g'_{\theta_2} \mathbf{e}_{r2} + y_2 \frac{\mathbf{N}'}{\|\mathbf{N}\|} \right)_1 \times \left(\frac{\mathbf{N}}{\|\mathbf{N}\|} \right)_2 - \left(g'_{\theta_1} \mathbf{e}_{r1} + g'_{\theta_2} \mathbf{e}_{r2} + y_2 \frac{\mathbf{N}'}{\|\mathbf{N}\|} \right)_2 \times \left(\frac{\mathbf{N}}{\|\mathbf{N}\|} \right)_1 \\ &= \left(g'_{\theta_1} \cos(\theta_1) + g'_{\theta_2} \cos(\theta_1 + \theta_2) + y_2 \frac{-g''_{\theta_1} \sin(\theta_1) - g''_{\theta_2} \sin(\theta_1 + \theta_2)}{\|\mathbf{N}\|} \right) \times \frac{g'_{\theta_1} \cos(\theta_1) + g'_{\theta_2} \cos(\theta_1 + \theta_2)}{\|\mathbf{N}\|} \\ &\quad - \left(g'_{\theta_1} \sin(\theta_1) + g'_{\theta_2} \sin(\theta_1 + \theta_2) + y_2 \frac{g''_{\theta_1} \cos(\theta_1) + g''_{\theta_2} \cos(\theta_1 + \theta_2)}{\|\mathbf{N}\|} \right) \times \frac{-g'_{\theta_1} \sin(\theta_1) - g'_{\theta_2} \sin(\theta_1 + \theta_2)}{\|\mathbf{N}\|} \\ &= \frac{1}{\|\mathbf{N}\|} \left((g'_{\theta_1})^2 + (g'_{\theta_2})^2 + 2g'_{\theta_1} g'_{\theta_2} (\sin(\theta_1) \sin(\theta_1 + \theta_2) + \cos(\theta_1) \cos(\theta_1 + \theta_2)) \right) \\ &\quad + \frac{y_2}{\|\mathbf{N}\|^2} \left(g''_{\theta_1} g'_{\theta_2} (\cos(\theta_1) \sin(\theta_1 + \theta_2) - \sin(\theta_1) \cos(\theta_1 + \theta_2)) \right. \\ &\quad \left. + g'_{\theta_1} g''_{\theta_2} (\cos(\theta_1 + \theta_2) \sin(\theta_1) - \cos(\theta_1) \sin(\theta_1 + \theta_2)) \right) \\ &= \frac{1}{\|\mathbf{N}\|} \left((g'_{\theta_1})^2 + (g'_{\theta_2})^2 + 2g'_{\theta_1} g'_{\theta_2} \cos(\theta_2) \right) + \frac{y_2}{\|\mathbf{N}\|^2} \left(g''_{\theta_1} g'_{\theta_2} \sin(\theta_2) + g''_{\theta_2} g'_{\theta_1} \sin(-\theta_2) \right) \\ &= \|\mathbf{N}\| + \frac{y_2}{\|\mathbf{N}\|^2} (g''_{\theta_1} g'_{\theta_2} - g''_{\theta_2} g'_{\theta_1}) \sin(\theta_2). \end{aligned}$$

B Proof of Lemma 3.2

This section is devoted to the proof of Lemma 3.2. We start with some intermediate lemmas that will be used to decompose the intricate terms of Lemma 3.2 in smaller pieces.

B.1 Technical Lemmas

The following lemma contains Lipschitz estimates on several terms.

Lemma B.1. *For $R > 0$, there exists a constant $C = C(R) > 0$ such that for every $T \in (0, T_0)$ and every $(\cdot, \cdot, \theta_1^j, \theta_2^j) \in \mathbb{B}_R(T)$, the following estimates hold*

$$\|\Phi(\theta_1^a, \theta_2^a) - \Phi(\theta_1^b, \theta_2^b)\|_{L^\infty(0,T;\mathbf{L}^\infty(\Omega))} \leq CT\|\theta^a - \theta^b\|_{\Theta_T}, \quad (\text{B.1})$$

$$\|\mathcal{J}_\Phi(\theta_1^a, \theta_2^a) - \mathcal{J}_\Phi(\theta_1^b, \theta_2^b)\|_{L^\infty(0,T;\mathbf{H}^2(\Omega))} \leq CT\|\theta^a - \theta^b\|_{\Theta_T}, \quad (\text{B.2})$$

$$\|\mathcal{J}_\Psi(\theta_1^a, \theta_2^a, \Phi^a) - \mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b)\|_{L^\infty(0,T;\mathbf{H}^2(\Omega))} \leq CT\|\theta^a - \theta^b\|_{\Theta_T}, \quad (\text{B.3})$$

$$\|(\partial_{x_j} \mathcal{J}_\Psi(\theta_1^a, \theta_2^a)) \circ \Phi(\theta_1^a, \theta_2^a) - (\partial_{x_j} \mathcal{J}_\Psi(\theta_1^b, \theta_2^b)) \circ \Phi(\theta_1^b, \theta_2^b)\|_{L^\infty(0,T;\mathbf{H}^1(\Omega))} \leq CT\|\theta^a - \theta^b\|_{\Theta_T}, \quad (\text{B.4})$$

$$\|(\partial_{x_j}^2 \mathcal{J}_\Psi(\theta_1^a, \theta_2^a)) \circ \Phi(\theta_1^a, \theta_2^a) - (\partial_{x_j}^2 \mathcal{J}_\Psi(\theta_1^b, \theta_2^b)) \circ \Phi(\theta_1^b, \theta_2^b)\|_{L^\infty(0,T;\mathbf{L}^2(\Omega))} \leq CT\|\theta^a - \theta^b\|_{\Theta_T}, \quad (\text{B.5})$$

$$\|\mathcal{M}_{\theta_1^a, \theta_2^a} - \mathcal{M}_{\theta_1^b, \theta_2^b}\|_{L^\infty(0,T)} \leq CT\|\theta^a - \theta^b\|_{\Theta_T}, \quad (\text{B.6})$$

$$\|\mathbf{n}_{\theta_1^a, \theta_2^a}(\Phi(\theta_1^a, \theta_2^a)) - \mathbf{n}_{\theta_1^b, \theta_2^b}(\Phi(\theta_1^b, \theta_2^b))\|_{L^\infty(0,T;\mathbf{L}^\infty(\partial S_0))} \leq CT\|\theta^a - \theta^b\|_{\Theta_T}, \quad (\text{B.7})$$

$$\|\det(\mathcal{J}_\Psi(\theta_1^a, \theta_2^a)) - \det(\mathcal{J}_\Psi(\theta_1^b, \theta_2^b))\|_{L^\infty(0,T;\mathbf{L}^\infty(\Omega))} \leq CT\|\theta^a - \theta^b\|_{\Theta_T}, \quad (\text{B.8})$$

$$\|\partial_{\theta_j} \Phi(\theta_1^a, \theta_2^a, \cdot) - \partial_{\theta_j} \Phi(\theta_1^b, \theta_2^b, \cdot)\|_{L^\infty(0,T;\mathbf{H}^2(\Omega))} \leq CT\|\theta^a - \theta^b\|_{\Theta_T}, \quad (\text{B.9})$$

$$\|\partial_{\theta_k \theta_j} \Phi(\theta_1^a, \theta_2^a, \cdot) - \partial_{\theta_k \theta_j} \Phi(\theta_1^b, \theta_2^b, \cdot)\|_{L^\infty(0,T;\mathbf{H}^2(\Omega))} \leq CT\|\theta^a - \theta^b\|_{\Theta_T}, \quad (\text{B.10})$$

$$\| |\mathcal{J}_\Phi(\theta_1^a, \theta_2^a) \mathbf{t}_0| - |\mathcal{J}_\Phi(\theta_1^b, \theta_2^b) \mathbf{t}_0| \|_{L^\infty(0,T;\mathbf{L}^\infty(\partial S_0))} \leq CT\|\theta^a - \theta^b\|_{\Theta_T}, \quad (\text{B.11})$$

and

$$\|\partial_t \mathcal{J}_\Phi(\theta_1^a, \theta_2^a) - \partial_t \mathcal{J}_\Phi(\theta_1^b, \theta_2^b)\|_{L^\infty(0,T;\mathbf{H}^2(\Omega))} \leq C\|\theta^a - \theta^b\|_{\Theta_T}, \quad (\text{B.12})$$

$$\|\partial_t(\Psi(\theta_1^a, \theta_2^a)) \circ \Phi(\theta_1^a, \theta_2^a) - \partial_t(\Psi(\theta_1^b, \theta_2^b)) \circ \Phi(\theta_1^b, \theta_2^b)\|_{L^\infty(0,T;\mathbf{L}^\infty(\Omega))} \leq C\|\theta^a - \theta^b\|_{\Theta_T}, \quad (\text{B.13})$$

$$\|\partial_t(\mathcal{J}_\Psi(\theta_1^a, \theta_2^a)) \circ \Phi(\theta_1^a, \theta_2^a) - \partial_t(\mathcal{J}_\Psi(\theta_1^b, \theta_2^b)) \circ \Phi(\theta_1^b, \theta_2^b)\|_{L^\infty(0,T;\mathbf{L}^\infty(\Omega))} \leq C\|\theta^a - \theta^b\|_{\Theta_T}, \quad (\text{B.14})$$

$$\|\partial_t(\partial_{\theta_j} \Phi(\theta_1^a, \theta_2^a, \cdot)) - \partial_t(\partial_{\theta_j} \Phi(\theta_1^b, \theta_2^b, \cdot))\|_{L^\infty(0,T;\mathbf{H}^2(\mathcal{F}_0))} \leq C\|\theta^a - \theta^b\|_{\Theta_T}. \quad (\text{B.15})$$

Moreover, for every $(\tilde{\mathbf{u}}^j, \cdot, \theta_1^j, \theta_2^j) \in \mathbb{B}_R(T)$, the following estimates hold on \mathcal{G} defined in (3.3)

$$\|\mathcal{G}(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a) - \mathcal{G}(\theta_1^b, \theta_2^b, \tilde{\mathbf{u}}^b)\|_{L^2(0,T;\mathbf{L}^2(\partial S_0))} \leq C(\|\theta^a - \theta^b\|_{\Theta_T} + \|\tilde{\mathbf{u}}^a - \tilde{\mathbf{u}}^b\|_{\mathbf{U}_T}), \quad (\text{B.16})$$

$$\|\nabla \tilde{\mathbf{u}}^a - \mathcal{G}(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a) - \nabla \tilde{\mathbf{u}}^b + \mathcal{G}(\theta_1^b, \theta_2^b, \tilde{\mathbf{u}}^b)\|_{L^2(0,T;\mathbf{L}^2(\partial S_0))} \leq CT(\|\theta^a - \theta^b\|_{\Theta_T} + \|\tilde{\mathbf{u}}^a - \tilde{\mathbf{u}}^b\|_{\mathbf{U}_T}). \quad (\text{B.17})$$

In particular, as a direct application of Lemma B.1, using that $(0, 0, 0, 0) \in \mathbb{B}_R(T)$, we obtain the following lemma.

Lemma B.2. *For $R > 0$, there exists a constant $C = C(R) > 0$, such that for every $T \in (0, T_0)$ and every $(\cdot, \cdot, \theta_1, \theta_2) \in \mathbb{B}_R(T)$, the following estimates hold*

$$\|\mathcal{J}_\Phi(\theta_1, \theta_2) - I\|_{L^\infty(0,T;\mathbf{H}^2(\Omega))} \leq CT, \quad (\text{B.18})$$

$$\|\mathcal{J}_\Psi(\theta_1, \theta_2, \Phi(\theta_1, \theta_2)) - I\|_{L^\infty(0,T;\mathbf{H}^2(\Omega))} \leq CT, \quad (\text{B.19})$$

$$\|\partial_{x_j} \mathcal{J}_\Psi(\theta_1, \theta_2) \circ \Phi(\theta_1, \theta_2)\|_{L^\infty(0,T;\mathbf{H}^1(\Omega))} \leq CT, \quad (\text{B.20})$$

$$\|\partial_{x_j}^2 \mathcal{J}_\Psi(\theta_1, \theta_2) \circ \Phi(\theta_1, \theta_2)\|_{L^\infty(0,T;\mathbf{L}^2(\Omega))} \leq CT, \quad (\text{B.21})$$

$$\|\mathcal{M}_{\theta_1, \theta_2} - \mathcal{M}_{0,0}\|_{L^\infty(0,T)} \leq CT, \quad (\text{B.22})$$

$$\|\mathbf{n}_{\theta_1, \theta_2}(\Phi(\theta_1, \theta_2)) - \mathbf{n}_0\|_{L^\infty(0,T;\mathbf{L}^\infty(\partial S_0))} \leq CT, \quad (\text{B.23})$$

$$\| |\mathcal{J}_\Phi \mathbf{t}_0| - 1 \|_{L^\infty(0,T;\mathbf{L}^\infty(\partial S_0))} \leq CT, \quad (\text{B.24})$$

and

$$\|\partial_t(\mathcal{J}_\Phi(\theta_1, \theta_2))\|_{L^\infty(0,T;\mathbf{H}^2(\Omega))} \leq C, \quad (\text{B.25})$$

$$\left\| \frac{\partial}{\partial t}(\Psi(\theta_1, \theta_2)) \circ \Phi(\theta_1, \theta_2) \right\|_{L^\infty(0,T;\mathbf{L}^\infty(\Omega))} \leq C, \quad (\text{B.26})$$

$$\left\| \frac{\partial}{\partial t}(\mathcal{J}_\Psi(\theta_1, \theta_2)) \circ \Phi(\theta_1, \theta_2) \right\|_{L^\infty(0,T;\mathbf{L}^\infty(\Omega))} \leq C. \quad (\text{B.27})$$

Moreover, for every $(\tilde{\mathbf{u}}, \cdot, \theta_1, \theta_2) \in \mathbb{B}_R(T)$, we have the following estimate on \mathcal{G}

$$\|\nabla \tilde{\mathbf{u}} - \mathcal{G}(\theta_1, \theta_2, \tilde{\mathbf{u}})\|_{L^2(0,T;\mathbf{L}^2(\partial S_0))} \leq CT. \quad (\text{B.28})$$

Proof of Lemma B.1. Three kinds of estimates have to be proven. First estimates (B.1)–(B.10) are of the type

$$\|\alpha(\theta_1^a, \theta_2^a) - \alpha(\theta_1^b, \theta_2^b)\|_{L^\infty(0,T;\mathbb{X})} \leq CT\|(\theta_1^a, \theta_2^a) - (\theta_1^b, \theta_2^b)\|_{\Theta_T},$$

where α is a differentiable function defined on \mathbb{D}_Θ and valued in \mathbb{X} . We thus use Taylor series and get

$$\|\alpha(\theta_1^a, \theta_2^a) - \alpha(\theta_1^b, \theta_2^b)\|_{L^\infty(0,T;\mathbb{X})} \leq \sup_{(\theta_1, \theta_2) \in \mathbb{D}_\Theta} \|\nabla_\theta \alpha(\theta_1, \theta_2)\|_{L^\infty(0,T;\mathbb{X})} \|\theta^a - \theta^b\|_{L^\infty(0,T)}.$$

According to the definition of $\mathbb{B}_R(T)$ in (3.6), $\theta^a(0) = \theta^b(0) = (0, 0)$, we finish with

$$\|\theta^a - \theta^b\|_{L^\infty(0,T)} \leq T\|\theta^a - \theta^b\|_{\Theta_T}.$$

The second type of estimates (B.12)–(B.15) is of the form

$$\|\alpha(\theta_1^a, \theta_2^a, \dot{\theta}_1^a, \dot{\theta}_2^a) - \alpha(\theta_1^b, \theta_2^b, \dot{\theta}_1^b, \dot{\theta}_2^b)\|_{L^\infty(0,T;\mathbb{X})} \leq C\|(\theta_1^a, \theta_2^a) - (\theta_1^b, \theta_2^b)\|_{\Theta_T},$$

where α is now a function defined on $\mathbb{D}_\Theta \times \mathbb{R}^2$ with values in \mathbb{X} . We use the same strategy and get

$$\begin{aligned} & \|\alpha(\theta_1^a, \theta_2^a, \dot{\theta}_1^a, \dot{\theta}_2^a) - \alpha(\theta_1^b, \theta_2^b, \dot{\theta}_1^b, \dot{\theta}_2^b)\|_{L^\infty(0,T;\mathbb{X})} \\ & \leq \sup_{\substack{(\theta_1, \theta_2) \in \mathbb{D}_\Theta \\ |\omega_1| + |\omega_2| \leq R}} \|\nabla_{\theta, \omega} \alpha(\theta_1, \theta_2, \omega_1, \omega_2)\|_{L^\infty(0,T;\mathbb{X})} (\|\theta^a - \theta^b\|_{L^\infty(0,T)} + \|\dot{\theta}^a - \dot{\theta}^b\|_{L^\infty(0,T)}). \end{aligned}$$

Note that contrary to the first type of estimates, we do not have the decay in T because we did not enforce $\dot{\theta}^a(0) = \dot{\theta}^b(0)$.

Estimate (B.16) is a direct consequence of (B.17). The last estimate to prove is (B.17). We do it via the computation

$$\begin{aligned} & (\mathcal{G}(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a) - \mathcal{G}(\theta_1^b, \theta_2^b, \tilde{\mathbf{u}}^b) - \nabla \tilde{\mathbf{u}}^a + \nabla \tilde{\mathbf{u}}^b)_{ij} \\ & = \sum_k (\text{cof}(\partial_{x_j} \mathcal{J}_\Psi(\theta_1^a, \theta_2^a, \cdot) \circ \Phi^a)_{ki} - \text{cof}(\partial_{x_j} \mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \cdot) \circ \Phi^b)_{ki}) \tilde{u}_k^a \\ & + \sum_k \text{cof}(\partial_{x_j} \mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \cdot) \circ \Phi^b)_{ki} (\tilde{u}_k^a - \tilde{u}_k^b) \\ & + \sum_{k, \ell} \text{cof}(\mathcal{J}_\Psi(\theta_1^a, \theta_2^a, \Phi^a) - \mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b))_{ki} \frac{\partial \tilde{u}_k^a}{\partial y_\ell} \frac{\partial \Psi_\ell}{\partial x_j}(\theta_1^a, \theta_2^a, \Phi^a) \\ & + \sum_{k, \ell} \left(\text{cof}(\mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b))_{ki} \frac{\partial \Psi_\ell}{\partial x_j}(\theta_1^a, \theta_2^a, \Phi^a) - \delta_{ki} \delta_{\ell j} \right) \left(\frac{\partial \tilde{u}_k^a}{\partial y_\ell} - \frac{\partial \tilde{u}_k^b}{\partial y_\ell} \right) \\ & + \sum_{k, \ell} \text{cof}(\mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b))_{ki} \frac{\partial \tilde{u}_k^b}{\partial y_\ell} \left(\frac{\partial \Psi_\ell}{\partial x_j}(\theta_1^a, \theta_2^a, \Phi^a) - \frac{\partial \Psi_\ell}{\partial x_j}(\theta_1^b, \theta_2^b, \Phi^b) \right), \end{aligned}$$

and with the use of estimates (B.3), (B.4), (B.19) and (B.20) we get estimate (B.17). \square

B.2 Detailed proof of Lemma 3.2

Proof. In all the following estimates we use Lemmas B.1 and B.2.

• **Estimate (3.7)** is a consequence of the following estimates

$$\|\mathbf{F}^1(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a) - \mathbf{F}^1(\theta_1^b, \theta_2^b, \tilde{\mathbf{u}}^b)\|_{\mathbb{F}_T} \leq CT(\|\theta^a - \theta^b\|_{\Theta_T} + \|\tilde{\mathbf{u}}^a - \tilde{\mathbf{u}}^b\|_{\mathbb{U}_T}), \quad (\text{B.29})$$

$$\|\mathbf{F}^2(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a) - \mathbf{F}^2(\theta_1^b, \theta_2^b, \tilde{\mathbf{u}}^b)\|_{\mathbb{F}_T} \leq CT^{1/2}(\|\theta^a - \theta^b\|_{\Theta_T} + \|\tilde{\mathbf{u}}^a - \tilde{\mathbf{u}}^b\|_{\mathbb{U}_T}), \quad (\text{B.30})$$

$$\|\mathbf{F}^3(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a) - \mathbf{F}^3(\theta_1^b, \theta_2^b, \tilde{\mathbf{u}}^b)\|_{\mathbb{F}_T} \leq CT(\|\theta^a - \theta^b\|_{\Theta_T} + \|\tilde{\mathbf{u}}^a - \tilde{\mathbf{u}}^b\|_{\mathbb{U}_T}), \quad (\text{B.31})$$

$$\|\mathbf{F}^4(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a) - \mathbf{F}^4(\theta_1^b, \theta_2^b, \tilde{\mathbf{u}}^b)\|_{\mathbb{F}_T} \leq CT^{1/4}(\|\theta^a - \theta^b\|_{\Theta_T} + \|\tilde{\mathbf{u}}^a - \tilde{\mathbf{u}}^b\|_{\mathbb{U}_T}), \quad (\text{B.32})$$

$$\|\mathbf{F}^5(\theta_1^a, \theta_2^a, \tilde{p}^a) - \mathbf{F}^5(\theta_1^b, \theta_2^b, \tilde{p}^b)\|_{\mathbb{F}_T} \leq CT(\|\theta^a - \theta^b\|_{\Theta_T} + \|\tilde{p}^a - \tilde{p}^b\|_{\mathbb{P}_T}), \quad (\text{B.33})$$

where C does not depend on T . We now prove all of them.

• **Estimate (B.29):** We use the decomposition

$$\mathbf{F}^1(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a) - \mathbf{F}^1(\theta_1^b, \theta_2^b, \tilde{\mathbf{u}}^b) = (I - \text{cof}(\mathcal{J}_\Psi(\theta_1^a, \theta_2^a, \Phi^a))^T) \left(\frac{\partial \tilde{\mathbf{u}}^a}{\partial t} - \frac{\partial \tilde{\mathbf{u}}^b}{\partial t} \right) + \text{cof}(\mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b) - \mathcal{J}_\Psi(\theta_1^a, \theta_2^a, \Phi^a))^T \frac{\partial \tilde{\mathbf{u}}^b}{\partial t},$$

and we use estimates (B.19) and (B.3).

• **Estimate (B.30):** We use the decomposition

$$\begin{aligned} \mathbf{F}^2(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a) - \mathbf{F}^2(\theta_1^b, \theta_2^b, \tilde{\mathbf{u}}^b) &= -\operatorname{cof}\left(\frac{\partial}{\partial t}(\mathcal{J}_\Psi(\theta_1^a, \theta_2^a, \cdot)) \circ \Phi^a\right)^T \tilde{\mathbf{u}}^a + \operatorname{cof}\left(\frac{\partial}{\partial t}(\mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \cdot)) \circ \Phi^b\right)^T \tilde{\mathbf{u}}^b \\ &\quad + \operatorname{cof}\left(\mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b) - \mathcal{J}_\Psi(\theta_1^a, \theta_2^a, \Phi^a)\right)^T \nabla_{\mathbf{y}} \tilde{\mathbf{u}}^a \left(\frac{\partial}{\partial t} \Psi(\theta_1^a, \theta_2^a, \cdot)\right) \circ \Phi^a \\ &\quad + \operatorname{cof}(\mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b))^T \nabla_{\mathbf{y}}(\tilde{\mathbf{u}}^b - \tilde{\mathbf{u}}^a) \left(\frac{\partial}{\partial t} \Psi(\theta_1^a, \theta_2^a, \cdot)\right) \circ \Phi^a \\ &\quad + \operatorname{cof}(\mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b))^T \nabla_{\mathbf{y}} \tilde{\mathbf{u}}^b \left(\left(\frac{\partial}{\partial t} \Psi(\theta_1^b, \theta_2^b, \cdot)\right) \circ \Phi^b - \left(\frac{\partial}{\partial t} \Psi(\theta_1^a, \theta_2^a, \cdot)\right) \circ \Phi^a\right), \end{aligned}$$

where, for $j \in \{a, b\}$, we denote $\Phi^j = \Phi(\theta_1^j, \theta_2^j)$. We use estimates (B.27), (B.26), (B.14), (B.13), (B.3) and the estimate $\|\tilde{\mathbf{u}}\|_{L^2(0, T; \mathbf{H}^1(\mathcal{F}_0))} \leq T^{1/2} \|\tilde{\mathbf{u}}\|_{L^\infty(0, T; \mathbf{H}^1(\mathcal{F}_0))}$.

• **Estimate (B.31):** Since $\tilde{\mathbf{u}}$ has a space regularity of $\mathbf{H}_\beta^2(\mathcal{F}_0)$ and not $\mathbf{H}^2(\mathcal{F}_0)$, we need the following estimate in the sequel: for $f \in L^\infty(0, T; L^\infty(\mathcal{F}_0))$ with support within $\mathcal{F}_0 \cap \Omega_\varepsilon$ (defined in Lemma 1.6), we have

$$\left\| f \times \frac{\partial^2 \tilde{u}_k}{\partial y_\ell \partial y_m} \right\|_{L^2(0, T; L^2(\mathcal{F}_0))} \leq C \|f\|_{L^\infty(0, T; L^\infty(\mathcal{F}_0))} \|\tilde{\mathbf{u}}\|_{L^2(0, T; \mathbf{H}_\beta^2(\mathcal{F}_0))}, \quad (\text{B.34})$$

where C does not depend on T . It is a consequence of

$$\left\| f \times \frac{\partial^2 \tilde{u}_k}{\partial y_\ell \partial y_m} \right\|_{L^2(0, T; L^2(\mathcal{F}_0))} \leq \left\| \frac{f}{\prod_{n \in \mathcal{I}_{d,n}} r_n^\beta} \right\|_{L^\infty(0, T; L^\infty(\Omega_\varepsilon \cap \mathcal{F}_0))} \left\| \prod_{n \in \mathcal{I}_{d,n}} r_n^\beta \frac{\partial^2 \tilde{u}_k}{\partial y_\ell \partial y_m} \right\|_{L^2(0, T; L^2(\mathcal{F}_0))}.$$

We now use the decomposition

$$\left(\mathbf{F}^3(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a) - \mathbf{F}^3(\theta_1^b, \theta_2^b, \tilde{\mathbf{u}}^b)\right)_i = A_{1,i} + A_{2,i} + A_{3,i} + A_{4,i},$$

where

$$\begin{aligned} A_{1,i} &= \nu \sum_{j,k,\ell,m} \left(\operatorname{cof}(\mathcal{J}_\Psi(\theta_1^a, \theta_2^a, \Phi^a))_{ki} \frac{\partial \Psi_\ell}{\partial x_j}(\theta_1^a, \theta_2^a, \Phi^a) \frac{\partial \Psi_m}{\partial x_j}(\theta_1^a, \theta_2^a, \Phi^a) \right. \\ &\quad \left. - \operatorname{cof}(\mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b))_{ki} \frac{\partial \Psi_\ell}{\partial x_j}(\theta_1^b, \theta_2^b, \Phi^b) \frac{\partial \Psi_m}{\partial x_j}(\theta_1^b, \theta_2^b, \Phi^b) \right) \frac{\partial^2 \tilde{u}_k^a}{\partial y_\ell \partial y_m} \\ &\quad + \nu \sum_{j,k,\ell,m} \left(\operatorname{cof}(\mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b))_{ki} \frac{\partial \Psi_\ell}{\partial x_j}(\theta_1^b, \theta_2^b, \Phi^b) \frac{\partial \Psi_m}{\partial x_j}(\theta_1^b, \theta_2^b, \Phi^b) - \delta_{ki} \delta_{j\ell} \delta_{mj} \right) \left(\frac{\partial^2 \tilde{u}_k^a}{\partial y_\ell \partial y_m} - \frac{\partial^2 \tilde{u}_k^b}{\partial y_\ell \partial y_m} \right), \\ A_{2,i} &= 2\nu \sum_{j,k,\ell} \left(\operatorname{cof}(\partial_{x_j} \mathcal{J}_\Psi(\theta_1^a, \theta_2^a, \Phi^a))_{ki} \frac{\partial \Psi_\ell}{\partial x_j}(\theta_1^a, \theta_2^a, \Phi^a) - \operatorname{cof}(\partial_{x_j} \mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b))_{ki} \frac{\partial \Psi_\ell}{\partial x_j}(\theta_1^b, \theta_2^b, \Phi^b) \right) \frac{\partial \tilde{u}_k^a}{\partial y_\ell} \\ &\quad + 2\nu \sum_{j,k,\ell} \operatorname{cof}(\partial_{x_j} \mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b))_{ki} \frac{\partial \Psi_\ell}{\partial x_j}(\theta_1^b, \theta_2^b, \Phi^b) \left(\frac{\partial \tilde{u}_k^a}{\partial y_\ell} - \frac{\partial \tilde{u}_k^b}{\partial y_\ell} \right), \\ A_{3,i} &= \nu \sum_{j,k,\ell} \left(\operatorname{cof}(\mathcal{J}_\Psi(\theta_1^a, \theta_2^a, \Phi^a))_{ki} \frac{\partial^2 \Psi_\ell}{\partial x_j^2}(\theta_1^a, \theta_2^a, \Phi^a) - \operatorname{cof}(\mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b))_{ki} \frac{\partial^2 \Psi_\ell}{\partial x_j^2}(\theta_1^b, \theta_2^b, \Phi^b) \right) \frac{\partial \tilde{u}_k^a}{\partial y_\ell} \\ &\quad + \nu \sum_{j,k,\ell} \operatorname{cof}(\mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b))_{ki} \frac{\partial^2 \Psi_\ell}{\partial x_j^2}(\theta_1^b, \theta_2^b, \Phi^b) \left(\frac{\partial \tilde{u}_k^a}{\partial y_\ell} - \frac{\partial \tilde{u}_k^b}{\partial y_\ell} \right), \end{aligned}$$

and

$$A_{4,i} = \nu \sum_{j,k} \left(\operatorname{cof}(\partial_{x_j}^2 \mathcal{J}_\Psi(\theta_1^a, \theta_2^a, \Phi^a) - \partial_{x_j}^2 \mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b))_{ki} \tilde{u}_k^a + \operatorname{cof}(\partial_{x_j}^2 \mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b))_{ki} (\tilde{u}_k^a - \tilde{u}_k^b) \right).$$

Now,

- to estimate $A_{1,i}$, we use estimates (B.3), (B.19) and (B.34), also note that $\mathcal{J}_\Psi(\theta_1^a, \theta_2^a, \Phi^a) - \mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b)$ has support within Ω_ε .
- To estimate $A_{2,i}$, we use estimates (B.3), (B.4), (B.19) and (B.20).

- To estimate $A_{3,i}$, we use estimates (B.3), (B.4), (B.19) and (B.20).
- To estimate $A_{4,i}$, we use (B.5) and (B.21).

• **Estimate (B.32):** Let us first prove, for $f \in L^\infty(0, T; H^1(\mathcal{F}_0))$ and $g \in L^\infty(0, T; L^2(\mathcal{F}_0)) \cap L^2(0, T; H^{1/2}(\mathcal{F}_0))$, the following estimate

$$\|fg\|_{L^2(0,T;L^2(\mathcal{F}_0))} \leq CT^{1/4} \|f\|_{L^\infty(0,T;H^1(\mathcal{F}_0))} \|g\|_{L^\infty(0,T;L^2(\mathcal{F}_0))}^{1/2} \|g\|_{L^2(0,T;H^{1/2}(\mathcal{F}_0))}^{1/2}, \quad (\text{B.35})$$

where C does not depend on T .

- Hölder inequality with respect to time applied to $fg \times 1$ gives

$$\|fg\|_{L^2(0,T;L^2(\mathcal{F}_0))} \leq T^{1/4} \|fg\|_{L^4(0,T;L^2(\mathcal{F}_0))}.$$

- We have the relation $1/10 + 2/5 = 1/2$, then Hölder inequality yields

$$\|fg\|_{L^4(0,T;L^2(\mathcal{F}_0))} \leq \|f\|_{L^\infty(0,T;L^{10}(\mathcal{F}_0))} \|g\|_{L^4(0,T;L^{5/2}(\mathcal{F}_0))}.$$

- We have the Sobolev embedding $H^1(\mathcal{F}_0) \hookrightarrow L^{10}(\mathcal{F}_0) : \|f\|_{L^\infty(0,T;L^{10}(\mathcal{F}_0))} \leq C \|f\|_{L^\infty(0,T;H^1(\mathcal{F}_0))}$.
- We have the Sobolev embedding $H^{1/4}(\mathcal{F}_0) \hookrightarrow L^{5/2}(\mathcal{F}_0)$ (see [1, Theorem 7.58]) and the Sobolev interpolation $[L^2(\mathcal{F}_0), H^{1/2}(\mathcal{F}_0)]_{1/2} = H^{1/4}(\mathcal{F}_0) : \|g\|_{L^{5/2}(\mathcal{F}_0)} \leq C \|g\|_{H^{1/4}(\mathcal{F}_0)} \leq C \|g\|_{L^2(\mathcal{F}_0)}^{1/2} \|g\|_{H^{1/2}(\mathcal{F}_0)}^{1/2}$.
- We end the proof of (B.35) with

$$\|g\|_{L^4(0,T;L^{5/2}(\mathcal{F}_0))}^4 = \int_0^T \|g\|_{L^{5/2}(\mathcal{F}_0)}^4 \leq C \int_0^T \|g\|_{L^2(\mathcal{F}_0)}^2 \|g\|_{H^{1/2}(\mathcal{F}_0)}^2,$$

which implies

$$\|g\|_{L^4(0,T;L^{5/2}(\mathcal{F}_0))} \leq C \|g\|_{L^\infty(0,T;L^2(\mathcal{F}_0))}^{1/2} \|g\|_{L^2(0,T;H^{1/2}(\mathcal{F}_0))}^{1/2}.$$

We then use the decomposition

$$\left(\mathbf{F}^4(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a) - \mathbf{F}^4(\theta_1^b, \theta_2^b, \tilde{\mathbf{u}}^b) \right)_i = B_{1,i} + B_{2,i} + B_{3,i} + B_{4,i},$$

where

$$B_{1,i} = - \sum_{j,k,r} \left(\text{cof}(\mathcal{J}_\Psi(\theta_1^a, \theta_2^a, \Phi^a))_{kj} \text{cof}(\partial_{x_j} \mathcal{J}_\Psi(\theta_1^a, \theta_2^a, \Phi^a))_{ri} - \text{cof}(\mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b))_{kj} \text{cof}(\partial_{x_j} \mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b))_{ri} \right) \tilde{u}_k^a \tilde{u}_r^a,$$

$$B_{2,i} = - \sum_{j,k,r} \text{cof}(\mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b))_{kj} \text{cof}(\partial_{x_j} \mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b))_{ri} \left(\tilde{u}_k^a \tilde{u}_r^a - \tilde{u}_k^b \tilde{u}_r^b \right),$$

$$B_{3,i} = - \sum_{k,r} \left(\det(\mathcal{J}_\Psi(\theta_1^a, \theta_2^a, \Phi^a))^2 \frac{\partial \Phi_i^a}{\partial y_r} - \det(\mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b))^2 \frac{\partial \Phi_i^b}{\partial y_r} \right) \tilde{u}_k^a \frac{\partial \tilde{u}_r^a}{\partial y_k},$$

$$B_{4,i} = - \sum_{k,r} \det(\mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b))^2 \frac{\partial \Phi_i^b}{\partial y_r} \left(\tilde{u}_k^a \frac{\partial \tilde{u}_r^a}{\partial y_k} - \tilde{u}_k^b \frac{\partial \tilde{u}_r^b}{\partial y_k} \right).$$

We use

- estimates (B.3), (B.4), (B.19), (B.20) and (B.35) with $f = \tilde{u}_k^a$, $g = \tilde{u}_r^a$ for $B_{1,i}$,
- estimates (B.19), (B.20), (B.35) with $f = \tilde{u}_k^a$, $g = \tilde{u}_r^a - \tilde{u}_r^b$ and (B.35) with $f = \tilde{u}_k^a - \tilde{u}_k^b$, $g = \tilde{u}_r^b$ for $B_{2,i}$,
- estimates (B.3), (B.2), (B.18), (B.19) and (B.35) with $f = \tilde{u}_k^a$, $g = \frac{\partial \tilde{u}_r^a}{\partial y_k}$ for $B_{3,i}$,
- estimates (B.18), (B.19) (B.35) with $f = \tilde{u}_k^a$, $g = \frac{\partial \tilde{u}_r^a}{\partial y_k} - \frac{\partial \tilde{u}_r^b}{\partial y_k}$ and (B.35) with $f = \tilde{u}_k^a - \tilde{u}_k^b$, $g = \frac{\partial \tilde{u}_r^b}{\partial y_k}$ for $B_{4,i}$.

• **Estimate (B.33):** We have

$$\mathbf{F}^5(\theta_1^a, \theta_2^a, \tilde{p}^a) - \mathbf{F}^5(\theta_1^b, \theta_2^b, \tilde{p}^b) = \left(I - \mathcal{J}_\Psi(\theta_1^a, \theta_2^a, \Phi^a) \right)^T \nabla_{\mathbf{y}}(\tilde{p}^a - \tilde{p}^b) + \left(\mathcal{J}_\Psi(\theta_1^b, \theta_2^b, \Phi^b) - \mathcal{J}_\Psi(\theta_1^a, \theta_2^a, \Phi^a) \right)^T \nabla_{\mathbf{y}} \tilde{p}^b,$$

we use estimates (B.3), (B.19) and we adapt estimate (B.34) to obtain a $H_\beta^1(\mathcal{F}_0)$ norm for \tilde{p} .

• **Estimate (3.8):** We use, for $f \in L^2(0, T)$, the Hölder inequality

$$\|f\|_{L^2(0, T)} \leq T^{1/2} \|f\|_{L^\infty(0, T)}. \quad (\text{B.36})$$

We have

$$\frac{\partial}{\partial t} \left(\mathbf{G}(\theta_1^a, \theta_2^a, \dot{\theta}_1^a, \dot{\theta}_2^a) - \mathbf{G}(\theta_1^b, \theta_2^b, \dot{\theta}_1^b, \dot{\theta}_2^b) \right) = \mathbf{D}_1 + \mathbf{D}_2 + \mathbf{D}_3 + \mathbf{D}_4,$$

where

$$\begin{aligned} \mathbf{D}_1 &= \sum_j (\ddot{\theta}_j^a - \ddot{\theta}_j^b) \left(\text{cof}(\mathcal{J}_\Phi(\theta_1^a, \theta_2^a, \mathbf{y}))^T \partial_{\theta_j} \Phi(\theta_1^a, \theta_2^a, \mathbf{y}) - \partial_{\theta_j} \Phi(0, 0, \mathbf{y}) \right), \\ \mathbf{D}_2 &= \sum_j \ddot{\theta}_j^b \left(\text{cof}(\mathcal{J}_\Phi(\theta_1^a, \theta_2^a, \mathbf{y}))^T \partial_{\theta_j} \Phi(\theta_1^a, \theta_2^a, \mathbf{y}) - \text{cof}(\mathcal{J}_\Phi(\theta_1^b, \theta_2^b, \mathbf{y}))^T \partial_{\theta_j} \Phi(\theta_1^b, \theta_2^b, \mathbf{y}) \right), \\ \mathbf{D}_3 &= \sum_j (\dot{\theta}_j^a - \dot{\theta}_j^b) \left(\text{cof}(\partial_t(\mathcal{J}_\Phi(\theta_1^a, \theta_2^a, \mathbf{y})))^T \partial_{\theta_j} \Phi(\theta_1^a, \theta_2^a, \mathbf{y}) + \text{cof}(\mathcal{J}_\Phi(\theta_1^a, \theta_2^a, \mathbf{y}))^T \partial_t(\partial_{\theta_j} \Phi(\theta_1^a, \theta_2^a, \mathbf{y})) \right), \\ \mathbf{D}_4 &= \sum_j \dot{\theta}_j^b \left(\text{cof}(\partial_t(\mathcal{J}_\Phi(\theta_1^a, \theta_2^a, \mathbf{y})))^T \partial_{\theta_j} \Phi(\theta_1^a, \theta_2^a, \mathbf{y}) + \text{cof}(\mathcal{J}_\Phi(\theta_1^a, \theta_2^a, \mathbf{y}))^T \partial_t(\partial_{\theta_j} \Phi(\theta_1^a, \theta_2^a, \mathbf{y})) \right. \\ &\quad \left. - \text{cof}(\partial_t(\mathcal{J}_\Phi(\theta_1^b, \theta_2^b, \mathbf{y})))^T \partial_{\theta_j} \Phi(\theta_1^b, \theta_2^b, \mathbf{y}) + \text{cof}(\mathcal{J}_\Phi(\theta_1^b, \theta_2^b, \mathbf{y}))^T \partial_t(\partial_{\theta_j} \Phi(\theta_1^b, \theta_2^b, \mathbf{y})) \right). \end{aligned}$$

We use

- (B.18) and (B.9) for \mathbf{D}_1 ,
- (B.2), (B.18) and (B.9) for \mathbf{D}_2 ,
- (B.36), (B.25), (B.9), (B.18) and (B.15) for \mathbf{D}_3 ,
- (B.36), (B.12), (B.9), (B.2), (B.15), (B.18) and (B.25) for \mathbf{D}_4 ,

this gives us the estimate on $\frac{\partial}{\partial t} (\mathbf{G}(\theta_1^a, \theta_2^a, \dot{\theta}_1^a, \dot{\theta}_2^a) - \mathbf{G}(\theta_1^b, \theta_2^b, \dot{\theta}_1^b, \dot{\theta}_2^b))$. Similar arguments give the estimate on $\mathbf{G}(\theta_1^a, \theta_2^a, \dot{\theta}_1^a, \dot{\theta}_2^a) - \mathbf{G}(\theta_1^b, \theta_2^b, \dot{\theta}_1^b, \dot{\theta}_2^b)$.

• **Estimate (3.9):** we use the following decomposition

$$\left(\mathbf{S}(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a, \tilde{p}^a) - \mathbf{S}(\theta_1^b, \theta_2^b, \tilde{\mathbf{u}}^b, \tilde{p}^b) \right)_i = E_{1,i} + E_{2,i} + E_{3,i} + E_{4,i} + E_{5,i} + E_{6,i},$$

where

$$\begin{aligned} E_{1,i} &= - \left(\left(\mathcal{M}_{\theta_1^a, \theta_2^a} - \mathcal{M}_{\theta_1^b, \theta_2^b} \right) \begin{pmatrix} \ddot{\theta}_1^a \\ \ddot{\theta}_2^a \end{pmatrix} - \left(\mathcal{M}_{\theta_1^b, \theta_2^b} - \mathcal{M}_{0,0} \right) \begin{pmatrix} \ddot{\theta}_1^a - \ddot{\theta}_1^b \\ \ddot{\theta}_2^a - \ddot{\theta}_2^b \end{pmatrix} \right)_i, \\ E_{2,i} &= \left(\mathbf{M}_I(\theta_1^a, \theta_2^a, \dot{\theta}_1^a, \dot{\theta}_2^a) - \mathbf{M}_I(\theta_1^b, \theta_2^b, \dot{\theta}_1^b, \dot{\theta}_2^b) \right)_i, \\ E_{3,i} &= \int_{\partial S_0} (|\mathcal{J}_\Phi^a \mathbf{t}_0| - |\mathcal{J}_\Phi^b \mathbf{t}_0|) \left(\tilde{p}^a I - \nu(\mathcal{G}(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a) + \mathcal{G}(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a)^T) \right) \mathbf{n}_{\theta_1^a, \theta_2^a}(\Phi^a) \cdot \partial_{\theta_i} \Phi^a, \\ E_{4,i} &= \int_{\partial S_0} |\mathcal{J}_\Phi^b \mathbf{t}_0| \left(\tilde{p}^a I - \nu(\mathcal{G}(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a) + \mathcal{G}(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a)^T) \right) \mathbf{n}_{\theta_1^a, \theta_2^a}(\Phi^a) \cdot (\partial_{\theta_i} \Phi^a - \partial_{\theta_i} \Phi^b), \\ E_{5,i} &= \int_{\partial S_0} |\mathcal{J}_\Phi^b \mathbf{t}_0| \left(\tilde{p}^a I - \nu(\mathcal{G}(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a) + \mathcal{G}(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a)^T) \right) (\mathbf{n}_{\theta_1^a, \theta_2^a}(\Phi^a) - \mathbf{n}_{\theta_1^b, \theta_2^b}(\Phi^b)) \cdot \partial_{\theta_i} \Phi^b, \\ E_{6,i} &= \int_{\partial S_0} |\mathcal{J}_\Phi^b \mathbf{t}_0| \left((\tilde{p}^a - \tilde{p}^b) I - \nu(\mathcal{G}(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a) + \mathcal{G}(\theta_1^a, \theta_2^a, \tilde{\mathbf{u}}^a)^T - \mathcal{G}(\theta_1^b, \theta_2^b, \tilde{\mathbf{u}}^b) - \mathcal{G}(\theta_1^b, \theta_2^b, \tilde{\mathbf{u}}^b)^T) \right) \mathbf{n}_{\theta_1^b, \theta_2^b} \cdot \partial_{\theta_i} \Phi^b \\ &\quad - \int_{\partial S_0} \left((\tilde{p}^a - \tilde{p}^b) I - \nu(\nabla \tilde{\mathbf{u}}^a - \nabla \tilde{\mathbf{u}}^b + (\nabla \tilde{\mathbf{u}}^a - \nabla \tilde{\mathbf{u}}^b)^T) \right) \mathbf{n}_0 \cdot \partial_{\theta_i} \Phi(0, 0, \gamma_y). \end{aligned}$$

Then we use

- (B.9) for $E_{1,i}$,
- (B.9), (B.10) and (B.36) for $E_{2,i}$,
- (B.11), (B.28), (B.23) and (B.9) for $E_{3,i}$,

- (B.24), (B.28), (B.23) and (B.9) for $E_{4,i}$,
 - (B.24), (B.28), (B.7) and (B.9) for $E_{5,i}$,
 - (B.24), (B.17), (B.23) and (B.9) for $E_{6,i}$.
- **Estimate (3.10):** we use the Lipschitz regularity of $\mathbf{f}_{\mathcal{F}}$ and estimate (B.1),

$$\begin{aligned} & \|\mathbf{f}_{\mathcal{F}}(t, \Phi(\theta_1^a, \theta_2^a, \mathbf{y})) - \mathbf{f}_{\mathcal{F}}(t, \Phi(\theta_1^b, \theta_2^b, \mathbf{y}))\|_{L^2(0,T;L^2(\mathcal{F}_0))} \\ & \leq C \|\mathbf{f}_{\mathcal{F}}\|_{L^2(0,T;W^{1,\infty}(\Omega))} \|\Phi(\theta_1^a, \theta_2^a, \mathbf{y}) - \Phi(\theta_1^b, \theta_2^b, \mathbf{y})\|_{L^\infty(0,T;L^\infty(\mathcal{F}_0))} \\ & \leq CT \|\theta^a - \theta^b\|_{\Theta_T}. \end{aligned}$$

□

References

- [1] R. A. Adams. *Sobolev spaces*. Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London, 1975. Pure and Applied Mathematics, Vol. 65.
- [2] Y. Bazilevs, K. Takizawa, and T. E. Tezduyar. *Computational Fluid-Structure Interaction, Methods and Applications*. Wiley, 2013.
- [3] H. Beirão da Veiga. On the existence of strong solutions to a coupled fluid-structure evolution problem. *J. Math. Fluid Mech.*, 6(1):21–52, 2004.
- [4] A. Bensoussan, G. Da Prato, M. C. Delfour, and S. K. Mitter. *Representation and control of infinite dimensional systems*. Systems & Control: Foundations & Applications. Birkhäuser Boston, Inc., Boston, MA, second edition, 2007.
- [5] M. Boulakia. Existence of weak solutions for the three-dimensional motion of an elastic structure in an incompressible fluid. *J. Math. Fluid Mech.*, 9(2):262–294, 2007.
- [6] M. Boulakia and S. Guerrero. A regularity result for a solid-fluid system associated to the compressible Navier-Stokes equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 26(3):777–813, 2009.
- [7] M. Boulakia and S. Guerrero. Regular solutions of a problem coupling a compressible fluid and an elastic structure. *J. Math. Pures Appl. (9)*, 94(4):341–365, 2010.
- [8] M. Boulakia and S. Guerrero. On the interaction problem between a compressible fluid and a Saint-Venant Kirchhoff elastic structure. *Adv. Differential Equations*, 22(1-2):1–48, 2017.
- [9] M. Boulakia, E. L. Schwindt, and T. Takahashi. Existence of strong solutions for the motion of an elastic structure in an incompressible viscous fluid. *Interfaces Free Bound.*, 14(3):273–306, 2012.
- [10] L. Bălilescu, J. San Martín, and T. Takahashi. Fluid-rigid structure interaction system with Coulomb’s law. *SIAM J. Math. Anal.*, 49(6):4625–4657, 2017.
- [11] A. Chambolle, B. Desjardins, M. J. Esteban, and C. Grandmont. Existence of weak solutions for the unsteady interaction of a viscous fluid with an elastic plate. *J. Math. Fluid Mech.*, 7(3):368–404, 2005.
- [12] G. Duvaut and J.-L. Lions. *Les inéquations en mécanique et en physique*. Dunod, Paris, 1972. Travaux et Recherches Mathématiques, No. 21.
- [13] G. P. Galdi. *An introduction to the mathematical theory of the Navier-Stokes equations*. Springer Monographs in Mathematics. Springer, New York, second edition, 2011.
- [14] C. Grandmont and Y. Maday. Existence for an unsteady fluid-structure interaction problem. *M2AN Math. Model. Numer. Anal.*, 34(3):609–636, 2000.
- [15] P. Grisvard. *Elliptic problems in nonsmooth domains*, volume 69 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2011.
- [16] M. Hillairet. Lack of collision between solid bodies in a 2D incompressible viscous flow. *Comm. Partial Differential Equations*, 32(7-9):1345–1371, 2007.
- [17] J.G. Houot, J. San Martin, and M. Tucsnak. Existence of solutions for the equations modeling the motion of rigid bodies in an ideal fluid. *J. Funct. Anal.*, 259(11):2856–2885, 2010.

- [18] J. Lequeurre. Existence of strong solutions to a fluid-structure system. *SIAM J. Math. Anal.*, 43(1):389–410, 2011.
- [19] J.-L. Lions and E. Magenes. *Problèmes aux limites non homogènes et applications. Vol. 1.* Travaux et Recherches Mathématiques, No. 17. Dunod, Paris, 1968.
- [20] Y. Liu, T. Takahashi, and M. Tucsnak. Single input controllability of a simplified fluid-structure interaction model. *ESAIM Control Optim. Calc. Var.*, 19(1):20–42, 2013.
- [21] D. Maity, T. Takahashi, and M. Tucsnak. Analysis of a system modelling the motion of a piston in a viscous gas. *J. Math. Fluid Mech.*, 19(3):551–579, 2017.
- [22] V. Maz’ya and J. Rossmann. *Elliptic equations in polyhedral domains*, volume 162 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2010.
- [23] Š. Nečasová, T. Takahashi, and M. Tucsnak. Weak solutions for the motion of a self-propelled deformable structure in a viscous incompressible fluid. *Acta Appl. Math.*, 116(3):329–352, 2011.
- [24] P. A. Nguyen and J.-P. Raymond. Boundary stabilization of the Navier-Stokes equations in the case of mixed boundary conditions. *SIAM J. Control Optim.*, 53(5):3006–3039, 2015.
- [25] A. Pazy. *Semigroups of linear operators and applications to partial differential equations*, volume 44 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1983.
- [26] J.-P. Raymond. Stokes and Navier-Stokes equations with nonhomogeneous boundary conditions. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 24(6):921–951, 2007.
- [27] J.-P. Raymond and M. Vanninathan. A fluid-structure model coupling the Navier-Stokes equations and the Lamé system. *J. Math. Pures Appl. (9)*, 102(3):546–596, 2014.
- [28] J. San Martín, J.-F. Scheid, T. Takahashi, and M. Tucsnak. An initial and boundary value problem modeling of fish-like swimming. *Arch. Ration. Mech. Anal.*, 188(3):429–455, 2008.
- [29] J.A. San Martín, V. Starovoitov, and M. Tucsnak. Global weak solutions for the two-dimensional motion of several rigid bodies in an incompressible viscous fluid. *Arch. Ration. Mech. Anal.*, 161(2):113–147, 2002.
- [30] H. Sohr. *The Navier-Stokes equations*. Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks]. Birkhäuser Verlag, Basel, 2001.
- [31] T. Takahashi. Analysis of strong solutions for the equations modeling the motion of a rigid-fluid system in a bounded domain. *Adv. Differential Equations*, 8(12):1499–1532, 2003.
- [32] T. Takahashi, M. Tucsnak, and G. Weiss. Stabilization of a fluid-rigid body system. *J. Differential Equations*, 259(11):6459–6493, 2015.