



HAL
open science

Effects of spatial aggregation of nests on population recruitment: the case of a small population of Atlantic salmon

Colin Bouchard, Agnes Bardonnnet, Mathieu Buoro, Cédric Tentelier

► **To cite this version:**

Colin Bouchard, Agnes Bardonnnet, Mathieu Buoro, Cédric Tentelier. Effects of spatial aggregation of nests on population recruitment: the case of a small population of Atlantic salmon. International Statistical Ecology Conference (ISEC 2018), Jul 2018, St Andrews, United Kingdom. 328 p., 2018, ISEC 2018 Conference Book. 10.13140/RG.2.2.35939.04643 . hal-01860020

HAL Id: hal-01860020

<https://hal.science/hal-01860020>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ISEC 2018 Conference Book



ISEC

International
Statistical
Ecology
Conference

University of St Andrews

2nd – 6th July 2018



University of
St Andrews

Organization

ISEC 2018 is organized by



CREEM

Centre for Research into Ecological
and Environmental Modelling

A research group based at



University of
St Andrews | FOUNDED
1413 |

We are delighted to receive you !

Effects of spatial aggregation of nests on population recruitment: the case of a small population of Atlantic salmon

Colin Bouchard¹, Agnès Bardonnnet², Mathieu Buoro², Cédric Tentelier¹
¹University of Pau and Pays de l'Adour, ²French National Institute of Agronomic Research (INRA)

Abstract ID: 2474
Poster Session (I)/27

In natural populations, individuals are often distributed in such a way that aggregation increases at some places, and decreases at others (Hoare et al. 2004, Broly et al. 2012). Aggregation may modify ecological processes acting at individual level such as breeding interactions or competition, and affect population demography (Chesson & Neuhauser 2002, Murrell et al. 2002). Including aggregation data into stock-recruitment relationships is expected to improve estimations of recruitment, but also to test the effects of aggregation on population recruitment.

Spatial aggregation of Atlantic salmon (*Salmo salar*) nests is expected to diminish recruitment through density-dependent mortality unless females aggregate in the best breeding sites (Hendry et al. 2001, Tentelier et al. 2016), and leads to steadier recruitment under hypothesis that females select safest breeding sites (Gauthey et al. 2017).

Yearly nest mapping in the small Atlantic salmon population of the Nivelle (France) was used to compute spatial aggregation under different scales (Lloyd 1967). This aggregation was incorporated in different stock-recruitment models, linking egg density to juvenile density over a 30 years period. Heteroscedastic hierarchical models without stock were also fitted to check whether accounting for the stock is important when testing the effect of aggregation on recruitment. We found that population recruitment was not impaired by aggregation of nests, whereas aggregation diminished the variability of population recruitment. In addition, stock-recruitment models provided better estimates than models ignoring the stock.

The originality of this work was to use local distribution to compute aggregation at different scales and add it in different stock-recruitment relationships. Altogether, our results indicated that salmon females select breeding sites on environmental risk and on habitat quality. In addition, our results suggest that females first aggregate in best sites increasing aggregation at a low number of breeders, but spread out in sites of low quality when population size increases. These results are consistent with previous studies on salmon (Hendry et al. 2001, Gauthey et al. 2017). Finally, we demonstrated that incorporating ecological data in stock-recruitment models is relevant to improve our understanding of demographic processes.

References

- Broly P, Mullier R, Deneubourg J-L, Devigne C (2012) Aggregation in woodlice: social interaction and density effects. *ZooKeys* 176:133–144
- Chesson P, Neuhauser C (2002) Intraspecific aggregation and species coexistence. *Trends Ecol Evol* 17:210–211
- Gauthey Z, Lang M, Elozegi A, Tentelier C, Rives J, Labonne J (2017) Brown trout spawning habitat selection and its effects on egg survival. *Ecol Freshw Fish* 26:133–140
- Hendry AP, Berg OK, Quinn TP (2001) Breeding location choice in salmon: causes (habitat, competition, body size, energy stores) and consequences (life span, energy stores). *Oikos* 93:407–418
- Hoare D., Couzin I., Godin J-G., Krause J (2004) Context-dependent group size choice in fish. *Anim Behav* 67:155–164
- Lloyd M (1967) Mean crowding. *J Anim Ecol* 36:1–30