



HAL
open science

A random Schrödinger operator associated with the Vertex Reinforced Jump Process on infinite graphs

Christophe Sabot, Xiaolin Zeng

► **To cite this version:**

Christophe Sabot, Xiaolin Zeng. A random Schrödinger operator associated with the Vertex Reinforced Jump Process on infinite graphs. *Journal of the American Mathematical Society*, 2019, 32, pp.311-349. 10.1090/jams/906 . hal-01863722

HAL Id: hal-01863722

<https://hal.science/hal-01863722>

Submitted on 12 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A RANDOM SCHRÖDINGER OPERATOR ASSOCIATED WITH THE VERTEX REINFORCED JUMP PROCESS ON INFINITE GRAPHS

CHRISTOPHE SABOT AND XIAOLIN ZENG

ABSTRACT. This paper concerns the Vertex reinforced jump process (VRJP), the Edge reinforced random walk (ERRW) and their link with a random Schrödinger operator. On infinite graphs, we define a 1-dependent random potential β extending that defined in [20] on finite graphs, and consider its associated random Schrödinger operator H_β . We construct a random function ψ as a limit of martingales, such that $\psi = 0$ when the VRJP is recurrent, and ψ is a positive generalized eigenfunction of the random Schrödinger operator with eigenvalue 0, when the VRJP is transient. Then we prove a representation of the VRJP on infinite graphs as a mixture of Markov jump processes involving the function ψ , the Green function of the random Schrödinger operator and an independent Gamma random variable. On \mathbb{Z}^d , we deduce from this representation a zero-one law for recurrence or transience of the VRJP and the ERRW, and a functional central limit theorem for the VRJP and the ERRW at weak reinforcement in dimension $d \geq 3$, using estimates of [10, 8]. Finally, we deduce recurrence of the ERRW in dimension $d = 2$ for any initial constant weights (using the estimates of Merkl and Rolles, [15, 17]), thus giving a full answer to the old question of Diaconis. We also raise some questions on the links between recurrence/transience of the VRJP and localization/delocalization of the random Schrödinger operator H_β .

1. INTRODUCTION

This paper concerns the Vertex Reinforced Jump Process (VRJP) and the Edge Reinforced Random Walk (ERRW) and their relation with a random Schrödinger operator associated with a stationary 1-dependent random potential (i.e. the potential is independent at distance larger or equal to 2).

The VRJP is a continuous time self-interacting process introduced in [5], investigated on trees in [3, 2] and on general graphs in [20, 21]. We first recall its definition. Let $\mathcal{G} = (V, E)$ be an undirected graph with finite degree at each vertex. We write $i \sim j$ if $i \in V$, $j \in V$ and $\{i, j\}$ is an edge of the graph. We always assume that the graph is connected and has no trivial loops (i.e. vertex i such that $i \sim i$). Let $(W_{i,j})_{i \sim j}$ be a set of positive conductances, $W_{i,j} > 0$, $W_{i,j} = W_{j,i}$. The VRJP is the continuous time process $(Y_s)_{s \geq 0}$ on V , starting at time 0 at some vertex $i_0 \in V$, which, conditionally on the past at time s , if $Y_s = i$, jumps to a neighbour j of i at rate

$$W_{i,j} L_j(s),$$

where

$$L_j(s) := 1 + \int_0^s \mathbb{1}_{\{Y_u=j\}} du.$$

This work was supported by the LABEX MILYON (ANR-10-LABX-0070) of Université de Lyon, within the program "Investissements d'Avenir" (ANR-11-IDEX-0007) operated by the French National Research Agency (ANR), and by the ANR/FNS project MALIN (ANR-16-CE93-0003). The second author is supported by ERC Starting Grant 678520.

In [20], Sabot and Tarrès introduced the following time change of the VRJP

$$(1.1) \quad Z_t = Y_{D^{-1}(t)},$$

where $D(s)$ is the following increasing function

$$D(s) = \sum_{i \in V} (L_i^2(s) - 1).$$

We call this process the VRJP in exchangeable time scale and denote by $\mathbb{P}_{i_0}^{\text{VRJP}}$ its law starting from the vertex i_0 . When the graph is finite it is proved in [20, Theorem 2] that the VRJP in exchangeable time scale $(Z_t)_{t \geq 0}$ is a mixture of Markov jump processes. More precisely, there exists a random field $(u_j)_{j \in V}$ such that Z is a mixture of Markov jump processes with jump rates from i to j

$$\frac{1}{2} W_{i,j} e^{u_j - u_i}.$$

The law of (u_j) is explicit, c.f. [20, Theorem 2], and forthcoming Theorem B. It appears to be a marginal of a supersymmetric sigma-field which had been investigated previously by Disertori, Spencer, Zirnbauer (c.f. [9], [10], [24]). As a consequence of this representation and of [9], [10], it was proved in [20] the following: when the graph has bounded degree, there exists a real $\lambda_0 > 0$ such that if $W_{i,j} \leq \lambda_0$ for all $i \sim j$, then the VRJP is positively recurrent, more precisely, Z is a mixture of positive recurrent Markov jump processes. When the graph is the grid \mathbb{Z}^d , with $d \geq 3$, there exists $\lambda_1 < +\infty$ such that if $W_{i,j} \geq \lambda_1$ for all $i \sim j$, the VRJP is transient. Hence, it shows a phase transition between recurrence and transience in dimension $d \geq 3$. The question of the representation of the VRJP on infinite graphs as a mixture of Markov jump processes is non trivial, especially in the transient case. It is possible to prove such a representation by a weak convergence argument, following [16], but it gives little information on the mixing law. In this paper we prove such a representation involving the Green function and a generalized eigenfunction of a random Schrödinger operator.

Let us give a flavour of the main results of the paper in the case of the VRJP on \mathbb{Z}^d with $W_{i,j} = W$ constant. We construct a positive 1-dependent random potential $(\beta_j)_{j \in \mathbb{Z}^d}$ (i.e. two subsets of the β 's are independent if their indices are at least at distance 2) and with marginal given by inverse of Inverse Gaussian law with parameters $(\frac{1}{dW}, 1)$. This field is a natural extension to infinite graphs of the field defined by Sabot, Tarrès, Zeng in [22]. We consider the random Schrödinger operator

$$H_\beta = -W\Delta + V,$$

where Δ is the usual discrete (non-positive) Laplacian and V is the multiplication operator defined by $V_j = 2\beta_j - 2dW$. Hence, it corresponds to the Anderson model with a random potential which is not i.i.d. but only stationary and 1-dependent. When the VRJP is transient we prove that there exists a positive generalized eigenfunction ψ of H_β with eigenvalue 0, stationary and ergodic. Let $(G(i, j))_{i \in \mathbb{Z}^d, j \in \mathbb{Z}^d}$ be defined by

$$G(i, j) = \widehat{G}(i, j) + \frac{1}{2} \gamma^{-1} \psi(i) \psi(j),$$

where $\widehat{G} = (H_\beta)^{-1}$ is the Green function (which happens to be well-defined in an appropriate sense) and γ is an extra random variable independent of the field β with law $\text{Gamma}(\frac{1}{2}, 1)$. We prove the following representation for the VRJP: the VRJP in exchangeable time scale

Z starting from the point i_0 is a mixture of Markov jump processes with jump rates from i to j

$$(1.2) \quad \frac{1}{2} W_{i,j} \frac{G(i_0, j)}{G(i_0, i)}.$$

When the VRJP is recurrent the same representation is valid with $\psi = 0$. In fact, the function ψ is the a.s. limit of a martingale, the limit being positive when the VRJP is transient and 0 when the VRJP is recurrent. It is remarkable that when the VRJP is recurrent it can be represented as a mixture with β -measurable jump rates, but when the VRJP is transient it involves an extra independent Gamma random variable. This representation extends to infinite graphs the representation given in [22] for finite graphs. A new feature appears in the transient case, where the generalized eigenfunction ψ is involved in the representation. We suspect that recurrence/transience of the VRJP is related to localization/delocalization of the random Schrödinger operator H_β at the bottom of the spectrum.

The representation (1.2) has several consequences on the VRJP and the ERRW. The ERRW is a reinforced process introduced by Diaconis and Coppersmith in 1986 (see Section 2.5 for a definition). The recurrence of the 2-dimensional ERRW is a famous open question raised by Diaconis, see [4, 18, 12, 17] for early references. Important progress has been done recently in the understanding of this process. In particular, in [20], an explicit relation between the ERRW and the VRJP was stated, thus somehow reducing the analysis of the ERRW to that of the VRJP. In [20, 1], it was proved by rather different methods that the ERRW on any graph with bounded degree at strong enough reinforcement is positive recurrent. In [8], it was proved that the ERRW is transient on \mathbb{Z}^d , $d \geq 3$, at weak reinforcement.

The representation (1.2) allows us to complete the picture both in dimension 2 and in the transient regime. More precisely, we prove a functional central limit theorem for the ERRW and for the discrete time process associated with the VRJP in dimension $d \geq 3$ at weak reinforcement, using the estimates of [10, 8]. Using the polynomial estimate provided by Merkl and Rolles, [17], we are able to prove recurrence of ERRW on \mathbb{Z}^2 for all initial constant weights, hence giving a full answer to the question of Diaconis.

2. STATEMENTS OF THE RESULTS

2.1. Notations. We denote by \mathbb{R}_+ (resp. \mathbb{R}_+^*) the set of non-negative (resp. positive) reals.

Let $\mathcal{G} = (V, E)$ be an undirected, locally finite, connected graph without trivial loops or multiples edges. For $i, j \in V$, write $i \sim j$ if i is a neighbor of j . We write $d_{\mathcal{G}}$ for the graph distance in \mathcal{G} , and for two subsets U, U' of V , we define $d_{\mathcal{G}}(U, U') = \inf_{i \in U, j \in U'} d_{\mathcal{G}}(i, j)$. We suppose given, for each edge $e = \{i, j\} \in E$, a positive real $W_{i,j} > 0$, understood as the conductance of the edge e . In this case we call $(\mathcal{G}, (W_e)_{e \in E})$ a graph with conductances.

Convention: We adopt the notation $\sum_{i \sim j}$ for the sum on all undirected edges $\{i, j\}$, counting only once each edge.

When $\beta = (\beta_i)_{i \in V} \in \mathbb{R}^V$ is a real vector indexed by the vertices and $U \subset V$, we write β_U for the restriction of β to U , i.e. $\beta_U = (\beta_i)_{i \in U}$. When $A = (A_{i,j})_{i,j \in V} \in \mathbb{R}^{V \times V}$ is a real function on $V \times V$ and $U \subset V$, $U' \subset V$, we write $A_{U,U'}$ for the restriction of A to $U \times U'$, i.e. $A_{U,U'} = (A_{i,j})_{i \in U, j \in U'}$.

It will be convenient to define the continuous time processes that appear in the text on the same canonical space. In the sequel, we will denote by $D([0, \infty), V)$ the space of càdlàg

functions from $[0, \infty)$ to V . The law of the VRJP in exchangeable time scale defined in (1.1), starting from i_0 , will be denoted by $\mathbb{P}_{i_0}^{\text{VRJP}}$, which is a probability on $D([0, \infty), V)$. The VRJP will always be defined on the canonical space and $(Z_t)_{t \in \mathbb{R}_+}$ will denote the canonical process defined by $Z_t(\omega) = \omega(t)$ for $\omega \in D([0, \infty), V)$.

Remark 1. *We do not allow multiple edges or trivial loops since it does not bring more generality to the VRJP. Indeed, from its definition, it follows that the VRJP on a graph with multiple edges and trivial loops has the same law as the VRJP on the graph where trivial loops are removed and multiple edges are replaced by a single edge by summing the conductances of the multiples edges. Similarly, the law on random potentials that appears in the sequel can always be reduced to graphs without multiple edges or trivial loops. Nevertheless, in Section 5 it simplifies notations to allow trivial loops.*

2.2. Representation of the VRJP on infinite graphs. Define the operator $P = (P_{i,j})_{i,j \in V}$ by

$$P_{i,j} = \begin{cases} W_{i,j}, & \text{if } i \sim j, \\ 0, & \text{otherwise.} \end{cases}$$

We define below a probability distribution on potentials on the graph. A potential on the graph will generically be denoted $\beta = (\beta_i)_{i \in V} \in \mathbb{R}^V$. With the potential $\beta \in \mathbb{R}^V$, we associate the Schrödinger operator on \mathcal{G}

$$(2.1) \quad H_\beta = -P + 2\beta,$$

where β represents the operator of multiplication by the potential (β_i) (or equivalently the diagonal operator with diagonal terms $(\beta_i)_{i \in V}$).

We denote by

$$(2.2) \quad \mathcal{D}_V^W = \{\beta \in \mathbb{R}^V, (H_\beta)_{U,U} > 0 \text{ for all finite subsets } U \subset V\},$$

where $(H_\beta)_{U,U} > 0$ means that the restriction of H_β to $U \times U$ is positive definite. Obviously, $\mathcal{D}_V^W \subset (\mathbb{R}_+^*)^V$ since when $U = \{i\}$ the restriction of H_β is the real $2\beta_i$. We endow \mathcal{D}_V^W with its Borelian σ -field denoted $\mathcal{B}(\mathcal{D}_V^W)$.

The following statement extends the random potential defined in [22, Theorem 1] to infinite graphs.

Proposition 1. *Let $(\mathcal{G}, (W_e)_{e \in E})$ be a graph with conductances as defined in Section 2.1. There exists a unique probability distribution ν_V^W defined on $(\mathcal{D}_V^W, \mathcal{B}(\mathcal{D}_V^W))$, such that for any finite subset $U \subset V$ and any $(\lambda_i)_{i \in U} \in \mathbb{R}_+^U$:*

$$\int e^{-\sum_{i \in U} \lambda_i \beta_i} \nu_V^W(d\beta) = e^{-\sum_{i \sim j, i, j \in U} W_{i,j}(\sqrt{(1+\lambda_i)(1+\lambda_j)}-1) - \sum_{i \sim j, i \in U, j \notin U} W_{i,j}(\sqrt{1+\lambda_i}-1)} \frac{1}{\prod_{i \in U} \sqrt{1+\lambda_i}}.$$

In particular, we have the following properties: on the probability space $(\mathcal{D}_V^W, \mathcal{B}(\mathcal{D}_V^W), \nu_V^W(d\beta))$,

- *(1-dependence) if $U, U' \subset V$ are such that $d_{\mathcal{G}}(U, U') \geq 2$, then the random variables $\beta \mapsto \beta_U$ and $\beta \mapsto \beta_{U'}$ are independent,*
- *(Reciprocal inverse Gaussian marginals) for $i \in V$, the random variable $\beta \mapsto \frac{1}{2\beta_i}$ has an inverse Gaussian distribution with parameter $(\frac{1}{W_i}, 1)$ where $W_i = \sum_{j \sim i} W_{i,j}$.*

Remark 2. *On finite graphs, the density of ν_V^W is explicit, c.f. [22, Theorem 1] and Theorem A below.*

In the sequel, the probability space $(\mathcal{D}_V^W, \mathcal{B}(\mathcal{D}_V^W), \nu_V^W)$ will be considered as the canonical space of random potentials on the graph. We write $\mathbb{E}_{\nu_V^W}$ for the expectation with respect to ν_V^W . We will introduce several random variables on this probability space, and adopt the following notation: when $\beta \mapsto X_\beta$ is a measurable function we will write X for the associated random variable and X_β for its realization on the potential β . In particular, we will write H for the random Schrödinger operator $\beta \mapsto H_\beta$ defined above. By abuse of notation, we sometimes consider β_i for $i \in V$ or β_U for $U \subset V$ as random variables (more precisely, the random variables are $\beta \mapsto \beta_i$ and $\beta \mapsto \beta_U$).

Definition 1. Let $(V_n)_{n \in \mathbb{N}}$ be an increasing sequence of finite connected subsets of V such that

$$\bigcup_{n=0}^{\infty} V_n = V.$$

For $n \in \mathbb{N}$, we define $\mathcal{F}^{(n)} \subset \mathcal{B}(\mathcal{D}_V^W)$ as the sub σ -field generated by the random variable $\beta \mapsto \beta_{V_n}$. For $n \in \mathbb{N}$ and $\beta \in \mathcal{D}_V^W$, we define a random operator $(\widehat{G}_\beta^{(n)}(i, j))_{i, j \in V}$ by

$$\widehat{G}_\beta^{(n)}(i, j) = \begin{cases} ((H_\beta)_{V_n, V_n})^{-1}(i, j), & \text{if } i, j \in V_n, \\ 0, & \text{otherwise.} \end{cases}$$

For $n \in \mathbb{N}$ and $\beta \in \mathcal{D}_V^W$, we define a random function $(\psi_\beta^{(n)}(i))_{i \in V}$ as the unique solution of the following equation:

$$\begin{cases} H_\beta(\psi_\beta^{(n)})(i) = 0, & \text{for } i \in V_n, \\ \psi_\beta^{(n)}(i) = 1, & \text{for } i \in V_n^c. \end{cases}$$

By definition, the random variables $\widehat{G}^{(n)} : \beta \mapsto \widehat{G}_\beta^{(n)}$ and $\psi^{(n)} : \beta \mapsto \psi_\beta^{(n)}$ are $\mathcal{F}^{(n)}$ -measurable.

The fact that there is a unique solution to the equation defining $\psi_\beta^{(n)}$ is elementary, see the proof in Section 4.2.

Our main theorem is the following.

Theorem 1. (i) For all $i, j \in V$, the sequence of random variables $\widehat{G}^{(n)}(i, j)$ is non-decreasing and converges a.s. to

$$\widehat{G}(i, j) := \lim_{n \rightarrow \infty} \widehat{G}^{(n)}(i, j).$$

Moreover, ν_V^W -almost surely, $0 < \widehat{G}(i, j) < \infty$ and the limit does not depend on the choice of the sequence of subsets V_n .

(ii) Under the probability ν_V^W , for all $i \in V$, $\psi^{(n)}(i)$ is a positive $\mathcal{F}^{(n)}$ -martingale. It converges a.s. to a random variable $\psi(i)$, such that $\psi(i) \geq 0$ a.s., and the limit does not depend on the choice of the increasing sequence (V_n) . Moreover, the quadratic variation of the vectorial martingale $(\psi^{(n)}(i))_{i \in V}$ is given a.s. by

$$\langle \psi(i), \psi(j) \rangle_n = \widehat{G}^{(n)}(i, j).$$

In particular, $\psi^{(n)}(i)$ is bounded in L^2 if and only if $\mathbb{E}_{\nu_V^W}(\widehat{G}(i, i)) < \infty$.

(iii) For any real $\gamma > 0$ and $\beta \in \mathcal{D}_V^W$, we define

$$G_{\beta, \gamma}(i, j) = \widehat{G}_\beta(i, j) + \frac{1}{2} \gamma^{-1} \psi_\beta(i) \psi_\beta(j).$$

For $i_0 \in V$ and $x \in V$, denote by P_x^{β, γ, i_0} the law of the Markov jump process which starts at $x \in V$ and jumps from i to j at rate

$$(2.3) \quad \frac{1}{2} W_{i,j} \frac{G_{\beta, \gamma}(i_0, j)}{G_{\beta, \gamma}(i_0, i)}.$$

Then the VRJP in exchangeable time scale, defined in section 2.1, with conductances $(W_{i,j})$ and starting from i_0 is a mixture of these Markov jump processes and has law

$$(2.4) \quad \mathbb{P}_{i_0}^{VRJP}(\cdot) = \int P_{i_0}^{\beta, \gamma, i_0}(\cdot) \nu_V^W(d\beta) \frac{\mathbb{1}_{\gamma > 0}}{\sqrt{\pi\gamma}} e^{-\gamma} d\gamma.$$

(iv) For ν_V^W -almost all β , all $\gamma > 0$ and all $i_0 \in V$, we have,

- the Markov process $P_{i_0}^{\beta, \gamma, i_0}$ is transient if and only if $\psi_\beta(j) > 0$ for all $j \in V$,
- the Markov process $P_{i_0}^{\beta, \gamma, i_0}$ is recurrent if and only if $\psi_\beta(j) = 0$ for all $j \in V$.

N.B.: Note that P_x^{β, γ, i_0} is well defined for ν_V^W -almost all β and all $\gamma > 0$ by (i) and (ii).

Notations . We denote by

$$(2.5) \quad \nu_V^W(d\beta, d\gamma) := \nu_V^W(d\beta) \otimes \frac{\mathbb{1}_{\gamma > 0}}{\sqrt{\pi\gamma}} e^{-\gamma} d\gamma$$

the probability distribution which appears in (2.4), under which γ is Gamma($\frac{1}{2}, 1$)-distributed and independent of β . In general, we simply write $G(i, j)$ for $G_{\beta, \gamma}(i, j)$ and consider it as a random variable on the probability space $(\mathcal{D}_V^W \times \mathbb{R}_+^*, \mathcal{B}(\mathcal{D}_V^W) \otimes \mathcal{B}(\mathbb{R}_+^*), \nu_V^W(d\beta, d\gamma))$.

Remark 3. When the VRJP is recurrent, $G = \widehat{G}$, and the representation of the VRJP (2.4) only involves the variable β and not γ .

Remark 4. The representation (2.3) extends to infinite graphs the representation provided in [22, Theorem 2] for finite graphs. An interesting new feature appears in the transient regime, where the generalized eigenfunction ψ and the extra gamma random variable enter the expression of $G(i, j)$. As it appears in the proof, the eigenfunction ψ can be interpreted as the mixing field of a VRJP starting from infinity.

Denote by $\tau_{i_0}^+ = \inf\{t \geq 0, Z_t = i_0, \exists s < t \text{ s.t. } Z_s \neq i_0\}$ the first return time to i_0 by $(Z_t)_{t \geq 0}$. The point (iv) of the previous theorem is in fact a consequence of the following more precise assertion.

Proposition 2. We have, for ν_V^W -almost all β , for all $\gamma > 0$ and $i_0, i \in V$,

$$P_i^{\beta, \gamma, i_0}(\tau_{i_0}^+ = \infty) = \begin{cases} \frac{\psi(i_0)^2}{4\gamma \widetilde{\beta}_{i_0} \widehat{G}(i_0, i_0) G(i_0, i_0)}, & \text{if } i = i_0, \\ \frac{\psi(i_0) \widehat{G}(i_0, i_0) \psi(i) - \widehat{G}(i_0, i) \psi(i_0)}{2\gamma \widehat{G}(i_0, i_0) G(i_0, i)}, & \text{if } i \neq i_0, \end{cases}$$

where $\widetilde{\beta}_{i_0} = \sum_{j \sim i_0} \frac{1}{2} W_{i_0, j} \frac{G(i_0, j)}{G(i_0, i_0)}$. In particular, $\psi(i_0) = 0$ if and only if $P_{i_0}^{\beta, \gamma, i_0}(\tau_{i_0}^+ = \infty) = 0$.

Using Doob's h transform, the law of the process (Z_t) conditioned on the event $\{\tau_0^+ < \infty\}$ or $\{\tau_0^+ = \infty\}$ can be computed and takes a rather nice form, both under the law $\mathbb{P}_{i_0}^{VRJP}$, or under the law $P_{i_0}^{\beta, \gamma, i_0}$ for ν_V^W -almost all β . We provide these formulae in Section 7.

A natural question that emerges from point (iv) of the theorem is that of a 0-1 law for transience/recurrence. We provide an answer below in the case of vertex transitive graphs with conductances. We say that (\mathcal{G}, W) is vertex transitive if the group of automorphisms of

\mathcal{G} that leaves invariant $(W_{i,j})$ is transitive on vertices. In particular, it is the case for the cubic lattice \mathbb{Z}^d with constant conductances $W_{i,j} = W$. Denote by \mathcal{A} the group of automorphisms that leave W invariant.

Proposition 3. *If (\mathcal{G}, W) is vertex transitive and \mathcal{G} is infinite, then under the distribution $\nu_V^W(d\beta)$, the random variables $(\beta_i)_{i \in V}$, $(\psi(i))_{i \in V}$, $(\widehat{G}(i, j))_{i, j \in V}$ are stationary and ergodic for the group of transformations \mathcal{A} . Moreover, the VRJP is either recurrent or transient, i.e.*

$$\mathbb{P}_{i_0}^{\text{VRJP}}(\text{ every vertex is visited i.o. }) = 1 \text{ or } \mathbb{P}_{i_0}^{\text{VRJP}}(\text{ every vertex is visited f.o. }) = 1.$$

In the first case $\psi(i) = 0$ for all $i \in V$, a.s., in the second case $\psi(i) > 0$ for all $i \in V$, a.s.

N.B: The action of \mathcal{A} on \widehat{G} is $(\tau\widehat{G})(i, j) = \widehat{G}(\tau i, \tau j)$ for $\tau \in \mathcal{A}$.

2.3. Relation with random Schrödinger operators. Let us now relate Theorem 1 to the properties of the random Schrödinger operator $H : \beta \mapsto H_\beta$ associated with the random potential (β_j) under the law ν_V^W , defined in (2.1) and Proposition 1.

Theorem 2. *Under $\nu_V^W(d\beta)$:*

(i) *The spectrum of H is a.s. included in $[0, \infty)$.*

(ii) *The operator \widehat{G} is the inverse of H in the following sense: for all $i, j \in V$, a.s.*

$$\widehat{G}(i, j) = \lim_{\epsilon > 0, \epsilon \rightarrow 0} (H + \epsilon)^{-1}(i, j).$$

(iii) *We have $(H\psi)(i) = 0$ a.s. for all $i \in V$.*

(iv) *In the case of the grid \mathbb{Z}^d and when $W_{i,j} = W$ is constant, $(\widehat{G}(i, j))$ and $(\psi(i))$ are stationary ergodic for the spacial shift. Moreover, in the transient case, ψ is a.s. a positive generalized eigenfunction with eigenvalue 0 in the sense that $H\psi = 0$ and ψ has at most polynomial growth. More precisely, for all $p > d$ and $C > 0$, a.s. there exists a random integer $K > 0$ such that*

$$|\psi(i)| \leq C \|i\|_\infty^p \quad \forall i \in \mathbb{Z}^d \text{ such that } \|i\|_\infty \geq K.$$

2.4. Functional central limit theorem. We denote by $(\widetilde{Z}_n)_{n \in \mathbb{N}}$ the discrete time process that describes the successive jumps of $(Z_t)_{t \in \mathbb{R}_+}$. From Theorem 1 (iii), under $\mathbb{P}_{i_0}^{\text{VRJP}}$, \widetilde{Z}_n is a mixture of Markov chains starting from i_0 and with conductances

$$(2.6) \quad W_{i,j} G(i_0, i) G(i_0, j),$$

under the probability distribution $\nu_V^W(d\beta, d\gamma)$.

We prove below a functional central limit theorem for the discrete time VRJP on \mathbb{Z}^d , $d \geq 3$, at weak reinforcement (i.e. for W large enough).

Theorem 3. *Consider the cubic graph \mathbb{Z}^d , $d \geq 3$, with constant conductances $W_{i,j} = W$. Denote*

$$\widetilde{Z}_t^{(n)} = \frac{\widetilde{Z}_{[nt]}}{\sqrt{n}}.$$

There exists $\lambda_2 > 0$ such that if $W > \lambda_2$, the discrete time VRJP satisfies a functional central limit theorem, i.e. under $\mathbb{P}_0^{\text{VRJP}}$, for any real $0 < T < \infty$, $(\widetilde{Z}_t^{(n)})_{t \in [0, T]}$ converges in law (for the Skorokhod topology) to a d -dimensional Brownian motion $(B_t)_{t \in [0, T]}$ with non degenerate isotropic diffusion matrix $\sigma^2 Id$, for some $0 < \sigma^2 < \infty$.

2.5. Consequences for the Edge Reinforced Random Walk (ERRW). The Edge Reinforced Random Walk (ERRW) is a famous discrete time process introduced in 1986 by Coppersmith and Diaconis, [4, 12].

Endow the edges of the graph $\mathcal{G} = (V, E)$ with some positive weights $(a_e)_{e \in E}$. Let $(X_n)_{n \in \mathbb{N}}$ be a random process that takes values in V , and let $\mathcal{F}_n = \sigma(X_0, \dots, X_n)$ be the filtration of its past. For any $e \in E$, $n \in \mathbb{N}$, let

$$(2.7) \quad N_n(e) = a_e + \sum_{k=1}^n \mathbb{1}_{\{\{X_{k-1}, X_k\} = e\}}$$

be the number of crossings of the (undirected) edge e up to time n plus the initial weight a_e .

Then $(X_n)_{n \in \mathbb{N}}$ is called Edge Reinforced Random Walk (ERRW) with starting point $i_0 \in V$ and weights $(a_e)_{e \in E}$, if $X_0 = i_0$ and, for all $n \in \mathbb{N}$,

$$(2.8) \quad \mathbb{P}(X_{n+1} = j \mid \mathcal{F}_n) = \mathbb{1}_{\{j \sim X_n\}} \frac{N_n(\{X_n, j\})}{\sum_{k \sim X_n} N_n(\{X_n, k\})}.$$

We denote by $\mathbb{P}_{i_0}^{ERRW}$ the law of the ERRW starting from the initial vertex i_0 . We will assume that the ERRW is defined on the canonical space $V^{\mathbb{N}}$, i.e. that $(X_n)_{n \in \mathbb{N}}$ is the canonical process on $V^{\mathbb{N}}$.

Important progress has been done in the last ten years in the understanding of this process, c.f. e.g. [1, 8, 17, 20]. In particular, it was proved in 2012 by Sabot, Tarrès, [20], and Angel, Crawford, Kozma, [1], on any graph with bounded degree at strong reinforcement (i.e. for $a_e < \tilde{\lambda}_0$ for some fixed $\tilde{\lambda}_0 > 0$) that the ERRW is a mixture of positive recurrent Markov chains. It was proved by Disertori, Sabot, Tarrès [8] that on \mathbb{Z}^d , $d \geq 3$, the ERRW is transient at weak reinforcement (i.e. for $a_e > \tilde{\lambda}_1$ for some fixed $\tilde{\lambda}_1 < \infty$).

From [20, Theorem 1], we know that the ERRW has the law of a VRJP in independent conductances. More precisely, consider $(W_e)_{e \in E}$ as independent random variables with gamma distribution with parameters $(a_e, 1)$. Consider the VRJP in conductances $(W_e)_{e \in E}$ and its underlying discrete time process (\tilde{Z}_n) . Then the annealed law of (\tilde{Z}_n) (after expectation with respect to W) is that of the ERRW (X_n) with initial weights (a_e) . Hence, we can apply Theorem 1 at fixed W and then integrate on W . We thus consider the joint law $\tilde{\nu}_V^a(dW, d\beta, d\gamma)$ on $(\mathbb{R}_+^*)^E \times (\mathbb{R}_+^*)^V \times \mathbb{R}_+^*$ obtained from $\nu_V^W(d\beta, d\gamma)$ after randomization with respect to W . More formally, let $\tilde{\nu}_V^a(dW)$ be the probability distribution on $(\mathbb{R}_+^*)^E$ such that under $\tilde{\nu}_V^a(dW)$ the random variables $W \mapsto W_e$ are independent with gamma distribution with parameters $(a_e, 1)$, then $\tilde{\nu}_V^a(dW, d\beta, d\gamma)$ is the probability distribution on $(\mathbb{R}_+^*)^E \times (\mathbb{R}_+^*)^V \times \mathbb{R}_+^*$ such that for any bounded measurable test function F ,

$$\int F(W, \beta, \gamma) \tilde{\nu}_V^a(dW, d\beta, d\gamma) = \int \left(\int F(W, \beta, \gamma) \nu_V^W(d\beta, d\gamma) \right) \tilde{\nu}_V^a(dW).$$

In the sequel, $\tilde{\nu}_V^a(dW, d\beta)$, $\tilde{\nu}_V^a(d\beta)$ will denote the corresponding marginal distributions, and $\tilde{\nu}_V^a(dW)$ is the W marginal. (By definition, $\tilde{\nu}_V^a(dW, d\beta)$ is supported on the set of (W, β) such that $\beta \in \mathcal{D}_V^W$.) From Theorem 1, we see that the ERRW starting from i_0 is a mixture of reversible Markov chains with conductances

$$(2.9) \quad x_{i,j} = W_{i,j} G(i_0, i) G(i_0, j),$$

where G is defined in Theorem 1, and (W, β, γ) are distributed according to $\tilde{\nu}_V^a(dW, d\beta, d\gamma)$. More formally, if $\tilde{P}_{i_0}^x$ denotes the law of the Markov chain starting at i_0 and with conductances

$(x_{i,j})_{i \sim j}$, then

$$\mathbb{P}_{i_0}^{ERRW}(\cdot) = \int \tilde{P}_{i_0}^x(\cdot) \tilde{\nu}_V^a(dW, d\beta, d\gamma).$$

An important point is that we keep the 1-dependence of the field β , after taking expectation with respect to W .

Proposition 4. *Under $\tilde{\nu}_V^a(d\beta)$, $(\beta_j)_{j \in V}$ is 1-dependent: if $U, U' \subset V$ are such that $d_G(U, U') \geq 2$, then $(\beta_i)_{i \in U}$ and $(\beta_j)_{j \in U'}$ are independent.*

Proof. Indeed, from Proposition 1, the Laplace transform of $(\beta_i)_{i \in U}$ under $\nu_V^W(d\beta)$ only involves the conductances $W_{i,j}$ for i or j in U . This implies that, if $d_G(U, U') \geq 2$, the joint Laplace transform of $(\beta_i)_{i \in U}$ and $(\beta_i)_{i \in U'}$ is still the product of Laplace transforms even after taking expectation with respect to the random variables (W_e) , i.e. under $\tilde{\nu}_V^a(d\beta)$. \square

This yields a counterpart of Proposition 3 for the ERRW.

Proposition 5. *Assume $(\mathcal{G}, (a_{i,j}))$ is vertex transitive with automorphism group \mathcal{A} , and \mathcal{G} infinite. Then under the distribution $\tilde{\nu}_V^a(dW, d\beta)$, the random variables $(W_e)_{e \in E}$, $(\beta_i)_{i \in V}$, $(\psi(i))_{i \in V}$, $(\widehat{G}(i, j))_{i, j \in V}$ are stationary and ergodic for the group of transformations \mathcal{A} . Moreover, the ERRW is either recurrent or transient, i.e.*

$$\mathbb{P}_{i_0}^{ERRW}(\text{every vertex is visited i.o.}) = 1, \text{ or } \mathbb{P}_{i_0}^{ERRW}(\text{every vertex is visited f.o.}) = 1.$$

In the first case $\psi(i) = 0$ for all $i \in V$, a.s., in the second case $\psi(i) > 0$ for all $i \in V$, a.s.

N.B: The action of \mathcal{A} on \widehat{G} and W is $(\tau\widehat{G})(i, j) = \widehat{G}(\tau i, \tau j)$, $\tau W_{i,j} = W_{\tau i, \tau j}$ for $\tau \in \mathcal{A}$.

Remark 5. *In [16], it was proved on infinite graphs that the ERRW is a mixture of Markov chains, obtained as a weak limit of the mixing law of the ERRW on finite approximating graphs. The difference in the representation we give in (2.9) is that the random variables ψ , \widehat{G} are obtained as almost sure limits and hence are measurable functions of the random variables β . This yields stationarity and ergodicity, which are the key ingredients in the 0-1 law, and in forthcoming Theorems 4 and 5.*

Remark 6. *It seems that this 0-1 law is new, both for the VRJP and the ERRW. In [16], it was proved that if the ERRW comes back with probability 1 to its starting point then it visits infinitely often all points, a.s., which is a weaker result. This was proved using the representation of the ERRW as mixture of Markov chains of [16]. (A short proof of this last result can also be given, c.f. [23].)*

We now give a counterpart of Theorem 3 for the ERRW. It is a consequence of Theorem 1 and of the delocalization result proved by Disertori, Sabot, Tarrès in [8].

Theorem 4. *Consider the cubic graph \mathbb{Z}^d , $d \geq 3$, with constant weights $a_{i,j} = a$. Denote*

$$X_t^{(n)} = \frac{X_{[nt]}}{\sqrt{n}}.$$

There exists $\tilde{\lambda}_2 > 0$ such that if $a > \tilde{\lambda}_2$, the ERRW satisfies a functional central limit theorem, i.e. under \mathbb{P}_0^{ERRW} , for any real $0 < T < \infty$, $(X_t^{(n)})_{t \in [0, T]}$ converges in law (for the Skorokhod topology) to a d -dimensional Brownian motion $(B_t)_{t \in [0, T]}$ with non degenerate isotropic diffusion matrix $\sigma^2 Id$, for some $0 < \sigma^2 < \infty$.

Finally, we can deduce recurrence of the ERRW in dimension 2 from Theorem 1, Proposition 5 and the estimates obtained by Merkl and Rolles in [15, 17]¹.

Theorem 5. *The ERRW $(X_n)_{n \geq 0}$ on \mathbb{Z}^2 with constant weights $a_{i,j} = a$ is a.s. recurrent, i.e.*

$$\mathbb{P}_0^{ERRW}(\text{every vertex is visited infinitely often}) = 1.$$

In [15, 17], by a Mermin-Wagner type argument, Merkl and Rolles proved a polynomial decrease of the form

$$(2.10) \quad \mathbb{E} \left(\left(\frac{x_\ell}{x_0} \right)^{\frac{1}{4}} \right) \leq c(a) |\ell|^{-\xi(a)},$$

for some constants $c(a) > 0$, $\xi(a) > 0$, depending only on a , and where x_ℓ is the conductance at the site ℓ for the mixing measure of the ERRW, uniformly for a sequence of finite approximating graphs. When $0 < \xi < 1$, it does not give by itself enough information to prove recurrence. It was used in the case of a diluted 2-dimensional graph to prove positive recurrent at strong reinforcement. The extra information given by the representation (2.9) and the stationarity of ψ , implies that the polynomial estimate (2.10) is incompatible with $\psi(i) > 0$ and hence is incompatible with transience. Detailed arguments are provided in Section 8.

Remark 7. *We expect similarly that the 2-dimensional VRJP with constant conductances $W_{i,j} = W > 0$ is recurrent. This would be implied by an estimate of the type (2.10) for the mixing field of the VRJP, which is still not available. More precisely, we can see from the proof of Theorem 5 in Section 8, that recurrence of the 2-dimensional VRJP would be implied by Theorem 1, Proposition 3, and an estimate of the type*

$$\mathbb{E} (e^{\eta(u_\ell - u_0)}) \leq \epsilon(\|\ell\|_\infty),$$

for $\eta > 0$ and $\epsilon(n)$ a positive function such that $\lim_{n \rightarrow \infty} \epsilon(n) = 0$, where (u_j) is the mixing field of the VRJP starting from 0 (c.f. Theorem B) on finite boxes with wired boundary condition as in Section 4.2. We learned from G. Kozma and R. Peled that they have a proof of such an estimate.

2.6. Open questions. The most important question certainly concerns the relation between the properties of the VRJP and the spectral properties of the random Schrödinger operator H_β . For example on \mathbb{Z}^d with constant weights $W_{i,j} = W$, is recurrence/transience of the VRJP related to the localized/delocalized regimes of H_β ? A more precise question would be: does the transient regime of the VRJP coincide with the existence of extended states at least at the bottom of the spectrum of H_β ? It might at first seem inconsistent to expect extended states at the bottom of the spectrum since the Anderson model with i.i.d. potential is expected to be localized at the edges of the spectrum (a fact which is proved in several cases). But this localization is a consequence of Lifshitz tails, and there are good reasons to expect that Lifshitz tails fail for the potential β , which is not i.i.d. but 1-dependent. Indeed, the bottom of the spectrum of H_β is 0, it does not coincide with the minimum of the support of the distribution of 2β translated by the spectrum of $-P$, as it is the case for i.i.d. potential. In fact, on a finite set, the minimum of the spectrum is reached on the set $\det(2\beta - P) = 0$ which is a set of codimension 1, hence it is "big".

¹We are grateful to Franz Merkl and Silke Rolles for a useful discussion on that subject

Another natural question concerns the uniform integrability of the martingale $\psi^{(n)}(i)$. Let us ask a more precise question: is it true (at least for \mathbb{Z}^d with constant weights) that transience of the VRJP implies that the martingale $\psi^{(n)}(i)$ is bounded in L^2 ? It is quite natural to expect such a property from relation (5.2) since $\widehat{G}^{(n)}(i, i)$ appears to be the quadratic variation of $\psi^{(n)}(i)$. This would have several consequences. Firstly, it would imply that in dimension $d \geq 3$, the VRJP satisfies a functional central limit theorem as soon as the VRJP is transient, by the same argument as that of the proof of Theorem 3. It would also imply directly that the VRJP is recurrent as soon as the reversible Markov chain in conductances $(W_{i,j})$ is recurrent, if the group of automorphisms of (\mathcal{G}, W) is transitive. Indeed, assume that the property is true and the VRJP is transient. By Theorem 1, the discrete time process (\widetilde{Z}_n) would be represented as a mixture of reversible Markov chains with conductances $W_{i,j}G(0, i)G(0, j)$. From Proposition 2 applied to $i_0 = 0$, we have that

$$\frac{\widehat{G}(0, i)}{\widehat{G}(0, 0)} \leq \frac{\psi(i)}{\psi(0)}.$$

Hence, (\widetilde{Z}_n) is equivalently a mixture of Markov chains with conductances

$$\frac{\psi(0)^2}{G(0, 0)^2} W_{i,j} G(0, i) G(0, j) \leq W_{i,j} \psi(i) \psi(j).$$

But $(\psi(i))$ is stationary ergodic, if ψ_0 is squared integrable, we would have

$$\mathbb{E}_{\nu_V^W}(W_{i,j} \psi(i) \psi(j)) \leq C W_{i,j}$$

for some constant $C > 0$. Usual arguments imply that the Markov chain in conductance $W_{i,j} \psi(i) \psi(j)$ is recurrent if the Markov chain in conductances $(W_{i,j})$ is recurrent (c.f. e.g. Exercise 2.75, [14]). We arrive at a contradiction.

2.7. Organization of the paper. In Section 3, we gather several results in the case of finite graphs, in particular we recall the main results of [22]. In Section 4, we define the important notion of restriction with wired boundary condition and the compatibility property. Section 5 is the key step in the paper where the martingale property is proved. In Section 6, we prove Theorem 1, Propositions 2 and 3 and Theorem 2. In Section 7, we provide extra computations of h -transforms. In section 8, we prove recurrence of ERRW in dimension 2 for all initial constant weights. In Section 9, we prove functional central limit theorems for the VRJP and the ERRW, Theorems 3 and 4.

3. THE RANDOM POTENTIAL β ON FINITE GRAPHS

In this section we assume that $\mathcal{G} = (V, E)$ is a finite graph and gather several results in this case. Recall that every undirected edge $e = \{i, j\}$ is labeled with a positive conductance $W_e = W_{i,j}$. In the case of a finite graph, the Schrödinger operator H_β defined in (2.1) can be represented by the $V \times V$ -matrix given by

$$H_\beta(i, j) = \begin{cases} 2\beta_i, & i = j, \\ -W_{i,j}, & i \neq j, i \sim j, \\ 0, & \text{otherwise,} \end{cases}$$

and the set \mathcal{D}_V^W defined in (2.2) is the set of potentials β such that H_β is positive definite.

3.1. The probability distribution ν_V^W on finite graphs and relation to the VRJP. We recall [22, Theorem 1], which defines the probability distribution $\nu_V^W(d\beta)$ by its density on any finite graph.

Theorem A (Theorem 1, Definition 1 and Proposition 1 of [22]). *Let $(\mathcal{G}, (W_e)_{e \in E})$ be a finite graph with conductances. The measure below is a probability on \mathcal{D}_V^W :*

$$(3.1) \quad \nu_V^W(d\beta) := \mathbf{1}_{H_\beta > 0} \left(\frac{2}{\pi} \right)^{|V|/2} \exp\left(-\sum_{i \in V} \beta_i + \sum_{e \in E} W_e\right) \frac{d\beta_V}{\sqrt{\det H_\beta}}$$

with $d\beta_V = \prod_{i \in V} d\beta_i$, and where $H_\beta > 0$ means that H_β is positive definite.

The Laplace transform of the probability distribution $\nu_V^W(d\beta)$ is given, for all $(\lambda_i) \in \mathbb{R}_+^V$, by

$$(3.2) \quad \int e^{-\langle \lambda, \beta \rangle} \nu_V^W(d\beta) = \exp\left(-\sum_{i \sim j} W_{i,j} (\sqrt{(\lambda_i + 1)(\lambda_j + 1)} - 1)\right) \prod_{i \in V} \frac{1}{\sqrt{\lambda_i + 1}}.$$

Moreover, we have the following properties: under $\nu_V^W(d\beta)$,

- (1-dependence): if $U, U' \subset V$ are such that $d_{\mathcal{G}}(U, U') \geq 2$, then the random variables $\beta \mapsto \beta_U$ and $\beta \mapsto \beta_{U'}$ are independent,
- (Reciprocal inverse Gaussian marginals) for $i \in V$, the random variable $\beta \mapsto \frac{1}{2\beta_i}$ has an inverse Gaussian distribution with parameter $(\frac{1}{W_i}, 1)$ where $W_i = \sum_{j \sim i} W_{i,j}$.

If we apply formula (3.2) to $(\lambda_i) \in \mathbb{R}_+^V$ such that $\lambda_{V \setminus U} = 0$ for a subset $U \subset V$, we find the expression of Proposition 1. Hence, it implies Proposition 1 in the case of a finite graph.

The field β is closely related to the VRJP, as shown in the next two theorems. In [20], it is shown that the VRJP in exchangeable time scale defined in Section 2.1 is a mixture of Markov jump processes, more precisely:

Theorem B (Theorem 2 of [20]). *Assume V finite. The following measure is a probability distribution on the set $\{(u_i)_{i \in V} \in \mathbb{R}^V, u_{i_0} = 0\}$:*

$$(3.3) \quad \mathcal{Q}_{i_0}^W(du) = \frac{1}{\sqrt{2\pi}^{|V|-1}} \exp\left(-\sum_{i \in V} u_i - \sum_{i \sim j} W_{i,j} (\cosh(u_i - u_j) - 1)\right) \sqrt{D(W, u)} du_{V \setminus \{i_0\}}$$

where $du_{V \setminus \{i_0\}} = \prod_{i \in V \setminus \{i_0\}} du_i$ and $D(W, u) = \sum_{T \in \mathcal{T}} \prod_{\{i,j\} \in T} W_{i,j} e^{u_i + u_j}$, where the sum is over \mathcal{T} , the set of spanning trees of the graph \mathcal{G} .

For $(u_i)_{i \in V} \in \mathbb{R}^V$, we denote by $P_{i_0}^{(u)}$ the law of the Markov jump process starting at vertex i_0 and with jump rates from i to j given by

$$\frac{1}{2} W_{i,j} e^{u_j - u_i}.$$

The law of the VRJP in exchangeable time scale starting at i_0 is a mixture of Markov jump processes, with mixing law given by

$$\mathbb{P}_{i_0}^{\text{VRJP}}(\cdot) = \int P_{i_0}^{(u)}(\cdot) \mathcal{Q}_{i_0}^W(du).$$

Remark 8. By the matrix-tree theorem, $D(W, u)$ is any diagonal minor of the $|V| \times |V|$ matrix $(m_{i,j})$ with coefficients

$$m_{i,j} = \begin{cases} 0, & \text{if } i \not\sim j, i \neq j, \\ -W_{i,j}e^{u_i+u_j}, & \text{if } i \sim j, i \neq j, \\ \sum_{k \in V, k \sim i} W_{i,k}e^{u_i+u_k}, & \text{if } i = j. \end{cases}$$

Remark 9. The probability measure $\mathcal{Q}_{i_0}^W(du)$ appeared previously to [20] in a rather different context in the work of Disertori, Spencer, Zirnbauer, [10]. In particular, the fact that $\mathcal{Q}_{i_0}^W(du)$ is a probability measure was proved there as a consequence of a Berezin identity applied to a supersymmetric extension of that measure.

On finite graphs, the random environment (u_i) of the previous theorem can be represented thanks to the Green function of the random potential $(\beta_i, i \in V)$ distributed according to $\nu_V^W(d\beta)$. Let us first recall [22, Proposition 1].

Proposition A (Proposition 1 of [22]). Assume V finite. For $\beta \in \mathcal{D}_V^W$, we denote by

$$G_\beta := (H_\beta)^{-1}$$

the Green function of the Schrödinger operator H_β . For $\beta \in \mathcal{D}_V^W$, $i, j \in V$, we define $u_\beta(i, j)$ by

$$(3.4) \quad e^{u_\beta(i,j)} = \frac{G_\beta(i, j)}{G_\beta(i, i)}.$$

For $i_0 \in V$, $(u_\beta(i_0, j))_{j \in V}$ is the unique solution of the equation

$$(3.5) \quad \begin{cases} u_\beta(i_0, i_0) = 0, \\ \sum_{j \sim i} W_{i,j}e^{u_\beta(i_0,j)-u_\beta(i_0,i)} = \beta_i, & \text{if } i \neq i_0. \end{cases}$$

In particular, the function $\beta \mapsto (u_\beta(i_0, j))_{j \in V}$ is $(\beta_j)_{j \in V \setminus \{i_0\}}$ -measurable. Moreover, for all $\beta \in \mathcal{D}_V^W$,

$$(3.6) \quad \beta_{i_0} = \frac{1}{2G_\beta(i_0, i_0)} + \frac{1}{2} \sum_{j \sim i_0} W_{i_0,j}e^{u_\beta(i_0,j)-u_\beta(i_0,i_0)}.$$

As usual, we simply denote by $G(i, j)$ and $u(i, j)$ the associated random variables on the probability space $(\mathcal{D}_V^W, \mathcal{B}(\mathcal{D}_V^W), \nu_V^W)$. Let us now recall [22, Theorem 3].

Theorem C (Theorem 3 of [22]). Assume V finite. For all $i_0 \in V$, under the probability $\nu_V^W(d\beta)$,

- (i) the random field $(u(i_0, j))_{j \in V}$ has the distribution $\mathcal{Q}_{i_0}^W$ of Theorem B,
- (ii) $\frac{1}{2G(i_0, i_0)}$ has a gamma distribution with parameters $(1/2, 1)$,
- (iii) $G(i_0, i_0)$ is independent of $(\beta_j)_{j \neq i_0}$, hence independent of the field $(u(i_0, j))_{j \in V}$.

Remark 10. Here we only consider the VRJP with initial local time 1, in fact, the above correspondence between β and VRJP still holds for the process starting with any positive local times $(\phi_i, i \in V)$, in such case, there is a corresponding density ν_V^{W, ϕ^2} , which is defined in [22], see Definition 1 and Theorem 3. We choose here to normalize the initial local time to 1 since it is equivalent to the general case by a change of time and W , see [22] Appendix B.

Combining Theorem B and Theorem C, it gives a representation of the VRJP in exchangeable time scale starting from different points in terms of the probability on random potentials ν_V^W . We state this representation below.

Corollary 1. *Assume V finite. For $\beta \in \mathcal{D}_V^W$, define P_x^{β, i_0} as the law of the Markov jump process starting from x and with jump rates from i to j given by*

$$\frac{1}{2} W_{i,j} \frac{G_\beta(i_0, j)}{G_\beta(i_0, i)}.$$

Then, the VRJP in exchangeable time scale is a mixture of these Markov jump processes,

$$(3.7) \quad \mathbb{P}_{i_0}^{VRJP}(\cdot) = \int P_{i_0}^{\beta, i_0}(\cdot) \nu_V^W(d\beta).$$

3.2. Representation as a sum on paths. We call *path* in \mathcal{G} from i to j a finite sequence $\sigma = (\sigma_0, \dots, \sigma_m)$ in V such that $\sigma_0 = i$, $\sigma_m = j$ and $\sigma_k \sim \sigma_{k+1}$, for $k = 0, \dots, m-1$. The length of σ is defined by $|\sigma| = m$. We denote by $\mathcal{P}_{i,j}^V$ be the collection of paths in V from i to j , and $\bar{\mathcal{P}}_{i,j}^V$ be the collection of paths $\sigma = (\sigma_0 = i, \dots, \sigma_m = j)$ in V from i to j such that $\sigma_k \neq j, k = 0, \dots, m-1$. For a path σ and for $\beta \in \mathcal{D}_V^W$, we set

$$(3.8) \quad W_\sigma = \prod_{k=0}^{m-1} W_{\sigma_k, \sigma_{k+1}}, \quad (2\beta)_\sigma = \prod_{k=0}^m (2\beta_{\sigma_k}), \quad (2\beta)_\sigma^- = \prod_{k=0}^{m-1} (2\beta_{\sigma_k}).$$

For the trivial path $\sigma = (\sigma_0)$, we define $W_\sigma = 1$, $(2\beta)_\sigma = 2\beta_{\sigma_0}$, $(2\beta_\sigma)^- = 1$. (Note that these definitions make sense also in the case of infinite graphs.)

The following representation of the Green function $G(\cdot, \cdot)$ as a sum on paths will be convenient.

Proposition 6. *Assume that V is finite. For all $\beta \in \mathcal{D}_V^W$, we have, with the notations of Theorem A,*

$$(3.9) \quad G_\beta(i, j) = \sum_{\sigma \in \mathcal{P}_{i,j}^V} \frac{W_\sigma}{(2\beta)_\sigma}, \quad \exp(u_\beta(i, j)) = \sum_{\sigma \in \bar{\mathcal{P}}_{j,i}^V} \frac{W_\sigma}{(2\beta)_\sigma^-}.$$

Proof. Write D_β for the diagonal $V \times V$ matrix with $(\beta_i)_{i \in V}$ as diagonal coefficients, then $H_\beta = (\text{Id} - PD_\beta^{-1})D_\beta$. Since $H_\beta > 0$, by Perron-Frobenius theorem, we have that $\rho(PD_\beta^{-1}) < 1$, where $\rho(PD_\beta^{-1})$ is the spectral radius of PD_β^{-1} . Hence, we can write the following convergent expansion,

$$G_\beta = H_\beta^{-1} = D_\beta^{-1} \sum_{k=0}^{\infty} (PD_\beta^{-1})^k,$$

which exactly corresponds to (3.9).

For the expansion of $\exp(u_\beta(i, j))$, note first that $\sum_{\sigma \in \bar{\mathcal{P}}_{j,i}^V} \frac{W_\sigma}{\beta_\sigma} \leq \beta_i G_\beta(j, i) < \infty$. A path in $\bar{\mathcal{P}}_{j,i}^V$ can be cut at its first visit to i , turning it into the concatenation of a path in $\bar{\mathcal{P}}_{j,i}^V$ and a path in $\mathcal{P}_{i,i}^V$, and this operation is bijective. It implies that

$$(3.10) \quad \left(\sum_{\sigma \in \bar{\mathcal{P}}_{j,i}^V} \frac{W_\sigma}{(2\beta)_\sigma^-} \right) G_\beta(i, i) = \left(\sum_{\sigma \in \bar{\mathcal{P}}_{j,i}^V} \frac{W_\sigma}{(2\beta)_\sigma^-} \right) \left(\sum_{\sigma \in \mathcal{P}_{i,i}^V} \frac{W_\sigma}{(2\beta)_\sigma} \right) = \sum_{\sigma \in \mathcal{P}_{j,i}^V} \frac{W_\sigma}{(2\beta)_\sigma} = G_\beta(i, j),$$

hence the result. \square

3.3. **A priori estimates on $\mathcal{Q}_{i_0}^W(du)$.** The following proposition is borrowed from [10, Lemma 3]. For convenience, we give a shorter proof of that estimate based on spanning trees instead of fermionic variables, following the proof of the corresponding result for the ERRW, c.f. [8, Lemma 7].

Proposition 7. *Let $(\mathcal{G} = (V, E), W)$ be a finite graph with conductances. Fix a vertex i_0 . Let $\eta > 0$ and let $e_1 = \{\underline{e}_1, \overline{e}_1\}, \dots, e_K = \{\underline{e}_K, \overline{e}_K\}$ be K distinct undirected edges such that $W_{e_k} \geq 2\eta$ for all $k = 1, \dots, K$. Then*

$$\int \exp\left(\eta \sum_{k=1}^K \cosh(u_{\overline{e}_k} - u_{\underline{e}_k})\right) \mathcal{Q}_{i_0}^W(du) \leq e^{\eta K} 2^{K/2}$$

where $\mathcal{Q}_{i_0}^W(du)$ is the probability distribution defined in Theorem B.

Proof. We remind that $\mathcal{Q}_{i_0}^W(du)$ is defined by

$$\mathcal{Q}_{i_0}^W(du) = \frac{1}{\sqrt{2\pi}^{|V|-1}} \exp\left(-\sum_i u_i - \sum_{i \sim j} W_{i,j}(\cosh(u_i - u_j) - 1)\right) \sqrt{D(W, u)} du_{V \setminus \{i_0\}},$$

with $du_{V \setminus \{i_0\}} = \prod_{i \neq i_0} du_i$ and $D(W, u) = \sum_{T \in \mathcal{T}} \prod_{\{i,j\} \in T} W_{i,j} e^{u_i + u_j}$ where the sum is on spanning trees.

Let $\widetilde{W} = W - \eta \sum_{k=1}^K \mathbb{1}_{e_k}$, (i.e. \widetilde{W} is equal to $W - \eta$ on the edges e_1, \dots, e_K , and unchanged on the other edges). By assumption, we have $\widetilde{W}_{i,j} > 0$ on the edges, and for all spanning trees T , since edges appear at most once:

$$\prod_{\{i,j\} \in T} W_{i,j} e^{u_i + u_j} \leq \left(\prod_{k=1}^K \frac{W_{e_k}}{W_{e_k} - \eta} \right) \prod_{\{i,j\} \in T} \widetilde{W}_{i,j} e^{u_i + u_j} \leq 2^K \prod_{\{i,j\} \in T} \widetilde{W}_{i,j} e^{u_i + u_j},$$

which implies $D(W, u) \leq 2^K D(\widetilde{W}, u)$. From the expression of $\mathcal{Q}_{i_0}^W(du)$, we deduce that

$$\exp\left(\eta \sum_{k=1}^K \cosh(u_{\overline{e}_k} - u_{\underline{e}_k})\right) \mathcal{Q}_{i_0}^W(du) \leq e^{\eta K} 2^{K/2} \mathcal{Q}_{i_0}^{\widetilde{W}}(du).$$

It implies that

$$\int \exp\left(\eta \sum_{k=1}^K \cosh(u_{\overline{e}_k} - u_{\underline{e}_k})\right) \mathcal{Q}_{i_0}^W(du) \leq e^{\eta K} 2^{K/2} \int \mathcal{Q}_{i_0}^{\widetilde{W}}(du) = e^{\eta K} 2^{K/2}.$$

□

4. THE WIRED BOUNDARY CONDITION AND KOLMOGOROV EXTENSION TO INFINITE GRAPHS

4.1. **Restriction with wired boundary condition.** Our objective is to extend the relations between the VRJP and the β field to the case of infinite graphs. To this end, we need an appropriate boundary condition, which turns out to be the wired boundary condition.

Definition 2. *Let $\mathcal{G} = (V, E)$ be a connected graph with finite degree at each site, and V_1 a strict finite subset of V . We define the restriction of \mathcal{G} to V_1 with wired boundary condition as the graph $\mathcal{G}_1 = (\widetilde{V}_1 = V_1 \cup \{\delta\}, E_1)$ where δ is an extra point and*

$$E_1 = \{\{i, j\} \in E, \text{ s.t. } i \in V_1, j \in V_1, i \sim j\} \cup \{\{i, \delta\}, i \in V_1 \text{ s.t. } \exists j \notin V_1, i \sim j\}.$$

If $(W_{i,j})_{\{i,j\} \in E}$ is a set of positive conductances, we define $(W_{i,j}^{(1)})_{\{i,j\} \in E_1}$ as the set of restricted conductances by

$$\begin{cases} W_{i,j}^{(1)} = W_{i,j}, & \text{if } i, j \in V_1, \{i, j\} \in E_1, \\ W_{i,\delta}^{(1)} = \sum_{j \notin V_1, j \sim i} W_{i,j}, & \text{if } \{i, \delta\} \in E_1, \\ 0, & \text{otherwise.} \end{cases}$$

Remark 11. *Intuitively, this restriction corresponds to identifying all points in $V \setminus V_1$ to a single point δ and to delete the edges connecting points of $V \setminus V_1$. The new weights are obtained by summing the weights of the edges identified by this procedure.*

The following lemma is fundamental and is the justification for the choice of this notion of restriction.

Lemma 1. *Let $(\mathcal{G} = (V, E), W)$ be a finite graph with conductances and ν_V^W the associated distribution on random potentials defined in Theorem A. Let V_1 be a strict subset of V and $(\mathcal{G}_1 = (\tilde{V}_1, E_1), W^{(1)})$ be the restriction of (\mathcal{G}, W) to V_1 with wired boundary condition. Let $\nu_{\tilde{V}_1}^{W^{(1)}}$ be the distribution of random potential associated with $(\mathcal{G}_1, W^{(1)})$. We denote by $(\nu_V^W)_{|V_1}$ and $(\nu_{\tilde{V}_1}^{W^{(1)}})_{|V_1}$ the marginal distributions on V_1 of ν_V^W and $\nu_{\tilde{V}_1}^{W^{(1)}}$. Then*

$$(\nu_V^W)_{|V_1} = (\nu_{\tilde{V}_1}^{W^{(1)}})_{|V_1}.$$

Remark 12. *Note that there is no such compatibility relation with the more usual notion of restriction of graph. The wired boundary condition is fundamental and in fact will be responsible for the extra gamma random variable that appears in the representation of the VRJP on the infinite graph.*

Proof. Taking $(\lambda_i)_{i \in V} \in \mathbb{R}_+^V$ such that $\lambda_{V \setminus V_1} = 0$ in Theorem A, we get that

$$\begin{aligned} & \int e^{-\sum_{i \in V_1} \lambda_i \beta_i} \nu_V^W(d\beta) \\ &= \exp \left(- \sum_{i \sim j, i, j \in V_1} W_{i,j} (\sqrt{(1 + \lambda_i)(1 + \lambda_j)} - 1) - \sum_{i \sim j, i \in V_1, j \notin V_1} (W_{i,j} (\sqrt{1 + \lambda_i} - 1)) \right) \prod_{i \in V_1} \frac{1}{\sqrt{1 + \lambda_i}}. \end{aligned}$$

Applying Theorem A to the graph \mathcal{G}_1 with $(\lambda_i)_{i \in \tilde{V}_1} \in \mathbb{R}_+^{\tilde{V}_1}$ such that $\lambda_\delta = 0$, we get

$$\begin{aligned} & \int e^{-\sum_{i \in V_1} \lambda_i \beta_i} \nu_{\tilde{V}_1}^{W^{(1)}}(d\beta) \\ &= \exp \left(- \sum_{i \sim j, i, j \in V_1} W_{i,j}^{(1)} (\sqrt{(1 + \lambda_i)(1 + \lambda_j)} - 1) - \sum_{i \in V_1, i \overset{\mathcal{G}_1}{\sim} \delta} (W_{i,\delta}^{(1)} (\sqrt{1 + \lambda_i} - 1)) \right) \prod_{i \in V_1} \frac{1}{\sqrt{1 + \lambda_i}}. \end{aligned}$$

By definition of $W_{i,j}^{(1)}$, these Laplace transforms are equal, hence the marginal distributions are equal. \square

4.2. Kolmogorov extension: proof of Proposition 1 and Definition 1. Let $\mathcal{G} = (V, E)$ be a connected infinite graph with finite degree at each site with conductances $(W_{i,j})$. We remind that $(V_n)_{n \geq 1}$ is an increasing sequence of finite strict subsets of V that exhausts V , i.e. $\cup_n V_n = V$.

Let $\mathcal{G}_n = (\tilde{V}_n = V_n \cup \{\delta_n\}, E_n)$ be the restriction of \mathcal{G} to V_n with wired boundary condition, and $(W^{(n)})$ the restricted conductances. By construction, if $n < m$, then $(\mathcal{G}_n, W^{(n)})$ is the restriction of $(\mathcal{G}_m, W^{(m)})$ to V_n with wired boundary condition. Lemma 1 implies that the sequence of marginal distributions $(\nu_{\tilde{V}_n}^{W^{(n)}})_{|V_n}$ is a compatible sequence of probabilities. By Kolmogorov extension theorem, it implies that there exists a probability measure ν_V^W such that

$$(\nu_V^W)_{|V_n} = (\nu_{\tilde{V}_n}^{W^{(n)}})_{|V_n},$$

for all integer n . By Theorem A, $\nu_V^W(d\beta)$ is supported by the set of potentials β such that $(H_\beta)_{V_n, V_n}$ is positive definite for all integers n , hence by \mathcal{D}_V^W . It also implies the other properties of $\nu_V^W(d\beta)$.

The solution of the equation defining $\psi_\beta^{(n)}$ in Definition 1 exists and is unique since it is equivalent to $(\psi_\beta^{(n)})_{V_n^c} = 1$ and

$$(4.1) \quad (H_\beta)_{V_n, V_n}(\psi_\beta^{(n)})_{V_n}(i) = \sum_{j \sim i, j \in V_n^c} W_{i,j}, \quad \text{for } i \in V_n.$$

Since $(H_\beta)_{V_n, V_n}$ is positive definite for $\beta \in \mathcal{D}_V^W$, it defines $\psi_\beta^{(n)}$ uniquely.

4.3. Coupling lemma. Definition of $G^{(n)}$, and relations with $\widehat{G}^{(n)}$, $\psi^{(n)}$ and γ . Consider the probability $\nu_V^W(d\beta, d\gamma)$ defined in (2.5). It will be convenient to couple the measure $\nu_V^W(d\beta, d\gamma)$ and the measure $\nu_{\tilde{V}_n}^{W^{(n)}}(d\beta)$ in the following way.

Lemma 2. For $\beta \in \mathcal{D}_V^W$ and $\gamma > 0$, we define $\beta^{(n)} \in \mathbb{R}^{\tilde{V}_n}$ by

$$(4.2) \quad \beta_{V_n}^{(n)} = \beta_{V_n}, \quad \beta_{\delta_n}^{(n)} = \sum_{j \in V_n, j \sim \delta_n} \frac{1}{2} W_{j, \delta_n} \psi_\beta^{(n)}(j) + \gamma.$$

Then, $\beta^{(n)} \in \mathcal{D}_{\tilde{V}_n}^{W^{(n)}}$ and under $\nu_V^W(d\beta, d\gamma)$, $\beta^{(n)}$ is distributed according to $\nu_{\tilde{V}_n}^{W^{(n)}}$.

Let $H_{\beta^{(n)}}^{(n)}$ be the Schrödinger operator associated with \mathcal{G}_n , $W^{(n)}$ and potential $\beta^{(n)}$. Let $G_{\beta^{(n)}}^{(n)} = (H_{\beta^{(n)}}^{(n)})^{-1}$, be its Green function. Then,

$$G_{\beta^{(n)}}^{(n)}(\delta_n, \delta_n) = \frac{1}{2\gamma},$$

and, for all $i \in V_n$,

$$\psi_\beta^{(n)}(i) = \frac{G_{\beta^{(n)}}^{(n)}(\delta_n, i)}{G_{\beta^{(n)}}^{(n)}(\delta_n, \delta_n)} = e^{u_{\beta^{(n)}}^{(n)}(\delta_n, i)},$$

where $u_{\beta^{(n)}}^{(n)}$ is the field defined in Proposition A for the graph \mathcal{G}_n and with the potential $\beta^{(n)}$.

As usual, we often omit the subscript β and write $H^{(n)}$, $G^{(n)}$, $u^{(n)}$, and consider them as random variables on $\mathcal{D}_V^W \times \mathbb{R}_+^*$ under $\nu_V^W(d\beta, d\gamma)$.

Proof. Let $\beta \in \mathcal{D}_V^W$ and $\gamma > 0$. Denote in this proof by $(u(j))_{j \in \tilde{V}_n}$ the vector defined by

$$u(j) = \begin{cases} 0, & \text{if } j = \delta_n, \\ \log \psi_\beta^{(n)}(j), & \text{if } j \in V_n. \end{cases}$$

Then, by definition of $\psi_\beta^{(n)}$ and $\beta^{(n)}$, we have $(H_{\beta^{(n)}}^{(n)}(e^u))_{V_n} = (H_\beta(\psi_\beta^{(n)}))_{V_n} = 0$ and

$$H_{\beta^{(n)}}^{(n)}(e^u)(\delta_n) = 2\beta_{\delta_n}^{(n)} - \sum_{j \in V_n, j \sim \delta_n} W_{\delta_n, j}^{(n)} \psi_\beta^{(n)}(j) = 2\gamma.$$

Since $(e^{u(j)})$ is a vector with positive coefficients, by general results on symmetric M-matrices, see Theorem 2.7 page 141 of [19], it implies that $H_{\beta^{(n)}}^{(n)} > 0$. Moreover, it implies that

$$\frac{1}{2\gamma} e^{u(\cdot)} = G_{\beta^{(n)}}^{(n)}(\delta_n, \cdot), \text{ hence that } G_{\beta^{(n)}}^{(n)}(\delta_n, \delta_n) = \frac{1}{2\gamma} \text{ and } e^{u(\cdot)} = \frac{G_{\beta^{(n)}}^{(n)}(\delta_n, \cdot)}{G_{\beta^{(n)}}^{(n)}(\delta_n, \delta_n)} = e^{u_{\beta^{(n)}}^{(n)}(\delta_n, \cdot)}.$$

Finally, by Theorem C, the law of $(\beta_{V_n}^{(n)}, G_{\beta^{(n)}}^{(n)}(\delta_n, \delta_n))$ is the same under $\nu_V^W(d\beta, d\gamma)$ and $\nu_{\tilde{V}_n}^{W^{(n)}}(d\beta^{(n)})$, and since $\beta^{(n)} \mapsto (\beta_{V_n}^{(n)}, G_{\beta^{(n)}}^{(n)}(\delta_n, \delta_n))$ is a bijection by Proposition A, it implies that under $\nu_V^W(d\beta, d\gamma)$, $\beta^{(n)}$ has law $\nu_{\tilde{V}_n}^{W^{(n)}}$. \square

Proposition 8. *With the definition of Proposition 2, for all $i, j \in V_n$, all $\beta \in \mathcal{D}_V^W$, all $\gamma > 0$,*

$$G_{\beta^{(n)}}^{(n)}(i, j) = \widehat{G}_\beta^{(n)}(i, j) + \frac{1}{2\gamma} \psi_\beta^{(n)}(i) \psi_\beta^{(n)}(j).$$

Proof. For simplicity, we omit the subscripts $\beta, \beta^{(n)}$ in $\widehat{G}_\beta^{(n)}, \psi_\beta^{(n)}, G_{\beta^{(n)}}^{(n)}$ in the expression below. By Proposition 6, Lemma 2, using $(\beta^{(n)})_{V_n} = \beta_{V_n}$, we find that

$$(4.3) \quad G^{(n)}(i, j) = \sum_{\sigma \in \mathcal{P}_{i, j}^{\tilde{V}_n}} \frac{W_\sigma^{(n)}}{(2\beta^{(n)})_\sigma}, \quad \widehat{G}^{(n)}(i, j) = \sum_{\sigma \in \mathcal{P}_{i, j}^{V_n}} \frac{W_\sigma}{(2\beta)_\sigma}$$

and

$$\psi^{(n)}(i) = \frac{G^{(n)}(\delta_n, i)}{G^{(n)}(\delta_n, \delta_n)} = \sum_{\sigma \in \overline{\mathcal{P}}_{i, \delta_n}^{\tilde{V}_n}} \frac{W_\sigma^{(n)}}{(2\beta)_\sigma^-}.$$

Therefore, if we denote $\mathcal{P}_{i, \delta_n, j}^{\tilde{V}_n}$ the collection of paths on \tilde{V}_n starting from i , visiting δ_n at least once, and ending at j , that is,

$$\mathcal{P}_{i, \delta_n, j}^{\tilde{V}_n} = \{\sigma = (\sigma_0, \dots, \sigma_m) \in \mathcal{P}_{i, j}^{\tilde{V}_n}, \text{ such that } \exists 0 \leq k \leq m, \sigma_k = \delta_n\}$$

then, since $(W^{(n)})_{V_n, V_n} = W_{V_n, V_n}$ and $(\beta^{(n)})_{V_n} = \beta_{V_n}$,

$$\begin{aligned} G^{(n)}(i, j) - \widehat{G}^{(n)}(i, j) &= \sum_{\sigma \in \mathcal{P}_{i, \delta_n, j}^{\tilde{V}_n}} \frac{W_\sigma^{(n)}}{(2\beta^{(n)})_\sigma} \\ &= \left(\sum_{\sigma \in \overline{\mathcal{P}}_{i, \delta_n}^{\tilde{V}_n}} \frac{W_\sigma^{(n)}}{(2\beta^{(n)})_\sigma^-} \right) \cdot \left(\sum_{\sigma \in \mathcal{P}_{\delta_n, j}^{\tilde{V}_n}} \frac{W_\sigma^{(n)}}{(2\beta^{(n)})_\sigma} \right) \\ &= \psi^{(n)}(i) G^{(n)}(\delta_n, j) = \psi^{(n)}(i) \psi^{(n)}(j) G^{(n)}(\delta_n, \delta_n) = \psi^{(n)}(i) \psi^{(n)}(j) \frac{1}{2\gamma} \end{aligned}$$

where we used Lemma 2 in the last equality. \square

5. THE MARTINGALE PROPERTY

We remind that $\mathcal{F}^{(n)} = \sigma(\beta_i, i \in V_n)$ is the sub σ -field generated by the random variables $\beta \mapsto \beta_i, i \in V_n$. The following proposition is the key property for the main theorem.

Proposition 9. *With the notations of Definition 1, for all $n \in \mathbb{N}$, $\psi^{(n)}$ has finite moments. Moreover, we have, ν_V^W -a.s.,*

$$(5.1) \quad \mathbb{E}_{\nu_V^W} (\psi^{(n+1)}(i) | \mathcal{F}^{(n)}) = \psi^{(n)}(i), \quad \forall i \in V,$$

and for all $i, j \in V$,

$$(5.2) \quad \mathbb{E}_{\nu_V^W} (\psi^{(n+1)}(i)\psi^{(n+1)}(j) - \psi^{(n)}(i)\psi^{(n)}(j) | \mathcal{F}^{(n)}) = \mathbb{E}_{\nu_V^W} (\widehat{G}^{(n+1)}(i, j) - \widehat{G}^{(n)}(i, j) | \mathcal{F}^{(n)}).$$

Remark 13. *In Theorem B, by the change of variables $\tilde{u}(\cdot) = u(\cdot) - \frac{\sum_{i \in V} u(i)}{|V|}$, the new variables $(\tilde{u}(i))_{i \in V}$ are in the space $\{\sum_{i \in V} \tilde{u}(i) = 0\}$ and the density becomes*

$$\tilde{\mathcal{Q}}_{i_0}^W(d\tilde{u}) = \frac{1}{\sqrt{2^{|V|-1}}} e^{\tilde{u}(i_0)} e^{-\sum_{i \sim j} W_{i,j} (\cosh(\tilde{u}(i) - \tilde{u}(j)) - 1)} \sqrt{D(W, \tilde{u})} d\tilde{u}_{V \setminus \{i_0\}}.$$

We see from this expression that $e^{\tilde{u}(i) - \tilde{u}(i_0)} \cdot \tilde{\mathcal{Q}}_{i_0}^W = \tilde{\mathcal{Q}}_i^W$, hence that $\int e^{\tilde{u}(i) - \tilde{u}(i_0)} \tilde{\mathcal{Q}}_{i_0}^W(d\tilde{u}) = 1$. Applied to $V = \tilde{V}_n, i_0 = \delta_n$, we get $\mathbb{E}_{\nu_V^W} (\psi^{(n)}(i)) = 1$ which is a particular case of (5.1).

The original proof of that property was rather technical (see the second arXiv version of the present paper). Some time after the first version of this paper was posted on arXiv, a simpler proof of the martingale property (5.1) was given in [7]. Moreover, using some supersymmetric arguments, the following more general property was proved.

Lemma 3 ([7]). *Let $\lambda \in (\mathbb{R}_+)^V$ be a non-negative function on V with bounded support, then*

$$\mathbb{E} \left(e^{-\langle \lambda, \psi^{(n+1)} \rangle - \frac{1}{2} \langle \lambda, \widehat{G}^{(n+1)} \lambda \rangle} | \mathcal{F}^{(n)} \right) = e^{-\langle \lambda, \psi^{(n)} \rangle - \frac{1}{2} \langle \lambda, \widehat{G}^{(n)} \lambda \rangle}.$$

We provide here a different proof of this assertion based on elementary computations on the measures ν_V^W on finite sets. It also provides a simpler proof of the original assertion Proposition 9 by differentiating in λ .

5.1. Marginal and conditional laws of ν_V^W . In this subsection, we suppose that $\mathcal{G} = (V, E)$ is finite. We state some identities on marginal and conditional laws of the distribution ν_V^W , which will be instrumental in the proof of the martingale property in the next subsection.

Let us first remark that the law ν_V^W defined in Theorem A can be extended to the case where $P = (W_{i,j})_{i,j \in V}$ has non-zero, diagonal coefficients. Indeed, if some diagonal coefficients of P are positive, then changing from variables (β_i) to variables $(\beta_i - \frac{1}{2}W_{i,i})$, we get the law $\nu_V^{\widetilde{W}}$ where $(\widetilde{W}_{i,j})$ is obtained from $(W_{i,j})$ by replacing all diagonal entries by 0. While it is not very natural from the point of view of the VRJP to allow non-zero diagonal coefficients, it is convenient in this section to allow this possibility since it simplifies the statements about conditional law.

Recall that for any function $\zeta : V \mapsto \mathbb{R}$ and any subset $U \subset V$, we write ζ_U for the restriction of ζ to the subset U . Similarly, if A is a $V \times V$ matrix and $U \subset V, U' \subset V$, we

write $A_{U,U'}$ for its restriction to the block $U \times U'$. We also write $d\beta_U = \prod_{i \in U} d\beta_i$ to denote integration on variables β_U .

In the next lemma we give an extension of the family of probability distributions ν_V^W . This extension was proposed by Letac, in the unpublished note [13] discussing the integral defined in [22]. We give a proof of this lemma using Theorem A.

Lemma 4 (Letac, [13]). *Let V be finite and $P = (W_{i,j})_{i,j \in V}$ be a symmetric matrix with non-negative coefficients. Let $(\eta_i)_{i \in V} \in \mathbb{R}_+^V$ be a vector with non-negative coefficients. Then the following measure on \mathcal{D}_V^W*

$$(5.3) \quad \begin{aligned} \nu_V^{W,\eta}(d\beta) &:= e^{-\frac{1}{2}\langle \eta, (H_\beta)^{-1} \eta \rangle} e^{\langle \eta, 1 \rangle} \nu_V^W(d\beta) \\ &= \mathbf{1}_{H_\beta > 0} \left(\frac{2}{\pi} \right)^{|V|/2} e^{-\frac{1}{2}\langle 1, H_\beta 1 \rangle - \frac{1}{2}\langle \eta, (H_\beta)^{-1} \eta \rangle} \frac{1}{\sqrt{\det H_\beta}} e^{\langle \eta, 1 \rangle} d\beta_V \end{aligned}$$

is a probability distribution, where 1 in the scalar products $\langle 1, H_\beta 1 \rangle$ and $\langle \eta, 1 \rangle$ is to be un-

derstood as the vector $\begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix}$. Its Laplace transform is, for any $\lambda \in \mathbb{R}_+^V$

$$(5.4) \quad \int e^{-\langle \lambda, \beta \rangle} \nu_V^{W,\eta}(d\beta) = e^{-\langle \eta, \sqrt{\lambda+1} - 1 \rangle - \sum_{i \sim j} W_{i,j} (\sqrt{(1+\lambda_i)(1+\lambda_j)} - 1)} \prod_{i \in V} \frac{1}{\sqrt{1+\lambda_i}}$$

where $\sqrt{\lambda+1} - 1$ should be considered as the vector $(\sqrt{\lambda_i+1} - 1)_{i \in V}$.

It appears in the following lemma that this extension describes all marginal laws of ν_V^W , and that the larger family $\nu_V^{W,\eta}$ is stable by taking marginals and conditional distributions.

Lemma 5. *Assume that V is finite and let $U \subset V$ be a subset. Under $\nu_V^{W,\eta}(d\beta)$,*

(i) β_U is distributed according to $\nu_U^{W_U, \hat{\eta}}$, where

$$(5.5) \quad \hat{\eta} = \eta_U + P_{U, U^c} \mathbf{1}_{U^c},$$

(ii) conditionally on β_U , β_{U^c} is distributed according to $\nu_{U^c}^{\tilde{W}, \tilde{\eta}}$, where $\tilde{P} = (\tilde{W}_{i,j})_{i,j \in U^c}$ and $\tilde{\eta} \in (\mathbb{R}_+)^{U^c}$ are the matrix and vector defined by

$$\tilde{P} = P_{U^c, U^c} + P_{U^c, U} ((H_\beta)_{U, U})^{-1} P_{U, U^c}, \quad \tilde{\eta} = \eta_{U^c} + P_{U^c, U} ((H_\beta)_{U, U})^{-1} (\eta_U).$$

Remark 14. *Note that \tilde{P} has non-zero diagonal coefficients.*

N.B. As we can observe, all the quantities with \checkmark are relative to vectors or matrices on U^c , while the quantities with $\hat{\cdot}$ are relative to vectors or matrices on U .

Lemma 6. *Let $\mathcal{G} = (V, E)$ be a finite connected graph endowed with conductances $P = (W_{i,j})_{i,j \in V}$. Let $(\eta_i)_{i \in V} \in \mathbb{R}_+^V$ be a vector with non-negative coefficients. Let $U \subset V$. For $\beta \in \mathcal{D}_V^W$, define $\psi_\beta = G_\beta \eta$ where $G_\beta = H_\beta^{-1}$; define $\hat{\eta} = \eta_U + P_{U, U^c} \mathbf{1}_{U^c}$, $\hat{G}_\beta^U = ((H_\beta)_{U, U})^{-1}$ and $\hat{\psi}_\beta = \hat{G}_\beta^U (\hat{\eta})$. For any $\lambda \in \mathbb{R}_+^V$, we have, $\nu_V^{W,\eta}$ a.s.,*

$$(5.6) \quad \mathbb{E}_{\nu_V^{W,\eta}}(e^{-\langle \lambda, \psi \rangle - \frac{1}{2}\langle \lambda, G \lambda \rangle} | \mathcal{F}_U) = e^{-\langle \lambda_U, \hat{\psi} \rangle - \langle \lambda_{U^c}, \mathbf{1}_{U^c} \rangle - \frac{1}{2}\langle \lambda_U, \hat{G}^U \lambda_U \rangle}$$

where $\mathcal{F}_U = \sigma(\beta_i, i \in U)$.

Proof of Lemma 4 and Lemma 5. Lemma 4 and the assertions (i) and (ii) of Lemma 5 are consequences of the same decomposition of the measure $\nu_V^{W,\eta}$. It is partially inspired by computations in [13]. We write H_β as block matrix

$$H_\beta = \begin{pmatrix} H_{U,U} & -P_{U,U^c} \\ -P_{U^c,U} & H_{U^c,U^c} \end{pmatrix} \text{ and define } \widehat{G}^U = (H_{U,U})^{-1}.$$

Now, define the Schur's complement

$$(5.7) \quad \check{H}^{U^c} = H_{U^c,U^c} - P_{U^c,U} \widehat{G}^U P_{U,U^c},$$

and

$$\check{G}^{U^c} = \left(\check{H}^{U^c} \right)^{-1}.$$

We have

$$(5.8) \quad H_\beta = \begin{pmatrix} I_U & 0 \\ -P_{U^c,U} \widehat{G}^U & I_{U^c} \end{pmatrix} \begin{pmatrix} H_{U,U} & 0 \\ 0 & \check{H}^{U^c} \end{pmatrix} \begin{pmatrix} I_U & -\widehat{G}^U P_{U,U^c} \\ 0 & I_{U^c} \end{pmatrix}.$$

Remark that with notations of (ii) of Lemma 5 we have

$$\check{H}^{U^c} = 2\beta_{U^c} - \check{P}.$$

By (5.8), we have

$$(5.9) \quad \begin{aligned} & \langle 1, H_\beta 1 \rangle \\ &= \langle 1_{U^c}, \check{H}^{U^c} 1_{U^c} \rangle + \langle 1_U, H_{U,U} 1_U \rangle + \langle 1_{U^c}, P_{U^c,U} \widehat{G}^U P_{U,U^c} 1_{U^c} \rangle - 2 \langle 1_U, P_{U,U^c} 1_{U^c} \rangle. \end{aligned}$$

On the other hand, by (5.8) again, we have

$$(5.10) \quad G_\beta = H_\beta^{-1} = \begin{pmatrix} I_U & \widehat{G}^U P_{U,U^c} \\ 0 & I_{U^c} \end{pmatrix} \begin{pmatrix} \widehat{G}^U & 0 \\ 0 & \check{G}^{U^c} \end{pmatrix} \begin{pmatrix} I_U & 0 \\ P_{U^c,U} \widehat{G}^U & I_{U^c} \end{pmatrix}$$

therefore, since

$$\begin{pmatrix} I_U & 0 \\ P_{U^c,U} \widehat{G}^U & I_{U^c} \end{pmatrix} \begin{pmatrix} \eta_U \\ \eta_{U^c} \end{pmatrix} = \begin{pmatrix} \eta_U \\ \check{\eta} \end{pmatrix},$$

we get,

$$(5.11) \quad \langle \eta, G_\beta \eta \rangle = \langle \eta_U, \widehat{G}^U \eta_U \rangle + \langle \check{\eta}, \check{G}^{U^c} \check{\eta} \rangle.$$

Combining (5.9) and (5.11) we have

$$(5.12) \quad \begin{aligned} \langle 1, H_\beta 1 \rangle + \langle \eta, G_\beta \eta \rangle - 2 \langle \eta, 1 \rangle &= \langle 1_{U^c}, \check{H}^{U^c} 1_{U^c} \rangle + \langle \check{\eta}, \check{G}^{U^c} \check{\eta} \rangle - 2 \langle \check{\eta}, 1_{U^c} \rangle \\ &+ \langle 1_U, H_{U,U} 1_U \rangle + \langle \widehat{\eta}, \widehat{G}^U \widehat{\eta} \rangle - 2 \langle \widehat{\eta}, 1_U \rangle. \end{aligned}$$

By (5.8), we also have

$$(5.13) \quad \det H_\beta = \det H_{U,U} \det \check{H}^{U^c}, \quad \mathbf{1}_{H_\beta > 0} = \mathbf{1}_{H_{U,U} > 0} \mathbf{1}_{\check{H}^{U^c} > 0}.$$

Combining (5.12) and (5.13), we have,

$$\begin{aligned}
(5.14) \quad & \left(\frac{2}{\pi}\right)^{|V|/2} e^{-\frac{1}{2}\langle 1, H_\beta 1 \rangle - \frac{1}{2}\langle \eta, G_\beta \eta \rangle + \langle \eta, 1 \rangle} \frac{\mathbb{1}_{H_\beta > 0}}{\sqrt{\det H_\beta}} \\
& = \left(\frac{2}{\pi}\right)^{|U|/2} e^{-\frac{1}{2}\langle 1_U, H_{U,U} 1_U \rangle - \frac{1}{2}\langle \hat{\eta}, \hat{G}^U \hat{\eta} \rangle + \langle \hat{\eta}, 1_U \rangle} \frac{\mathbb{1}_{H_{U,U} > 0}}{\det H_{U,U}} \\
& \quad \cdot \left(\frac{2}{\pi}\right)^{|U^c|/2} e^{-\frac{1}{2}\langle 1_{U^c}, \check{H}^{U^c} 1_{U^c} \rangle - \frac{1}{2}\langle \check{\eta}, \check{G}^{U^c} \check{\eta} \rangle + \langle \check{\eta}, 1_{U^c} \rangle} \frac{\mathbb{1}_{\check{H}^{U^c} > 0}}{\sqrt{\det \check{H}^{U^c}}}.
\end{aligned}$$

We remark that the left-hand side is the density of $\nu_V^{W,\eta}(d\beta)$, that the first term of the right-hand side corresponds to the density of $\nu_U^{W_{U,U},\hat{\eta}}(d\beta_U)$ and that, β_U being fixed, the second term of the right-hand side is the density of $\nu_{U^c}^{\check{W},\check{\eta}}(d\beta_{U^c})$. (Indeed, as remarked above, $\check{H}^{U^c} = 2\beta_{U^c} - \check{P}$ and $\check{P}, \check{\eta}$ are β_U -measurable).

Proof of Lemma 4. Take $\eta = 0$. Then $\check{\eta} = 0$. Integrating on $d\beta_{U^c}$ on both sides of (5.14), with β_U fixed, gives

$$\int \left(\frac{2}{\pi}\right)^{|V|/2} e^{-\frac{1}{2}\langle 1, H_\beta 1 \rangle} \frac{\mathbb{1}_{H_\beta > 0}}{\sqrt{\det H_\beta}} d\beta_{U^c} = \left(\frac{2}{\pi}\right)^{|U|/2} e^{-\frac{1}{2}\langle 1_U, H_{U,U} 1_U \rangle - \frac{1}{2}\langle \hat{\eta}, \hat{G}^U \hat{\eta} \rangle + \langle 1_U, \hat{\eta} \rangle} \frac{\mathbb{1}_{H_{U,U} > 0}}{\det H_{U,U}}$$

since $\int \nu_{U^c}^{\check{W},\check{\eta}}(d\beta_{U^c}) = 1$ by Theorem A. Integrating on $d\beta_U$, it gives $\int \nu_U^{W_{U,U},\hat{\eta}}(d\beta_U) = 1$, since ν_V^W is a probability. Hence, $\nu_U^{W_{U,U},\hat{\eta}}$ is a probability. This implies Lemma 4 since this restriction procedure allows to obtain all possible parameters of the family of measures $\nu_V^{W,\eta}$. Indeed, for V, W, η , consider $\tilde{V} = V \cup \{\delta\}$ the set obtained by adding an extra point, and define $(\tilde{W}_{i,j})_{i,j \in \tilde{V}}$ by $\tilde{W}_{V,V} = W$ and $W_{i,\delta} = \eta_i$ for $i \in V$. Then if we apply the previous identity to \tilde{V} and $U := V \subset \tilde{V}$, we get $\hat{\eta} = \eta$ and $\nu_V^{W,\eta}$ is a probability by the previous argument.

Proof of Lemma 5. Integrating on $d\beta_{U^c}$ on both sides of (5.14), with β_U fixed, gives

$$\begin{aligned}
& \int \left(\frac{2}{\pi}\right)^{|V|/2} e^{-\frac{1}{2}\langle 1, H_\beta 1 \rangle - \frac{1}{2}\langle \eta, G_\beta \eta \rangle + \langle \eta, 1 \rangle} \frac{\mathbb{1}_{H_\beta > 0}}{\sqrt{\det H_\beta}} (d\beta_{U^c}) \\
& = \left(\frac{2}{\pi}\right)^{|U|/2} e^{-\frac{1}{2}\langle 1_U, H_{U,U} 1_U \rangle - \frac{1}{2}\langle \hat{\eta}, \hat{G}^U \hat{\eta} \rangle + \langle 1_U, \hat{\eta} \rangle} \frac{\mathbb{1}_{H_{U,U} > 0}}{\det H_{U,U}}
\end{aligned}$$

since $\int \nu_{U^c}^{\check{W},\check{\eta}}(d\beta_{U^c}) = 1$ by Lemma 4. Hence, the marginal distribution of β_U is $\nu_U^{W_{U,U},\hat{\eta}}$, proving (i). Finally, (ii) is a consequence of the conditional probability density formula. Indeed, if we denote temporarily by $f(\beta)$ the density of $\nu_V^W(d\beta)$, by $f_U(\beta_U)$ its marginal density on U and by $f_{U^c}^{\beta_U}(\beta_{U^c})$ the conditional density of β_{U^c} conditioned on β_U , we have by (5.14) and (i),

$$f_{U^c}^{\beta_U}(\beta_{U^c}) = \frac{f(\beta)}{f_U(\beta_U)} = \left(\frac{2}{\pi}\right)^{|U^c|/2} e^{-\frac{1}{2}\langle 1_{U^c}, \check{H}^{U^c} 1_{U^c} \rangle - \frac{1}{2}\langle \check{\eta}, \check{G}^{U^c} \check{\eta} \rangle + \langle \check{\eta}, 1_{U^c} \rangle} \frac{\mathbb{1}_{\check{H}^{U^c} > 0}}{\sqrt{\det \check{H}^{U^c}}}.$$

Since $\check{H}^{U^c} = 2\beta_{U^c} - \check{P}$, $\check{G}^{U^c} = (\check{H}^{U^c})^{-1}$ and \check{P} , $\check{\eta}$ are β_U -measurable, it implies that the right-hand side is the density of $\nu_{U^c}^{\check{W}, \check{\eta}}(d\beta_{U^c})$. \square

Proof of Lemma 6. We take the same notations as in the proof of Lemma 5. By Lemma 5, under $\nu_V^W(d\beta)$, the law of β_{U^c} , conditionally on β_U , is $\nu_{U^c}^{\check{W}, \check{\eta}}$. Now, we set

$$\check{\psi} = \check{G}^{U^c} \check{\eta}.$$

By (5.10), we have

$$\begin{aligned} \langle \lambda, \psi \rangle + \frac{1}{2} \langle \lambda, G\lambda \rangle &= \langle \lambda, G\eta \rangle + \frac{1}{2} \langle \lambda, G\lambda \rangle \\ &= \left(\lambda_U, \lambda_U \widehat{G}^U P_{U, U^c} + \lambda_{U^c} \right) \begin{pmatrix} \widehat{G}^U & 0 \\ 0 & \check{G}^{U^c} \end{pmatrix} \begin{pmatrix} \eta_U \\ P_{U^c, U} \widehat{G}^U \eta_U + \eta_{U^c} \end{pmatrix} \\ &\quad + \frac{1}{2} \left(\lambda_U, \lambda_U \widehat{G}^U P_{U, U^c} + \lambda_{U^c} \right) \begin{pmatrix} \widehat{G}^U & 0 \\ 0 & \check{G}^{U^c} \end{pmatrix} \begin{pmatrix} \lambda_U \\ P_{U^c, U} \widehat{G}^U \lambda_U + \lambda_{U^c} \end{pmatrix} \end{aligned}$$

If we define $\check{\lambda} = \lambda_{U^c} + P_{U^c, U} \widehat{G}^U \lambda_U \in \mathbb{R}_+^{U^c}$, we have

$$\begin{aligned} \langle \lambda, \psi \rangle + \frac{1}{2} \langle \lambda, G\lambda \rangle &= \langle \check{\lambda}, \check{\psi} \rangle + \frac{1}{2} \langle \check{\lambda}, \check{G}^{U^c} \check{\lambda} \rangle + \langle \lambda_U, \widehat{G}^U \eta_U \rangle + \frac{1}{2} \langle \lambda_U, \widehat{G}^U \lambda_U \rangle \\ &= \langle \check{\lambda}, \check{\psi} \rangle + \frac{1}{2} \langle \check{\lambda}, \check{G}^{U^c} \check{\lambda} \rangle + \langle \lambda_U, \widehat{\psi} \rangle + \frac{1}{2} \langle \lambda_U, \widehat{G}^U \lambda_U \rangle - \langle 1_{U^c}, \check{\lambda} - \lambda_{U^c} \rangle. \end{aligned}$$

Now, remark that

$$\langle \check{\lambda}, \check{\psi} \rangle + \frac{1}{2} \langle \check{\lambda}, \check{G}^{U^c} \check{\lambda} \rangle + \frac{1}{2} \langle \check{\eta}, \check{G}^{U^c} \check{\eta} \rangle = \frac{1}{2} \langle \check{\lambda} + \check{\eta}, \check{G}^{U^c} (\check{\lambda} + \check{\eta}) \rangle.$$

We get,

$$\begin{aligned} \mathbb{E}_{\nu_V^W} \left(e^{-\langle \lambda, \psi \rangle - \frac{1}{2} \langle \lambda, G\lambda \rangle} \mid \mathcal{F}_U \right) &= e^{-\langle \lambda_U, \widehat{\psi} \rangle - \langle \lambda_{U^c}, 1_{U^c} \rangle - \frac{1}{2} \langle \lambda_U, \widehat{G}^U \lambda_U \rangle} \mathbb{E}_{\nu_{U^c}^{\check{W}, \check{\eta}}} \left(e^{-\langle \check{\lambda}, \check{\psi} \rangle - \frac{1}{2} \langle \check{\lambda}, \check{G}^{U^c} \check{\lambda} \rangle + \langle 1_{U^c}, \check{\lambda} \rangle} \right) \\ &= e^{-\langle \lambda_U, \widehat{\psi} \rangle - \langle \lambda_{U^c}, 1_{U^c} \rangle - \frac{1}{2} \langle \lambda_U, \widehat{G}^U \lambda_U \rangle} \mathbb{E}_{\nu_{U^c}^{\check{W}, \check{\lambda} + \check{\eta}}} (1) \end{aligned}$$

which concludes the proof of the lemma, using that $\nu_{U^c}^{\check{W}, \check{\lambda} + \check{\eta}}$ is a probability \square

5.2. Proof of Lemma 3. Remark that since $\psi^{(n)}$ is defined for all n by

$$\begin{cases} (H_\beta \psi^{(n)})_{V_n} = 0, \\ \psi_{V_n^c}^{(n)} = 1, \end{cases}$$

we have $\psi_{V_n}^{(n)} = ((H_\beta)_{V_n, V_n})^{-1}(\eta^{(n)})$, where $\eta^{(n)} = P_{V_n, V_n^c}(1_{V_n^c})$. Moreover, by Lemma 5 (i), under $\nu_V^W(d\beta)$, we know that β_{V_n} has law $\nu_{V_n}^{W, \eta^{(n)}}$. Using Lemma 6 applied to $V = V_{n+1}$ and $U = V_n$, we have that $\widehat{G}_{V_{n+1}, V_{n+1}}^{(n+1)}$ corresponds to G_β in Lemma 6 and $\widehat{G}_{V_n, V_n}^{(n)}$ to \widehat{G}^U , $\eta^{(n+1)}$ to η , and $\eta^{(n)}$ to $\widehat{\eta}$. Hence, we get that a.s.

$$\begin{aligned} \mathbb{E}_{\nu_V^W} \left(e^{-\langle \lambda_{V_{n+1}}, \psi_{V_{n+1}}^{(n+1)} \rangle - \frac{1}{2} \langle \lambda_{V_{n+1}}, \widehat{G}^{(n+1)} \lambda_{V_{n+1}} \rangle} \mid \mathcal{F}^{(n)} \right) &= e^{-\langle \lambda_{V_n}, \psi_{V_n}^{(n)} \rangle - \langle \lambda_{V_{n+1} \setminus V_n}, 1_{V_{n+1} \setminus V_n} \rangle - \frac{1}{2} \langle \lambda_{V_n}, \widehat{G}^{(n)} \lambda_{V_n} \rangle} \\ &= e^{-\langle \lambda_{V_{n+1}}, \psi_{V_{n+1}}^{(n)} \rangle - \frac{1}{2} \langle \lambda_{V_n}, \widehat{G}^{(n)} \lambda_{V_n} \rangle} \end{aligned}$$

since $\psi_{V_{n+1} \setminus V_n}^{(n)} = 1$. This concludes the proof since $\psi^{(n)}$ and $\psi^{(n+1)}$ both equal 1 on V_{n+1}^c .

6. PASSING TO THE LIMIT: PROOF OF THEOREM 1, PROPOSITION 2, PROPOSITION 3

6.1. Proof of Theorem 1 i) and ii).

Proof of Theorem 1 i) and ii). By the path representation (4.3), we know that $\widehat{G}_\beta^{(n)}(i, j)$ is non-decreasing for all $i, j \in V$ since the set of paths $\mathcal{P}_{i,j}^{V_n}$ is non-decreasing, and that $\widehat{G}_\beta^{(n)}(i, j) \leq G^{(n)}(i, j)$ since $\mathcal{P}_{i,j}^{V_n} \subset \widetilde{\mathcal{P}}_{i,j}^{V_n}$. Hence, it converges a.s. to a random variable $\widehat{G}(i, j)$. Since $\widehat{G}^{(n)}(i, j) > 0$ as soon as $i, j \in V_n$ (indeed, V_n is connected), we have $\widehat{G}(i, j) > 0$ a.s.. It remains to prove that $\widehat{G}(i, j) < \infty$. As $\widehat{G}^{(n)}(i, i)$ converges a.s. to $\widehat{G}(i, i)$ and is non-decreasing, for any $h \geq 0$,

$$\begin{aligned} \nu_V^W(\widehat{G}(i, i) \leq h) &= \nu_V^W(\lim_{n \rightarrow \infty} \widehat{G}^{(n)}(i, i) \leq h) \\ &= \lim_{n \rightarrow \infty} \nu_V^W(\widehat{G}^{(n)}(i, i) \leq h) \\ &\geq \lim_{n \rightarrow \infty} \nu_V^W(G^{(n)}(i, i) \leq h) \\ &= \mathbb{P}\left(\frac{1}{2\Gamma} \leq h\right), \end{aligned}$$

where Γ is a gamma random variable with parameters $(\frac{1}{2}, 1)$. In the last equality, we used that $\frac{1}{2G^{(n)}(i, i)}$ has gamma law with parameters $(\frac{1}{2}, 1)$ by Theorem C. Therefore, $\widehat{G}(i, i) < \infty$ a.s. For the off diagonal term, since $(H_\beta)_{V_n, V_n}$ is positive definite, we have by Cauchy-Schwarz inequality

$$\widehat{G}^{(n)}(i, j) = \left\langle \delta_i, \widehat{G}^{(n)} \delta_j \right\rangle \leq \sqrt{\left\langle \delta_i, \widehat{G}^{(n)} \delta_i \right\rangle \left\langle \delta_j, \widehat{G}^{(n)} \delta_j \right\rangle} = \sqrt{\widehat{G}^{(n)}(i, i) \widehat{G}^{(n)}(j, j)}$$

therefore, $\widehat{G}(i, j) \leq \sqrt{\widehat{G}(i, i) \widehat{G}(j, j)}$ and $\widehat{G}(i, j)$ is a.s. finite.

From Proposition 9, we know that $\psi^{(n)}(i)$ is a positive martingale for all $i \in V$. As a positive martingale, $\psi^{(n)}(i)$ converges a.s. to some non-negative integrable random variable $\psi(i)$.

It remains to show that the limits ψ and \widehat{G} do not depend on the choice of the exhausting sequence (V_n) . Assume that (Ω_n) is another increasing exhausting sequence, we can similarly construct the martingale $\phi^{(n)}(i)$ associated with Ω_n . As (Ω_n) and (V_n) are exhausting, we can construct a subsequence n_k such that the alternating sequence $V_{n_1}, \Omega_{n_2}, V_{n_3}, \dots$ is increasing and thus the alternating sequence $\psi^{(n_1)}(i), \phi^{(n_2)}(i), \psi^{(n_3)}(i), \dots$ is a martingale for all $i \in V$. This martingale converges a.s. and this identifies the limits of $\psi^{(n)}(i)$ and $\phi^{(n)}(i)$. The argument is the same for \widehat{G} since the sequence of Green functions associated with the alternating sequence of subsets is non-decreasing and converges a.s. \square

6.2. Representation of the VRJP as a mixture on the infinite graphs: proof of iii).

With the coupling of Section 4.3, by Proposition 8, we have for $\beta \in \mathcal{D}_V^W$ and $\gamma > 0$,

$$G^{(n)}(i, j) = \widehat{G}^{(n)}(i, j) + \frac{1}{2\gamma} \psi^{(n)}(i) \psi^{(n)}(j).$$

By Theorem 1 (i) and (ii), we have that $\nu_V^W(d\beta, d\gamma)$ -a.s.

$$(6.1) \quad \lim_{n \rightarrow \infty} G^{(n)}(i, j) = G(i, j),$$

where $G(i, j)$ is defined in Theorem 1 (iii).

The next corollary of Proposition 7 gives the necessary uniform integrability on jump rates to extend the representation of the VRJP for finite graphs to infinite graphs.

Corollary 2. *For any $i, j \in V$, there exists $n_0 \in \mathbb{N}$, such that the family of random variables $\left\{ \frac{G^{(n)}(i_0, j)}{G^{(n)}(i_0, i)} \right\}_{n \geq n_0}$ is uniformly integrable under $\nu_V^W(d\beta, d\gamma)$.*

Proof. Choose n_0 such that $i, j \in V_{n_0}$, and i and j are connected by a path in V_{n_0} . Denote by K the distance between i and j for the graph distance in V_{n_0} and let $(\sigma_0 = i, \sigma_1, \dots, \sigma_K = j)$ be a directed path from i to j in V_{n_0} . Note that it is also a directed path in any V_n for $n \geq n_0$ since V_n is increasing. Let

$$\eta := \frac{1}{2} \min_{k=0, \dots, K-1} (W_{\sigma_k, \sigma_{k+1}}) > 0.$$

Let $c(\eta) > 0$ be a positive constant depending only on η such that $e^{2x} \leq c(\eta)e^{\eta \cosh(x)}$ for all reals x (which exists since $2|x| \leq \eta \cosh(x)$ for x large enough and since $\cosh(x) \geq 1$ for all x). We can write with Notation (3.4), for $n \geq n_0$,

$$\begin{aligned} \left(\frac{G^{(n)}(i_0, j)}{G^{(n)}(i_0, i)} \right)^2 &= e^{2(u^{(n)}(i_0, j) - u^{(n)}(i_0, i))} = \prod_{k=0}^{K-1} e^{2(u^{(n)}(i_0, \sigma_{k+1}) - u^{(n)}(i_0, \sigma_k))} \\ &\leq c(\eta)^K \prod_{k=0}^{K-1} e^{\eta \cosh((u^{(n)}(i_0, \sigma_{k+1}) - u^{(n)}(i_0, \sigma_k)))}. \end{aligned}$$

By Theorem C and Lemma 2, under $\nu_V^W(d\beta, d\gamma)$, $u^{(n)}(i_0, \cdot)$ has law $\mathcal{Q}_{i_0}^W$. Proposition 7 then implies that for $n \geq n_0$,

$$\mathbb{E}_{\nu_V^W} \left(\left(\frac{G^{(n)}(i_0, j)}{G^{(n)}(i_0, i)} \right)^2 \right) \leq e^{\eta K} 2^{K/2} c(\eta)^K.$$

The family is uniformly bounded in L^2 , in particular uniformly integrable. \square

Consider now a connected finite subset $\Lambda \subset V$ containing i_0 and set

$$\partial^+ \Lambda = \{j \in \Lambda^c, \exists i \in \Lambda \text{ such that } i \sim j\}.$$

Consider also a real $t_0 > 0$. Let T be the following stopping time

$$T = t_0 \wedge \inf\{t \geq 0, Z_t \notin \Lambda\}.$$

By construction, the law of the VRJP in exchangeable time scale on \mathcal{G} up to time T equals the law of the VRJP in exchangeable time scale on \mathcal{G}_n up to time T , for all n such that $\Lambda \cup \partial^+ \Lambda \subset V_n$. For convenience, in this proof we write $\mathbb{P}_{i_0}^{\text{VRJP}, \mathcal{G}}$ for its law on \mathcal{G} and $\mathbb{P}_{i_0}^{\text{VRJP}, \mathcal{G}_n}$ for its law on \mathcal{G}_n . Hence, our previous discussion formally means that $\mathbb{P}_{i_0}^{\text{VRJP}, \mathcal{G}}((Z_t)_{t \leq T} \in \cdot) = \mathbb{P}_{i_0}^{\text{VRJP}, \mathcal{G}_n}((Z_t)_{t \leq T} \in \cdot)$, for n large enough. We denote by

$$\ell_i(T) = \int_0^T \mathbb{1}_{Z_u=i} du$$

the local time of Z up to time T . Using Corollary 1 and the coupling in Lemma 2, the VRJP in exchangeable time scale on \mathcal{G}_n , starting at i_0 , is a mixture of Markov jump processes with

jump rates from i to j

$$(6.2) \quad \frac{1}{2} W_{i,j}^{(n)} \frac{G^{(n)}(i_0, j)}{G^{(n)}(i_0, i)}$$

under the law $\nu_V^W(d\beta, d\gamma)$. We denote by

$$\tilde{\beta}_i^{(n)} = \sum_{j \sim i} \frac{1}{2} W_{i,j}^{(n)} \frac{G^{(n)}(i_0, j)}{G^{(n)}(i_0, i)}$$

the holding time at site i (note that $\tilde{\beta}_i^{(n)} = \beta_i^{(n)}$ for $i \neq i_0$). We denote by $P_{i_0}^{\text{MJP}}$ the law of the Markov jump process with jump rates $\frac{1}{2} W_{i,j}$ starting at i_0 . The Radon-Nykodim derivative of the law of $(Z_t)_{t \leq T}$ under the law of the Markov jump process with jump rates (6.2) with respect to its law under $P_{i_0}^{\text{MJP}}$ is

$$(6.3) \quad e^{-\sum_{i \in \Lambda} \ell_i(T) (\tilde{\beta}_i^{(n)} - \frac{1}{2} W_i)} \frac{G^{(n)}(i_0, Z_T)}{G^{(n)}(i_0, i_0)},$$

where as usual $W_i = \sum_{j \sim i} W_{i,j}$. We postpone the proof of this formula to the end this subsection.

Formula (6.3) implies that for all positive bounded test functions F , for n large enough,

$$(6.4) \quad \begin{aligned} & \mathbb{E}_{i_0}^{\text{VRJP}, \mathcal{G}} (F((Z_t)_{t \leq T})) = \mathbb{E}_{i_0}^{\text{VRJP}, \mathcal{G}_n} (F((Z_t)_{t \leq T})) \\ & = \int \sum_{j \in \Lambda \cup \partial^+ \Lambda} E_{i_0}^{\text{MJP}} \left(\mathbf{1}_{Z_T=j} F((Z_t)_{t \leq T}) e^{-\sum_{i \in \Lambda} \ell_i(T) (\tilde{\beta}_i^{(n)} - \frac{1}{2} W_i)} \frac{G^{(n)}(i_0, j)}{G^{(n)}(i_0, i_0)} \right) \nu_V^W(d\beta, d\gamma). \end{aligned}$$

From (6.1), we have a.s.

$$\lim_{n \rightarrow \infty} \tilde{\beta}_i^{(n)} = \tilde{\beta}_i := \sum_{j \sim i} \frac{1}{2} W_{i,j} \frac{G(i_0, j)}{G(i_0, i)}.$$

Remark that in (6.4), the term $e^{-\sum_{i \in \Lambda} \ell_i(T) (\tilde{\beta}_i^{(n)} - \frac{1}{2} W_i)}$ is bounded since $\Lambda \cup \partial^+ \Lambda$ is finite and $T \leq t_0$. Using the uniform integrability of $\frac{G^{(n)}(i_0, j)}{G^{(n)}(i_0, i_0)}$, Corollary 2, we get, letting n tend to ∞ , that

$$\begin{aligned} & \mathbb{E}_{i_0}^{\text{VRJP}, \mathcal{G}} (F((Z_t)_{t \leq T})) \\ & = \int \sum_{j \in \Lambda \cup \partial^+ \Lambda} E_{i_0}^{\text{MJP}} \left(\mathbf{1}_{Z_T=j} F((Z_t)_{t \leq T}) e^{-\sum_{i \in \Lambda} \frac{1}{2} \ell_i(T) (\tilde{\beta}_i - \frac{1}{2} W_i)} \frac{G(i_0, j)}{G(i_0, i_0)} \right) \nu_V^W(d\beta, d\gamma) \\ & = \int E_{i_0}^{\beta, \gamma, i_0} (F((Z_t)_{t \leq T})) \nu_V^W(d\beta, d\gamma) \end{aligned}$$

where $E_{i_0}^{\beta, \gamma, i_0}$ is the expectation associated with the probability $P_{i_0}^{\beta, \gamma, i_0}$ defined in Theorem 1. Since Λ and t_0 can be chosen arbitrarily, the previous identity characterizes the law of $(Z(t))_{t \geq 0}$. This concludes the proof of Theorem 1 (iii).

Proof of formula (6.3). Consider the Markov jump process on the graph \mathcal{G}_n with jump rates $\frac{1}{2} W^{(n)}$, denote by $P_{i_0}^{\text{MJP}, (n)}$ its law starting from i_0 . At each vertex i , it waits an exponential random time with parameter $\frac{1}{2} W_i^{(n)}$, then jumps to $j \sim i$ with probability proportional

to $W_{i,j}^{(n)}$. On time interval $[0, t]$, the probability that it follows the discrete path $(\sigma_0 = i_0, \sigma_1, \dots, \sigma_n)$ and jumps at times $0 < s_1 < \dots < s_n < t$ has distribution

$$\begin{aligned} & \left(\prod_{k=0}^{n-1} \frac{W_{\sigma_k, \sigma_{k+1}}^{(n)}}{W_{\sigma_k}^{(n)}} \right) \left(e^{-\frac{1}{2} W_{\sigma_n}^{(n)} (t-s_n)} \prod_{k=0}^{n-1} W_{\sigma_k}^{(n)} e^{-\sum_{k=0}^{n-1} \frac{1}{2} W_{\sigma_k}^{(n)} (s_{k+1} - s_k)} \right) ds_1 \cdots ds_n \\ &= \left(\prod_{k=0}^{n-1} W_{\sigma_k, \sigma_{k+1}}^{(n)} \right) e^{-\frac{1}{2} \sum_{i \in \tilde{V}_n} W_i^{(n)} \ell_i((\sigma_k), (s_k))} ds_1 \cdots ds_n, \end{aligned}$$

where $\ell_i((\sigma_k), (s_k))$ is the total time spent at position i by the trajectory with discrete path (σ_k) and jump times (s_k) . The same formula is true for the Markov jump process with jump rates (6.2) with $\frac{1}{2} W_{i,j}^{(n)}$ replaced by $\frac{1}{2} W_{i,j}^{(n)} \frac{G^{(n)}(i_0, j)}{G^{(n)}(i_0, i)}$. By cancellation of the ratios $\frac{G^{(n)}(i_0, j)}{G^{(n)}(i_0, i)}$ along the trajectory, it gives that on time interval $[0, t]$, the Radon-Nikodym derivative of the law of the Markov jump process with jump rates (6.2) starting at i_0 with respect to $P_{i_0}^{MJP, (n)}$ is

$$M_t := e^{-\sum_{i \in \tilde{V}_n} \ell_i(t) (\tilde{\beta}_i^{(n)} - \frac{1}{2} W_i^{(n)}) \frac{G^{(n)}(i_0, Z_t)}{G^{(n)}(i_0, i_0)}}.$$

Moreover, M_t is a martingale and is bounded on finite time intervals. Since T is a bounded stopping time, $T \leq t_0$, we have (6.3) for the stopping time T since $W_{V_n, V_n}^{(n)} = W_{V_n, V_n}$. \square

6.3. Proof of Proposition 2, and iv) of Theorem 1.

Proof of Proposition 2. Recall notation of Section 3.2 and identities (3.9). As $n \mapsto \widehat{G}^{(n)}(i, j)$ is increasing, we have

$$(6.5) \quad \widehat{G}(i, j) = \sum_{\sigma \in \mathcal{P}_{i,j}^V} \frac{W_\sigma}{(2\beta)_\sigma}.$$

By arguments similar to (3.10), we have

$$\frac{\widehat{G}(i_0, i)}{\widehat{G}(i_0, i_0)} = \sum_{\sigma \in \mathcal{P}_{i_0, i}^V} \frac{W_\sigma}{(2\beta)_\sigma}.$$

We recall that $(\tilde{Z}_n)_{n \in \mathbb{N}}$ denotes the discrete time process which describes successive jumps of the process $(Z_t)_{t \in \mathbb{R}_+}$. Clearly, $\{\tau_{i_0}^+ < \infty\} = \{\exists n \geq 1, \text{ s.t. } \tilde{Z}_n = i_0\}$. Therefore, if we denote $\{(\tilde{Z}_n) \sim \sigma\} = \{\tilde{Z}_0 = \sigma_0, \dots, \tilde{Z}_m = \sigma_m\}$ with $m = |\sigma|$, then for $i \neq i_0$

$$\begin{aligned} (6.6) \quad h(i) &:= P_i^{\beta, \gamma, i_0}(\tau_{i_0}^+ < \infty) = \sum_{\sigma \in \mathcal{P}_{i, i_0}^V} P_i^{\beta, \gamma, i_0}((\tilde{Z}_n) \sim \sigma) \\ &= \sum_{\sigma \in \mathcal{P}_{i, i_0}^V} \frac{W_\sigma}{(2\beta)_\sigma} \frac{G(i_0, i_0)}{G(i_0, i)} = \frac{\widehat{G}(i_0, i)}{\widehat{G}(i_0, i_0)} \cdot \frac{G(i_0, i_0)}{G(i_0, i)}. \end{aligned}$$

It follows from $G(i, j) = \widehat{G}(i, j) + \frac{1}{2\gamma} \psi(i) \psi(j)$ that, for $i \neq i_0$,

$$P_i^{\beta, \gamma, i_0}(\tau_{i_0}^+ = \infty) = 1 - h(i) = \frac{\psi(i_0) \widehat{G}(i_0, i_0) \psi(i) - \widehat{G}(i_0, i) \psi(i_0)}{2\gamma \widehat{G}(i_0, i_0) G(i_0, i)}.$$

Therefore,

$$\begin{aligned}
P_{i_0}^{\beta, \gamma, i_0}(\tau_{i_0}^+ = \infty) &= \sum_{j \sim i_0} \frac{W_{i_0, j} G(i_0, j)}{2\tilde{\beta}_{i_0} G(i_0, i_0)} P_j^{\beta, \gamma, i_0}(\tau_{i_0}^+ = \infty) \\
(6.7) \qquad \qquad \qquad &= \sum_{j \sim i_0} \frac{\psi(i_0) W_{i_0, j} \widehat{G}(i_0, i_0) \psi(j) - \widehat{G}(i_0, j) \psi(i_0)}{4\gamma \tilde{\beta}_{i_0} \widehat{G}(i_0, i_0) G(i_0, i_0)}.
\end{aligned}$$

By definition, for n large enough, we have $H\widehat{G}^{(n)}(i_0, \cdot) = \mathbf{1}_{i_0}(\cdot)$. Taking the limit $n \rightarrow \infty$, we have $H\widehat{G}(i_0, \cdot) = \mathbf{1}_{i_0}(\cdot)$. By (iii) of Theorem 2 (proved in section 6.5), we have $H\psi(\cdot) = 0$, therefore,

$$\sum_{j \sim i_0} W_{i_0, j} [\psi(j) \widehat{G}(i_0, i_0) - \psi(i_0) \widehat{G}(i_0, j)] = \psi(i_0),$$

$$\text{hence } P_{i_0}^{\beta, \gamma, i_0}(\tau_{i_0}^+ = \infty) = \frac{\psi(i_0)^2}{4\gamma \tilde{\beta}_{i_0} \widehat{G}(i_0, i_0) G(i_0, i_0)}. \quad \square$$

Proof of Theorem 1, (iv). From Proposition 2, we see that $P_{i_0}^{\beta, \gamma, i_0}(\tau_{i_0}^+ = \infty) > 0$ if and only if $\psi(i_0) > 0$. Since the Markov jump process $P_{i_0}^{\beta, \gamma, i_0}$ is irreducible (\mathcal{G} is connected), (iv) follows. \square

6.4. Ergodicity and the 0-1 law: proof of Proposition 3 and 5.

Proof of Proposition 3. From the expression of the Laplace transform of β , c.f. Proposition 1, we see that under $\nu_V^W(d\beta)$, $(\beta_i)_{i \in V}$ is stationary for the action of \mathcal{A} .

By 1-dependence, c.f. Proposition 1, it is also ergodic. Indeed, assume that $(\tau_n) \in \mathcal{A}^{\mathbb{N}}$ is a sequence of automorphisms such that $d_{\mathcal{G}}(i_0, \tau_n(i_0)) \rightarrow \infty$ for some vertex i_0 . We prove that (τ_n) is mixing in the sense that for all $A, B \in \sigma(\beta_i, i \in V)$

$$(6.8) \qquad \qquad \qquad \lim_{n \rightarrow \infty} \nu_V^W(\tau_n^{-1}(B) \cap A) = \nu_V^W(A) \nu_V^W(B).$$

Assume that $V_1 \subset V$ is finite and that $A, B \in \sigma(\beta_j, j \in V_1)$. By 1-dependence, $\tau_n^{-1}(B)$ is independent of A for n large enough. This implies that (6.8) is true for all A, B in the algebra $\mathcal{P} = \cup_{V_1 \text{ finite}} \sigma(\beta_j, j \in V_1)$. Now, any measurable set in $\sigma\{\beta_j, j \in V\}$ can be approximated by an element in \mathcal{P} , i.e. for $\epsilon > 0$ and A, B in $\sigma\{\beta_j, j \in V\}$ we can find $A_0, B_0 \in \mathcal{P}$ such that $\nu_V^W(A \Delta A_0) < \epsilon$ and $\nu_V^W(B \Delta B_0) < \epsilon$ where Δ is the symmetric difference (see e.g. [11, Theorem D, Section 13, page 56]). Hence, also $\nu_V^W(\tau_n^{-1}(B) \Delta \tau_n^{-1}(B_0)) < \epsilon$ since τ_n is measure preserving. This easily imply (6.8) for all A and B . Finally, if A is τ -invariant, (6.8) implies $(\nu_V^W(A))^2 = \nu_V^W(A)$, hence $\nu_V^W(A)$ equals 0 or 1.

Let us prove stationarity and ergodicity of the random variables $(\widehat{G}(i, j))_{i, j \in V}$. We also denote by τ the transformation on $\mathbb{R}^{V \times V}$ given by, for $(M(i, j)) \in \mathbb{R}^{V \times V}$, $\tau M(i, j) = M(\tau i, \tau j)$. Since the limit $\widehat{G}_\beta(i, j)$ does not depend on the choice of the sequence V_n , a.s., we have that $\tau(\widehat{G}_\beta) = \widehat{G}_{\tau(\beta)}$, by choosing the approximating sequences V_n and τV_n . It implies that $(\widehat{G}(i, j))$ is stationary. Moreover, if $A \in \mathcal{B}(\mathbb{R}^{V \times V})$ is τ -invariant, then the set $\{\beta, \widehat{G}_\beta \in A\}$ is a.s. τ -invariant. Hence, A has measure 0 or 1 under the law of \widehat{G} . The proof is similar for the random variables $(\psi(i))_{i \in V}$.

The event $\{\psi(i) = 0, \forall i \in V\}$ is clearly invariant by \mathcal{A} , hence has probability 0 or 1. Together with (iv) of Theorem 1 it concludes the proof of the proposition. \square

Proof of Proposition 5. The proof is similar to the proof of Proposition 3. \square

6.5. Proof of Theorem 2: relation with spectral properties of the random schrödinger operator.

Proof of Theorem 2 (i). By Proposition 1, since ν_V^W is supported on \mathcal{D}_V^W , we have a.s. that $H_{V_n \times V_n} > 0$ and passing to the limit, we get $H \geq 0$. Hence, $\sigma(H) \subset [0, +\infty)$. \square

Proof of Theorem 2 (ii). As $-\varepsilon$ is strictly outside the spectrum of H a.s., the equation $(H + \varepsilon)\widehat{G}^\varepsilon = \text{Id}$ has a unique finite solution, we can verify that $\sum_{\sigma \in \mathcal{P}_{i,j}^V} \frac{W_\sigma}{(2\beta + \varepsilon)_\sigma}$ is a solution to this equation. Now by (6.5) we have

$$(H + \varepsilon)^{-1}(i, j) = \sum_{\sigma \in \mathcal{P}_{i,j}^V} \frac{W_\sigma}{(2\beta + \varepsilon)_\sigma} \leq \sum_{\sigma \in \mathcal{P}_{i,j}^V} \frac{W_\sigma}{(2\beta)_\sigma} = \widehat{G}(i, j) < \infty.$$

Therefore, as $\sum_{\sigma \in \mathcal{P}_{i,j}^V} \frac{W_\sigma}{(2\beta + \varepsilon)_\sigma}$ is increasing as $\varepsilon \rightarrow 0$, it converges a.s. to $\widehat{G}(i, j)$. \square

Proof of Theorem 2 (iii). We have, for all $i \in V_n$, $\beta \in \mathcal{D}_V^W$, $\psi_\beta^{(n)}(i) = \sum_{j \sim i} \frac{W_{i,j}}{2\beta_i} \psi_\beta^{(n)}(j)$. As $\psi_\beta^{(n)}(i)$ converges a.s. to $\psi(i)$, the above equality holds in the limit, i.e., for all $i \in V$, a.s.

$$\psi(i) = \sum_{j \sim i} \frac{W_{i,j}}{2\beta_i} \psi(j),$$

this exactly means $(H\psi)(i) = 0$. \square

Proof of Theorem 2 (iv). By Fatou's Lemma, the limit $\psi(i)$ satisfies $\mathbb{E}_{\nu_V^W}(\psi(i)) \leq 1$. By Markov inequality

$$\nu_V^W(\psi(i) \geq C\|i\|_\infty^p) \leq \frac{1}{C\|i\|_\infty^p}.$$

Let $\partial B(0, n)$ be the sphere of radius n for $\|\cdot\|_\infty$, i.e. $\partial B(0, n) = \{i \in \mathbb{Z}^d, \|i\|_\infty = n\}$. When $p > d$,

$$\begin{aligned} \sum_{i \in \mathbb{Z}^d, i \neq 0} \nu_V^W(\psi(i) \geq C\|i\|_\infty^p) &= \sum_{n \geq 1} \sum_{i \in \partial B(0, n)} \nu_V^W(\psi(i) \geq C\|i\|_\infty^p) \\ &\leq \sum_{n \geq 1} \sum_{i \in \partial B(0, n)} \frac{1}{C\|i\|_\infty^p} \\ &\leq C' \sum_n \frac{n^{d-1}}{n^p} < \infty \end{aligned}$$

for some constant $C' > 0$. By Borel-Cantelli lemma, a.s. only a finite number of i satisfies $\psi(i) \geq C\|i\|_\infty^p$. \square

7. h -TRANSFORMS

Corollary 3. Recall that $\tau_{i_0}^+ = \inf\{t \geq 0, Z_t = i_0, \exists s < t \text{ s.t. } Z_s \neq i_0\}$ is the first return time to i_0 of the process $(Z_t)_{t \geq 0}$.

(i) For almost all β and $i_0 \in V$, denote by $\widehat{P}_{i_0}^{\beta, i_0}$ the law of the Markov jump process with jump rate from i to j

$$\begin{cases} \frac{1}{2} W_{i,j} \frac{\widehat{G}(i_0, j)}{\widehat{G}(i_0, i)}, & i \neq i_0, \\ \widetilde{\beta}_{i_0} \frac{W_{i_0, j} \widehat{G}(i_0, j)}{\sum_{k \sim i_0} W_{i_0, k} \widehat{G}(i_0, k)}, & i = i_0, j \sim i_0, \end{cases}$$

where as before $\tilde{\beta}_{i_0} = \sum_{j \sim i_0} \frac{1}{2} W_{i_0, j} \frac{G(i_0, j)}{G(i_0, i_0)}$. Then, for ν_V^W -almost all β , for $\gamma > 0$,

$$P_{i_0}^{\beta, \gamma, i_0} \left((Z_t)_{t \leq \tau_{i_0}^+} \in \cdot \mid \tau_{i_0}^+ < \infty \right) = \widehat{P}_{i_0}^{\beta, i_0} \left((Z_t)_{t \leq \tau_{i_0}^+} \in \cdot \right).$$

(ii) The VRJP in exchangeable time scale, conditionally on $\{\tau_{i_0}^+ < \infty\}$ and up to its first return time to i_0 , is given by the following mixture:

$$\mathbb{P}_{i_0}^{VRJP} \left((Z_t)_{t \leq \tau_{i_0}^+} \in \cdot \mid \tau_{i_0}^+ < \infty \right) = \int \widehat{P}_{i_0}^{\beta, i_0} \left((Z_t)_{t \leq \tau_{i_0}^+} \in \cdot \right) \frac{P_{i_0}^{\beta, \gamma, i_0}(\tau_{i_0}^+ < \infty)}{\mathbb{P}_{i_0}^{VRJP}(\tau_{i_0}^+ < \infty)} \nu_V^W(d\beta, d\gamma).$$

(iii) Let $i_0 \in V$. A.s. on the event $\{\psi(i) > 0, \forall i \in V\}$, $\check{G}(i_0, j) := \widehat{G}(i_0, i_0)\psi(j) - \widehat{G}(i_0, j)\psi(i_0)$ is positive for all $j \neq i_0$, and we define $\check{P}_{i_0}^{\beta, \gamma, i_0}$ as the law of the Markov jump process starting at i_0 and with jump rate from i to j

$$\begin{cases} \frac{1}{2} W_{i, j} \frac{\check{G}(i_0, j)}{\check{G}(i_0, i)}, & i \neq i_0, j \neq i_0, \\ \tilde{\beta}_{i_0} \frac{W_{i_0, j} \check{G}(i_0, j)}{\sum_{k \sim i_0} W_{i_0, k} \check{G}(i_0, k)}, & i = i_0, j \sim i_0, \\ 0, & i \sim i_0, j = i_0. \end{cases}$$

Then, ν_V^W -almost surely on this event, for $\gamma > 0$,

$$P_{i_0}^{\beta, \gamma, i_0} \left((Z_t)_{t \geq 0} \in \cdot \mid \tau_{i_0}^+ = \infty \right) = \check{P}_{i_0}^{\beta, \gamma, i_0} \left((Z_t)_{t \geq 0} \in \cdot \right).$$

(iv) The VRJP in exchangeable time scale, conditionally on the event $\{\tau_{i_0}^+ = \infty\}$, is a mixture of Markov jump processes with mixing law

$$\mathbb{P}_{i_0}^{VRJP} \left(\cdot \mid \tau_{i_0}^+ = \infty \right) = \int \check{P}_{i_0}^{\beta, \gamma, i_0}(\cdot) \frac{P_{i_0}^{\beta, \gamma, i_0}(\tau_{i_0}^+ = \infty)}{\mathbb{P}_{i_0}^{VRJP}(\tau_{i_0}^+ = \infty)} \nu_V^W(d\beta, d\gamma).$$

Remark 15. Note that in the case (i), the conditional jump rates do not depend on γ .

Proof of Corollary 3. (i) Recall from (6.6) that for $i \neq i_0$

$$h(i) = P_i^{\beta, \gamma, i_0}(\tau_{i_0}^+ < \infty) = \frac{\widehat{G}(i_0, i)G(i_0, i_0)}{\widehat{G}(i_0, i_0)G(i_0, i)}.$$

For $i \neq i_0$, we have

$$P_{i_0}^{\beta, \gamma, i_0} \left(X_{t+dt} = j \mid X_t = i, t \leq \tau_{i_0}^+ < \infty \right) \sim \frac{h(j)}{h(i)} P_{i_0}^{\beta, \gamma, i_0} \left(X_{t+dt} = j \mid X_t = i \right).$$

Hence, the jump rate of $P_{i_0}^{\beta, \gamma, i_0}(\cdot \mid \tau_{i_0}^+ < \infty)$, up to time $\tau_{i_0}^+$, from i to j is

$$\frac{1}{2} W_{i, j} \frac{G(i_0, j)}{G(i_0, i)} \frac{h(j)}{h(i)} = \frac{1}{2} W_{i, j} \frac{\widehat{G}(i_0, j)}{\widehat{G}(i_0, i)}.$$

The jump rate of $P_{i_0}^{\beta, \gamma, i_0}(\cdot \mid \tau_{i_0}^+ < \infty)$, up to time $\tau_{i_0}^+$, from i_0 to j is given by

$$\frac{1}{2} W_{i_0, j} \frac{G(i_0, j)}{G(i_0, i_0)} \frac{h(j)}{P_{i_0}^{\beta, \gamma, i_0}(\tau_{i_0}^+ < \infty)} = \tilde{\beta}_{i_0} \frac{W_{i_0, j} \widehat{G}(i_0, j)}{\sum_{k \sim i_0} W_{i_0, k} \widehat{G}(i_0, k)},$$

where $\tilde{\beta}_{i_0} = \sum_{l \sim i_0} \frac{1}{2} W_{i_0, l} \frac{G(i_0, l)}{G(i_0, i_0)}$.

(ii) From (i), we have

$$\begin{aligned} & \mathbb{P}_{i_0}^{\text{VRJP}}((Z_t)_{t \leq \tau_{i_0}^+} \in \cdot \mid \tau_{i_0}^+ < \infty) \\ &= \int P_{i_0}^{\beta, \gamma, i_0}((Z_t)_{t \leq \tau_{i_0}^+} \in \cdot \mid \tau_{i_0}^+ < \infty) \frac{P_{i_0}^{\beta, \gamma, i_0}(\tau_{i_0}^+ < \infty)}{\mathbb{P}_{i_0}^{\text{VRJP}}(\tau_{i_0}^+ < \infty)} \nu_V^W(d\beta, d\gamma) \\ &= \int \widehat{P}_{i_0}^{\beta, \gamma, i_0}((Z_t)_{t \leq \tau_{i_0}^+} \in \cdot) \frac{P_{i_0}^{\beta, \gamma, i_0}(\tau_{i_0}^+ < \infty)}{\mathbb{P}_{i_0}^{\text{VRJP}}(\tau_{i_0}^+ < \infty)} \nu_V^W(d\beta, d\gamma). \end{aligned}$$

(iii) The fact that $\check{G}(i_0, j)$ is positive for $j \neq i_0$ when $\psi > 0$ is a consequence of Proposition 2, and Theorem 1 (iv). Similarly to (i), for $i \neq i_0$, we have

$$P_{i_0}^{\beta, \gamma, i_0}(X_{t+dt} = j \mid X_t = i, \tau_{i_0}^+ = \infty) \sim \frac{1 - h(j)}{1 - h(i)} P^{\beta, \gamma, i_0}(X_{t+dt} = j \mid X_t = i).$$

Hence, the jump rate of $P_{i_0}^{\beta, \gamma, i_0}(\cdot \mid \tau_{i_0}^+ = \infty)$, from $i \neq i_0$ to j is

$$\frac{1}{2} W_{i,j} \frac{G(i_0, j)}{G(i_0, i)} \frac{1 - h(j)}{1 - h(i)} = \frac{1}{2} W_{i,j} \frac{\widehat{G}(i_0, i_0)\psi(j) - \widehat{G}(i_0, j)\psi(i_0)}{\widehat{G}(i_0, i_0)\psi(i) - \widehat{G}(i_0, i)\psi(i_0)} = \frac{1}{2} W_{i,j} \frac{\check{G}(i_0, j)}{\check{G}(i_0, i)}.$$

The jump rate of $P_{i_0}^{\beta, \gamma, i_0}(\cdot \mid \tau_{i_0}^+ = \infty)$, from i_0 to j is given by

$$\frac{1}{2} W_{i_0, j} \frac{G(i_0, j)}{G(i_0, i_0)} \frac{1 - h(j)}{P_{i_0}^{\beta, \gamma, i_0}(\tau_{i_0}^+ = \infty)} = \widetilde{\beta}_{i_0} \frac{W_{i_0, j} \check{G}(i_0, j)}{\sum_{k \sim i_0} W_{i_0, k} \check{G}(i_0, k)},$$

where $\widetilde{\beta}_{i_0} = \sum_{l \sim i_0} \frac{1}{2} W_{i_0, l} \frac{G(i_0, l)}{G(i_0, i_0)}$.

(iv) follows easily from (iii) in the same way as in (ii). □

8. PROOF OF RECURRENCE OF 2-DIMENSIONAL ERRW: THEOREM 5

Consider the square grid $\mathcal{G} = (\mathbb{Z}^2, E)$ with constant edge weight $a_e = a > 0$. From (2.9) in Section 2.5, we know that the ERRW on \mathbb{Z}^2 is a mixture of reversible Markov chains with conductances

$$(8.1) \quad x_{i,j} = W_{i,j} G(0, i) G(0, j)$$

where (W, β, γ) are distributed according to $\widetilde{\nu}_V^a(dW, d\beta, d\gamma)$. We will use [15] to prove the following lemma.

Lemma 7. *There exists $c(a) > 0$ and $\xi(a) > 0$, depending only on a , such that for $\ell \in \mathbb{Z}^2$,*

$$(8.2) \quad \mathbb{E}_{\widetilde{\nu}_V^a} \left(\left(\frac{x_\ell}{x_0} \right)^{\frac{1}{4}} \right) \leq c(a) \|\ell\|_\infty^{-\xi(a)},$$

where $x_i = \sum_{j \sim i} x_{i,j}$ and $(x_{i,j})$ is defined in (8.1).

Proof. This estimate follows from Theorem 2.8 of [15] (it can also be deduced from [17, Lemma 2.5]) which gives a similar estimate on finite boxes. In [15, Theorem 2.8], the estimate is stated for a periodic torus, but it is clear in the proof that the only necessary ingredient is that the finite graph with conductances is invariant by the reflection exchanging 0 and ℓ . For this reason we choose the approximating sequence $V_n = B(\frac{\ell}{2}, n) \cap \mathbb{Z}^2$, where $B(\frac{\ell}{2}, n)$ is the ball with center $\ell/2$ and radius n . Consider as in Section 4.2 the graph

$\mathcal{G}_n = (\tilde{V}_n = V_n \cup \{\delta_n\}, E_n)$, and the associated weights $(a_e^{(n)})_{e \in E_n}$ obtained by restriction of $(\mathcal{G}, (a_e)_{e \in E})$ to V_n with wired boundary condition. Clearly, central symmetry with respect to $\frac{\ell}{2}$ (mapping δ_n to itself) leaves $(\mathcal{G}_n, a^{(n)})$ invariant and exchanges 0 and ℓ .

With the coupling defined in Section 4.3, we define for $i \sim j$, i, j in \tilde{V}_n ,

$$x_{i,j}^{(n)} = W_{i,j}^{(n)} G^{(n)}(0, i) G^{(n)}(0, j).$$

where $W^{(n)}$ is obtained by restriction with wired boundary condition from W . By additivity of Gamma random variables, under $\tilde{\nu}_V^a$, $(W_e^{(n)})_{e \in E_n}$ are independent Gamma random variables with parameters $(a_e^{(n)})_{e \in E_n}$. Hence, the ERRW on \mathcal{G}_n , with initial weights $a^{(n)}$, starting from 0, is a mixture of reversible Markov chains with conductances $(x_e^{(n)})_{e \in E_n}$.

From Theorem 1, with the coupling defined in Section 4.3, we have that for all $i, j \in \mathbb{Z}^2$, $i \sim j$, a.s.

$$(8.3) \quad \lim_{n \rightarrow \infty} x_{i,j}^{(n)} = x_{i,j}.$$

The proof of Theorem 2.8 of [15], can be readily adapted to prove the following estimate.

Lemma 8. *There exists $c(a) > 0$ and $\xi(a) > 0$ only depending on a such that for $\ell \in \mathbb{Z}^2$ and n large enough,*

$$\mathbb{E}_{\tilde{\nu}_V^a} \left(\left(\frac{x_\ell^{(n)}}{x_0^{(n)}} \right)^{\frac{1}{4}} \right) \leq c(a) \|\ell\|_\infty^{-\xi(a)},$$

where, with the usual convention, $x_\ell^{(n)} = \sum_{j \sim \ell} x_{\ell,j}^{(n)}$.

Then, Lemma 7 follows from Lemma 8, (8.3) and Fatou's lemma. \square

We now deduce recurrence of the ERRW from the estimate (8.2) and from Theorem 1 and Proposition 5. We have, for $\ell \neq 0$,

$$x_\ell = \sum_{j \sim \ell} W_{\ell,j} G(0, \ell) G(0, j) = 2\beta_\ell G(0, \ell)^2 \geq \frac{\beta_\ell}{2\gamma^2} \psi(0)^2 \psi(\ell)^2.$$

Similarly,

$$x_0 = \sum_{j \sim 0} W_{0,j} G(0, 0) G(0, j) = G(0, 0)(2\beta_0 G(0, 0) - 1).$$

Hence,

$$(8.4) \quad \frac{x_\ell}{x_0} \geq \frac{\psi(0)^2}{2\gamma^2 G(0, 0)(2\beta_0 G(0, 0) - 1)} \beta_\ell \psi(\ell)^2.$$

Assume the ERRW is transient. By Proposition 5 it implies that, a.s., $\psi(i) > 0$ for all i . Choose first $\eta > 0$ such that

$$\tilde{\nu}_V^a \left(\frac{\psi(0)^2}{2\gamma^2 G(0, 0)(2\beta_0 G(0, 0) - 1)} \leq \eta \right) \leq \frac{1}{2}.$$

For all $\epsilon > 0$, we have by (8.2)

$$(8.5) \quad \tilde{\nu}_V^a \left(\frac{x_\ell}{x_0} \geq \epsilon \right) \leq \frac{1}{\epsilon^{\frac{1}{4}}} c(a) \|\ell\|_\infty^{-\xi(a)}.$$

On the other hand, we have by (8.4)

$$\begin{aligned}
\tilde{\nu}_V^a \left(\frac{x_\ell}{x_0} \geq \epsilon \right) &\geq \tilde{\nu}_V^a \left(\frac{\psi(0)^2}{2\gamma^2 G(0,0)(2\beta_0 G(0,0) - 1)} > \eta, \beta_\ell \psi(\ell)^2 > \frac{\epsilon}{\eta} \right) \\
&= 1 - \tilde{\nu}_V^a \left(\left\{ \frac{\psi(0)^2}{2\gamma^2 G(0,0)(2\beta_0 G(0,0) - 1)} \leq \eta \right\} \cup \left\{ \beta_\ell \psi(\ell)^2 \leq \frac{\epsilon}{\eta} \right\} \right) \\
(8.6) \quad &\geq \frac{1}{2} - \tilde{\nu}_V^a \left(\beta_\ell \psi(\ell)^2 \leq \frac{\epsilon}{\eta} \right).
\end{aligned}$$

By Proposition 5, $\beta_\ell \psi(\ell)^2$ is stationary with respect to translations. Together with (8.6) and (8.5), it implies that

$$\tilde{\nu}_V^a \left(\beta_0 \psi(0)^2 \leq \frac{\epsilon}{\eta} \right) = \tilde{\nu}_V^a \left(\beta_\ell \psi(\ell)^2 \leq \frac{\epsilon}{\eta} \right) \geq \frac{1}{2} - \frac{1}{\epsilon^{\frac{1}{4}}} c(a) \|\ell\|_\infty^{-\xi(a)}.$$

By sending ℓ to infinity, we get $\tilde{\nu}_V^a \left(\beta_0 \psi(0)^2 \leq \frac{\epsilon}{\eta} \right) \geq \frac{1}{2}$. Letting $\epsilon \rightarrow 0$, this is incompatible with $\psi(0) > 0$ a.s., hence with transience of ERRW.

9. PROOF OF FUNCTIONAL CENTRAL LIMIT THEOREMS FOR THE VRJP AND THE ERRW: THEOREM 3 AND 4

Proof of Theorem 3 and Theorem 4. Let us start by the VRJP on \mathbb{Z}^d , $d \geq 3$, with constant weights $W_{i,j} = W$. Assume that the VRJP is transient.

Recall that $(X_n)_{n \in \mathbb{N}}$ is the canonical discrete process on $(\mathbb{Z}^d)^\mathbb{N}$. For ν_V^W -almost all β , let us define \tilde{P}_x^ψ to be the law of the reversible Markov chain, starting at x , with conductances $W_{i,j} \psi(i) \psi(j)$, i.e. with transition probabilities

$$\tilde{P}_x^\psi(X_{n+1} = j | X_n = i) = \frac{W_{i,j} \psi(j)}{\sum_{l \sim i} W_{i,l} \psi(l)}.$$

Denote by $\tilde{P}_x^{\beta, \gamma, 0}$ the law of the underlying discrete time process associated with the Markov jump process $P_x^{\beta, \gamma, 0}$, so that for $i \sim j$

$$\tilde{P}_x^{\beta, \gamma, 0}(X_{n+1} = j | X_n = i) = \frac{W_{i,j} G(0, j)}{\sum_{l \sim i} W_{i,l} G(0, l)}.$$

As ψ is a generalized eigenfunction of H_β , for any $i \in V$,

$$\beta_i = \sum_{j \sim i} \frac{1}{2} W_{i,j} \frac{\psi(j)}{\psi(i)}.$$

It then follows by Proposition 6 that, for $i \neq 0$,

$$h^\psi(i) := \tilde{P}_i^\psi(\tau_0^+ < \infty) = \sum_{\sigma \in \mathcal{P}_{i,0}^V} \tilde{P}_i^\psi(X_n \sim \sigma) = \sum_{\sigma \in \mathcal{P}_{i,0}^V} \frac{W_\sigma \psi(0)}{(2\beta)_\sigma^- \psi(i)} = \frac{\widehat{G}(0, i) \psi(0)}{\widehat{G}(0, 0) \psi(i)}.$$

(recall that $\bar{\mathcal{P}}_{i,0}^V$ is defined in Section 3.2.) Consider the Markov chain $\tilde{P}_0^\psi(\cdot | \tau_0^+ = \infty)$ (Doob's $(1 - h^\psi)$ -transform). By similar computation as in the proof of Proposition 3, we

have that the transition probability of $\tilde{P}_0^\psi(\cdot | \tau_0^+ = \infty)$ from i to j is

$$\frac{W_{i,j}\psi(j)(1-h^\psi(j))}{\sum_{l \sim i} W_{i,l}\psi(l)(1-h^\psi(l))} = \frac{W_{i,j}\check{G}(0,j)}{\sum_{l \sim i} W_{i,l}\check{G}(0,l)}$$

for $j \neq 0$, and 0 when $j = 0$. Therefore, we see that the transition probabilities of $\tilde{P}_0^\psi(\cdot | \tau_0^+ = \infty)$ are the same as those of $\tilde{P}_0^{\beta,\gamma,0}(\cdot | \tau_0^+ = \infty)$, c.f. iii) of Proposition 3. Moreover, if we denote

$$\xi_0 = \sup\{n; X_n = 0\},$$

then, by strong Markov property

$$\begin{aligned} \tilde{P}_0^\psi(X_n \in \cdot | \tau_0^+ = \infty) &= \tilde{P}_0^\psi((X \circ \theta_{\xi_0})_n \in \cdot) \\ \tilde{P}_0^{\beta,\gamma,0}(X_n \in \cdot | \tau_0^+ = \infty) &= \tilde{P}_0^{\beta,\gamma,0}((X \circ \theta_{\xi_0})_n \in \cdot) \end{aligned}$$

where θ_n is the shift in time by n . It follows that $(X \circ \theta_{\xi_0})_n$ has the same law under \tilde{P}_0^ψ and under $\tilde{P}_0^{\beta,\gamma,0}$.

Remark also, from Proposition 3, that $W_{i,j}\psi(i)\psi(j)$ are stationary and ergodic conductances under $\nu_V^W(d\beta)$. We can thus apply Theorem 4.5 and Theorem 4.6 of [6]. In order to have a functional central limit theorem we need to show that, c.f. Theorem 4.5 of [6],

$$(9.1) \quad \mathbb{E}_{\nu_V^W}(W_{i,j}\psi(i)\psi(j)) < \infty.$$

In order to show that it has non-degenerate asymptotic covariance we need to show that, c.f. Theorem 4.6 and identity (4.20) of [6],

$$(9.2) \quad \mathbb{E}_{\nu_V^W} \left(\frac{1}{W_{i,j}\psi(i)\psi(j)} \right) < \infty.$$

By invariance of the law of the conductances by symmetries of \mathbb{Z}^d , we know that the limit diffusion matrix is of the form $\sigma^2 \text{Id}$.

The same reasoning works in the case of the ERRW with constant weights $a_{i,j} = a$: in this case $(W_{i,j})$ are i.i.d., but as shown in Proposition 5, $W_{i,j}\psi(i)\psi(j)$ is also stationary and ergodic under $\tilde{\nu}_V^a(dW, d\beta)$.

Estimates (9.1) and (9.2) are provided by [10] in the VRJP case, and by [8] in the ERRW case. This is summarized in the following lemma.

Lemma 9. (i) (VRJP case) Consider the VRJP on \mathbb{Z}^d , for $d \geq 3$, with constant weights $W_{ij} = W$. There exists $0 < \lambda_2 < \infty$ such that for $W > \lambda_2$, the VRJP is transient and such that (9.1), (9.2) are true under $\nu_V^W(d\beta)$.

(ii) (ERRW case) Consider the ERRW on \mathbb{Z}^d , for $d \geq 3$, with constant weights $a_{ij} = a$. There exists $0 < \tilde{\lambda}_2 < \infty$ such that for $a > \tilde{\lambda}_2$, the ERRW is transient and (9.1), (9.2) are true under $\tilde{\nu}_V^a(dW, d\beta)$.

The proof of that lemma is given below. We first apply it to prove the functional central limit theorem. Consider the VRJP case. Assume that the condition of the lemma is satisfied. Define

$$X_t^{(n)} = \frac{X_{\lfloor nt \rfloor}}{\sqrt{n}}.$$

From [6], we know that there exists $0 < \sigma^2 < \infty$ such that for all bounded Lipschitz function F for the Skorokhod topology, for all $\epsilon > 0$, for all $0 < T < \infty$,

$$(9.3) \quad \lim_{n \rightarrow \infty} Q^* \left(\left| \tilde{E}_0^\psi(F((X_{0 \leq t \leq T}^{(n)})) - \mathbb{E}(F((B_{0 \leq t \leq T}))) \right| \geq \epsilon \right) = 0,$$

where B_t is a d -dimensional Brownian motion with covariance $\sigma^2 \text{Id}$, and where Q^* is the invariant measure for the processes viewed from the particle

$$Q^*(d\beta) = \frac{\sum_{j \sim 0} W_{0,j} \psi(0) \psi(j)}{\mathbb{E}_{\nu_V^W}(\sum_{j \sim 0} W_{0,j} \psi(0) \psi(j))} \cdot \nu_V^W(d\beta).$$

It is clear, since Q^* and ν_V^W are equivalent probability distributions that (9.3) is also true when Q^* is replaced by ν_V^W . This implies an annealed functional central limit theorem for the process (X_n) under the annealed law $\mathbb{E}_{\nu_V^W}(\tilde{P}_0^\psi(\cdot))$:

$$(9.4) \quad \lim_{n \rightarrow \infty} \left| \mathbb{E}_{\nu_V^W} \left(\tilde{E}_0^\psi(F((X_{0 \leq t \leq T}^{(n)})) - \mathbb{E}(F((B_{0 \leq t \leq T}))) \right) \right| = 0.$$

Let $\Upsilon_t^{(n)} := \frac{1}{\sqrt{n}}(X \circ \theta_{\xi_0})_{[nt]}$. Denote d° the Skorohod metric on $D([0, \infty), \mathbb{R}^d)$, the space of càdlag functions $f : [0, \infty) \rightarrow \mathbb{R}^d$. As

$$|X_t^{(n)} - \Upsilon_t^{(n)}| = \frac{1}{\sqrt{n}} |X_{[nt]} - X_{[nt+\xi_0]}| \leq \frac{|\xi_0|}{\sqrt{n}} \xrightarrow{n \rightarrow \infty} 0,$$

we have

$$(9.5) \quad d^\circ(X^{(n)}, \Upsilon^{(n)}) \rightarrow 0.$$

Recall that F is a bounded Lipschitz function for the Skorohod topology, therefore,

$$|F(X_t^{(n)}) - F(\Upsilon_t^{(n)})| \rightarrow 0$$

and (9.4) is valid for $X^{(n)}$ replaced by $\Upsilon^{(n)}$. But $\Upsilon^{(n)}$ has the same law under \tilde{P}_0^ψ and $\tilde{P}_0^{\beta, \gamma, 0}$. This implies the functional central limit theorem (9.4), for the law $\mathbb{E}_{\nu_V^W}(\tilde{P}_0^{\beta, \gamma, 0}(\cdot))$ in place of $\mathbb{E}_{\nu_V^W}(\tilde{P}_0^\psi(\cdot))$ starting from 0. By Theorem 1, the law $\mathbb{E}_{\nu_V^W}(\tilde{P}_0^{\beta, \gamma, 0}(\cdot))$ is that of the discrete time process (\tilde{Z}_n) under \mathbb{P}_0^{VRJP} .

The proof is exactly the same for the ERRW, one just needs to replace the law $\nu_V^W(d\beta)$ by the law $\tilde{\nu}_V^a(dW, d\beta)$. \square

Proof of Lemma 9. Let us start by the ERRW case, ii). Consider the sequence of subsets of \mathbb{Z}^d , $V_n = [-n, n]^d$. Recall that

$$\psi^{(n)}(j) = e^{u^{(n)}(\delta_n, j)},$$

when $j \in V_n$. Consider the point $y_n = (-n, 0, \dots, 0)$, so that y_n is at the boundary of the set, $y_n \sim \delta_n$. By [8, Lemma 7] (which is the ERRW's counterpart of Proposition 7, Section 3.3), we have for $a > 16$,

$$(9.6) \quad \mathbb{E}_{\tilde{\nu}_V^a}((\cosh(u(\delta_n, y_n)))^8) \leq 2.$$

(Indeed, the proof does not depend on the graph structure, nor on the choice of the rooting.)

From, [8, Theorem 4], there exists $0 < \tilde{\lambda}_2 < \infty$ such that if $a > \tilde{\lambda}_2$, then for all i, j in V_n ,

$$(9.7) \quad \mathbb{E}_{\tilde{\nu}_V^a} \left(\left((\cosh(u^{(n)}(\delta_n, i)) - u^{(n)}(\delta_n, j)) \right)^8 \right) \leq 2.$$

Remark that in [8], the rooting of the field is at 0 and the graph is the restriction of the graph \mathbb{Z}^d to V_n . But an attentive reading of the proof shows that the result is also valid for the graph $\mathcal{G}_n = (V_n \cup \{\delta_n\}, E_n)$ and rooting δ_n as well. Indeed, the estimate is based on the protected Ward's estimates, [8, Lemma 4], which remain valid for diamonds inside the set V_n , and on the estimate on effective conductances, [8, Proposition 3], which is in fact an estimate inside a "diamond". Remark that the estimate (9.7) is also valid when i or j is at the boundary of the set V_n (in fact the proof is written in the case where the diamond $R_{i,j}$ is inside the set V_n , which is the case when $j = y_n$ and $i \in \mathbb{Z}^d$ fixed for n large enough). Specified to $j = y_n$ and $i \in \mathbb{Z}^d$ fixed, it gives for n large enough

$$(9.8) \quad \mathbb{E}_{\tilde{\nu}_V^a} \left((\cosh(u^{(n)}(\delta_n, i) - u^{(n)}(\delta_n, y_n)))^8 \right) \leq 2.$$

By Cauchy-Schwartz inequality, and by (9.6) and (9.8), we get that

$$\mathbb{E}_{\tilde{\nu}_V^a} \left((\psi^{(n)}(i))^{\pm 4} \right) \leq \mathbb{E}_{\tilde{\nu}_V^a} \left(e^{\pm 8u^{(n)}(\delta_n, y_n)} \right)^{\frac{1}{2}} \mathbb{E}_{\tilde{\nu}_V^a} \left(e^{\pm 8(u^{(n)}(\delta_n, i) - u^{(n)}(\delta_n, y_n))} \right)^{\frac{1}{2}} \leq C_{\pm}$$

for some constant $C_{\pm} > 0$ independent of n . From this we deduce by Fatou's lemma for all i, j in \mathbb{Z}^d , $i \sim j$,

$$\mathbb{E}_{\tilde{\nu}_V^a} \left(((W_{i,j}\psi(i)\psi(j))^{\pm 1}) \right) \leq \mathbb{E}_{\tilde{\nu}_V^a} \left((W_{i,j})^{\pm 2} \right)^{\frac{1}{2}} \mathbb{E}_{\tilde{\nu}_V^a} \left((\psi(0))^{\pm 4} \right)^{\frac{1}{2}} < \infty,$$

for a large enough.

The proof is very similar in the VRJP case, and uses Theorem 1 of [10]. As previously, the estimate is valid in the case we are interested in, that is for the graph \mathcal{G}_n , rooted at δ_n , and for $x \in \mathbb{Z}^d$, $y = y_n$ for n large enough. \square

Acknowledgment. We are very grateful to an anonymous referee for valuable comments.

REFERENCES

- [1] Omer Angel, Nicholas Crawford, and Gady Kozma. Localization for linearly edge reinforced random walks. *Duke Mathematical Journal*, 163(5):889–921, 2014.
- [2] Anne-Laure Basdevant and Arvind Singh. Continuous-time vertex reinforced jump processes on Galton–Watson trees. *The Annals of Applied Probability*, 22(4):1728–1743, 2012.
- [3] Andrea Collecchio. Limit theorems for vertex-reinforced jump processes on regular trees. *Electron. J. Probab*, 14(66):1936–1962, 2009.
- [4] Don Coppersmith and Persi Diaconis. Random walk with reinforcement. *Unpublished manuscript*, pages 187–220, 1987.
- [5] Burgess Davis and Stanislav Volkov. Vertex-reinforced jump processes on trees and finite graphs. *Probability theory and related fields*, 128(1):42–62, 2004.
- [6] Anna De Masi, Pablo A Ferrari, Sheldon Goldstein, and William David Wick. An invariance principle for reversible Markov processes. Applications to random motions in random environments. *Journal of Statistical Physics*, 55(3-4):787–855, 1989.
- [7] Margherita Disertori, Franz Merkl, and Silke W. W. Rolles. A supersymmetric approach to martingales related to the vertex-reinforced jump process. *preprint arXiv:1511.07157*, 2015.
- [8] Margherita Disertori, Christophe Sabot, and Pierre Tarres. Transience of edge-reinforced random walk. *Communications in Mathematical Physics*, 339(1):121–148, 2015.
- [9] Margherita Disertori and Tom Spencer. Anderson localization for a supersymmetric sigma model. *Communications in Mathematical Physics*, 300(3):659–671, 2010.
- [10] Margherita Disertori, Tom Spencer, and Martin R Zirnbauer. Quasi-diffusion in a 3D supersymmetric hyperbolic sigma model. *Communications in Mathematical Physics*, 300(2):435–486, 2010.
- [11] Paul R. Halmos. *Measure Theory*. D. Van Nostrand Company, Inc., New York, N. Y., 1950.

- [12] M. S. Keane and S. W. W. Rolles. *Edge-reinforced random walk on finite graphs*, volume 52 of *Verh. Afd. Natuurkd. 1. Reeks. K. Ned. Akad. Wet. R. Neth. Acad. Arts Sci.*, Amsterdam, 2000.
- [13] Gérard Letac. Personal communication. 2015.
- [14] Russell Lyons and Yuval Peres. *Probability on Trees and Networks*. Cambridge University Press. In preparation. Current version available at <http://pages.iu.edu/~rdlyons/>, 2015.
- [15] Franz Merkl and Silke W. W. Rolles. Bounding a random environment for two-dimensional edge-reinforced random walk. *Electron. J. Probab.*, 13:no. 19, 530–565, 2008.
- [16] Franz Merkl and Silke WW Rolles. A random environment for linearly edge-reinforced random walks on infinite graphs. *Probability theory and related fields*, 138(1-2):157–176, 2007.
- [17] Franz Merkl and Silke WW Rolles. Recurrence of edge-reinforced random walk on a two-dimensional graph. *The Annals of Probability*, pages 1679–1714, 2009.
- [18] Robin Pemantle. Phase transition in reinforced random walk and RWRE on trees. *The Annals of Probability*, pages 1229–1241, 1988.
- [19] R.J. Plemmons and A Berman. *Nonnegative matrices in the mathematical sciences*. Academic Press, New York, 1979.
- [20] Christophe Sabot and Pierre Tarrès. Edge-reinforced random walk, vertex-reinforced jump process and the supersymmetric hyperbolic sigma model. *J. Eur. Math. Soc.*, 17(9):2353–2378, 2015.
- [21] Christophe Sabot and Pierre Tarres. Inverting Ray-Knight identity. *Probab. Theory Related Fields*, 165(3-4):559–580, 2016.
- [22] Christophe Sabot, Pierre Tarrès, and Xiaolin Zeng. The vertex reinforced jump process and a random Schrödinger operator on finite graphs. *Ann. Probab.*, 45(6A):3967–3986, 2017.
- [23] Laurent Tournier. A note on the recurrence of edge reinforced random walks. *arXiv preprint arXiv:0911.5255*, 2009.
- [24] Martin R Zirnbauer. Fourier analysis on a hyperbolic supermanifold with constant curvature. *Communications in mathematical physics*, 141(3):503–522, 1991.

UNIVERSITÉ DE LYON, UNIVERSITÉ LYON 1, INSTITUT CAMILLE JORDAN, CNRS UMR 5208, 43,
BOULEVARD DU 11 NOVEMBRE 1918, 69622 VILLEURBANNE CEDEX, FRANCE
E-mail address: `sabot@math.univ-lyon1.fr`

108 SCHREIBER BUILDING, SCHOOL OF MATHEMATICS, TEL AVIV UNIVERSITY, P.O.B. 39040, RAMAT
AVIV, TEL AVIV 69978, ISRAEL
E-mail address: `xzeng@math.univ-lyon1.fr`