



HAL
open science

The Horn Problem for Real Symmetric and Quaternionic Self-Dual Matrices

Robert Coquereaux, Jean-Bernard Zuber

► **To cite this version:**

Robert Coquereaux, Jean-Bernard Zuber. The Horn Problem for Real Symmetric and Quaternionic Self-Dual Matrices. 2018. hal-01871610v1

HAL Id: hal-01871610

<https://hal.science/hal-01871610v1>

Preprint submitted on 11 Sep 2018 (v1), last revised 1 Feb 2019 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The Horn Problem for Real Symmetric and Quaternionic Self-Dual Matrices

Robert Coquereaux

*Aix Marseille Univ, Université de Toulon, CNRS, CPT, Marseille, France
Centre de Physique Théorique*

and

Jean-Bernard Zuber

*Sorbonne Université, UMR 7589, LPTHE, F-75005, Paris, France
& CNRS, UMR 7589, LPTHE, F-75005, Paris, France*

Abstract

Horn's problem, i.e., the study of the eigenvalues of the sum $C = A + B$ of two matrices, given the spectrum of A and of B , is re-examined, comparing the case of real symmetric, complex Hermitian and self-dual quaternionic 3×3 matrices. In particular, what can be said on the probability distribution function (PDF) of the eigenvalues of C if A and B are independently and uniformly distributed on their orbit under the action of, respectively, the orthogonal, unitary and symplectic group ?

While the two latter cases (Hermitian and quaternionic) may be studied by use of explicit formulae for the relevant orbital integrals, the case of real symmetric matrices is much harder. It is also quite intriguing, since numerical experiments reveal the occurrence of singularities where the PDF of the eigenvalues diverges.

Here we show that the computation of the PDF of the symmetric functions of the eigenvalues for traceless 3×3 matrices may be carried out in terms of algebraic functions –roots of quartic polynomials– and their integrals. The computation is carried out in detail in a particular case, and reproduces the expected singular patterns. The divergences are of logarithmic or inverse power type.

We also relate this PDF to the (rescaled) structure constants of zonal polynomials and introduce a zonal analogue of the Weyl $SU(n)$ characters.

Keywords: Horn problem. Honeycombs. Polytopes. Zonal polynomials. Littlewood–Richardson coefficients.

Mathematics Subject Classification 2010: 17B08, 17B10, 22E46, 43A75, 52Bxx

Introduction

Recall what Horn's problem is: given two $n \times n$ matrices A and B of given spectrum of eigenvalues, what can be said about the spectrum of their sum $C = A + B$? The problem has been addressed by many authors, see [1] for a review and references. For Hermitian matrices, the support of the spectrum of C has been completely determined after some crucial work by Klyashko [2] and by Knutson and Tao [3, 4, 5].

In a recent paper [6], a more specific question was considered: given $\alpha = \{\alpha_1 \geq \alpha_2 \geq \dots \alpha_n\}$ and $\beta = \{\beta_1 \geq \beta_2 \geq \dots \beta_n\}$, take the Hermitian matrices A and B uniformly and independently distributed on the orbit of $\text{diag}(\alpha)$ and $\text{diag}(\beta)$ under the action of the $SU(n)$ group. The probability distribution function (PDF) of the eigenvalues γ of $C = A + B$ may then be computed, see also [7, 8].

The same question may, however, be raised if instead of Hermitian matrices, one considers other classes of matrices and the action of an appropriate group. The case of skew-symmetric real matrices was considered in [6], but more interesting is that of real symmetric matrices and the action of the orthogonal group $SO(n)$.

Let us start with a numerical experiment with specific 3×3 matrices. Since $\text{tr} C = \text{tr} A + \text{tr} B$, only two eigenvalues of C , say γ_1 and γ_2 are linearly independent. Then compare the distribution of points in the (γ_1, γ_2) -plane for the three cases of

- (a) orbits of real symmetric matrices A and B equivalent to $J_z := \text{diag}(1, 0, -1)$ under conjugation by the orthogonal group $SO(3)$;
- (b) orbits of such matrices, regarded as Hermitian, under conjugation by the unitary group $SU(3)$;
- (c) orbits of such matrices, regarded as quaternionic self-dual, (i.e., $A \otimes \mathbb{I}_2, B \otimes \mathbb{I}_2$ as 6×6 matrices), under conjugation by the unitary symplectic group $USp(3)$.

Following the nomenclature introduced by Dyson, we label these three cases by the index $\underline{\beta} = 1, 2$ or 4 , respectively¹.

Some features appear clearly on the plots and histograms of Fig. 1:

- (i) the PDF vanishes faster on the boundaries of the Horn domain as $\underline{\beta}$ increases;
 - (ii) the non-analyticities are stronger and stronger as $\underline{\beta}$ decreases;
 - (iii) these singularities seem to appear at the same place in the (γ_1, γ_2) plane (for α and β fixed).
- In the Hermitian case (and in the quaternionic case as well, see below), it is known that the PDF is a piece-wise polynomial function. The plots of Fig. 1(a) show that this cannot be true for real symmetric matrices.

It should be emphasized that these general features do not depend on the explicit case we have chosen. Similar singular patterns have been observed in numerical experiments with other matrices A and B , see Fig. 7 in [6].

The aim of this paper is to compare the three cases, to reproduce analytically the previous empirical observations and in particular to analyse the location and nature of the singularities that occur in the symmetric case. After a brief review of known results on the relevant orbital integrals (sect. 1), we treat rapidly the easy case of quaternionic self-dual matrices (sect. 2), before turning to the more challenging case of real symmetric matrices in sect. 3 and 4. In sect. 3, it is shown that for 3×3 traceless matrices, the introduction of the two symmetric functions p and q of the three eigenvalues (of vanishing sum) simplifies matters: one may express the PDF $\rho(p, q)$ in terms of roots of some polynomial equations and integral thereof, see (33) below. The actual computation is carried out again for our pet example of $\alpha = \beta = (1, 0, -1)$ in sect. 4. In particular we reproduce and analyze in sect. 4.4 and 4.5 the singularities that are apparent in Fig. 1(a). Finally in sect. 5, we show that this function $\rho(p, q)$ is related to the distribution of the (rescaled) structure constants of zonal polynomials, thus elaborating on a claim of [9].

¹The reader should not confuse this index $\underline{\beta}$ with the multiplet β of eigenvalues of the B matrix.




Figure 1: Plots and histograms of (γ_1, γ_2) for $A \sim B \sim J_z$ in the three cases: (a) orbits under $\text{SO}(3)$, (b) under $\text{SU}(3)$, (c) $\text{USp}(3)$. Plots made of 10^4 points, histograms of 10^6 points.

1 The orbital integrals for $\underline{\beta} = 1, 2, 4$

Consider the *orbital integrals*

$$\mathcal{I}_{\underline{\beta}}(X, A) = \int_{G_{\underline{\beta}}} d\Omega \exp[\text{tr } X\Omega A\Omega^\dagger] \quad (1)$$

1. $\underline{\beta} = 1$, $\Omega \in G_1 := \text{SO}(n)$, X, A real symmetric matrices;
2. $\underline{\beta} = 2$, $\Omega \in G_2 := \text{U}(n)$, X, A complex Hermitian matrices;
3. $\underline{\beta} = 4$, $\Omega \in G_4 := \text{USp}(n)$, X, A real quaternionic self-dual matrices.

In fact each of these integrals depends only on the eigenvalues α_i , resp. x_i , of A , resp. X , and by a small abuse of notations, we write them $\mathcal{I}_{\underline{\beta}}(x, \alpha)$ in the following.

For $\underline{\beta} = 2$, this orbital integral is well known, [10, 11]

$$\mathcal{I}_2(x, \alpha) = \prod_{p=1}^{n-1} p! \sum_{P \in S_n} \frac{1}{\Delta(x)\Delta(\alpha_P)} e^{\sum x_j \alpha_{Pj}} = \prod_{p=1}^{n-1} p! \frac{\det e^{x_i \alpha_j}}{\Delta(x)\Delta(\alpha)}. \quad (2)$$

Here and below, Δ with no subscript denotes the Vandermonde determinant $\Delta(x) = \prod_{i < j} (x_i - x_j)$.

For $\underline{\beta} = 4$ and for a given n , as shown by [13], we have closed formulae

$$\mathcal{I}_4(x, \alpha) = \hat{\kappa}_n \sum_{P \in S_n} \frac{1}{\Delta(x)^3 \Delta(\alpha_P)^3} e^{\sum x_j \alpha_{Pj}} f_n(\tau(\alpha, P)), \quad (3)$$

where f_n a polynomial in the variables

$$\tau(\alpha, P) = \{\tau_{ij} := (x_i - x_j)(\alpha_{Pi} - \alpha_{Pj}) \mid 1 \leq i < j \leq n\}, \quad (4)$$

and $\hat{\kappa}_n$ is an α - and x -independent constant. For example for $n = 2$, $\hat{\kappa}_2 = -12$ and $f_2(\tau(\alpha, P)) = 1 - \frac{1}{2}\tau_{12}$ while for $n = 3$, $\hat{\kappa}_3 = -2 \times 3! \times 6!$ and

$$f_3(\tau(\alpha, P)) = 1 - \frac{1}{3}(\tau_{12} + \tau_{13} + \tau_{23}) + \frac{1}{6}(\tau_{12}\tau_{13} + \tau_{13}\tau_{23} + \tau_{23}\tau_{12}) - \frac{1}{12}\tau_{12}\tau_{13}\tau_{23}.$$

These closed formulae allow an explicit computation of the PDF, see [6] and below, sect. 2. In contrast, for $\beta = 1$, i.e., for real symmetric matrices under the action of the orthogonal group $\text{SO}(n)$, the best that can be achieved is an expansion in zonal polynomials [12, 13, 9]

$$\mathcal{I}_1(x, \alpha) = \sum_{m=0}^{\infty} \frac{1}{m! \prod_{q=0}^{m-1} (1 + 2q)} \sum_{\kappa \vdash m} c(\kappa) \frac{Z(\kappa)(x)Z(\kappa)(\alpha)}{Z(\kappa)(I)} \quad (5)$$

where the second sum runs over partitions κ of m with no more than n parts, and $c(\kappa)$ is a constant which depends on the normalization of the $Z(\kappa)$'s, see sect. 5. If the zonal polynomials in the above formula are written with the so-called James normalization, one can use the values of $Z(\kappa)(I)$ and of the coefficient $c(\kappa)$ tabulated by James up to $m = 4$, for all n (the dimension of I) in the Appendix of [12], p.157; one can also use commands contained in the Mathematica package [14]. The infinite sum in the previous expansion, however, makes the computation of the PDF intractable (as far as we can see), and we will have to follow another route, see below section 3.

In section 5, however, we return to this formulation in terms of zonal polynomials and show a connection between the PDF and the distribution of (rescaled) ‘‘zonal multiplicities’’ i.e., appropriate structure constants of zonal polynomials.

2 Quaternionic case for $n = 3$

We start with a warming up exercise: compute the PDF of the eigenvalues γ for a sum of two self-dual quaternionic matrices that are independently and uniformly distributed on their orbit under the action of the unitary symplectic group. The computation of the PDF of the γ 's follows the same lines as that in the Hermitian case. We will therefore be a bit sketchy in its derivation, referring the reader to [6] for details of the computation.

Up to an overall factor, this PDF is given by the integral of three orbital integrals of the type \mathcal{I}_4 in (3).

$$\mathbf{p}(\gamma|\alpha, \beta) = \frac{\kappa_n^2}{(2\pi)^N} \Delta(\gamma)^4 \int \prod_{i=1}^n dx_i \Delta(x)^4 \mathcal{I}_4(x, i\alpha) \mathcal{I}_4(x, i\beta) \mathcal{I}_4(x, -i\gamma), \quad (6)$$

where $N = n(2n-1)$ is the number of independent matrix elements of a self-dual quaternion matrix and $\kappa_n = \frac{(2\pi)^{(N-n)/2}}{n!(2n-1)!(\frac{1}{2}n(n-1))!}$ stems from the change from those N variables to the n eigenvalues. Thus

$$\mathbf{p}(\gamma|\alpha, \beta) = \frac{2\kappa_n^2 \hat{\kappa}_n^3}{(2\pi)^N} (-1)^{n(n-1)/2} \pi^n \delta\left(\sum_k (\alpha_k + \beta_k - \gamma_k)\right) \frac{\Delta(\gamma)}{\Delta(\alpha)^3 \Delta(\beta)^3} \mathcal{J}_n \quad (7)$$

$$\mathcal{J}_n = \frac{n!}{i^{n(n-1)/2} \pi^{n-1}} \sum_{P, P' \in \mathcal{S}_n} \varepsilon_P \varepsilon_{P'} \int \frac{d^{n-1}u}{\Delta'(u)^5} \prod_{j=1}^{n-1} e^{i u_j A_j(P, P', I)} f_n(\tau(i\alpha, P)) f_n(\tau(i\beta, P')) f_n(\tau(-i\gamma, I)) \quad (8)$$

with $u_j := x_j - x_{j+1}$, $\Delta'(u) := \prod_{1 \leq i < j \leq n} (u_i + u_{i+1} + \dots + u_{j-1})$ and

$$A_j(P, P', P'') = \sum_{k=1}^j (\alpha_{P(k)} + \beta_{P'(k)} - \gamma_{P''(k)}). \quad (9)$$




Figure 2: Comparing the histogram of 10^4 points with the PDF of (10) for (left) $\alpha = \beta = (1, -1)$ and (right) $\alpha = (1, -1)$, $\beta = (2, -2)$.

The power 5 of $\Delta'(u)$ results from 3×3 (three denominators $\Delta^3(x)$), minus 4 from the Jacobian $\Delta^4(x)$.

With the little extra complication caused by the f_n factors of (3), the integration may be carried out as in [6], namely by partial fraction decomposition of the integrand and repeated use of the Dirichlet formula for the Cauchy principal value

$$P \int_{\mathbb{R}} \frac{du}{u^r} e^{i u A} = i \pi \frac{(i A)^{r-1}}{(r-1)!} \epsilon(A),$$

with ϵ the sign function. The result, though cumbersome, is completely explicit.

For $n = 2$, we leave as an exercise for the reader to check that for $\alpha_2 = -\alpha_1$, $\beta_2 = -\beta_1$, we have $\mathbf{p}(\gamma|\alpha, \beta) = 6 \frac{\Delta(\gamma)}{\Delta^3(\alpha)\Delta^3(\beta)} \delta(\gamma_1 + \gamma_2) \mathcal{J}_2$ with

$$\begin{aligned} \mathcal{J}_2 &= \frac{1}{4} \left(-(\alpha_1^2 - \beta_1^2)^2 + 2(\alpha_1^2 + \beta_1^2)\gamma_1^2 - \gamma_1^4 \right) \left(\epsilon(\gamma_1 - \alpha_1 + \beta_1) + \epsilon(\gamma_1 + \alpha_1 - \beta_1) - \epsilon(\gamma_1 - \alpha_1 - \beta_1) - \epsilon(\gamma_1 + \alpha_1 + \beta_1) \right) \\ &= \frac{1}{2} \left(-(\alpha_1^2 - \beta_1^2)^2 + 2(\alpha_1^2 + \beta_1^2)\gamma_1^2 - \gamma_1^4 \right) (\mathbf{1}_I(\gamma_1) - \mathbf{1}_{-I}(\gamma_1)). \end{aligned} \quad (10)$$

showing that the spectrum of γ_1 is supported by two segments $I := [|\alpha_1 - \beta_1|, \alpha_1 + \beta_1]$ and $-I$, or only by the former if one imposes $\gamma_2 \leq \gamma_1$. Compare with the analogous formulae obtained for Hermitian and real symmetric matrices in [6]. In Fig. 2, we show the agreement with the histogram made of 10^4 for two pairs $\alpha = \beta = (1, -1)$ and $\alpha = (1, -1)$, $\beta = (2, -2)$.

For $n = 3$,

$$\mathbf{p}(\gamma|\alpha, \beta) = 8640 \frac{\Delta(\gamma)}{\Delta^3(\alpha)\Delta^3(\beta)} \delta\left(\sum_k \gamma_k - \alpha_k - \beta_k\right) \mathcal{J}_3 \quad (11)$$

where \mathcal{J}_3 is a piecewise polynomial of degree 13 in γ_1 and γ_2 ; indeed

$$\mathcal{J}_3 = \frac{3!}{i^3 \pi^2} \sum_{P, P' \in S_3} \varepsilon_P \varepsilon_{P'} \int \frac{du_1 du_2}{(u_1 u_2 (u_1 + u_2))^5} e^{i(u_1 A_1 + u_2 A_2)} F(u_1, u_2), \quad (12)$$

($F(u_1, u_2)$ stands for the product $f_3(\tau(i\alpha, P))f_3(\tau(i\beta, P'))f_3(\tau(-i\gamma, I))$ in (8), see also (4)), whence, by homogeneity, a behavior as $[\gamma]^{15-2=13}$.

\mathcal{J}_3 is of differentiability class C^2 : indeed as u_1 , say, goes to infinity, the integrand behaves as $\frac{1}{u_1} e^{i A_1 u_1}$ (up to subdominant terms), so that the integral is twice continuously differentiable with respect to A_1 (or γ), hence of class C^2 . Non-analyticities are expected (and do occur) along the boundaries of the Horn polygon and across the same singular lines (or half-lines) as in the Hermitian case, namely

$$\begin{aligned} \gamma_2 &= \alpha_2 + \beta_2; \\ \gamma_1 &= \alpha_1 + \beta_2, \gamma_2 \geq \alpha_3 + \beta_1; \gamma_2 = \alpha_3 + \beta_1, \gamma_1 \leq \alpha_1 + \beta_2; \gamma_3 = \alpha_2 + \beta_3, \gamma_1 \geq \alpha_1 + \beta_2, \\ &\text{the same with } \alpha \leftrightarrow \beta. \end{aligned} \quad (13)$$




Figure 3: The four sectors in the Horn polygon of the quaternionic case for $\alpha = \beta = \{1, 0, -1\}$.

For our favorite example of $A \sim B \sim J_z = \text{diag}(1, 0, -1)$, the function has four sectors of piecewise polynomiality, labelled by $i = 1, \dots, 4$ according to Fig. 3. In each of these four sectors the function takes the form

$$\mathcal{J}_3|_{\text{sector } i} = -\frac{2 \times 5!}{13!} \mathcal{P}_i(\gamma_1, \gamma_2) \quad (14)$$

where

$$\mathcal{P}_i(\gamma_1, \gamma_2) = p_i(\gamma_1, \gamma_2)^3 \tilde{p}_i(\gamma_1, \gamma_2), \quad (15)$$

$$p_i(\gamma_1, \gamma_2) = \begin{cases} (2 - \gamma_1 - \gamma_2) & \text{in sector 1, i.e., } \gamma_1 \geq 1, \gamma_2 \geq 0, 2 - \gamma_1 - \gamma_2 \geq 0 \\ (2 - \gamma_1) & \text{in sector 2, i.e., } \gamma_1 \leq 2, \gamma_2 \leq 0, \gamma_1 + \gamma_2 \geq 1 \\ (\gamma_2 - \gamma_3) = (\gamma_1 + 2\gamma_2) & \text{in sector 3, i.e., } \gamma_2 \leq 0, \gamma_1 + \gamma_2 \leq 1, \gamma_1 + 2\gamma_2 \geq 0 \\ (\gamma_1 - \gamma_2) & \text{in sector 4, i.e., } \gamma_1 \leq 1, \gamma_2 \geq 0, \gamma_1 - \gamma_2 \geq 0, \end{cases}$$

and each of the $\tilde{p}_i(\gamma_1, \gamma_2) = 40\gamma_1^{10} + \dots$ is too cumbersome to be given here ².

Note that the form (15) guarantees that \mathcal{J}_3 vanishes “cubically” on each boundary of the polygon. This explains one of the features observed on the plots of Fig. 1 (c). As a side remark, we note that $\tilde{p}_4(\gamma_1, \gamma_2)$ is symmetric under $\gamma_1 \leftrightarrow \gamma_2$, and $\tilde{p}_3(\gamma_1, \gamma_2)$ is symmetric under $\gamma_2 \leftrightarrow \gamma_3 = -\gamma_1 - \gamma_2$. One may also compute the transition functions between adjacent domains:

$$\begin{aligned} \mathcal{P}_1 - \mathcal{P}_2 &= \gamma_2^3 (\dots) \\ \mathcal{P}_2 - \mathcal{P}_3 &= (1 - \gamma_1 - \gamma_2)^3 (\dots) \\ \mathcal{P}_3 - \mathcal{P}_4 &= \gamma_2^3 (\dots) \\ \mathcal{P}_4 - \mathcal{P}_1 &= (1 - \gamma_1)^3 (\dots) \end{aligned} \quad (16)$$

where in each case, the ellipsis stands for a polynomial of degree 10 that we refrain from displaying. Just like on the boundary, the transition functions vanish cubically along the non-analyticity lines. This is in agreement with the differentiability argument above.

The resulting PDF is plotted in Fig. 4 (a) and compared to the “experimental” histogram of Fig. 4 (b).

A good check of our calculation is that the integral of the PDF over the whole domain of γ 's, made of $3!$ copies of the Horn polygon $\tilde{\mathbf{H}}_{\alpha, \beta}$, (see below in (19)), is indeed 1, thanks to $\int_{\tilde{\mathbf{H}}_{\alpha, \beta}} d\gamma_1 d\gamma_2 \Delta(\gamma) \mathcal{J}_3 = \frac{1}{810}$.

² Suggestion : This may be found on the web site http://www.lpthe.jussieu.fr/~zuber/Z_Unpub2.html. Other data that are not given explicitly in this paper may be found either on this web site or on <http://www.cpt.univ-mrs.fr/~coque/Varia/JackZonalSchurResults.html>




Figure 4: Comparing the result of (7–15) with the “experimental” histogram.

3 Computing the PDF for a sum for two real symmetric matrices

As explained above, in contrast with the unitary or the symplectic groups, there is no closed expression for the orbital integral on the orthogonal group and the formula

$$p(\gamma|\alpha, \beta) = \text{const.} |\Delta(\gamma)| \int \prod_{i=1}^n dx_i |\Delta(x)| \mathcal{I}_1(x, i\alpha) \mathcal{I}_1(x, i\beta) \mathcal{I}_1(x, -i\gamma). \quad (17)$$

remains intractable. We thus have to resort to a different approach. We first trade the PDF of eigenvalues γ_i of the sum $C = A + B$ for the PDF $\rho(p, q)$ of symmetric functions of the γ 's, see (20) below. The relation between the two is given below in (24).

3.1 The support of the symmetric functions P and Q

For 3×3 real symmetric *traceless* matrices A and B , the characteristic polynomial of their sum C reads³

$$\det(z\mathbb{I} - C) = \det(z\mathbb{I} - A - B) = z^3 + Pz + Q. \quad (18)$$

The support of the eigenvalues γ_i of C is a convex polygon $\tilde{\mathbf{H}}_{\alpha, \beta}$ in \mathbb{R}^2 , defined in terms of the eigenvalues α_i and β_i of A and B by the same Horn inequalities as in the Hermitian case [1]:

$$\begin{aligned} \gamma_{3min} := \alpha_3 + \beta_3 &\leq \gamma_3 \leq \min(\alpha_1 + \beta_3, \alpha_2 + \beta_2, \alpha_3 + \beta_1) =: \gamma_{3max} \\ \gamma_{2min} := \max(\alpha_2 + \beta_3, \alpha_3 + \beta_2) &\leq \gamma_2 \leq \min(\alpha_1 + \beta_2, \alpha_2 + \beta_1) =: \gamma_{2max} \\ \gamma_{1min} := \max(\alpha_1 + \beta_3, \alpha_2 + \beta_2, \alpha_3 + \beta_1) &\leq \gamma_1 \leq \alpha_1 + \beta_1 =: \gamma_{1max} \\ \text{as well as (by convention)} &\quad \gamma_3 \leq \gamma_2 \leq \gamma_1. \end{aligned} \quad (19)$$

This translates for the symmetric functions $P = \gamma_1\gamma_2 + \gamma_2\gamma_3 + \gamma_3\gamma_1$ and $Q = -\gamma_1\gamma_2\gamma_3$ into a curvilinear polygon, whose sides are either segments of lines $cp + q = -c^3$ (for sides of $\mathbf{H}_{\alpha, \beta}$ where some $\gamma_i = c$) or arcs of the cubic $4p^3 + 27q^2 = 0$ for sides of $\tilde{\mathbf{H}}_{\alpha, \beta}$ where $\gamma_i = \gamma_j$. See Fig. 5 for an example with $\alpha = \{11, -1, -10\}$, $\beta = \{7, 4, -11\}$.

It has been suggested [6] that the Horn problem for real symmetric matrices not only shares the same support as in the Hermitian case (for given α and β), as proved by Fulton [1], but has also singularities at the same locations, although the latter look much stronger. Thus we expect

³The assumption of tracelessness is harmless: one may always translate A and B by a multiple of the identity matrix so as to enforce it, at the price of shifting the eigenvalues of their sum C by a common real number.




Figure 5: The Horn-tensor polygon $\tilde{\mathbf{H}}_{\alpha\beta}$ in the (γ_1, γ_2) plane, and the curlinear polygon in the (p, q) plane, drawn here for $\alpha = \{11, -1, -10\}$, $\beta = \{7, 4, -11\}$. The dashed (blue, red and green) lines are the expected loci of singularities of the PDF. (A histogram of eigenvalues γ for an equivalent configuration of (α, β) has appeared in [6], Fig. 7.)

singularities to occur for 3×3 matrices along the same lines (or half-lines) as in (13)⁴. These lines are illustrated in Fig. 5 for $\alpha = \{11, -1, -10\}$, $\beta = \{7, 4, -11\}$ in the (γ_1, γ_2) and in the (p, q) planes.

3.2 The statistics of the symmetric functions P and Q

For 3×3 real *traceless* symmetric matrices A and B , of respective eigenvalues α and β , the characteristic polynomial of their sum C reads

$$\det(z \mathbb{I}_3 - C) = \det(z \mathbb{I}_3 - \text{diag}(\alpha) - \mathcal{R} \text{diag}(\beta) \mathcal{R}^T) = z^3 + P(\mathcal{R})z + Q(\mathcal{R}).$$

For given α 's and β 's, and \mathcal{R} regarded as a random variable uniformly distributed in $\text{SO}(3)$ (in the sense of the Haar measure), $P(\mathcal{R})$ and $Q(\mathcal{R})$ are also random variables, whose PDF may be written as

$$\rho(p, q) = \mathbb{E}(\delta(P - p)\delta(Q - q)) = \int D\mathcal{R} \delta(P(\mathcal{R}) - p) \delta(Q(\mathcal{R}) - q). \quad (20)$$

We parametrize \mathcal{R} in terms of Euler angles

$$\mathcal{R} = \mathcal{R}_z(\phi) \mathcal{R}_y(\theta) \mathcal{R}_z(\psi)$$

with

$$0 \leq \phi \leq 2\pi, \quad 0 \leq \theta \leq \pi, \quad 0 \leq \psi \leq 2\pi$$

and the normalized Haar measure is then

$$D\mathcal{R} = \frac{1}{8\pi^2} \sin \theta d\theta d\phi d\psi. \quad (21)$$

⁴There may be non-analyticities at other places, though.

$P_p := P - p$ and $Q_q := Q - q$ are then degree 2 polynomials of the variable $c := \cos \theta$, of the form

$$\begin{aligned}
P_p(c) &= -\alpha_1^2 - \alpha_1\alpha_2 - \alpha_2^2 - \beta_1^2 - \beta_1\beta_2 - \beta_2^2 + \alpha_1(\beta_2 - \beta_1) + \alpha_2(\beta_1 + 2\beta_2) \\
&\quad + x^2(\alpha_1 - \alpha_2)(\beta_2 + 2\beta_1) + y^2(\alpha_2 + 2\alpha_1)(\beta_1 - \beta_2) - x^2y^2(\alpha_1 - \alpha_2)(\beta_1 - \beta_2) - p \\
&\quad + 2xy(\alpha_1 - \alpha_2)(\beta_1 - \beta_2) \sin \phi \sin \psi \mathbf{c} \\
&\quad - (\alpha_1 + 2\alpha_2 + \alpha_1x^2 - \alpha_2x^2) (\beta_1 + 2\beta_2 + \beta_1y^2 - \beta_2y^2) \mathbf{c}^2 \\
Q_q(c) &= (\alpha_1 + \beta_1)(\alpha_1 + \alpha_2 - \beta_2)(\alpha_2 - \beta_1 - \beta_2) + (\alpha_1 - \alpha_2)(\alpha_1 + \alpha_2 - \beta_2)(2\beta_1 + \beta_2)x^2 \\
&\quad - (2\alpha_1 + \alpha_2)(\alpha_2 - \beta_1 - \beta_2) (\beta_1 - \beta_2)y^2 - (\alpha_1 - \alpha_2)(\beta_1 - \beta_2)(\alpha_1 + \alpha_2 + \beta_1 + \beta_2)x^2y^2 - q \\
&\quad + 2xy(\alpha_1 - \alpha_2)(\beta_1 - \beta_2)(\alpha_1 + \alpha_2 + \beta_1 + \beta_2) \sin \phi \sin \psi \mathbf{c} \\
&\quad + ((\alpha_1 + 2\alpha_2)(\alpha_1 + \beta_1)(\beta_1 + 2\beta_2) - (\alpha_1 - \alpha_2)(\alpha_1 + \alpha_2 - \beta_1)(\beta_1 + 2\beta_2)x^2 \\
&\quad + (\alpha_1 + 2\alpha_2)(\alpha_1 - \beta_1 - \beta_2)(\beta_1 - \beta_2)y^2 - (\alpha_1 - \alpha_2)(\beta_1 - \beta_2)(\alpha_1 + \alpha_2 + \beta_1 + \beta_2)x^2y^2) \mathbf{c}^2
\end{aligned} \tag{22}$$

where $x := \cos \phi$ and $y := \cos \psi$.

Note the peculiar feature of these polynomials in c : their degree 0 and 2 terms are (degree 1) polynomials in the variables x^2 and y^2 , while their degree 1 term in c is of the form $xy \sin \phi \sin \psi$, up to a (α - and β - dependent) factor.

Note also that these expressions are π -periodic in ϕ and ψ , making it possible to restrict these angles to the interval $(0, \pi)$ where their sine is non negative. This will be implicit in the following.

Thus

$$\rho(p, q) = \frac{1}{2\pi^2} \int_0^\pi d\phi \int_0^\pi d\psi \int_{-1}^1 dc \delta(Q_q)\delta(P_p). \tag{23}$$

The PDF for the independent variables γ_1, γ_2 then follows simply:

$$\mathbf{p}(\gamma_1, \gamma_2) = |\Delta(\gamma)| \rho(p, q). \tag{24}$$

3.3 Reducing $\delta(Q_q)\delta(P_p)$ to $\delta(R)$

In this subsection, we show that $\int dc \delta(Q_q)\delta(P_p)$ may be reduced, up a factor, to a single $\delta(R)$, where R is the *resultant* of the two polynomials $P_p(c)$ and $Q_q(c)$. We shall make repeated use of two classical identities [15]:

– for $f(t)$ a function with a finite number of “simple” zeros t_i (i.e., such that $f'(t_i) \neq 0$),

$$\delta(f(t)) = \sum_{t_i} \frac{\delta(t - t_i)}{|f'(t_i)|}; \tag{25}$$

– for f and g two functions with no common zero,

$$\delta(f(t)g(t)) = \frac{\delta(f(t))}{|g(t)|} + \frac{\delta(g(t))}{|f(t)|}, \tag{26}$$

to which we may then apply the previous identity. Then starting from the product $\delta(Q_q)\delta(P_p)$, we assume that the discriminant Δ_Q of Q_q is positive, in such a way that the roots $c_{1,2}$ are real and distinct, and we may write

$$\delta(Q_q) = \frac{1}{\sqrt{\Delta_Q}}(\delta(c - c_1) + \delta(c - c_2))$$

and

$$\int dc \delta(Q_q)\delta(P_p) = \frac{1}{\sqrt{\Delta_Q}}(\delta(P_p(c_1)) + \delta(P_p(c_2)))$$

(where it is understood that the delta’s act on functions of the remaining variables ϕ and ψ or x and y).

We want to compare this expression with $\delta(R)$ where, as said above, R is the resultant of the two polynomials $P_p(c)$ and $Q_q(c)$. If a and a' are the coefficients of terms of degree 2 of the polynomials Q_q and P_p , respectively, and c'_1, c'_2 the roots of the latter, thus

$$P_p(c) = a'(c - c'_1)(c - c'_2) \quad Q_q(c) = a(c - c_1)(c - c_2),$$

the resultant R defined as $a^2 a'^2 \prod_{i,j=1,2} (c_i - c'_j)$ may also be written as

$$R = a^2 P_p(c_1) P_p(c_2).$$

(For the polynomials of (22), this is a quite cumbersome polynomial of degree 5 in $u = x^2$ and in $z = y^2$, with 4089 α - and β -dependent terms².)

According to (26), one writes

$$\delta(R) = \frac{1}{a^2} \left(\frac{\delta(P_p(c_1))}{|P_p(c_2)|} + \frac{\delta(P_p(c_2))}{|P_p(c_1)|} \right). \quad (27)$$

But $P_p(c_{1,2})$ has the general form

$$P_p(c_{1,2}) = A \pm B\sqrt{\Delta_Q}$$

with A and B functions of x and y . Thus whenever $P_p(c_1)$ vanishes for some (x, y) , i.e., $A = -B\sqrt{\Delta_Q}$, we have $P_p(c_2) = -2B\sqrt{\Delta_Q}$ for those values. And vice versa, if $P_p(c_2)$ vanishes, then $P_p(c_1) = 2B\sqrt{\Delta_Q}$. One may thus rewrite (27) as

$$\delta(R) = \frac{1}{2a^2|B|\sqrt{\Delta_Q}} (\delta(P_p(c_1)) + \delta(P_p(c_2))) = \frac{1}{2a^2|B|} \int dc \delta(Q_q) \delta(P_p), \quad (28)$$

a non trivial and useful identity.

For the polynomials P_p and Q_q of (22), it is easy to compute that $2a^2 B = (xy \sin \phi \sin \psi) \tilde{B}$, with

$$\begin{aligned} \tilde{B} &= 2(\alpha_1 - \alpha_2)(\beta_1 - \beta_2) \times \\ &[(2\alpha_1 + \alpha_2)(\alpha_1 + 2\alpha_2)((\beta_1 - \beta_2)z + (\beta_1 + 2\beta_2)) + (2\beta_1 + \beta_2)(\beta_1 + 2\beta_2)((\alpha_1 - \alpha_2)u + (\alpha_1 + 2\alpha_2))] \end{aligned} \quad (29)$$

while the prefactor $|x||y| \sin \phi \sin \psi$ (recalling that $x = \cos \phi$, $y = \cos \psi$) enables us to change variables to $u = x^2$, $z = y^2$, with the result that

$$\rho(p, q) = \frac{1}{2\pi^2} \int_0^1 dz \int_0^1 du |\tilde{B}| \delta(R). \quad (30)$$

The final transformation of this expression follows from (25). If $u_i(z)$ denote those roots of the polynomial $R(u, z)$ that belong to the interval $[0, 1]$, we may write

$$\rho(p, q) = \frac{1}{2\pi^2} \int_0^1 dz \sum_i \frac{|\tilde{B}(u_i, z)|}{|R'_u(u_i, z)|}. \quad (31)$$

This integral will be studied more explicitly in a particular example in the next section.

Three remarks are in order:

- we have assumed from the start that the discriminant Δ_Q is positive, and this led us to (30). Conversely, the vanishing of R for real values of the variables u and z encompassed in (30) implies that $P_p(c)$ and $Q_q(c)$ have a common root, and this may only be possible if that root is a root of $aP_p - a'Q_q$ which is a degree 1 polynomial in c with real coefficients. Thus the common roots of $P_p(c)$ and $Q_q(c)$ are necessarily real, justifying our assumption that $\Delta_Q \geq 0$;
- the roots c_1 and c_2 have to lie in $[-1, 1]$ for the consistency of the derivation;
- the two functions $P_p(c_1)$ and $P_p(c_2)$ have to have no common zero. Otherwise if that happened at some values (x, y) , both A and B would vanish.

The two latter points will be verified in the particular case that we discuss now.




Figure 6: Left, the curvilinear polygon, bounded by the lines $2p + 8 \pm q = 0$ and two arcs of the cubic $(p/3)^3 + (q/2)^2 = 0$; (right) the distribution of 8000 points in the (p, q) plane for $A = B = \text{diag}(1, 0, -1)$. The (red and black) dashed lines along which singularities are expected are $q = 0$ and $p = -1 \pm q$

4 The particular case $A \sim B \sim J_z$

4.1 Symmetries and reduction to algebraic equations

From now on, we restrict ourselves to the particular case of $\alpha = \beta = \{1, 0, -1\}$. The polynomials P_p and Q_q then reduce to

$$\begin{aligned} P_p(c) = P(R) - p &= -c^2(1+x^2)(1+y^2) + 2cxy \sin \phi \sin \psi - (x^2 y^2 - 2x^2 - 2y^2 + 3) - p \\ Q_q(c) = Q(R) - q &= 2c^2(1-x^2 y^2) + 4cxy \sin \phi \sin \psi - 2(1-x^2)(1-y^2) - q, \end{aligned}$$

where we recall that $c = \cos \theta$, $x = \cos \phi$, $y = \cos \psi$, and we could substitute $\sin \phi = \sqrt{1-x^2}$, $\sin \psi = \sqrt{1-y^2}$ since ϕ and ψ are restricted to $(0, \pi)$. Note that these expressions are invariant under the exchange of ϕ and ψ , or of x and y .

Another symmetry of the problem will be quite useful. Because $-J_z$ is conjugate to J_z by the action of $R_y(\pi/2)$, it is clear that the distribution of the γ 's will be invariant under change of sign, and the distribution of the (p, q) variables will be invariant under change of sign of q . This is apparent in Fig. 6 where we see that both the support of ρ , here a curvilinear quadrangle, and the distribution of points (here a simulation with 8000 points), are symmetric under $q \rightarrow -q$. Our formalism, however, does not have that manifest symmetry, and we will find it useful in the following to choose q *negative*, as we see shortly. The PDF for all values of q will then be reconstructed by symmetry.

The discriminant of Q_q is $\Delta_Q = 16(1-x^2)(1-y^2) + 8q(1-x^2 y^2)$. As explained previously, its positivity follows by consistency from the vanishing of the resultant R . For $-2 \leq q \leq 0$, one may check that, if $\Delta_Q \geq 0$, the real roots obey $|c_{1,2}| \leq 1$ for all $x, y \in [-1, 1]$. **We shall hereafter assume that $-2 \leq q \leq 0$.**

Following the discussion of sect. 3.3, we then determine the resultant of the two polynomials $P_p(c)$ and $Q_q(c)$, a degree 4 polynomial in u and z

$$\begin{aligned} R = \text{Res}(P_p, Q_q; c) &= 4p^2(1-uz)^2 + 4pq(1+u)(1+z)(1-uz) \\ &\quad - 8p(-4 + 2(u+z) + (u+z)^2 - 2uz(u+z) + uz(u-z)^2) \\ &\quad + q^2((1+uz)^2 + (u+z)^2 + 2uz(u+z) + 2(u+z)) \\ &\quad + 4q((4-u^3 + 2u^2 z^2 - z^3 + 2(u+z) - uz(u+z) - 3(u+z)^2 + 3uz(u^2+z^2))) \\ &\quad + 4(u^4 - 8u^3 z - 2u^2 z^2 - 8uz^3 + z^4) + 16(u+z)^3 - 16(u-z)^2 + 64(-u-z+1) \end{aligned} \quad (32)$$

Of course, this polynomial R is also symmetric under the swapping of u and z , since P and Q were under $x \leftrightarrow y$. The factor \tilde{B} appearing in (30) is $\tilde{B} = 4(2+x^2+y^2)$. Then, according to (31), the

PDF reads

$$\rho(p, q) = \frac{2}{\pi^2} \int_0^1 dz \sum_{\substack{\text{roots } u_i \text{ of } R \\ 0 \leq u_i \leq 1}} \frac{(2 + u + z)}{|R'_u|} \Big|_{u=u_i}. \quad (33)$$

4.2 Roots u_i of the resultant R and their singularities

Let us start with some general features of the roots $u_i(z)$ of R :

– The polynomial $R(u, z)$ being symmetric in u, z , its roots $u_i(z)$ or $z_i(u)$ are built by the same function: $z \xrightarrow{S_i} u_i(z)$, $u \xrightarrow{S_i} z_i(u)$. This means that their graph is symmetric with respect to the first diagonal, see Fig. 10 below.

– Within the full domain $q \leq 0$, there are either 0, two or four roots $u_i(z) \in [0, 1]$ for $z \in [0, 1]$. When z varies in $(0, 1)$, this number may change: either some of these roots may evade the interval $[0, 1]$ through one of its end points 0 or 1, but they always do it pairwise; or a pair of real roots “pops out” of the complex plane or disappears into it, but this may occur only when they coalesce. In both cases, the discriminant Δ_R of R with respect to u vanishes. We compute

$$\Delta_R(z, p, q) = 8192(8 + 2p - q)^2(1 - z)^2 z^2 \Delta_R^{(1)}(z, p, q) \quad (34)$$

where $\Delta_R^{(1)}(z, p, q)$ is a fairly cumbersome polynomial of degree 8 in z that we refrain from writing here ². The vanishing of the first factor does not occur for $q < 0$ and $p > -4$. In the following, we denote the ordered roots z_s of Δ_R belonging to $[0, 1]$ as

$$z_{s_0} = 0 \leq z_{s_1} \leq \dots \leq z_{s_r} \leq 1. \quad (35)$$

It turns out there are up to five roots z_s of $\Delta_R^{(1)}(z, p, q)$ in the open interval $]0, 1[$, hence seven in $[0, 1]$. Some of these roots z_s may be *irrelevant*, in the sense that they are associated with the merging of *irrelevant roots* $u_i(z)$ of R , i.e., roots that *do not* satisfy $u_i \in [0, 1]$.

At some particular values of (p, q) , the number of z_s roots may change. Either some z_s evade the interval $[0, 1]$ or enter it, through 0 or 1: computing $\Delta_R^{(1)}(0, p, q)$ and $\Delta_R^{(1)}(1, p, q)$, one finds that this happens along the lines $p \pm q + 1 = 0$ and $q = 0$ which are the dashed lines in Fig. 6. Or two roots of $\Delta_R^{(1)}(z, p, q)$ coalesce, and this occurs for values of (p, q) that are roots of the discriminant $\Delta_R^{(2)}$ of $\Delta_R^{(1)}(z, p, q)$ with respect to z ,

$$\begin{aligned} \Delta_R^{(2)}(p, q) &= q^4(2 - q)^2(1 + p - q)^2(4 + p - q)^2(5 + p - q)^2(8 + 2p - q)^2(4p^3 + 27q^2) \times \\ &(64 + 16p - 48q - 8pq + 13q^2 + pq^2 - q^3)^2(8 + 10p + 2p^2 - 19q - 10pq - p^2q + 8q^2 + 2pq^2 - q^3)^2(T(p, q))^3 \end{aligned} \quad (36)$$

where $T(p, q)$ is a horrendous polynomial of degree 21 in p and 18 in q ². The relevant roots of $\Delta_R^{(2)}$ (in fact of T) for our discussion define the “horned” (red) curve in the upper right part of Fig. 7. The cusp of that curve occurs at $(p_c, q_c) = (-1.37657\dots, -0.234765\dots)$. Note that this p_c is a root of the “third generation discriminant” $\Delta_R^{(3)}(p) := \Delta_T$, in fact of its factor $(2 - 3p - p^2 + 3p^3 + p^4)$. The horned curve intersects the q -axis at $p = p_0 := -1.21891\dots$, a root of $1328 + 1325p + 171p^2 - 17p^3 + p^4$, which is a factor of $T(p, 0)$. Also, it intersects the dotted line $p + q + 1 = 0$ at $p = -0.910988\dots$, $q = -0.089012\dots$; it intersects the dashed line $p - q + 1 = 0$ at $p = -1.14617\dots$, $q = -0.146174\dots$.

A detailed analysis shows that one has to distinguish six regions in the domain of (p, q) , $q \leq 0$, see Fig. 8. These regions differ by the subset of relevant values z_s . Note that we have to carry out the z -integration of (33) in each interval $(z_{s_i}, z_{s_{i+1}})$, after one another, because the integrand is singular at each z_{s_j} , as we discuss in the next subsection. The properties of these six regions are summarized in Table 1; the pattern of z_s when q varies while p is fixed at some value are displayed in Fig. 9; and the various scenarii for the roots $u_i(z)$, which describe several branches of a closed curve in the (z, u) plane, are illustrated in Fig. 10, where colors refer to points of Fig. 7.




Figure 7: The $q \leq 0$ part of the domain in the (p, q) -plane. The special curves are the lines $p - q + 1 = 0$ (dashed) and $p + q + 1 = 0$ (dotted), and an arc of the “horned” curve where $T = 0$ (red). The colored dots refer to the plots of Fig. 10




Figure 8: Zoom on the 6 regions




Figure 9: Plot of the roots z_s of $\Delta_R^{(1)}$ as a function of q : (a) for $p = -3.5$; (b) for $p = -1.5$; (c) for $p = -1.25$, for $q_{\min}(p) \leq q \leq 0$; (d) zoom of the latter on $-0.25 \leq q \leq 0$. The cusps at $q = -0.179503$ and $q = -0.0868984$ lie on the boundary of the horned domain. In each case, the largest z_s is irrelevant.

Region	Number of $z_s \in]0, 1[$	Relevant z_s	Intervals	Roots u_i
I	2	$z_{s_0} = 0, z_{s_1}$	$0 \leq z \leq z_{s_1}$	$u_1 \leq u_2$
II	4	$z_{s_0} = 0, z_{s_1}, z_{s_2}, z_{s_3}$	$0 \leq z < z_{s_1}$ $z_{s_1} < z < z_{s_2}$ $z_{s_2} < z < z_{s_3}$	$u_1 \leq u_2$ $u_1 \leq u_3 \leq u_4 \leq u_2$ $u_4 \leq u_2$
III	5	$z_{s_1}, z_{s_2}, z_{s_3}, z_{s_4}$	$z_{s_1} \leq z < z_{s_2}$ $z_{s_2} < z < z_{s_3}$ $z_{s_3} < z < z_{s_4}$	$u_1 \leq u_2$ $u_1 \leq u_3 \leq u_4 \leq u_2$ $u_4 \leq u_2$
IV	5	$z_{s_2}, z_{s_3}, z_{s_4}, z_{s_5}$	$z_{s_2} \leq z < z_{s_3}$ $z_{s_3} < z < z_{s_4}$ $z_{s_4} < z < z_{s_5}$	$u_1 \leq u_2$ $u_1 \leq u_3 \leq u_4 \leq u_2$ $u_1 \leq u_3$
V	3	z_{s_1}, z_{s_2}	$z_{s_1} \leq z \leq z_{s_2}$	$u_1 \leq u_2$
VI	3	z_{s_2}, z_{s_3}	$z_{s_2} \leq z \leq z_{s_3}$	$u_1 \leq u_2$

Table 1. Pattern of roots z_s of Δ_R and of roots $u_i(z)$ of R in the various regions

For example, in region II, (shaded triangle \cap horned region), there are four roots z_{s_j} , $j = 1, \dots, 4$, of $\Delta_R^{(1)}$ but only the first three are relevant, and four relevant roots for Δ_R , namely $z_{s_0} = 0, z_{s_1}, z_{s_2}, z_{s_3}$:

- * for $0 \leq z \leq z_{s_1}$, there are two roots $0 \leq u_1 \leq u_2 \leq 1$;
- * at $z = z_{s_1}$, a new pair of roots (u_3, u_4) pops out of the complex plane;
- * for $z_{s_1} < z < z_{s_2}$, we have four roots $0 \leq u_1 \leq u_3 \leq u_4 \leq u_2 < 1$;
- * at $z = z_{s_2}$, the pair (u_1, u_3) merges and disappears into the complex plane;
- * for $z_{s_2} \leq z \leq z_{s_3}$, we are left with two roots $u_4 \leq u_2$, which merge at $z = z_{s_3}$ and disappear in the complex plane;
- * for $z > z_{s_3}$, there is no root in the interval $u \in [0, 1]$. The z -integration must be carried out separately on the three intervals $(0, z_{s_1})$, (z_{s_1}, z_{s_2}) , and (z_{s_2}, z_{s_3}) .

4.3 The integrand $\varphi(z)$

Consider the integrand in (33)

$$\varphi(z) := \sum_{\substack{\text{roots } u_i \text{ of } R \\ |u_i| \leq 1}} \frac{(2 + u + z)}{|R'_u|} \Big|_{u=u_i} \quad (37)$$

It has both integrable and non-integrable singularities, the latter where the integral diverges. Typical plots of $\varphi(z)$ in the various regions are displayed in Fig. 11.




Figure 10: A portrait gallery. At various points of the (p, q) domain, 1st line: plot of the roots $u_i(z)$ of R , which describe several branches of a closed curve; 2nd line: the value of (p, q) and the region number; 3d line: the values of the *relevant* z_s .




Figure 11: Typical plots of $\varphi(z)$ in regions I, V or VI (left) or II, III, IV (right). In the latter, the middle and right intervals have been dilated for clarity. The discontinuity of $\varphi(z)$ at $z_{s_{i+1}}, z_{s_{i+2}}$ is due to the contribution of two new roots u_3 and u_4 in that interval.

The singularities of φ as a function of z come either from singularities of $u_i(z)$, or from zeros of the denominator $|R'_u(u_i)|$. Both cases are associated with the merging of roots $u_i(z)$ of R , which occurs at some relevant root z_s of its discriminant Δ_R . The singularity of u_i in the numerator is, however, at worst of square root type and gives rise to no divergence of the integral. We thus concentrate on the possible vanishing of the denominator R'_u .

If we write the polynomial R in a factorized form, $R = c \prod_j (u - u_j(z))$, its derivative at u_i reads $R'_u(u_i) = c \prod_{j \neq i} (u_i(z) - u_j(z))$, and vanishes when u_i coalesces with some u_j , thus at some value $z = z_s$, a root of the discriminant Δ_R .

– Either the pair of roots (u_i, u_j) belonging to $[0, 1]$ emerges from or disappears into the complex plane at z_s with a square root behaviour, $u_{i,j}(z) \sim u_{i,j}(z_s) \pm \frac{1}{2}a|z - z_s|^{1/2} + \dots$, hence $|u_i(z) - u_j(z)| \sim a|z - z_s|^{1/2} + \dots$. Graphically, it manifests itself as a smooth curvature of the “portraits” of $u_i(z)$ and $u_j(z)$ at the points of vertical tangent in Fig. 10. This is what happens at $z = z_{s_0} = 0$ above the dashed line; and generically at the various relevant roots z_s of Δ_R . At such a point, the singularity of $\varphi(z)$ is integrable.

– Or the two roots u_i, u_j cross at a finite angle at z_s : $u_i(z) - u_j(z) \sim b(z - z_s) + o(z - z_s)$ with a finite coefficient b . This is what happens along the lines $p - q + 1 = 0$, or $q = 0$, and translates graphically into an angular point in Fig. 10, see for example cases (a), (d), (g). At such a point, the integral of $\varphi(z)$ diverges “logarithmically” at $z = z_s$, which supposes the introduction of some cut-off that measures the departure of (p, q) from the singular point. This explains the growths of the PDF observed in Fig. 6 along the lines $q = 0$, $p - q + 1 = 0$ and (by symmetry $q \leftrightarrow -q$) $p + q + 1 = 0$, as we discuss in the next subsection.

– Or at exceptional points, the difference $u_i(z) - u_j(z)$ may vanish faster at z_s . This is what happens at the point $(-1, 0)$, where it vanishes as $|z_s - z|^{3/2}$. Graphically, the two curves $u_i(z)$ and $u_j(z)$ form a cusp, see for instance Fig. 10.(f). In that case, the integral of $\varphi(z)$ diverges as an inverse power of the cut-off, see below.

But there is another source of divergence of ρ . The function φ itself may diverge as (p, q) approaches a singular point. This is what happens at the three corners $(-4, 0)$, $(-3, -2)$, $(0, 0)$ of the domain, where we shall see that $R'_u(u_i)$ vanishes for all z in the integration interval.

4.4 The PDF, plots and divergences

We are now in position to draw the plot of the PDF $\rho(p, q)$ resulting from the integration in (33) for $q \leq 0$, supplemented by its mirror image by $q \rightarrow -q$, and to compare it with the histogram obtained by a simulation with 10^6 points, see Fig. 12 and 13.

An important check consists in comparing the probability of occurrence of (p, q) in a finite domain computed by integrating the PDF $\rho(p, q)$ over that domain to that estimated from a big




Figure 12: Left: plot of the function ρ computed on a grid of mesh 10^{-2} ; the regions I to VI, $q < 0$ have been plotted in different colors (deep blue, orange, green, red, violet and brown), while their symmetric partners, for $q > 0$, are all in light blue. Right: histogram of 10^6 points in the (p, q) plane.




Figure 13: The same, in the (γ_1, γ_2) plane.

sample of “random events”. For example, $\mathbb{P}(-3.6 \leq p \leq -3.5, q \leq 0) \Big|_{\text{computed}} = 0.04496$ while the estimate from a sample of size 10^6 gives 0.044886.

As anticipated, the computed ρ exhibits singularities along the lines $q = 0$ and $p \pm q + 1 = 0$. Note that the computation of ρ is carried out point by point on a grid of mesh 10^{-2} in the (p, q) -plane, cutting off the vicinity of the singular lines, while the histogram uses bins of width 0.02 throughout the domain. This explains the slight difference of appearance of the singularities.

By a long (and fairly tedious) case by case analysis, we may assert that the singularities are logarithmic in the approach of *generic* points of the singular lines $q = 0$; $p - q + 1 = 0$ for $q \leq 0$; and $p + q + 1 = 0$ for $q \geq 0$. At the end points and intersection of these lines, i.e., at the corners of the (p, q) -domain, as well as the point $(p, q) = (-1, 0)$, the divergence is stronger, as an inverse power. This is summarized in the following Table, which gathers results obtained in the detailed discussion of the next subsection. The reader will also find in that subsection numerical verifications of the asserted divergences.

In most cases, we proceed as follows: as (p, q) approaches a singular point, some z_s approaches a limiting value z_{s*} while the common value of a pair of coinciding roots, say, $u_s = u_1(z_s) = u_2(z_s)$ approaches $u_{s*} = u_{1,2}(z_{s*})$. Series expansions of z_s and u_s in powers of a “distance” $\epsilon \ll 1$ to the singularity may be computed. On the other hand, the roots $u_1(z)$ and $u_2(z)$ as well as the denominator R'_u of $\varphi(z)$ in (33) may be expanded in powers of $\zeta = \sqrt{|z - z_s|}$. Finally, a double series expansion in powers of ϵ and ζ is obtained for $R'_u(u_{1,2}(z))$, which upon integration in the vicinity of z_s , yields the singular contribution to ρ of z_{s*} . This program is carried out in detail in

the next subsection. Three particular singular points are treated separately.

Position of singularity and (z_{s^*}, u_{s^*})	Approach to singularity	Divergent part of ρ
$p - q + 1 = 0, q \leq 0$ $z_{s^*} = 0, u_{s^*} = (p + 3)/2$ $z_{s^*} = (p + 3)/2, u_{s^*} = 0$	$p - q + 1 \rightarrow 0$	$\rho_{\text{div}} = \frac{1}{\pi^2 \sqrt{2(p+3)} \sqrt{-(1+p)}} \log p - q + 1 $
$q = 0, -4 < p < -1$ $z_{s^*} = 1, u_{s^*} = 1$	$q \rightarrow 0, p$ fixed	$\rho_{\text{div}} = \frac{1}{2\pi^2 p \sqrt{p+4}} \log q $
$q = 0, -1 < p < 0$ $z_{s^*} = p + 1, u_{s^*} = 1$ $z_{s^*} = 1, u_{s^*} = 1$ $z_{s^*} = 1, u_{s^*} = p + 1$	$q \rightarrow 0, p$ fixed	$\rho_{\text{div}} = \frac{1}{2\pi^2 p } ((1 + p)^{-\frac{1}{2}} + (4 + p)^{-\frac{1}{2}}) \log q $
$(p, q) = (-1, 0)$ $z_{s^*} = 0, u_{s^*} = 1$ $z_{s^*} = 1, u_{s^*} = 0$ $z_{s^*} = u_{s^*} = 1$: subdominant	$q \rightarrow 0, p = -1$	$\rho_{\text{div}} = \frac{1}{2\sqrt{6}\pi q ^{\frac{1}{2}}} + O(\log q)$
$(p, q) = (-4, 0)$ $z_{s^*} = 0, u_{s^*} = 0$ $z_{s^*} = u_{s^*} = \frac{2-\kappa}{3\kappa+2}$	$\kappa(p + 4) = -q \rightarrow 0,$ $0 < \kappa < 2$	$\rho_{\text{div}} = C(\kappa)/ q ^{\frac{1}{2}}$
$(p, q) = (0, 0)$ $z_{s^*} = u_{s^*} = 1$	$\kappa p^3 = -27q^2 \rightarrow 0$ $0 < \kappa < 4$	$\rho_{\text{div}} = C''(\kappa)/ q ^{2/3} \sim 1/ p $
$(p, q) = (-3, -2)$ $z_{s^*} = u_{s^*} = 0$	$(p + 3) = \kappa(q + 2) \rightarrow 0$ $-\frac{1}{2} \leq \kappa \leq 1$	$\rho_{\text{div}} = C'(\kappa)/(q + 2)^{\frac{1}{2}}$

Table 2. Position and expression of the singularities of $\rho(p, q)$ in the $q < 0$ part of the domain of Fig. 6. The expressions of the coefficient functions $C(\kappa)$ etc are given in the next subsection.

Finally, note the PDF $\rho(p, q)$ does not vanish along the boundaries of the Horn domain in the (p, q) -plane. In particular, on the lower or upper sides of the domain, i.e., along the arcs of the cubic $q = \mp 2(-p/3)^{3/2}$, $\rho(p, q)$ has a finite limit. This is in no contradiction with the expected vanishing of the PDF $\mathbf{p}(\gamma_1, \gamma_2)$ on the left boundaries $\gamma_1 = \gamma_2$ and $\gamma_2 = \gamma_3$ of the Horn domain in the (γ_1, γ_2) -plane, because of (24) and of the vanishing of the Vandermonde determinant Δ along those curves.

4.5 Analysis of the singularities

In this subsection, we proceed to a detailed –and lengthy– case-by-case analysis of the divergent singularities of ρ .

A preliminary observation is that, due to the $u \leftrightarrow z$ symmetry of our particular case, the $u < z$

and $u > z$ sectors will contribute equally:

$$\begin{aligned} \int \int dudz \delta(R(u, z))(2 + u + z) &= \int \int_{u < z} dudz \delta(R(u, z))(2 + u + z) + \int \int_{u > z} dudz \delta(R(u, z))(2 + u + z) \\ &= 2 \int \int_{u > z} dudz \delta(R(u, z))(2 + u + z). \end{aligned}$$

This will be apparent in the following.

As explained in the previous subsection, we proceed heuristically, making appropriate Taylor expansions close to the singularities. We do it in detail in the first case (singularity along the dashed line), and are then more sketchy.

1. Along the dashed line $p + 1 - q = 0$, the integrand φ has two non-integrable singularities: $\varphi \sim 1/z$ for $z \rightarrow z_{s_0} = 0$, corresponding to $u_{1,2}(z) \rightarrow u_{s_0} = (p + 3)/2$, and, by symmetry between u and z , $\varphi \sim 1/(z_s - z)$ for $z \rightarrow z_s = (p + 3)/2$. Thus the integral diverges and the PDF is infinite along the line.

- For $q = p + 1 + \epsilon$, (i.e., close to and above the line), z_s remains equal to zero, and we determine the common value of the two roots at that point by plugging a series expansion of the form $u_s = \frac{p+3}{2} + \alpha\epsilon + \dots$ in the equation $R(u_s, z_s = 0) = 0$, whence

$$u_{1,2}(0) = u_s = \frac{p+3}{2} + \frac{5+p}{2(9+p)}\epsilon + \dots$$

- For z close to $z_{s_0} = 0$, we write $z = \zeta^2$, approximate $u_{1,2}(z) = u_s + \beta_{1,2}\zeta + \gamma_{1,2}\zeta^2 + O(\zeta^3)$, and determine the coefficients of that expansion by plugging it again in the equation $R(u_{1,2}(z), z = \zeta^2) = 0$. β_1 (resp. β_2) is the negative (positive) root of an equation which, for $q = p + 1 + \epsilon$ reduces to

$$\beta^2 = -\frac{2(3+p)(1+p)(7+p)^2}{(9+p)^3}\epsilon + O(\epsilon^2) \quad (38)$$

and the coefficient γ is given by

$$\gamma_{1,2} = -\frac{p^3 + 5p^2 - 13p + 15 \pm 4\sqrt{2}(p+3)(p+7)\sqrt{-p-1}}{2(p+9)^2} + O(\epsilon). \quad (39)$$

- We then expand the denominator R'_u of (37), for z close to $z_{s_0} = 0$ and ϵ small and of order ζ^2 , as

$$R'_u(u_{1,2}, z = \zeta^2) = 2\sqrt{2}\sqrt{-p-1}(p+3)(p+7) \left(\sqrt{\frac{p+9}{p+3}}\sqrt{\epsilon} \mp 2\zeta \right) \zeta + o(\epsilon, \zeta\sqrt{\epsilon}, \zeta^2) \quad (40)$$

- and we finally derive the divergence of $\int_0^{z_s} dz \varphi$ at the lower end point 0 as

$$\int_0 dz \varphi \Big|_{\text{div}} = \sum_{u_1, u_2} \int_0 \frac{2\zeta d\zeta (p+7)/2}{|R'_u(u_{1,2}, z)|} \Big|_{\text{div}} = \frac{1}{4\sqrt{-2(p+1)}(p+3)} |\log \epsilon|.$$

As explained above, the divergence of the integral at its other end point, obtained by the symmetry $u \leftrightarrow z$, contributes the same amount.

Thus the total divergence of $\rho(p, q)$ as the dashed line is approached from above is

$$\rho(p, q) \Big|_{p-q+1 \rightarrow 0} \approx \rho_{\text{div}} := \frac{1}{\pi^2 \sqrt{2}(p+3)\sqrt{-(1+p)}} |\log |p - q + 1||. \quad (41)$$




Figure 14: $\rho - \rho_{\text{div}}$ above (blue disks) or below (yellow squares) the dashed line for $p = -2.5$, $q = p + 1 \pm \epsilon$, $\epsilon = 3^{-j-3}$, plotted against $j = 1, \dots, 20$.

This formula is well verified on numerical data, see Fig. 14. Note that the logarithmic behavior is enhanced in the approach to $p = -1$, $q = 0$ and to $p = -3$, $q = -2$, at the southern end of the plot. This will be reconsidered in detail in items 4 and 7 below.

One also checks that the same formula applies to the approach to the dashed line from below ($\epsilon < 0$). The relevant expressions are

$$z_{s_1} = \frac{(p+9)|\epsilon|}{4(p+3)}; \quad u_{s_1} = \frac{p+3}{2} - \frac{(p+5)|\epsilon|}{2(p+9)}; \quad u_{1,2}(z = z_{s_1} + \zeta^2) = u_{s_1} \mp \beta\zeta + \gamma_{1,2}\zeta^2 + \dots$$

with $\beta > 0$ and $\gamma_{1,2}$ as given in (38) and (39), and the same expression for R'_u as in (40):

$$R'_u(u_{1,2}, z_{s_1} + \zeta^2) \approx \pm 2\zeta \sqrt{-2(1+p)(p+7)(p+3)} (-\sqrt{(p+9)/(3+p)} \sqrt{|\epsilon|} + 2\zeta)$$

and hence the same divergence as above the line, see Fig. 14 for an illustration at $p = -2.5$.

- By a similar discussion, one finds that when $q \rightarrow 0$ with $-4 \leq p \leq p_0 = -1.21891$, (i.e., in region I), the singularity at $z_{s_0} = 0$ is integrable while that at $z_{s_1} \rightarrow 1$ gives rise to a divergence of the integral. We write for short $z_s = z_{s_1}$. For $|q|$ small,

$$z_s = 1 + 2q/(4+p) + O(q^2), \quad u_s = u_1(z_s) = u_2(z_s) = 1 - 8q/(p(4+p)) + O(q^2);$$

as $z \rightarrow z_s$, we write $z = z_s - \zeta^2$, and the two roots $u_{1,2} \approx u_s \mp \beta_1\zeta - \gamma_{1,2}\zeta^2$, for some computable coefficients $\beta_1, \gamma_{1,2}$, so that $R'_u(u_{1,2}, z_s - \zeta^2) \approx \mp 32|p|(\sqrt{4+p}\zeta + \sqrt{-2q})\zeta$, whence a divergence of the integral $\int^{z_s} dz \varphi(z) \sim \frac{1}{4|p|\sqrt{4+p}} |\log |q||$.

The same applies in region II, $p_0 = -1.21891 \leq p < 1$, where now four values of z_s exist, (see sect. 4.2 and Table 1), but the singularities of R'_u at the points $z_{s_0} = 0$, z_{s_1} and z_{s_2} are of inverse square root type, hence integrable, and only the linear vanishing of R'_u at z_{s_3} matters. Hence

$$\rho(p, q) \Big|_{-4 < p < -1}^{q \rightarrow 0} \approx \rho_{\text{div}} := \frac{1}{2\pi^2 |p| \sqrt{p+4}} |\log |q||. \quad (42)$$

See Fig. 15 for comparison with numerical data.

As $p \rightarrow -1$, however, the singularities at $z_{s_0} = 0$ and $z_{s_2} \rightarrow 1$ become sharper and sharper, resulting in a stronger divergence at $z = 1$, see below item 4.

- In region IV, ($-1 < p < 0$), q is small ($|q| < 1/10$), and the relevant values of (z_s, u_s) are




Figure 15: $\rho - \rho_{\text{div}}$ for $p = -3.5, \dots, -1.1$, $q = -3^{-j-1}$, plotted against $j = 1, \dots, 30$.

well approximated, up to higher powers of q , by

$$\begin{aligned}
z_{s_2} &\approx p + 1 + \alpha_{z_2} q \\
\text{with } \alpha_{z_2} &\text{ the largest root of } -4 + p + p^2 + 2\alpha p(p+1) - 4p\alpha^2, \\
\text{i.e., } \alpha_{z_2} &= (p + p^2 - \sqrt{-16p + 5p^2 + 6p^3 + p^4}) / (4p) \\
u_{s_2} &\approx 1 + \alpha_{u_2} q, \quad \alpha_{u_2} = \frac{1}{4} \left(1 + (8 + 5p + p^2) / \sqrt{-16p + 5p^2 + 6p^3 + p^4} \right); \\
z_{s_4} &\approx 1 + \frac{2q}{p+4}, \quad u_{s_4} \approx 1 - \frac{8}{p(4+p)} q; \\
z_{s_5} &\approx 1 + \frac{1}{2} q, \quad u_{s_5} \approx p + 1 + \frac{1}{2} q \left(p + 3 + \frac{4}{p} \right).
\end{aligned}$$

(The value of z_{s_3} where u_3 and u_4 merge, see Fig. 10 (g), is of no concern to us here, as φ is integrable there.)

As $q \rightarrow 0$ three divergences occur:

– the pair of roots (u_1, u_2) merge toward u_{s_2} as $z \rightarrow z_{s_2}$; setting $\zeta^2 = z - z_{s_2}$, as $\zeta \rightarrow 0$, $u_1 = u_{s_2} \mp \beta \zeta + \gamma_1 \zeta^2 + \dots$ with β^2 and $\gamma_{1,2}$ determined by a series expansion of the equation $R(u, z) = 0$. One finds that $\beta = O(|q|^{\frac{1}{2}})$, while $\gamma_{1,2} = O(1)$. Consequently

$$R'_u(u_1, z_{s_2} + \zeta^2) \approx \mp 8(p+4)(p+1)^{\frac{1}{2}} (\sqrt{-2p|q|^{\frac{1}{2}}(p^4 + 6p^3 + 5p^2 - 16p)^{\frac{1}{4}} + 2p\zeta}) \zeta$$

whence a divergence of the integral at its lower bound $\int_{z_{s_2}} \varphi(z) dz \Big|_{\text{div}} = \frac{1}{8|p|(p+1)^{\frac{1}{2}}} |\log |q||$;

– the pair of roots (u_1, u_3) converges to u_{s_5} as $z \rightarrow z_{s_5}$, and $R'_u(u_{1,3}, z) \approx 16|p|((p+4)\sqrt{1+p}(1-z) + |q|)$: this is just the image by the $u \leftrightarrow z$ symmetry of the latter, whence another contribution $|\log |q|| / (8|p|\sqrt{p+1})$ to the integral of φ ;

– the pair of roots (u_2, u_4) converges to u_{s_4} as $z \rightarrow z_{s_4}$. By similar expansions near z_{s_4} one finds that $R'_u(u_2, z_{s_4} - \zeta^2) = 16|p|(\mp 2\sqrt{-2q} + (-(p+8) \mp 2(p+4)^{\frac{1}{2}} \zeta)) \zeta$, whence a divergence of the integral $\int^{z_{s_4}} dz \varphi(z) \Big|_{\text{div}} = |\log |q|| / (4|p|\sqrt{p+4})$.

Thus as above in regions I and II, for $q \approx 0$, $\rho \sim C \log |q|$, but with a larger value of the coefficient, $C = \frac{1}{2\pi^2|p|} ((1+p)^{-\frac{1}{2}} + (4+p)^{-\frac{1}{2}})$. The agreement between numerical results and that coefficient C is illustrated in Fig. 16 for $p = -0.8$. It deteriorates at small values of $|q|$ where the convergence of the integral is bad.

4. At $p = -1$, $q = 0$, one can see that, as $z \rightarrow 0$, u_1 and u_2 approach 1 with the same slope, so that we have now $|u_1 - u_2| \sim \alpha z^{3/2}$, causing a strong divergence of ρ . More precisely, for




Figure 16: $\rho - \rho_{\text{div}}$ for $p = -0.8$, $q = -3^{-j-3}$, plotted against $j = 1, \dots, 18$.

$p = -1$ and q small, $z_{s_1} = -q + O(q^2)$ and if $z - z_{s_1} = \zeta^2 \rightarrow 0$, one finds

$$R'_u| \approx 36\sqrt{2}(\zeta^2 + \frac{4}{3}|q|)\zeta$$

whence a contribution to the divergent part of $\int_{z_{s_1}} dz\varphi(z)$ equal to $\frac{\pi}{4\sqrt{6}|q|^{\frac{1}{2}}}$. The coalescence of roots u_1 and u_3 towards $u_{s_4} \approx 0$ at $z_{s_4} \approx 1$ gives rise to the same divergence (by the $u - z$ symmetry once again), while there is a weaker (logarithmic) divergence coming from u_2 and u_4 merging to $u_{s_3} \approx 1$ as $z \rightarrow z_{s_3} \approx 1$ (see Fig. 10 f). In total, we have

$$|\rho(p, q)| \Big|_{\substack{p=-1 \\ q \rightarrow 0_-}} \approx \rho_{\text{div}} := \frac{1}{\sqrt{6}\pi|q|^{\frac{1}{2}}} \quad (43)$$

up to a subdominant $\log|q|$ term. See Fig. 17-left for a numerical plot of ρ/ρ_{div} converging to 1.

5. Divergence at $(p, q) = (-4, 0)$. Let us approach that singular point in a linear way, letting $p + 4 = \epsilon$, $q = -\kappa\epsilon$, $\epsilon \rightarrow 0$, with $0 < \kappa < 2$ (so as to remain within the Horn domain); then to leading order in ϵ , the z -integration runs between $z_{s_0} = 0$ and $z_{s_1} = (2 - \kappa)/(2 + 3\kappa)$. Solving $R = 0$ in that limit, one finds that the portrait $(u_1(z), u_2(z))$ forms a vanishingly thin ellipse-like curve along the diagonal: $u_1(z) = z \mp \sqrt{2 + 3\kappa} \left(\frac{(1-z)z(z_{s_1}-z)}{1+z} \right)^{\frac{1}{2}} \epsilon^{\frac{1}{2}}$ to the first non trivial order in ϵ , and plugging this expression into R'_u yields

$$|R'_u(u_{1,2})| = 32\sqrt{2 + 3\kappa} (1+z)^{\frac{3}{2}} \left((1-z)z(z_{s_1}-z) \right)^{\frac{1}{2}} \epsilon^{\frac{1}{2}} + O(\epsilon)$$




Figure 17: Left: ρ/ρ_{div} for $p = -1$, $q = -3^{-j} \rightarrow 0$, $j = 3, \dots, 10$, plotted against q .

Right: Plot of ρ/ρ_{div} as a function of p , in the approach of $(-4, 0)$ along the line $p + q/\kappa = -4$ for three values of κ .




Figure 18: (Left) Plot of $\rho(p, q)/\rho_{\text{div}}$ in the approach of $(0, 0)$ along the cubic $\kappa p^3 + 27q^2 = 0$ for $\kappa = 1, 2, 3$ (in region VI). Convergence of the integral deteriorates for small values of p . (Right) Plot of $\rho(p, q)/\rho_{\text{div}}$ in the approach of $(-3, -2)$ along the line $p - q\kappa = -3 + 2\kappa$ for five values of κ .

whence a divergence of ρ as $|q|^{-\frac{1}{2}}$

$$\rho_{\text{div}} := \frac{1}{4\pi^2} \sqrt{\frac{\kappa}{2 + 3\kappa}} \frac{1}{|q|^{\frac{1}{2}}} \int_0^{z_{s_1}} \frac{dz}{\sqrt{z(1-z)(1+z)(z_{s_1} - z)}}. \quad (44)$$

in very good agreement with numerical data, see Fig. 17-right.

6. Divergence at $p = q = 0$. We let (p, q) approach $(0, 0)$ in region VI, for example along the cubic $\kappa p^3 + 27q^2 = 0$, with $\kappa < 4$; then the two end points z_{s_2} and z_{s_3} of the integral go to 1 as $p = -\epsilon^2$ goes to 0, i.e., $z_{s_i} \approx 1 - \alpha_{z,i}\epsilon^2 + O(\epsilon^3)$ where $\alpha_{z,i}$, $i = 2, 3$, are the second and third largest roots of $27\alpha_z(1 - \alpha_z)^2 - \kappa$. Both z and u are thus confined in an interval of size ϵ^2 near 1, and solving the equation $R = 0$ in the rescale variable $z = 1 - \zeta\epsilon^2$ and plugging into R'_u , one finds that the latter has a limiting shape described by the elliptic curve

$$R'_u(u_{2,3}) = \pm \frac{2^6}{3\sqrt{3}} \epsilon^4 \sqrt{\zeta(27\zeta(1 - \zeta)^2 - \kappa)}$$

so that

$$\rho_{\text{div}} := \frac{\sqrt{3} \kappa^{\frac{1}{3}}}{4\pi^2 |q|^{\frac{2}{3}}} \int_{\alpha_{z,3}}^{\alpha_{z,2}} \frac{d\zeta}{\sqrt{\zeta(27\zeta(\zeta - 1)^2 - \kappa)}}. \quad (45)$$

This behavior is again well supported by numerical calculations, see Fig. 18-left.

7. Divergence at $p = -3, q = -2$. If one approaches that corner of the domain along lines $p = -3 + \kappa\epsilon, q = -2 + \epsilon$ with $-\frac{1}{2} < \kappa < 1$ so as to remain in region I, one finds that $z_{s_0} = 0$ and $z_{s_1} = \frac{1}{6}(1 + 2\kappa)\epsilon$ to the lowest order in ϵ , so that both z and u remain small of order ϵ . Solving the equation $R = 0$ to order ϵ^3 , one finds that $u_1(z) = z_{s_1} - z \pm \frac{2}{9} \left(12\epsilon(1 - \kappa)(z(z_{s_1} - z)) \right)^{\frac{1}{2}}$ so that $|R'_u(u_{1,2})| \approx 32\sqrt{3(1 - \kappa)}(z(z_{s_1} - z))^{\frac{1}{2}}\epsilon^{\frac{1}{2}}$, and the z -integration may be carried out, leading to

$$\rho_{\text{div}} := \frac{1}{\pi\sqrt{48(1 - \kappa)}} \frac{1}{|q + 2|^{\frac{1}{2}}}. \quad (46)$$

This is corroborated by the numerical calculation at various values of κ , see Fig. 18-right.

The alert reader may wonder why the singularity along the line $p + q + 1 = 0$ of the *upper* half-plane, (a reflection of the singularity along the dashed line of the $q < 0$ half-plane) does not manifest itself along the *dotted* line of the lower half-plane. The reason is that, in that lower half-plane, the two z_s that merge there are in fact irrelevant for $q < 0$.

5 Zonal polynomials

In the Hermitian case, it is known that the Horn problem discussed so far has a discrete counterpart, involving Littlewood–Richardson multiplicities, and may be regarded as a semi-classical limit of the latter. The PDF $p(\gamma|\alpha, \beta)$, or rather the “volume function” \mathcal{J} equal to the latter up to a Vandermonde factor, measures the distribution of (rescaled) Littlewood–Richardson multiplicities, i.e., structure constants of Schur polynomials, in the large scale limit, see [5, 16]. In the real symmetric case, one expects similarly the PDF, or rather some “volume like function” \mathcal{J} proportional to it to measure, at least in the generic case, the distribution of (rescaled) “zonal multiplicities” i.e., appropriate structure constants of zonal polynomials, see [9]. This motivates the discussion of the present section, where we pay special attention to the normalization and specializations of the Jack and zonal polynomials. We shall in particular use $SU(n)$ reduction⁵ of zonal (or Jack) polynomials and introduce a notion of $SU(n)$ zonal characters very similar to the usual Weyl characters.

5.1 About Jack and zonal polynomials

5.1.1 Jack polynomials and their normalizations

Zonal polynomials can be defined in many ways and we refer the reader to the abundant literature (see for instance [17], [18]). One possible approach is to start from Jack polynomials (themselves a particular case of the larger family called Macdonald polynomials). Sentence re-written A Jack polynomial with n variables is labelled by an integer partition κ and a real parameter α . When one specializes the value of α , Jack polynomials, in turn, give rise to various interesting families, in particular the Schur polynomials (case $\alpha = 1$), the zonal polynomials (case $\alpha = 2$), and the quaternionic polynomials (case $\alpha = 1/2$).

Actually there are three variants of the Jack polynomials, denoted J_X^α with $X = P, C, J$ ⁶ differing by an overall normalization (an overall α -dependent and partition-dependent numerical factor): for example, one writes $J_J = c_{PJ} J_P$, etc. When $\alpha = 2$ one has therefore also three kinds of zonal polynomials respectively denoted Z_P, Z_J and Z_C . When studying zonal or Jack polynomials, many authors – in particular in old papers – use the James normalization (polynomials Z_J) without saying so explicitly. As we shall see, the most interesting family, for us, is the family of the zonal polynomials (and also the Jack polynomials) defined with the P normalization, the reason, that will be discussed below, is that this normalization is compatible with $SU(n)$ -reduction and with the conjugation of irreducible representations (irreps) of $SU(n)$ —the latter being described by integer partitions with at most $n - 1$ parts. When expanded in terms of monomial symmetric polynomials, the J normalization of the Jack polynomial defined by the partition κ makes the coefficient of the lowest order monomial $[1^n]$ equal to $n!$, whereas, using the same expansion, the P normalization makes the coefficient of the monomial relative to the highest partition (i.e., κ) equal to 1. For a given partition κ , the normalization factor c_{PJ} is the lower α -hook coefficient of κ .

Note: Zonal polynomials, with an un-specified normalization, were denoted $Z(\kappa)(x)$ in (5).

5.1.2 Packages

Zonal polynomials and, more generally, Jack polynomials (variables are called x_j), are usually written in terms of monomial symmetric functions, or in terms of power sums, not very often in terms of the variables x_j themselves because this would take too much space. To the authors’ knowledge there are very few computer algebra packages devoted to the manipulation of those polynomials; we should certainly mention [23], written for Mapple (that we did not use), and the

⁵i.e., eliminate integer partitions κ when $\iota(\kappa) > n$, ι being the length of κ , and, when $\iota(\kappa) = n$, replace κ by $\kappa \bmod \{1, 1 \dots, 1\}$, the latter being therefore an *extended partition* (see footnote 7) with a last part equal to 0.

⁶the capital subscripts J and C , for both Jack and zonal polynomials, refer to the original papers of James [20, 21], and Constantine [22], see also [19].

small package [24] written for Mathematica (but it is slow, unstable (division by 0), and uses an obsolete version of the language). For those reasons we developed our own, using Mathematica: the definition chosen for Jack polynomials uses a recurrence algorithm in terms of skew Young diagrams and a modified Pieri's formula described by Macdonald in [18], see also [25]; our code, which also contains commands to convert Jack, Schur and zonal polynomials to several other basis (elementary symmetric polynomials, power sums, monomial sums, complete sums), and commands for calculating structure constants in each basis, is freely available on the web site [14]; the same package contains commands giving the coefficients $Z(\kappa)(I)$ and $c(\kappa)$ that appear in formula (5), with various normalization choices.

5.1.3 Structure constants

Zonal polynomials form a basis of the space of the ring of symmetric polynomials in n variables. Structure constants in this basis are only *rational*s (by way of contrast, the coefficients of Schur polynomials in the expansion of a product of two Schur polynomials are non-negative integers). For illustration, let us consider the zonal polynomial(s) for the extended⁷ partition $\{2, 1, 0\}$ (i.e., three variables x_1, x_2, x_3), and the decomposition of its square, using the three standard normalizations; we also give the decomposition obtained for the *square* of the Schur polynomial $s(\{2, 1, 0\})$.

$$\begin{aligned} Z_P(\{2, 1, 0\})^2 &= \frac{25}{12}Z_P(\{2, 2, 2\}) + \frac{12}{5}Z_P(\{3, 2, 1\}) + \frac{4}{3}Z_P(\{3, 3, 0\}) + \frac{4}{3}Z_P(\{4, 1, 1\}) + Z_P(\{4, 2, 0\}) \\ Z_J(\{2, 1, 0\})^2 &= \frac{5}{54}Z_J(\{2, 2, 2\}) + \frac{12}{35}Z_J(\{3, 2, 1\}) + \frac{4}{135}Z_J(\{3, 3, 0\}) + \frac{32}{405}Z_J(\{4, 1, 1\}) + \frac{1}{27}Z_J(\{4, 2, 0\}) \\ Z_C(\{2, 1, 0\})^2 &= \frac{3}{4}Z_C(\{2, 2, 2\}) + \frac{12}{25}Z_C(\{3, 2, 1\}) + \frac{21}{25}Z_C(\{3, 3, 0\}) + \frac{12}{25}Z_C(\{4, 1, 1\}) + \frac{63}{125}Z_C(\{4, 2, 0\}) \\ s(\{2, 1, 0\})^2 &= s(\{2, 2, 2\}) + 2s(\{3, 2, 1\}) + s(\{3, 3, 0\}) + s(\{4, 1, 1\}) + s(\{4, 2, 0\}). \end{aligned}$$

Specifying the number of variables matters. In the previous example for instance, the square of $Z_P(\{2, 1, 0, 0\})$, i.e., four variables, reads: $\frac{125}{63}Z_P(\{2, 2, 1, 1\}) + \frac{25}{12}Z_P(\{2, 2, 2, 0\}) + \frac{25}{18}Z_P(\{3, 1, 1, 1\}) + \frac{12}{5}Z_P(\{3, 2, 1, 0\}) + \frac{4}{3}Z_P(\{3, 3, 0, 0\}) + \frac{4}{3}Z_P(\{4, 1, 1, 0\}) + Z_P(\{4, 2, 0, 0\})$. Of course $Z_P(\{\kappa_1, \kappa_2, \kappa_3, 0\})$ restricts to $Z_P(\{\kappa_1, \kappa_2, \kappa_3\})$ if $x_4 = 0$, but the last decomposition also contains new terms like $Z_P(\{2, 2, 1, 1\})$ that vanish when $x_4 = 0$.

5.1.4 A particular feature of structure constants in the Z_P basis

We pause here to notice that the coefficients of $Z_P(\{3, 3, 0\})$ and of $Z_P(\{4, 1, 1\})$ are the same (both equal to $4/3$), with the same remark for the coefficients of $s(\{3, 3, 0\})$ and of $s(\{4, 1, 1\})$, which are both equal to 1. This is not so for Z_J and Z_C .

In the Schur case this remark is not surprising: indeed, in terms of irreps of $SU(3)$, the Schur decomposition of $s(\{2, 1, 0\})^2$ corresponds to the tensor decomposition of the square of the adjoint representation labelled⁸ by its highest weight $[1, 1]$; in other words one recovers the well-known decomposition $[1, 1]^{\otimes 2} = [0, 0] \oplus 2[1, 1] \oplus [0, 3] \oplus [3, 0] \oplus [2, 2]$. The two integer partitions $\{3, 3\}$ and $\{4, 1, 1\}$ determine, *once they are reduced to $SU(3)$* , two conjugate Young diagrams of respective shapes $\{3, 3\}$ and $\{3\}$ describing the two complex conjugate irreducible representations $[0, 3]$ and $[3, 0]$ of $SU(3)$. The irrep $[1, 1]$, associated with the partition $\{2, 1, 0\}$, is self-conjugate, actually real, and the coefficients of those complex conjugate irreps that appear in its square, in particular $[0, 3]$ and $[3, 0]$, are, of course, equal.

⁷Given an integer partition κ , say of length s , of the integer m , it is convenient, in order to specify the number of variables in symmetric polynomials, to call "extended partition" of length n , assuming that $n \geq s$, the partition that is obtained from κ by padding $n - s$ zeros to the right of κ . The length of the obtained partition (no longer an integer partition in the strict sense) is then equal to n , the chosen number of variables. Example: $Z_P(\{2, 1, 0, 0\}) = Z_P(\{2, 1\})(x_1, x_2, x_3, x_4)$.

⁸The components of a highest weight $\lambda = [\lambda_1, \lambda_2]$ are written in the Dynkin basis (the basis of fundamental weights).

More generally we observe (the proof of this conjecture is left to the reader) that the following property holds in the zonal P case⁹: If λ is a self-conjugate irreducible representation of $SU(n)$ described by a Young diagram of shape χ (an integer partition, or an extended partition of length n), and if χ' and χ'' are two partitions appearing in the decomposition of the square of the zonal polynomial $Z_P(\chi)$ in the Z_P basis that give rise, after $SU(n)$ reduction, to complex conjugate representations, the coefficients (structure constants) of $Z_P(\chi')$ and of $Z_P(\chi'')$ are equal.

Actually, the same property seems to hold for all values of the Jack parameter α , when using the P normalization. It does not hold for the normalizations J and C of zonal polynomials.

Remember that the notion of complex conjugation on $SU(n)$ irreps can be described in purely combinatorial terms: if κ is the integer partition describing some irrep of $SU(n)$, its length (number of parts) obeys $\iota(\kappa) < n$; then one obtains the partition describing the complex conjugate representation by taking the complement of (the Young diagram of) κ in a rectangle which is $\kappa(1)$ units wide ($\kappa(1)$ being the largest part of κ) and $(\iota(\kappa) + 1)$ units deep.

5.2 From zonal polynomials to $SU(n)$ zonal characters

Although we have in mind applications to the zonal case $\alpha = 2$, or to the quaternionic (zonal) case $\alpha = 1/2$, most of our considerations, in the section that follows, apply to arbitrary values α of the Jack parameter.

5.2.1 $SU(n)$ -zonal characters

An irrep of $SU(n)$, is characterized by its highest weight (hw) λ . Its components in the basis of fundamental weights (Dynkin labels) are denoted $[\lambda_1, \dots, \lambda_{n-1}]$. When considering irreps of $U(n)$ one adds a last index λ_n ; here we only consider the case $SU(n)$, but it is often handy to keep this last index, while setting $\lambda_n = 0$. One can also characterize the same irrep λ by the Young diagram defined by the extended partition $\alpha = \ell(\lambda)$ with components $\ell_i(\lambda) = \sum_{j=i}^n \lambda_j$, $i = 1, \dots, n$, obeying the constraint $\alpha_i \geq \alpha_{i+1}$ for all i . Conversely, given a partition δ of length n , extended or not, one obtains a highest weight λ for $SU(n)$ by setting $\lambda_i = \delta_i - \delta_{i+1}$; such a partition δ differs from $\alpha = \ell(\lambda)$ by a constant shift. We denote by $N_{\lambda\mu}^\nu$ the multiplicity¹⁰ of the irrep ν in the tensor product of the irreps of $SU(n)$ defined by λ and μ .

Given a dominant weight λ of $SU(n)$, i.e., a non-negative integer combination of the fundamental weights, call $\ell(\lambda)$ its associated partition of length n (i.e., $\ell(\lambda)_n = 0$) and take the Jack-P polynomial $J_P^\alpha(\ell(\lambda))(x_1, \dots, x_n)$ determined by the partition $\ell(\lambda)$. We then consider the following Laurent polynomial in the variables y_1, \dots, y_{n-1} :

$$J_P^\alpha(\ell(\lambda))(x_1 = y_1, x_2 = \frac{y_2}{y_1}, \dots, x_j = \frac{y_j}{y_{j-1}}, \dots, x_{n-1} = \frac{y_{n-1}}{y_{n-2}}, x_n = \frac{1}{y_{n-1}})$$

If $\alpha = 1$, J_P^α is a Schur polynomial and the previous Laurent polynomial is recognized as the Weyl character $\chi(\lambda)$ of the irrep λ , for the Lie group $SU(n)$.

If $\alpha = 2$, i.e., when J_P^α is a zonal polynomial Z_P (with the normalization P), we introduce, by analogy, and for lack of a better name, the following notation and definition¹¹:

⁹We remind the reader that Schur polynomials can be obtained from Jack polynomials, with the P normalization, just by setting $\alpha = 1$ (no pre-factors).

¹⁰ $N_{\lambda\mu}^\nu$ is sometimes called the Littlewood-Richardson (LR) multiplicity, although, strictly speaking, the latter refers to the coefficient of $\ell(\nu)$ in the decomposition in the Schur basis of the product of two Schur polynomials respectively defined by the partitions $\ell(\lambda)$ and $\ell(\mu)$. This decomposition often contains terms labelled by integer partitions of length larger than n , therefore not contributing to the tensor product of $SU(n)$ representations—a Young diagram of $SU(n)$ cannot have more than $n - 1$ lines.

¹¹Another notion of “zonal character” can be found in the literature, [26], but it is related to the symmetric group, not to irreducible representations of $SU(n)$. It differs from the notion that we consider here.

Definition 1. The zonal character $\chi_Z(\lambda)$ of $SU(n)$ associated with the dominant weight λ is defined as the Laurent polynomial

$$\chi_Z(\lambda)(y_1, \dots, y_{n-1}) = Z_P(\ell(\lambda))(y_1, \frac{y_2}{y_1}, \dots, \frac{y_j}{y_{j-1}}, \dots, \frac{y_{n-1}}{y_{n-2}}, \frac{1}{y_{n-1}})$$

Now, moving to the Lie algebra $\mathfrak{su}(n)$ – or, equivalently, to trigonometric characters, we start from the same Z_P polynomial expressed in terms of x_j variables but this time perform the following transformations on its arguments: $x_i \rightarrow e^{i(a_i - \frac{1}{n} \sum_j a_j)}$, then $a_j \rightarrow a_1 - \sum_{i=1}^{j-1} u_i$. The result is a trigonometrical expression in the variables u_j , that we call the *Lie algebra zonal character* of $\mathfrak{su}(n)$ associated with the hw λ , or the *trigonometric zonal character* of $SU(n)$ associated with the hw λ .

Let us give one example. Take $n = 3$ and $\lambda = [1, 1]$. The associated (extended) partition is $\ell(\lambda) = \{2, 1, 0\}$. The Jack polynomial $J_P^\alpha(\{2, 1, 0\})$ in terms of the variables x_1, x_2, x_3 , the associated Laurent polynomial, and its trigonometric version are given below. The Schur polynomial $s(\{2, 1, 0\})$ and the Zonal-P polynomial $Z_P(\{2, 1, 0\})$ are obtained from the first expression by setting respectively $\alpha = 1$ and $\alpha = 2$. The corresponding $SU(n)$ and $\mathfrak{su}(n)$ zonal characters are obtained from the last two expressions by setting $\alpha = 2$

$$\begin{aligned} J_P^\alpha(\{2, 1, 0\}) &= x_1^2 x_2 + x_1 x_2^2 + x_2^2 x_3 + x_2 x_3^2 + x_1^2 x_3 + x_1 x_3^2 + \frac{6x_1 x_2 x_3}{2 + \alpha} \\ \chi_\alpha^{\text{SU}(n)}([1, 1]) &= \frac{6}{\alpha + 2} + \frac{y_1^2}{y_2} + \frac{y_2^2}{y_1} + \frac{y_1}{y_2^2} + \frac{y_2}{y_1^2} + y_1 y_2 + \frac{1}{y_1 y_2} \\ \chi_\alpha^{\mathfrak{su}(n)}([1, 1]) &= 2 \left(\frac{3}{\alpha + 2} + \cos u_1 + \cos u_2 + \cos(u_1 + u_2) \right) \end{aligned}$$

Taking $\alpha = 1$ in the second expression, one recognizes the Weyl character of the adjoint representation of $SU(3)$, the powers (positive or negative) of the y_j being, as usual, the components of the weights of the weight system of this representation in the basis of fundamental weights. What happens in the zonal case, and more generally when $\alpha \neq 1$ is that the “multiplicities” of the weights are no longer integers. For this particular irrep, only the multiplicity of the weight at the origin of the weight system is modified by α , its value being 2 in the usual (Schur) case but $3/2$ in the zonal case. Notice that the trigonometric expression is real —it is so because the hw $[1, 1]$ is self-conjugate, otherwise the obtained expression would be complex. The arguments being specified (partitions or Dynkin labels), we shall denote $\chi_\alpha^{\text{SU}(n)}$, in the cases $\alpha = 2$ and $\alpha = 1/2$, by χ_Z and χ_Q .

5.2.2 Structure constants for $SU(n)$ -zonal characters

Given two $SU(n)$ irreps, there are many ways to obtain the decomposition of their tensor products into a sum of irreps. The honeycomb technique, for instance, is very fast¹² but it is not available in the zonal case that we consider. However, we can replace the multiplication of the associated $SU(n)$ Weyl characters by the multiplication of the associated $SU(n)$ zonal characters as defined above, which amounts to use the structure constants for the appropriate product of zonal polynomials. Let us illustrate this with our favorite example, the square of $[1, 1]$. From the already given decomposition of the square of $Z_P(\{2, 1, 0\})$ we obtain immediately:

$$\chi_Z([1, 1])^2 = \frac{25}{12} \chi_Z([0, 0]) + \frac{12}{5} \chi_Z([1, 1]) + \frac{4}{3} \chi_Z([0, 3]) + \frac{4}{3} \chi_Z([3, 0]) + \chi_Z([2, 2])$$

¹²and the semi-magic square algorithm, valid for $SU(3)$, is even faster, see [27].

The same decomposition can be obtained by using Laurent polynomials since the associated SU(3) zonal characters are as follows (the reader can then check that the previous equality holds):

$$\begin{aligned}
\{2, 1, 0\} &\rightarrow \chi_Z([1, 1]) = \frac{y_1^2}{y_2} + \left(y_2 + \frac{1}{y_2}\right) y_1 + \frac{y_2^2 + \frac{1}{y_2}}{y_1} + \frac{y_2}{y_1^2} + \frac{3}{2} = \frac{2y_2 y_1^4 + 2y_2^3 y_1^3 + 2y_1^3 + 3y_2^2 y_1^2 + 2y_2^4 y_1 + 2y_2 y_1 + 2y_2^3}{2y_1^2 y_2^2} \\
\{2, 2, 2\} &\rightarrow \chi_Z([0, 0]) = 1 \\
\{3, 2, 1\} &\rightarrow \chi_Z([1, 1]) = \text{already given} \\
\{3, 3, 0\} &\rightarrow \chi_Z([0, 3]) = \frac{1}{5} \left(\frac{5y_1^3}{y_2^2} + \frac{3y_1^2}{y_2} + 3y_2 y_1 + \frac{3y_1}{y_2^2} + 5y_2^3 + \frac{3y_2^2}{y_1} + \frac{3}{y_2 y_1} + \frac{3y_2}{y_1^2} + \frac{5}{y_1^3} + 2 \right) \\
\{4, 1, 1\} &\rightarrow \chi_Z([3, 0]) = \frac{1}{5} \left(\frac{5y_1^3}{y_2} + \frac{3y_1^2}{y_2} + 3y_2 y_1 + \frac{3y_1}{y_2^2} + \frac{5}{y_2^2} + \frac{3y_2^2}{y_1} + \frac{3}{y_2 y_1} + \frac{3y_2}{y_1^2} + \frac{5y_2^3}{y_1^3} + 2 \right) \\
\{4, 2, 0\} &\rightarrow \chi_Z([2, 2]) = \frac{1}{6} \left(\frac{6y_1^4}{y_2^2} + \frac{4y_1^3}{y_2^2} + 4y_1^3 + 6y_2^2 y_1^2 + \frac{6y_1^2}{y_2} + \frac{6y_1^2}{y_2^4} + 6y_2 y_1 + \frac{6y_1}{y_2^2} + 4y_2^3 + \frac{4}{y_2^2} + \frac{6y_2^2}{y_1} + \frac{6}{y_2 y_1} + \frac{6y_2^4}{y_1^2} + \frac{6y_2}{y_1^2} + \frac{6}{y_2^2 y_1^2} + \frac{6}{y_1^3} + \frac{4y_2^3}{y_1^3} + \frac{4}{y_1^3} + \frac{6y_2^2}{y_1^4} + 9 \right)
\end{aligned}$$

The previous result – and more generally any decomposition of a product of such characters – can be checked by using a concept of dimension. The dimension of an irreducible representation is the value taken by the associated Weyl SU(n) character at $y_j = 1$ (or the value taken by the su(n) character at $u_j = 0$), for all j . In the same way one can define a “zonal dimension” for an irrep of hw λ as the value taken by the SU(n) zonal character $\chi(\lambda)$ for $y_j = 1$ (or the value taken by the su(n) character for $u_j = 0$). This is a slight terminological abuse since the obtained number is not an integer in general, but this dimension function is obviously compatible with addition and multiplication, as it should. In the case¹³ of SU(3), we conjecture that

$$\dim_Z([\lambda_1, \lambda_2]) = \frac{\lambda_1! \lambda_2! (2\lambda_1 + 2\lambda_2 + 1)!!}{(2\lambda_1 - 1)!! (2\lambda_2 - 1)!! (\lambda_1 + \lambda_2)!}$$

Written $\frac{2\sqrt{\pi}\Gamma(\lambda_1+1)\Gamma(\lambda_2+1)\Gamma(\lambda_1+\lambda_2+\frac{3}{2})}{\Gamma(\lambda_1+\frac{1}{2})\Gamma(\lambda_2+\frac{1}{2})\Gamma(\lambda_1+\lambda_2+1)}$ this expression looks very similar to the standard SU(3) dimension $\dim([\lambda_1, \lambda_2]) = \frac{(1+\lambda_1)(1+\lambda_2)(2+\lambda_1+\lambda_2)}{2} = \frac{\Gamma(\lambda_1+2)\Gamma(\lambda_2+2)\Gamma(\lambda_1+\lambda_2+3)}{2\Gamma(\lambda_1+1)\Gamma(\lambda_2+1)\Gamma(\lambda_1+\lambda_2+2)}$. This zonal dimension is (non-surprisingly) related to the normalization coefficient $Z_p(I)$ that enters (5), and was introduced by A.T. James in [12]:

$$\dim_Z(\lambda) = \frac{Z_{\ell(\lambda)}(I)}{c_{PJ}(\ell(\lambda), \alpha = 2)}$$

It may be interesting to notice that both the numerator and the denominator of this formula are not invariant under a global shift (translation of the partition $\ell(\lambda)$ by an arbitrary integer), but their ratio is invariant —this can be interpreted as a kind of “gauge freedom” in the writing of the SU(n) highest weight λ as a partition.

Going back to our favorite SU(3) example, we can check the consistency of the obtained result for $\chi_Z([1, 1])^2$ in terms of dimensions. The (usual) dimensions of irreps labelled [0,0], [1,1], [0,3], [3,0], [2,2] are 1, 8, 10, 10, 27, and the usual decomposition of $\chi([1, 1])^2$ is compatible with the identity $8 \times 8 = (1 + 2 \times 8 + 1 \times 10 + 1 \times 10 + 1 \times 27)$. The zonal dimensions of the same irreps are 1, 15/2, 7, 7, 35/2, and the decomposition of $\chi_Z([1, 1])^2$ implies $(\frac{15}{2})^2 = \frac{25}{12} + \frac{12}{5} \times \frac{15}{2} + \frac{4}{3} \times 7 + \frac{4}{3} \times 7 + 1 \times \frac{35}{2}$.

Finally, still another way to calculate a structure constant is to use the (α -dependent) Hall inner product, see for instance [18] or [28], for which Jack polynomials – and in particular zonal polynomials – are orthogonal. Using it, one can see for instance that the “multiplicity” of [1, 1] in the decomposition of the square of [1, 1], or of {3, 2, 1} in the square of {2, 1, 0}, which is 2 in the usual case ($\alpha = 1$), and 12/5 in the zonal-P case ($\alpha = 2$) is more generally (i.e., in the Jack-P case) equal to

$$\frac{\langle J_P(\{2, 1, 0\}) J_P(\{2, 1, 0\}), J_P(\{3, 2, 1\}) \rangle}{\langle J_P(\{3, 2, 1\}), J_P(\{3, 2, 1\}) \rangle} = \frac{6\alpha(2\alpha^2 + 11\alpha + 2)}{(\alpha + 2)(2\alpha + 1)(3\alpha + 2)}.$$

¹³In the case of SU(2) we find that the zonal dimension is given by $\dim_Z[\lambda] = \frac{\sqrt{\pi}\Gamma(\lambda+1)}{\Gamma(\lambda+\frac{1}{2})} = \frac{(2\lambda)!!}{(2\lambda-1)!!}$, a formula which is the zonal analog of $\dim[\lambda] = \Gamma[(\lambda + 2)]/\Gamma[(\lambda + 1)] = \lambda + 1$.

In particular we can also find in this way the decomposition of $J_P[\{2, 1, 0\}]^2$:

$$\frac{3\alpha^2(\alpha+3)(2\alpha+1)}{(\alpha+1)^2(\alpha+2)^2} J_P[2, 2, 2] + \frac{6\alpha(2\alpha^2+11\alpha+2)}{(\alpha+2)(2\alpha+1)(3\alpha+2)} J_P[3, 2, 1] + \frac{2\alpha}{\alpha+1} J_P[3, 3, 0] + \frac{2\alpha}{\alpha+1} J_P[4, 1, 1] + J_P[4, 2, 0]$$

and we recover the first and last decompositions already given in sect. 5.1.3 by setting $\alpha = 2$ or 1.

5.3 Back to the PDF (symmetric case) and to the “volume function”

In the Hermitian case it is known that one may associate with a given admissible triple (λ, μ, ν) a convex polytope $\mathcal{H}_{\lambda\mu}^\nu$, the “polytope of honeycombs”, in a $d \leq (n-1)(n-2)/2$ -dimensional space [3]. As recalled above, it is known that the function called \mathcal{J} in [16], which differs from the PDF $\mathbf{p}(\gamma)$ mainly by a Vandermonde factor Δ , measures the volume of $\mathcal{H}_{\lambda\mu}^\nu$ and that it is also a good approximation of the LR multiplicity $N_{\lambda\mu}^\nu$ of that triple¹⁴. More precisely \mathcal{J} is equal to the highest degree coefficient of the stretching (or LR) polynomial that gives the multiplicity when the triple (λ, μ, ν) is scaled by a factor s , i.e., \mathcal{J} is also the dominant coefficient of the Ehrhart polynomial of the polytope $\mathcal{H}_{\lambda\mu}^\nu$. Since this multiplicity is known to be given by the number of *integral* points inside the polytope [3], this property is just expressing that in the large s limit, a semi-classical picture approximates well this number of points by the volume¹⁵ of the polytope.

In the symmetric case multiplicities are not integral, there are no honeycombs (at least the concept was not (yet?) generalized to cover this case), no polytope $\mathcal{H}_{\lambda\mu}^\nu$, and no volume function either. However, as already mentioned, one expects the PDF $\mathbf{p}(\gamma)$, or rather the “volume like function” $\mathcal{J} = \mathbf{p}(\gamma)/\Delta = \rho$, (see (17,24)), to measure, at least in the *generic case* where \mathcal{J} does not vanish, the behavior of “zonal multiplicities” (i.e., appropriate zonal structure constants) under scaling.

A claim going in that direction was actually already made in [9]. Some of the tools developed in the previous subsection could certainly be developed further and we hope to return to this problem someday but we are happy, in the present paper, to show “experimentally” that the overall features of the PDF ρ computed in section 4 are consistent with the values obtained for zonal multiplicities, when the argument (written as a highest weight), is scaled. Remember that the list of eigenvalues $(1, 0, -1)$ chosen for the example studied in sections 2, 3 and 4 differs from the partition $\{2, 1, 0\}$ (aka $[1, 1]$ if reinterpreted in terms of $SU(3)$ highest weights) only by a constant shift¹⁶; we are therefore led to consider the behavior of the zonal structure constants that appear in the reduction of the square of $s\{2, 1, 0\}$, with a scaling factor $s = 1, 2, \dots$, with the plot of the function ρ in terms of Dynkin labels. Using [14] we could perform exact calculations of multiplicities, i.e., obtain the decomposition of the square of $\chi_Z(s[1, 1])$ up to the value $s = 6$ of the scaling factor¹⁷. The same considerations and calculations extend to the quaternionic case, where we compare multiplicities in the decomposition of $\chi_Q([6, 6])^2$ with the function \mathcal{J}_3 computed in sect. 2.

In Fig. 19, we compare the multiplicities obtained for the decomposition of $\chi_Z([6, 6])^2$, $\chi([6, 6])^2$ and $\chi_Q([6, 6])^2$ with the plot of the volume functions ρ or \mathcal{J}_3 . We conclude that already with $s = 6$, the classical limit provided by the “volume” approximates very well the distribution of multiplicities.

¹⁴provided the triple is generic, i.e., provided \mathcal{J} does not vanish.

¹⁵For $n \times n$ matrices, and in the generic case, this is its $(n-1)(n-2)/2$ volume.

¹⁶ In the same way, the partitions that appear in the decomposition of the square of $Z_P(\{2, 1, 0\})$, namely $\{2, 2, 2\}$, $\{3, 2, 1\}$, $\{3, 3, 0\}$, $\{4, 1, 1\}$, $\{4, 2, 0\}$, whose reduction to $SU(3)$ are $\{0, 0, 0\}$, $\{2, 1, 0\}$, $\{3, 3, 0\}$, $\{3, 0, 0\}$, $\{4, 2, 0\}$, and read $[0, 0]$, $[1, 1]$, $[0, 3]$, $[3, 0]$, $[2, 2]$ in terms of Dynkin labels, differ by a constant shift from the following lists of eigenvalues (traceless condition for $\gamma_1, \gamma_2, \gamma_3$ see section 3.1): $\{0, 0, 0\}$, $\{1, 0, -1\}$, $\{1, 1, -2\}$, $\{2, -1, -1\}$, $\{2, 0, -2\}$, and appear in the Horn polygon of Fig. 6 as (special) points with respective coordinates (p, q) given by $(0, 0)$, $(-1, 0)$, $(-3, 2)$, $(-3, -2)$, $(4, 0)$.

¹⁷The explicit reduction of $\chi_Z([s, s]) \times \chi_Z([s, s])$, for $s = 1, 2, \dots$ can be obtained from the web site <http://www.cpt.univ-mrs.fr/~coque/Varia/JackZonalSchurResults.html>




Figure 19: First line: Interpolating function surfaces representing an array of “multiplicities”.
 Left: Decomposition of $\chi_Z([6, 6])^2$ on SU(3) zonal characters.
 Middle: Decomposition of $\chi([6, 6])^2$ on SU(3) Weyl characters.
 Right : Decomposition of $\chi_Q([6, 6])^2$ on SU(3) (zonal) quaternionic characters.
 Second line: Plot of the “volume” function in the three cases. The four colors refer to the four sectors of Fig. 3, but here in Dynkin labels.

6 Conclusion

To summarize:

- we have reproduced the main features of the Horn problem for symmetric matrices and understood the analytic origin of the singularities, at least for $n = 3$, and in detail for the particular case of $A \sim B \sim J_z$;
- we have confirmed, at least in that particular case, that the divergences of the PDF occur on the same locus of non-analyticities as in the Hermitian and quaternionic cases;
- we have also confirmed numerically the connection between the “volume functions” and the (asymptotic) distribution of multiplicities in the product of zonal/Schur/quaternionic polynomials.

This leaves, however, open several issues and room for further progress:

- a more synthetic and general discussion of the singularities in the symmetric case would clearly be desirable. Can one understand their origin from a geometric point of view and assert a priori their location and nature without detailed calculations? Beside the divergences analyzed in the present paper, are there other non-analyticities?
- what happens for higher n and/or for generic $\underline{\beta}$ (or $\alpha = 2/\underline{\beta}$)? The methods developed recently in [29] should be helpful in that respect.

The points we find most challenging are the following:

- There is an enhancement of particular eigenvalues in the Horn spectrum of real symmetric matrices, due to the divergences of the PDF. Is this enhancement observable in some physical process?
- The discussion of sect. 5 has pointed to an analogue of the volume function for real symmetric or quaternionic matrices: is there an underlying geometric interpretation to this “volume”? is there a geometric object generalizing the polytope $\mathcal{H}_{\lambda\mu}^\nu$ of the Hermitian case, whose volume is com-

puted there? on a representation theoretic side, what is the origin of the enhancement of certain multiplicities ?

We leave these questions to the sagacity of our readers...

Acknowledgements

Many thanks to Michèle Vergne for challenging us to carry out the $n = 3$ calculation and to Michel Bauer for his encouragement and in particular for his constructive criticism of identity (28). Stimulating conversations with O. Babelon, É. Brézin, P. Di Francesco, J. Faraut, V. Gorin, S. Majumdar and G. Schehr are also acknowledged.

References

- [1] W. Fulton, Eigenvalues, invariant factors, highest weights, and Schubert calculus, *Bull. Amer. Math. Soc.* **37** (2000) 209–249; <http://arxiv.org/abs/math/9908012>
- [2] A. A. Klyachko, Stable vector bundles and Hermitian operators, *Selecta Math.* (N.S.) **4** (1998) 419–445
- [3] A. Knutson and T. Tao, The honeycomb model of $GL_n(\mathbb{C})$ tensor products I: proof of the saturation conjecture, *J. Amer. Math. Soc.* **12** (1999) 1055–1090, <http://arxiv.org/abs/math/9807160>
- [4] A. Knutson, T. Tao and C. Woodward, The honeycomb model of $GL_n(\mathbb{C})$ tensor products II: Puzzles determine facets of the Littlewood-Richardson cone, *J. Amer. Math. Soc.* **17** (2004), 19–48, <http://arxiv.org/abs/math/0107011>
- [5] A. Knutson and T. Tao, Honeycombs and sums of Hermitian matrices, *Notices Amer. Math. Soc.* **48** no 2, (2001) 175–186; <http://arxiv.org/abs/math/0009048>
- [6] J.-B. Zuber, Horn’s problem and Harish-Chandra’s integrals. Probability distribution functions, *Ann. Inst. Henri Poincaré Comb. Phys. Interact.*, **5** (2018) 309–338, <http://arxiv.org/abs/1705.01186>
- [7] A. Frumkin and A. Goldberger, On the distribution of the spectrum of the sum of two Hermitian or real symmetric matrices, *Adv. Appl. Math.* **37** (2006) 268–286
- [8] J. Faraut, Horn’s problem, and Fourier analysis, April 2018 preprint
- [9] A. Frumkin and A. Goldberger, Diagonals of real symmetric matrices of given spectra as a measure space, <http://arxiv.org/abs/1505.06418v1>
- [10] Harish-Chandra, Differential Operators on a Semisimple Algebra, *Amer. J. Math.* **79** (1957) 87–120
- [11] C. Itzykson, J.-B. Zuber, The planar approximation II, *J. Math. Phys.* **21** (1980) 411–421
- [12] A.T. James, The distribution of the latent roots of the covariance matrix. *The Annals of Mathematical Statistics*, **31** (1960) 151–158
- [13] É. Brézin and S. Hikami, An extension of the Harish-Chandra–Itzykson–Zuber integral, <http://arxiv.org/abs/math-ph/0208002v1>; WKB-expansion of the Harish-Chandra–Itzykson–Zuber integral for arbitrary β , *Progress of Theoretical Physics*, **116** (2006) 441–502, <http://arxiv.org/abs/math-ph/0604041v1>

- [14] R. Coquereaux, Mathematica package “SymPol\$Package”, available on the web page http://www.cpt.univ-mrs.fr/~coque/Computer_programs/index.html and on Github, see <https://github.com/RobertCoquereaux/Lie-AffineLie-Representations>.
- [15] I.M. Gelfand and G.E. Shilov, *Generalized Functions*, vol.1, chap. 3, Academic Press 1964
- [16] R. Coquereaux and J.-B. Zuber, From orbital measures to Littlewood–Richardson coefficients and hive polytopes, *Ann. Inst. Henri Poincaré Comb. Phys. Interact.*, **5** (2018) 339–386, <http://arxiv.org/abs/1706.02793>
- [17] A.M. Mathai, S. B. Provost, and T. Hayakawa, *Bilinear Forms and Zonal Polynomials*, Lecture Notes in Statistics 102, Springer Verlag, 1995
- [18] I. G. Macdonald, *Symmetric Functions and Hall Polynomials*, Oxford Science Publications 1998
- [19] Hua Lo-Ken, Harmonic analysis of functions of several complex variables in the classical domains, *AMS Translations*, **6** (1963) (1958 in Chinese)
- [20] A.T. James, Normal multivariate analysis and the orthogonal group, *The Annals of Mathematical Statistics*, **25** (1954) 40–75
- [21] A.T. James, Zonal polynomials of the real positive definite symmetric matrices. *Annals of Mathematics*, **74** (1961) 456–469
- [22] A. G. Constantine, Some noncentral distribution problems in multivariate analysis, *The Annals of Mathematical Statistics*, **34** (1963) 1270–1285
- [23] I. Dumitriu, A. Edelman and G. Shuman, MOPS: Multivariate Orthogonal Polynomials (symbolically), preprint 2004, <http://arxiv.org/abs/math-ph/0409066>
- [24] W. Baratta, 2008, <http://www.ms.unimelb.edu.au/~wbaratta/index.html>
- [25] J. Demmel and P. Koev, Accurate and efficient evaluation of Schur and Jack functions, *Mathematics of Computation*, **75** (2005) 223–239
- [26] V. Feray and P. Sniady, Zonal polynomials via Stanley’s coordinates and free cumulants, *Journal of Algebra* **334** (2010) 338–373
- [27] R. Coquereaux and J.-B. Zuber, *On some properties of SU(3) Fusion Coefficients*, Nucl. Phys. **B 912** (2016) 119–150 , <http://arxiv.org/abs/1605.05864>
- [28] R.P. Stanley, Some Combinatorial Properties of Jack Symmetric Functions, *Advances in Mathematics* **77** (1989) 76–115
- [29] V. Gorin and A. W. Marcus, Crystallization of random matrix orbits, <http://arxiv.org/abs/1706.07393> ;
A. Bufetov and V. Gorin, Fourier transform on high-dimensional unitary groups with applications to random tilings, <http://arxiv.org/abs/1712.09925>