



**HAL**  
open science

# Some new inequalities of the Grüss type for conformable fractional integrals

Gauhar Rahman, Sooppy Nisar, Feng Qi

► **To cite this version:**

Gauhar Rahman, Sooppy Nisar, Feng Qi. Some new inequalities of the Grüss type for conformable fractional integrals: Inequalities for conformable fractional integrals. 2018. hal-01871682v1

**HAL Id: hal-01871682**

**<https://hal.science/hal-01871682v1>**

Preprint submitted on 11 Sep 2018 (v1), last revised 19 Jan 2020 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# SOME NEW INEQUALITIES OF THE GRÜSS TYPE FOR CONFORMABLE FRACTIONAL INTEGRALS

GAUHAR RAHMAN, KOTTAKKARAN SOOPPY NISAR, AND FENG QI

ABSTRACT. In the paper, the authors establish some new inequalities of the Grüss type for conformable fractional integrals. These inequalities generalize some known results.

## 1. INTRODUCTION

In [7], Grüss showed an integral inequality which connects the integral of the product of two functions and the product of integrals for these two function. This inequality reads that, if  $f$  and  $g$  are two continuous functions on  $[a, b]$  satisfying  $m \leq f(\tau) \leq M$  and  $n \leq g(\tau) \leq N$  for all  $\tau \in [a, b]$  and  $m, M, n, N \in \mathbb{R}$ , then

$$\left| \frac{1}{b-a} \int_a^b f(\tau)g(\tau) dt - \frac{1}{(b-a)^2} \int_a^b f(\tau) dt \int_a^b g(\tau) dt \right| \leq \frac{1}{4}(M-m)(N-n). \quad (1)$$

For more information on the Grüss inequality (1), please refer to [11, Chapter X] and closely related references therein.

In the latest decades, the fractional integral inequalities involving the Riemann–Liouville fractional integrals have been widely studied by various researchers. The interested readers can refer to the work in [1, 2, 3, 5, 15, 20, 22]. In [3], Dahmani introduced the following fractional integral inequalities for the Riemann–Liouville fractional integrals: if  $f$  and  $g$  are two integrable functions on  $[0, \infty)$  satisfying  $m \leq f(\tau) \leq M$  and  $n \leq g(\tau) \leq N$  for all  $\tau \in [0, \infty)$  and  $m, M, n, N \in \mathbb{R}$ , then, for all  $\tau, \alpha, \beta > 0$ , we have

$$\left| \frac{\tau^\alpha}{\Gamma(\alpha+1)} \mathfrak{J}^\alpha(fg)(\tau) - \mathfrak{J}^\alpha f(\tau) \mathfrak{J}^\alpha g(\tau) \right| \leq \left[ \frac{\tau^\alpha}{\Gamma(\alpha+1)} \right]^2 (M-m)(N-n)$$

and

$$\begin{aligned} & \left[ \frac{\tau^\alpha}{\Gamma(\alpha+1)} \mathfrak{J}^\beta(fg)(\tau) - \frac{\tau^\beta}{\Gamma(\beta+1)} \mathfrak{J}^\alpha(fg)(\tau) - \mathfrak{J}^\alpha f(\tau) \mathfrak{J}^\beta g(\tau) - \mathfrak{J}^\beta f(\tau) \mathfrak{J}^\alpha g(\tau) \right]^2 \\ & \leq \left[ \left( M \frac{\tau^\alpha}{\Gamma(\alpha+1)} - \mathfrak{J}^\alpha f(\tau) \right) \left( \mathfrak{J}^\beta f(\tau) - m \frac{\tau^\beta}{\Gamma(\beta+1)} \right) + \left( \mathfrak{J}^\alpha f(\tau) - m \frac{\tau^\alpha}{\Gamma(\alpha+1)} \right) \right. \\ & \quad \times \left. \left( M \frac{\tau^\beta}{\Gamma(\beta+1)} - \mathfrak{J}^\beta f(\tau) \right) \right] \left[ \left( N \frac{\tau^\alpha}{\Gamma(\alpha+1)} - \mathfrak{J}^\alpha g(\tau) \right) \left( \mathfrak{J}^\beta g(\tau) - n \frac{\tau^\beta}{\Gamma(\beta+1)} \right) \right. \\ & \quad \left. + \left( \mathfrak{J}^\alpha g(\tau) - n \frac{\tau^\alpha}{\Gamma(\alpha+1)} \right) \left( N \frac{\tau^\beta}{\Gamma(\beta+1)} - \mathfrak{J}^\beta g(\tau) \right) \right], \end{aligned}$$

---

2010 *Mathematics Subject Classification.* Primary 26A33; Secondary 26D10, 26D15.

*Key words and phrases.* Riemann–Liouville fractional integral; inequality of the Grüss type; conformable fractional integral; integral inequality; fractional integral operator.

This paper was typeset using  $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$ .

where  $\Gamma$  is the classical Euler gamma function [12, 13, 19] and the Riemann–Liouville fractional integral  $\mathfrak{J}^\mu$  of order  $\mu \geq 0$  for a function  $f \in L^1((0, \infty), \mathbb{R})$  is defined [10] by  $\mathfrak{J}^0 f(x) = f(x)$  and

$$\mathfrak{J}^\mu f(x) = \frac{1}{\Gamma(\mu)} \int_0^x (x - \tau)^{\mu-1} f(\tau) d\tau, \quad \Re(\mu) > 0.$$

The Riemann–Liouville fractional integrals  $\mathfrak{J}_{\alpha+}^\mu$  and  $\mathfrak{J}_{\beta-}^\mu$  of order  $\mu > 0$  are defined respectively by

$$\mathfrak{J}_{\alpha+}^\mu f(x) = \frac{1}{\Gamma(\mu)} \int_\alpha^x (x - \tau)^{\mu-1} f(\tau) d\tau, \quad x > \alpha, \quad \Re(\mu) > 0 \quad (2)$$

and

$$\mathfrak{J}_{\beta-}^\mu f(x) = \frac{1}{\Gamma(\mu)} \int_x^\beta (x - \tau)^{\mu-1} f(\tau) d\tau, \quad x < \beta, \quad \Re(\mu) > 0. \quad (3)$$

For more details about fractional integral operators (2) and (3), please refer to [4, 6, 9, 10, 14, 16] and closely related references.

The left and right sided fractional conformable integral operators are respectively defined [8] by

$${}^\lambda \mathfrak{J}_{a+}^\mu f(x) = \frac{1}{\Gamma(\lambda)} \int_a^x \left[ \frac{(x-a)^\mu - (\tau-a)^\mu}{\mu} \right]^{\lambda-1} \frac{f(\tau)}{(\tau-a)^{1-\mu}} d\tau \quad (4)$$

and

$${}^\lambda \mathfrak{J}_{b-}^\mu f(x) = \frac{1}{\Gamma(\lambda)} \int_x^b \left[ \frac{(b-x)^\mu - (b-\tau)^\mu}{\mu} \right]^{\lambda-1} \frac{f(\tau)}{(b-\tau)^{1-\mu}} d\tau, \quad (5)$$

where  $\Re(\lambda) > 0$ . Obviously, if taking  $a = 0$  and  $\mu = 1$ , then (4) reduces to the Riemann–Liouville fractional integral (2). Similarly, if setting  $b = 0$  and  $\mu = 1$ , then (5) becomes the Riemann–Liouville fractional integral (3).

In [17] the conformable fractional integral

$${}^\beta \mathfrak{J}^\mu f(x) = \frac{1}{\Gamma(\beta)} \int_0^x \left( \frac{x^\mu - \tau^\mu}{\mu} \right)^{\beta-1} \frac{f(\tau)}{\tau^{1-\mu}} d\tau. \quad (6)$$

was defined. From (6), One can obtain easily that  ${}^\alpha \mathfrak{J}^\mu {}^\beta \mathfrak{J}^\mu f(x) = {}^{\beta+\alpha} \mathfrak{J}^\mu f(x)$  and  ${}^\alpha \mathfrak{J}^\mu {}^\beta \mathfrak{J}^\mu f(x) = {}^\beta \mathfrak{J}^\mu {}^\alpha \mathfrak{J}^\mu f(x)$ .

In this paper, we will employ the conformable fractional integral (6) to establish some new inequalities of the Grüss type for conformable fractional integrals.

## 2. MAIN RESULTS

We are now in a position to state and prove our main results.

**Theorem 2.1.** *Let  $f$  be an integrable function on  $[0, \infty)$ . Assume that there exist two integrable functions  $\phi_1, \phi_2$  on  $[0, \infty)$  such that*

$$\phi_1(x) \leq f(x) \leq \phi_2(x), \quad x \in [0, \infty). \quad (7)$$

Then, for  $x, \alpha, \beta > 0$ , we have

$${}^\beta \mathfrak{J}^\mu \phi_1(x) {}^\alpha \mathfrak{J}^\mu f(x) + {}^\alpha \mathfrak{J}^\mu \phi_2(x) {}^\beta \mathfrak{J}^\mu f(x) \geq {}^\alpha \mathfrak{J}^\mu \phi_2(x) {}^\beta \mathfrak{J}^\mu \phi_1(x) + {}^\alpha \mathfrak{J}^\mu f(x) {}^\beta \mathfrak{J}^\mu f(x). \quad (8)$$

*Proof.* From (7), for all  $\tau, \rho \geq 0$ , it follows that

$$[\phi_2(\tau) - f(\tau)][f(\rho) - \phi_1(\rho)] \geq 0.$$

Therefore, we have

$$\phi_2(\tau)f(\rho) + \phi_1(\rho)f(\tau) \geq \phi_1(\rho)\phi_2(\tau) + f(\tau)f(\rho). \quad (9)$$

Multiplying both sides of (9) by  $\frac{1}{\Gamma(\alpha)}\left(\frac{x^\mu - \tau^\mu}{\mu}\right)^{\alpha-1}\tau^{\mu-1}$  and integrating over  $\tau \in (0, x)$  lead to

$$\begin{aligned} f(\rho) \int_0^x \frac{1}{\Gamma(\alpha)} \left(\frac{x^\mu - \tau^\mu}{\mu}\right)^{\alpha-1} \tau^{\alpha-1} \phi_2(\tau) d\tau \\ + \phi_1(\rho) \int_0^x \frac{1}{\Gamma(\alpha)} \left(\frac{x^\mu - \tau^\mu}{\mu}\right)^{\alpha-1} \tau^{\mu-1} f(\tau) d\tau \\ \geq \phi_1(\rho) \int_0^x \frac{1}{\Gamma(\alpha)} \left(\frac{x^\mu - \tau^\mu}{\mu}\right)^{\alpha-1} \tau^{\mu-1} \phi_2(\tau) d\tau \\ + f(\rho) \int_0^x \frac{1}{\Gamma(\alpha)} \left(\frac{x^\mu - \tau^\mu}{\mu}\right)^{\alpha-1} \tau^{\mu-1} f(\tau) d\tau \end{aligned}$$

which gives

$$f(\rho)^\alpha \mathfrak{J}^\mu \phi_2(x) + \phi_1(\rho)^\alpha \mathfrak{J}^\mu f(x) \geq \phi_1(\rho)^\alpha \mathfrak{J}^\mu \phi_2(x) + f(\rho)^\alpha \mathfrak{J}^\mu f(x). \quad (10)$$

Multiplying both sides of (10) by  $\frac{1}{\Gamma(\beta)}\left(\frac{x^\mu - \rho^\mu}{\mu}\right)^{\beta-1}\rho^{\mu-1}$  and integrating over  $\rho \in (0, x)$  result in

$$\begin{aligned} {}^\alpha \mathfrak{J}^\mu \phi_2(x) \int_0^x \frac{1}{\Gamma(\beta)} \left(\frac{x^\mu - \rho^\mu}{\mu}\right)^{\beta-1} \rho^{\mu-1} f(\rho) d\rho \\ + {}^\alpha \mathfrak{J}^\mu f(x) \int_0^x \frac{1}{\Gamma(\beta)} \left(\frac{x^\mu - \rho^\mu}{\mu}\right)^{\beta-1} \rho^{\mu-1} \phi_1(\rho) d\rho \\ \geq {}^\alpha \mathfrak{J}^\mu \phi_2(x) \int_0^x \frac{1}{\Gamma(\beta)} \left(\frac{x^\mu - \rho^\mu}{\mu}\right)^{\beta-1} \rho^{\mu-1} \phi_1(\rho) d\rho \\ + {}^\alpha \mathfrak{J}^\mu f(x) \int_0^x \frac{1}{\Gamma(\beta)} \left(\frac{x^\mu - \rho^\mu}{\mu}\right)^{\beta-1} \rho^{\mu-1} f(\rho) d\rho. \end{aligned}$$

which gives the required inequality (8).  $\square$

From Theorem 2.1, we can derive the following two corollaries.

**Corollary 2.1.** *Let  $f$  be an integrable function on  $[0, \infty)$  satisfying  $m \leq f(x) \leq M$  for all  $x \in [0, \infty)$  and  $m, M \in \mathbb{R}$ . Then, for  $x, \alpha, \beta > 0$ , we have*

$$\begin{aligned} m \frac{x^{\mu\beta}}{\mu^\beta \Gamma(\beta+1)} {}^\alpha \mathfrak{J}^\mu f(x) + M \frac{x^{\mu\alpha}}{\mu^\alpha \Gamma(\alpha+1)} {}^\beta \mathfrak{J}^\mu f(x) \\ \geq mM \frac{x^{\mu(\alpha+\beta)}}{\mu^{\alpha+\beta} \Gamma(\alpha+1) \Gamma(\beta+1)} + {}^\alpha \mathfrak{J}^\mu f(x) {}^\beta \mathfrak{J}^\mu f(x). \end{aligned}$$

**Corollary 2.2.** *Let  $f$  be an integrable function on  $[0, \infty)$  satisfying  $x^\mu \leq f(x) \leq x^\mu + 1$  for all  $x \in [0, \infty)$ . Then, for  $x, \alpha > 0$ , we have*

$$\begin{aligned} & \left[ \frac{2x^{\mu(\alpha+1)}}{\mu^\alpha \Gamma(\alpha+2)} + \frac{x^{\mu\alpha}}{\mu^\alpha \Gamma(\alpha+1)} \right] {}^\alpha \mathcal{J}^\mu f(x) \\ & \geq \left[ \frac{x^{\mu(\alpha+1)}}{\mu^\alpha \Gamma(\alpha+2)} + \frac{x^{\mu\alpha}}{\mu^\alpha \Gamma(\alpha+1)} \right] \left[ \frac{x^{\mu(\alpha+1)}}{\mu^\alpha \Gamma(\alpha+2)} \right] + [{}^\alpha \mathcal{J}^\mu f(x)]^2. \end{aligned}$$

**Theorem 2.2.** *Let  $f$  and  $g$  be two integrable function on  $[0, \infty)$ . Suppose that the inequality (7) holds and that there exist two integrable functions  $\psi_1$  and  $\psi_2$  on  $[0, \infty)$  such that*

$$\psi_1(x) \leq g(x) \leq \psi_2(x), \quad t \in [0, \infty). \quad (11)$$

Then, for  $x, \alpha, \beta > 0$ , the following four inequalities hold:

$${}^\beta \mathcal{J}^\mu \psi_1(x) {}^\alpha \mathcal{J}^\mu f(x) + {}^\alpha \mathcal{J}^\mu \phi_2(x) {}^\beta \mathcal{J}^\mu g(x) \geq {}^\alpha \mathcal{J}^\mu \phi_2(x) {}^\beta \mathcal{J}^\mu \psi_1(x) + {}^\alpha \mathcal{J}^\mu f(x) {}^\beta \mathcal{J}^\mu g(x), \quad (12)$$

$${}^\beta \mathcal{J}^\mu \phi_1(x) {}^\alpha \mathcal{J}^\mu g(x) + {}^\alpha \mathcal{J}^\mu \psi_2(x) {}^\beta \mathcal{J}^\mu f(x) \geq {}^\alpha \mathcal{J}^\mu \phi_1(x) {}^\beta \mathcal{J}^\mu \psi_2(x) + {}^\alpha \mathcal{J}^\mu f(x) {}^\beta \mathcal{J}^\mu g(x), \quad (13)$$

$${}^\alpha \mathcal{J}^\mu \phi_2(x) {}^\beta \mathcal{J}^\mu \psi_2(x) + {}^\alpha \mathcal{J}^\mu f(x) {}^\beta \mathcal{J}^\mu g(x) \geq {}^\alpha \mathcal{J}^\mu \phi_2(x) {}^\beta \mathcal{J}^\mu g(x) + {}^\beta \mathcal{J}^\mu \psi_2(x) {}^\alpha \mathcal{J}^\mu f(x), \quad (14)$$

$${}^\alpha \mathcal{J}^\mu \phi_1(x) {}^\beta \mathcal{J}^\mu \psi_1(x) + {}^\alpha \mathcal{J}^\mu f(x) {}^\beta \mathcal{J}^\mu g(x) \geq {}^\alpha \mathcal{J}^\mu \phi_1(x) {}^\beta \mathcal{J}^\mu g(x) + {}^\alpha \mathcal{J}^\mu f(x) {}^\beta \mathcal{J}^\mu \psi_1(x). \quad (15)$$

*Proof.* From (7) and (11) and for  $x \in [0, \infty)$ , we have

$$[\phi_2(\tau) - f(\tau)][g(\rho) - \psi_1(\rho)] \geq 0.$$

Therefore, it follows that

$$\phi_2(\tau)g(\rho) + \psi_1(\rho)f(\tau) \geq \psi_1(\rho)\phi_2(\tau) + f(\tau)g(\rho). \quad (16)$$

Multiplying both sides of (16) by  $\frac{1}{\Gamma(\alpha)} \left( \frac{x^\mu - \tau^\mu}{\mu} \right)^{\alpha-1} \tau^{\mu-1}$  and integrating over  $\tau \in (0, x)$  arrive at

$$\begin{aligned} g(\rho) \int_0^x \frac{1}{\Gamma(\alpha)} \left( \frac{x^\mu - \tau^\mu}{\mu} \right)^{\alpha-1} \tau^{\mu-1} \phi_2(\tau) d\tau \\ + \psi_1(\rho) \int_0^x \frac{1}{\Gamma(\alpha)} \left( \frac{x^\mu - \tau^\mu}{\mu} \right)^{\alpha-1} \tau^{\mu-1} f(\tau) d\tau \\ \geq \psi_1(\rho) \int_0^x \frac{1}{\Gamma(\alpha)} \left( \frac{x^\mu - \tau^\mu}{\mu} \right)^{\alpha-1} \tau^{\mu-1} \phi_2(\tau) d\tau \\ + g(\rho) \int_0^x \frac{1}{\Gamma(\alpha)} \left( \frac{x^\mu - \tau^\mu}{\mu} \right)^{\alpha-1} \tau^{\mu-1} f(\tau) d\tau \end{aligned}$$

which gives

$$g(\rho) {}^\alpha \mathcal{J}^\mu \phi_2(x) + \psi_1(\rho) {}^\alpha \mathcal{J}^\mu f(x) \geq \psi_1(\rho) {}^\alpha \mathcal{J}^\mu \phi_2(x) + g(\rho) {}^\alpha \mathcal{J}^\mu f(x). \quad (17)$$

Multiplying both sides of (17) by  $\frac{1}{\Gamma(\beta)} \left( \frac{x^\mu - \rho^\mu}{\mu} \right)^{\beta-1} \rho^{\mu-1}$  and integrating over  $\rho \in (0, x)$  reveal

$$\begin{aligned} {}^\alpha \mathcal{J}^\mu \phi_2(x) \int_0^x \frac{1}{\Gamma(\beta)} \left( \frac{x^\mu - \rho^\mu}{\mu} \right)^{\beta-1} \rho^{\mu-1} g(\rho) d\rho \\ + {}^\alpha \mathcal{J}^\mu f(x) \int_0^x \frac{1}{\Gamma(\beta)} \left( \frac{x^\mu - \rho^\mu}{\mu} \right)^{\beta-1} \rho^{\mu-1} \psi_1(\rho) d\rho \end{aligned}$$

$$\begin{aligned} &\geq {}^\alpha \mathfrak{J}^\mu \phi_2(x) \int_0^x \frac{1}{\Gamma(\beta)} \left( \frac{x^\mu - \rho^\mu}{\mu} \right)^{\beta-1} \rho^{\mu-1} \psi_1(\rho) \, d\rho \\ &\quad + {}^\alpha \mathfrak{J}^\mu f(x) \int_0^x \frac{1}{\Gamma(\beta)} \left( \frac{x^\mu - \rho^\mu}{\mu} \right)^{\beta-1} \rho^{\mu-1} g(\rho) \, d\rho. \end{aligned}$$

which gives the required inequality (12).

Making use of the inequalities

$$[\psi_2(\tau) - g(\tau)][f(\rho) - \phi_1(\rho_1)] \geq 0, \quad [\psi_2(\tau) - f(\tau)][g(\rho) - \psi_2(\rho_1)] \leq 0,$$

and  $[\phi_1(\tau) - f(\tau)][g(\rho) - \psi_1(\rho_1)] \leq 0$ , we can prove (13) to (15).  $\square$

**Corollary 2.3.** *Let  $f$  and  $g$  be two integrable functions on  $[0, \infty)$ . Assume that there exist  $m, M, n, N \in \mathbb{R}$  such that*

$$m \leq f(x) \leq M, \quad n \leq g(x) \leq N, \quad t \in [0, \infty).$$

Then we have the following four inequalities

$$\begin{aligned} &n \frac{x^{\mu\beta}}{\mu^\beta \Gamma(\beta+1)} {}^\alpha \mathfrak{J}^\mu f(x) + M \frac{x^{\mu\alpha}}{\mu^\alpha \Gamma(\alpha+1)} {}^\beta \mathfrak{J}^\mu g(x) \\ &\geq nM \frac{x^{\mu(\alpha+\beta)}}{\mu^{\alpha+\beta} \Gamma(\alpha+1) \Gamma(\beta+1)} + {}^\alpha \mathfrak{J}^\mu f(x) {}^\beta \mathfrak{J}^\mu g(x), \\ &m \frac{x^{\mu\beta}}{\mu^\beta \Gamma(\beta+1)} {}^\alpha \mathfrak{J}^\mu g(x) + N \frac{x^{\mu\alpha}}{\mu^\alpha \Gamma(\alpha+1)} {}^\beta \mathfrak{J}^\mu f(x) \\ &\geq mN \frac{x^{\mu(\alpha+\beta)}}{\mu^{\alpha+\beta} \Gamma(\alpha+1) \Gamma(\beta+1)} + {}^\beta \mathfrak{J}^\mu f(x) {}^\alpha \mathfrak{J}^\mu g(x), \\ &MN \frac{x^{\mu(\alpha+\beta)}}{\mu^{\alpha+\beta} \Gamma(\alpha+1) \Gamma(\beta+1)} + {}^\beta \mathfrak{J}^\mu f(x) {}^\alpha \mathfrak{J}^\mu g(x) \\ &\geq M \frac{x^{\mu\alpha}}{\mu^\alpha \Gamma(\alpha+1)} {}^\beta \mathfrak{J}^\mu g(x) + N \frac{x^{\mu\beta}}{\mu^\beta \Gamma(\beta+1)} {}^\alpha \mathfrak{J}^\mu f(x), \end{aligned}$$

and

$$\begin{aligned} &mn \frac{x^{\mu(\alpha+\beta)}}{\mu^{\alpha+\beta} \Gamma(\alpha+1) \Gamma(\beta+1)} + {}^\beta \mathfrak{J}^\mu f(x) {}^\alpha \mathfrak{J}^\mu g(x) \\ &\geq m \frac{x^{\mu\alpha}}{\mu^\alpha \Gamma(\alpha+1)} {}^\beta \mathfrak{J}^\mu g(x) + n \frac{x^{\mu\beta}}{\mu^\beta \Gamma(\beta+1)} {}^\alpha \mathfrak{J}^\mu f(x). \end{aligned}$$

**Theorem 2.3.** *Let  $f$  be an integrable function on  $[0, \infty)$  and let  $\phi_1$  and  $\phi_2$  be two integrable functions on  $[0, \infty)$ . Assume that the inequality (7) holds. Then, for  $x, \alpha > 0$ , we have*

$$\begin{aligned} &\frac{x^{\mu\alpha}}{\mu^\alpha \Gamma(\alpha+1)} {}^\alpha \mathfrak{J}^\mu f^2(x) - [{}^\alpha \mathfrak{J}^\mu f(x)]^2 \\ &= [{}^\alpha \mathfrak{J}^\mu \phi_2(x) - {}^\alpha \mathfrak{J}^\mu f(x)][{}^\alpha \mathfrak{J}^\mu f(x) - {}^\alpha \mathfrak{J}^\mu \phi_1(x)] \\ &\quad - \frac{x^{\mu\alpha}}{\mu^\alpha \Gamma(\alpha+1)} {}^\alpha \mathfrak{J}^\mu [\phi_2(x) - f(x)][f(x) - \phi_1(x)] \\ &\quad + \frac{x^{\mu\alpha}}{\mu^\alpha \Gamma(\alpha+1)} {}^\alpha \mathfrak{J}^\mu \phi_1 f(x) - {}^\alpha \mathfrak{J}^\mu \phi_1(x) {}^\alpha \mathfrak{J}^\mu f(x) \end{aligned}$$

$$\begin{aligned}
& + \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} {}^\alpha\mathcal{J}^\mu\phi_2 f(x) - {}^\alpha\mathcal{J}^\mu\phi_2(x) {}^\alpha\mathcal{J}^\mu f(x) \\
& \quad + {}^\alpha\mathcal{J}^\mu\phi_1(x) {}^\alpha\mathcal{J}^\mu\phi_2(x) - \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} {}^\alpha\mathcal{J}^\mu\phi_1\phi_2(x). \quad (18)
\end{aligned}$$

*Proof.* For  $\tau, \rho > 0$ , we have

$$\begin{aligned}
& [\phi_2(\rho) - f(\rho)][f(\tau) - \phi_1(\tau)] + [\phi_2(\tau) - f(\tau)][f(\rho) - \phi_1(\rho)] \\
& \quad - [\phi_2(\tau) - f(\tau)][f(\tau) - \phi_1(\tau)] - [\phi_2(\rho) - f(\rho)][f(\rho) - \phi_1(\rho)] \\
& = f^2(\tau) + f^2(\rho) - 2f(\tau)f(\rho) + \phi_2(\rho)f(\tau) + \phi_1(\tau)f(\rho) - \phi_1(\tau)\phi_2(\rho) \\
& \quad + \phi_2(\tau)f(\rho) + \phi_1(\rho)f(\tau) - \phi_1(\rho)\phi_2(\tau) - \phi_2(\tau)f(\tau) + \phi_1(\tau)\phi_2(\tau) \\
& \quad - \phi_1(\tau)f(\tau) - \phi_2(\rho)f(\rho) + \phi_1(\rho)\phi_2(\rho) - \phi_1(\rho)f(\rho). \quad (19)
\end{aligned}$$

Multiplying both sides of (19) by  $\frac{1}{\Gamma(\alpha)}\left(\frac{x^\mu - \tau^\mu}{\mu}\right)^{\alpha-1}\tau^{\mu-1}$  and integrating over  $\tau \in (0, x)$  yield

$$\begin{aligned}
& [\phi_2(\rho) - f(\rho)][{}^\alpha\mathcal{J}^\mu f(x) - {}^\alpha\mathcal{J}^\mu\phi_1(x)] + [{}^\alpha\mathcal{J}^\mu\phi_2(x) - {}^\alpha\mathcal{J}^\mu f(x)][f(\rho) - \phi_1(\rho)] \\
& \quad - {}^\alpha\mathcal{J}^\mu[\phi_2(x) - f(x)][f(x) - \phi_1(x)] - [\phi_2(\rho) - f(\rho)][f(\rho) - \phi_1(\rho)] \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} \\
& \quad = {}^\alpha\mathcal{J}^\mu f^2(x) + f^2(\rho) \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} - 2f(\rho) {}^\alpha\mathcal{J}^\mu f(x) + \phi_2(\rho) {}^\alpha\mathcal{J}^\mu f(x) \\
& \quad + f(\rho) {}^\alpha\mathcal{J}^\mu\phi_1(x) - \phi_2(\rho) {}^\alpha\mathcal{J}^\mu\phi_1(x) + f(\rho) {}^\alpha\mathcal{J}^\mu\phi_2(x) + \phi_1(\rho) {}^\alpha\mathcal{J}^\mu f(x) \\
& \quad \quad - \phi_1(\rho) {}^\alpha\mathcal{J}^\mu\phi_2(x) - {}^\alpha\mathcal{J}^\mu\phi_2 f(x) + {}^\alpha\mathcal{J}^\mu\phi_1\phi_2(x) - {}^\alpha\mathcal{J}^\mu\phi_1 f(x) \\
& \quad - \phi_2(\rho)f(\rho) \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} + \phi_1(\rho)\phi_2(\rho) \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} - \phi_1(\rho)f(\rho) \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)}. \quad (20)
\end{aligned}$$

Multiplying both sides of (20) by  $\frac{1}{\Gamma(\alpha)}\left(\frac{x^\mu - \rho^\mu}{\mu}\right)^{\alpha-1}\rho^{\mu-1}$  and integrating the resultant identity with respect to  $\rho$  from 0 to  $x$  bring out

$$\begin{aligned}
& [{}^\alpha\mathcal{J}^\mu\phi_2(x) - {}^\alpha\mathcal{J}^\mu f(x)][{}^\alpha\mathcal{J}^\mu f(x) - {}^\alpha\mathcal{J}^\mu\phi_1(x)] + [{}^\alpha\mathcal{J}^\mu\phi_2(x) - {}^\alpha\mathcal{J}^\mu f(x)] \\
& \quad \times [{}^\alpha\mathcal{J}^\mu f(x) - {}^\alpha\mathcal{J}^\mu\phi_1(x)] - {}^\alpha\mathcal{J}^\mu[\phi_2(x) - f(x)][f(x) - \phi_1(x)] \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} \\
& \quad \quad - {}^\alpha\mathcal{J}^\mu[\phi_2(\rho) - f(\rho)][f(\rho) - \phi_1(\rho)] \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} \\
& \quad = \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} {}^\alpha\mathcal{J}^\mu f^2(x) + \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} {}^\alpha\mathcal{J}^\mu f^2(x) - 2{}^\alpha\mathcal{J}^\mu f(x) {}^\alpha\mathcal{J}^\mu f(x) \\
& \quad \quad + {}^\alpha\mathcal{J}^\mu\phi_2(x) {}^\alpha\mathcal{J}^\mu f(x) + {}^\alpha\mathcal{J}^\mu f(x) {}^\alpha\mathcal{J}^\mu\phi_1(x) - {}^\alpha\mathcal{J}^\mu\phi_2(x) {}^\alpha\mathcal{J}^\mu\phi_1(x) \\
& \quad \quad + {}^\alpha\mathcal{J}^\mu f(x) {}^\alpha\mathcal{J}^\mu\phi_2(x) + {}^\alpha\mathcal{J}^\mu\phi_1(x) {}^\alpha\mathcal{J}^\mu f(x) - {}^\alpha\mathcal{J}^\mu\phi_1(x) {}^\alpha\mathcal{J}^\mu\phi_2(x) \\
& \quad - \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} {}^\alpha\mathcal{J}^\mu\phi_2 f(x) + \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} {}^\alpha\mathcal{J}^\mu\phi_1\phi_2(x) - \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} {}^\alpha\mathcal{J}^\mu\phi_1 f(x) \\
& \quad - \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} {}^\alpha\mathcal{J}^\mu\phi_2 f(x) + \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} {}^\alpha\mathcal{J}^\mu\phi_1\phi_2(x) - \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} {}^\alpha\mathcal{J}^\mu\phi_1 f(x)
\end{aligned}$$

which yields the required inequality (18).  $\square$

**Corollary 2.4.** *Let  $f$  be an integrable function on  $[0, \infty)$  satisfying  $m \leq f(t) \leq M$  for all  $x \in [0, \infty)$ . Then, for all  $x, \alpha > 0$ , we have*

$$\begin{aligned} \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} \alpha\mathfrak{J}^\mu f^2(x) - [\alpha\mathfrak{J}^\mu f(x)]^2 &= \left[ M \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} - \alpha\mathfrak{J}^\mu f(x) \right] \\ &\times \left[ \alpha\mathfrak{J}^\mu f(x) - m \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} \right] - \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} \alpha\mathfrak{J}^\mu [M - f(x)][f(x) - m]. \end{aligned}$$

**Theorem 2.4.** *Let  $f, g, \phi_1, \phi_2, \psi_1$ , and  $\psi_2$  be integrable functions on  $[0, \infty)$  satisfying (7) and (11) on  $[0, \infty)$ . Then, for all  $x, \alpha > 0$ , we have*

$$\left| \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} \alpha\mathfrak{J}^\mu fg(x) - \alpha\mathfrak{J}^\mu f(x) \alpha\mathfrak{J}^\mu g(x) \right| \leq \sqrt{T(f, \phi_1, \phi_2)T(g, \psi_1, \psi_2)}, \quad (21)$$

where

$$\begin{aligned} T(u, v, w) &= [\alpha\mathfrak{J}^\mu w(x) - \alpha\mathfrak{J}^\mu u(x)][\alpha\mathfrak{J}^\mu u(x) - \alpha\mathfrak{J}^\mu v(x)] \\ &+ \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} \alpha\mathfrak{J}^\mu vu(x) - \alpha\mathfrak{J}^\mu v(x) \alpha\mathfrak{J}^\mu u(x) + \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} \alpha\mathfrak{J}^\mu wu(x) \\ &- \alpha\mathfrak{J}^\mu w(x) \alpha\mathfrak{J}^\mu u(x) + \alpha\mathfrak{J}^\mu v(x) \alpha\mathfrak{J}^\mu w(x) - \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} \alpha\mathfrak{J}^\mu vw(x). \end{aligned}$$

*Proof.* Define

$$H(\tau, \rho) = [f(\tau) - f(\rho)][g(\tau) - g(\rho)], \quad \tau, \rho \in (0, x), \quad x > 0. \quad (22)$$

Multiplying both sides of (22) by  $\frac{1}{\Gamma^2(\alpha)} \left(\frac{x^\mu - \tau^\mu}{\mu}\right)^{\alpha-1} \left(\frac{x^\mu - \rho^\mu}{\mu}\right)^{\alpha-1} \tau^{\mu-1} \rho^{\mu-1}$  for  $\tau, \rho \in (0, x)$  and integrating the desired inequality with respect to  $\tau, \rho$  from 0 to  $x$  yield

$$\begin{aligned} \frac{1}{2\Gamma^2(\alpha)} \int_0^t \int_0^t \left(\frac{x^\mu - \tau^\mu}{\mu}\right)^{\alpha-1} \left(\frac{x^\mu - \rho^\mu}{\mu}\right)^{\alpha-1} \tau^{\mu-1} \rho^{\mu-1} H(\tau, \rho) d\tau d\rho \\ = \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} \alpha\mathfrak{J}^\mu fg(x) + \alpha\mathfrak{J}^\mu f(x) \alpha\mathfrak{J}^\mu g(x). \end{aligned} \quad (23)$$

Applying the Cauchy-Schwartz inequality to (23) leads to

$$\begin{aligned} \left[ \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} \alpha\mathfrak{J}^\mu fg(x) + \alpha\mathfrak{J}^\mu f(x) \alpha\mathfrak{J}^\mu g(x) \right]^2 \\ \leq \left[ \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} \alpha\mathfrak{J}^\mu f^2(x) - [\alpha\mathfrak{J}^\mu f(x)]^2 \right] \left[ \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} \alpha\mathfrak{J}^\mu g^2(x) - [\alpha\mathfrak{J}^\mu g(x)]^2 \right]. \end{aligned} \quad (24)$$

Since  $[\phi_2(x) - f(x)][f(x) - \phi_1(x)] \geq 0$  and  $[\psi_2(x) - g(x)][g(x) - \psi_1(x)] \geq 0$  for  $x \in [0, \infty)$ , we have

$$\frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} [\phi_2(x) - f(x)][f(x) - \phi_1(x)] \geq 0$$

and

$$\frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} [\psi_2(x) - g(x)][g(x) - \psi_1(x)] \geq 0.$$

Thus, from Theorem 2.3, we obtain

$$\begin{aligned} \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} \alpha\mathfrak{J}^\mu f^2(x) - [\alpha\mathfrak{J}^\mu f(x)]^2 &\leq [\alpha\mathfrak{J}^\mu \phi_2(x) - \alpha\mathfrak{J}^\mu f(x)][\alpha\mathfrak{J}^\mu f(x) \\ &- \alpha\mathfrak{J}^\mu \phi_1(x)] + \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} \alpha\mathfrak{J}^\mu \phi_1 f(x) - \alpha\mathfrak{J}^\mu \phi_1(x) \alpha\mathfrak{J}^\mu f(x) \end{aligned}$$



$$\begin{aligned}
& + \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} {}^\alpha\mathcal{J}^\mu\phi_2 f(x) - {}^\alpha\mathcal{J}^\mu\phi_2(x) {}^\alpha\mathcal{J}^\mu f(x) \\
& + {}^\alpha\mathcal{J}^\mu\phi_1(x) {}^\alpha\mathcal{J}^\mu\phi_2(x) - \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} {}^\alpha\mathcal{J}^\mu\phi_1\phi_2(x) = T(f, \phi_1, \phi_2). \quad (25)
\end{aligned}$$

Similarly, we have

$$\begin{aligned}
& \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} {}^\alpha\mathcal{J}^\mu g^2(x) - [{}^\alpha\mathcal{J}^\mu g(x)]^2 \leq [{}^\alpha\mathcal{J}^\mu\psi_2(x) - {}^\alpha\mathcal{J}^\mu g(x)][{}^\alpha\mathcal{J}^\mu g(x) - {}^\alpha\mathcal{J}^\mu\psi_1(x)] \\
& + \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} {}^\alpha\mathcal{J}^\mu\psi_1 g(x) - {}^\alpha\mathcal{J}^\mu\psi_1(x) {}^\alpha\mathcal{J}^\mu g(x) + \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} {}^\alpha\mathcal{J}^\mu\psi_2 g(x) \\
& - {}^\alpha\mathcal{J}^\mu\psi_2(x) {}^\alpha\mathcal{J}^\mu g(x) + {}^\alpha\mathcal{J}^\mu\psi_1(x) {}^\alpha\mathcal{J}^\mu\psi_2(x) - \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} {}^\alpha\mathcal{J}^\mu\psi_1\psi_2(x) = T(g, \psi_1, \psi_2). \quad (26)
\end{aligned}$$

Combining (24), (25), and (26), we obtain the desired inequality (21).  $\square$

*Remark 2.1.* For  $m, M, n, N \in \mathbb{R}$ , if  $T(f, \phi_1, \phi_2) = T(g, m, M)$  and  $T(g, \psi_1, \psi_2) = T(g, n, N)$ , then the inequality (21) reduces to

$$\left| \frac{x^{\mu\alpha}}{\mu^\alpha\Gamma(\alpha+1)} {}^\alpha\mathcal{J}^\mu f g(x) - {}^\alpha\mathcal{J}^\mu f(x) {}^\alpha\mathcal{J}^\mu g(x) \right| \leq \left[ \frac{x^{\mu\alpha}}{2\mu^\alpha\Gamma(\alpha+1)} \right]^2 (M-m)(N-n).$$

in [18, Theorem 1].

*Remark 2.2.* In this paper, we presented some new conformable fractional integral inequalities which generalize those corresponding ones in [21].

#### REFERENCES

- [1] G. A. Anastassiou, *Opial type inequalities involving fractional derivatives of two functions and applications*, *Comput. Math. Appl.* **48** (2004), no. 10-11, 1701–1731; Available online at <https://doi.org/10.1016/j.camwa.2003.08.013>.
- [2] S. Belarbi and Z. Dahmani, *On some new fractional integral inequalities*, *J. Inequal. Pure Appl. Math.* **10** (2009), no. 3, Article 86, 5 pp. Available online at <https://www.emis.de/journals/JIPAM/article1142.html>.
- [3] Z. Dahmani, *New inequalities in fractional integrals*, *Int. J. Nonlinear Sci.* **9** (2010), no. 3, 493–497.
- [4] Z. Dahmani, L. Tabharit, and S. Taf, *New generalization of Grüss inequality using Riemann-Liouville fractional integrals*, *Bull. Math. Anal. Appl.* **2** (2010), no. 3, 93–99.
- [5] Z. Denton and A. S. Vatsala, *Fractional integral inequalities and applications*, *Comput. Math. Appl.* **59** (2010), no. 3, 1087–1094; Available online at <https://doi.org/10.1016/j.camwa.2009.05.012>.
- [6] R. Gorenflo and F. Mainardi, *Fractional Calculus: Integral and Differential Equations of Fractional order*, *Fractals and Fractional Calculus in Continuum Mechanics* (Udine, 1996), 223–276, CISM Courses and Lectures, 378, Springer, Vienna, 1997.
- [7] G. Grüss, *Über das Maximum des absoluten Betrages von  $\frac{1}{b-a} \int_a^b f(x)g(x) dx - \frac{1}{(b-a)^2} \int_a^b f(x) dx \int_a^b g(x) dx$* , *Math. Z.* **39** (1935), no. 1, 215–226; Available online at <https://doi.org/10.1007/BF01201355>. (German)
- [8] F. Jarad, E. Uğurlu, T. Abdeljawad, and D. Baleanu, *On a new class of fractional operators*, *Adv. Difference Equ.* **2017**, Paper No. 247, 16 pp. Available online at <https://doi.org/10.1186/s13662-017-1306-z>.
- [9] A. A. Kilbas, *Hadamard-type fractional calculus*, *J. Korean Math. Soc.* **38** (2001), no. 6, 1191–1204.
- [10] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, North-Holland Mathematics Studies, 204. Elsevier Science B.V., Amsterdam, 2006.

- [11] D. S. Mitrinović, J. E. Pečarić, and A. M. Fink, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, 1993; Available online at <http://dx.doi.org/10.1007/978-94-017-1043-5>.
- [12] K. S. Nisar, F. Qi, G. Rahman, S. Mubeen, and M. Arshad, *Some inequalities involving the extended gamma function and the Kummer confluent hypergeometric k-function*, J. Inequal. Appl. (2018), **2018**:135, 12 pages; Available online at <https://doi.org/10.1186/s13660-018-1717-8>.
- [13] F. Qi and B.-N. Guo, *Integral representations and complete monotonicity of remainders of the Binet and Stirling formulas for the gamma function*, Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Math. RACSAM **111** (2017), no. 2, 425–434; Available online at <https://doi.org/10.1007/s13398-016-0302-6>.
- [14] S. G. Samko, A. A. Kilbas, and O. I. Marichev, *Fractional Integrals and Derivatives: Theory and Applications*, Edited and with a foreword by S. M. Nikol'skiĭ. Translated from the 1987 Russian original. Revised by the authors. Gordon and Breach Science Publishers, Yverdon, 1993.
- [15] M. Z. Sarikaya and H. Ogunmez, *On new inequalities via Riemann-liouville fractional integrations*, Abstr. Appl. Anal. **2012**, Art. ID 428983, 10 pp. Available online at <http://dx.doi.org/10.1155/2012/428983>.
- [16] E. Set, İ. İşcan, M. Z. Sarikaya, and M. E. Özdemir, *On new inequalities of Hermite-Hadamard-Fejér type for convex functions via fractional integrals*, Appl. Math. Comput. **259** (2015), 875–881; Available online at <https://doi.org/10.1016/j.amc.2015.03.030>.
- [17] E. Set, İ. Mumcu, and S. Demirbaş, *Chebyshev type inequalities involving new conformable fractional integral operators*, ResearchGate Preprint (2018), available online at <https://www.researchgate.net/publication/323880498>.
- [18] E. Set, İ. Mumcu, and M. E. Özdemir, *Grüss type inequalities involving new conformable fractional integral operators*, ResearchGate Preprint (2018), available online at <https://www.researchgate.net/publication/323545750>.
- [19] H. M. Srivastava and J. Choi, *Zeta and q-Zeta Functions and Associated Series and Integrals*, Elsevier, Inc., Amsterdam, 2012; Available online at <http://dx.doi.org/10.1016/B978-0-12-385218-2.00001-3>.
- [20] W. T. Sulaiman, *Some new fractional integral inequalities*, J. Math. Anal. **2** (2011), no. 2, 23–28.
- [21] J. Tariboon, S. K. Ntouyas, and W. Sudsutad, *Some new Riemann-Liouville fractional integral inequalities*, Int. J. Math. Math. Sci. **2014**, Art. ID 869434, 6 pp. Available online at <https://doi.org/10.1155/2014/869434>.
- [22] C. Zhu, W. Yang, and Q. Zhao, *Some new fractional q-integral Grüss-type inequalities and other inequalities*, J. Inequal. Appl. **2012**, 2012:299, 15 pp. Available online at <https://doi.org/10.1186/1029-242X-2012-299>.

(Rahman) DEPARTMENT OF MATHEMATICS, INTERNATIONAL ISLAMIC UNIVERSITY, ISLAMABAD, PAKISTAN

*Email address:* [gauhar55uom@gmail.com](mailto:gauhar55uom@gmail.com)

(Nisar) DEPARTMENT OF MATHEMATICS, COLLEGE OF ARTS AND SCIENCE AT WADI ALDAWASER, 11991, PRINCE SATTAM BIN ABDULAZIZ UNIVERSITY, ALKHARJ, KINGDOM OF SAUDI ARABIA

*Email address:* [ksnisar1@gmail.com](mailto:ksnisar1@gmail.com), [n.soopy@psau.edu.sa](mailto:n.soopy@psau.edu.sa)

(Qi) INSTITUTE OF MATHEMATICS, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO, HENAN, 454010, CHINA; DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCE, TIANJIN POLYTECHNIC UNIVERSITY, TIANJIN, 300387, CHINA

*Email address:* [qifeng618@gmail.com](mailto:qifeng618@gmail.com), [qifeng618@hotmail.com](mailto:qifeng618@hotmail.com), [qifeng618@qq.com](mailto:qifeng618@qq.com)

*URL:* <http://qifeng618.wordpress.com>