



# KEY POLYNOMIALS, SEPARATE AND IMMEDIATE VALUATIONS, AND SIMPLE EXTENSIONS OF VALUED FIELDS

Gérard Leloup

## ► To cite this version:

Gérard Leloup. KEY POLYNOMIALS, SEPARATE AND IMMEDIATE VALUATIONS, AND SIMPLE EXTENSIONS OF VALUED FIELDS. 2019. hal-01876056v2

**HAL Id: hal-01876056**

**<https://hal.science/hal-01876056v2>**

Preprint submitted on 18 Dec 2019 (v2), last revised 16 May 2022 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# KEY POLYNOMIALS, SEPARATE AND IMMEDIATE VALUATIONS, AND SIMPLE EXTENSIONS OF VALUED FIELDS.

G. LELOUP

**ABSTRACT.** In order to study simple extensions of valued fields, notions of key polynomials were developed. Model theoretical properties of extensions of valued fields were also studied. The properties of valuations used in model theory shed a new light on key polynomials and they make it possible to obtain underlying properties of these extensions. Key polynomials are used for defining separate valuations which approximate a valuation on an extension  $K(X)$ . A valuation  $\nu_\lambda$  on  $K(X)$  is separate if there is a  $K$ -basis  $\mathcal{B}_\lambda$  of  $K[X]$  such that  $\nu_\lambda$  is determined by its restrictions to  $K$  and  $\mathcal{B}_\lambda$ . For every valuation  $\nu$  the aim is to find a family of monic polynomials of  $K[X]$ , which are called key polynomials, and a family  $\nu_\lambda$  of separate valuations such that for every  $\lambda$  the elements of  $\mathcal{B}_\lambda$  are products of key polynomials, and, for every  $f \in K[X]$ ,  $\nu(f)$  is the maximum of the family  $(\nu_\lambda(f))$ . The approach of the present paper shows the links between some properties of valuations used in model theory and properties of key polynomials. Our definitions rely on euclidean division of polynomials, on bases of vector spaces and on classical properties of valuations.

<sup>1</sup> The aim of this work is twofold. On the one hand, it is to show the links between the model theoretical study of extensions of valued fields of W. Baur, F. Delon and the author (see [B 81], [B 82], [D 82], [D 88], [D 91], [L 89], [L 03]) and the study of simple extensions of valued fields of S. MacLane ([ML 36a], [ML 36b]), M. Vaquié ([V 07]), J. Decaup, F. J. Herrera Govantes, W. Mahloud, J. Novakoski, M. A. Olalla Acosta, M. Spivakovsky ([HOS 07], [HMOS 14], [DMS 18], [NS 18], [N 19]), and others. The purposes of these last authors are related to algebraic geometry. On the other hand, it is to give a different approach to the study of key polynomials. In above papers the families of key polynomials are constructed by induction on the degrees. Then two cases arise: the set  $\nu(X^d - K_{d-1}[X])$  having or not a greatest element (where  $K_{d-1}[X]$  is the  $K$ -module of polynomials of degree at most  $d - 1$ ). We will show that in the first case we have a separate subextension, and in the last one this subextension is immediate. Separate and immediate extensions play a central role in the model theoretical study of extensions of valued fields. So we found interesting to revisit key polynomials in the light of another background. In particular, we define key polynomials by first order formulas (in the language of valued rings together with a predicate interpreted by  $X$ ), and we do not require to define families of key polynomials by induction. The approach of the present paper relies on euclidean division of polynomials, basis of vector spaces and the notions of separate and immediate extensions used in model theoretical study of extensions of valuations. In the following, we use the word “module” which is shorter than “vector space”. We do not require the reader having any previous knowledge of key polynomials. As much as possible, we try to use only elementary properties of valuations. However, we use the graded algebra associated to a valuation in two proofs and, in the last section, an approximation theorem of Ribenboim in one proof, and pseudo-Cauchy sequences in the study of immediate extensions (all the definitions will be given below). On the one side, graded algebras are a good tool for providing nice proofs of some properties of separate extensions. On the other side, pseudo-Cauchy sequences play an important role in the study of immediate extensions. This work is self-contained and when we use results on key polynomials of preceding papers, we generally give a proof. This work started before the publication of the papers [DMS 18], [NS 18], [N 19], so the first version (online on September 2018) did not include their results. In the present version we had to change some definitions and make the links with the results of those papers. We will show that our definition of key polynomials is equivalent to the definition of these last papers.

Let  $(K, \nu)$  be a valued field. S. MacLane introduced key polynomials to define families of separate valuations which approximate an extension of  $\nu$  to the field  $K(X)$ , where  $X$  is algebraic or transcendental over  $K$ . They are also used for defining the different extensions of a fixed valuation to a given simple algebraic extension of a field. Here we will focus on the first purpose. In the works of S. MacLane and of M. Vaquié, the key polynomials  $\Phi$  are constructed by induction, starting from the degree 1 key polynomials. In [HOS 07] the authors do not define key polynomials but families of key polynomials, by means of properties that these families have to satisfy. Then they construct such families by induction, starting from the degree 1, in a different way from the constructions of S. MacLane and M. Vaquié. A definition

---

*Date:* December 11, 2019.

<sup>1</sup>2010 *Mathematics Subject Classification.* 12J10, 12J20, 12F99.

*Keywords:* simple extension, valuation, key polynomial.

of key polynomials appears in [NS 18], [DMS 18] and [N 19] in the case of a transcendental extension (in [DMS 18] they are called abstract key polynomials). Here, we do not restrict to transcendental extensions and we give a characterization of key polynomials by means of first order formulas (in the language of valued fields together with a predicate interpreted by the generator of the simple extension) which are satisfied by the abstract key polynomials. A slight modification in our definition gives rise to the definition of polynomials that we will call weak key polynomials (in short w-key polynomials). The key polynomials are also w-key polynomials. In Proposition 3.24 we show that  $\Phi$  being a w-key polynomial for  $\nu$  is equivalent to being a MacLane key polynomial for the generalized Gauss valuation  $\nu_\Phi$  which we will define below.

### General properties.

A *valuation* on a field  $K$  is a morphism  $\nu$  from the multiplicative group  $(K^*, \cdot)$  to an abelian linearly ordered group  $(\nu K, +)$  called the *valuation group*. We add an element  $\infty$  to  $\nu K$  ( $\infty > \nu K$ ), we set  $\nu(0) := \infty$ , and we assume that  $\nu(x + y) \geq \min(\nu(x), \nu(y))$ , for every  $x, y$  in  $K$ . It follows that if  $\nu(x) \neq \nu(y)$ , then  $\nu(x + y) = \min(\nu(x), \nu(y))$ . If  $\nu(x) = \nu(y)$ , then  $\nu(x + y)$  can be arbitrarily large. The set  $\{x \in K \mid \nu(x) \geq 0\}$  is a local domain and  $\{x \in K \mid \nu(x) > 0\}$  is its maximal ideal, they are called respectively the *valuation ring* and the *maximal ideal* of the valued field. The quotient of the valuation ring by the maximal ideal is a field which is denoted by  $K_\nu$ , and is called the *residue field* of the valued field. The *residue characteristic* of  $(K, \nu)$  is the characteristic of  $K_\nu$ . We have  $\text{char } K_\nu = 0 \Rightarrow \text{char } K = 0$  and  $\text{char } K > 0 \Rightarrow \text{char } K_\nu = \text{char } K$ . For  $x$  in the valuation ring of  $(K, \nu)$ , its class modulo the maximal ideal will be denoted by  $x_\nu$ . For every subset  $M$  of  $K$  we denote by  $\nu(M)$ , or  $\nu M$ , the set  $\{\nu(x) \mid x \in M, x \neq 0\}$ .

We assume that  $\nu$  is a finite rank valuation (the *rank* of  $\nu$  is the number of proper convex subgroups of  $\nu K$ ), so that  $\nu K$  has countable cofinality. If the rank is 1, then  $\nu K$  embeds in the ordered group  $(\mathbb{R}, +)$ ; we also say that  $\nu$  is *archimedean*.

An extension  $(L|K, \nu)$  of valued fields consists in an extension  $L|K$  of fields, where  $L$  is equipped with a valuation  $\nu$ . If  $L|K$  is a field extension and  $\nu$  is a valuation on  $K$ , then there is a least one extension of  $\nu$  to  $L$ .

Assume that  $M$  is a  $K$ -module (where  $K$  is a field). A mapping  $\nu$  from  $M$  to a linearly ordered group together with an element  $\infty$  will be called a  *$K$ -module valuation*, if for every  $y_1, y_2$  in  $M$  and  $x \in K$ :  $\nu(y_1) = \infty \Leftrightarrow y_1 = 0$ ,  $\nu(y_1 + y_2) \geq \min(\nu(y_1), \nu(y_2))$ ,  $\nu(xy_1) = \nu(x) + \nu(y_1)$  (see. [FVK]). It follows that its restriction to  $K$  is a valuation of field.

### Separate and immediate extensions.

For  $n \in \mathbb{N}^*$ , we denote by  $K_n[\chi]$  the  $K$ -submodule of  $K[\chi]$  of all polynomials of degree at most  $n$ , where  $\chi$  is algebraic or transcendental over  $K$ . We let  $K_0[\chi] := K$  and if  $\chi$  is transcendental over  $K$ , then we let  $K_\infty[\chi] := K[\chi]$ . In the classification of key polynomials, and in the construction of families of key polynomials will appear the sets  $\nu(\chi^n - K_{n-1}[\chi])$ . We will show that these sets are related to characterizations of separate and immediate extensions.

Assume that  $\nu$  is a  $K$ -module valuation on  $M$ . In general, for every  $x_1, \dots, x_n$  in  $K$ , pairwise distinct  $y_1, \dots, y_n$  in  $M$ ,  $\nu(x_1 y_1 + \dots + x_n y_n)$  is at least equal to  $\min(\nu(x_1 y_1), \dots, \nu(x_n y_n))$ , and it can be arbitrarily large. The  $K$ -module valuation  $\nu$  is said to be *separate* if there exists a basis  $\mathcal{B}$  of  $M$  such that for every  $x_1, \dots, x_n$  in  $K$ , pairwise distinct  $y_1, \dots, y_n$  in  $\mathcal{B}$ ,  $\nu(x_1 y_1 + \dots + x_n y_n) = \min(\nu(x_1 y_1), \dots, \nu(x_n y_n))$ . If this holds, then the basis  $\mathcal{B}$  is said to be  *$\nu$ -separate* (in short *separate*). Note that if  $\mathcal{B}$  is a basis of  $L$ , then we can define a separate  $K$ -module valuation  $\nu'$  by setting, for every  $x_1, \dots, x_n$  in  $K$ , pairwise distinct  $y_1, \dots, y_n$  in  $\mathcal{B}$ ,  $\nu'(x_1 y_1 + \dots + x_n y_n) = \min(\nu(x_1 y_1), \dots, \nu(x_n y_n))$ . If  $\nu$  is separate, then one can compute the valuation of any element of  $M$  by means of the restrictions of  $\nu$  to  $K$  and  $\mathcal{B}$ .

We will see in Proposition 2.19 that for  $d \in \mathbb{N} \cup \{\infty\}$  such that  $0 < d \leq [K[\chi]:K] - 1$ , the extension  $(K_d[\chi]|K, \nu)$  is separate if, and only if, for every integer  $n$ ,  $1 \leq n \leq d$ ,  $\nu(\chi^n - K_{n-1}[\chi]) = \{\nu(\chi^n - y) \mid y \in K_{n-1}[\chi]\}$  has a maximal element. Furthermore, if  $\chi$  is transcendental over  $K$ , and  $\nu$  is a field valuation on  $K(\chi)$ , then  $(K(\chi)|K, \nu)$  is separate if, and only if, for every  $n \in \mathbb{N}^*$ ,  $\nu(\chi^n - K_{n-1}[\chi])$  has a maximal element.

Let  $M \subset N$  be  $K$ -submodules of  $L$ . We say that the  $K$ -module  $N$  is *immediate* over  $M$  if for every  $l \in N$ , the set  $\nu(l - M) = \{\nu(l - x) \mid x \in M \setminus \{l\}\}$  is a subset of  $\nu M$  and it has no maximal element. We say that it is *dense* over  $M$  if for every  $l \in N$ , the set  $\nu(l - M)$  is equal to  $\nu M$ . One can prove that saying that the extension of valued field  $(L|K, \nu)$  is immediate is equivalent to saying that  $\nu L = \nu K$  and  $L_\nu = K_\nu$ . We will show in Proposition 1.8 that for  $d, d'$  in  $\mathbb{N} \cup \{\infty\}$  such that  $0 \leq d < d' \leq [K[\chi]:K] - 1$ , (that is, if  $\chi$  is algebraic over  $K$ , then  $d' < [K[\chi]:K]$ ), the extension  $(K_{d'}[\chi]|K_d[\chi], \nu)$  is immediate if, and only if, for every  $n \in \{d+1, \dots, d'\}$ ,  $\nu(\chi^n - K_{n-1}[\chi])$  has no maximal element. Furthermore, if  $\nu$  is a field valuation and  $\nu K[\chi]$  is a group, then it is dense if, and only if, for every  $n \in \{d+1, \dots, d'\}$ ,  $\nu(\chi^n - K_{n-1}[\chi]) = \nu K$ . Propositions 1.8 and 2.19, show that the separate and the immediate cases are in some sense opposite ones. This difference will also be illustrated by Theorem 2.7, which implies that

a valued fields extension  $(L|K, \nu)$  is separate if, and only if, for every finitely generated  $K$ -submodule  $M$  of  $L$  and  $l \in L \setminus M$ , the set  $\nu(l - M)$  has a maximal element, and the definition of an immediate extension. Furthermore, Theorem 2.8 implies that if  $(L|K, \nu)$  is a finite algebraic extension of valued fields, then  $(L|K, \nu)$  is separate if, and only if,  $[L:K] = [L_\nu:K_\nu](\nu L:\nu K)$ , although Remark 1.6, implies that  $(L|K, \nu)$  is immediate if, and only if,  $\nu L = \nu K$  and  $L_\nu = K_\nu$ .

### Families of key polynomials.

Now, let  $(K(\chi)|K, \nu)$  be a simple extension of valued fields. If there is no confusion, then for every polynomial  $f(\chi)$  of  $K[\chi]$  we write  $f$  instead of  $f(\chi)$ . S. MacLane defined families  $(\nu_i)$  of separate  $K$ -module valuations to approximate  $\nu$  (where  $i$  runs over a well-ordered set). Note that if  $\mathcal{B}_i$  is a separate basis for  $\nu_i$ , and the restrictions of  $\nu$  and  $\nu_i$  to  $K$  and  $\mathcal{B}_i$  are equal, then for every  $f \in K[\chi]$  we have  $\nu_i(f) \leq \nu(f)$  (in short,  $\nu_i \leq \nu$ ). Assume that  $\chi$  is algebraic (so that  $K(\chi) = K[\chi]$ ), and that there exists  $f \in K[\chi]$  such that  $\nu_i(f) < \nu(f)$ . Since  $\nu_i \leq \nu$ , we have  $\nu_i(1/f) \leq \nu(1/f)$ . Hence  $\nu_i(f) < \nu(f) = -\nu(1/f) \leq -\nu_i(1/f)$ . So  $\nu_i(1/f) \neq -\nu_i(f)$ . It follows that  $\nu_i$  doesn't satisfy the rule  $\nu_i(fg) = \nu_i(f) + \nu_i(g)$ . Now, we will prove in Proposition 3.16 that, in some cases, if  $\deg(f) + \deg(g) < [K[\chi]:K]$ , then  $\nu_i(fg) = \nu_i(f) + \nu_i(g)$ . This motivates the following definitions.

If for every  $f, g$  in  $K[\chi]$  such that  $\deg(f) + \deg(g) < [K[\chi]:K]$  we have  $\nu(fg) = \nu(f) + \nu(g)$ , then we say that  $\nu$  is *partially multiplicative*, or a *p-m valuation*. In the case where  $[K[\chi]:K] = \infty$ , we assume in addition that  $g \neq 0$  implies  $\nu(f/g) = \nu(f) - \nu(g)$ , that is,  $\nu$  is a field valuation in the usual sense. If  $\nu$  is a field valuation (in the usual sense), then we will also say that it is *multiplicative*.

Let  $\Phi$  be a monic polynomial of degree  $d$ . We say that  $\Phi$  is a *key polynomial* if for every  $f, g$  in  $K_{d-1}[\chi]$  with  $\deg(f) + \deg(g) < [K[\chi]:K]$  we have  $\nu(fg) = \nu(r) < \nu(q\Phi)$ , where  $fg = q\Phi + r$  is the euclidean division. A positive integer  $d$  is said to be a *key degree* if there is a key polynomial of degree  $d$ . Trivially, 1 is a key degree, and we see that any key polynomial is irreducible. Let  $d_1 := 1 < d_2 < \dots < d_n$  be the first  $n$  key degrees, and  $d_{n+1} := [K(\chi):K]$ . For  $i \leq n$ , let  $\Phi_{d_i}$  be a key polynomial of degree  $d_i$ . We let  $\mathcal{B}$  be the family of the  $\Phi_{d_1}^{e_1} \dots \Phi_{d_n}^{e_n}$ , where, for  $1 \leq j \leq n$ ,  $e_1 + e_2 d_2 + \dots + e_j d_j < d_{j+1}$ . Since the degree mapping is one-to-one from  $\mathcal{B}$  onto  $[0, [K(\chi):K]]$ ,  $\mathcal{B}$  is indeed a basis of the  $K$ -module  $K[\chi]$ . We let  $\nu_{\Phi_{d_1}, \dots, \Phi_{d_n}}$  be the separate  $K$ -module valuation defined by this basis. Then, for every  $f \in K[\chi]$ , all the  $\nu_{\Phi_{d_1}, \dots, \Phi_{d_n}}(f)$  are bounded above by  $\nu(f)$ . We prove that there exists a family  $\mathcal{F}$  of key polynomials such that, for every  $f \in K[\chi]$ , by letting  $d_n$  be the greatest key degree which is at most equal to  $\deg(f)$ , there exist  $\Phi_{d_1}, \dots, \Phi_{d_n}$  in  $\mathcal{F}$  such that  $\nu_{\Phi_{d_1}, \dots, \Phi_{d_n}}(f) = \nu(f)$ .

The family  $\mathcal{F}$  is defined in the following way. Let  $d$  be a key degree. We say that  $d$  is a *separate* key degree if the set  $\nu(\chi^d - K_{d-1}[\chi])$  has a maximal element. We will show in Proposition 3.27 that every monic polynomial  $\Phi_d$  of degree  $d$  such that  $\nu(\Phi_d) = \max \nu(\chi^d - K_{d-1}[\chi])$  is a key polynomial. We say that  $d$  is an *immediate* key degree if the set  $\nu(\chi^d - K_{d-1}[\chi])$  has no maximal element. We will show in Proposition 3.27 that there exists a sequence  $(\Phi_{d,n})$  of key polynomials of degree  $d$  such that the sequence  $(\nu(\Phi_{d,n}))$  is increasing and cofinal in  $\nu(\chi^d - K_{d-1}[\chi])$ .

Since key polynomials was first constructed by induction on the degree, some authors introduced the definition of limit key polynomial. In fact, a key polynomial is a limit key polynomial if, and only if, the preceding key degree is an immediate one. Here we prefer make a distinction between the properties of the key polynomials and of their degrees, instead of the previous key degree. So we make a distinction between immediate and separate key degrees, which seems more relevant.

Now, let  $1 := d_1 < d_2 < \dots$  be the sequence of key degrees. If  $d_i$  is a separate key degree, then we let  $\Phi_{d_i}$  be a key polynomial such that  $\nu(\Phi_{d_i}) = \max \nu(\chi^{d_i} - K_{d_i-1}[\chi])$ . For notational convenience, for every non-negative integer  $n$  we set  $\Phi_{d_i,n} := \Phi_{d_i}$ . If  $d_i$  is an immediate key degree, then we let  $(\Phi_{d_i,n})$  be a sequence of key polynomials of degree  $d_i$  such that the sequence  $(\nu(\Phi_{d_i,n}))$  is increasing and cofinal in  $\nu(\chi^{d_i} - K_{d_i-1}[\chi])$ . Then we can let  $\mathcal{F}$  be the family of the  $\nu_{\Phi_{d_1,n_1}, \dots, \Phi_{d_k,n_k}}$ 's, where  $k$  and the  $n_i$ 's run over  $\mathbb{N}$ . In Theorem 3.44 we prove that if the degree of  $f$  is less than  $d_{k+1}$ , then  $\nu(f)$  is the maximum of the family  $\nu_{\Phi_{d_1,n_1}, \dots, \Phi_{d_k,n_k}}(f)$ , and infinitely many  $\nu_{\Phi_{d_1,n_1}, \dots, \Phi_{d_k,n_k}}(f)$ 's are equal to  $\nu(f)$ .

If there is at most one immediate key degree, then we get an algorithm to calculate  $\nu(f)$  for every polynomial  $f$ . Let  $d_k$  be the greatest key degree which is at most equal to the degree of  $f$ . Then  $\nu_{\Phi_{d_1,n}, \dots, \Phi_{d_k,n}}(f) = \nu(f) \Leftrightarrow \nu_{\Phi_{d_1,n}, \dots, \Phi_{d_k,n}}(f) = \nu_{\Phi_{d_1,n+1}, \dots, \Phi_{d_k,n+1}}(f)$ . Hence we compute  $\nu_{\Phi_{d_1,n}, \dots, \Phi_{d_k,n}}(f)$ , and we stop when this condition holds.

Now we set a new definition. If we have  $\nu(r) \leq \nu(q\Phi)$  instead of  $\nu(r) < \nu(q\Phi)$  in the definition of key polynomials, then we say that  $\Phi$  is a *weak key polynomial* (in short a *w-key polynomial*). The w-key polynomials can also be characterized in the following way. Let  $\Phi$  be a monic irreducible polynomial and  $d$  be the degree of  $\Phi$ . Assume that  $d \leq [K[\chi]:K]/2$ . Since the  $K$ -modules  $K_{d-1}[\chi]$ ,  $K_{d-1}[\chi]$  and  $K[\chi]/(\Phi)$  are isomorphic (where  $K[\chi]$  denotes the ring of formal polynomials),  $\nu$  induces a  $K$ -module valuation  $\bar{\nu}$  on  $K[\chi]/(\Phi)$  by setting, for  $f(X) \in K_{d-1}[\chi]$ ,  $\bar{\nu}(f(X) + (\Phi)) = \nu(f(X))$ . Note that  $\bar{\nu}(K[\chi]/(\Phi)) = \nu(K_{d-1}[\chi])$ . Now, we will prove in Proposition 3.7 that  $\Phi$  is a w-key polynomial if, and only if,  $(K[\chi]/(\Phi), \bar{\nu})$  is a valued field (i.e.  $\bar{\nu}$  is multiplicative). If  $\Phi$  is a key polynomial, then in addition

the residue field  $(K[X]/(\Phi))_{\bar{\nu}}$  is canonically isomorphic to  $(K_{d-1}[\chi])_{\nu}$  (so, it embeds in  $(K[\chi])_{\nu}$ ).

### Some model theory.

In above study, we see that the simpler case is the separate one. In the model theoretical study of extensions of valued fields, separate extensions also give interesting results. A valued field  $(K, \nu)$  is said to be *algebraically maximal* if no extension of  $\nu$  to an algebraic extension of  $K$  is immediate. If the residue characteristic is 0, then being algebraically maximal is equivalent to being *henselian*, i.e.  $\nu$  having a unique extension to any algebraic extension of  $K$ . A famous theorem of J. Ax, S. Kochen and Y. Ershov ([AK 65]) says that the elementary theory of a henselian valued field  $(K, \nu)$  of residue characteristic 0 is determined by the elementary theory of its residue field and the elementary theory of its value group. Next, this result was extended to other families of algebraically maximal valued fields. In the model theoretical study of extensions of valued fields, we try to get similar results. Now, in [D 91] F. Delon proved that given a theory  $T_F$  of fields of characteristic 0 and a theory  $T_V$  of non trivial linearly ordered abelian groups, the theory of immediate henselian extensions  $(L|K, \nu)$ , where  $K_{\nu}$  is a model of  $T_F$  and  $\nu K$  is a model of  $T_V$ , is undecidable and admits  $2^{\aleph_0}$  completions. The failure comes from the sets  $\nu(l - K)$ , where  $l \in L$ . Now, if  $K, L$  are henselian,  $\text{char} K_{\nu} = 0$ , and  $(L|K, \nu)$  is separate, then the first-order theory of the extension is determined by the theories of the residual extension and of the extension of valued groups. The same holds if  $(L|K, \nu)$  is an extension of algebraically maximal Kaplansky fields or of real-closed fields (see [B 81], [B 82], [L 89], [L 03]). Furthermore, we have similar results with dense extensions, and some extensions can be decomposed into separate and dense subextensions. In the case where  $(L|K, \nu)$  is an extension of valued fields of residue characteristic 0, there exists a henselian subfield  $H$ ,  $K \subseteq H \subseteq L$ , such that  $(H|K, \nu)$  is separate and  $(L|H, \nu)$  is immediate (see [D 88]). This shows that it can be interesting to focus on separate and immediate extensions.

### Summary of the paper.

In Section 1 we generalize the definitions of immediate and dense extensions to extensions of modules, equipped with  $K$ -module valuations. Then we focus on the case of simple extensions. We prove Proposition 1.8, which characterizes immediate extensions by the condition that the sets  $\nu(\chi^n - K_{n-1}[\chi])$  have no greatest elements. Section 2 is devoted to separate extensions. In the same way as in Section 1, we focus on the case of simple extensions, and we prove Proposition 2.19, which characterizes separate extensions by the condition that the sets  $\nu(\chi^n - K_{n-1}[\chi])$  have greatest elements. Then we recall definitions and properties of the graded algebras associated to valuations, since they are useful in the study of separate extensions. Furthermore, we prove that a family is separate if, and only if, its image in the graded algebra is linearly independent (Fact 2.20). In Section 3, we characterize w-key polynomials, key polynomials and key degrees. We compare the definition of w-key polynomials with the definition of S. MacLane and M. Vaquié (Proposition 3.24). We study the  $K$ -module valuations defined by w-key polynomials, and the associated bases generated by these w-key polynomials. Next we set properties of key degrees. We prove, for example, that if  $d$  is an immediate key degree and  $d'$  is the next key degree, then the extension  $(K_{d'-1}[\chi]|K_{d-1}[\chi], \nu)$  is immediate (Theorem 3.36). We also give characterizations of the successor of a given key degree (Theorems 3.36 and 3.37). In particular, we look at conditions for being the greatest key degree. For example, we show that this holds if  $\nu(\chi^d - K_{d-1}[\chi]) = \nu K_{d-1}[\chi] = \nu K(\chi)$  (Proposition 3.41). We use w-key polynomials to define the  $K$ -module valuations which approximate a given valuation of  $K(\chi)|K$ . In Subsection 3.7 we show that the key polynomials defined in the transcendental case in [NS 18], [DMS 18] and [N 19] are the same as the key polynomials defined here (Corollaries 3.50 and 3.52). Next, we look at the links between the separate key polynomials and the graded algebra of a valuation. Then, in Section 4, we focus on the particular cases of immediate and separate extensions. For this purpose, we generalize a result of F. Delon which shows that if the residue characteristic is 0 and  $\nu$  is archimedean, then any simple immediate algebraic extension of valued field is dense (Theorem 4.4).

## 1. IMMEDIATE EXTENSIONS

In this section,  $L|K$  is an extension of fields and  $\nu$  is a  $K$ -module valuation on  $L$ .

If  $M$  is a  $K$ -module, then we assume that  $\nu M$  has no greatest element. This holds if  $\nu K$  is not trivial. Indeed, for every  $x \in K$  with  $\nu(x) > 0$  and  $y \in M$ , we have  $xy \in M$  and  $\nu(xy) = \nu(x) + \nu(y) > \nu(y)$ .

We generalize the definitions of immediate and dense extensions to extensions of  $K$ -modules. We list some basic properties, then we characterize the immediate and dense extensions by means of the sets  $\nu(\chi^n - K_{n-1}[\chi])$  (Proposition 1.8).

**Notations 1.1.** Let  $l \in L$  and  $M$  be a  $K$ -submodule of  $L$ . We denote by  $\nu(l - M)$  the subset  $\{\nu(l - x) \mid x \in M \setminus \{l\}\}$  of  $\nu M$ . For any polynomial  $f$ , we denote by  $\nu(f(M))$ , or  $\nu f(M)$ , the subset  $\{\nu(f(x)) \mid x \in M\} \setminus \{\infty\}$ .

Note that  $\nu(l - M) \cap \nu M$  is an initial segment of  $\nu M$ .

**Definitions 1.2.** Let  $M \subseteq N$  be  $K$ -submodules of  $L$ , and  $l \in L$ .

We say that  $l$  is *pseudo-limit* over  $(M, \nu)$  if  $\nu(l - M) \subseteq \nu M$ , and  $\nu(l - M)$  has no maximal element.

We say that  $l$  is *limit* over  $(M, \nu)$  if  $\nu(l - M) = \nu M$ .

The extension  $(N|M, \nu)$  is said to be *immediate* if every element of  $N$  is pseudo-limit over  $(M, \nu)$ .

The extension  $(N|M, \nu)$  is said to be *dense* if every element of  $N$  is limit over  $M$ .

**Remark 1.3.** It follows that if  $(N|M, \nu)$  is dense, then it is immediate.

**Lemma 1.4.** Let  $l \in L$  and  $M$  be a  $K$ -submodule of  $L$ . The element  $l$  is pseudo-limit over  $(M, \nu)$  if, and only if, for every  $x \in M$  there exists  $y \in M$  such that  $\nu(l - y) > \nu(l - x)$ .

*Proof.* If  $l$  is pseudo-limit, then  $\nu(l - M)$  has no maximal element. Hence, for every  $x \in M$  there exists  $y \in M$  such that  $\nu(l - y) > \nu(l - x)$ . Conversely, if for every  $x \in M$  there exists  $y \in M$  such that  $\nu(l - y) > \nu(l - x)$ , then  $\nu(l - M)$  has no maximal element. Now, let  $x, y \in M$  such that  $\nu(l - y) > \nu(l - x)$ . Then, since  $\nu(l - x) = \nu(l - y + y - x) < \nu(l - y)$ , it follows that  $\nu(l - y) = \nu(y - x) \in \nu M$ . Hence  $\nu(l - M) \subseteq \nu M$ . Consequently,  $l$  is pseudo-limit over  $M$ .  $\square$

**Notations 1.5.** For  $\gamma$  in  $\nu L$  and  $M$  a  $K$ -submodule of  $L$ , let  $M_{\gamma, \nu}$  be the  $K_\nu$ -module  $\{x \in M \mid \nu(x) \geq \gamma\} / \{x \in M \mid \nu(x) > \gamma\}$ . In the case where  $\gamma = 0$ , we often write  $M_\nu$  instead of  $M_{0, \nu}$ . For  $f \in K[l]$  with  $\nu(f) \geq \gamma$ , we denote by  $f_{\gamma, \nu}$  the class of  $f$  modulo the ideal  $\{g \in K[l] \mid \nu(g) > \gamma\}$  of the ring  $K[l]$ .

**Remark 1.6.**

- 1)  $(N|M, \nu)$  is immediate if, and only if,  $\nu N = \nu M$  and, for every  $\gamma \in \nu M$ ,  $N_{\gamma, \nu} = M_{\gamma, \nu}$ .
- 2)  $(N|K, \nu)$  is immediate if, and only if,  $\nu N = \nu K$  and  $N_\nu = K_\nu$ .

*Proof.*

1) We assume that  $(N|M, \nu)$  is immediate. Let  $l \in N$ . Since  $l$  is pseudo-limit over  $M$ , there is  $x \in M$  such that  $\nu(l - 0) < \nu(l - x)$ . Hence  $\nu(l) = \nu(x) \in \nu M$ . Let  $\gamma \in \nu M$  and  $l \in N$  such that  $\nu(l) = \gamma$ . There exists  $x \in M$  such that  $\gamma = \nu(l - 0) < \nu(l - x)$ . Hence  $l_{\gamma, \nu} = x_{\gamma, \nu} \in M_{\gamma, \nu}$ .

Assume that  $\nu N = \nu M$  and, for every  $\gamma \in \nu M$ ,  $N_{\gamma, \nu} = M_{\gamma, \nu}$ . Let  $l \in N$  and  $x \in M$ . Let  $y_1 \in M$  such that  $\nu(y_1) = \nu(l - x)$  and  $(y_1)_{\nu(y_1), \nu} = l_{\nu(y_1), \nu}$ . Therefore,  $\nu(l - x - y_1) > \nu(y_1) = \nu(l - x)$ . We let  $y := x + y_1$ . By Lemma 1.4,  $(N|M, \nu)$  is immediate.

2) It remains to prove that if  $\nu N = \nu K$  and  $N_\nu = K_\nu$ , then for every  $\gamma \in \nu M$ ,  $N_{\gamma, \nu} = M_{\gamma, \nu}$ . Let  $\gamma \in \nu K$  and  $l \in N$  such that  $\nu(l) = \gamma$ . We take  $x_1 \in K \setminus \{0\}$  such that  $\nu(x_1) = \nu(l)$ . Hence  $\nu(l/x_1) = 0$ . Since  $N$  is a  $K$ -module,  $l/x_1$  belongs to  $N$ . Now, let  $x_2 \in K$  such that  $(l/x_1)_\nu = (x_2)_\nu$ . Therefore,  $\nu(l/x_1 - x_2) > 0$ . Set  $x := x_1 x_2$ . Then,  $\nu(l - x) = \nu(x_1(l/x_1 - x_2)) = \nu(x_1) + \nu(l/x_1 - x_2) > \nu(x_1) = \nu(l)$ . It follows that  $l_{\gamma, \nu} = x_{\gamma, \nu}$ .  $\square$

**Definitions 1.7.** We say that  $\nu$  is *partially multiplicative*, or a *p-m valuation*, if for every  $f, g$  in  $K[X]$  such that  $\deg(f) + \deg(g) < [K[X]:K]$  we have  $\nu(fg) = \nu(f) + \nu(g)$ . In the case where  $[K[X]:K] = \infty$ , we assume in addition that  $g \neq 0$  implies  $\nu(f/g) = \nu(f) - \nu(g)$ , that is,  $\nu$  is a field valuation in the usual sense. If  $\nu$  is a field valuation (in the usual sense), then we will also say that it is *multiplicative*.

**Proposition 1.8.** Let  $K(X)|K$  be a simple extension of valued fields, where  $X$  is algebraic or transcendental over  $K$ , and let  $d, d'$  in  $\mathbb{N} \cup \{\infty\}$  such that  $0 \leq d < d' \leq [K[X]:K] - 1$ , (that is, if  $X$  is algebraic over  $K$ , then  $d' < [K[X]:K]$ ). Recall that we defined  $K_0[X] = K$  and  $K_\infty[X] = K[X]$ .

a)  $(K_{d'}[X]|K_d[X], \nu)$  is immediate if, and only if, for every  $n \in \{d+1, \dots, d'\}$ ,  $\nu(X^n - K_{n-1}[X])$  has no maximal element.

b) Assume that  $\nu$  is a p-m valuation on  $K(X)$  and that  $\nu K_d[X]$  is a subgroup of  $\nu K(X)$ . Then the extension  $(K_{d'}[X]|K_d[X], \nu)$  is dense if, and only if, for every  $n \in \{d+1, \dots, d'\}$ ,  $\nu(X^n - K_{n-1}[X]) = \nu K_d[X]$ .

*Proof.* a)  $\Rightarrow$ . Assume that, for some  $n \in \{d+1, \dots, d'\}$ ,  $\nu(X^n - K_{n-1}[X])$  has a greatest element  $\nu(f)$ . If  $\nu(f)$  is not in  $\nu K_d[X]$ , then  $\nu K_{d'}[X] \neq \nu K_d[X]$ , and, by Remark 1.6 1), the extension is not immediate. If  $\nu(f) \in \nu K_d[X]$ , say  $\nu(f) = \gamma$ . For every  $g \in K_d[X]$  we have  $\nu(f - g) \leq \nu(f)$ , hence  $f_{\gamma, \nu} \neq g_{\gamma, \nu}$ . It follows that  $(K_{d'}[X])_{\gamma, \nu} \neq (K_d[X])_{\gamma, \nu}$ , so the extension is not immediate.

$\Leftarrow$ . Assume that  $(K_{d'}[X]|K_d[X], \nu)$  is not immediate, and let  $n$  be the smallest integer such that, for some polynomial  $f$  of degree  $n$ , either  $\nu(f) \notin \nu K_d[X]$  or, for every  $g \in K_d[X]$ ,  $f_{\gamma, \nu} \neq g_{\gamma, \nu}$ , where  $\gamma = \nu(f)$ . Note that, by dividing  $f$  by an element of  $K$ , we can assume that  $f$  is a monic polynomial of degree  $n$ . First assume that  $\nu(f(X)) \notin \nu K_d[X]$ , and let  $g$  be a monic polynomial of degree  $n$ . Then  $\deg(f - g) < n$ , hence, by minimality of  $n$ ,  $\nu(f - g) \in \nu K_d[X]$ . So  $\nu(f - g) \neq \nu(f)$ . It follows:  $\nu(g) = \min(\nu(g - f), \nu(f)) \leq \nu(f)$ , which proves that  $\nu(f)$  is the greatest element of  $\nu(X^n - K_{n-1}[X])$ .

Assume that  $\gamma = \nu(f) \in \nu K_d[X]$ , and, for every  $g \in K_d[X]$ ,  $f_{\gamma, \nu} \neq g_{\gamma, \nu}$ . Let  $g$  be a monic polynomial of degree  $n$ . If  $\nu(g - f) \neq \gamma$ , then  $\nu(g) = \min(\nu(g - f), \nu(f)) \leq \nu(f)$ . Now, assume that  $\nu(g - f) = \gamma$ . By minimality of  $n$ , we have  $f_{\gamma, \nu} \neq (g - f)_{\gamma, \nu}$ . So  $\nu(g) = \nu(g - f + f) = \min(\nu(g - f), \nu(f)) \leq \nu(f)$ . Consequently,  $\nu(f)$  is the greatest element of  $\nu(X^n - K_{n-1}[X])$ .

b) If  $(K_{d'}[X]|K_d[X], \nu)$  is dense, then it is immediate. Hence for every  $n \in \{d+1, \dots, d'\}$  we have:  $\nu(\chi^n - K_{n-1}[X]) \subseteq \nu K_d[X]$ . Furthermore:  $\nu(\chi^{d+1} - K_d[X]) = \nu K_d[X]$ . Now, for every  $n \in \{d+2, \dots, d'\}$  we have  $\nu(\chi^n - K_{n-1}[X]) \supseteq \nu(\chi^n - \chi^{n-(d+1)}K_d[X]) = (n - (d+1))\nu(\chi) + \nu(\chi^{d+1} - K_d[X]) = \nu K_d[X]$  (since  $\nu K_d[X]$  is a subgroup). It follows:  $\nu(\chi^n - K_{n-1}[X]) = \nu K_d[X]$ .

Conversely, assume that for every  $n \in \{d+1, \dots, d'\}$  we have  $\nu(\chi^n - K_{n-1}[X]) = \nu K_d[X]$ . Then in particular it has no greatest element. So by a)  $(K_{d'}[X]|K_d[X], \nu)$  is immediate. Let  $f \in K_{d'}[X]$ ,  $n$  be its degree and  $x_n$  be the coefficient of  $\chi^n$  in  $f$ . Without loss of generality we can assume that  $n \in \{d+1, \dots, d'\}$ . We show that for every  $\gamma \in K_d[X]$  there is  $g \in K_d[X]$  such that  $\nu(f - g) > \gamma$ . Since  $\nu(\chi^n - K_{n-1}[X]) = \nu K_d[X]$  which is a group, there is  $g_{n-1} \in K_{n-1}[X]$  such that  $\nu((f/x_n) - (g_{n-1}/x_n)) > \gamma - \nu(x_n)$ . Hence  $\nu(f - g_{n-1}) > \gamma$ . In the same way, for  $d \leq j \leq n-2$  we get  $g_j \in K_j[X]$  such that  $\nu(g_{j+1} - g_j) > \gamma$ . Therefore  $\nu(f - g_d) = \nu(f - g_{n-1} + \dots + g_{d+1} - g_d) \geq \min(\nu(f - g_{n-1}), \dots, \nu(g_{d+1} - g_d)) > \gamma$ . So every element of  $K_{d'}[X]$  is limit over  $K_d[X]$ . This implies that  $(K_{d'}[X]|K_d[X], \nu)$  is dense.  $\square$

## 2. SEPARATE EXTENSIONS

In this section,  $L|K$  is an extension of fields and  $\nu$  is a  $K$ -module valuation on  $L$ .

In subsection 2.1 we list definitions and properties of separate extensions and we prove Theorem 2.7 (which implies that a valued fields extension  $(L|K, \nu)$  is separate if, and only if, for every finitely generated  $K$ -submodule  $M$  of  $L$  and  $l \in L \setminus M$ , the set  $\nu(l - M)$  has a maximal element), and Theorem 2.8 (which implies that if  $(L|K, \nu)$  is a finite algebraic extension of valued fields, then  $(L|K, \nu)$  is separate if, and only if,  $[L:K] = [L_\nu:K_\nu](\nu L:\nu K)$ ).

In subsection 2.2 we prove Proposition 2.19, which characterizes separateness by means of the sets  $\nu(\chi^n - K_{n-1}[X])$ .

Subsection 2.3 is dedicated to the graded algebra associated to a valuation, and to the links with separate extensions.

### 2.1. Basic properties.

**Definitions 2.1.** (Baur, [B 82]) Let  $M$  be a  $K$ -submodule of  $L$ .

1) A sequence  $(l_1, \dots, l_n)$  of  $L$  is said to be *separate over  $M$*  (or  $\nu$ -separate if necessary) if for every  $x_1, \dots, x_n$  in  $M$ , we have:  $\nu(x_1 l_1 + \dots + x_n l_n) = \min_{1 \leq i \leq n} \nu(x_i l_i)$ . If  $M = K$ , then we say separate instead of separate over  $K$ .

2) The extension  $(L|K, \nu)$  is said to be *separate* if every finitely generated  $K$ -submodule of  $L$  admits a basis which is separate over  $K$ . If this holds, then we say that  $\nu$  is separate (or *separate over  $K$* ).

**Definition 2.2.** Let  $M$  be a finitely generated  $K$ -submodule of  $L$ . If  $M$  admits a basis which is separate over  $K$ , then we say that  $M$  is *separate*.

**Remark 2.3.** ([B 81]) If the sequence  $(l_1, \dots, l_n)$  of  $L$  is separate over  $K$ , then  $l_1, \dots, l_n$  are linearly independent over  $K$ .

The following properties provide examples of separate extensions.

**Proposition 2.4.** ([B 81]) If  $(K, \nu)$  is a maximal valued field, then every multiplicative extension of  $(K, \nu)$  is separate.

**Theorem 2.5.** ([D 88, Corollaire 7]) Assume that  $\nu$  is multiplicative on  $L$  and that  $(K, \nu)$  is henselian of residue characteristic 0. Then any algebraic extension of  $(K, \nu)$  is separate.

The remainder of this subsection is dedicated to properties which make clear the difference between immediate and separate extensions.

**Theorem 2.6.** ([D 88, p. 421]) Assume that  $\nu$  is multiplicative on  $L$  and that  $(K, \nu)$  is henselian of residue characteristic 0. Then  $(L|K, \nu)$  is separate if, and only if,  $L$  is linearly disjoint over  $K$  from every immediate extension of  $(K, \nu)$ .

**Theorem 2.7.** (Delon) Let  $N$  be a  $K$ -submodule of  $L$ . Then,  $(N|K, \nu)$  is a separate extension if, and only if, for every finitely generated  $K$ -submodule  $M$  of  $N$  and  $l \in N \setminus M$ , the set  $\nu(l - M)$  has a maximal element.

This theorem has been stated in [D 88, p. 421], assuming that  $(K, \nu)$  is henselian and  $\text{char}(K_\nu) = 0$ . So we give the proof for completeness. We will also prove the next theorem.

We know that if  $L|K$  is finite and  $\nu$  is multiplicative, then  $1 \leq [L_\nu:K_\nu](\nu L:\nu K) \leq [L:K]$ . Furthermore, by Remark 1.6,  $(L|K, \nu)$  is immediate if, and only if,  $1 = [L_\nu:K_\nu](\nu L:\nu K)$ . The following theorem proves that  $(L|K, \nu)$  being separate can be seen as the opposite case.

**Theorem 2.8.** Assume that  $L|K$  is a finite algebraic extension of fields and that  $\nu$  is multiplicative on  $L$ . Then  $(L|K, \nu)$  is separate if, and only if,  $[L:K] = [L_\nu:K_\nu](\nu L:\nu K)$ .

**Definition 2.9.** Let  $(L|K, \nu)$  be a finite extension of valued fields (where  $\nu$  is multiplicative). Then  $(L|K, \nu)$  is *defectless* if  $\frac{[L:K]}{[L_\nu:K_\nu](\nu L:\nu K)}$  is equal to the number of extensions of  $\nu|_K$  to  $L$ .

We recall that  $(K, \nu)$  is *henselian* if  $\nu$  admits a unique extension to every algebraic extension of  $K$ .

**Corollary 2.10.** *If  $(K, \nu)$  is henselian, then every defectless finite algebraic extension of  $(K, \nu)$  is separate.*

We start with properties a separate sequences.

**Lemma 2.11.** ([B 82] p. 676) *Let  $(l_1, \dots, l_n)$  be a separate sequence of elements of  $L$ ,  $y$  in  $L$  and  $k_1, \dots, k_n$  in  $K$ . The following holds.*

*Every subsequence of  $(l_1, \dots, l_n)$  is separate.*

*The sequence  $(k_1 l_1, \dots, k_n l_n)$  is separate.*

*If  $\nu$  is multiplicative on  $L$  then the sequence  $(y l_1, \dots, y l_n)$  is separate.*

**Lemma 2.12.** ([B 81], [B 82] (S4), p. 676) *Let  $(l_1, \dots, l_n)$  be a sequence of elements of  $L$  such that:  $\forall i, 1 \leq i \leq n, \nu(l_i) = 0$ . Then  $(l_1, \dots, l_n)$  is separate if, and only if,  $(l_1)_\nu, \dots, (l_n)_\nu$  are linearly independent over  $K_\nu$ . This can be generalized in the following way. If  $\nu(l_1) = \dots = \nu(l_n) = g$ , then  $(l_1, \dots, l_n)$  is separate if, and only if, for every  $x_1, \dots, x_n$  in  $\{x \in K \mid \nu(x) = 0\} \cup \{0\}$ , either  $\nu(x_1 l_1 + \dots + x_n l_n) = g$  or  $x_1 = \dots = x_n = 0$ .*

**Proposition 2.13.** *Let  $l_{i1}, \dots, l_{in_i}, 1 \leq i \leq p$ , be sequences which satisfy:*

$$\forall i, 1 \leq i \leq p, \forall j, 1 \leq j \leq n_i, \nu(l_{ij}) = \nu(l_{i1}) < \infty$$

*and the  $\nu(l_{i1})$  are pairwise non-congruent modulo  $\nu K$ . The following assertions are equivalent.*

*The sequence  $l_{11}, \dots, l_{1n_1}, l_{21}, \dots, l_{2n_2}, \dots, l_{p1}, \dots, l_{pn_p}$  is separate.*

*For every  $i$  in  $\{1, \dots, p\}$ ,  $l_{i1}, l_{i2}, \dots, l_{in_i}$  is separate.*

*If  $\nu$  is multiplicative on  $L$  then this condition is equivalent to:*

*for every  $i$  in  $\{1, \dots, p\}$ ,  $1, (l_{i2} l_{i1}^{-1})_\nu, \dots, (l_{in_i} l_{i1}^{-1})_\nu$  are linearly independent over  $K_\nu$ .*

*Proof.* Assume that the sequence is separate. Then by Lemma 2.11, for  $1 \leq i \leq p$ , the sequence  $l_{i1}, l_{i2}, \dots, l_{in_i}$  is separate.

Conversely, let  $x_{11}, \dots, x_{1n_1}, x_{21}, \dots, x_{2n_2}, \dots, x_{p1}, \dots, x_{pn_p}$  in  $K$ . For  $1 \leq i \leq p$ , set  $y_i = x_{i1} l_{i1} + \dots + x_{in_i} l_{in_i}$ . Since  $l_{i1}, \dots, l_{in_i}$  is separate, we have:  $\nu(y_i) = \min\{\nu(x_{ij}) + \nu(l_{ij}) \mid 1 \leq j \leq n_i\}$ . Therefore, the  $\nu(y_i)$ 's are pairwise non-congruent modulo  $\nu K$ . In particular, they are pairwise distinct, and  $\nu(y_1 + \dots + y_p) = \min\{\nu(y_i) \mid 1 \leq i \leq p\}$ . This proves that the sequence  $l_{11}, \dots, l_{1n_1}, l_{21}, \dots, l_{2n_2}, \dots, l_{p1}, \dots, l_{pn_p}$  is separate.

If  $\nu$  is multiplicative on  $L$ , then

$$l_{i1}, l_{i2}, \dots, l_{in_i} \text{ is separate if, and only if, } 1, (l_{i2} l_{i1}^{-1}), \dots, (l_{in_i} l_{i1}^{-1}) \text{ is separate.}$$

By Lemma 2.12, this in turn is equivalent to  $1, (l_{i2} l_{i1}^{-1})_\nu, \dots, (l_{in_i} l_{i1}^{-1})_\nu$  are linearly independent over  $K_\nu$ .  $\square$

**Lemma 2.14.** ([D 88] Lemme 5) *Let  $M \subseteq N$  be two  $K$ -submodules of  $L$  such that  $M$  is finitely generated and  $N$  admits a separate basis. Then  $M$  admits a separate basis (in other words, it is separate).*

The following theorem is an immediate consequence of Lemma 2.14.

**Theorem 2.15.** *Assume that  $L$  is a finite algebraic extension of  $K$ . Then  $(L|K, \nu)$  is separate if, and only if, the  $K$ -module  $L$  admits a separate basis.*

**Remark.** Theorem 2.15 is not true if  $(L|K)$  is not a finite algebraic extension, (see [D 88] p. 426). However, we will see in Proposition 2.17 that it remains true if  $L$  is generated by one transcendental element and  $\nu$  is multiplicative.

*Proof of Theorem 2.8.* Set  $r := [L_\nu:K_\nu]$  and  $q := (\nu L:\nu K)$ . By properties of valuations, we have that  $rq \leq [L:K]$ .

Assume that  $[L:K] = rq$ . Let  $x_1, \dots, x_r, y_1, \dots, y_q$  be elements of  $L$  such that  $\nu(x_1) = \dots = \nu(x_r) = 0$ ,  $(x_1)_\nu, \dots, (x_r)_\nu$  are linearly independent over  $K_\nu$ , and  $\nu(y_1), \dots, \nu(y_q)$  are pairwise non-congruent modulo  $\nu K$ . By Proposition 2.13, the sequence  $\{x_i y_j \mid 1 \leq i \leq r, 1 \leq j \leq q\}$  is separate. It follows that they are linearly independent over  $K$ . Since its cardinal is  $rq$ , it is a basis of  $L$  over  $K$ . Now, by Theorem 2.15,  $(L|K, \nu)$  is separate.

Assume that  $(L|K, \nu)$  is separate, so  $L$  admits a separate basis  $\mathcal{B}$ . We define an equivalence relation over  $\mathcal{B}$  by setting  $y_1 \sim y_2 \Leftrightarrow \nu(y_1) \equiv \nu(y_2)$  modulo  $\nu K$ . By Lemma 2.11, we can assume that all the elements of every class of  $\mathcal{B}$  modulo  $\sim$  have the same valuation. Let  $\mathcal{C} := \{l_1, \dots, l_p\}$  be a class of  $\mathcal{B}$  modulo  $\sim$ . By Lemma 2.11  $1, (l_2/l_1), \dots, (l_p/l_1)$  is a separate sequence of elements of  $L$  with valuation



0. We deduce from Lemma 2.12 that  $p \leq r$ . Now, if  $p < r$ , then there exists  $l'_{p+1}$  in  $L$  such that  $1_\nu, (l_2/l_1^{-1})_\nu, \dots, (l_p/l_1^{-1})_\nu, (l'_{p+1})_\nu$  are linearly independent over  $K_\nu$ . Then  $1, (l_2/l_1^{-1}), \dots, (l_p/l_1^{-1}), l'_{p+1}$  is a separate sequence, hence so is  $l_1, l_2, \dots, l_p, l_{p+1}$ , where  $l_{p+1} = l'_{p+1}l_1$ . By Proposition 2.13,  $\mathcal{B} \cup \{l_{p+1}\}$  is a separate sequence, hence  $\mathcal{B}$  is not a maximal subset of linearly independent elements, so it is not a basis: a contradiction. It follows that  $p = r$ . Now, there are at most  $(\nu L : \nu K) = q$  classes modulo  $\sim$ . Since  $\mathcal{B}$  is a separate basis, for every  $l \in L$  there exists  $x \in K$  and  $b \in \mathcal{B}$  such that  $\nu(l) = \nu(xb)$ . It follows that there are exactly  $q$  classes modulo  $\sim$ . Hence  $\mathcal{B}$  is the disjoint union of  $q$  classes, each one contains  $r$  elements. It follows:  $[L : K] = \text{card}(\mathcal{B}) = rq$ .  $\square$

**Lemma 2.16.** *Assume that  $(L|K, \nu)$  is separate. Let  $M$  be a finite  $K$ -submodule of  $L$  and  $l \in L \setminus M$ . Then every separate basis of  $M$  extends to a separate basis of the  $K$ -submodule generated by  $M$  and  $l$ .*

*Proof.* Consider a separate basis  $\mathcal{B}$  of  $M$  and a separate basis  $\mathcal{B}'$  of  $N := M \oplus K \cdot l$ . Since  $\nu(M) \subseteq \nu(N)$ , the number of classes of  $\mathcal{B}'$  modulo the relation  $\sim$  defined in the proof of Theorem 2.8 is greater or equal to the number of classes of  $\mathcal{B}$  modulo  $\sim$ . If it is greater, then we add to  $\mathcal{B}$  an element of the additional class, and we get the separate basis of  $N$ . Otherwise, one of the classes of  $\mathcal{B}'$  has more elements than the corresponding class of  $\mathcal{B}$ . Say  $l'_1, \dots, l'_{k+1}$  and  $l_1, \dots, l_k$ . By Lemma 2.11, we can assume that all the element of these classes have the same valuation  $\gamma$ . Assume that there exists a family  $(x_{ij})_{1 \leq i \leq k+1, 1 \leq j \leq k}$  in  $\{x \in K \mid \nu(x) = 0\} \cup \{0\}$  such that for  $i \in \{1, \dots, k+1\}$ :  $(*)i \nu(l'_i - (x_{i1}l_1 + \dots + x_{ik}l_k)) > \gamma$ . We show that we get a contradiction. Without loss of generality we can assume that  $x_{11} \neq 0$ . Then:  $\nu(l_1 + x_{12}x_{11}^{-1}l_2 + \dots + x_{1k}x_{11}^{-1}l_k - x_{11}^{-1}l'_1) > \gamma$ . For  $i \geq 2$  we put  $x_{11}^{-1}l'_1 - (x_{12}x_{11}^{-1}l_2 + \dots + x_{1k}x_{11}^{-1}l_k)$  in place of  $l_1$ . So we get an inequality  $\nu(l'_i + x_{11}^{-1}l'_1 - (x_{12}x_{11}^{-1}l_2 + \dots + x_{1k}x_{11}^{-1}l_k)) > \gamma$ . We can eliminate  $(x_{1j}x_{11}^{-1} + x_{ij})l_j$  if  $\nu(x_{1j}x_{11}^{-1} + x_{ij}) > 0$ , so we can assume that all the coefficients belong to  $\{x \in K \mid \nu(x) = 0\} \cup \{0\}$ . We proceed in the same way with  $l_2, \dots, l_k$ . Since there are  $k+1$  inequalities  $(*)i$ , finally we get some  $\nu(y_1l'_1 + \dots + y_{k+1}l'_{k+1}) > \gamma = \min \nu(y_i l'_i)$ : a contradiction. Hence there is some  $l'_i$ , say  $l'_{k+1}$  such that for every  $x_j$  in  $\{x \in K \mid \nu(x) = 0\} \cup \{0\}$  ( $1 \leq j \leq k$ ):  $\nu(l'_{k+1} + x_1l_1 + \dots + x_kl_k) = \gamma$ . It follows that the sequence  $l_1, \dots, l_k, l'_{k+1}$  is separate. By Proposition 2.13, the sequence  $\mathcal{B} \cup \{l'_{k+1}\}$  is separate.  $\square$

*Proof of Theorem 2.7.* Assume that  $(N|K, \nu)$  is separate. Let  $M$  be a finitely generated  $K$ -submodule of  $N$ , and  $l \in N \setminus M$ . By Lemma 2.16, there exist a basis  $l_1, \dots, l_k$  of  $M \oplus K \cdot l$  such that  $l_1, \dots, l_{k-1}$  belong to  $M$ . Now,  $l$  can be written as  $l = x_1l_1 + \dots + x_kl_k$ , with  $x_1, \dots, x_k$  in  $K$ . Since  $l \notin M$ , we have  $x_k \neq 0$ . Hence for every  $y$  in  $M$  there exist  $y_1, \dots, y_{k-1}$  in  $K$  such that  $l - y = y_1l_1 + \dots + y_{k-1}l_{k-1} + x_kl_k$ , hence  $\nu(l - y) \leq \nu(x_kl_k)$ . So  $\nu(x_kl_k) = \max \nu(l - M)$ . Conversely, we prove by induction on the dimension of the submodule  $M$  that it contains a separate basis. If  $\dim(M) = 1$ , then the result is trivial. Assume that  $M$  admits a separate basis  $l_1, \dots, l_k$  and let  $l \notin M$ . Let  $y \in M$  such that  $\nu(l - y) = \max \nu(l - M)$ , and set  $l_{k+1} = l - y$ . We show that the family  $l_1, \dots, l_k, l_{k+1}$  is separate. Let  $x_1, \dots, x_k, x_{k+1}$  in  $K$ , with  $x_{k+1} \neq 0$ , and  $\gamma := \min(\nu(x_1l_1), \dots, \nu(x_kl_k))$ . If  $\gamma < \nu(x_{k+1}l_{k+1})$ , then  $\nu(x_1l_1 + \dots + x_kl_k + x_{k+1}l_{k+1}) = \gamma = \min(\nu(x_1l_1), \dots, \nu(x_kl_k), \nu(x_{k+1}l_{k+1}))$ . If  $\gamma > \nu(x_{k+1}l_{k+1})$ , then  $\nu(x_1l_1 + \dots + x_kl_k + x_{k+1}l_{k+1}) = \nu(x_{k+1}l_{k+1}) = \min(\nu(x_1l_1), \dots, \nu(x_kl_k), \nu(x_{k+1}l_{k+1}))$ . Assume that  $\gamma = \nu(x_{k+1}l_{k+1})$ . Since  $\nu(l_{k+1})$  is the maximum of  $\nu(l - M)$ ,  $\gamma \leq \nu(x_1l_1 + \dots + x_kl_k + x_{k+1}l_{k+1}) = \nu(x_{k+1}) + \nu(x_1x_{k+1}^{-1}l_1 + \dots + x_kx_{k+1}^{-1}l_k + l_{k+1}) \leq \nu(x_{k+1}) + \nu(l_{k+1}) = \gamma$ . So,  $\nu(x_1l_1 + \dots + x_kl_k + x_{k+1}l_{k+1}) = \min(\nu(x_1l_1), \dots, \nu(x_kl_k), \nu(x_{k+1}l_{k+1}))$ .  $\square$

**2.2. Extensions generated by one element.** In this subsection,  $K(\chi)|K$  is a simple extension of fields, where  $\chi$  is algebraic or transcendental over  $K$ .

**Proposition 2.17.** *Let  $d \in \mathbb{N} \cup \{\infty\}$  such that  $0 < d \leq [K[\chi] : K] - 1$ . Then  $(K_d[\chi]|K, \nu)$  is separate if, and only if, the  $K$ -module  $K_d[\chi]$  admits a separate basis. Furthermore, we can assume that the degree mapping is one-to-one, and that every polynomial of this basis is monic.*

*Assume that  $\chi$  is transcendental over  $K$ , and that  $\nu$  is multiplicative on  $K(\chi)$ . Then  $(K(\chi)|K, \nu)$  is separate if, and only if, the  $K$ -module  $K[\chi]$  admits a separate basis. ( $\Rightarrow$  holds even if  $\nu$  is not multiplicative).*

*Proof.* Assume that  $(K_d[\chi]|K, \nu)$  is separate. By Lemma 2.16, the separate basis 1 of  $K$  can be completed in a separate basis of the module generated by 1 and  $\chi$ . Necessarily, the second element of this basis has degree 1. Let  $n \geq 1$  and assume that the  $K$ -module  $K_n[\chi]$  of polynomials of degree at most  $n$  has a separate basis of  $(n+1)$  elements of respective degrees  $0, 1, \dots, n$ . By Lemma 2.16, this separate basis can be completed in a separate basis of  $K_{n+1}[\chi]$ , and the degree of the new element is  $n+1$ . So we get the required separate basis by induction. By Lemma 2.11 we can assume that every polynomial of this basis is monic.

Conversely, assume that  $K_d[\chi]$  contains a separate basis, and let  $M$  be a finitely generated  $K$ -submodule. By Lemma 2.14,  $M$  has a separate basis.

Assume that  $\chi$  is transcendental over  $K$ , and that  $\nu$  is multiplicative on  $K(\chi)$  and let  $M$  be a finitely generated submodule of  $K(\chi)$ . Then there is a polynomial  $f(\chi) \neq 0$  such that  $f(\chi) \cdot M \subseteq K[\chi]$ . We take

a separate basis of  $f(\chi) \cdot M$ , and we divide all its elements by  $f(\chi)$  so that, since  $\nu$  is multiplicative, by Lemma 2.11, we get a separate basis of  $M$ .  $\square$

**Example 2.18.** Let  $(K, \nu)$  be a valued field. Pick some  $x$  in  $K$ , some  $\gamma$  in an extension of  $\nu K$ , and, for every  $x_0, x_1, \dots, x_n$  in  $K$ , set  $\nu'(x_n(\chi - x)^n + \dots + x_1(\chi - x) + x_0) := \min(\nu(x_n) + n\gamma, \dots, \nu(x_1) + \gamma, \nu(x_0))$ . Then one can check that  $\nu'$  defines a p-m valuation on the ring  $K[\chi]$ . We say that  $\nu'$  is a *Gauss valuation*. The p-m valuation  $\nu'$  is separate and  $1, (\chi - x), \dots, (\chi - x)^n, \dots$  is a separate basis of  $(K[\chi], \nu')$ .

Note that if  $\nu''$  is another  $K$ -module valuation on  $K[\chi]$  which extends  $\nu$  and such that  $\nu(\chi - x) = \gamma$ , then, for every  $f$  in  $K[\chi]$ ,  $\nu'(f) \leq \nu''(f)$ .

We state a refinement of Theorem 2.7, which characterizes separate extensions by means of initial segments. This proposition completes Proposition 1.8.

**Proposition 2.19.** *Let  $d \in \mathbb{N} \cup \{\infty\}$  such that  $0 < d \leq [K[\chi] : K] - 1$ . The extension  $(K_d[\chi] | K, \nu)$  is separate if, and only if, for every integer  $n$ ,  $1 \leq n \leq d$ ,  $\nu(\chi^n - K_{n-1}[\chi])$  has a maximal element. Assume that  $\chi$  is transcendental over  $K$ , and that  $\nu$  is multiplicative on  $K(\chi)$ . Then  $(K(\chi) | K, \nu)$  is separate if, and only if, for every  $n \in \mathbb{N}^*$ ,  $\nu(\chi^n - K_{n-1}[\chi])$  has a maximal element.*

*Proof.* In both equivalences,  $\Rightarrow$  follows from Theorem 2.7. In order to prove the converse, we construct by induction a separate basis such that the degree mapping is one-to-one. Then, by Proposition 2.17,  $(K_d[\chi] | K, \nu)$  is separate. The case where  $\chi$  is transcendental also follows from Proposition 2.17. Trivially, 1 is a separate basis of  $K_0[\chi] = K$ . Assume that we have a separate basis  $(f_0, \dots, f_{n-1})$  of  $K_{n-1}[\chi]$ . Let  $f_n$  be a monic polynomial such that  $\nu(f_n) = \max(\nu(\chi^n - K_{n-1}[\chi]))$ . Since the degree of  $f_n$  is  $n$ ,  $(f_0, \dots, f_{n-1}, f_n)$  is a basis of  $K_n[\chi]$ . Let  $f := x_n f_n + \dots + x_0$  in  $K_n[\chi]$ . If  $x_n = 0$ , then by induction hypothesis  $\nu(f) = \min_{0 \leq i \leq n-1} \nu(x_i f_i) = \min_{0 \leq i \leq n} \nu(x_i f_i)$ . Now we assume:  $x_n \neq 0$ . Since  $\nu(f_n)$  is maximal, we

have  $\nu\left(\frac{f}{x_n}\right) \leq \nu(f_n)$ . If  $\nu\left(\frac{f}{x_n} - f_n\right) < \nu(f_n)$ , then  $\nu\left(\frac{f}{x_n} - f_n\right) = \nu\left(\frac{f}{x_n}\right)$ . So:

$$\begin{aligned} \nu(f) &= \nu(x_n) + \nu\left(\frac{f}{x_n}\right) = \nu(x_n) + \nu\left(\frac{f}{x_n} - f_n\right) = \\ &= \nu(x_n) + \min_{0 \leq i \leq n-1} \nu\left(\frac{x_i f_i}{x_n}\right) = \min_{0 \leq i \leq n-1} \nu(x_i f_i) = \min_{0 \leq i \leq n} \nu(x_i f_i). \end{aligned}$$

If  $\nu\left(\frac{f}{x_n} - f_n\right) \geq \nu(f_n)$ , then  $\min_{0 \leq i \leq n-1} \nu(x_i f_i) \geq \nu(x_n f_n)$ . Furthermore, since  $\nu(f_n)$  is maximal we have  $\nu\left(\frac{f}{x_n}\right) = \nu(f)$ . Therefore:  $\nu(f) = \min_{0 \leq i \leq n} \nu(x_i f_i)$ .  $\square$

**2.3. Graded algebra associated to a valuation.** In the proofs of Remark 3.35 and Theorem 3.37 we will introduce the graded algebra associated to a valuation. We will also show more properties in Subsection 3.8 because they are used in the definition of key polynomials by F. J. Herrera Govantes, W. Mahloud, M. A. Olalla Acosta and M. Spivakovsky. Now, we review some observations.

Let  $(K, \nu)$  be a valued field. Recall that, for every  $\gamma \in \nu K$ ,  $K_{\gamma, \nu}$  denotes the  $K_\nu$ -module  $\{x \in K \mid \nu(x) \geq \gamma\} / \{x \in K \mid \nu(x) > \gamma\}$ . Now, let  $G_\nu(K)$  be the graded algebra  $G_\nu(K) = \bigoplus_{\gamma \in \nu K} K_{\gamma, \nu}$ .

In the case where  $K$  is the valued field  $k((\Gamma))$  of generalized formal power series with coefficients in a field  $k$  and exponents in a linearly ordered abelian group  $\Gamma$ :

$$k((\Gamma)) = \left\{ \sum_{\gamma \in \Lambda} x_\gamma X^\gamma \mid \Lambda \text{ is a well-ordered subset of } \Gamma, \text{ and } \forall \gamma \in \Lambda \ x_\gamma \in k \right\},$$

then  $G_\nu(K)$  is isomorphic to the ring of generalized polynomials  $k[\Gamma]$ , the subring of all elements of  $k((\Gamma))$  with finite support. More generally, the  $K$ -module  $G_\nu(K)$  is isomorphic to the  $K$ -module  $K_\nu[\nu K]$  of polynomials with coefficients in  $K_\nu$  and exponents in  $\nu K$ .

If  $K$  contains a lifting of  $\nu K$ , then we can assume that these graded algebras are isomorphic. In particular, if  $\nu K = \mathbb{Z}$ , then they are isomorphic. If  $(K', \nu')$  is an  $\aleph_1$ -saturated elementary extension of  $(K, \nu)$ , then it contains a lifting of its value group (see [K 75]). Hence  $G_{\nu'}(K')$  is isomorphic to the ring of polynomials  $K'_{\nu'}(\nu' K')$ . Therefore every graded algebra  $G_\nu(K)$  embeds in a ring of polynomials. If  $(K, \nu)$  contains a lifting  $K_0$  of its residue field and a lifting  $\Gamma$  of  $\nu K$ , then it contains the algebra  $K_0[\Gamma]$ , which is isomorphic to  $G_\nu(K)$ . Now, if  $(K, \nu)$  is henselian and  $\text{char}(K_\nu) = 0$ , then we know that it admits a lifting of  $K_\nu$ . It follows that every valued field  $(K, \nu)$  of residue characteristic 0 admits an extension  $(K', \nu')$  which contains a subalgebra which is isomorphic to  $G_\nu(K)$ . Furthermore,  $(K', \nu')$  embeds in the power series field  $K'_0((\nu' K'))$  equipped with the canonical valuation.

For every  $x \in K$ , let  $in_\nu(x) := x_{\nu(x), \nu}$  be the image of  $x$  in  $K_{\nu(x), \nu}$ , which is also its image in  $G_\nu(K)$ . In

the case of a subfield of a power series field, we have  $in_\nu \left( \sum_{\gamma \in \Lambda} x_\gamma \chi^\gamma \right) = x_{\gamma_0} \chi^{\gamma_0}$ , where  $\gamma_0$  is the smallest element of the *support* of the serie (i.e. the well ordered subset  $\Lambda$  of  $\nu K$  such that  $x_{\gamma_0} \neq 0$ ). In general, for every  $x, y$  in  $K$ , we have  $in_\nu(x)in_\nu(y) = in_\nu(xy)$ . Assume that  $\nu(x) = \nu(y)$ . If  $in_\nu(x) = -in_\nu(y)$ , then  $in_\nu(x + y) = 0$ . Otherwise,  $in_\nu(x + y) = in_\nu(x) + in_\nu(y)$ .

An element of  $G_\nu(K)$  is called *homogeneous* if it belongs to  $\bigcup_{\gamma \in \nu K} K_{\gamma, \nu}$ . In the case of a polynomial ring, this is equivalent to being a monomial. One can see that the inversible elements of  $G_\nu(K)$  are the homogeneous ones.

For further purposes, if  $M$  is a  $K$ -submodule of  $L$ , we denote by  $G_\nu(M)$  the additive group  $\bigoplus_{\gamma \in \nu M} M_{\gamma, \nu}$ .

By Remark 1.6, an extension of valued fields  $(L|K, \nu)$  is immediate if, and only if,  $G_\nu(L) = G_\nu(K)$ .

Now we turn to the separate case. Let  $(L|K, \nu)$  be an extension of valued fields, and  $l_1, \dots, l_n$  in  $L$ .

**Fact 2.20.** *The family  $(l_1, \dots, l_n)$  is separate over  $(K, \nu)$  if, and only if,  $in_\nu(l_1), \dots, in_\nu(l_n)$  are linearly independent over  $G_\nu(K)$ .*

*If  $(l_1, \dots, l_n)$  is a maximal separate family, then  $(in_\nu(l_1), \dots, in_\nu(l_n))$  is a basis of  $G_\nu(L)$ .*

*The extension  $(L|K, \nu)$  is separate if, and only if,  $G_\nu(L)$  is a  $G_\nu(K)$ -module of dimension  $[L:K]$ .*

*Proof.* Recall that the family  $(l_1, \dots, l_n)$  is a separate over  $(K, \nu)$  if, and only if, for every  $x_1, \dots, x_n$  in  $K$ ,  $\nu(x_1 l_1, \dots, x_n l_n) = \min(\nu(x_1 l_1), \dots, \nu(x_n l_n))$ . Now, this equivalent to saying that for every  $x_1, \dots, x_n$  in  $K$  with  $\nu(x_1 l_1) = \dots = \nu(x_n l_n)$ , we have  $\nu(x_1 l_1, \dots, x_n l_n) = \nu(x_1 l_1)$ . This last equality is equivalent to  $in_\nu(x_1)in_\nu(l_1) + \dots + in_\nu(x_n)in_\nu(l_n) \neq 0$ . So, if  $in_\nu(l_1), \dots, in_\nu(l_n)$  are linearly independent in the  $G_\nu(K)$ -module  $G_\nu(L)$ , then the family  $(l_1, \dots, l_n)$  is separate. Now, assume that for every  $x_1, \dots, x_n$  in  $K$  with  $\nu(x_1 l_1) = \dots = \nu(x_n l_n)$ , we have  $\nu(x_1 l_1, \dots, x_n l_n) = \nu(x_1 l_1)$ . Let  $y_1, \dots, y_n$  in  $G_\nu(K)$ . Every  $y_j$  can be written as a finite sum of homogeneous elements:  $y_j = in_\nu(x_{j,1}) + \dots + in_\nu(x_{j,i_j})$ . It follows that  $y_1 in_\nu(l_1) + \dots + y_n in_\nu(l_n)$  can be written as a sum of  $in_\nu(x_{1,k_1})in_\nu(l_1) + \dots + in_\nu(x_{n,k_n})in_\nu(l_n)$ , where the non-zero  $in_\nu(x_{j,k_j})in_\nu(l_j)$  have the same valuation. Therefore,  $y_1 in_\nu(l_1) + \dots + y_n in_\nu(l_n) \neq 0$ . Consequently, the family  $(l_1, \dots, l_n)$  is separate over  $(K, \nu)$  if, and only if,  $in_\nu(l_1), \dots, in_\nu(l_n)$  are linearly independent over  $G_\nu(K)$ . Furthermore, if  $(l_1, \dots, l_n)$  is a maximal separate family, then  $(in_\nu(l_1), \dots, in_\nu(l_n))$  is a basis of  $G_\nu(L)$ . Now, if  $[L:K]$  is finite, then the dimension of the  $G_\nu(K)$ -module  $G_\nu(L)$  is  $[L_\nu:K_\nu] \cdot (\nu L:\nu K)$ . Hence, by Theorem 2.8,  $(L|K, \nu)$  is separate if, and only if,  $G_\nu(L)$  is a  $G_\nu(K)$ -module of dimension  $[L:K]$ .  $\square$

The following lemma shows that if, for  $l \in L$ ,  $in_\nu(l)$  satisfies a relation of algebraic dependence over  $G_\nu(K)$ , then we can define its irreducible polynomial.

**Lemma 2.21.** *Let  $l \in L$ . Assume that  $in_\nu(l)$  satisfies a relation of algebraic dependence over  $G_\nu(K)$ , and let  $n$  be the smallest degree such that such a relation exists. Then,  $in_\nu(l)$  satisfies a relation of the form  $in_\nu(l)^n + in_\nu(x_{n-1})in_\nu(l)^{n-1} + \dots + in_\nu(x_0) = 0$ , where  $x_0, \dots, x_{n-1}$  belong to  $K$  and  $\nu(x_0) = \dots = \nu(x_{n-1}l^{n-1}) = \nu(l^n)$ .*

*Proof.* See for example [HOS 07].  $\square$

We sometimes call *homogeneous* a polynomial  $in_\nu(\chi)^n + in_\nu(x_{n-1})in_\nu(\chi)^{n-1} + \dots + in_\nu(x_0)$ , where  $x_0, \dots, x_{n-1}$  belong to  $K$ , such that  $\nu(x_0) = \dots = \nu(x_{n-1}\chi^{n-1}) = \nu(\chi^n)$ .

### 3. KEY POLYNOMIALS.

In this section,  $(K(\chi)|K, \nu)$  is an extension of valued fields, where  $\chi$  is algebraic or transcendental over  $K$ .

In Subsection 3.1 we state the definitions of key polynomials, w-key polynomials and key degrees and we also state basic facts. In particular we prove Proposition 3.7 ( $\Phi$  is a w-key polynomial if, and only if,  $(K[X]/(\Phi), \bar{\nu})$  is a valued field, where  $\bar{\nu}$  is defined by  $\bar{\nu}(f(X) + \Phi) = \nu(f(X))$ ).

In Subsection 3.2 we compare Definition 3.2 and S. MacLane's definition ([ML 36a] and [ML 36b]).

In Subsection 3.3 we define the valuations  $\nu_\Phi$ . We show that it follows from properties of extensions of valuations that these valuation are not, in general, multiplicative. We also deduce characterizations of w-key polynomials and of MacLane's key polynomials (Propositions 3.20 and 3.24).

The bases defined by polynomials are introduced in Subsection 3.4.

In Subsection 3.5 we focus on properties of key degrees. Given a key polynomial, Proposition 3.27 allows to construct other key polynomials of the same degree. Proposition 3.34 gives a sufficient condition for  $\nu$  being equal to  $\nu_\Phi$  for some w-key polynomial  $\Phi$ . This holds in particular if  $\chi$  is transcendental and Abhyankar inequality is an equality. In Theorems 3.36 and 3.37 we characterize the successor of a given key degree, in the immediate case and in the separate case. Proposition 3.39 states a necessary and

sufficient condition for an integer being a valuational key degree. Next we show that if  $d$  is a key degree such that  $\chi^d$  is limit over  $K_{d-1}[\chi]$ , then  $d$  is the greatest key degree (Proposition 3.41).

In Subsection 3.6, we define families of key polynomials and associated separate valuations (in the same way as S. MacLane and M. Vaquié). In Theorem 3.44 we show that the valuation of any polynomial is equal to its valuation with respect to some separate valuations so constructed.

Subsection 3.7 is dedicated to proving that the definitions of key polynomials and of abstract key polynomials are equivalent.

In [HOS 07] and [HMOS 14], families of key polynomials are constructed by lifting some polynomials of the graded algebra of the valuation. So, in Subsection 3.8 we see links between key polynomials and their images in the graded algebra associated to the valuation.

### 3.1. Definitions.

**Notation 3.1.** Let  $\Phi$  be a monic polynomial of degree  $d \geq 1$ . For  $f, g$  in  $K_{d-1}[\chi]$ , we will denote by  $q_\Phi(f, g)$  and  $r_\Phi(f, g)$  respectively (in short  $q(f, g)$  and  $r(f, g)$ ) the quotient and the remainder of the euclidean division of  $fg$  by  $\Phi$ . In other words,  $q(f, g)$  and  $r(f, g)$  belong to  $K_{d-1}[\chi]$  and  $fg = \Phi \cdot q(f, g) + r(f, g)$ .

**Definitions 3.2.** Let  $\nu$  be a  $K$ -module valuation on  $K(\chi)$  and  $\Phi$  be a monic polynomial of degree  $d \geq 1$ . We say that  $\Phi$  is a *weak key polynomial* for  $\nu$  (or a *w-key polynomial* for  $\nu$ ) if, for every  $f, g$  in  $K_{d-1}[\chi]$  with  $\deg(f) + \deg(g) < [K[\chi] : K]$ , we have  $\nu(fg) = \nu(r(f, g))$ .

We say that  $\Phi$  is a *key polynomial* for  $\nu$  if, for every  $f, g$  in  $K_{d-1}[\chi]$  with  $\deg(f) + \deg(g) < [K[\chi] : K]$ , we have  $\nu(fg) = \nu(r(f, g)) < \nu(q(f, g) \cdot \Phi)$ .

Let  $d$  be a positive integer.

We say that  $d$  is a *key degree* of  $(K(\chi)|K, \nu)$  if there exists a key polynomial of degree  $d$ .

Assume that  $d$  is a key degree. If  $\nu(\chi^d - K_{d-1}[\chi])$  has no maximal element, then we say that  $d$  is an *immediate* key degree. Otherwise, we say that  $d$  is a *separate* key degree. If this maximum does not belong to  $\nu K_{d-1}[\chi]$ , then we say that  $d$  is a *valuational* key degree. If this maximum belongs to  $\nu K_{d-1}[\chi]$ , then we say that  $d$  is a *residual* key degree.

**Remarks 3.3.** 1) The integer 1 is a key degree. Furthermore, every monic polynomial of degree 1 is a key polynomial.

2) Every w-key polynomial is irreducible.

3) If  $\nu$  is partially multiplicative and  $\Phi$  is a w-key polynomial (resp. a key polynomial) for  $\nu$  of degree  $d$ , then, for every p-m valuation  $\nu'$  such that the restriction of  $\nu'$  to  $K_{d-1}[\chi]$  is equal to the restriction of  $\nu$  and  $\nu'(\Phi) \geq \nu(\Phi)$ ,  $\Phi$  is a w-key polynomial (resp. a key polynomial) for  $\nu'$ .

4) Assume that  $\nu$  is partially multiplicative. If  $\Phi$  is a w-key polynomial, then  $\nu(\Phi) \geq \{\nu(r(f, g)) - \nu(q(f, g)) \mid f \in K_{d-1}[\chi], g \in K_{d-1}[\chi], \deg(f) + \deg(g) < [K[\chi] : K]\}$ . If  $\Phi$  is a monic polynomial, then  $\Phi$  is a key polynomial if, and only if,  $\nu(\Phi) > \{\nu(r(f, g)) - \nu(q(f, g)) \mid f \in K_{d-1}[\chi], g \in K_{d-1}[\chi], \deg(f) + \deg(g) < [K[\chi] : K]\}$ .

*Proof.* 1) and 3) are trivial.

2) Assume that there exist two polynomials  $f, g$  in  $K_{d-1}[\chi]$  such that  $fg = \Phi$ , then  $r(f, g) = 0$ , and  $\nu(r(f, g)) = \infty > \nu(fg)$ . Hence  $\Phi$  is not a w-key polynomial for  $\nu$ .

4) Clearly, if  $\Phi$  is a w-key polynomial (resp. a key polynomial), then  $\nu(\Phi) \geq \{\nu(r(f, g)) - \nu(q(f, g)) \mid f \in K_{d-1}[\chi], g \in K_{d-1}[\chi], \deg(f) + \deg(g) < [K[\chi] : K]\}$  (resp.  $\nu(\Phi) > \{\nu(r(f, g)) - \nu(q(f, g)) \mid f \in K_{d-1}[\chi], g \in K_{d-1}[\chi], \deg(f) + \deg(g) < [K[\chi] : K]\}$ ). Now, if  $\nu(\Phi) > \nu(r(f, g)) - \nu(q(f, g))$ , then  $\nu(q(f, g)\Phi) > \nu(r(f, g)) = \nu(fg - q(f, g)\Phi)$ . Hence  $\nu(r(f, g)) - \nu(fg)$ .  $\square$

**Remark 3.4.** Let  $d$  be a key degree. Then, by Proposition 1.8,  $d$  is an immediate key degree if, and only if, the extension  $(K[\chi]|K_{d-1}[\chi], \nu)$  is immediate.

The following lemma explains the distinction that we make between the valuational and the residual key degrees.

**Lemma 3.5.** Let  $d$  be a positive integer. Assume that  $\nu(\chi^d - K_{d-1}[\chi])$  has a maximum  $\nu(\Phi)$ . Then either  $\nu(\Phi) \notin \nu K_{d-1}[\chi]$ , or  $\Phi_{\nu(\Phi), \nu} \notin (K_{d-1}[\chi])_{\nu(\Phi), \nu}$ .

*Proof.* Assume that  $\nu(\Phi) \in \nu K_{d-1}[\chi]$ . Since  $\nu(\Phi)$  is maximal in  $\nu(\chi^d - K_{d-1}[\chi])$ , for every  $f \in K_{d-1}[\chi]$  such that  $\nu(f) = \nu(\Phi)$  we have  $\nu(\Phi - f) = \nu(\Phi) = \nu(f)$ . Hence  $(\Phi)_{\nu(\Phi), \nu} \neq f_{\nu(\Phi), \nu}$ . It follows that  $\Phi_{\nu(\Phi), \nu} \notin (K_{d-1}[\chi])_{\nu(\Phi), \nu}$ .  $\square$

Let  $\Phi$  be a monic irreducible polynomial of  $K[X]$  of degree  $d \geq 1$  ( $K[X]$ , the ring of formal polynomials with coefficients in  $K$ ). Then  $K[X]/(\Phi)$  is a field, such that the canonical epimorphism  $\rho : K[X] \rightarrow K[X]/(\Phi)$  is an isomorphism from the  $K$ -module  $K_{d-1}[X]$  onto the  $K$ -module  $K[X]/(\Phi)$ . Now, for  $f, g$  in  $K_{d-1}[X]$ , we set  $f * g := r(f, g)$ . Then  $\rho(f * g) = \rho(r(f, g)) = \rho(fg)$ . Hence  $(K_{d-1}[X], +, *)$  is a field which is isomorphic to  $K[X]/(\Phi)$ . The same operation can be defined in  $K[\chi]$  whenever  $d \leq [K(\chi) : K]/2$ .

**Notation 3.6.** Let  $\Phi$  be a monic irreducible polynomial of  $K[\chi]$  of degree  $d$ ,  $1 \leq d \leq [K(\chi):K]/2$ . The field  $(K_{d-1}[\chi], +, *)$  defined above will be denoted by  $K_\Phi$ .

Note that if  $\nu$  is a p-m valuation on the field  $K(\chi)$ , then its restriction to  $K_{d-1}[\chi]$  induces a valuation of the  $K$ -modules  $K_\Phi$  and  $K[\chi]/(\Phi)$ . If  $\mathcal{Y}$  is a root of  $\Phi(X)$  in some algebraic extension, then the fields  $K_\Phi$  and  $K[\mathcal{Y}]$  are isomorphic.

**Proposition 3.7.** *Let  $\Phi$  be an irreducible monic polynomial of degree  $d$ ,  $1 \leq d \leq [K(\chi):K]/2$ , and  $\nu$  be a p-m valuation on  $K(\chi)$ . Then  $\Phi$  is a w-key polynomial for  $\nu$  if, and only if, the valued  $K$ -module  $(K_\Phi, \nu)$  is a valued field.*

*Proof.* Let  $f, g$  in  $K_{d-1}[\chi]$ . Then  $\nu(f * g) = \nu(r(f, g))$  and  $\nu(fg) = \nu(f) + \nu(g)$ . Hence  $(K_\Phi, \nu)$  is a valued field if, and only if, for every  $f, g$  in  $K_{d-1}[\chi]$  we have:  $\nu(r(f, g)) = \nu(fg)$ . This in turn is equivalent to saying that  $\Phi$  is a w-key polynomial.  $\square$

**Corollary 3.8.** *Let  $\Phi$  be a w-key polynomial of degree  $d$ ,  $1 \leq d \leq [K(\chi):K]/2$ , and  $\nu$  be a p-m valuation on  $K(\chi)$ . Then  $\nu K_{d-1}[\chi]$  is a subgroup of  $\nu K(\chi)$ .*

*Proof.* Indeed, if  $\Phi$  is a w-key polynomial, then for every  $f, g$  in  $K_{d-1}[\chi]$  we have  $\nu(f * g) = \nu(r_\Phi(f, g)) = \nu(fg) = \nu(f) + \nu(g)$ . Hence  $\nu K_\Phi = \nu K_{d-1}[\chi]$ .  $\square$

**Remark 3.9.** Let  $\Phi$  be a monic polynomial of degree  $d$ ,  $1 \leq d \leq [K(\chi):K]/2$ , and  $\nu$  be a p-m valuation on  $K(\chi)$ .

(1) The polynomial  $\Phi$  is a key polynomial if, and only if, for every  $f, g$  in  $K_{d-1}[\chi]$ ,  $\nu(fg) = \nu(r_\Phi(f, g))$  and  $(fg)_{\nu(fg), \nu} = (r_\Phi(f, g))_{\nu(fg), \nu}$ . Indeed, if  $\nu(fg) = \nu(r_\Phi(f, g))$ , then  $\nu(fg - r_\Phi(f, g)) > 0$  is equivalent to  $(fg)_{\nu(fg), \nu} = (r_\Phi(f, g))_{\nu(fg), \nu}$ .

(2) If  $\Phi$  is a key polynomial, then the group  $G_\nu(K_{d-1}[\chi])$  is a subalgebra of  $G_\nu(K(\chi))$ . Indeed, since  $(fg)_{\nu(fg), \nu} = f_{\nu(f), \nu} g_{\nu(g), \nu}$ ,  $\Phi$  is a key polynomial if, and only if, for every  $f, g$  in  $K_{d-1}[\chi]$ ,  $(f * g)_{\nu(fg), \nu} = f_{\nu(f), \nu} g_{\nu(g), \nu}$ . Therefore, the group  $G_\nu(K_{d-1}[\chi])$  is a subalgebra of  $G_\nu(K(\chi))$ , and it is isomorphic to  $G_\nu(K_\Phi)$ . Hence  $G_\nu(K)$  embeds in  $G_\nu(K_\Phi)$  and  $G_\nu(K_\Phi)$  embeds in  $G_\nu(K(\chi))$ .

(3) It follows from Proposition 3.7 that if  $\Phi$  is a key polynomial, then  $K_{d-1}[\chi]_\nu$  is a subfield of  $K(\chi)_\nu$ .

**3.2. MacLane's key polynomials.** In [ML 36a] and [ML 36b] S. MacLane defined key polynomials in the case of discrete valuations. M. Vaquié ([V 07]) generalized this definition to arbitrary valuations. In [V 07] the key polynomials are defined on the ring of formal polynomials. The case of algebraic extensions is obtained by means of a pseudo-valuation, by quotienting  $K[\chi]$  by the socle of  $\nu$ . A *pseudo-valuation*  $\nu$  of  $K$  is a mapping from  $K$  onto a linearly ordered group  $\nu K$  together with an element  $\infty$  which shares the properties of multiplicative valuations except  $\nu(x) = \infty \Rightarrow x = 0$ . In this case the set  $I := \{x \in K \mid \nu(x) = \infty\}$  is a prime ideal which is called the *socle* of  $\nu$ . Then  $\nu$  induces a p-m valuation on the integral domain  $K/I$ . So, in [V 07]  $\chi$  is transcendental over  $K$  and  $\nu$  is a valuation or a pseudo-valuation. Now, in Definition 3.2, we can assume that  $\nu$  is a pseudo-valuation and that  $\chi$  is transcendental. Here we extend the definition of [V 07] to the case of an algebraic extension, and we do not require  $\chi$  being transcendental.

In this subsection,  $\nu$  is a  $K$ -module valuation on  $K(\chi)$  or a pseudo-valuation.

**Definition 3.10.** Let  $\Phi$  be a monic polynomial of degree  $d$  and  $d' := [K[\chi]:K] - d$ . We say that  $\Phi$  is  $\nu$ -minimal if for every  $f \in K_{d-1}[\chi]$  and every  $h \in K_{d'-1}[\chi]$ ,  $\nu(f - h\Phi) = \min(\nu(f), \nu(h\Phi))$ . We say that  $\Phi$  is  $\nu$ -irreducible if for every  $f, g$  in  $K[\chi]$  with  $\deg(f) + \deg(g) < [K[\chi]:K]$  such that, for every  $h \in K_{d'-1}[\chi]$ ,  $\nu(f - h\Phi) = \min(\nu(f), \nu(h\Phi))$  and  $\nu(g - h\Phi) = \min(\nu(g), \nu(h\Phi))$ , we have:  $\forall h \in K_{d'-1}[\chi] \nu(fg - h\Phi) = \min(\nu(fg), \nu(h\Phi))$ .

**Remark 3.11.** By setting  $h := 1$  in above definition, we see that every monic polynomial, which is  $\nu$ -minimal and  $\nu$ -irreducible, is irreducible.

**Proposition 3.12.** *Assume that  $\nu$  is a p-m valuation or a pseudo-valuation. Let  $d < [K[\chi]:K]$  in  $\mathbb{N}^*$ ,  $d' := [K[\chi]:K] - d$ ,  $\Phi$  be a non constant monic polynomial in  $K[\chi]$  of degree  $d$ . The following assertions are equivalent.*

- 1)  $\Phi$  is  $\nu$ -minimal and  $\nu$ -irreducible.
- 2) For every  $f, g$  in  $K_{d-1}[\chi]$  with  $\deg(f) + \deg(g) < [K[\chi]:K]$  and every  $h$  in  $K_{d'-1}[\chi]$ , we have  $\nu(fg + h\Phi) = \min(\nu(fg), \nu(h\Phi))$ .
- 3)  $\Phi$  is a w-key polynomial such that the sequence  $(\Phi^m)$  ( $md < [K(\chi):K]$ ) is separate over  $K_{d-1}[\chi]$ .

Before proving Proposition 3.12 we state a lemma.

**Lemma 3.13.** ([ML 36a] Lemma 4.3) *Assume that  $\nu$  is a p-m valuation or a pseudo-valuation. Let  $d < [K[\chi]:K]$  in  $\mathbb{N}^*$ ,  $d' := [K[\chi]:K] - d$ ,  $\Phi$  be a  $\nu$ -minimal non constant monic polynomial in  $K[\chi]$  of degree  $d$ . Let  $f \in K[\chi]$  and  $f = q\Phi + r$  be the euclidean division of  $f$  by  $\Phi$ . The following assertions are equivalent.*

- a)  $\nu(r) > \nu(f)$
- b)  $\nu(r) > \min(\nu(f), \nu(q\Phi))$
- c)  $\exists h \in K_{d'-1}[\chi] \ \nu(f - h\Phi) > \min(\nu(f), \nu(h\Phi))$ .

*Proof.* Trivially we have: a)  $\Rightarrow$  b) and b)  $\Rightarrow$  c). We prove c)  $\Rightarrow$  a). Let  $h \in K_{d'-1}[\chi]$ . By hypothesis,  $\deg(r) < d$  and  $\deg(q) < d'$ . Hence  $\nu(f - h\Phi) = \nu((q - h)\Phi + r) = \min(\nu((q - h)\Phi), \nu(r)) \leq \nu(r)$ . Therefore  $\nu(f - h\Phi) > \min(\nu(f), \nu(h\Phi)) \Rightarrow \nu(r) > \min(\nu(f), \nu(h\Phi))$ . Now,  $\nu(f - h\Phi) > \min(\nu(f), \nu(h\Phi)) \Rightarrow \nu(f) = \nu(h\Phi)$ , hence  $\nu(r) > \nu(f)$ .  $\square$

*Proof of Proposition 3.12.*

2)  $\Rightarrow$  1). Assume that  $\Phi$  satisfies the hypothesis of 2). By setting  $g = 1$  it follows that  $\Phi$  is a non constant monic polynomial in  $K[\chi]$  of degree  $d$  such that for every  $f$  in  $K_{d-1}[\chi]$  and every  $h$  in  $K_{d'-1}[\chi]$  we have  $\nu(f + h\Phi) = \min(\nu(f), \nu(h\Phi))$ . Hence  $\Phi$  is  $\nu$ -minimal. In order to prove that  $\Phi$  is  $\nu$  irreducible, let  $f$  and  $g$  in  $K[\chi]$  such that  $\deg(f) + \deg(g) < [K[\chi] : K]$  and for every  $h \in K_{d'-1}[\chi]$ ,  $\nu(f - h\Phi) = \min(\nu(f), \nu(h\Phi))$  and  $\nu(g - h\Phi) = \min(\nu(g), \nu(h\Phi))$ . By euclidean division,  $f$  et  $g$  can be written as  $f = q\Phi + r$  and  $g = q'\Phi + r'$ . By Lemma 3.13, we have  $\nu(f) = \nu(r)$  and  $\nu(g) = \nu(r')$ . Let  $h \in K_{d'-1}[\chi]$ . Then  $\nu(fg - h\Phi) = \nu((qq'\Phi + qr' + q'r - h)\Phi + rr') = \min(\nu((qq'\Phi + qr' + q'r - h)\Phi), \nu(rr'))$ , because both of  $r$  and  $r'$  belong to  $K_{d-1}[\chi]$ . Hence  $\nu(fg - h\Phi) \leq \nu(rr') = \nu(fg)$ . Now, we have  $\min(\nu(fg), \nu(h\Phi)) \leq \nu(fg - h\Phi) \leq \nu(fg)$ . So,  $\nu(fg - h\Phi) = \min(\nu(fg), \nu(h\Phi))$ . Hence  $\Phi$  is  $\nu$ -irreducible.

1)  $\Rightarrow$  2). We assume that  $\Phi$  be a monic,  $\nu$ -minimal and  $\nu$ -irreducible polynomial. Let  $f, g$  in  $K_{d-1}[\chi]$  with  $\deg(f) + \deg(g) < [K[\chi] : K]$ . Since  $\Phi$  is  $\nu$ -minimal, for every  $h$  in  $K[\chi]$  we have  $\nu(f + h\Phi) = \min(\nu(f), \nu(h\Phi))$  and  $\nu(g + h\Phi) = \min(\nu(g), \nu(h\Phi))$ . Now,  $\Phi$  is  $\nu$ -irreducible, hence  $\nu(fg + h\Phi) = \min(\nu(fg), \nu(h\Phi))$ .

2)  $\Rightarrow$  3). Assume that  $\Phi$  satisfies the hypothesis of 1) and 2). Let  $f, g$  in  $K_{d-1}[\chi]$  with  $\deg(f) + \deg(g) < [K[\chi] : K]$ . By letting  $h := -q_\Phi(f, g)$ , 2) implies  $\nu(r_\Phi(f, g)) = \min(\nu(fg), \nu(q_\Phi(f, g)\Phi))$ . Since  $\Phi$  is  $\nu$ -minimal, by Lemma 3.13 we have  $\nu(r_\Phi(f, g)) = \nu(fg)$ . Hence  $\Phi$  is a w-key polynomial. Now, let  $m \in \mathbb{N}^*$  with  $dm < [K[\chi] : K]$ ,  $f_0, \dots, f_m$  in  $K_{d-1}[\chi]$ . We have:  $\nu(f_m\Phi^m + \dots + f_0) = \min(\nu((f_m\Phi^{n-1} + \dots + f_1)\Phi), \nu(f_0))$ ,  $\nu(f_m\Phi^{n-1} + \dots + f_1) = \min(\nu((f_m\Phi^{n-2} + \dots + f_2)\Phi), \nu(f_1))$ , and so on. So by induction we have  $\nu(f_m\Phi^m + \dots + f_0) = \min(\nu(f_m\Phi^{n-1}), \dots, \nu(f_0))$ . Hence the family  $(\Phi^m)_{m \in \mathbb{N}}$  is separate over  $K_{d-1}[\chi]$ .

3)  $\Rightarrow$  2). We take  $f, g$  in  $K_{d-1}[\chi]$  and  $h$  in  $K_{d'-1}[\chi]$  with  $\deg(f) + \deg(g) < [K[\chi] : K]$ . The polynomial  $h$  can be written as  $h = h_m\Phi^m + \dots + h_1\Phi + h_0$ , where  $h_m, \dots, h_1, h_0$  belong to  $K_{d-1}[\chi]$ . Let  $q := q(f, g)$  and  $r := r(f, g)$ ; since  $f$  and  $g$  belong to  $K_{d-1}[\chi]$ , we have  $\deg(q) < d$ , i.e.  $q \in K_{d-1}[\chi]$ . We have:  $\nu(fg + h\Phi) = \nu(h_m\Phi^{m+1} + \dots + h_1\Phi^2 + (q + h_0)\Phi + r) = \min(\nu(h_m\Phi^{m+1}), \dots, \nu(h_1\Phi^2), \nu((q + h_0)\Phi), \nu(r))$ . Since  $\Phi$  is a w-key polynomial, we have  $\nu(fg) = \nu(r) \leq \nu(q\Phi)$ . If  $\nu(h_0\Phi) \geq \nu(r)$ , then  $\nu((q + h_0)\Phi) \geq \nu(r)$ . Hence  $\min(\nu((q + h_0)\Phi), \nu(r)) = \nu(r) = \min(\nu(h_0\Phi), \nu(r))$ . If  $\nu(h_0\Phi) < \nu(r)$ , then  $\min(\nu((q + h_0)\Phi), \nu(r)) = \nu(h_0\Phi) = \min(\nu(h_0\Phi), \nu(r))$ . Therefore:  $\nu(fg + h\Phi) = \min(\nu(h_m\Phi^{m+1}), \dots, \nu(h_1\Phi^2), \nu(h_0\Phi), \nu(fg)) = \min(\nu(h\Phi), \nu(fg))$ .  $\square$

**Remark 3.14.** We use the hypothesis “ $\nu$  is a p-m valuation” for proving 2)  $\Rightarrow$  1). For proving 2)  $\Rightarrow$  3) the condition “for every  $f \in K_{d'-1}[\chi]$ ,  $\nu(\Phi f) = \nu(\Phi) + \nu(f)$ ” is sufficient. The remainder of the proof remains true with a  $K$ -module valuation.

In MacLane’s definition, the key polynomials are the  $\nu$ -minimal and  $\nu$ -irreducible polynomials. Proposition 3.12 shows that this definition is stronger than Definition 3.2. The difference will appear more clearly in Subsection 3.3 (for example Remark 3.19). Now, we extend the definition of S. MacLane to  $K$ -module valuations.

**Definition 3.15.** Let  $\Phi$  be a polynomial of degree  $d$ . We say that  $\Phi$  is a *ML key polynomial* for  $\nu$  if  $\Phi$  is a w-key polynomial such that the sequence  $(\Phi^m)$  ( $md < [K(\chi) : K]$ ) is separate over  $K_{d-1}[\chi]$ .

**3.3. Separate valuations defined by key polynomials.** Assume that  $\Phi$  is a monic irreducible polynomial of degree  $d$ , and let  $\gamma$  be an element of an extension of  $\nu K(\chi)$ .

For every  $f := f_0 + f_1\Phi + \dots + f_m\Phi^m$  in  $K[\chi]$  (with  $f_0, f_1, \dots, f_m$  in  $K_{d-1}[\chi]$  and  $\deg(f_m) + dm < [K[\chi] : K]$ ), set  $\nu'(f) = \min_{0 \leq i \leq m} \nu(f_i) + i\gamma$ .

Assume that  $\nu$  is a p-m valuation on  $K(\chi)$  and that  $\Phi$  is a w-key polynomial for  $\nu$  of degree  $d$ . We saw in Remarks 3.3 4) that the set  $\{\nu(r(f, g)) - \nu(q(f, g)) \mid f \in K_{d-1}[\chi], g \in K_{d-1}[\chi], \deg(f) + \deg(g) < [K[\chi] : K]\}$  is bounded above by  $\nu(\Phi)$ . We extend the addition of elements of  $\nu K$  to the addition of Dedekind cuts in the usual way. We also define an element  $-\infty < \nu K(\chi)$ , and we let  $\delta + \infty = \infty$ ,  $\delta - \infty = -\infty$ , for every  $\delta$  in the Dedekind completion of  $\nu K(\chi)$ .

**Proposition 3.16.** Let  $\Phi$  be a monic irreducible polynomial of degree  $d$ ,  $\gamma$  be an element of an extension of  $\nu K(\chi)$ ,  $\nu$  be a  $K$ -module valuation defined on  $K_{d-1}[\chi]$ , and  $\nu'$  be defined as above.

1) The application  $\nu'$  is a  $K$ -module valuation and the family  $(\Phi^m)$  ( $dm < [K[\chi] : K]$ ) is separate over  $K_{d-1}[\chi]$ .

2) Assume that  $\nu$  is a  $p$ - $m$  valuation and that  $\Phi$  is a  $w$ -key polynomial for  $\nu$ . Denote by  $\beta$  the upper-bound of the set  $\{\nu(r(f, g)) - \nu(q(f, g)) \mid f \in K_{d-1}[X], g \in K_{d-1}[X], \deg(f) + \deg(g) < [K[X]:K]\}$ , and assume that  $\gamma \geq \beta$ .

a) For every  $f = f_m\Phi^m + \dots + f_1\Phi + f_0$ ,  $g = g_m\Phi^m + \dots + g_1\Phi + g_0$ , with  $f_0, \dots, f_m, g_0, \dots, g_m$  in  $K_{d-1}[X]$  such that  $\deg(f) + \deg(g) < [K[X]:K]$  we have:

$$\nu'(fg) = \nu' \left( \sum_{j=0}^{2m} \left( \sum_{i=0}^j r(f_i, g_{j-i}) \right) \Phi^j \right) \leq \nu' \left( fg - \sum_{j=0}^{2m} \left( \sum_{i=0}^j r(f_i, g_{j-i}) \right) \Phi^j \right) - (\gamma - \beta).$$

(Here if  $i > m$ , then we let  $f_i = g_i = 0$ .) In particular,  $\nu'$  is a  $p$ - $m$  valuation.

b) The polynomial  $\Phi$  is a ML key polynomial for  $\nu'$ . Furthermore it is a key polynomial if, and only if,  $\gamma > \beta$  or  $\beta$  is not a maximum.

*Proof.* 1) Clearly, if  $\nu'$  is a  $K$ -module valuation, then the family  $(\Phi^m)$  is separate over  $K_{d-1}[X]$ . Let  $f_0, \dots, f_m, g_0, \dots, g_m$  in  $K_{d-1}[X]$  and let  $f = f_m\Phi^m + \dots + f_1\Phi + f_0$ ,  $g = g_m\Phi^m + \dots + g_1\Phi + g_0$ . We have trivially

$$\begin{aligned} \nu'(f+g) &= \min_{0 \leq j \leq m} (\nu(f_j + g_j) + j\gamma) \geq \min_{0 \leq j \leq m} (\min(\nu(f_j) + j\gamma, \nu(g_j)) + j\gamma) \\ &\geq \min(\nu'(f), \nu'(g)). \end{aligned}$$

We have that  $\nu'(f) = \infty \Leftrightarrow f = 0$ . Now, for  $x \in K$ ,  $\nu'(xf) = \nu'(x) + \nu'(f)$ , since  $\nu$  is a  $K$ -module valuation.

2) a) Since  $\deg(f) + \deg(g) < [K[X]:K]$ , for every  $i, j$  we have  $\deg(f_i) + \deg(g_j) < [K(X):K]$ . Hence  $\nu(f_i g_j) = \nu(f_i) + \nu(g_j)$ . Denote by  $i_0$  (resp.  $j_0$ ) the smallest index such that  $\nu'(f) = \nu(f_{i_0} + i_0\gamma)$  (resp.  $\nu'(g) = \nu(g_{j_0} + j_0\gamma)$ ), and for  $i > m$  set  $f_i = g_i = 0$ . We have

$$\begin{aligned} \nu'(fg) &= \nu' \left( \sum_{j=0}^{2m} \left( \sum_{i=0}^j f_i g_{j-i} \right) \Phi^j \right) \geq \min_{0 \leq i \leq j \leq 2m} (\nu'(f_i g_{j-i}) + j\gamma) \\ &\geq \min_{0 \leq i \leq j \leq 2m} (\nu(f_i) + i\gamma + \nu(g_{j-i}) + (j-i)\gamma) \geq \nu'(f) + \nu'(g). \end{aligned}$$

For every  $i, j$ , let  $q_{i,j-i} = q(f_i, g_{j-i})$  and  $r_{i,j-i} = r(f_i, g_{j-i})$ .

$$\begin{aligned} fg &= \sum_{j=0}^{2m} \left( \sum_{i=0}^j q_{i,j-i} \Phi + r_{i,j-i} \right) \Phi^j = \\ &= \sum_{j=1}^{2m+1} \left( \sum_{i=0}^{j-1} q_{i,j-i-1} \right) \Phi^j + \sum_{j=0}^{2m} \left( \sum_{i=0}^j r_{i,j-i} \right) \Phi^j. \end{aligned}$$

Since  $\max(\deg(f_i), \deg(g_{j-i})) < d$ ,  $\nu$  is a  $p$ - $m$  valuation and  $\Phi$  is a  $w$ -key polynomial, we have:

$$\begin{aligned} \nu'(f_i) + \nu'(g_{j-i}) &= \nu(f_i) + \nu(g_{j-i}) = \nu(r_{i,j-i}) = \nu'(r_{i,j-i}) = \nu'(f_i g_{j-i}), \text{ and} \\ \nu(r_{i,j-i}) &\leq \nu(q_{i,j-i}) + \beta \leq \nu'(q_{i,j-i}) + \gamma. \\ \text{Furthermore, } \nu'(f) + \nu'(g) &= \nu(f_{i_0}) + i_0\gamma + \nu(g_{j_0}) + j_0\gamma \leq \\ &\leq \nu(f_i) + i\gamma + \nu(g_{j-i-1}) + (j-i-1)\gamma = \nu(f_i g_{j-i-1}) + (j-1)\gamma = \\ &= \nu(r_{i,j-i-1}) + (j-1)\gamma \leq \nu(q_{i,j-i-1}) + \beta + (j-1)\gamma = \nu(q_{i,j-i-1}) + j\gamma - (\gamma - \beta). \end{aligned}$$

$$\text{Consequently: } \nu' \left( fg - \sum_{j=0}^{2m} \left( \sum_{i=0}^j r(f_i, g_{j-i}) \right) \Phi^j \right) \geq \nu'(f) + \nu'(g) + (\gamma - \beta).$$

Now, let  $h_{i_0+j_0}$  be the coefficient of  $\Phi^{i_0+j_0}$  in the decomposition of  $fg$  by  $\Phi$ . We have

$$h_{i_0+j_0} = r_{0,i_0+j_0} + r_{1,i_0+j_0-1} + \dots + r_{i_0,j_0} + \dots + r_{i_0+j_0,0} + q_{0,i_0+j_0-1} + \dots + q_{i_0+j_0-1,0}.$$

By hypotheses, for  $0 \leq i \leq i_0 + j_0 - 1$  we have:

$$\begin{aligned} \nu(q_{i,i_0+j_0-1-i}) + (i_0+j_0)\gamma &\geq \nu(q_{i,i_0+j_0-1-i}) + \beta + (i_0+j_0-1)\gamma \geq \nu(f_i g_{i_0+j_0-1-i}) + (i_0+j_0-1)\gamma = \\ &= \nu(f_i) + i\gamma + \nu(g_{i_0+j_0-1-i}) + (i_0+j_0-1-i)\gamma. \\ \text{If } i \leq i_0-1, \text{ then } \nu(f_i) + i\gamma &> \nu'(f). \text{ Otherwise, } i_0+j_0-1-i \leq j_0-1, \text{ and } \nu(g_{i_0+j_0-1-i}) + (i_0+j_0-1-i)\gamma > \\ &> \nu'(g). \text{ In any case, } \nu(q_{i,i_0+j_0-1-i}) + (i_0+j_0)\gamma > \nu'(f) + \nu'(g). \text{ In the same way, for } 0 \leq i \leq i_0+j_0: \\ \nu(r_{i,i_0+j_0-i}) + (i_0+j_0)\gamma &= \nu(f_i) + i\gamma + \nu(g_{i_0+j_0-i}) + (i_0+j_0-i)\gamma \geq \nu'(f) + \nu'(g), \\ \text{and equality holds if, and only if, } i &= i_0 \text{ and } j = j_0. \text{ Hence the minimum is carried by a unique term, so} \\ \nu(h_{i_0+j_0}) &= \nu(f_{i_0} g_{j_0}) = \nu(f_{i_0}) + \nu(g_{j_0}). \text{ Consequently: } \nu'(fg) = \nu'(f) + \nu'(g). \end{aligned}$$

Assume that  $X$  is transcendental over  $K$ . For all nonzero  $f, g$  in  $K[X]$ , set  $\nu'(f/g) = \nu'(f) - \nu'(g)$ . Then, for every  $f, f', g, g'$  in  $K[X]$ , with  $g \neq 0 \neq g'$ , we have:

$$\nu \left( \frac{f}{g} \cdot \frac{f'}{g'} \right) = \nu \left( \frac{ff'}{gg'} \right) = \nu(f) + \nu(f') - \nu(g) - \nu(g') = \nu \left( \frac{f}{g} \right) + \nu \left( \frac{f'}{g'} \right), \text{ and}$$

$$\begin{aligned} \nu \left( \frac{f}{g} + \frac{f'}{g'} \right) &= \nu \left( \frac{fg' + f'g}{gg'} \right) = \nu(fg' + f'g) - \nu(g) - \nu(g') \geq \\ &\geq \min(\nu(fg'), \nu(f'g)) - \nu(g) - \nu(g') = \min \left( \nu \left( \frac{f}{g} \right), \nu \left( \frac{f'}{g'} \right) \right). \end{aligned}$$

Hence  $\nu'$  is a multiplicative valuation.

2) b) If  $f, g$  belong to  $K_{d-1}[\chi]$ , then  $f = f_0, g = g_0$  and  $\nu'(fg) = \nu'(r(f, g)) \leq \nu'(fg - r(f, g)) - (\gamma - \beta) = \nu'(q(f, g)) - (\gamma - \beta)$ . Hence  $\Phi$  is a w-key polynomial for  $\nu'$ . If  $\gamma > \beta$  or  $\beta$  is not a maximum, then the inequality is strict. If  $\beta$  is a maximum, say  $\beta = \nu(r_\Phi(f, g)) - \nu(q_\Phi(f, g))$ , then  $\nu(q_\Phi(f, g)\Phi) = \nu(q_\Phi(f, g)) + \beta = \nu(r_\Phi(f, g))$ . Hence  $\Phi$  is not a key polynomial. Since the family  $(\Phi^m)$  ( $dm < [K[\chi]:K]$ ) is separate over  $K_{d-1}[\chi]$ , by 3) of Proposition 3.12,  $\Phi$  is a ML key polynomial for  $\nu'$ .  $\square$

**Notations 3.17.** The  $K$ -module valuation  $\nu'$  defined in Proposition 3.16 will be denoted by  $\nu_{\Phi, \gamma}$ . We set  $\nu_\Phi = \nu_{\Phi, \nu(\Phi)}$ . If  $\Phi_1$  and  $\Phi_2$  are irreducible polynomials such that  $\deg(\Phi_1) < \deg(\Phi_2)$ , we denote by  $\nu_{\Phi_1, \Phi_2}$  the  $K$ -module valuation  $(\nu_{\Phi_1, \nu(\Phi_1)})_{\Phi_2, \nu(\Phi_2)}$ . By induction, for every irreducible polynomials  $\Phi_1, \dots, \Phi_n$ , with  $\deg(\Phi_1) < \dots < \deg(\Phi_n)$ , we define the  $K$ -module valuation  $\nu_{\Phi_1, \dots, \Phi_n}$ .

These valuations  $\nu_\Phi$  generalize the augmented valuations of S. MacLane ([ML 36a] and [ML 36b]).

**Remark 3.18.** If the degree of  $\Phi$  is 1, then for every  $f, g$  in  $K_{d-1}[\chi] = K$ , we have  $q(f, g) = 0$ . Hence the set  $\{\nu(r(f, g)) - \nu(q(f, g)) \mid f \in K_{d-1}[\chi], g \in K_{d-1}[\chi], \deg(f) + \deg(g) < [K[\chi]:K]\}$  is equal to  $\{-\infty\}$  and is bounded above by any element. Hence Proposition 3.16 shows that the Gauss valuations defined in Example 2.18 are p-m valuations. So, for every monic  $\Phi \in K[\chi]$ , the p-m valuation  $\nu_\Phi$  can be called a *generalized Gauss valuation*.

**Remark 3.19.** If  $\nu$  is a p-m valuation, then it follows from Proposition 3.16 that  $\Phi$  is a ML key polynomial for  $\nu$  if, and only if,  $\Phi$  is a w-key polynomial for  $\nu$  and  $\nu = \nu_\Phi$ .

**Proposition 3.20.** Let  $\Phi$  be a monic polynomial of degree  $d$  and  $\nu$  be a p-m valuation on  $K[\chi]$ . Then  $\Phi$  is a w-key polynomial for  $\nu$  if, and only if, there exists a p-m valuation  $\nu'$  of  $K[\chi]$ , such that its restriction to  $K_{d-1}[\chi]$  is equal to the restriction of  $\nu$ , and  $\nu'(\Phi) > \nu(\Phi)$ .

If this holds, then for every  $\gamma \geq \nu(\Phi)$  in an extension of  $\nu K[\chi]$ ,  $\nu_{\Phi, \gamma}$  is a p-m valuation of  $K[\chi]$  such that its restriction to  $K_{d-1}[\chi]$  is equal to the restriction of  $\nu$ ,  $\nu_{\Phi, \gamma}(\Phi) = \gamma$  and  $\Phi$  is a ML key polynomial for  $\nu_{\Phi, \gamma}$ .

*Proof.* Assume that  $\nu'$  is a p-m valuation of  $K[\chi]$  such that its restriction to  $K_{d-1}[\chi]$  is equal to the restriction of  $\nu$ , and  $\nu'(\Phi) > \nu(\Phi)$ . Let  $f, g$  in  $K_{d-1}[\chi]$  such that  $\deg(f) + \deg(g) < [K[\chi]:K]$ ,  $q = q(f, g)$  and  $r = r(f, g)$ . Without loss of generality we can assume that  $q \neq 0$ . We have  $\nu(fg) = \nu(f) + \nu(g) = \nu'(f) + \nu'(g) = \nu'(fg)$ . Therefore,  $\nu(q\Phi + r) = \nu'(q\Phi + r)$  is greater or equal to both of  $\min(\nu(q\Phi), \nu(r))$  and  $\min(\nu'(q\Phi), \nu'(r))$ , where  $\nu(q\Phi) < \nu'(q\Phi)$ . It follows that  $\nu(q\Phi) \neq \nu(r)$  or  $\nu'(q\Phi) \neq \nu'(r)$ , hence  $\nu(q\Phi + r) = \min(\nu(q\Phi), \nu(r))$  or  $\nu'(q\Phi + r) = \min(\nu'(q\Phi), \nu'(r))$ . In any case, since  $\nu(q\Phi) = \nu(q) + \nu(\Phi) < \nu'(q) + \nu'(\Phi) = \nu'(q\Phi)$  we see that this minimum is  $\nu(r)$  and that  $\nu(r) \leq \nu(q\Phi) - (\nu'(\Phi) - \nu(\Phi))$ . Now, if  $\Phi$  is a w-key polynomial for  $\nu$ , then the hypotheses of Proposition 3.16 2) are satisfied. Hence, the proof of the converse follows the proof of Proposition 3.16.

The remainder also follows from Proposition 3.16.  $\square$

**Notation 3.21.** If  $\nu$  and  $\nu'$  are  $K$ -module valuations on  $K(\chi)$ , then we set  $\nu \leq \nu'$  if for every  $f \in K[\chi]$  we have  $\nu(f) \leq \nu'(f)$ .

**Remark 3.22.** 1) By the definition of  $\nu_\Phi$ , for every  $K$ -module valuation  $\nu'$  such that the restrictions of  $\nu'$  and  $\nu_\Phi$  to  $K_{d-1}[\chi]$  are equal and  $\nu'(\Phi) = \nu_\Phi(\Phi)$ , we have  $\nu_\Phi \leq \nu'$ .

2) Assume that  $\chi$  is algebraic over  $K$ , that  $\nu$  is multiplicative, and  $\nu'(f) < \nu(f)$ . If  $\nu' \leq \nu$ , then  $\nu'(1/f) \leq \nu(1/f) = -\nu(f) < -\nu'(f)$ . Hence  $\nu'(1/f) \neq -\nu'(f)$ . It follows that  $\nu'$  is not multiplicative. Hence we cannot improve the conclusion that  $\nu'$  is partially multiplicative in Proposition 3.16.

**Remark 3.23.** In valuation theory, we say that  $\nu'$  is *finer* than  $\nu$  if  $\forall x \nu'(x) \geq 0 \Rightarrow \nu(x) \geq 0$  (see [R 68, p. 54]). Assume that  $\chi$  is algebraic over  $K$ . Then  $K(\chi) = K[\chi]$ , so, if  $\nu' \leq \nu$ , then  $\nu'$  is finer than  $\nu$ . Now, any two distinct extensions of a valuation to an algebraic extension are incomparable (see Corollaire 5, p. 158 in [R 68]). Therefore, this also proves that if  $\nu' \neq \nu$ , then  $\nu$  or  $\nu'$  is not multiplicative. In the case where  $\chi$  is transcendental over  $K$  and  $\nu, \nu'$  are valuations such that  $\nu' \leq \nu$ , then we cannot deduce that  $\nu$  and  $\nu'$  are comparable in the sense of Ribenboim. Indeed, assume that  $\nu(f) = \nu'(f) = \nu'(g) < \nu(g)$ . Then,  $\nu'(f/g) = 0 > \nu(f/g)$ . Assume that  $\nu'(f) < \nu(f) = \nu(g) = \nu(g)$ . Then  $\nu'(f/g) < 0 = \nu(f/g)$ .

**Proposition 3.24.** Let  $\nu$  be a p-m valuation on  $K[\chi]$ , and  $\Phi$  be a non constant monic polynomial in  $K[\chi]$  of degree  $d$ . Then  $\Phi$  is a w-key polynomial for  $\nu$  if, and only if, there exists a p-m valuation  $\nu' \leq \nu$  such that the restrictions of  $\nu$  and  $\nu'$  to  $K_{d-1}[\chi]$  are equal, and  $\Phi$  is a ML key polynomial for  $\nu'$ . Furthermore, we can take  $\nu' = \nu_\Phi$ .



*Proof.*  $\Leftarrow$ . If  $\Phi$  is a ML key polynomial for  $\nu'$ , then it is a w-key polynomial for  $\nu'$ . Now, by 3) of Remark 3.3, it is a w-key polynomial for  $\nu$ .

$\Rightarrow$ . By Proposition 3.20,  $\nu_\Phi$  is a p-m valuation such that  $\Phi$  is a ML key polynomial for  $\nu_\Phi$ . By construction, the restrictions of  $\nu$  and  $\nu_\Phi$  to  $K_{d-1}[\chi]$  are equal. By Remark 3.22 1) we have  $\nu_\Phi \leq \nu$ .  $\square$

**3.4. Bases generated by polynomials.** Let  $\Phi_1, \dots, \Phi_n, \dots$  be monic irreducible polynomials of  $K[\chi]$  of degrees  $d_1 = 1, d_2, \dots, d_n$ , respectively, where  $d_{n-1}$  divides  $d_n$  (if the family has a maximal element  $d_n$ , we set  $d_{n+1} = [K(\chi):K]$ ). We let  $\mathcal{B}$  be the family of the  $\Phi_1^{e_1} \cdots \Phi_n^{e_n}$ , where for  $0 \leq i \leq n-1$  we have:  $0 \leq e_i < \frac{d_{i+1}}{d_i}$ . Since the degree mapping is one-to-one from  $\mathcal{B}$  onto  $[0, [K(\chi):K]]$ ,  $\mathcal{B}$  is a basis of the  $K$ -module  $K[\chi]$ . Furthermore, for every  $m$  in  $\mathbb{N}$ ,  $\mathcal{B} \cap K_m[\chi]$  is a basis of the  $K$ -module  $K_m[\chi]$ . Now, we can define a basis even if some degree does not divide the following one. Indeed, in the case where we have only  $d_1 = 1 < d_2 < \dots < d_n < \dots$ , we require: for every  $n$ ,  $e_1 + e_2 d_2 + \dots + e_n d_n < d_{n+1}$ .

**Definition 3.25.** Let  $\mathcal{B}$  be a  $K$ -basis of  $K[\chi]$ . We say that  $\mathcal{B}$  is *generated by polynomials* if it is constructed in the above way. If so, then  $\Phi_1, \Phi_2, \dots, \Phi_n, \dots$  are called the *generating polynomials* for  $\mathcal{B}$ .

**Remark 3.26.** Let  $\nu$  be a  $K$ -module valuation on  $K(\chi)$ ,  $\Phi_1, \dots, \Phi_k, \dots$  be generating polynomials for a basis  $\mathcal{B}$ . For  $k \geq 1$ , fix  $\nu'(\Phi_k)$ . For every  $e_1, \dots, e_k$  with  $e_1 + e_2 d_2 + \dots + e_k d_k < d_{k+1}$ , let  $\nu'(\Phi_1^{e_1} \cdots \Phi_k^{e_k}) = e_1 \nu'(\Phi_1) + e_2 \nu'(\Phi_2) + \dots + e_k \nu'(\Phi_k)$  and for every pairwise distinct  $y_1, \dots, y_k$  in  $\mathcal{B}$ ,  $x_1, \dots, x_k$  in  $K$  set  $\nu'(x_1 y_1 + \dots + x_k y_k) = \min_{1 \leq i \leq k} \nu(x_i) \nu'(y_i)$ . Then  $\nu'$  is a separate  $K$ -module valuation. If for every  $k \geq 1$  we have  $\nu'(\Phi_k) = \nu(\Phi_k)$ , then the  $K$ -module valuation  $\nu'$  defined above is the  $K$ -module valuation  $\nu_{\Phi_{d_1}, \dots, \Phi_{d_k}}$  defined in Notations 3.17.

### 3.5. Properties of key degrees.

**Proposition 3.27.** Let  $d$  be an integer and  $\nu$  be a p-m valuation on  $K[\chi]$ .

- 1) Let  $\Phi$  be a w-key polynomial of degree  $d$ . Then every monic polynomial  $\Phi'$  of degree  $d$ , such that  $\nu(\Phi') > \nu(\Phi)$ , is a key polynomial. In particular  $d$  is a key degree.
- 2) Assume that  $d$  is a separate key degree and let  $\Phi$  be a key polynomial of degree  $d$ . Then every monic polynomial  $\Phi'$  of degree  $d$ , such that  $\nu(\Phi') \geq \nu(\Phi)$ , is a key polynomial. In particular, any monic polynomial  $\Phi'$  of degree  $d$ , such that  $\nu(\Phi')$  is maximal in  $\nu(\chi^d - K_{d-1}[\chi])$ , is a key polynomial.
- 3) Assume that  $\Phi$  is a w-key polynomial of degree  $d$  and that the set  $\nu(\chi^d - K_{d-1}[\chi])$  has a greatest element. Then every w-key polynomial of degree  $d$  has valuation  $\nu(\Phi)$ , and  $\nu(\Phi)$  is maximal in  $\nu(\chi^d - K_{d-1}[\chi])$ .

This proposition is a consequence of the following lemma.

**Lemma 3.28.** Let  $d$  be a positive integer,  $\nu$  be a p-m valuation on  $K[\chi]$ ,  $\Phi$  and  $\Phi'$  be monic polynomials of degree  $d$  such that  $\Phi$  is a w-key polynomial for  $\nu$ .

- 1) If  $\nu(\Phi') > \nu(\Phi)$ , then  $\Phi'$  is a w-key polynomial for  $\nu_\Phi$ , a key polynomial for  $\nu$ , and  $\nu_\Phi \leq \nu_{\Phi'}$ .
- 2) If  $\nu(\Phi') = \nu(\Phi)$  and  $\Phi$  is a key polynomial for  $\nu$ , then  $\Phi'$  is a key polynomial for both of  $\nu$  and  $\nu_\Phi$ . Furthermore,  $\nu_\Phi = \nu_{\Phi'}$ .
- 3) If  $\nu(\Phi') = \nu(\Phi)$  and  $\Phi, \Phi'$  are w-key polynomials for  $\nu$ , then  $\nu_\Phi = \nu_{\Phi'}$ .

*Proof.* Set  $h = \Phi' - \Phi$ . Assume that  $\nu(\Phi') \geq \nu(\Phi)$ . Then we have  $\nu(h) \geq \nu(\Phi)$ . Furthermore,  $\Phi$  and  $\Phi'$  are monic polynomials, so the degree of  $h$  is lower than  $d$ . Now,  $\Phi' = \Phi + h$ , hence  $\nu_\Phi(\Phi') = \min(\nu(\Phi), \nu(h)) = \nu(\Phi)$ . Let  $f, g$  in  $K_{d-1}[\chi]$  with  $\deg(f) + \deg(g) < [K[\chi]:K]$ . We have:  $\nu_\Phi(fg) = \nu_\Phi(f) + \nu_\Phi(g) = \nu(f) + \nu(g) = \nu(fg)$ . Let  $r' = r_{\Phi'}(f, g)$ ,  $q' = q_{\Phi'}(f, g)$ .

1) We assume that  $\nu(\Phi') > \nu(\Phi)$ . Hence  $\nu(f) = \nu(\Phi)$ . If  $\nu_\Phi(q'\Phi') < \nu_\Phi(r') (= \nu(r'))$ , then  $\nu_\Phi(fg) = \nu_\Phi(q'\Phi' + r') = \nu_\Phi(q'\Phi') = \nu(q'\Phi) = \nu_\Phi(q'\Phi)$ . Now,  $\nu(q'\Phi') > \nu(q'\Phi) = \nu(fg)$  and  $\nu(r') > \nu_\Phi(q'\Phi') = \nu_\Phi(fg) = \nu(fg)$ . Hence  $\nu(fg) \geq \min(\nu(q'\Phi'), \nu(r')) > \nu(fg)$ : a contradiction. Therefore,  $\nu_\Phi(q'\Phi') \geq \nu_\Phi(r')$ . Now,  $\nu(r') = \nu_\Phi(r')$  and  $\nu_\Phi(q'\Phi') = \nu(q'\Phi) < \nu'(q'\Phi')$ . Hence  $\Phi'$  is a key polynomial for  $\nu$ . Since  $\nu_\Phi(fg) = \nu(fg) = \nu(r') = \nu_\Phi(r')$ ,  $\Phi'$  is a w-key polynomial for  $\nu_\Phi$ . Furthermore,  $\nu_\Phi$  and  $\nu_{\Phi'}$  are equal on  $K_{d-1}[\chi]$ , and  $\nu_{\Phi'}(\Phi) = \nu_{\Phi'}(\Phi' - h) = \min(\nu_{\Phi'}(\Phi'), \nu_{\Phi'}(h)) = \min(\nu(\Phi'), \nu(h)) = \nu(h) = \nu(\Phi) = \nu_\Phi(\Phi)$ . By Remark 3.22, we have  $\nu_\Phi \leq \nu_{\Phi'}$ .

2) We assume that  $\nu(\Phi') = \nu(\Phi)$  and  $\Phi$  is a key polynomial for  $\nu$ . Let  $q_1 = q_\Phi(q', h)$  and  $r_1 = r_\Phi(q', h)$ . Since  $\Phi$  is a key polynomial for  $\nu$ , we have  $\nu(q_1\Phi) > \nu(r_1) = \nu(q'h) \geq \nu(q'\Phi)$ . Hence  $\nu(q') < \nu(q_1)$  and  $\nu(q' + q_1) = \nu(q')$ . We have:  $fg = q'\Phi' + r' = q'\Phi + q'h + r' = (q' + q_1)\Phi + r' + r_1$ , hence  $q' + q_1 = q_\Phi(f, g)$ ,  $r' + r_1 = r_\Phi(f, g)$  and  $\nu(fg) = \nu(r' + r_1) < \nu((q' + q_1)\Phi) = \nu(q'\Phi) \leq \nu(r_1)$ . It follows:  $\nu(r' + r_1) = \nu(r')$ , so  $\nu(fg) = \nu(r') < \nu(q'\Phi) = \nu(q'\Phi')$ . This proves that  $\Phi'$  is a key polynomial for  $\nu$ . Now,  $\nu_\Phi(fg) = \nu(fg) = \nu(r') = \nu_\Phi(r')$  and  $\nu_\Phi(q'\Phi') = \nu_\Phi(q'\Phi) = \nu(q'\Phi) > \nu(r')$ . Hence  $\Phi'$  is a key polynomial for  $\nu_\Phi$ . In the same way as in 1), we have:  $\nu_\Phi \leq \nu_{\Phi'}$ . Now, since  $\Phi'$  is a key polynomial, we have in a symmetric way:  $\nu_{\Phi'} \leq \nu_\Phi$ .

3) We assume that  $\nu(\Phi') = \nu(\Phi)$  and  $\Phi, \Phi'$  are w-key polynomials for  $\nu$ . We have:  $\Phi' = \Phi + h$ , with

$\nu(h) \geq \nu(\Phi)$  and  $\deg(h) < d$ . Hence  $\nu_{\Phi}(\Phi') = \min(\nu(\Phi), \nu(h)) = \nu(\Phi) = \nu(\Phi') = \nu_{\Phi'}(\Phi')$ . So by Remark 3.22 we have:  $\nu_{\Phi'} \leq \nu_{\Phi}$ . In the same way,  $\nu_{\Phi} \leq \nu_{\Phi'}$ , hence  $\nu_{\Phi} = \nu_{\Phi'}$ .  $\square$

**Lemma 3.29.** *Let  $\nu$  be a  $p$ - $m$  valuation on  $K[X]$  and  $\Phi$  be a monic polynomial of degree  $d$ . Then,  $\nu = \nu_{\Phi}$  on  $K_d[X]$  if, and only if,  $\nu(\Phi) = \max \nu(\chi^d - K_{d-1}[X])$ .*

*Proof.* By the definition of  $\nu_{\Phi}$ ,  $\nu$  and  $\nu_{\Phi}$  are equal on  $K_{d-1}[X]$ . Hence, we can consider polynomials of degree  $d$ . So, we let  $f$  be a polynomial of degree  $d$ . Without loss of generality we can assume that  $f$  is a monic polynomial. Hence  $f - \Phi$  has degree less than  $\Phi$ . Assume that  $\nu(f) = \nu_{\Phi}(f)$ . Then  $\nu(f) = \nu_{\Phi}(\Phi + f - \Phi) = \min(\nu(\Phi), \nu(f - \Phi)) \leq \nu(\Phi)$ . This proves that  $\nu(\Phi)$  is the maximum of  $\nu(\chi^d - K_{d-1}[X])$ . Conversely, assume that  $\nu(\Phi) = \max \nu(\chi^d - K_{d-1}[X])$ . We have  $\nu(f - \Phi) \geq \min(\nu(f), \nu(\Phi)) = \nu(f)$ . Therefore,  $\nu(f) \geq \nu_{\Phi}(f) = \nu_{\Phi}(\Phi + f - \Phi) = \min(\nu(\Phi), \nu(f - \Phi)) \geq \nu(f)$ . Hence  $\nu_{\Phi}(f) = \nu(f)$ .  $\square$

**Remark 3.30.** The valuation of a key polynomial is not necessarily maximal. Indeed, we saw in Remark 3.3 1) that every monic polynomial of degree 1 is a key polynomial. This holds whether 1 is a separate key degree or not.

The following two lemmas give useful criteria for being a  $w$ -key polynomial or a key polynomial.

**Lemma 3.31.** *Let  $\nu, \nu'$  be  $p$ - $m$  valuations on  $K(X)$ , and  $\Phi$  be a monic polynomial of degree  $d \geq 1$ . Assume that their restrictions to  $K_{d-1}[X]$  are equal, that  $\nu' \leq \nu$  and  $\nu'(\Phi) < \nu(\Phi)$ .*

- 1) *Let  $f \in K[X]$  and  $f = q\Phi + r$  be the euclidean division of  $f$  by  $\Phi$ . Then  $\nu'(f) < \nu'(r) \Leftrightarrow \nu'(f) < \nu(f)$  and  $\nu'(f) = \nu'(r) \Leftrightarrow \nu'(f) = \nu(f)$ .*
- 2)  *$\Phi$  is a  $w$ -key polynomial for  $\nu'$  and a key polynomial for  $\nu$ .*
- 3)  *$\nu' = \nu'_{\Phi} = \nu_{\Phi, \nu'(\Phi)}$ .*

*Proof.* 1) We have  $\nu'(r) = \nu(r)$  and  $\nu'(q\Phi) < \nu(q\Phi)$ . Assume that  $\nu'(r) \leq \nu'(q\Phi)$ . Then  $\nu(f) = \min(\nu(q\Phi), \nu(r)) = \nu(r)$ . Furthermore,  $\nu(f) \geq \nu'(f) \geq \min(\nu'(q\Phi), \nu(r)) \geq \nu(r) = \nu(f)$ . Hence  $\nu'(f) = \nu(f) = \nu(r)$ . Assume that  $\nu'(q\Phi) < \nu(r) < \nu(q\Phi)$ . Hence  $\nu'(f) = \min(\nu'(q\Phi), \nu(r)) = \nu'(q\Phi) < \nu'(r) = \min(\nu(q\Phi), \nu(r)) = \nu(f)$ . Assume that  $\nu(q\Phi) \leq \nu(r)$ . Then  $\nu'(f) = \min(\nu'(q\Phi), \nu(r)) = \nu'(q\Phi) < \nu(q\Phi) = \min(\nu(q\Phi), \nu(r)) \leq \nu(f)$ , and  $\nu'(f) < \nu'(r)$ .

2) Let  $f, g$  in  $K_{d-1}[X]$  with  $\deg(f) + \deg(g) < [K[X]:K]$ . Since  $\deg(f) < \deg(\Phi)$  and  $\deg(g) < \deg(\Phi)$ , we have:  $\nu'(f) = \nu(f)$  and  $\nu'(g) = \nu(g)$ , therefore:  $\nu'(fg) = \nu'(f) + \nu'(g) = \nu(f) + \nu(g) = \nu(fg)$ . By 1) this implies  $\nu'(fg) = \nu'(r(f, g))$  and  $\Phi$  is a  $w$ -key polynomial for  $\nu'$ . Since the restrictions of  $\nu$  and  $\nu'$  to  $K_{d-1}[X]$  are equal, it follows that  $\Phi$  is also a  $w$ -key polynomial for  $\nu$ . Now, for  $f, g$  in  $K_{d-1}[X]$  we have  $\nu(r(f, g)) = \nu'(r(f, g)) \leq \nu'(q(f, g) \cdot \Phi) < \nu(q(f, g) \cdot \Phi)$ . Hence  $\Phi$  is a key polynomial for  $\nu$ .

3) First we prove that for every  $f, g$  in  $K_{d-1}[X]$  with  $\deg(f) + \deg(g) < [K[X]:K]$  and  $h \in K[X]$  such that  $\deg(h) < [K[X]:K] - d$  we have:  $\nu'(fg + h\Phi) = \min(\nu'(fg), \nu'(h\Phi))$ . If  $\nu'(fg) \neq \nu'(h\Phi)$ , then the result is trivial. Assume that  $\nu'(fg) = \nu'(h\Phi)$ , and let  $q = q(f, g)$ ,  $r = r(f, g)$ . Since  $\Phi$  is a  $w$ -key polynomial for  $\nu'$ , we have  $\nu'(fg) = \nu'(r)$ . Hence  $\nu'(fg + h\Phi) \geq \nu'(r)$ . Note that  $fg + h\Phi = (q + h)\Phi + r$ , hence by 1) it follows:  $\nu'(fg + h\Phi) = \nu(fg + h\Phi)$ . Now,  $\nu(fg) = \nu'(fg) = \nu'(h\Phi) < \nu(h\Phi)$ , hence  $\nu(fg + h\Phi) = \nu(fg) = \nu'(fg) = \min(\nu'(fg), \nu'(h\Phi))$ . By 2)  $\Rightarrow$  3) of Proposition 3.12 and Remark 3.19, we have  $\nu' = \nu'_{\Phi}$ . Now, since  $\nu'$  and  $\nu$  coincide on  $K_d[X]$  we have  $\nu'_{\Phi} = \nu_{\Phi, \nu'(\Phi)}$ .  $\square$

**Remark 3.32.** It follows from Lemma 3.31 that if  $\nu$  is a valuation on  $K(X)$ , then every  $p$ - $m$  valuation  $\nu' \leq \nu$ ,  $\nu' \neq \nu$ , which coincide on  $K$  with  $\nu$ , can be written as  $\nu' = \nu_{\Phi, \nu'(\Phi)}$  where  $\Phi$  is a monic polynomial of minimal degree such that  $\nu'(\Phi) < \nu(\Phi)$ .

**Lemma 3.33.** *Let  $\nu$  be a  $p$ - $m$  valuation on  $K(X)$ ,  $\Phi$  be a  $w$ -key polynomial for  $\nu$  and  $\Phi'$  be a monic polynomial such that  $d = \deg(\Phi) < \deg(\Phi') = d'$ .*

- 1) *If  $\nu(\Phi') = \nu_{\Phi}(\Phi')$ , then  $\Phi'$  is not a key polynomial for  $\nu$ .*
- 2) *Assume that  $\nu_{\Phi} = \nu$  on  $K_{d'-1}[X]$ . Then,  $\Phi'$  is a key polynomial for  $\nu$  if, and only if,  $\nu(\Phi') > \nu_{\Phi}(\Phi')$ .*

*Proof.* 1) Write  $\Phi'$  as  $\Phi' = f_m\Phi^m + \dots + f_1\Phi + f_0$ , with  $f_0, \dots, f_m$  in  $K_{d-1}[X]$ . By the definition of  $\nu_{\Phi}$ , we have  $\nu(\Phi') = \nu_{\Phi}(\Phi') = \min(f_0, f_1\Phi, \dots, f_m\Phi^m)$ . Let  $f = f_m\Phi^{m-1} + \dots + f_1$  and  $g = \Phi$ . Then  $q_{\Phi'}(f, g) = 1$ ,  $r_{\Phi'}(f, g) = -f_0$ , and  $\nu(q_{\Phi'}(f, g)\Phi') = \nu(\Phi') = \min(\nu(r_{\Phi'}(f, g)), \nu(fg)) \leq \nu(fg)$ . This proves that  $\Phi'$  is not a key polynomial.

2)  $\Leftarrow$  follows from Lemma 3.31,  $\Rightarrow$  follows from 1).  $\square$

**Proposition 3.34.** *Let  $d$  be a positive integer,  $\nu$  be a  $p$ - $m$  valuation on  $K[X]$ .*

- 1) *If  $\nu = \nu_{\Phi}$  for some monic polynomial  $\Phi$ , then there is no key degree greater than  $d$ .*
- 2) *If  $d$  is a separate key degree such that there is no key degree greater than  $d$ , and  $\Phi$  is a  $w$ -key polynomial of degree  $d$ , with  $\nu(\Phi) = \max(\nu(\chi^d - K_{d-1}[X]))$ , then  $\nu = \nu_{\Phi}$ .*

*Proof.* 1) Follows from Lemma 3.33.

2) By Lemma 3.29,  $\nu$  and  $\nu_{\Phi}$  coincide on  $K_d[X]$ . Now, by Lemma 3.33,  $\nu = \nu_{\Phi}$  on  $K[X]$ .  $\square$

**Remark 3.35.** Let  $\nu$  be a multiplicative valuation on  $K(X)$ . If the irrational rank of  $\nu K(X)$  over  $\nu K$  is 1, or the transcendence degree of  $K(X)_\nu$  over  $K_\nu$  is 1 (in other words, if  $X$  is transcendental and Abhyankar's inequality is an equality), then there is a finite number of key degrees, and  $\nu = \nu_\Phi$  for some w-key polynomial  $\Phi$ .

*Proof.* Note that in this case  $X$  is transcendental. Abhyankar's inequality states that the transcendence degree of  $K(X)|K$  is at least equal to the product of the transcendence degree of  $(K(X))_\nu|K_\nu$  by the irrational rank of  $\nu(K[X])|\nu K$ . Assume that the irrational rank of  $\nu K(X)$  over  $\nu K$  is 1. Let  $d$  be the smallest integer such that there exists a monic polynomial  $\Phi$  of degree  $d$  such that  $\nu(\Phi)$  is not rational over  $\nu K$ . If  $d = 1$ , then we know that  $\Phi$  is a key polynomial. Otherwise, clearly,  $\nu(\Phi) \neq \nu(X^d) = d\nu(X)$ . If  $\nu(\Phi) < \nu(X^d)$ , then  $\nu(X^d - \Phi) = \nu(\Phi)$ , with  $\deg(X^d - \Phi) < d$ : a contradiction (since  $d$  is minimal). Hence  $\nu(\Phi) > \nu(X^d)$ . Then  $\nu(\Phi) = \max \nu(X^d - K_{d-1}[X])$ . Let  $f, g$  in  $K_{d-1}[X]$  with  $\deg(f) + \deg(g) < [K[X]:K]$  and  $q = q(f, g)$ ,  $r = r(f, g)$ . Since  $d$  is minimal, we have:  $\nu(r) \neq \nu(q\Phi)$  and  $\nu(q\Phi) \neq \nu(f) + \nu(g) = \nu(fg)$ . It follows that  $\nu(fg) = \nu(r) < \nu(q\Phi)$  (the other cases lead to a contradiction). So  $\Phi$  is a key polynomial for  $\nu$ . Now, the elements  $\nu(\Phi^k) = k\nu(\Phi)$  are pairwise non-congruent modulo  $\nu K_{d-1}[X]$ . In the same way as in Proposition 2.13, this implies that the family  $(\Phi^k)$  is separate over  $K_{d-1}[X]$ . Therefore,  $\nu = \nu_\Phi$ . Now, by 1) of Proposition 3.34, if  $d' > d$ , then  $d$  is not a key degree.

Now, assume that the transcendence degree of  $K(X)_\nu$  over  $K_\nu$  is 1. Let  $\Phi$  be a polynomial such that  $\nu(\Phi) = 0$ . If  $\text{in}_\nu(\Phi)$  is algebraic over  $G_\nu(K)$ , then by Lemma 2.21  $\text{in}_\nu(\Phi)$  is algebraic over  $K_\nu$  (the converse is trivial). Consequently,  $\text{in}_\nu(\Phi)$  is transcendental over  $K_\nu$  if, and only if, it is transcendental over  $G_\nu(K)$ . In particular, if the transcendence degree of  $K(X)$  over  $K$  is 1, then there is  $\Phi$  in  $K[X]$  such that  $\text{in}_\nu(\Phi)$  is transcendental over  $G_\nu(K)$ . Let  $d$  be the smallest integer such that there exists a polynomial  $\Phi$  of degree  $d$  such that  $\text{in}_\nu(\Phi)$  is transcendental over  $G_\nu(K)$ . Without loss of generality we can assume that  $\Phi$  is a monic polynomial. Let  $f, g$  in  $K_{d-1}[X]$ ,  $q = q_\Phi(f, g)$ ,  $r = r_\Phi(f, g)$ . We have:  $\text{in}_\nu(fg) = \text{in}_\nu(q\Phi) \Leftrightarrow 0 = \text{in}_\nu(fg) - \text{in}_\nu(q\Phi) \Leftrightarrow \nu(fg - q\Phi) > \min(\nu(fg), \nu(q\Phi)) \Leftrightarrow \nu(r) > \min(\nu(fg), \nu(q\Phi))$ . Now,  $\text{in}_\nu(fg) = \text{in}_\nu(f)\text{in}_\nu(g)$  is algebraic over  $G_\nu(K)$ . Hence  $\text{in}_\nu(fg) \neq \text{in}_\nu(q\Phi)$ , which is transcendental over  $G_\nu(K)$ . It follows that  $\nu(r) \leq \min(\nu(fg), \nu(q\Phi))$ . This proves that  $\Phi$  is a w-key polynomial. Let  $f_0, \dots, f_n$  in  $K_{d-1}[X]$ . By hypothesis,  $\text{in}_\nu(f_0), \dots, \text{in}_\nu(f_n)$  are algebraic over  $G_\nu(K)$ . Hence  $\text{in}_\nu(f_n)\text{in}_\nu(\Phi)^n + \dots + \text{in}_\nu(f_0) \neq 0$ . It follows that  $\nu(f_0 + \dots + f_n\Phi^n) = \min(\nu(f_0), \dots, \nu(f_n\Phi^n))$ . This proves that the sequence  $(\Phi^n)$  is separate over  $K_{d-1}[X]$ . Therefore,  $\nu = \nu_\Phi$ . By 1) of Proposition 3.34, if  $d' > d$ , then  $d$  is not a key degree.  $\square$

The following two theorems characterize the successor of a key degree. They will be useful for proving the equivalence of the definitions of key polynomials and abstract key polynomials.

**Theorem 3.36.** Let  $\nu$  be a  $p$ - $m$  valuation on  $K(X)$ ,  $d$  be an immediate key degree, and let  $(\Phi_i)$  be a sequence of key polynomials of degree  $d$  such that the sequence  $(\nu(\Phi_i))$  is increasing and cofinal in  $\nu(X^d - K_{d-1}[X])$ . Then.

- 1) The sequence  $(\nu_{\Phi_i})$  converges to  $\nu$  on  $K_d[X]$  in the sense that for every  $f \in K_d[X]$  the sequence  $(\nu_{\Phi_i}(f))$  is eventually equal to  $\nu(f)$ .
- 2) Let  $d'$  be the smallest degree (if any) such that there exists a monic polynomial  $\Phi'$  of degree  $d'$  satisfying  $\nu_{\Phi_i}(\Phi') < \nu(\Phi')$  for every  $i$ . Then,  $\Phi'$  is a key polynomial and  $d'$  is the next key degree.
- 3) The extension  $(K_{d'-1}[X]|K_{d-1}[X], \nu)$  is immediate.

*Proof.* 1) The family  $(\nu_{\Phi_i})$  is increasing, and for every  $f \in K[X]$  we have  $\nu_{\Phi_i}(f) \leq \nu(f)$ . Assume that  $f$  is a polynomial of degree  $d$ . Without loss of generality we can assume that  $f$  is monic. Since the sequence  $(\nu(\Phi_i))$  is cofinal in  $\nu(X^d - K_{d-1}[X])$ , there is an  $i$  such that  $\nu(\Phi_i) > \nu(f)$ . Now, for every  $i$  such that  $\nu(\Phi_i) > \nu(f)$  we have  $\nu(f - \Phi_i) = \nu(f)$ , and  $\nu_{\Phi_i}(f) = \min(\nu(\Phi_i), \nu(f - \Phi_i)) = \nu(f)$ .

2) Let  $f, g$  in  $K_{d'-1}[X]$  with  $\deg(f) + \deg(g) < [K[X]:K]$  and  $q = q_{\Phi'}(f, g)$ ,  $r = r_{\Phi'}(f, g)$ . By hypothesis there exists  $i$  such that  $\nu_{\Phi_i}(f) = \nu(f)$ ,  $\nu_{\Phi_i}(g) = \nu(g)$ ,  $\nu_{\Phi_i}(q) = \nu(q)$  and  $\nu_{\Phi_i}(r) = \nu(r)$ . Then  $\nu(q\Phi' + r) = \nu(fg) = \nu(f) + \nu(g) = \nu_{\Phi_i}(f) + \nu_{\Phi_i}(g) = \nu_{\Phi_i}(fg) = \nu_{\Phi_i}(q\Phi' + r)$ , with  $\nu(q\Phi') > \nu_{\Phi_i}(q\Phi')$ . Assume that  $\nu(q\Phi') \leq \nu(r)$ . Hence  $\nu_{\Phi_i}(q\Phi') < \nu(r) = \nu_{\Phi_i}(r)$ , and  $\nu(fg) = \nu_{\Phi_i}(fg) = \nu_{\Phi_i}(q\Phi') < \nu(q\Phi') = \min(\nu(q\Phi'), \nu(r)) \leq \nu(fg)$ : a contradiction. Hence  $\nu(q\Phi') > \nu(r)$ , which proves that  $\Phi'$  is a key polynomial for  $\nu$ . Let  $f$  be a monic polynomial of degree  $d'' < d'$ . Then, there exists  $i$  such that  $\nu(f) = \nu_{\Phi_i}(f)$ . By Lemma 3.33 1),  $f$  is not a key polynomial.

3) Let  $f$  be a monic polynomial of degree  $n$ ,  $d < n \leq d' - 1$ . We show that  $\nu(f)$  is not the maximum of  $\nu(X^n - K_{n-1}[X])$ . It will follow by Proposition 1.8 that the extension  $(K_{d'-1}[X]|K_{d-1}[X], \nu)$  is immediate. Let  $\Phi_i$  be a w-key polynomial of degree  $d$  such that  $\nu(f) = \nu_{\Phi_i}(f)$ , and let  $f$  be written as  $f = f_k\Phi_i^k + \dots + f_1\Phi_i + f_0$ , where  $f_0, f_1, \dots, f_k$  belong to  $K_{d-1}[X]$ ,  $f_k$  is monic, and  $\deg(f_k) + kd = n$ . Then  $\nu(f) = \min(\nu(f_k\Phi_i^k), \dots, \nu(f_1\Phi_i), \nu(f_0))$ . Assume that  $\nu(f_0) > \nu(f)$ . Then  $\nu(f) = \min(\nu(f_k\Phi_i^k), \dots, \nu(f_1\Phi_i))$ . Let  $\Phi_j$  be a w-key polynomial of degree  $d$  such that  $\nu(\Phi_j) > \nu(\Phi_i)$ , and set  $g = f_k\Phi_j^k + \dots + f_1\Phi_j$ . Then  $g$  is a monic polynomial of degree  $n$ , and  $\nu(g) \geq \nu_{\Phi_j}(g) = \min(\nu(f_k\Phi_j^k), \dots, \nu(f_1\Phi_j)) > \nu(f)$ . Now, assume that  $\nu(f_0) = \nu(f)$ . Then  $\nu(f - f_0) \geq \nu(f_0)$ . If  $\nu(f - f_0) > \nu(f_0)$ , then we can let  $g = f - f_0$ . If

$\nu(f - f_0) = \nu(f_0) = \nu(f)$ , then we let  $\Phi_j$  be a w-key polynomial of degree  $d$  such that  $\nu(\Phi_j) > \nu(\Phi_i)$ , and  $g = f_k \Phi_j^k + \dots + f_1 \Phi_j$ . Then  $\nu(g) \geq \nu_{\Phi_j}(g) = \min(\nu(f_k \Phi_j^k), \dots, \nu(f_1 \Phi_j)) > \min(\nu(f_k \Phi_i^k), \dots, \nu(f_1 \Phi_i)) = \nu(f)$ .  $\square$

**Theorem 3.37.** *Let  $\nu$  be a  $p$ - $m$  valuation on  $K(X)$ ,  $d$  be a separate key degree,  $\Phi$  be a monic polynomial of degree  $d$  such that  $\nu(\Phi)$  is the maximum of  $\nu(X^d - K_{d-1}[X])$ . Let  $d'$  be the next key degree if it exists or  $d' = [K[X]:K]$  otherwise.*

- 1)  $\Phi$  is a key polynomial, and if  $d' < [K[X]:K]$ , then  $d'$  is the smallest degree such that there exists a monic polynomial  $\Phi'$  of degree  $d'$  with  $\nu_{\Phi}(\Phi') < \nu(\Phi')$ .
- 2) The restrictions of  $\nu$  and  $\nu_{\Phi}$  to  $K_{d'-1}[X]$  are equal.
- 3) The fraction  $d'/d$  is equal to  $[(K_{d'-1}[X])_{\nu} : (K_{d-1}[X])_{\nu}] \cdot (\nu(K_{d'-1}[X]) : \nu(K_{d-1}[X]))$  (in particular  $d \leq [K[X]:K]/2$ ).
- 4) If  $(K_{d-1}[X]|K, \nu)$  is separate, then  $(K_{d'-1}[X]|K, \nu)$  is separate.

*Proof.* 1) Since  $\nu(\Phi)$  is the maximum of  $\nu(X^d - K_{d-1}[X])$ , by (2) of Proposition 3.27,  $\Phi$  is a key polynomial. By Lemma 3.33, if  $d' < [K[X]:K]$ , then  $d'$  is the smallest degree such that there exists a monic polynomial  $\Phi'$  of degree  $d'$  with  $\nu_{\Phi}(\Phi') < \nu(\Phi')$ .

2) Since  $\nu_{\Phi} \leq \nu$ , by 1) the restrictions of  $\nu$  and  $\nu_{\Phi}$  to  $K_{d'-1}[X]$  are equal.

3) By Lemma 3.5  $in_{\nu}(\Phi) \notin G_{\nu}(K(X))$ . If  $in_{\nu}(\Phi)$  is transcendental over the graded algebra generated by  $G_{\nu}(K_{d-1}[X])$ , then  $X$  is transcendental over  $K$ . It follows that  $[K(X):K]$  is infinite, and in the same way as in the proof of Remark 3.35 the family  $(\Phi^n)$  is separate over  $K_{d-1}[X]$  and  $\nu = \nu_{\Phi}$ . So,  $d' = \infty$ . The dimension of the  $G_{\nu}(K)$ -module  $G_{\nu}(K(X))$  is also infinite, so  $[(K_{d'-1}[X])_{\nu} : (K_{d-1}[X])_{\nu}] \cdot (\nu(K_{d'-1}[X]) : \nu(K_{d-1}[X]))$  is infinite. Assume that  $in_{\nu}(\Phi)$  is algebraic over  $G_{\nu}(K_{d-1}[X])$ , and that  $G_{\nu}(K_{d-1}[X])$  is a graded algebra (by Remark 3.9 this holds if  $d \leq [K[X]:K]/2$ ). Let  $X^n + in_{\nu}(f_{n-1})X^{n-1} + \dots + in_{\nu}(f_0)$  be its irreducible polynomial, with  $f_0, \dots, f_{n-1}$  in  $K_{d-1}[X]$  and  $\nu(f_0) = \dots = \nu(f_{n-1}\Phi^{n-1}) = \nu(\Phi^n)$ , as in Lemma 2.21. Note that  $dn < [K[X]:K]$ , since, in  $K[X]$ ,  $\Phi^n$  is a polynomial of degree less than  $[K[X]:K]$ . By minimality of  $n$ , this implies that  $in_{\nu}(1), in_{\nu}(\Phi), \dots, in_{\nu}(\Phi^{n-1})$  are linearly independent over  $G_{\nu}(K_{d-1}[X])$ . We saw in Subsection 2.3 that this is equivalent to saying that the sequence  $1, \Phi, \dots, \Phi^{n-1}$  is separate over  $K_{d-1}[X]$ . Since  $\nu_{\Phi}(\Phi^n + f_{n-1}\Phi^{n-1} + \dots + f_0) = \min(\nu(\Phi^n), \nu(f_{n-1}\Phi^{n-1}), \dots, \nu(f_0)) < \nu(\Phi^n + f_{n-1}\Phi^{n-1} + \dots + f_0)$ , we have  $d' = dn$ . This also shows that the group  $G_{\nu}(K_{d'-1}[X])$  is equal to  $G_{\nu}(K_{d-1}[X])(in_{\nu}(\Phi))$ , so it is a subalgebra of  $G_{\nu}(K(X))$ . In particular,  $(K_{d-1}[X])_{\nu}$  is a subfield of  $(K[X])_{\nu}$  and  $\nu K_{d-1}[X]$  is a subgroup of  $\nu(X)$ . Now, this also proves that  $(in_{\nu}(1), in_{\nu}(\Phi), \dots, in_{\nu}(\Phi^{n-1}))$  is a basis of the  $G_{\nu}(K_{d-1}[X])$ -module  $G_{\nu}(K_{d'-1}[X])$ . Hence its dimension is  $n$ . Since this dimension is also equal to  $[(K_{d'-1}[X])_{\nu} : (K_{d-1}[X])_{\nu}] \cdot (\nu(K_{d'-1}[X]) : \nu(K_{d-1}[X]))$ , the result follows.

If  $[K[X]:K]$  is infinite, then by Remark 3.9 for, every key degree  $d$ ,  $G_{\nu}(K_{d-1}[X])$  is a graded algebra. We assume that  $[K[X]:K]$  is finite and we show by induction that, for every key degree  $d$ ,  $G_{\nu}(K_{d-1}[X])$  is a graded algebra. If  $d = 1$ , then  $G_{\nu}(K_{d-1}[X]) = G_{\nu}(K)$  and the result is trivial. Now, assume that  $d \geq 1$ , that  $G_{\nu}(K_{d-1}[X])$  is a graded algebra and that  $d$  is not the greatest key degree. Let  $d'$  be the next key degree. If  $d$  is immediate, then, by Theorem 3.36 3), The extension  $(K_{d'-1}[X]|K_{d-1}[X], \nu)$  is immediate. Hence  $G_{\nu}(K_{d'-1}[X]) = G_{\nu}(K_{d-1}[X])$  is an algebra. We assume that  $d$  is a separate key degree, and we let  $\Phi$  be a monic polynomial of degree  $d$  such that  $\nu(\Phi)$  is the maximum of  $\nu(X^d - K_{d-1}[X])$ . We already proved above that  $G_{\nu}(K_{d'-1}[X]) = G_{\nu}(K_{d-1}[X])(in_{\nu}(\Phi))$  is a subalgebra of  $G_{\nu}(K(X))$ .

4) By Proposition 2.19, it is sufficient to show that for every integer  $n \leq d' - 1$ ,  $\nu(X^n - K_{n-1}[X])$  has a maximum. Let  $f = f_k \Phi^k + \dots + f_1 \Phi + f_0 \in X^n - K_{n-1}[X]$ , where  $f_0, f_1, \dots, f_k$  belong to  $K_{d-1}[X]$ ,  $f_k$  is monic, and  $\deg(f_k) + kd = n$ . Then  $\nu(f) = \min(\nu(f_k \Phi^k), \dots, \nu(f_1 \Phi), \nu(f_0)) \leq \nu(f_k \Phi^k)$ . Let  $j = n - dk$  be the degree of  $f_k$  and  $g \in X^j - K_{j-1}[X]$  be such that  $\nu(g)$  is the maximum of  $\nu(X^j - K_{j-1}[X])$ . Then  $\nu(f) \leq \nu(g \Phi^k)$ , which proves that  $\nu(g \Phi^k)$  is the maximum of  $\nu(X^n - K_{n-1}[X])$ .  $\square$

**Remark 3.38.** In the proof of 3) of Theorem 3.37 we showed that, for every key degree  $d$ ,  $G_{\nu}(K_{d-1}[X])$  is a graded algebra. So, Remark 3.9 2) remains true without the restriction  $d \leq [K[X]:K]/2$ . Furthermore, if  $d$  is a separate key degree, then  $d \leq [K[X]:K]/2$ . It follows that the field  $K_{\Phi}$  of Notation 3.6 is defined (where  $\Phi$  is a monic polynomial of degree  $d$  such that  $\nu(\Phi)$  is the maximum of  $\nu(X^d - K_{d-1}[X])$ ). Note that the graded algebra  $G_{\nu}(K_{d'-1}[X])$  is greater than  $G_{\nu}(K_{d-1}[X])$  if, and only if,  $d$  is a separate key degree.

We deduce a characterization of valuational key degrees by means of  $\nu K_{d-1}[X]$ .

**Proposition 3.39.** *Let  $\nu$  be a  $p$ - $m$  valuation on  $K[X]$  and  $d$  be a positive integer such that  $1 \leq d \leq [K(X):K]/2$ . Then,  $d$  is a valuational key degree if, and only if,  $\nu K_{d-1}[X]$  is a group and  $\nu K_d[X] \neq \nu K_{d-1}[X]$ . If this holds, then every monic polynomial  $\Phi$  of degree  $d$  such that  $\nu(\Phi)$  is the maximum of  $\nu(X^d - K_{d-1}[X])$  (which is equivalent to saying that  $\nu(\Phi) \notin \nu K_{d-1}[X]$ ) is a key polynomial.*

*Proof.*  $\Rightarrow$  follows from the definition and Corollary 3.8. Assume that  $\nu K_{d-1}[X]$  is a group and  $\nu K_d[X] \neq \nu K_{d-1}[X]$ . By hypothesis, there is a polynomial  $\Phi$  of degree  $d$  such that  $\nu(\Phi) \notin \nu K_{d-1}[X]$ . Since

$\nu K_{d-1}[\chi]$  is a group, by dividing  $\Phi$  by an element of  $K$  we can assume that  $\Phi$  is a monic polynomial. Let  $f \in K_{d-1}[\chi]$ . Then  $\nu(f) \neq \nu(\Phi)$ . Hence  $\nu(\Phi - f) = \min(\nu(\Phi), \nu(f)) \leq \nu(\Phi)$ . Consequently,  $\nu(\Phi)$  is the maximum of  $\nu(\chi^d - K_{d-1}[\chi])$ .

Let  $f, g$  in  $K_{d-1}[\chi]$ ,  $q = q_\Phi(f, g)$  and  $r = r_\Phi(f, g)$ . Since  $\nu(q\Phi) \notin \nu K_{d-1}[\chi]$ , we have that  $\nu(q\Phi) \neq \nu(r)$ . Hence  $\nu(fg) = \min(\nu(q\Phi), \nu(r))$ . Now, since  $\nu K_{d-1}[\chi]$  is a group,  $\nu(fg) = \nu(f) + \nu(g) \in \nu K_{d-1}[\chi]$ . Hence  $\nu(fg) \neq \nu(q\Phi)$ . So,  $\nu(q\Phi) > \nu(r)$ . This proves that  $\Phi$  is a key polynomial. Since  $\nu(\phi)$  is the maximum of  $\nu(\chi^d - K_{d-1}[\chi])$ ,  $d$  is a separate key degree.  $\square$

**Corollary 3.40.** *Let  $\nu$  be a  $p$ - $m$  valuation on  $K[\chi]$  and  $d$  be a positive integer. Assume that  $\nu K_{d-1}[\chi]$  is a group and let  $d'$  be the smallest integer such that  $\nu K_{d'}[\chi] \neq \nu K_{d-1}[\chi]$  (if any). Then  $d'$  is a valuational key degree. In particular, the smallest degree  $d$  such that  $\nu K_d[\chi] \neq \nu K$  (if any) is a valuational key degree.*

Now we turn to a particular case of immediate key degrees that we can call *dense* key degrees. It follows that if a key degree is dense, then it is the greatest key degree.

**Proposition 3.41.** *Let  $d \geq 2$ ,  $\nu' \leq \nu$  be  $p$ - $m$  valuations on  $K[\chi]$ . Assume that  $\chi^d$  is limit over  $(K_{d-1}[\chi], \nu)$ , that the restrictions of  $\nu$  and  $\nu'$  to  $K_{d-1}[\chi]$  are equal, and that there exists a monic polynomial  $\Phi$  of degree  $d$  such that  $\nu'(\Phi) < \nu(\Phi)$ . Let  $(\Phi_i)$  be a sequence of monic polynomials of degree  $d$  such that the sequence  $(\nu(\Phi_i))$  is increasing and cofinal in  $\nu K[\chi]$ , and  $\nu(\Phi) < \nu(\Phi_i)$ . Then.*

- 1)  *$d$  is an immediate key degree for  $\nu$ ,  $\Phi$  is a  $w$ -key polynomial for  $\nu$  and  $\nu'$ , and  $\nu' = \nu'_\Phi$ .*
- 2) *The  $\Phi_i$ 's are key polynomials for  $\nu$ , and for every  $f \in K[\chi]$  the sequence  $(\nu_{\Phi_i}(f))$  is eventually equal to  $\nu(f)$ .*
- 3) *The extension  $(K[\chi]|K_{d-1}[\chi], \nu)$  is dense.*
- 4) *There is no key degree greater than  $d$ .*

*Proof.* 1) By Lemma 3.31,  $\Phi$  is a  $w$ -key polynomial for  $\nu'$ ,  $\nu' = \nu'_\Phi$ , and  $\Phi$  is a key polynomial for  $\nu$ . Hence  $d$  is a key degree for  $\nu$ . Since  $\nu(\chi^d - K_{d-1}[\chi]) = \nu K[\chi]$ ,  $d$  is an immediate key degree.

2) Since  $\deg(\Phi_i - \Phi) < d$ , we have  $\nu(\Phi_i - \Phi) = \nu'(\Phi_i - \Phi)$ , and  $\nu'(\Phi_i) = \nu'_\Phi(\Phi_i) = \nu'_\Phi(\Phi + \Phi_i - \Phi) = \min(\nu'(\Phi), \nu'(\Phi_i - \Phi)) = \min(\nu'(\Phi), \nu(\Phi_i - \Phi)) = \min(\nu'(\Phi), \nu(\Phi)) = \nu'(\Phi) < \nu(\Phi) < \nu(\Phi_i)$ . By Lemma 3.31,  $\Phi_i$  is a key polynomial for  $\nu$ . We have also:  $\nu' = \nu'_\Phi$ . Let  $f \in K[\chi]$ ,  $k$  be the quotient of the euclidean division of  $\deg(f)$  by  $d$ ,  $\lambda = \nu'(f) - k\nu'(\Phi)$ , and  $i$  be such that  $\nu(\Phi_i) > \max(\nu(f) - \lambda, \nu(f))$ . Decompose  $f$  as  $f = f_{i,k}\Phi_i^k + \dots + f_{i,1}\Phi_i + f_{i,0}$ , where the  $f_{i,j}$ 's belong to  $K_{d-1}[\chi]$ . Then  $\nu'(f) = \nu'_\Phi(f) = \min(\nu'(f_{i,j}\Phi_i^j))$ . Therefore, for every  $j$  we have:  $\nu(f_{i,j}) = \nu'(f_{i,j}) \geq \nu'(f) - j\nu'(\Phi_i) \geq \nu'(f) - k\nu'(\Phi) = \nu'(f) - k\nu'(\Phi) = \lambda$ . Then for every  $j$  we have  $\nu(f_{i,j}\Phi_i^j) = \nu(f_{i,j}) + j\nu(\Phi_i) > \lambda + j \max(\nu(f) - \lambda, \nu(f))$ . If  $j \geq 1$ , then  $\nu(f_{i,j}\Phi_i^j) > \nu(f)$ . It follows that  $\nu(f) = \nu(f_{i,0}) = \nu_{\Phi_i}(f)$ . Since the sequence  $(\nu_{\Phi_i})$  is increasing, this proves that the sequence  $(\nu_{\Phi_i}(f))$  is eventually equal to  $\nu(f)$ .

3) If  $\nu(\Phi_i) > \max(\nu(f) - \lambda, \nu(f))$ , then

$$\nu(f - f_{i,0}) = \nu \left( \sum_{j=1}^k f_{i,j}\Phi_i^j \right) \geq \min_{1 \leq j \leq k} (\nu(f_{i,j}) + j\nu(\Phi_i)) \geq \lambda + \nu(\Phi_i)$$

is cofinal in  $\nu K[\chi]$ , since the sequence  $(\nu(\Phi_i))$  is. It follows that  $(K[\chi]|K_{d-1}[\chi], \nu)$  is dense.

4) Let  $f$  be a monic polynomial of degree  $d' > d$ . Then, there exists  $i$  such that  $\nu(f) = \nu_{\Phi_i}(f)$ . By Lemma 3.33 1),  $f$  is not a key polynomial for  $\nu$ . Hence  $d'$  is not a key degree.  $\square$

### 3.6. Approximations of valuations of $K(\chi)|K$ .

**Definition 3.42.** Let  $\nu$  be a  $p$ - $m$  valuation on  $K(\chi)$ ,  $1 = d_1 < d_2 < \dots < d_k < \dots$  be the sequence of key degrees of  $\nu$ . Let  $\mathcal{F}$  be a family of key polynomials which satisfies the following properties for every  $k \geq 1$ .

If  $d_k$  is a separate key degree, then  $\mathcal{F}$  contains exactly one key polynomial  $\Phi_{d_k}$  of degree  $d_k$ , and  $\nu(\Phi_{d_k}) = \max(\nu(\chi^{d_k} - K_{d_k-1}[\chi]))$ . For notational convenience, for every integer  $m$  we set  $\Phi_{d_k,m} = \Phi_{d_k}$ .

If  $d_k$  is an immediate key degree, then the key polynomials of degree  $d_k$  of  $\mathcal{F}$  form a sequence  $(\Phi_{d_k,m})$  such that the sequence  $(\nu(\Phi_{d_k,m}))$  is increasing, cofinal in  $\nu(\chi^{d_k} - K_{d_k-1}[\chi])$ .

Then we say that  $\mathcal{F}$  is a *complete family of key polynomials* for  $\nu$ .

**Remark 3.43.** For every  $p$ - $m$  valuation on  $K[\chi]$  there exists a complete family of key polynomials.

Note that in the case of transcendental extension, this has been asserted in [NS 18, Theorem 1.1].

**Theorem 3.44.** *Let  $\nu$  be a  $p$ - $m$  valuation on  $K(\chi)$ ,  $1 = d_1 < d_2 < \dots < d_k < \dots$  be the sequence of key degrees of  $\nu$  and  $\mathcal{F}$  be a complete family of key polynomials for  $\nu$ . If  $d_k$  is a separate key degree, then we let  $\Phi_{d_k}$  be the unique key polynomial of degree  $d_k$  in  $\mathcal{F}$ , and for every integer  $m$  we set  $\Phi_{d_k,m} = \Phi_{d_k}$ . If  $d_k$  is an immediate key degree, then we let  $(\Phi_{d_k,m})$  be the sequence of key polynomials of degree  $d_k$  in  $\mathcal{F}$  such that the sequence  $(\nu(\Phi_{d_k,m}))$  is increasing and cofinal in  $\nu(\chi^{d_k} - K_{d_k-1}[\chi])$ .*

For every  $k$ ,  $m_1, \dots, m_k$ , we let  $\nu_{(m_1, \dots, m_k)} = \nu_{\Phi_{d_1, m_1}, \dots, \Phi_{d_k, m_k}}$  (see Notations 3.17 and Remark 3.26). Then  $\nu$  is the supremum of the family  $(\nu_{(m_1, \dots, m_k)})$  of separate  $K$ -module valuations. For every  $f$ ,  $\nu(f)$  is the maximum of the family  $(\nu_{(m_1, \dots, m_k)}(f))$ , and there are infinitely many  $(m_1, \dots, m_k)$ 's such that  $(\nu_{(m_1, \dots, m_k)}(f)) = \nu(f)$ .

*Proof.* Note that, for every  $k$ ,  $m_1, \dots, m_k$ , we have  $\nu_{(m_1, \dots, m_k)} \leq \nu$ . Let  $f \in K[\chi]$  and  $d_k$  be the greatest key degree such that the degree of  $f$  is at least equal to  $d_k$ . By Theorems 3.37 and 3.36, there exists  $m_k$  such that  $\nu(f) = \nu_{\Phi_{d_k, m_k}}(f)$ . Furthermore, since the family  $(\nu_{\Phi_{d_k, m_k}})$  is increasing, we have  $m'_k \geq m_k \Rightarrow \nu_{\Phi_{d_k, m_k}}(f) = \nu(f)$ . Let  $f = f_j \Phi_{d_k, m_k}^j + \dots + f_1 \Phi_{d_k, m_k} + f_0$ , where  $f_0, f_1, \dots, f_j$  belong to  $K_{d_k-1}[\chi]$ . We know that there is some  $m_{k-1}$  such that  $\nu(f_0) = \nu_{\Phi_{d_{k-1}, m_{k-1}}}(f_0), \dots, \nu(f_j) = \nu_{\Phi_{d_{k-1}, m_{k-1}}}(f_j)$ . Then,  $\nu(f) = \nu_{\Phi_{d_{k-1}, m_{k-1}}, \Phi_{d_k, m_k}}(f)$ . So by induction we get a  $k$ -uple  $(m_1, \dots, m_k)$  such that  $\nu(f) = \nu_{(m_1, \dots, m_k)}(f)$ .

Now, we can do the same construction with any  $m'_k \geq m_k$ , which proves that there are infinitely many such  $k$ -uples.  $\square$

**Remark 3.45.** Let  $d$  be the degree of  $f$  and  $d_1 < \dots < d_k$  be the sequence of key degrees which are at most equal to  $d$ . If  $d_1, \dots, d_k$  are separate key degrees, then  $\nu_{(m_1, \dots, m_k)} = \nu_{(m'_1, \dots, m'_k)}$  for every  $k$ -uples  $(m_1, \dots, m_k)$  and  $(m'_1, \dots, m'_k)$ . Hence, in fact all the  $K$ -module valuations  $\nu_{(m_1, \dots, m_k)}$  are equal. Note that the restrictions of  $\nu$  and  $\nu_{(m_1, \dots, m_k)}$  to  $K_d[\chi]$  are equal. Now, if at least one of  $d_1, \dots, d_k$  is an immediate key degree, then there are indeed infinitely many distinct  $K$ -module valuations  $\nu_{(m_1, \dots, m_k)}$  such that  $\nu_{(m_1, \dots, m_k)}(f) = \nu(f)$ .

In order to get an algorithm for calculating  $\nu(f)$ , for any  $f \in K[\chi]$ , by means of the valuations  $\nu_{(m_1, \dots, m_k)}$ , we need a criterion to know whether some  $\nu_{(m_1, \dots, m_k)}(f)$  is the maximum of the family  $(\nu_{(m_1, \dots, m_k)}(f))$ . If the extension is separate, or dense, we saw in Subsection 4.1 and 4.2 that this criterion exists. Now, the dense case can be generalized, as shows the following proposition.

**Proposition 3.46.** *With the same hypothesis as in Theorem 3.44, assume that there is one immediate key degree  $d_k$ . Then, there is an algorithm for calculating  $\nu(f)$ , for any  $f \in K[\chi]$ .*

*Proof.* By Theorem 4.1 the restrictions of  $\nu$  and  $\nu_{\Phi_{d_1}, \dots, \Phi_{d_{k-1}}}$  to  $K_{d_k-1}[\chi]$  are equal. By 2) of Theorem 3.36, for every  $f \in K_{d_{k+1}-1}[\chi]$  there is an integer  $m$  such that  $\nu(f) = \nu_{\Phi_{d_k, m}}(f)$ . Now, since the sequence  $(\nu_{\Phi_{d_k, m}})$  is increasing, the family  $(\nu_{\Phi_{d_k, m}}(f))$  is eventually equal to  $\nu(f)$ . Hence  $\nu_{\Phi_{d_k, m}}(f) = \nu(f)$  if, and only if,  $\nu_{\Phi_{d_k, m}}(f) = \nu_{\Phi_{d_k, m+1}}(f)$ . Now, since the restrictions of  $\nu$  and  $\nu_{\Phi_{d_1}, \dots, \Phi_{d_{k-1}}}$  to  $K_{d_k-1}[\chi]$  are equal,  $\nu_{\Phi_{d_k, m}} = \nu_{\Phi_{d_1}, \dots, \Phi_{d_{k-1}}, \Phi_{d_k, m}}$ . For  $j > k$ , there is only one key polynomial  $\Phi_{d_j}$  of degree  $d_j$  in the family, and for every  $f$  of degree in  $\{d_j, \dots, d_{j+1} - 1\}$  we have  $\nu_{\Phi_{d_j}}(f) = \nu(f)$ . Hence we can follow in the same way as in the proof of Theorem 3.44. The algorithm is the following: we let  $d_n$  be the greatest key degree which is at most equal to the degree of  $f$ . We compute  $\nu_{\Phi_{d_1, m}, \dots, \Phi_{d_n, m}}(f)$  until  $\nu_{\Phi_{d_1, m}, \dots, \Phi_{d_n, m}}(f) = \nu_{\Phi_{d_1, m-1}, \dots, \Phi_{d_n, m-1}}(f)$  holds.  $\square$

Now, in general, a priori  $m'_1 \geq m_1, \dots, m'_k \geq m_k$  and  $\nu_{(m_1, \dots, m_k)}(f) = \nu(f)$  does not imply  $\nu_{(m'_1, \dots, m'_k)}(f) = \nu(f)$ . So we cannot get a similar criterion.

In the case of a discrete archimedean valuation (in other words,  $\nu K \simeq \mathbb{Z}$ ), every increasing sequence of  $\nu K$  is cofinal. Hence, if  $\nu(\chi^d - K_{d-1}[\chi])$  has no maximal element, then it is cofinal. Therefore,  $\chi^d$  is limit over  $K_{d-1}[\chi]$  and by Proposition 3.41  $(K[\chi]|K_{d-1}[\chi], \nu)$  is dense and  $d$  is the greatest key degree. Consequently, the first key degrees are separate and there is at most one immediate key degree. So, as we noted above, there exists an algorithm for calculating  $\nu(f)$ , for every  $f \in K[\chi]$ . This is the case studied by S. MacLane ([ML 36a] and [ML 36b]).

Now, Theorems 3.36 and 3.37 show that, given the key polynomials associated to a key degree, we can define the next key degree. So, we can construct key polynomials by induction on the degrees. We get a construction similar to the construction of M. Vaquié in [V 07]. Now, in Theorem 3.44 the definition of the key polynomials of a given key degree is independent from the key polynomials of preceding key degrees. We will not go into the details of the constructions of families of key polynomials of [HOS 07] and [V 07], because it is not the purpose of this paper and it would take up too much space.

In the case of a separate key degree  $d$ , M. Vaquié takes a monic polynomial  $\Phi$  such that  $\nu(\Phi)$  is maximal in  $\nu(\chi^d - K_{d-1}[\chi])$ . In the immediate case, he takes a sequence of monic polynomials such that the sequence of their valuations is cofinal in  $\nu(\chi^d - K_{d-1}[\chi])$ . So, our construction of the key polynomials is the same as the construction of M. Vaquié.

**3.7. Abstract key polynomials.** In this subsection  $\chi$  is assumed to be transcendental over  $K$ .

We start with some notations.

Let  $f$  be a polynomial degree of  $n$ . For  $1 \leq i \leq n$ , set  $f_{(i)} = (1/i!) f^{(i)}$ , where  $f^{(i)}$  is the  $i$ -th formal derivative of  $f$ . If the characteristic of  $K$  is  $p > 0$ , then we do not replace  $px$  by 0; the simplification holds,

if necessary, after the division by  $i!$ . For example, if  $f(X) = X^p$ , then we have  $f_{(1)}(X) = pX^{p-1} = 0$ , and  $f_{(p)}(X) = (1/p!) \cdot p! = 1$ .

**Definition 3.47.** ([DMS 18]) For any polynomial  $f$ , let  $\varepsilon_\mu(f)$  be the maximum of the set

$$\left\{ \frac{\nu(f) - \nu(f_{(i)})}{i} \mid i \in \mathbb{N}^* \right\}.$$

We say that  $\Phi$  is an abstract key polynomial (or a key polynomial) for  $\nu$  if it is monic and for every  $f \in K[X]$  such that  $\varepsilon_\nu(f) \geq \varepsilon_\nu(\Phi)$  we have  $\deg(f) \geq \deg(\Phi)$ .

**Remark 3.48.** ([NS 18, Remark 2.1]) The monic polynomials of degree 1 are abstract key polynomials.

**Proposition 3.49.** ([NS 18, Lemma 2.3 (iii)]) Let  $\Phi$  be an abstract key polynomial for  $\nu$ ,  $h_1, \dots, h_s$  be polynomials with degrees less than  $\deg(\Phi)$  and  $q$  be the quotient of the euclidean division of  $\prod_{i=1}^s h_i$  by  $\Phi$ .

$$\text{Then } \nu\left(\prod_{i=1}^s h_i - q\Phi\right) = \nu\left(\prod_{i=1}^s h_i\right) < \nu(q\Phi).$$

**Corollary 3.50.** The abstract key polynomials are key polynomials as defined in Definition 3.2.

The converse is a consequence of the following theorem. In this theorem for a key polynomial  $\Phi$  we let  $\square(\Phi) := \{f \in K[X] \mid f \text{ is monic, } \nu_\Phi(f) < \nu(f)\}$ , and  $\Xi(\Phi)$  be the subset of elements of  $\square(\Phi)$  with minimum degree.

**Theorem 3.51.** ([NS 18, Theorem 2.12]) Let  $\Phi$  be a monic polynomial. Then  $\Phi$  is an abstract key polynomial if, and only if, there exists an abstract key polynomial  $\Phi_-$  such that:

- a) either  $\Phi$  belongs to  $\Xi(\Phi_-)$ ,
- b) or
- i) the elements in  $\Xi(\Phi_-)$  have the same degree as  $\Phi_-$ ,
- ii) the set of valuations of all elements in  $\Xi(\Phi_-)$  has no maximum element,
- iii) for every  $\Phi'$  in  $\Xi(\Phi_-)$ ,  $\nu_{\Phi'}(\Phi) < \nu(\Phi)$ ,
- iv) the degree of  $\Phi$  is minimum among the degrees of polynomials satisfying iii).

**Corollary 3.52.** The key polynomials as defined in Definition 3.2 are abstract key polynomials.

*Proof.* We prove this by induction on the degrees. By Remarks 3.3 1) and 3.48 this is true for polynomials of degree 1. Let  $\Phi$  be a key polynomial of degree  $d > 1$ , and assume that every key polynomial of degree less than  $d$  is an abstract key polynomial. We denote by  $d'$  the preceding key degree. If  $d'$  is a separate key degree, then by a) of Theorem 3.51 and Theorem 3.37,  $\Phi$  is an abstract key polynomial. If  $d'$  is an immediate key degree, then by b) of Theorem 3.51 and Theorem 3.36,  $\Phi$  is an abstract key polynomial.  $\square$

**3.8. Key polynomials and graded algebras of valuations.** Here we make some remarks about the construction of families of key polynomials by lifting polynomials of the graded algebra. Let  $\Phi$  be a key polynomial, and  $d$  be its degree. We saw in Remark 3.9 (2) that  $G_\nu((K_d - 1)[X])$  is a subalgebra of  $G_\nu(K[X])$ .

For  $f := f_n\Phi^n + f_{n-1}\Phi^{n-1} + \dots + f_1\Phi + f_0$ , where  $f_0, \dots, f_{n-1}$  belong to  $K_{d-1}[X]$ , set

$$S_\Phi(f) := \{i \in \{0, \dots, n\} \mid \nu(f_i) + i\nu(\Phi) = \nu_\Phi(f) = \min(\nu(f_n) + n\nu(\Phi), \dots, \nu(f_1) + \nu(\Phi), \nu(f_0))\},$$

and  $\text{in}_{\nu, \Phi}(f)(X) = \sum_{i \in S_\Phi(f)} \text{in}_\nu(f_i)X^i$ . Then,  $\nu_\Phi(f) < \nu(f) \Leftrightarrow \text{in}_{\nu, \Phi}(f)(\text{in}_\nu(\Phi)) = 0$ . This in turn is equivalent to saying that the irreducible polynomial of  $\text{in}_\nu(\Phi)$  over  $G_\nu(K_{d-1}[X])$  divides  $\text{in}_{\nu, \Phi}(f)(X)$ .

With above notations, let  $g = \sum_{i \in S_\Phi(f)} f_i\Phi^i$ . Then,  $\nu_\Phi(g) < \nu(g)$ , and we can say that  $g$  is homogeneous

with respect to  $\Phi$ . We have  $\text{in}_{\nu, \Phi}(g)(X) = \text{in}_{\nu, \Phi}(f)(X)$ .

Assume that  $\nu(\Phi)$  is the maximum of  $\nu(\chi^d - K_{d-1}[X])$  (so  $d$  is a separate key degree). By Theorem 3.37,  $d$  divides the next key degree  $d'$ , and in the proof of this theorem we saw that  $\frac{d'}{d}$  is the degree of the irreducible polynomial of  $\text{in}_\nu(\Phi)$  over  $G_\nu(K_{d-1}[X])$ .

If we let  $\Phi'$  be a lifting of the irreducible polynomial of  $\text{in}_\nu(\Phi)$  over  $G_\nu(K_{d-1}[X])$ , then  $\Phi'$  is a key polynomial of degree  $d'$ .

**Fact 3.53.** A priori we can't assume that  $\nu(\Phi')$  is maximal, as required in Theorem 3.37.

*Proof.* Assume that  $]\nu_{\Phi_{d_k}}(g), \nu(g)[$  is nonempty and let  $\gamma \in ]\nu_{\Phi_{d_k}}(g), \nu(g)[$ ,  $h_i\Phi_{d_k}^i$  be a monomial of valuation  $\gamma$ , and  $h := g + h_i\Phi^i$ . Then,  $\text{in}_{\nu, \Phi}(h)(\text{in}_\nu(\Phi)) = \text{in}_{\nu, \Phi}(f)(\text{in}_\nu(\Phi))$ ,  $\nu_\Phi(h) = \nu(f)$ , and  $\nu(h) = \gamma$ .  $\square$

Trivially,  $\nu(\Phi)$  is not the maximum of  $\nu(\chi^d - K_{d-1}[\chi])$  if, and only if, there is  $f \in K_{d-1}[\chi]$  such that  $\nu(\Phi - f) > \nu(\Phi)$ . Now,  $\nu(\Phi - f) > \nu(\Phi)$  is equivalent to  $\nu(\Phi) = \nu(f) = \nu_\Phi(\Phi - f) < \nu(\Phi - f)$ . Therefore, the irreducible polynomial of  $\text{in}_\nu(\Phi)$  over  $G_\nu(K_{d-1}[\chi])$  has degree 1 if, and only if,  $\nu(\Phi)$  is not the maximum of  $\nu(\chi^d - K_{d-1}[\chi])$ .

Assume that  $\nu(\Phi) < \max \nu(\chi^d - K_{d-1}[\chi])$ , and let  $X - \text{in}_\nu(f)$  be its irreducible polynomial of  $\text{in}_\nu(\Phi)$  over  $G_\nu(K_{d-1}[\chi])$ , with  $f \in K_{d-1}[\chi]$ . So,  $\nu(\Phi) < \nu(\Phi - f)$ . We let  $\Phi_2 = \Phi - f$ . If  $d$  is an immediate key degree, then in this way we can construct a sequence  $(\Phi_i)$  of key polynomials such that the sequence  $(\nu(\Phi_i))$  is increasing.

**Remark 3.54.** If  $(K(\chi)|K, \nu)$  is an immediate extension, then  $G_\nu(K(\chi)) = G_\nu(K)$ . So we see that if  $d$  is an immediate key degree and  $\Phi$  is a key polynomial of degree  $d$ , then its image  $\text{in}_\nu(\Phi)$  in  $G_\nu(K(\chi))$  is already in  $G_\nu(K)$ . More generally, if  $\Phi$  is a key polynomial of degree  $d$  where  $d$  is an immediate key degree, then  $\text{in}_\nu(\Phi)$  belongs to  $G_\nu(K_{d-1}[\chi])$ .

**Fact 3.55.** Let  $d_{k_1} < \dots < d_{k_i} < \dots$  be the separate key degrees, and for every  $k_i$  let  $\Phi_{d_{k_i}}$  be a key polynomial of degree  $d_{k_i}$  such that  $\nu(\Phi_{d_{k_i}})$  is the maximum of  $\nu(\chi^{d_{k_i}} - K_{d_{k_i}-1}[\chi])$ . Then, the family of all  $\text{in}_\nu(\Phi_{d_{k_1}}^{e_1} \dots \Phi_{d_{k_i}}^{e_i})$ 's is a basis of the  $G_\nu(K)$ -module  $G_\nu(K(\chi))$ .

*Proof.* We saw in the proof of Proposition 3.37 3) that the family  $(\text{in}_\nu(\Phi_{d_{k_i}})^{e_j})$ , where  $0 \leq e_j \leq \frac{d_{k_i}+1}{d_{k_i}}$ , is a basis of the  $G_\nu(K_{d_{k_i}-1}[\chi])$ -module  $G_\nu(K_{d_{k_i}+1}-1[\chi])$ . Now, if  $d_k$  is an immediate key degree, we also noticed in the proof of Proposition 3.37 3) that  $G_\nu(K_{d_k-1}[\chi]) = G_\nu(K_{d_{k+1}-1}[\chi])$ . Assume that  $k_i + 1 \neq k_{(i+1)}$ , i.e.  $k_i + 1$  is an immediate key degree. Then, we get by induction:  $G_\nu(K_{d_{k_i+1}-1}[\chi]) = G_\nu(K_{d_{k_i+2}-1}[\chi]) = \dots = G_\nu(K_{d_{k_{i+1}}-1}[\chi])$ . Hence the family  $(\text{in}_\nu(\Phi_{d_{k_i}})^{e_j})$ , where  $0 \leq e_j \leq \frac{d_{k_i}+1}{d_{k_i}}$ , is a basis of the  $G_\nu(K_{d_{k_i}-1}[\chi])$ -module  $G_\nu(K_{d_{k_{i+1}}-1}[\chi])$ . Therefore, the family  $(\text{in}_\nu(\Phi_{d_{k_1}}^{e_1} \dots \Phi_{d_{k_i}}^{e_i}))$ , where

$$0 \leq e_1 \leq \frac{d_{k_1}+1}{d_{k_1}}, \dots, 0 \leq e_i \leq \frac{d_{k_i}+1}{d_{k_i}},$$

is a basis of the  $G_\nu(K)$ -module  $G_\nu(K_{d_{k_i}-1}(\chi))$ . It follows that the family of all  $\text{in}_\nu(\Phi_{d_{k_1}}^{e_1} \dots \Phi_{d_{k_i}}^{e_i})$ 's is a basis of the  $G_\nu(K)$ -module  $G_\nu(K(\chi))$ .  $\square$

#### 4. APPROXIMATIONS OF VALUATIONS IN THE CASE OF SEPARATE OR IMMEDIATE EXTENSIONS.

##### 4.1. Separate extensions.

**Theorem 4.1.** Let  $\nu$  be a multiplicative valuation on  $K(\chi)$ ,  $d \geq 1$  such that  $d < [K(\chi) : K]$ , and  $d_1 < \dots < d_k$  be the sequence of key degrees which are at most equal to  $d$ . Then  $(K_d[\chi]|K, \nu)$  is separate over  $K$  if, and only if,  $d_1, \dots, d_k$  are separate key degrees. Assume that this holds, and let  $\Phi_{d_1}, \dots, \Phi_{d_k}$  be key polynomials associated to the key degrees  $d_1, \dots, d_k$ , with  $\nu(\Phi_i) = \max(\nu(\chi^{d_i} - K_{d_i-1}[\chi]))$  ( $1 \leq i \leq k$ ). Then the restrictions of  $\nu$  and  $\nu_{\Phi_{d_1}, \dots, \Phi_{d_k}}$  to  $K_d[\chi]$  are equal.

*Proof.* By Proposition 2.19, if  $(K_d[\chi]|K, \nu)$  is separate, then  $d_1, \dots, d_k$  are separate key degrees. Now, assume that  $d_1, \dots, d_k$  are separate key degrees. By Theorem 3.37 2), the restrictions of  $\nu$  and  $\nu_{\Phi_1}$  to  $K_{d_2-1}[\chi]$  are equal. This is equivalent to saying that  $(1, \Phi, \dots, \Phi^{d_2-1})$  is a separate  $K$ -basis of  $K_{d_2-1}[\chi]$ . It follows by induction and by Theorem 3.37 4) that  $(K_d[\chi]|K, \nu)$  is separate. By Theorem 3.37 2), for every  $j$ ,  $1 \leq j \leq k$ , the restrictions of  $\nu$  and  $\nu_{\Phi_j}$  to  $K_{d_{j+1}-1}[\chi]$  are equal. Hence by induction  $\nu = \nu_{\Phi_1, \dots, \Phi_k}$ .  $\square$

**Remark 4.2.** Let  $(K(\chi)|K, \nu)$  be a separate algebraic extension of valued fields. Then the valuation  $\nu$  is determined by its restriction to  $K$  and by the couples  $(\Phi_1, \nu(\Phi_1)), \dots, (\Phi_k, \nu(\Phi_k))$ .

Theorem 4.1 holds for example if  $(K, \nu)$  is maximal (see Proposition 2.4). It also holds if  $\chi$  is algebraic over  $K$  and  $(K, \nu)$  is henselian with residue characteristic 0 (see Theorem 2.5).

**4.2. Immediate and dense extensions.** By definition, if  $(K(\chi)|K, \nu)$  is immediate, then  $\chi$  is pseudo-limit over  $K$ . In Paragraph 4.2.1 we show that, with some additional conditions, a limit element can generate a dense extension. Then we prove that any simple algebraic extension of a valued field is dense (Theorem 4.4). In Paragraph 4.2.2 we recall some properties of pseudo-Cauchy sequences, that we will also need in Paragraph 4.2.3. If  $\chi$  is pseudo-limit over  $(K, \nu)$ , then defining a pseudo-Cauchy sequence  $(x_i)$  with pseudo-limit  $\chi$  is equivalent to defining a sequence of key polynomials  $\Phi_i = \chi - x_i$  such that the sequence  $(\nu(\Phi_i))$  is increasing and cofinal in  $\nu(\chi - K[\chi])$ . So, the key polynomials can be seen as generalizations of the pseudo-Cauchy sequences, as noted M. Vaquié in [V 07]. In Paragraph 4.2.3 we deepen the links between these two notions. In particular, we show that if  $(K(\chi)|K, \nu)$  is a dense extension of valued fields, then 1 is the unique key degree (Theorem 4.26).



## 4.2.1. Complements to section 1.

**Proposition 4.3.** *Assume that  $\nu$  is a  $p$ - $m$  valuation on  $K(X)$ . Let  $A$  be the valuation ring of  $(K, \nu)$  and  $n \geq 2$ ,  $n < [K(X) : K]$ . If  $\nu(X^n - (K_{n-1}[X] \cap A[X])) = \nu(K_{n-1}[X] \cap A[X])$  and  $\nu K[X] = \nu K_{n-1}[X]$ , then  $(K[X]|K_{n-1}[X], \nu)$  is dense. In particular, if  $\nu(X - A) = \nu(A)$  and  $\nu K[X] = \nu K$ , then  $(K[X] : K, \nu)$  is dense.*

*Proof.* Let  $(f_i)$  be a sequence of polynomials of  $K_{n-1}[X] \cap A[X]$  such that the sequence  $(\nu(X^n - f_i(X)))$  is increasing and cofinal in  $\nu K_{n-1}[X]$ . Let  $f \in K[X]$ . By multiplying all the coefficients of  $f$  by an element of  $K$ , we can assume that  $f \in A[X]$ . If  $f \in K_{n-1}[X]$ , then  $f(X)$  is limit over  $K_{n-1}[X]$ . Assume that  $\deg(f) \geq n$ , and for every  $i$  let  $f(X) = g_i(X)(X^n - f_i(X)) + h_i(X)$  be the euclidean division. Since  $X^n - f_i(X)$  is a monic polynomial, when we do the euclidean division we see that the valuations of all the coefficients of  $g_i$  and of  $h_i$  belong to  $A$ . In particular,  $\nu(g_i(X)) \geq \min(n\nu(X), 0)$  and  $\nu(h_i(X)) \geq \min(n\nu(X), 0)$ . Now, for  $i$  large enough we have  $\nu(X^n - f_i(X)) > -\min(n\nu(X), 0) + \nu(f(X))$ , hence  $\nu(f(X)) = \nu(h_i(X)) < \nu(g_i(X)(X^n - f_i(X)))$ . It follows that  $\nu(f(X) - h_i(X)) = \nu(g_i(X)(X^n - f_i(X))) \geq \nu(X^n - f_i(X)) + \min(n\nu(X), 0)$  is cofinal in  $\nu K_{n-1}[X]$ , and that  $\nu(f(X) - K_{n-1}[X]) = \nu K[X]$ . So  $f(X)$  is limit over  $(K, \nu)$ . Consequently,  $(K[X]|K_{n-1}[X], \nu)$  is dense.  $\square$

We defined defectless extensions in Definition 2.9. It follows that a finite immediate extension  $(L|K, \nu)$  of valued fields is defectless if the restriction of  $\nu$  to  $K$  admits  $[L : K]$  distinct extensions to  $L$ .

**Theorem 4.4.** *Assume that  $\nu$  is multiplicative on  $L$  and that  $(L|K, \nu)$  is a finite algebraic immediate defectless and Galois extension of valued fields, such that  $\nu$  is an archimedean valuation (i.e.  $\nu L$  embeds in  $\mathbb{R}$ ). Then  $(L|K, \nu)$  is dense.*

To prove this theorem, we need to state more properties. Recall that if  $f$  be a polynomial degree of  $n$ , then for  $1 \leq i \leq n$ ,  $f_{(i)} := (1/i!) f^{(i)}$ , where  $f^{(i)}$  is the  $i$ -th formal derivative of  $f$ , see Definition 3.47.

**Proposition 4.5.** *Assume that  $X$  is limit over  $(K, \nu)$ , and that  $\nu$  is a  $p$ - $m$  valuation on  $K(X)$ . Then  $(K(X)|K, \nu)$  is dense.*

*Proof.* Let  $f \in K[X]$ . If  $f(X) \in K$ , then it is limit over  $(K, \nu)$ . Otherwise, for every  $x \in K$ ,  $f(X) \neq f(x)$ . Let  $n$  be the degree of  $f$ . For  $x \in K$ , we let  $f(x) := (x - X)^n f_{(n)}(X) + \dots + (x - X)f_{(1)}(X) + f(X)$  be the Taylor expansion of  $f(x)$ . Since  $f(X) \neq f(x)$ , one of the  $f_{(j)}$ ,  $1 \leq j \leq n$ , is different from 0. Then  $\nu(f(x) - f(X)) \geq \min_{1 \leq j \leq n} \nu((x - X)^j f_{(j)}(X)) = \min_{1 \leq j \leq n} j\nu(x - X) + \nu(f_{(j)}(X))$ . Since the set of  $\nu(x - X)$ 's is cofinal in  $\nu K$  and  $f_{(1)}(X), \dots, f_{(n)}(X)$  are fixed elements, this proves that the set of  $\nu(f(x) - f(X))$ 's is cofinal in  $\nu K$ . Hence  $f(X)$  is limit over  $(K, \nu)$ . Consequently  $(K[X]|K, \nu)$  is dense. Now, assume that  $X$  is transcendental over  $K$ . Let  $f, g$  in  $K[X]$  and  $x, y$  in  $K$  such that  $\nu(f(X) - f(x)) > \nu(f(X)) = \nu(f(x))$  and  $\nu(g(X) - g(y)) > \nu(g(X)) = \nu(g(y))$ . Then

$$\begin{aligned} \nu\left(\frac{f(X)}{g(X)} - \frac{f(x)}{g(y)}\right) &= \nu(f(X)g(y) - g(X)f(x) - \nu(g(X)) - \nu(g(y))) = \\ &= \nu(f(X)(g(y) - g(X)) + g(X)(f(X) - f(x)) - 2\nu(g(X))). \end{aligned}$$

Now, the sets of  $\nu(g(y) - g(X))$ 's and  $\nu(f(X) - f(x))$ 's are cofinal in  $\nu K$ . Hence the set of  $\nu\left(\frac{f(X)}{g(X)}\right)$ 's is cofinal in  $\nu K$ . It follows that  $(K(X)|K, \nu)$  is dense.  $\square$

The following proposition generalizes a result of [D 82] (p. 103) to the case when the residue field need not have characteristic 0. The proof is based on the same idea. We give it for completeness. We get a sufficient condition for being dense, by proving that some initial segment is closed under addition.

**Proposition 4.6.** (Delon) *Let  $A$  be the valuation ring of  $K$ . Assume that  $X$  is separable algebraic and pseudo-limit over  $K$ , and let  $f$  be the irreducible polynomial of  $X$  over  $K$ . Assume in addition that  $\nu$  is a  $p$ - $m$  valuation on  $K(X)$ , that  $f(X) \in A[X]$  and  $\nu(f'(X)) = 0$ . We have the following.*

- 1) *The initial segment of  $\nu K$  generated by  $\nu f(K)$  is closed under addition.*
- 2) *If  $(K, \nu)$  is archimedean, then  $(K(X)|K, \nu)$  is dense.*

*Proof.* We keep the notations of the proof of Proposition 4.5.

1) Since  $f(X)$  is a monic polynomial of  $A[X]$ , by properties of extensions of valued fields we have  $\nu(x) \geq 0$ . We show that for every  $x \in K$ , such that  $\nu(x - X) > 0$ , we have  $\nu(f'(x)) = 0$  (note that  $f' = f_{(1)}$ ). Since  $f(X) \in A[X]$ , for every  $i \geq 0$  we have  $f_{(i)}(X) \in A[X]$ . Hence  $\nu(f_{(i)}(X)) \geq 0$ . Set  $h(X) := f'(X)$ , and let  $h(x) - h(X) := (x - X)h_{(1)}(X) + \dots + (x - X)^d h_{(d)}(X)$  be the Taylor expansion. Then

$$\nu(f(x) - f(X)) \geq \min_{1 \leq i \leq d} (i\nu(x - X) + \nu(h_{(i)}(X))) \geq \nu(x - X) > 0.$$

Hence  $\nu(h(x)) = \nu(h(X)) = 0$ . Now, let  $x_0 \in A$ , such that  $\nu(X - x_0) > 0$ . First we show that  $\nu(f(x_0)) = \nu(X - x_0)$ . Indeed, we can write  $f(x_0)$  as  $f(x_0) = (x_0 - X)f_{(1)}(x_0) + (x_0 - X)^2 g(x_0, X)$ , where the coefficients

of  $g$  belong to  $A$  (since the coefficients of  $f$  belong to  $A$ ). Now,  $\nu(\chi) \geq 0$  and  $\nu(\chi - x_0) > 0$ . Hence  $\nu(x_0) \geq 0$ . Therefore,  $\nu(g(x_0, \chi)) \geq 0$ . Consequently,  $\nu((x_0 - \chi)f_{(1)}(x_0)) = \nu(x_0 - \chi) < 2\nu(x_0 - \chi) \leq \nu((x_0 - \chi)^2 g(x_0, \chi))$ . So  $\nu(f(x_0)) = \nu(x_0 - \chi)$ .

Let  $x_1 \in K$  be such that

$$\nu\left(x_1 - x_0 + \frac{f(x_0)}{f_{(1)}(x_0)}\right) > 2\nu(x_0 - \chi). \text{ Since } \nu\left(\frac{f(x_0)}{f_{(1)}(x_0)}\right) = \nu(f(x_0)) = \nu(x_0 - \chi),$$

this implies that  $\nu(x_1 - x_0) = \nu(x_0 - \chi)$ . Using Taylor expansion,  $f(x_1)$  can be written as  $f(x_1) := f(x_0) + (x_1 - x_0)f_{(1)}(x_0) + (x_1 - x_0)^2\lambda$ , with  $\nu(\lambda) \geq 0$ . Hence  $\nu(f(x_1)) \geq \min(\nu(f(x_0) + (x_1 - x_0)f_{(1)}(x_0)), \nu((x_1 - x_0)^2\lambda)) \geq 2\nu(x_0 - \chi)$ . Since the set  $\nu(f(K))$  is an initial segment of  $\nu K$ , it follows that  $\nu(f(K))$  is closed under addition.

2) Let  $x_1$  be as in 1). Since  $\nu(x_1 - x_0) = \nu(x_0 - \chi)$ , we have  $\nu(x_1 - \chi) \geq \nu(x_0 - \chi)$ . Hence  $\nu(x_1 - \chi) = \nu(f(x_1))$ . It follows that  $\nu(\chi - K)$  contains a nontrivial initial segment which is closed under addition. Assume that  $\nu K$  is archimedean. Then  $\nu(\chi - K) = \nu K$ , hence  $\chi$  is limit over  $K$ . Now, it follows from Proposition 4.5 that  $(K(\chi)|K, \nu)$  is dense.  $\square$

Now, we give sufficient conditions for hypothesis of Proposition 4.6 being satisfied.

**Proposition 4.7.** *Assume that  $\nu$  is multiplicative on  $L$  and that  $(L|K, \nu)$  is a finite immediate defectless Galois extension of valued fields. Then, there is  $\chi \in L$  with irreducible polynomial  $f$  in  $A[X]$  such that  $\nu(f'(\chi)) = 0$  and  $L = K[\chi]$  (where  $A$  is the valuation ring of  $(K, \nu)$ ).*

Before proving this proposition, we recall some definitions of [R 68] and [E 72]. Let  $L$  be a field together with valuations  $\nu_1, \dots, \nu_n$ , and for  $i \in \{1, \dots, n\}$  let  $A_i$  be the valuation rings of  $(L, \nu_i)$  and  $U_i := \nu_i^{-1}(\{0\})$  be the group of units of  $A_i$ . For  $i \neq j$  in  $\{1, \dots, n\}$ , the valuations  $\nu_i$  and  $\nu_j$  are said to be *incomparable* if nor  $A_i \subseteq A_j$  nor  $A_j \subseteq A_i$ . They are *independent* if  $A_i \cdot A_j = L$ . Note that if these valuations are archimedean, then they are independent if, and if, they are incomparable, which in turn is equivalent to:  $\nu_i \neq \nu_j$  (see [E 72, p. 82]). Now, we assume that  $\nu_1, \dots, \nu_n$  are pairwise incomparable. Then one can prove that, for  $i \neq j$  in  $\{1, \dots, n\}$ ,  $\nu_j(U_i)$  is a nontrivial convex subgroup of  $\nu_j L$ . A  $n$ -tuple  $(\gamma_1, \dots, \gamma_n) \in \nu_1 L \times \dots \times \nu_n L$  is *compatible* if there exists  $l \in L$  such that  $\nu_1(l) = \gamma_1, \dots, \nu_n(l) = \gamma_n$  (Théorème 1 p. 135 in [R 68]). One can prove that if, for every  $i \in \{1, \dots, n\}$ ,  $\gamma_i \in \bigcap_{j \neq i} \nu_i(U_j)$ , then  $(\gamma_1, \dots, \gamma_n)$  is compatible. Since  $\bigcap_{j \neq i} \nu_i(U_j)$  is an intersection of finitely many non trivial convex subgroups, it is non trivial. Now, if the  $\nu_i L$  are embedded in the same ordered group, then there exists a compatible  $n$ -tuple  $(\gamma_1, \dots, \gamma_n)$  such that  $0 < \gamma_1 < \dots < \gamma_n$ . By the approximation Theorem (Théorème 3, p. 136, in [R 68]), if  $(\gamma_1, \gamma_2, \dots, \gamma_n)$  is compatible and  $l_1, l_2, \dots, l_n$  are elements of  $L$ , such that  $\forall i, 1 \leq i \leq n, \nu_i(l_i) < \gamma_i \Rightarrow \gamma_i - \nu_i(l_i) \in \bigcap_{j \neq i} \nu_i(U_j)$ , then there exists  $l \in L$  such that

$$\forall i, 1 \leq i \leq n, \nu_i(l - l_i) = \gamma_i.$$

*Proof of Proposition 4.7.* Let  $\nu_1 := \nu, \dots, \nu_n$  be the extensions to  $L$  of the restriction of  $\nu$  to  $K$ , and  $(\gamma_1, \dots, \gamma_n) \in \nu_1 L \times \dots \times \nu_n L$  be a compatible  $n$ -tuple such that  $0 < \gamma_1 < \dots < \gamma_n$ . Let  $x_1, \dots, x_n$  in  $K$  such that  $x_i \neq 0 \Rightarrow \nu(x_i) = 0$ ,  $(x_i)_\nu \neq (x_1)_\nu$  (for  $i > 1$ ). If  $K_\nu$  is infinite, then we can assume that the  $(x_i)_\nu$ 's are pairwise distinct. Otherwise, if  $(x_i)_\nu = (x_j)_\nu$ , then we assume  $x_i = x_j$ . By the approximation Theorem, there exists  $l \in L$  such that, for every  $i \in \{1, \dots, n\}$ ,  $\nu_i(l - x_i) = \gamma_i$ . Denote by  $f$  its irreducible polynomial over  $K$ . Let  $\sigma_1, \dots, \sigma_n$  be the elements of the Galois group of  $L|K$ , and  $l_i = \sigma_i(l)$ . We know that we can assume that for  $i \in \{1, \dots, n\}$  we have:  $\nu_i = \nu_1 \circ \sigma_i$  ([R 68, p. 166]). By hypothesis,  $\nu_1(l - x_1) > 0$  and, for  $i \in \{2, \dots, n\}$ ,  $\nu_1(l_i - l) = \nu_1(l_i - x_i + x_i - l)$ . Since  $l_{\nu_1} = (x_1)_{\nu_1} \neq (x_i)_{\nu_1}$ , it follows:  $\nu_1(x_i - l) = 0$ . Now,  $\nu_1(l_i - x_i) = \nu_1(\sigma_i(l) - \sigma_i(x_i))$  (because  $x_i \in K$ , hence  $x_i = \sigma_i(x_i)$ ) and  $\nu_1(l_i - x_i) = \nu_i(l - x_i) > 0$ . Therefore  $\nu_1(l_i - l) = 0$  and  $(l_i)_{\nu_1} \neq l_{\nu_1}$ .

Since the roots of  $f$  belong to the set  $\{l_1, \dots, l_n\} \subseteq A_1$  (indeed,  $\nu_1(l_i - x_j) = 0$  and  $\nu_1(x_j) \geq 0 \Rightarrow \nu_1(l_i) \geq 0$ ), it follows that  $f \in A_1[X]$ . Consequently,  $f \in A[X]$  (because  $f \in K[X]$ ).

Let  $f(X) := X^n + a_{n-1}X^{n-1} + \dots + a_1X + a_0$  and  $f_\nu(X) := X^n + (a_{n-1})_\nu X^{n-1} + \dots + (a_1)_\nu X + (a_0)_\nu$ . The element  $l_{\nu_1}$  is a root of  $f_\nu$  and  $f'_\nu = (f_\nu)'$ . Now,  $f(X) = \prod_{i=1}^n (X - l_i)$  hence  $f'(X) = \sum_{i=0}^n \prod_{i \neq i_0} (X - l_i)$ .

$$\text{We have } f'(X) = (X - l_2)(X - l_3) \dots (X - l_n) + \sum_{i_0=2}^n \prod_{i \neq i_0} (X - l_i),$$

$$(f'_\nu)(X) = (X - (l_2)_\nu)(X - (l_3)_\nu) \dots (X - (l_n)_\nu) + \sum_{i_0=2}^n \prod_{i \neq i_0} (X - (l_i)_\nu)$$

$$(f'_\nu)((l_1)_\nu) = ((l_1)_\nu - (l_2)_\nu)((l_1)_\nu - (l_3)_\nu) \dots ((l_1)_\nu - (l_n)_\nu) + 0.$$

Hence  $(f'_\nu)((l_1)_\nu) \neq 0$  because we proved:  $\forall i, 2 \leq i \leq n, (l_i)_\nu \neq (l_1)_\nu$ . So,  $(l_1)_\nu$  is a simple root of  $f_\nu$  and  $\nu(f'(l_1)) = 0$ .

We show that  $\nu_1, \nu_2, \dots, \nu_n$  are pairwise distinct on  $K(l)$ . Let  $1 \leq i < j \leq n$ .

If  $x_i \neq x_j$ , then by hypothesis  $(x_i)_\nu \neq (x_j)_\nu$ , and  $l_{\nu_i} = (x_i)_\nu \neq (x_j)_\nu = l_{\nu_j}$ . Consequently  $\nu_i \neq \nu_j$  on  $K(l)$ .

If  $x_i = x_j$ ,  $\nu_i(l - x_i) = \gamma_i \neq \gamma_j = \nu_j(l - x_j)$ . Therefore  $\nu_i \neq \nu_j$  on  $K(l)$ .

This proves that  $\nu$  admits  $n$  distinct extensions to  $K(l)$ , hence  $[K(l) : K] \geq n$ . Since  $K(l) \subseteq L$ , it follows that  $[K(l) : K] = n = [L : K]$ . Therefore:  $K(l) = L$ . We let  $\chi = l$ .  $\square$

*Proof of Theorem 4.4.* Follows from Propositions 4.6 and 4.7.  $\square$

**Remark 4.8.** If  $L|K$  is not algebraic, then the extension  $(L|K, \nu)$  need not be dense, even if it is archimedean and separable. For example, let  $k$  be the field  $\mathbb{Q}$  or  $\mathbb{F}_p$  with  $p$  prime. Let  $K$  be the field of generalized polynomials  $k(\mathbb{Q}) := \{f = \sum_{i=1}^n x_i X^{\gamma_i} \mid n \in \mathbb{N}^*, x_1, \dots, x_n \in k, \gamma_1, \dots, \gamma_n \in \mathbb{Q}\}$ , and for  $f \in K$  let  $\nu(f)$  be the minimum of the set of  $\gamma_i$ 's such that  $x_i \neq 0$ . Denote by

$$k((\mathbb{Q})) = \left\{ \sum_{\gamma \in \Lambda} x_\gamma X^\gamma \mid \Lambda \text{ is a well-ordered subset of } \mathbb{Q}, \text{ and } \forall \gamma \in \Lambda, x_\gamma \in k \right\}$$

the field of generalized formal power series with coefficients in  $k$  and exponents in  $\mathbb{Q}$ . If  $\gamma \notin \Lambda$ , then we set  $x_i := 0$ . Recall that for  $f \in k((\mathbb{Q}))$ , the set  $\{\gamma \in \Gamma \mid x_\gamma \neq 0\}$  is called the *support* of  $f$ . It follows that the support of  $f$  is well-ordered. We let  $\nu(f)$  be the minimum of the support of  $f$  if  $f \neq 0$ , and  $\nu(0) := \infty$ . By properties of valued fields,  $k((\mathbb{Q}))$  is a valued field such that the extension  $(k((\mathbb{Q}))|K, \nu)$  is immediate. Now, let  $l := \sum_{i=1}^\infty X^{1-\frac{1}{i}} \in k((\mathbb{Q}))$ . Then for every positive integer  $n$ ,  $\nu(l - \sum_{i=1}^n X^{1-\frac{1}{i}}) = 1 - \frac{1}{n+1}$ , and this sequence is cofinal in  $\nu(l - K)$ , but not in  $\mathbb{Q}$ . Therefore,  $\nu(l - K) = \{\gamma \in \mathbb{Q} \mid \gamma < 1\}$  is bounded. So  $(K(l)|K, \nu)$  is not dense.

We saw in Proposition 4.5 that if  $\chi$  is limit over  $(K, \nu)$  and  $\nu$  is a p-m valuation on  $K(\chi)$ , then the extension  $(K(\chi)|K, \nu)$  is dense. We will show in the following subsection that if  $\chi$  is pseudo-limit on  $(K, \nu)$ , then the extension  $(K(\chi)|K, \nu)$  is not necessarily immediate. Now, with additional conditions, if  $l$  is pseudo-limit over  $(K, \nu)$ , then the extension  $(K(l)|K, \nu)$  is immediate.

**4.2.2. Pseudo-Cauchy sequences.** We assume that  $\chi$  is pseudo-limit over  $(K, \nu)$ , and we let  $(x_i)$  be a sequence of elements of  $K$  such that the sequence  $(\nu(\chi - x_i))$  is increasing and cofinal in  $\nu(\chi - K)$ . We say that  $(x_i)$  is a *pseudo-Cauchy sequence* which *pseudo-converges* to  $\chi$ , and that  $\chi$  is a *pseudo-limit* of  $(x_i)$ . Pseudo-Cauchy sequences were introduced by Kaplansky in [K 42]. The reader can find definitions and properties online in the Book of F-V Kuhlmann [FVK].

**Definitions 4.9.** A *pseudo-Cauchy sequence* of the valued field  $(K, \nu)$  is a sequence  $(x_i)$  of elements of  $K$  (where  $i$  runs over a well-ordered set) such that for every  $i < j < k$ ,  $\nu(x_i - x_j) < \nu(x_j - x_k)$ . An element  $x$  of  $K$  is a *pseudo-limit* of  $(x_i)$  if for every  $i$  we have:  $\nu(x - x_i) = \nu(x_{i+1} - x_i)$ .

**Remark 4.10.** Let  $x$  be a pseudo-limit of a pseudo-Cauchy sequence  $(x_i)$ , and  $x'$  be another element. Then  $x'$  is a pseudo-limit of  $(x_i)$  if, and only if, for every  $i$ ,  $\nu(x - x') > \nu(x - x_i)$ . Now, if the sequence  $(\nu(x - x_i))$  is cofinal in  $\nu K$ , then there is no other pseudo-limit. So, we can say that  $x$  is the limit of the pseudo-Cauchy sequence  $(x_i)$ .

**Proposition 4.11.** Let  $(L|K, \nu)$  be an extension of valued fields.

- 1) The extension  $(L|K, \nu)$  is immediate if, and only if, every element of  $L$  is pseudo-limit of a pseudo-Cauchy sequence of  $K$  which has no pseudo-limit in  $K$ .
- 2) The extension  $(L|K, \nu)$  is dense if, and only if, it is immediate and every element of  $L \setminus K$  is limit of a pseudo-Cauchy sequence of  $K$  without pseudo-limit in  $K$ .

**Proposition 4.12.** Let  $(x_i)$  be a pseudo-Cauchy sequence of a valued field  $(K, \nu)$ . For every polynomial  $f(X) \in K[X]$  (the ring of formal polynomials), the sequence  $(\nu(f(x_i)))$  is either increasing or increasing then constant. In this last case, if  $x$  is a pseudo-limit of  $(x_i)$ , then  $(\nu(f(x_i)))$  is eventually equal to  $\nu(f(x))$ . Furthermore, there is a unique monic polynomial  $f$  of minimal degree such that the sequence  $(\nu(f(x_i)))$  is not eventually constant, and  $f$  is irreducible.

**Definitions 4.13.** Let  $(x_i)$  be a pseudo-Cauchy sequence of the valued field  $(K, \nu)$ .

- 1) If for every  $f \in K[X]$  the sequence  $(\nu(f(x_i)))$  is eventually constant, then  $(x_i)$  is said to be of *transcendental type*.
- 2) Otherwise,  $(x_i)$  is said to be of *algebraic type*. The monic polynomial  $f$  of degree minimal such that the sequence  $(\nu(f(x_i)))$  is not eventually constant is called the *irreducible polynomial* of the sequence  $(x_i)$  over  $(K, \nu)$ .

**Remark 4.14.** Assume that  $\chi$  is algebraic over  $K$  and is pseudo-limit of a pseudo-Cauchy sequence of  $(K, \nu)$  without pseudo-limit in  $K$ . Then  $(x_i)$  is of algebraic type, and its irreducible polynomial has degree at most equal to the degree of the irreducible polynomial of  $\chi$ .

One can define the extension of a valuation, to an immediate extension, by means of pseudo-Cauchy sequences.

**Proposition 4.15.** *Let  $(x_i)$  be a pseudo-Cauchy sequence of a valued field  $(K, \nu)$ , without pseudo-limit in  $K$ .*

- 1) *Assume that  $(x_i)$  is of transcendental type and that  $\chi$  is transcendental over  $K$ . There is a unique extension of  $\nu$  to  $K(\chi)$  such that  $(K(\chi)|K, \nu)$  is immediate and  $\chi$  is a pseudo-limit of  $(x_i)$ . We know that for every  $f(X) \in K[X]$ , the sequence  $(\nu(f(x_i)))$  is eventually equal to some  $\gamma$ . We set  $\nu(f(\chi)) := \gamma$ .*
- 2) *Assume that  $(x_i)$  is of algebraic type and that  $\chi$  is a root of the irreducible polynomial of  $(x_i)$ . Let  $d$  be the degree of this irreducible polynomial. Then  $K(\chi) = K_{d-1}[\chi]$  and we can define  $\nu(f(\chi))$ , for every  $f(X) \in K_{d-1}[X]$ , in the same way as in 1).*

**Lemma 4.16.** *Let  $\nu$  be a  $p$ - $m$  valuation of  $K(\chi)|K$ . Assume that  $\chi$  is pseudo-limit of a pseudo-Cauchy sequence  $(x_i)$  of  $K$  (without pseudo-limit in  $K$ ). Let  $f(X) \in K[X]$ , and assume that the sequence  $(\nu(f(x_i)))$  is eventually equal to some  $\gamma$ . Then  $\gamma = \nu(f(\chi))$ , and  $f(\chi)_{\gamma, \nu} \in K_{\gamma, \nu}$ .*

*Proof.* Let  $f(X) - f(\chi) = (X - \chi)^n f_{(n)}(\chi) + \cdots + (X - \chi) f_{(1)}(\chi)$  be the Taylor expansion of  $f(X)$ , as in Proposition 4.5. For every  $x_i$  we have  $\nu(f(x_i) - f(\chi)) = \nu((x_i - \chi)^n f_{(n)}(\chi) + \cdots + (x_i - \chi) f_{(1)}(\chi)) \geq \min_{1 \leq j \leq n} \nu((x_i - \chi)^j f_{(j)}(\chi)) = \min_{1 \leq j \leq n} (j\nu(x_i - \chi) + \nu(f_{(j)}(\chi)))$ . Now, since the sequence  $(\nu(x_i - \chi))$  is increasing, the  $j\nu(x_i - \chi) + \nu(f_{(j)}(\chi))$ 's are eventually pairwise distinct. Hence the minimum is carried by only one index, say  $j_0$ . Consequently,  $\nu(f(x_i) - f(\chi)) = \nu((x_i - \chi)^n f_{(n)}(\chi) + \cdots + (x_i - \chi) f_{(1)}(\chi)) = j_0\nu(x_i - \chi) + \nu(f_{(j_0)}(\chi))$  is increasing. Since  $\nu(f(x_i)) = \gamma$  and  $\nu(f(\chi))$  are constant, it follows that  $\nu(f(\chi)) = \nu(f(x_i)) < \nu(f(\chi) - f(x_i))$ . Therefore,  $f(\chi)_{\nu(f(\chi)), \nu} = f(x_i)_{\nu(f(\chi)), \nu} \in K_{\nu(f(\chi)), \nu}$ .  $\square$

**Definitions 4.17.** A valued field  $(K, \nu)$  is said to be *maximal* if  $(K, \nu)$  admits no immediate extension. It is said to be *algebraically maximal* if  $(K, \nu)$  admits no immediate algebraic extension.

**Proposition 4.18.** *Let  $(K, \nu)$  be a valued field.*

*The field  $(K, \nu)$  is maximal if, and only if, every pseudo-Cauchy sequence of  $(K, \nu)$  has a pseudo-limit in  $(K, \nu)$ .*

*The field  $(K, \nu)$  is algebraically maximal if, and only if, every pseudo-Cauchy sequence of  $(K, \nu)$  of algebraic type has a pseudo-limit in  $(K, \nu)$ .*

**Proposition 4.19.** *Let  $\nu$  be a  $p$ - $m$  valuation of valued field on  $K(\chi)$ . Assume that  $\chi$  is pseudo-limit of a pseudo-Cauchy sequence  $(x_i)$  of  $K$  (without pseudo-limit in  $K$ ) and that  $(x_i)$  is of transcendental type over  $(K, \nu)$ . Then the extension  $(K(\chi)|K, \nu)$  is immediate.*

*Proof.* By Lemma 4.16, for every  $f(X) \in K[X]$  we have  $f(\chi) \in \nu K$  and  $f(\chi)_{\nu(f(\chi)), \nu} \in K_{\nu(f(\chi)), \nu}$ . By Remark 1.6, the extension  $(K(\chi)|K, \nu)$  is immediate.  $\square$

**Corollary 4.20.** *Let  $\nu$  be  $p$ - $m$  valuation of valued fields on  $K(\chi)$ . Assume that  $(K, \nu)$  is algebraically maximal and that  $\chi$  is pseudo-limit over  $(K, \nu)$ . Then  $(K[\chi]|K, \nu)$  is immediate.*

The following example shows that in Proposition 4.19 we cannot delete the condition  $(x_i)$  being of transcendental type.

**Example 4.21.** Assume that  $(L|K, \nu)$  is an extension of valued fields such that  $\nu L > \nu K$ , and that  $L$  contains an element  $l'$  which is pseudo-limit and algebraic over  $(K, \nu)$ . We let  $f(X)$  be its irreducible polynomial,  $d$  be the degree of  $f(X)$ ,  $(x_i)$  be a pseudo-Cauchy sequence of  $(K, \nu)$  without pseudo-limit in  $K$  and which pseudo-converges to  $l'$ . We assume that  $f$  is also the irreducible polynomial of the sequence  $(x_i)$ . For every monic polynomial  $g(X)$  of degree  $n < d$ , we know that the sequence  $(\nu(g(x_i)))$  is eventually constant. Now, let  $g(x_i) - g(l') = (x_i - l')^{n-1} g_{(1)}(l') + \cdots + (x_i - l') g_{(n)}(l')$  be its Taylor expansion. Since  $l'$  is pseudo-limit, the set of  $\nu(x_i - l')$ 's is infinite. Hence for  $\nu(x_i - l')$  large enough, the minimum of the  $\nu((x_i - l')^j g_{(j)}(l')) = j\nu(x_i - l') + \nu(g_{(j)}(l'))$  is carried by only one  $j$ , and this index is fixed, say  $j_0$ . Then,  $\nu(g(x_i) - g(l')) = j_0\nu(x_i - l') + \nu(g_{(j_0)}(l'))$  is increasing. By hypothesis, the sequence  $(\nu(g(x_i)))$  is eventually constant. Hence  $(\nu(g(x_i)))$  is eventually equal to  $\nu(g(l'))$ , and  $\nu(g(l')) < \nu(g(l') - g(x_i))$ , with  $g(x_i) \in K$ . Hence  $g(l')$  is not the maximum of  $\nu((l')^n - K_{n-1}[l'])$ . By Proposition 1.8 the extension  $(K_{d-1}[l']|K, \nu)$  is immediate. Now, let  $l''$  in  $L$  such that  $\nu(l'') > \nu K$ , and  $l := l' + l''$ . Let  $g$  be a monic polynomial of degree  $n < d$ , and  $g(l) - g(l') = (l - l')^{n-1} g_{(1)}(l') + \cdots + (l - l') g_{(n)}(l')$  be its Taylor expansion. Then,  $g(l) - g(l') = (l'')^{n-1} g_{(1)}(l') + \cdots + (l'') g_{(n)}(l')$  has valuation greater than  $\nu K$ . Consequently,  $\nu(g(l)) = \nu(g(l'))$ . This proves that the extension  $(K_{d-1}[l]|K, \nu)$  is immediate. Now,  $f(l) = f(l) - f(l') = (l'')^{d-1} f_{(1)}(l') + \cdots + (l'') f_{(d)}(l')$  is greater than  $\nu K$ . It follows that  $f(l)$  is the maximum of  $\nu(l^d - K_{d-1}[l])$ , and that the extension  $(K_d[l]|K_{d-1}[l], \nu)$  is not immediate.

**Remark 4.22.** Example 4.21 shows that in Proposition 3.41 we cannot take  $\chi^d$  pseudo-limit over  $(K_{d-1}[\chi], \nu)$  instead of  $\chi^d$  limit over  $(K_{d-1}[\chi], \nu)$ . Furthermore, in Example 4.21, 1 is an immediate key degree, and the following key degree is  $d$ , which is a separate key degree. So, the immediate key degrees are not necessarily greater than the separate ones.

### 4.2.3. Approximation of valuations.

**Theorem 4.23.** *Let  $(K(X)|K, \nu)$  be an extension of valued fields and  $(x_i)$  be a pseudo-Cauchy sequence without pseudo-limit in  $K$  and which pseudo-converges to  $\chi$ . For every  $i$  we set  $\Phi_i = \chi - x_i$ .*

1) *If  $(x_i)$  is of transcendental type, then  $(K(X)|K, \nu)$  is immediate,  $\chi$  is transcendental over  $K$  and 1 is the unique key degree. For every  $i$  and  $f \in K[X]$ , we have:  $\nu(f) = \nu_{\Phi_i}(f) \Leftrightarrow \nu_{\Phi_i}(f) = \nu_{\Phi_{i+1}}(f)$ .*

2) *Assume that  $(K(X)|K, \nu)$  is immediate, that  $(x_i)$  is of algebraic type, and let  $d$  be the degree of its irreducible polynomial.*

a) *If  $d = [K[X]:K]$ , then 1 is the unique key degree.*

b) *Otherwise,  $d$  is the second key degree.*

c) *In any case, for every  $i \geq 1$  and  $f \in K_{d-1}[X]$ , we have:  $\nu(f) = \nu_{\Phi_i}(f) \Leftrightarrow \nu_{\Phi_i}(f) = \nu_{\Phi_{i+1}}(f)$ .*

*Proof.* If  $(x_i)$  is of transcendental type, then, by Remark 4.14,  $\chi$  is transcendental, and we deduce from Lemma 4.16 that  $(K(X)|K, \nu)$  is immediate. We now assume that the extension  $(K(X)|K, \nu)$  is immediate. Let  $f(X) \in K[X]$  (the ring of formal polynomials). In the same way as in Proposition 4.6, we let  $f(X) = (X - x_i)^n f_{(n)}(x_i) + \dots + (X - x_i) f_{(1)}(x_i) + f(x_i)$  be the Taylor expansion of  $f(X)$ . Then  $f(X) = (\chi - x_i)^n f_{(n)}(x_i) + \dots + (\chi - x_i) f_{(1)}(x_i) + f(x_i)$ . Hence  $\nu_{\Phi_i}(f(X)) = \min((n-1)\nu(\chi - x_i) + \nu(f_{(n-1)}(x_i)), \dots, \nu(\chi - x_i) + \nu(f_{(1)}(x_i)), \nu(f(x_i)))$ . Since the sequences  $(\nu(f_{(j)}(x_i)))$  are increasing or eventually constant, the sequences  $(\nu((\chi - x_i)^n f_{(n)}(x_i))), \dots, (\nu((\chi - x_i) f_{(1)}(x_i)))$  are increasing. Hence, if  $(\nu(f(x_i)))$  is eventually equal to  $\nu(f(X))$ , then  $\nu_{\Phi_i}(f(X))$  is eventually equal to  $\nu(f(x_i)) = \nu(f(X))$ . Furthermore, if the minimum of  $(n-1)\nu(\chi - x_i) + \nu(f_{(n-1)}(x_i)), \dots, \nu(\chi - x_i) + \nu(f_{(1)}(x_i)), \nu(f(x_i))$  is not  $\nu(f(x_i))$ , then  $\nu_{\Phi_{i+1}}(f) > \nu_{\Phi_i}(f)$ . If the minimum is  $\nu(f(x_i))$ , then at the next step it will be  $\nu(f(x_{i+1}))$ . It follows:  $\nu_{\Phi_i}(f) = \nu(f) \Leftrightarrow \nu_{\Phi_i}(f) = \nu_{\Phi_{i+1}}(f)$ . This proves 1), 2) a) and 2) c).

2) b) In the same way as above, if  $n < d$ , then  $\nu_{\Phi_i}(f(X))$  is eventually equal to  $\nu(f(X))$ . So, the second key degree is at least equal to  $d$ . Now, let  $g(X)$  be the irreducible polynomial of the sequence  $(x_i)$ . Since the sequence  $(\nu(g(x_i)))$  is increasing, the sequence  $(\nu_{\Phi_i}(g(X)))$  is increasing. It follows that, for every  $i$ ,  $\nu_{\Phi_i}(g(X)) < \nu(g(X))$ . By Theorem 3.36 2),  $d$  is the second key degree.  $\square$

**Corollary 4.24.** *Assume that  $(K, \nu)$  is algebraically maximal and that  $\chi$  is pseudo-limit over  $(K, \nu)$ . Then 1 is the unique key degree.*

*Proof.* Let  $(x_i)$  be a pseudo-Cauchy sequence without pseudo-limit in  $K$  and which pseudo-converges to  $\chi$ . By Proposition 4.18  $(x_i)$  is of transcendental type. We conclude by Theorem 4.23 1).  $\square$

A similar result has been obtained in [NS 18, Theorem 1.2].

Now, we turn to dense extensions. Recall that by Proposition 4.5 saying that  $(K(X)|K, \nu)$  is a dense extension is equivalent to saying that  $\chi$  is limit over  $(K, \nu)$ .

**Lemma 4.25.** *Let  $(K(X)|K, \nu)$  be a dense extension of valued fields and  $(x_i)$  be a pseudo-Cauchy sequence without pseudo-limit in  $K$  and which pseudo-converges to  $\chi$ . Then  $(x_i)$  is of algebraic type over  $(K, \nu)$  if, and only if,  $\chi$  is algebraic over  $K$ . If this holds, then the irreducible polynomial of the sequence  $(x_i)$  over  $(K, \nu)$  is equal to the irreducible polynomial of  $\chi$  over  $K$ .*

*Proof.* Assume that  $\chi$  is algebraic over  $K$  and let  $f$  be its irreducible polynomial. We let  $f(x) = (x_i - \chi)^n f_{(n)}(\chi) + \dots + (x_i - \chi) f_{(1)}(\chi) + f(\chi)$  be the Taylor expansion of  $f(x_i)$ . Since  $f(X)$  is not constant, one of the  $f_{(j)}(\chi)$ ,  $1 \leq j \leq n$ , is different from 0. Then

$$\nu(f(x_i)) \geq \min_{1 \leq j \leq n} \nu((x_i - \chi)^j f_{(j)}(\chi)) = \min_{1 \leq j \leq n} j\nu(x_i - \chi) + \nu(f_{(j)}(\chi))$$

which is cofinal in  $\nu K$ . Hence  $(x_i)$  is of algebraic type over  $(K, \nu)$ .

Assume that  $(x_i)$  is of algebraic type over  $(K, \nu)$ , let  $g$  be the irreducible polynomial of the sequence  $(x_i)$  over  $(K, \nu)$  and  $g(X) = (\chi - x_i)^d g_{(d)}(x_i) + \dots + (\chi - x_i) g_{(1)}(x_i) + g(x_i)$  be the Taylor expansion of  $g(X)$ . By hypothesis, the sequences  $(g_{(d)}(x_i)), \dots, (g_{(1)}(x_i))$  are eventually constant and  $(\nu(\chi - x_i))$  is cofinal in  $\nu K$ . Hence  $(\nu(g(X) - g(x_i)))$  is cofinal in  $\nu K$ . Since  $(\nu(g(x_i)))$  is increasing, it follows that  $g(X) = 0$ . Indeed, otherwise  $(\nu(g(X) - g(x_i)))$  is eventually equal to  $\nu(g(X))$ : a contradiction. Hence,  $\chi$  is algebraic over  $K$ , and its irreducible polynomial divides  $g$ . Now,  $g$  is irreducible, hence  $g$  is the irreducible polynomial of  $\chi$ .  $\square$

**Theorem 4.26.** *Let  $(K(X)|K, \nu)$  be a dense extension of valued fields. Then 1 is the unique key degree.*

*Proof.* Let  $(x_i)$  be a pseudo-Cauchy sequence without pseudo-limit in  $K$  and which pseudo-converges to  $\chi$ . If  $\chi$  is transcendental over  $K$ , then by Lemma 4.25  $(x_i)$  is of transcendental type. Now, by Theorem 4.23 1), 1 is the unique key degree. Assume that  $\chi$  is algebraic over  $K$ . By Lemma 4.25,  $(x_i)$  is of algebraic type and the degree of the irreducible polynomial of the sequence  $(x_i)$  is  $[K(X):K]$ . Hence by Theorem 4.23 2) a), 1 is the unique key degree.  $\square$

**Proposition 4.27.** *Assume that  $\chi$  is separable algebraic over  $K$  and let  $\nu$  be an archimedean valuation on the Galois extension  $L$  generated by  $K(\chi)$  such that  $(L|K, \nu)$  is immediate and defectless. We let  $(x_i)$  be a pseudo-Cauchy sequence of  $K$ , without a pseudo-limit in  $(K, \nu)$  and with pseudo-limit  $\chi$ . For every  $i$  we set  $\Phi_i := \chi - x_i$ . Then  $\nu$  is the limit of the sequence of separate  $p$ - $m$  valuations  $\nu_{\Phi_i}$ . For  $f \in K[\chi]$ , the sequence  $\nu_{\Phi_i}(f)$  is eventually equal to  $\nu(f)$ . Furthermore,  $\nu_{\Phi_i}(f) = \nu(f)$  if, and only if,  $\nu_{\Phi_i}(f) = \nu_{\Phi_{i+1}}(f)$ .*

*Proof.* Since  $(K(\chi)|K, \nu)$  is archimedean, immediate and defectless, it is dense (see Theorem 4.4). Now the result follows from Theorems 4.26 and 4.23.  $\square$

**Theorem 4.28.** *Assume that  $\nu$  is an archimedean valuation on  $K(\chi)$  and that every algebraic extension of  $(K, \nu)$  is Galois and defectless. We assume that  $\chi$  is pseudo-limit over  $(K, \nu)$  and we let  $(x_i)$  be a pseudo-Cauchy sequence of  $(K, \nu)$  without limit in  $(K, \nu)$  and which pseudo-converges to  $\chi$ . For every  $i$  we set  $\Phi_i := \chi - x_i$ . Let  $f \in K[\chi]$ . Then the sequence  $(\nu(f(x_i)))$  is eventually equal to  $\nu(f(\chi))$ , and for every  $i$  we have:  $\nu(f) = \nu_{\Phi_i}(f) \Leftrightarrow \nu_{\Phi_i}(f) = \nu_{\Phi_{i+1}}(f)$ .*

*Proof.* If  $(x_i)$  is of transcendental type, then this follows from Theorem 4.23 1). Assume that  $(x_i)$  is of algebraic type and let  $g(X)$  be its irreducible polynomial over  $(K, \nu)$ . Let  $y$  be a root of  $g$  in any algebraic extension of  $K$ . By Proposition 4.15,  $\nu$  extends to  $K(y)$  in such a way that  $(K(y)|K, \nu)$  is immediate. By hypothesis,  $(K(y)|K, \nu)$  is Galois and defectless. Hence by Theorem 4.4  $(K(y)|K, \nu)$  is dense, so the sequence  $(\nu(y - x_i))$  is cofinal in  $\nu K$ . Now, for every  $i$  we have  $\nu(y - x_i) = \nu(x_{i+1} - x_i) = \nu(\chi - x_i)$ . It follows that  $\chi$  is limit over  $(K, \nu)$ . Now, by Proposition 4.5, the extension  $(K(\chi)|K, \nu)$  is dense. So, the result follows from Proposition 4.27. Note that by Lemma 4.25,  $\chi$  is algebraic over  $K$ , and  $g$ , is its irreducible polynomial.  $\square$

The condition that every algebraic extension of  $(K, \nu)$  is Galois and defectless holds if the residue characteristic is 0. Now, it can hold for other fields, for example the fields which are called tame. For more details see the online book [FVK].

## REFERENCES

- [AK 65] J. Ax, S. Kochen, *Diophantine problems over local fields 1 & 2*, American Journal of Mathematics 87 (1965), pp. 605-630 & 631-648.
- [B 81] W. Baur, *Die Theorie der Paare reell abgeschlossener Körper*, Logic and Algorithmic (in honour of E. Specker), Monographies de l'Enseignement Mathématique, n° 30, Université de Genève, Geneva, 1982, pp. 25-34.
- [B 82] W. Baur, *On the theory of pairs of real closed fields*, The Journal of Symbolic Logic, vol 47 (1982), pp. 669-679.
- [Bo 59] N. Bourbaki, *Algèbre*, Chapitre 5, Corps Commutatifs, Hermann, Paris, 1959.
- [DMS 18] J. Decaup, W. Mahboub, M. Spivakovski, Abstract key polynomials and comparison theorems with the key polynomials of Mac Lane-Vaquié, Illinois J. Math. 62 (2018), pp. 253-270.
- [D 82] F. Delon, *Quelques propriétés des corps valués en théorie des modèles*, thèse d'état, Paris 7, 1982.
- [D 88] F. Delon, *Extensions séparées et immédiates de corps valués*, The Journal of Symbolic Logic, vol 53 (1988), pp. 421-428.
- [D 91] F. Delon, *Indécidabilité de la théorie des paires de corps valués henséliens*, The Journal of Symbolic Logic, vol 56 (1991), pp. 1236-1242.
- [E 72] O. Endler, *Valuation Theory*, Springer, Berlin, 1972.
- [HMOS 14] F. J. Herrera Govantes, W. Mahlout, M. A. Olalla Acosta, M. Spivakovsky, *Key polynomials for simple extensions of valued fields* arXiv:1406.0657v2 [math.AG] 12 Jun 2014.
- [HOS 07] F. J. Herrera Govantes, M. A. Olalla Acosta, M. Spivakovsky, *Valuations in algebraic field extensions*, J. of Algebra, 312 (2007), pp. 1033-1074.
- [K 42] I. Kaplansky, *Maximal fields with valuations*, Duke Math Journal 9 (1942), pp. 303-321.
- [K 75] S. Kochen, *The model theory of local fields*, Logic Conference, Kiel 1974, Lecture Notes in Mathematics (499), Springer Verlag, Berlin 1975.
- [La 65] S. Lang, *Algebra*, Addison-Wesley, Reading, Massachusetts, 1965.
- [L 89] G. Leloup, *Théories complètes de paires de corps valués henséliens*, The Journal of Symbolic Logic, vol 55, n° 1, March 1990, pp. 323-339.
- [L 03] G. Leloup, *Properties of extensions of algebraically maximal fields*, Collectanea mathematica, Vol 54, n° 1, 2003, pp. 53-72.
- [L 18] G. Leloup, *Key polynomials, separate and immediate valuations, and simple extensions of valued fields*, arxiv.org/abs/1809.07092.
- [ML 36a] S. MacLane, *A construction for absolute values in polynomials rings*, Trans. Amer. Math. Soc 40 (1936), pp. 363-395.
- [ML 36b] S. MacLane, *A construction for prime ideals as absolute values of an algebraic field*, Duke Math. j. 2 (1936), pp. 492-510.
- [N 19] J. Novakoski, *Key polynomials and minimal pairs*, J. Algebra 523 (2019), pp. 1-14.
- [NS 18] J. Novakoski, M. Spivakovski, *Key polynomials and pseudo-convergent sequences*, J. Algebra 495 (2018), pp. 199-219.
- [R 68] P. Ribenboim, *Théorie des valuations*, Les Presses de l'Université de Montréal, Montréal, 1968.
- [V 07] M. Vaquié, *Extension d'une valuation*, Trans. Amer. Math. Soc. 359 (7) (2007), pp. 3439-3481.
- [FVK] <https://math.usask.ca/fvk/bookch8.pdf>

## ADDRESS

G rard LELOUP, Laboratoire Manceau de Math matiques, Le Mans Universit , avenue Olivier Messiaen,  
72085 LE MANS CEDEX, FRANCE, [gerard.leloup-At-univ-lemans.fr](mailto:gerard.leloup-At-univ-lemans.fr)