



**HAL**  
open science

## Early Paleozoic or Early-Middle Triassic collision between the South China and Indochina Blocks: The controversy resolved? Structural insights from the Kon Tum massif (Central Vietnam)

Michel Faure, van Vuong Nguyen, Luong Thi Thu Hoai, Claude Lepvrier

### ► To cite this version:

Michel Faure, van Vuong Nguyen, Luong Thi Thu Hoai, Claude Lepvrier. Early Paleozoic or Early-Middle Triassic collision between the South China and Indochina Blocks: The controversy resolved? Structural insights from the Kon Tum massif (Central Vietnam). *Journal of Asian Earth Sciences*, 2018, 166, pp.162 - 180. 10.1016/j.jseaes.2018.07.015 . hal-01878007

**HAL Id: hal-01878007**

**<https://hal.science/hal-01878007>**

Submitted on 12 Mar 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Early Paleozoic or Early-Middle Triassic collision between the South China and  
2 Indochina Blocks: the controversy resolved? Structural insights from the Kon  
3 Tum massif (Central Vietnam)

4

5

6

7 Michel Faure <sup>1\*</sup>, Van Vuong Nguyen <sup>2</sup>, Luong Thi Thu Hoai<sup>2</sup>, Claude Lepvrier <sup>3</sup>

8 1: Institut des Sciences de la Terre d'Orléans, UMR 7327 CNRS-Université d'Orléans,

9 Campus géosciences, 1A rue de la Férollerie, 45071 Orléans cedex 2, France

10 Mail: michel.faure@univ-orleans.fr Tel: + 33 (0)2 38 41 73 06

11 2: Faculty of Geology, Vietnam National University of Science, Hanoi, Vietnam

12 3: Institut des Sciences de la Terre de Paris, UMR CNRS 7193, Université Pierre-et-Marie-

13 Curie, 75252 Paris cedex 5, France

14 \* corresponding author

15

16

17 Abstract

18 In Central Vietnam, the Kontum massif is subdivided from North to South into: i)  
19 Kham Duc, ii) Ngoc Linh, iii) Kan Nack complexes. The Kham Duc complex consists of  
20 metapelite, metapsammite, paragneiss, metagabbro, amphibolite, serpentized ultramafic,  
21 and orthogneiss blocks. The southern part of this complex experienced an Early Paleozoic  
22 crustal melting event responsible for migmatites and anatectic granites. The Ngoc Linh  
23 complex, the largest one, is formed by metatexite and granitoid enclosing blocks of eclogite,  
24 high pressure (HP) granulite, orthogneiss, metagabbro, amphibolite, granodiorite, and  
25 migmatite. The Kan Nack complex is formed by migmatites with (ultra) high temperature

26 (U)HT granulite, charnockites and enderbites. The three metamorphic complexes are intruded  
27 by per-aluminous granite, and subordinate mafic plutons. In spite of local variations, the  
28 foliation of the Ngoc Linh complex defines a bulk domal structure, and the stretching  
29 lineation consistently trends NW-SE. The NW-SE lineation is also observed in the  
30 tectonically overlying the Kan Nack complex. The Ngoc Linh and Kan Nack complexes are  
31 interpreted here as parts of a metamorphic core complex (MCC), or "Ngoc Linh MCC",  
32 exhumed by a top-to-the-NW detachment fault. Along the northern margin of the Ngoc Linh  
33 MCC, upright folds with E-W striking axes, and E-W stretching lineation deform the  
34 foliation. These structures are developed in a transpressive regime coeval with the activity of  
35 plurikilometer-scale dextral strike-slip faults. Zircon and monazite U/Pb radiometric datings  
36 document a 250-240 Ma age for the crustal melting, and the top-to-the-NW ductile shearing  
37 coeval with the formation of the Ngoc Linh MCC. This extensional tectonics is followed by a  
38 dextral strike-slip faulting, at ca 240-230 Ma, before the emplacement of the two-mica  
39 granitic plutons at ca 240-224 Ma. Furthermore in the Ngoc Linh and Kan Nack complexes,  
40 zircon and monazite yield U-Pb Early Paleozoic ages. In the Kham Duc complex, a ca 460  
41 Ma MP/MT, garnet-biotite-staurolite-kyanite metamorphism, followed by migmatites at ca  
42 450 Ma, is also recognized. The 420-400 Ma age of the Dai Loc plutonic suite, and the 450-  
43 425 Ma age of the Dien Binh calc-alkaline granodiorite document an Early Paleozoic event.  
44 The Early Paleozoic mafic and ultramafic masses of the Hiep Duc complex are ophiolites  
45 included into a metasedimentary matrix. Thus, the Kham Duc complex is interpreted as a  
46 tectonic mélangé formed during a collisional orogeny that subsequently underwent crustal  
47 melting giving rise to the migmatites. A part of the Early Paleozoic rocks are enclosed as  
48 xenoliths in the Triassic migmatite of the Ngoc Linh MCC. These petrologic, structural, and  
49 chronological features allow us to propose the following geodynamic evolution for the  
50 Kontum massif. During the Early Paleozoic, a continental collision, which was

51 accommodated by a north-directed subduction of a southern block (S. Vietnam, or Viet-  
52 Cambodia block) below a northern block (N. Vietnam or Viet-Lao block), occurred along the  
53 Tam Ky-Phuoc Son suture. During the Early Triassic, at ca 250-245 Ma, the S. Vietnam block  
54 experienced a crustal melting giving rise to the Ngoc Linh MCC that reworked the Early  
55 Paleozoic orogen. Lastly, during the Middle Triassic, the dextral faulting superimposed upon  
56 the Ngoc Linh migmatite. The geodynamic significance of the Kontum massif is discussed in  
57 the general framework of Indochina- South China relationships. The Early Paleozoic collision  
58 in Vietnam is viewed as the driving force responsible for the development of the  
59 intracontinental orogeny of S. China. In order to account for the important heat flow required  
60 for the formation of UHT metamorphic rocks, a possible interaction between the SCB-  
61 Indochina collision, and the Emeishan mantle plume is proposed.

62

63

64 Keywords: Early Paleozoic orogeny, Triassic tectonics, South China-Indochina connection,  
65 Kon Tum massif, Metamorphic Core Complex, Plume-modified orogeny

66

67

## 68 1. Introduction

69 Continent or microcontinent collisions that follow the closure of oceanic basins played  
70 a major role in the formation of Asia since the Neoproterozoic. In this general frame, the  
71 collision between the Indochina block and the South China block (SCB) is an important  
72 feature that accounts for the Asian continental assembly. In North Vietnam, the Late Triassic  
73 (Norian) unconformity documents an Early Triassic tectonic event, called "Indosinian  
74 orogeny" (Fromaget, 1927, 1941). Nevertheless, the location of the ophiolitic suture, the age  
75 of the collision, and the subduction polarity have been debated for a long time (for a recent

76 discussion of these issue see Faure et al., 2014, and enclosed references). Concerning the age  
77 of the Indochina block-SCB collision, two main opinions are currently proposed. Some  
78 authors (e.g. Lepvrier et al., 1997, 2004, 2008, 2011; Nakano et al., 2010; Liu et al., 2012;  
79 Faure et al., 2014, 2016a) argue that the collision took place in Early to Middle Triassic (ca  
80 250-230 Ma). The Ailaoshan-Song Ma-Song Chay-Hainan ophiolitic suture of SW China and  
81 N. Vietnam represents the plate boundary along which the Paleotethys ocean was subducted  
82 (Fig. 1). The collision was preceded by a South (or SW in the present geographic coordinate  
83 frame) directed subduction giving rise to a magmatic arc on the Indochina block side, and  
84 followed by a North (or NE) propagation of the folds and low-angle thrusts. In the Truong  
85 Son belt of N. Vietnam (Fig. 1), the late collisional Triassic deformation was accommodated  
86 by NW-SE striking dextral strike-slip faults (Lepvrier et al., 1997, 2004). Another group of  
87 authors proposes the view that the collision occurred in Early Paleozoic on the basis of ca 400  
88 Ma plutons, Devonian unconformity, and biostratigraphic correlations between Devonian  
89 shallow water terrigenous deposits and similar fossil assemblages in NE Vietnam and South  
90 China (e.g. Janvier et al., 1996, 1997; Thanh et al., 1996; Metcalfe, 1996, 2013; Findlay,  
91 1997; Findlay and Pham, 1997; Carter et al., 2001; Carter and Clift, 2008; Roger et al., 2008,  
92 2012; Tran and Vu, 2011). For these authors, the pre-Late Triassic tectonic-metamorphic and  
93 plutonic events were interpreted as an intracontinental tectonics due to a far-field effect of the  
94 collision between the Indochina and Sibumasu blocks.

95         Indeed, most of the controversial points arose from a lack of detail knowledge, or from  
96 over-interpretations of a local geological record extrapolated to remote areas. As shown  
97 below, the divergent points can be easily reconciled if a polyorogenic evolution is considered.  
98 This paper presents new structural data from the Kon Tum massif of Central Vietnam that  
99 was first considered as the Precambrian core of Indochina (e.g. Hoffet, 1933; Saurin, 1935,  
100 Hutchison, 1989; Lan et al., 2003; Tran and Vu, 2011) but more recently, Early Paleozoic and

101 Early to Middle Triassic ages have been documented there (Nam et al., 2001; Osanai et al.,  
102 2004; Maluski et al., 2005; Roger et al., 2007; Usuki et al., 2009; Nakano et al., 2007, 2009,  
103 2013). Our new field and laboratory analyses allow us to propose a crustal scale view of the  
104 architecture of the Kon Tum massif. The place of the Kon Tum massif in the Paleozoic to  
105 Triassic geodynamic evolution of the Indochina-South China collisional framework is also  
106 discussed.

107

## 108 2. Regional setting

109 Below the Late Triassic and younger formations, the study area (Fig. 2) can be divided  
110 into two domains, namely: i) a northern one, between Da Nang and Hue, forms the south  
111 Truong Son area (Lepvrier et al., 1997), or the Binh Tri Thien zone (Thanh et al., 2013),  
112 which is occupied by weakly or unmetamorphosed formations, and ii) the Kon Tum massif  
113 composed of metamorphic and plutonic rocks. Both domains also expose several generations  
114 of granitic plutons. The post-tectonic formations include, from the younger to the older,  
115 Quaternary to Cenozoic basaltic flows, and Cretaceous to Late Triassic terrigenous  
116 formations. SW of Da Nang, the Late Triassic to Jurassic sandstone, conglomerate, and  
117 siltstone deposits (Nong Son Formation) that unconformably overlie the folded Paleozoic  
118 formations in the northern domain provide an upper time limit for the tectono-metamorphic  
119 events of the study area. In the Kon Tum massif, Jurassic and Late Cretaceous red sandstones  
120 and conglomerates were deposited in N-S striking half-grabens that overlie the metamorphic  
121 rocks. The detailed structure of these formations that post-date the main tectonic events is not  
122 documented here.

123

### 124 2.1. Lithotectonic units of the Kon Tum massif

125           The granitic and metamorphic Kon Tum massif, formerly considered as the  
126 Precambrian nucleus of the Indochina block (Lan et al., 2003; Tran and Vu, 2011), is  
127 subdivided into three litho-tectonic units, namely from North to South, the Kham Duc, Ngoc  
128 Linh, and Kan Nack complexes (Tran and Vu, 2011; Fig. 2). The radiometric ages that are  
129 crucial to discuss the tectonic and geodynamic significance of the syn- to post-metamorphic  
130 events will be presented below in section 4.

### 131 *2.1.1. The Kham Duc complex.*

132           This unit is bounded to the North by a granitic intrusion belonging to the Ben Giang-  
133 Que Son plutonic suite, and to the South by the E-W striking dextral Tra Bong strike-slip fault  
134 (Lepvrier et al., 1997). The Kham Duc complex can be subdivided in two parts. The northern  
135 one consists of metapelite, metagraywacke, quartz schist, and chlorite actinolite volcanic-  
136 sedimentary rocks. Amphibolite, locally some of them preserving a gabbroic texture, and  
137 serpentinized ultramafics are exposed as scattered blocks in the metasedimentary suite (Fig.  
138 3A, B). The mafic-ultramafic rocks, called the Hiep Duc complex, are interpreted as a  
139 dismembered ophiolitic series (Izokh et al. 2006; Pham et al., 2006; Tran and Vu, 2011). To  
140 the South, the second part of the Kham Duc complex is characterized by the occurrence of  
141 migmatites and anatectic granites, called the Chu Lai migmatitic complex. In these rocks, the  
142 leucosomes are well foliated and folded (Fig 3C,D), and the amphibolite, and metagabbro  
143 bodies represent xenoliths. Due to poor outcrop quality, and progressive transition from non-  
144 melted to completely melted rocks, it is impossible to clearly delineate the boundary between  
145 the two parts (Fig. 2). In the following, they will be collectively referred to as the Kham Duc  
146 complex. The age of the protoliths is unknown, however the metapelite yields detrital zircon  
147 grains dated at  $558\pm 7$  Ma (Usuki et al., 2009) showing that at least a part of the Kham Duc  
148 complex is younger than Late Neoproterozoic. The Tra Bong fault system separates two  
149 different migmatitic complexes, the Chu Lai and Ngoc Linh to the North and South,

150 respectively. As they partly experienced the same deformation events, and in absence of  
151 precise dating, their distinction remains difficult.

152

### 153 2. 1. 2. *The Ngoc Linh complex.*

154 This rhombic shaped complex is limited to the North and South by the Tra Bong and  
155 Ba To faults, respectively. Westward, it probably extends into Laos (Fig. 2). The Ngoc Linh  
156 complex consists of metatexites and biotite-garnet-sillimanite gneiss (Fig. 4A, B) with  
157 foliated quartz-feldspar layers corresponding to leucosome. Mafic rocks, such as garnet  
158 amphibolite, eclogites, HP granulite, and biotite-hornblende orthogneiss, are widespread in  
159 the complex (Fig. 4C; Nakano et al., 2007; Osanai et al., 2008; Yonemura et al., 2013). These  
160 rocks are interpreted as refractory restites enclosed in the migmatite. The biotite-hornblende  
161 orthogneiss represent remnants of the pre-Ngoc Linh migmatite plutons, such as the Dai Loc  
162 or Ben Giang-Que Son bodies. Furthermore, South of Di Lang, along the Song Re, the Ngoc  
163 Linh migmatite encloses blocks of metatexite with a folded foliation unrelated to the  
164 deformation of the host rock migmatite (Fig. 4D). This feature argues for two generations of  
165 migmatites.

166

### 167 2. 1. 3. *The Kan Nack complex.*

168 This complex that crops out in the southeastern part of the Kon Tum massif, is  
169 composed of metagraywacke, metapelite, quartzite, marble, and mafic rocks metamorphosed  
170 under granulite facies conditions (Roger et al., 2007; Nakano et al., 2007, 2013). These HT  
171 metamorphic rocks, belonging to the khondalite-kinzigite series, are associated with garnet-  
172 orthopyroxene granite, or charnockites, and gabbro-tonalites, or enderbites, attesting to the  
173 melting of the lower continental crust under HT conditions (Roger et al., 2007). Zircon,  
174 monazite U-Pb, and biotite, muscovite radiometric datings document two metamorphic and



175 magmatic events at ca 465-422 Ma, and 255-240 Ma (Carter et al., 2001; Nagy et al., 2001;  
176 Maluski et al., 2005; Roger et al., 2007; Nakano et al., 2007, 2013).

177

## 178 2.2. The Northern Domain (South Truong Son area)

179 This area between Hue and Da Nang consists of Early Paleozoic to Triassic rocks. The  
180 Early Paleozoic (Cambrian-Ordovician) A Vuong Formation is composed of weakly  
181 metamorphosed quartz-sericite schist, sandstone and quartzite. It is unconformably covered  
182 by conglomerate, sandstone, and siltstone of Devonian age. Although globally non-  
183 metamorphic, these terrigenous formations are deformed by upright folds with the local  
184 development of an axial planar slaty cleavage. This event is related to the dextral strike-slip  
185 faulting widespread between Da Nang and Hue (Lepvrier et al., 1997). Permian and Triassic  
186 rocks crop out in the western part of the Northern Domain (Fig. 2). Similar rocks are also  
187 exposed in the southwestern part of the Kon Tum massif. These formations consist of  
188 unmetamorphosed terrigenous deposits and felsic and mafic volcanites. The basaltic rocks  
189 sometimes exhibit a pillowed structure. To the SW of the Kon Tum massif, near Sa Thay,  
190 Permo-Triassic felsic lava flows are also exposed.

191

## 192 2.3. The plutonic rocks

193 In the study area, the widely exposed plutonic rocks are gathered into two groups,  
194 namely the post-Ngoc Linh and the pre-Ngoc Linh suites (Fig. 2).

### 195 2.3.1. *The post Ngoc Linh plutons.*

196 These rocks are also collectively called the Hai Van complex. The Ba Na and Van  
197 Canh suites are included in this group (Tran and Vu, 2011). The Hai Van, Ba Na, and Van  
198 Canh plutons consist of biotite granite and two-mica cordierite  $\pm$  sillimanite  $\pm$  garnet  
199 granitoids with locally a migmatitic appearance and numerous aplitic dykes well exposed

200 North of Da Nang. Furthermore, the Hai Van granite intrudes migmatite in the central part of  
201 the Ngoc Linh Complex. Zircon yields LA-ICP-MS ages ranging between 242 Ma and 224  
202 Ma (Hieu et al., 2015). Monazite U-Pb chemical ages distribute between 240 Ma and 220 Ma  
203 (Nakano et al., 2013). This range of ages is globally older than the Late Triassic sandstone, at  
204 ca 220-200 Ma that unconformably overlie the plutonic rocks. The peraluminous geochemical  
205 signature of these Middle to Late Triassic plutonism document an important crustal melting  
206 event. The presence of Early Paleozoic inherited zircon grains, dated at  $438\pm 4$  Ma, and  $380\pm 4$   
207 Ma, also indicates the crustal contribution in the magma genesis (Shi et al., 2015). Small  
208 bodies of gabbro and diorite (Cha Val and Phu Loc magmatic suites) associated with the Hai  
209 Van plutons suggest that a mantle magmatic input might have triggered the crustal melting  
210 (Tran and Vu, 2011).

211 From a structural point of view, the Hai Van, Ba Na, and Van Canh plutons sometimes  
212 exhibit a planar mineral preferred orientation. Although a magmatic mineral fabric may be  
213 locally observed, a post-solidus one is lacking. In agreement with the radiometric ages, these  
214 plutons were emplaced after the development of the main foliation recorded in the Ngoc Linh,  
215 Kan Nack and Kham Duc metamorphic complexes.

216

### 217 2.3.2. *The pre-Ngoc Linh plutons.*

218 On the basis of their age and geochemistry, three earlier magmatic suites are  
219 distinguished. From the younger to the older they are the Ben Giang-Que Son, Dai Loc, Dien  
220 Binh, and Tra Bong suites (Tran and Vu, 2011):

221 The Ben Giang-Que Son granites are well exposed both in the Northern Domain, NW  
222 of Tam Ky, and in the Kon Tum metamorphic domain, where they are exposed as an  
223 unmelted blocks within the migmatite. Several petrographic facies including diorite,  
224 quartzdiorite, gabbro, porphyritic granodiorite, and syenite are recognized (Tran and Vu,

225 2011). Zircon U-Pb (SHRIMP and LA ICP-MS) dating from granodiorite yields ages ranging  
226 between 295 Ma and 261 Ma, with a peak at 294 Ma (Dinh, 2011; Nguyen Trong Giang  
227 unpublished in Tran and Vu, 2011). Furthermore, zircon separated from a quartz diorite rock  
228 in the large pluton that separated the southern part of the Truong Son belt) and the Kham Duc  
229 complex, yields LA-ICP-MS age at  $479\pm 3$  Ma (Hieu and Trung, 2016a). On the basis of its  
230 calc-alkaline geochemical features, the Ben Giang-Que Son plutonic suite is interpreted as  
231 mantle derived magmas contaminated by crustal materials. These rocks are considered as  
232 emplaced in a magmatic arc formed in a continental active margin (Tran et al., 2008; Hieu  
233 and Trung, 2016a).

234 The Dai Loc magmatic suite is well developed SW of Da Nang in the northern  
235 Domain, but it may also crop out as xenolith in the Ngoc Linh and Kan Nack complexes (Fig.  
236 2). The dominant rocks are porphyritic biotite granite with subordinate two-mica granite,  
237 syenogranite, and monzogranite. Most of the plutons of the northern domain exhibit a well  
238 developed planar and linear fabric related to the dextral strike-slip fault system widespread  
239 there (Lepvrier et al., 1997; 2008). The per-aluminous and subalkaline geochemical features  
240 of these plutons indicate that they result of the partial melting of continental crust (Hieu et al.,  
241 2016b). Zircon SHRIMP dating gives  $418\pm 18$  Ma and  $407\pm 11$  Ma ages (Carter et al., 2001),  
242 whereas zircon LA-ICP-MS analyzes indicate  $427\pm 10$  Ma and  $423\pm 2$  Ma ages (Hieu et al.,  
243 2016b). These late Silurian ages argue for an important Early Paleozoic magmatic event.

244 The Dien Binh and Tra Bong magmatic suites are scattered in the Western and in the  
245 Northern parts of the Ngoc Linh metamorphic complex. They mainly crop out close to the Po  
246 Ko and Tra Bong faults, respectively (Fig. 2). These rocks are dominantly diorite, and  
247 porphyritic granodiorite with limited gabbro. The Tra Bong pluton experienced a pervasive  
248 post-solidus ductile deformation associated with the dextral strike-slip shearing (Lepvrier et  
249 al., 1997, 2004, 2008). Zircon fractions from a Dien Binh granodiorite yield a U-Pb age at

250 415±3 Ma (Nagy et al., 2001), and SHRIMP dating of a gneissic diorite gives an Early  
251 Paleozoic age at 444±17 Ma (Carter et al., 200). The calc-alkaline geochemical character of  
252 the Dien Binh plutons suggests that they formed in a magmatic arc setting (Nagy et al., 2001).

253

### 254 3. Structural analysis of the polyphase deformation

255 In contrast to the numerous petrological and geochronological studies, the structural  
256 investigations of the Kon Tum massif remain rare (Lepvrier et al., 1997, 2004, 2008; Tran et  
257 al., 2014). Our study allows us to recognize three ductile and synmetamorphic events called in  
258 the following D<sub>1</sub>, D<sub>2</sub>, and D<sub>3</sub> in a younging relative chronological order. Furthermore, some  
259 rare lines of evidence argue for an early "pre-D<sub>1</sub>" event, but the regional foliation, ascribed to  
260 the D<sub>1</sub> event, represents the main deformation observed in all metamorphic complexes.

261

#### 262 3.1. The syn-migmatitic D<sub>1</sub> main structural event.

##### 263 3.1.1. *Fabric elements*

264 The three metamorphic complexes exhibit a pervasive foliation (S<sub>1</sub>) defined by the  
265 preferred orientation of platy minerals, such as biotite and muscovite, and also by the  
266 alternation of quartz and feldspar layers. This planar fabric is well developed in the Ngoc  
267 Linh complex where foliated migmatites exhibit a mylonitic foliation. S<sub>1</sub> often contains a  
268 mineral and stretching lineation, L<sub>1</sub>, represented by elongated biotite streaks, and feldspar  
269 porphyroclasts. As observed by planar joints and triple junctions at ca 120° of the quartz and  
270 feldspar grains, a thermal annealing induced a static recrystallization that often erased L<sub>1</sub>. In  
271 the eastern part of the Ngoc Linh complex, between Quang Ngai and Ba To, the foliation dips  
272 to the East or SE (Figs. 2, 5 stereograms G, I). In the southwestern part of the Ngoc Linh  
273 complex, from Kon Tum to Dak To, S<sub>1</sub> dips to the southwest, in its western part, North of  
274 Dak To, along the Po Ko fault, and up to the Laos border, S<sub>1</sub> dips westward (Fig. 2, 5

275 stereograms D, E). Therefore, as already recognized by Lepvrier et al., (2004, 2008), the  
276 foliation of the Ngoc Linh complex defines a domal structure. The antiformal geometry of  $S_1$   
277 clearly appears along the Song Re, between Di Lang and Ba To (Fig. 2, 5). Along the  
278 northern boundary of the Ngoc Linh dome,  $S_1$  dips alternatively northward or southward due  
279 to the post-folial folding ascribed to the  $D_2$  event. In several places,  $S_1$  is folded by NE-  
280 trending, and NW-verging folds. Although obviously post-dating the  $S_1$  formation, these  
281 structures are distinct from the upright folds ascribed below to the  $D_2$  event. They are  
282 interpreted here as a late increment of the  $D_1$  top-to-the NW sense of shear, and shown as  $F_{1-2}$   
283 folds in Fig. 5 stereograms D and J. Furthermore, the variable dip and strike attitude of  $S_1$   
284 suggests that the Ngoc Linh dome may be composed by several sub-domes, however the  
285 present state of knowledge does not allow us to precisely depict the geometry.

286 In contrast to the domal architecture of the Ngoc Linh complex defined by the  
287 variously oriented  $S_1$ , the  $L_1$  stretching and mineral lineation presents a well organized  
288 pattern, since whatever the attitude of  $S_1$ ,  $L_1$  trends NW-SE (Figs 2, 5). However, SW of Tra  
289 My, and in the NE corner of the dome, the stretching lineation trends NE-SW (Figs. 2, 5  
290 stereogram K). This local feature is interpreted as the result of recumbent folding, in which  
291 the dominantly NW-SE trending lineation trends locally NE-SW in the inverted limb (Fig. 6).  
292 Indeed, in the NE part of the Ngoc Linh dome, small-scale folds, overturned to the NW, and  
293 associated to a low angle axial plane, are recognized (Fig. 7C). They correspond to NW-  
294 verging collapse folds indicating a subvertical shortening. These folds deform  $S_1$ , but as their  
295 NW-ward vergence is consistent with the syn- $S_1$  ductile shearing, they are ascribed here to a  
296 late increment of the  $D_1$  event. As a matter of fact, when  $S_1$  is overturned,  $L_1$  strikes NE-SW  
297 instead of NW-SE (Fig. 6).

298 The same  $S_1$  foliation is observed in the Kan Nack complex with variable attitude (Fig  
299 5 stereogram H). Globally,  $S_1$  is folded into a NW-SE trending structure with a general

300 southward or southeastward dip, well observed in the SE part of the Kan Nack complex (Fig.  
301 2). Geometrically, in spite of undulation of  $S_1$ , the Kan Nack complex appears as a multi-km  
302 scale migmatitic and granulitic gneiss slice dipping to the SE. Most of the present boundaries  
303 between the Kan Nack and Ngoc Linh complexes are late high-angle faults (Fig. 2), and thus  
304 the syn-folial tectonic contact between the two units is not observed. From the field  
305 observation of the SE-ward dip of the  $S_1$  foliation in both complexes, we infer that the Kan  
306 Nack complex overlies the Ngoc Linh complex, in agreement with previous works (Nagy et  
307 al., 2001; Cau et al., 2004; Lepvrier et al., 2004; Tran et al. 2006). Due to the jump in the  
308 metamorphic grade between the two complexes, a flat-lying tectonic contact should separate  
309 them, however, this boundary is not precisely recognized in the field since the two units are  
310 presently separated by NE-SW striking high angle faults. Alike for the Ngoc Linh complex,  
311 the Kan Nack complex contains a E-W to NW-SE striking mineral and stretching lineation,  
312 also ascribed to  $L_1$  (Figs. 2, 5 stereogram H). Due to the intense recrystallization,  $L_1$  is less  
313 developed than in the Ngoc Linh complex.

314 The Kham Duc complex exhibits also a pervasive foliation  $S_1$  globally striking E-W or  
315 NE-SW but with a variable dip due to the late  $D_2$  folding (Fig. 5 stereograms B, and L). Near  
316 Phuoc Son,  $L_1$  is well developed, however, eastwards, the  $D_1$  event is no more observed and  
317  $D_2$  predominates (Fig. 5 stereogram L).

### 318 *3.1.2. Kinematic analysis*

319 In spite of the static recrystallization that often erased the kinematic indicators, in  
320 section perpendicular to  $S_1$  and parallel to  $L_1$ , (i.e. in the XZ plane of the finite strain  
321 ellipsoid), ductile shear criteria, coeval with the development of the SL fabric, can be  
322 recognized both in the field and in thin sections. In the field, the most common criteria are  
323 sigmoidal leucosomes, asymmetric boudins, and drag folds (Figs 7A,B, C, E, F, 8A, B). In  
324 thin section, cut in the XZ plane of the finite strain ellipsoid, sigmoidal biotite and muscovite,

325 sigma-type porphyroclast systems, oblique quartz shape fabric, and pressure shadows are the  
326 commonest kinematic indicators (Fig. 9). From these observations, a top-to-the-NW shearing  
327 is widely distributed in the migmatites, gneiss, amphibolites, micaschists and quartzites of the  
328 Ngoc Linh, Kan Nack, and Kam Duc complexes. The D<sub>1</sub> ductile shearing is coeval with the  
329 HT/MP metamorphism and migmatization recorded in the Ngoc Linh and Kan Nack  
330 complexes. It is worth to note that when L<sub>1</sub> strikes NE-SW, due to the recumbent folding, as  
331 observed at the cm- and m-scale (Fig. 7C), and inferred for larger scale, the same kinematic  
332 indicators show a top-to-the-NE sense of shear can be observed in the inverted limb (Fig. 6).  
333 3.2. The post-migmatitic D<sub>2</sub> event.

334 As mentioned above, the domal geometry of the Ngoc Linh migmatite is modified  
335 along its northern boundary by a set of ductile dextral strike-slip faults, collectively called the  
336 Tra Bong fault system (Fig. 2, Lepvrier et al., 1997, 2004). In this area, the S<sub>1</sub> foliation is  
337 deformed by E-W striking upright folds (F<sub>2</sub>, Fig. 5 stereogram L, 7D). Locally an axial planar  
338 foliation may develop. These F<sub>2</sub> folds formed during a transpressional component of the  
339 strike-slip faulting. It is worth to note that the Tra Bong fault system belongs to an important  
340 network of dextral strike-slip faults well developed in the entire Truong Son belt (Lepvrier et  
341 al., 1997, 2004). In the study area, the Dai Loc orthogneiss, and their country rocks, west of  
342 Da Nang, are intensely deformed by the dextral shearing (Fig. 5 stereograms A and M, see  
343 Lepvrier et al., 1997, 2004 for details). From a structural point of view, the D<sub>1</sub> features (S<sub>1</sub>,  
344 L<sub>1</sub>, and the top-to-the-NW shearing) are clearly deformed by the dextral shearing D<sub>2</sub> event. As  
345 indicated by the <sup>40</sup>Ar/<sup>30</sup>Ar radiometric ages at ca 245 Ma, and 230 Ma for D<sub>1</sub> and D<sub>2</sub>, the two  
346 events developed successively but in a short time span (Lepvrier et al., 2007, 2004).  
347 Furthermore, north of the Tra Bong fault, the F<sub>2</sub> folds involve the Chu Lai migmatite, the  
348 foliation of which is deformed by isoclinal folds coeval with melting (Fig. 8C, D) that must be  
349 related to an older event.

350

351 3.3. The late D<sub>3</sub> event.

352           The S<sub>1</sub> foliation of the Ngoc Linh dome is also locally cut by high angle normal faults.  
353 These structures are well documented to the NE of Kon Tum and in the southeastern part of  
354 the dome (Lepvrier et al., 2004, 2008). Since the Hai Van two mica granites are involved in  
355 the faulting, the deformation should have occurred in the Late Triassic or after. Furthermore,  
356 the eastern side of the N-S to NNE-SSW striking normal fault that reworks the Ngoc Linh and  
357 Kan Nack complexes is moving downward. The hangingwall of the fault is a half-graben  
358 filled by Jurassic continental deposits (Fig. 2). Thus, a Jurassic age is tentatively assumed for  
359 these brittle-ductile normal faults. In the following, this post-migmatitic tectonic event will  
360 not be considered.

361

362 3.4. The pre-D<sub>1</sub> inherited structures in the Kham Duc complex.

363           E-W striking upright folding, attributed to the D<sub>2</sub> event, is well developed in the Kham  
364 Duc complex (Fig. 8B,C). In the southern part of the complex, upright or NW-verging folds  
365 deform the isoclinally folded leucosomes parallel to the migmatitic foliation of the Chu Lai  
366 migmatite (Figs. 3D, 8D). Near Tra My, the axes of the isoclinal folds trends N-S with  
367 variable dip, depending on their location within F<sub>2</sub> (Fig. 10). Since these intrafolial folds are  
368 coeval with the crustal melting, the formation time of this isoclinal folding is that of the Chu  
369 Lai migmatite. Several lines of evidence argue for an Early Paleozoic age (ca 450 Ma) for the  
370 Chu Lai migmatite. Therefore, these structures have to be ascribed to a pre-D<sub>1</sub> tectonic-  
371 metamorphic event. This interpretation is supported by petrological and geochronological  
372 constraints presented in the next section.

373           Another line of evidence for a pre-D<sub>1</sub> event is provided by the Song Re section, south  
374 of Di Lang (Fig. 2). There, the Ngoc Linh migmatite, coeval with the D<sub>1</sub> event, encloses



375 highly foliated and folded migmatitic blocks, the block internal foliation and fold axes oblique  
376 to the regional ones (Fig 4D) indicate that they belong to an older pre-D<sub>1</sub> event, in agreement  
377 with zircon U-Pb ages.

378

#### 379 4. Metamorphic and geochronological constraints

380 The knowledge of the thermo-barometric conditions and ages of the tectonic-  
381 metamorphic events described in the previous section is a key issue in attempting to  
382 understand the tectonic evolution of the Kon Tum massif. This section synthesizes the  
383 available radiometric and metamorphic constrains available in the literature for the Kham  
384 Duc, Ngoc Linh, and Kan Nack complexes.

##### 385 2.1. The Kham Duc complex.

386 Thermo-barometric investigations from metapelites and metagreywackes of the Kham  
387 Duc complex indicate that these rocks underwent a polymetamorphic evolution (e.g. Nakano  
388 et al., 2007, 2009, 2013; Osanai et al., 2008; Usuki, 2009; Yonemura et al., 2013). The main  
389 metamorphic assemblage in metapelite is characterized by biotite-garnet-sillimanite, yielding  
390 P and T conditions of 0.7 to 0.8 GPa, and 720-750°C, respectively. Monazite U-Th-Pb  
391 chemical dating of the Kham Duc metapelite gives Middle Triassic ages ranging from 246±3  
392 Ma to 236±9 Ma (Nakano et al., 2009, 2013). Similar Triassic ages, between 257±4 Ma and  
393 245±3 Ma, have been obtained by LA-ICP-MS method on zircon and monazite from  
394 paragneiss and orthogneiss around the Bong Mieu gold mine, between Tra My and Tam Ky  
395 (Tran et al., 2014). Lastly, biotite, muscovite, and amphibole from the Kham Duc  
396 metamorphic rocks, and the mylonites developed along the Tra Bong and Tam Ky shear  
397 zones yield two biotite and muscovite <sup>40</sup>Ar/<sup>39</sup>Ar ages at ca 245 Ma and 230 Ma that are  
398 interpreted as those of the main D<sub>1</sub> metamorphic event, and the D<sub>2</sub> dextral shearing,  
399 respectively (Lepvrier et al., 1997, 2004, Vu et al., 2007).

400           Moreover, garnet, kyanite, and staurolite inclusions in plagioclase document an older  
401 MP/MT event under ca 1.2 GPa and 500-600°C P and T conditions. Furthermore, in this  
402 metamorphic complex, quartz-feldspar aggregates, foliated and lineated during the D<sub>1</sub> event,  
403 represent highly strained magmatic batches formed during an incipient stage of anatexis older  
404 than D<sub>1</sub> (Fig. 3C). Near Phuoc Son, in these migmatitic rocks, zircon rims yield SHRIMP U-  
405 Pb ages at 447±6 Ma, and 452±6 Ma interpreted as those of the migmatization (Usuki et al.,  
406 2009). Thus the age of the HP/MT event took place before 450 Ma, but after 548±7 Ma,  
407 which is the age of the youngest detrital zircon grain found in the metapelite (Usuki et al.,  
408 2009). A Late Ordovician age, at ca 460 Ma, seems likely. An Early Paleozoic, pre-D<sub>1</sub>,  
409 tectonic and metamorphic event, represented by rare microstructures but with clear  
410 metamorphic relicts, was overprinted by the D<sub>1</sub> Early to Middle Triassic event (Nakano et al.,  
411 2009, 2013).

412

#### 413 4.2. The Ngoc Linh complex.

414           The main metamorphic event in the Ngoc Linh complex is a HT/MP one, represented  
415 by biotite-garnet-sillimanite-cordierite migmatite that encloses mafic blocks of HP granulite  
416 with a garnet-clinopyroxene-orthopyroxene-plagioclase-quartz assemblage, and rare eclogites,  
417 (Osanai et al., 2004, 2009; Nakano et al., 2004, 2007, 2009, 2013). The thermo-barometric  
418 conditions for these two metamorphic facies are estimated at ca 900 to 1000°C and 1.3 to 1.6  
419 GPa, for the HT/MP crustal melting, and 900°C-1.6 GPa for the preceding HP/MT one.

420           A wealth of radiometric ages is available for the Ngoc Linh complex (e.g. Nakano et  
421 al., 2013, and enclosed references). Monazite yields U-Pb chemical ages ranging from 247  
422 Ma to 228 Ma. However, old ages between 462 Ma and 432 Ma are also measured in  
423 migmatitic gneiss from the western and southwestern parts of the Ngoc Linh complex. Zircon  
424 from this migmatite yields U-Pb SHRIMP ages at 436±10 Ma (Tran, 2004). Thus, two

425 migmatitic events, namely an Early Paleozoic, and a late Early Triassic one are recognized in  
426 the Ngoc Linh complex. Moreover, NE of Dak To, at Tu Mo Rong, Sm-Nd dating of HT  
427 granulite and gneiss gives isochron ages at  $240\pm 26$  Ma, and  $247\pm 11$  Ma, respectively  
428 (Nakano et al., 2007).

429

#### 430 4.3. The Kan Nack complex.

431 The thermo-barometric conditions for the formation of the Kan Nack metamorphic  
432 rocks have been estimated at ca 1000°C and 1.2 GPa for the mafic granulite (Nakano et al.,  
433 2007, 2013), and 750°C and 0.3-0.4 GPa for the granulitic migmatite (Roger et al., 2007). As  
434 for the Ngoc Linh complex, two groups of ages are recognized there. The central and western  
435 parts of the complex yield Triassic ages. Zircon U-Pb, biotite  $^{40}\text{Ar}/^{39}\text{Ar}$ , and monazite U-Pb  
436 chemical ages range from 255 Ma to 240 Ma (e.g. Maluski et al., 2005; Roger et al., 2007;  
437 Nakano et al., 2007, 2013). Furthermore, in the eastern part of the Kan Nack complex (Song  
438 Bien area), zircon and monazite from migmatites and gneiss yield Early Paleozoic ages,  
439 ranging from 465 Ma to 422 Ma (Roger et al., 2007; Nakano et al., 2013 and enclosed  
440 references).

441

#### 442 4.4. A polycyclic metamorphic evolution of the Kon Tum massif

443 The petrological and geochronological studies from the three metamorphic complexes  
444 allow us to draw their representative P-T-t paths, although the prograde parts of the paths are  
445 poorly documented (Fig. 11A, B). The Kham Duc metamorphic rocks document a kyanite-  
446 garnet-staurolite assemblage indicative of a MP/MT event, followed by a HT one, associated  
447 with crustal melting. This metamorphic evolution occurred during the Early Paleozoic, from  
448 460 Ma to 445 Ma. Moreover, a late Early Triassic thermal event, dated around 250-246 Ma,

449 overprinted the Early Paleozoic cycle. These two different events argue for a polycyclic  
450 evolution of the Kham Duc complex.

451 High temperature, even ultra-high temperature (UHT), metamorphic rocks, coeval  
452 with migmatites, are widespread in the Ngoc Linh and Kan Nack complexes. The later  
453 complex reached higher temperatures (ca 1000°C) than the former. The UHT event was coeval  
454 with melting of the lower crust under HP conditions that produced charnockite and enderbite  
455 (Fig. 11B). The migmatite encloses mafic granulite and eclogite xenoliths that document an  
456 early HP event. The numerous multi-method geochronological studies support a late Early  
457 Triassic to early Middle Triassic age, at ca 250-230 Ma, for the crustal melting. Moreover,  
458 some migmatites also yield Early Paleozoic ages, between 460 and 430 Ma. Thus, the  
459 existence of a polycyclic evolution, as in the Kham Duc complex, is well documented in the  
460 Ngoc Linh and Kan Nack complexes. The accuracy of both the thermo-barometric estimates,  
461 and the radiometric dating, does not allow us to discriminate between the Triassic and Early  
462 Paleozoic migmatites on the basis of their P-T conditions of formation. The two P-T paths  
463 superimposes on the Fig. 11A and B. Moreover, the Early Paleozoic deformed xenolith  
464 migmatite enclosed in the Early Triassic migmatite, observed in the Song Re section (Fig.  
465 4D), provides a field evidence for a polycyclic evolution.

466

## 467 5. Tectonic and geodynamic interpretations

### 468 5.1. The architecture of the Kon Tum massif

469 The structural, petrological, and geochronological data presented in the previous  
470 sections allow us to draw an interpretative crustal scale cross section of the Kon Tum massif,  
471 and the northern domain (Fig. 12, located in figure 2). The metamorphic rocks of the Ngoc  
472 Linh and Kan Nack complexes document a retrograde P-T path coeval with crustal melting,  
473 and top-to-the-North ductile shearing. These two complexes are interpreted here as forming a

474 metamorphic core complex (MCC) exhumed in the footwall of the Ngoc Linh detachment  
475 fault. In spite of some post-S<sub>1</sub> folds, the Kan Nack complex is interpreted as a flat lying unit  
476 exhumed from the granulitic lower crust, as indicated by the HT (or UHT) migmatite,  
477 charnockite, and enderbite overlying the Ngoc Linh complex. The Kan Nack complex  
478 exhumed from the deepest crust overlies the Ngoc Linh migmatitic complex. This extensional  
479 tectonics occurred between 250 Ma and 230 Ma. Both the Ngoc Linh and Kan Nack Triassic  
480 migmatite encloses Early Paleozoic xenoliths of migmatite, HP granulite, and Dai Loc or Ben  
481 Giang-Que Son granitoids. The northern boundary of the Ngoc Linh dome is cut by the D<sub>2</sub>  
482 dextral strike-slip faults of the Tra Bong fault system that mylonitized the Early Paleozoic Tra  
483 Bong granodiorite, and erased the microstructures related to the Ngoc Linh detachment.

484         More to the North, the Kham Duc complex is interpreted as an Early Paleozoic  
485 accretionary complex containing mafic and ultramafic blocks. The southern part of the  
486 complex experienced an Early Paleozoic anatexis, represented by the Chu Lai migmatite. As a  
487 whole, the Kham Duc metamorphic rocks, and the Chu Lai migmatite belong to the Tam Ky-  
488 Phuoc Son suture (Lepvrier et al., 2008; Tran and Vu, 2011; Tran et al., 2014). North of the  
489 Tam Ky-Phuoc Son suture, the Early Paleozoic sedimentary formations are devoid of any  
490 significant metamorphism. A Devonian unconformity, though often reworked by the Triassic  
491 tectonics, argues for an Early Paleozoic folding event (Tran and Vu, 2011; Than et al., 2012).  
492 Upright folds, associated to the Middle Triassic D<sub>2</sub> dextral ductile shearing, deform the entire  
493 Paleozoic succession. The Early Paleozoic Dai Loc plutons are also changed into orthogneiss  
494 during this D<sub>2</sub> shearing (Lepvrier et al., 1997, 2004). Therefore, the Kon Tum massif, and its  
495 northern domain, resulted from a polycyclic evolution. A HT/UHT extensional event,  
496 responsible for the late Early to early Middle Triassic exhumation of the Ngoc Linh and Kan  
497 Nack complexes, followed by a Middle Triassic strike-slip faulting, overprinted the Early  
498 Paleozoic orogen. At the crust-scale, the Tam Ky-Phuoc Son suture separates two different

499 basements. The northern block, tentatively called here the "North Vietnam" or "Viet-Lao  
500 block" underlies the northern domain, and to the South, a southern block, called here the  
501 "South Vietnam" or "Viet-Cambodia" block underlies the suture zone and was partly melted  
502 to form the Ngoc Linh and Kan Nack complexes during the Triassic.

503         On the western side of the Kon Tum massif, an Early Paleozoic ophiolitic suture has  
504 been proposed along the Po Ko fault (Lepvrier et al., 2008). Mafic and ultramafic rocks are  
505 exposed SW of Kon Tum in the Plei Wak complex, and Early Paleozoic Dien Binh  
506 granodiorite crops out west of the fault (Nagy et al., 2011). Our field observations suggest that  
507 the Po Ko fault is a Neogene sinistral strike-slip fault, but older ductile and synmetamorphic  
508 structures are not identified yet. Moreover, migmatite widely develops west of the Po Ko fault  
509 at least up the Laos border. Furthermore, the chemical composition of the main constitutive  
510 minerals (olivine, clinopyroxene, spinel) of the Plei Wak ultramafic body does not support an  
511 ophiolitic setting for this massif (Pham et al., 2006). More detailed work is necessary to  
512 clarify this point. In the following, the Po Ko suture will not be considered. NE of Dak To  
513 (Fig. 2), a dextral Dak To Kan shear zone was proposed (Osanai et al., 2008, Nakano et al.,  
514 2007, 2012). North of Dak To, the NW-SE striking, SW dipping  $S_1$  foliation contains a  
515 subhorizontal mineral and stretching  $L_1$  lineation (Fig.5, stereogram E). This structure appears  
516 different from the other strike-slip faults described in the northern domain both by its "old"  
517 age, around 250 Ma, at variance with the ca 230 Ma age of the Tra Bong shear zone, and by  
518 its HT metamorphic conditions. An alternative interpretation can be proposed if considering  
519 that the present SW dip of the  $S_1$  foliation is not primary but due to the late normal faulting  $D_3$   
520 event. In this case, if  $S_1$  is restored to its subhorizontal initial geometry, the apparent dextral  
521 sense of shear becomes a top-to-the-NW one, consistent with the  $D_1$  kinematics observed in  
522 other parts of the Ngoc Linh dome (Fig. 5).

523

524 5.2. The place of the Kon Tum massif in the Triassic (Indosinian) orogeny.

525           In order to discuss the Paleozoic and Mesozoic evolution of the Indochina block, it is  
526 necessary to reconstruct its Early Triassic paleogeography with respect to South China.  
527 Presently, the Indochina block is bounded to the North by the Ailaoshan-Song Ma ophiolitic  
528 suture that can be correlated to the Song Chay-Hainan, and Jinshajiang suture to the East and  
529 Northwest, respectively (Fig. 13; Faure et al., 2014, 2016b; Xu et al., 2015). Several lines of  
530 evidence lead us to argue that the Indosinian collision occurred in Early to Middle Triassic. It  
531 was preceded by a southwestward (in the present coordinate) subduction, that is to say that the  
532 South China Block (SCB) subducted below the Indochina block that represented the upper  
533 plate of the collision system (see Faure et al., 2014, 2016a, b for details). During the late  
534 orogenic evolution of the Indosinian belt, a series of NW-SE striking dextral shear zones  
535 developed (Lepvrier et al., 1997, 2004).

536           It is well acknowledged that the Indian collision was responsible for an important  
537 disturbance in the Triassic architecture of Indochina. The left-lateral Red River fault (RRF)  
538 accommodated the SE-ward extrusion of Sundaland with respect to the SCB (Tapponnier et  
539 al., 1990). A controversy remains on the amount of displacement along the RRF. Our field  
540 based tectonic reconstruction suggests an offset of ca 500 km close to that proposed by many  
541 authors on geological and paleomagnetic bases (e. g. Leloup et al., 1995; Royden et al., 2008;  
542 Otofujii et al., 2012; Li et al., 2017). Furthermore, the total amount of Triassic dextral motion  
543 that widely distributes between the Song Ma fault, Song Ca, Truong Son, and Tra Bong fault  
544 system, can be estimated to ca 250 to 350 km on the basis of the offset of geological markers,  
545 the km-scale width of each ductile shear zone, and the pervasive deformation associated to  
546 each shear zone. Thus, an Early Triassic palinspastic reconstruction of the Indochina and SCB  
547 relative position just after the collision, but before the late Middle to Early Triassic dextral  
548 faulting, is proposed in figure 14. The ca 500 km Cenozoic sinistral offset along the RRF is

549 partly balanced by the ca 300 km Triassic dextral one. In this qualitative reconstruction, the  
550 Kon Tum massif appears in the upper plate of the Indochina orogen, about 600 km southward  
551 of the Song Ma-Song Chay ophiolitic suture zone. The question of the heat source able to  
552 supply a sufficient temperature elevation to generate the UHT granulite facies metamorphism  
553 must be addressed.

554         A late orogenic thermal overprint is common in many collisional orogens such as the  
555 Variscan belt of Western Europe (e. g. Ledru et al., 2001; Martinez-Catalan et al., 2009;  
556 Schulman et al., 2008; Faure et al., 2010; Lardeaux et al. 2014), the Himalaya (Le Fort et al.,  
557 1987), or the Qinling-Dabie belt of central China (e.g. Lin et al., 2015; Ji et al., 2017). The  
558 collision related crustal thickening is often acknowledged as the source of the radiogenic heat  
559 responsible for the crustal melting, and subsequent MCC development in the subducted  
560 continental crust of the lower plate. In such a setting, an UHT metamorphism is rare. It also  
561 does not account for the Ngoc Linh MCC as this structure developed in the upper plate of the  
562 collision system. Another proposed case is heat advection in the upper plate lower crust  
563 triggered by asthenospheric mantle upwelling in a subduction setting. There, a slab breakoff  
564 of the subducting lithosphere is a prerequisite to promote UHT granulite facies metamorphism  
565 and crustal melting (e. g. Kelsey and Hand, 2015; Touret et al., 2016, and enclosed  
566 references). In this respect, the Kon Tum massif appears as unusual, as a linear trend of the  
567 UHT metamorphic rocks, parallel to the Triassic belt that would better comply with a slab  
568 breakoff mechanism is expected instead of the "punctual" occurrence observed for the Ngoc  
569 Linh MCC.

570         It is worth noting that during the Late Permian, between 263 Ma and 257 Ma, the  
571 Emeishan-Song Da-Tu L  Large Igneous Province (LIP, Ali et al., 2005; Schellnutt et al.,  
572 2015; Fig 14) was emplaced in the western part of the SCB. The present separation between  
573 the Emeishan LIP and the Song Da-Tu L  area is a consequence of the Cenozoic sinistral



574 faulting along the RRF (Fig. 13). Assuming that the mantle plume that produced the  
575 Emeishan-Song Da-Tu L<sup>ê</sup> LIP at ca. 260 Ma (e.g. Schellnutt et al., 2015) remained fixed with  
576 respect to the asthenospheric mantle, and that the SCB-Indochina lithospheric system moved  
577 northward with a 80 km/Ma velocity, as currently measured for the present Pacific or  
578 Philippine Sea plates (e.g. Gripp and Gordon, 2002), the Kon Tum massif was overlying the  
579 mantle plume in late Early Triassic, at ca 245 Ma (Fig. 15). The mantle plume-related heat  
580 flow triggered the melting of the lower crust, giving rise to the charnockite and enderbites of  
581 the Kan Nack complex, and subsequently the middle crustal melting in the Ngoc Linh  
582 complex. The tectonically assisted upward migration of the melted continental crust led to the  
583 extensional emplacement of the Ngoc Linh MCC. The top-to-the-NW sense of shear might be  
584 a consequence of the N-dipping Early Paleozoic inherited crustal structure. The importance of  
585 a mantle source for the heat flux required to produce the UHT metamorphism of the Kon Tum  
586 massif was also documented by petrography and geochemistry (Owada et al., 2006, 2016;  
587 Osanai et al., 2008). Thus, the role of the Emeishan-Song Da LIP, related to a mantle plume,  
588 as a possible heat source for the melting of the continental crust in the Kon Tum massif must  
589 be seriously considered.

590         The interaction between the rise of a mantle plume, and a subduction zone is a  
591 complex issue, still poorly documented. This question has been addressed for an oceanic  
592 subduction in several natural examples, such as the Cordilleran Laramide belt of western US  
593 (Murphy et al., 1998), in the W. Gondwana region (Dalziel et al., 2000), and in the Central  
594 Asia Orogenic Belt (Betts et al., 2015). Numerical and analogue modelings shed light on the  
595 active thermo-mechanical processes (Fletcher and Wyman, 2015; Betts et al., 2015, Mériaux  
596 et al., 2016). These studies emphasize the development of extensional tectonics in the upper  
597 crust of the upper plate overlying the plume. The case of continental subduction, as in SCB-  
598 Indochina area, is more complex than that of an oceanic subduction, as asthenospheric magma

599 upwelling must cross through the subducting continental lithosphere. A slab break off, as also  
600 assumed in the Western US (Murphy et al., 1988), would probably permit heat transfer from  
601 the mantle plume to the lithosphere necessary to enhance the magmatic underplating. Osanai  
602 et al., (2008) already suggested such a mechanism, but the northward subduction of Indochina  
603 block below SCB invoked by the authors is not in agreement with the magmatic arc on the  
604 Indochina block side. Presently, the geochemical and geophysical records that would allow us  
605 to test this interpretation are lacking. Thus, this geodynamic scenario, schematically depicted  
606 in Figure 15, must be considered as a working hypothesis.

607

### 608 5.3. The Early Paleozoic evolution of the Kon Tum massif.

609 Early Paleozoic ages are widespread in the Ngoc Linh MCC, but contemporaneous  
610 structures are rare, except in the Kham Duc complex where intrafolial folds might be ascribed  
611 to the pre-D<sub>1</sub> event. In the Kham Duc complex, various blocks of sedimentary or volcani-  
612 clastic rocks, amphibolite, metagabbro, and serpentinites enclosed into a metasandstone matrix  
613 has been recognized as an Early Paleozoic accretionary complex, also called the Tam Ky-  
614 Phuoc Son suture (Izokh et al., 2006; Pham et al., 2006; Lepvrier et al., 2008; Tran and Vu,  
615 2011; Tran et al., 2014). This plate boundary represents the place where the N. Vietnam (or  
616 Viet-Lao) block and the S. Vietnam (or Viet-Cambodia) block were welded together. Thus  
617 the question of the subduction polarity arises. The existence of an Early Paleozoic magmatic  
618 arc is supported by the calc-alkaline granodiorites of the Dien Binh and Tra Bong suites (Fig.  
619 2), though the available ages, between 445 Ma and 415 Ma (Nagy et al., 2001; Carter et al.,  
620 2001), are younger than the late orogenic crustal melting of the Kham Duc, and Ngoc Linh  
621 complexes dated at ca 450 Ma (Roger et al., 2007; Usuki et al., 2009; Nakano et al., 2013). In  
622 Laos, close to the study area, a calc-alkaline granodiorite similar to the Dien Binh suite  
623 yielded a zircon U-Pb LA-ICP-MS age of 470 Ma (Gardner et al., 2017). Moreover, a zircon

624 LA-ICP-MS age at  $479\pm 3$  Ma (Pham and Huynh, 2016) from a calc-alkaline quartz diorite in  
625 an Early Permian Ben Giang-Que Son pluton could be interpreted as a relict grain inherited  
626 from an Early Paleozoic magmatic arc formed in a continental active margin (Tran et al.,  
627 2008; Pham and Huynh, 2016).

628 From the litho-stratigraphic point of view, south of Hue (Fig. 2), in the Binh Tri Thien  
629 zone, or South Truong Son area, the Ordovician rocks of the Long Dai Formation consist of  
630 rhythmic alternations of sandstone, siltstone, mudstone, with intercalations of radiolarian  
631 chert, and andesitic layers (Tran and Vu, 2011, Thanh et al., 2013). The Early Silurian  
632 deposits consist of sandstone and tuffaceous shales, but lack andesitic rocks. Lastly, the Late  
633 Silurian (Ludlow, 423-419 Ma) Dai Giang Formation unconformably covers the Ordovician-  
634 Early Silurian rocks. Thus, an Ordovician-Early Silurian magmatic arc located to the North  
635 of the Kon Tum massif seems likely, although its precise location remains uncertain yet.

636 The Early Paleozoic (ca 450 Ma) late orogenic anatexis is documented by the Chu Lai  
637 migmatite in the southern part of the Kham Duc complex (Usuki et al., 2009). The Early  
638 Paleozoic age, at  $436\pm 10$  Ma, of the foliated migmatite xenoliths (Fig 4D; Nam et al. 2004)  
639 enclosed in the Triassic Ngoc Ling dome is also considered here as an evidence for the Early  
640 Paleozoic orogeny. During the late orogenic stage, after the collision, the melting of  
641 metapelites that formed the Dai Loc S-type plutons is dated between 427 Ma and 423 Ma  
642 (Fig. 16). In the same way, the Early Paleozoic HP rocks, granulitic and eclogitic relics,  
643 enclosed in the Ngoc Linh and Kan Nack complexes (Roger et al., 2007; Nakano et al., 2004,  
644 2007, 2013), might represent remnants of the deep-seated part of the Early Paleozoic belt of  
645 Central Vietnam. In most of orogens in the world, the highly metamorphosed and deformed  
646 rocks belong to the lower plate that underwent subduction. In Central Vietnam, the Early  
647 Paleozoic migmatites and metamorphic rocks are located to the south of the Tam Ky-Phuoc  
648 Son suture, whereas, north of the suture, the pre-Devonian rocks are represented by

649 sedimentary and volcanic formations. Therefore, in agreement with the presence of an Early  
650 Paleozoic andesitic magmatism North of the Tam Ky-Phuoc Son suture, a north-directed  
651 subduction is preferred here (Fig. 16). In the present state of knowledge, except the N-S  
652 striking mineral lineation observed in the Chu Lai migmatite, the microstructures and  
653 associated kinematic indicators related to the Early Paleozoic orogen have not been  
654 recognized yet.

655

656 5.4. A possible interaction between the Early Paleozoic belts of Indochina and South China.

657         The collision between the N. Vietnam (or Viet-Lao) block and the S. Vietnam (or  
658 Viet-Cambodia) block was responsible for the development of the Tam Ky-Phuoc Son suture  
659 (Fig. 13). This event might have also an important significance for the geodynamic evolution  
660 of Asia. This section explores some possible relationships between the pre-Devonian belts of  
661 South China and the Tam Ky-Phuoc Son belt. In South China, two Early Paleozoic orogenic  
662 belts are recognized (Figs. 13, 14). To the north, the Qinling-Qilian orogen resulted of the  
663 collision between the North China and South China blocks. The age of the collision remains  
664 disputed, although a Triassic collision is often put forward, several lines of evidence argue for  
665 a pre-Devonian collisional event, at ca 420 Ma (e.g. Mattauer et al., 1985; Faure et al., 2008,  
666 Dong et al., 2015; Shi et al., 2016). Nevertheless, whatever its real age, the SCB-NCB  
667 collision is younger than the Early Paleozoic tectonics in Indochina, and it will not be  
668 considered in the following.

669         On the contrary, the central part of the SCB is occupied by a NE-SW striking  
670 intracontinental orogen characterized by a south-directed fold-and-thrust belt, a top-to-the-  
671 South ductile decollement, and a polyphase metamorphism. An Early MP/MT event, dated at  
672 ca 460-455 Ma, reworked by a HT/LP event coeval with migmatites and peraluminous  
673 granites, dated between 440 Ma and 400 Ma, has been identified (for details see Faure et al.,

674 2009; Charvet et al., 2010, Li et al. 2010). This belt, devoid of ophiolites, subduction  
675 complex, and magmatic arc is an intracontinental orogen that resulted from the closure of a  
676 Neoproterozoic-Cambrian rift accommodated by a North-directed continental subduction  
677 (Faure et al., 2009; Charvet et al., 2010). The cause of this intracontinental orogeny was not  
678 understood yet. A collision between the main part of SCB and Australia, a piece of which is  
679 supposed to be located in Hainan island, has been proposed (Xu et al., 2016). However, the  
680 geological record for Early Paleozoic arc magmatism, and ophiolitic suture are missing in  
681 Hainan island. As no other Early Paleozoic ophiolitic suture is recognized between the SCB  
682 intracontinental orogen and the Tam Ky-Phuoc Son suture, the SCB and N. Vietnam block  
683 belonged to the same continent until the Devonian (Metcalf, 2013). In late Cambrian-Early  
684 Ordovician, the S. Vietnam (Viet-Khmer) continent subducted below the N. Vietnam (Viet-  
685 Lao) that was a part of the SCB, within which an intracontinental rift developed (Fig. 16). The  
686 Middle to Late Ordovician (460-450Ma) collision between the N. Vietnam and the S.  
687 Vietnam blocs led to the closure of the intracontinental rift of the SCB. During the Silurian,  
688 the late- to post-collisional evolution was responsible for the crustal melting experienced by  
689 the S. Vietnam block, and SCB. Therefore, we propose here that the collision between the N.  
690 Vietnam (Viet-Lao) and S. Vietnam (Viet-Cambodia) blocks induced stress propagation in the  
691 SCB, and was responsible for the closure of the Neoproterozoic-Cambrian rift of SCB, and  
692 the related deformation and metamorphism. The Early Paleozoic intracontinental orogeny of  
693 South China may be considered as a consequence of a far-field effect of the central Vietnam  
694 collision (Fig. 16), in the same way as the Central Asia Cenozoic structures resulted from the  
695 Himalayan collision. This SCB-Viet-Lao continental mass was split during the Carboniferous  
696 by the opening of a branch of the Paleotethys Ocean (Nguyen et al., 2013) that was closed in  
697 the late Early to early Middle Triassic by the Indosinian orogeny.  
698

699

## 700 6. Conclusion

701           The Kon Tum massif of Central Vietnam provides important clues for understanding  
702 the restless geodynamic movements between the SCB and Indochina block. The Kon Tum  
703 massif of Central Vietnam cannot be simply considered as the Precambrian nucleus of  
704 Indochina. This massif experienced a polycyclic evolution. Although structural evidence is  
705 poorly preserved, the geochronological and petrological records indicate that an Early  
706 Paleozoic (at ca 460-450 Ma) collision, along the Tam Ky-Phuoc Son suture. The syn-  
707 metamorphic structures preserved in the Kon Tum massif developed during Early Triassic (at  
708 ca 250-240 Ma) within an extensional tectonic setting, giving rise to pervasive crustal melting  
709 and the formation of the Ngoc Linh metamorphic core complex.

710           The Early Paleozoic collision was responsible for the welding of the S. Vietnam (or  
711 Viet-Cambodia) block and the N. Vietnam (or Viet-Lao) block that was at that time a part of  
712 the SCB. This collision is also interpreted here as the driving force that closed the  
713 Neoproterozoic-Early Paleozoic Nanhua continental rift basin, and triggered the formation of  
714 the intracontinental orogen of South China. Since the Early Devonian, the SCB, Viet-Lao, and  
715 Viet-Cambodia blocks formed a single continental mass characterized by similar faunistic and  
716 floristic fossil assemblages and sedimentary facies (Thanh et al., 1996; Janvier et al., 1997,  
717 1998; Metcalfe, 2002, 2013). This continent was subsequently disrupted by the opening of the  
718 Paleotethys ocean in Carboniferous times (Nguyen et al., 2013). Since that time, the SCB and  
719 Indochina became separated. The Indochina block appears as a composite continent formed,  
720 at least by the Viet-Lao and Viet-Cambodia blocks. The Ailaoshan-Song Ma-Song Chay-  
721 Hainan ophiolitic suture is now acknowledged as the plate boundary, the collision age is no  
722 longer a matter of debate. The closure of the Paleotethys ocean was accommodated by a  
723 south-directed subduction, as shown by the Permian Sam Nua magmatic arc, followed by the

724 continental subduction of the SCB below Indochina. The Song Ma and Song Chay ophiolitic  
725 sutures belong to a thousands of km long collisional belt that extends, at least, from Eastern  
726 Tibet to Hainan, and probably farther East (see discussion in Faure et al., 2016a).  
727 Within this general scenario, the tectonic survey of the Kon Tum massif indicates that  
728 the Early to Middle Triassic Ngoc Linh MCC developed in the upper plate of the collision  
729 system. The radiogenic crust of Indochina thickened by the Early Triassic collision was not  
730 able to supply the important heat input required to form the UHT metamorphic rocks, such as  
731 granulites, charnockites, and enderbites, in the lower continental crust. The mantle plume that  
732 formed the Emeishan LIP in Late Permian might have represented the main heat source, as  
733 this plume afterwards underlain Indochina in late Early to early Middle Triassic. Forthcoming  
734 structural, petrological, and geochemical investigations will provide new lines of evidence to  
735 refine the geodynamic scenario of this continental ballet.

736

737 Acknowledgements. This work was supported by the Research Program of the Vietnam  
738 National University (grant QG. 18.14). A part of the expenses for M. F. have been also  
739 provided by UMR 7327 CNRS-Université d'Orléans, and CNRS INSU Syster-TelluS  
740 program. J. Charvet and Y. Chen are acknowledged for fruitful discussions, and suggestions  
741 to improve the manuscript. Professor C. Morley, and two anonymous reviewers are  
742 thanked for their thorough and insightful reviews, and valuable suggestions to improve  
743 the quality of the manuscript, and Professor Khin Zaw for handling the manuscript and  
744 editorial input.

745

746 References

747 Ali, J., Thompson, G.M., Zhou, M.F., Song, X., 2005. Emeishan large igneous province,  
748 SW China. *Lithos* 79, 475–499.

749

750 Betts, P. G., Moresi, L., Miller, M. S., David Willis, D., 2015. Geodynamics of oceanic  
751 plateau and plume head accretion and their role in Phanerozoic orogenic systems of China.  
752 *Geoscience Frontiers*, 6, 49-59.

753

754 Carter, A., Roques, D., Bristow, C., Kinny, P., 2001. Understanding Mesozoic accretion  
755 in Southeast Asia: significance of Triassic thermotectonism (Indosinian orogeny) in Vietnam.  
756 *Geology* 29, 211–214.

757

758 Carter, A., Clift, P.D., 2008. Was the Indosinian orogeny a Triassic mountain building  
759 or a thermo-tectonic reactivation event? *Comptes Rendus Geoscience* 340, 83-93.

760

761 Cau D.V., Cau, N.C., Chanh, D.D., Linh D.V., Long T.V., Son, N.N., Vinh, M.K and Vu,  
762 N.C 2004. Report on mapping and mineral resources investigation of Ba To sheet group,  
763 scale 1:50.000. General Department of Geology and Minerals of Vietnam (in Vietnamese).

764

765 Charvet, J., Shu, L.S., Faure, M., Choulet, F., Wang, B., Le Breton, N., 2010.  
766 Structural development of the Lower Paleozoic belt of South China: genesis of an  
767 intracontinental orogen. *J. Asian Earth Sci.* 39, 309–330.

768

769 Dalziel, I. W. D., Lawver, L. A., Murphy, J. B., 2000. Plumes, orogenesis, and super-  
770 continental fragmentation. *Earth and Planetary Science Letters*, 178, 1-11

771



- 772 Dinh, Q. S. 2011. Petrographic Characteristics and Zircon U-Pb Geochronology of granitoid  
773 rocks in the Southern B n Giang, Quang Nam Province, Science & Technology  
774 Development, 14, 17-30.  
775
- 776 Dong, Y., Zhang, X., Liu, X., Li, W., Chen, Q., Zhang, G., Zhang, H., Yang, Z., Sun, S.,  
777 Zhang, F., Propagation tectonics and multiple accretionary processes of the Qinling Orogen,  
778 Journal of Asian Earth Sciences 104, 84–98.  
779
- 780 Faure, M., Lin, W., Moni , P., Meffre, S., 2008. Palaeozoic collision between the North and  
781 South China blocks, Triassic intracontinental tectonics, and the problem of the ultrahigh-  
782 pressure metamorphism, C. R. Geoscience 340, 139–150.  
783
- 784 Faure, M., Shu, L.S., Wang, B., Charvet, J., Choulet, F., Moni , P., 2009. Intracontinental  
785 subduction: a possible mechanism for the Early Paleozoic Orogen of SE China. Terra Nova  
786 21, 360–368.  
787
- 788 Faure, M., Cocherie, A., B -M z me, E., Charles, N., Rossi P., 2010. Middle Carboniferous  
789 crustal melting in the Variscan Belt: New insights from U-Th-Pb<sub>tot</sub>. monazite and U-Pb zircon  
790 ages of the Montagne Noire Axial Zone (southern French Massif Central). Gondwana  
791 Research, 18, 633-673.  
792
- 793 Faure, M., Lepvrier, C., Nguyen, VV., Vu, VT, Lin, W, Chen, Z. 2014. The South China  
794 block-Indochina collision: Where, when, and how? J. Asian Earth Sci. 79, 260–274.  
795

- 796 Faure, M., Lin, W., Yang, C., Lèpvrier, C. 2016a. Triassic tectonics of the southern margin of  
797 the South China Block. *Comptes Rendus Geoscience*, 328, 5-14.  
798
- 799 Faure M., Lin W., Chu Y. Lèpvrier C., 2016b. Triassic tectonics of the Ailaoshan Belt (SW  
800 China): Early Triassic collision between the South China and Indochina Blocks, and Middle  
801 Triassic intracontinental shearing. *Tectonophysics*, 683, 27–42.  
802
- 803 Fletcher, M., Wyman, D., A., 2015. Mantle plume–subduction zone interactions over the past  
804 60 Ma. *Lithos*, 233, 162–173.  
805
- 806 Findlay, R.H., 1997. The Song Ma anticlinorium, northern Vietnam: the structure of an  
807 allochthonous terrane containing an early Paleozoic island arc sequence. *Journal of Asian  
808 Earth Sciences* 15, 453–464.  
809
- 810 Findlay, R.H., Pham, T.T., 1997. The structural setting of the Song Ma region, Vietnam and  
811 the Indochina-South China plate boundary problem. *Gondwana Research* 1, 11–33.  
812
- 813 Fromaget, J. 1927. J., Études géologiques sur le Nord de l’Indochine centrale, *Bull. Serv.  
814 Geol. Indochine* 16, 1-368.  
815
- 816 Fromaget, J., 1941. L’Indochine française, sa structure géologique, ses roches, ses mines et  
817 leurs relations possibles avec la tectonique. *Bulletin du Service Géologique de l’Indochine* 26,  
818 1–140.  
819

- 820 Gripp, A.E., Gordon, R.G., 2002. Young tracks of hotspots and current plate velocities.  
821 Geophys. J. Int., 150, 321–361  
822
- 823 Hieu P. T., Trung H., 2016. U-Pb zircon age of quartz diorite from Ben Giang intrusive  
824 complex in the Ben Giang area, Quang Nam Province. Vietnam Journal of Earth Sciences, 37,  
825 156-162.  
826
- 827 Hieu P. T., Yang, Y. Z., Binh, D. Q., Nguyen, T. B. T. N., Dung, L.T., Chen, F., 2015. Late  
828 Permian to Early Triassic crustal evolution of the Kontum massif, central Vietnam: zircon U–  
829 Pb ages and geochemical and Nd–Hf isotopic composition of the Hai Van granitoid complex.  
830 International Geology Review, 57, 1877–1888.  
831
- 832 Hieu P. T., Trung H., 2016. U-Pb zircon age of quartz diorite from Ben Giang intrusive  
833 complex in the Ben Giang area, Quang Nam Province. Vietnam Journal of Earth Sciences, 37,  
834 156-162.  
835
- 836 Hieu, P. T., Dung N. T., Nguyen T. B. T., Minh, N. T. Minh, P. 2016. U–Pb ages and Hf  
837 isotopic composition of zircon and bulk rock geochemistry of the Dai Loc granitoid complex  
838 in Kontum massif: Implications for early Paleozoic crustal evolution in Central Vietnam.  
839 Journal of Mineralogical and Petrological Sciences, 1-16.  
840
- 841 Hoffet, J.H., 1933. Etudes géologiques sur le centre de l'Indochine entre Tourane et le  
842 Mékong (Annam Central et Bas Mékong). Bull. Service. Geol. Indochine, 20, pp. 154.  
843

- 844 Hutchison, C.S. 1989. Geological evolution of South-East Asia, Oxford Monographs on  
845 Geology and Geophysics, 13, Clarendon Press, Oxford, UK, 368pp.  
846
- 847 Izokh, A.E., Tran, T.H., Ngo, T.P., Tran, Q.H., 2006. Ophiolite ultramafic-mafic associations  
848 in the northern structure of the Kon Tum block (Central Vietnam). *J. of Geology, Hanoi*, 28,  
849 20-26.  
850
- 851 Janvier, P., Pham Kim, N., Ta Hoa, P., 1996. Une faune de vertébrés de type sud chinois  
852 dans le Dévonien inférieur de la basse Rivière Noire. *Comptes Rendus Académie des*  
853 *Sciences* 323 (II), 539-546.  
854
- 855 Janvier, P., Tong-Dzuy, T., Ta Hoa, P., Doan Nhât, T., 1997. The Devonian vertebrates  
856 (Placodermi, Sarcopterygii) from Central Vietnam and their bearing on the Devonian  
857 paleogeography of Southeast Asia. *Journal of Asian Earth Sciences* 15, 393–406.  
858
- 859 Ji, W., Lin, W., Faure, M., Shi, Y., Wang, Q., 2017. The early Cretaceous orogen-scale  
860 Dabieshan metamorphic core complex: implications for extensional collapse of the Triassic  
861 HP–UHP orogenic belt in east-central China. *Int. J. Earth Sci.*, 106, 1311–1340.  
862
- 863 Kelsey, D.E., Hand, M., 2015. On ultrahigh temperature crustal metamorphism: phase  
864 equilibria, trace element thermometry, bulk composition, heat sources, timescales and tectonic  
865 settings. *Geoscience Frontiers* 6, 311-356.  
866

- 867 Lan, C.Y., Chung, S.L., Long, T.V., Lo, C.H., 2003. Geochemical and Sr–Nd isotopic  
868 constraints from the Kontum massif, central Vietnam on the crustal evolution of the Indochina  
869 block. *Precambrian Research*, 22, 7–27.
- 870
- 871 Lardeaux JM., Schulmann K., Faure M., Janousek V., Lexa O., Skrzypek E., Edel J-B.,  
872 Stipska P., 2014. The Moldanubian Zone in French Massif Central, Vosges/ Schwarzwald and  
873 Bohemian Massif revisited: Differences and similarities. In « The Variscan Orogeny: Extent,  
874 Timescale and the Formation of the European Crust ». Schulmann, K., Martínez Catalán, J.  
875 R., Lardeaux, J. M., Janousek, V. & Oggiano, G. (eds). Geological Society, London, Special  
876 Publications, 405, 1-39
- 877
- 878 Ledru, P., Courrioux, G., Dallain, C., Lardeaux, J. M., Montel, J. M., Vanderhaeghe, O. &  
879 Vitel, G., 2001. The Velay Dome (French Massif Central): melt generation and granite  
880 emplacement during orogenic evolution. *Tectonophysics* 342, 207–237.
- 881
- 882 Le Fort, P., Cuney, M., Deniel, C., France-Lanord, C., Sheppard, S.M.F., Upreti, B.N., Vidal,  
883 P. 1987. Crustal generation of the Himalayan leucogranites. *Tectonophysics* 134, 39-57.
- 884
- 885 Leloup, P.H., Lacassin, R., Tapponnier, P., Schärer, U., Zhong, D.L., Liu, X.H., Zhang, L.S.,  
886 Ji, S.C., Phan, T.T., 1995. The Ailao Shan-Red River shear zone (Yunnan, China), Tertiary  
887 transform boundary of Indochina. *Tectonophysics* 251, 3–84.
- 888
- 889 Lepvrier, C., Maluski, H., Nguyen Van Vuong, Roques, D., Axente, V., Rangin, C., 1997.  
890 Indosinian NW-trending shear zones within the Truong Son belt (Vietnam):  $^{40}\text{Ar}$ - $^{39}\text{Ar}$   
891 Triassic ages and Cretaceous to Cenozoic overprints. *Tectonophysics* 283, 105–127.

892

893 Lepvrier, C. Maluski, H., Vu Van Tich, Leyreloup, A., Phan Truong Thi, Nguyen Van  
894 Vuong, 2004. The Early Triassic Indosinian orogeny in Vietnam (Truong Son Belt and  
895 Kontum Massif): Implications for the geodynamic evolution of Indochina, *Tectonophysics*  
896 393, 87-118.

897

898 Lepvrier, C., Nguyen Van Vuong, Maluski, H., Thi, Phan Truong, Vu Van Tich, 2008.  
899 Indosinian tectonics in Vietnam. *Comptes Rendus Geoscience* 340, 94–111.

900

901 Lepvrier, C., Faure, M., Nguyen Van Vuong, Vu Van Tich, Lin, W., Thang Ta Trong,  
902 Phuong Ta Hoa, 2011. North-directed Triassic nappes in Northeastern Vietnam  
903 (East Bac Bo). *Journal of Asian Earth Sciences* 41, 56–68.

904

905 Li S., Advokaat, E., van Hinsbergen, D., Koymans, M., Deng, C., Zhu, R., 2017.  
906 Paleomagnetic constraints on the Mesozoic-Cenozoic paleolatitudinal and rotational history  
907 of Indochina and South China: Review and updated kinematic reconstruction. *Earth-Science*  
908 *Reviews*, 171, 58-77.

909

910 Li, Z.X., Li, X.H., Wartho, J.A., Clark, C., Li, W.X., Zhang, C.L., Bao, C.M., 2010. Magmatic  
911 and metamorphic events during the Early Paleozoic Wuyi–Yunkai orogeny, southeastern  
912 South China: new age constraints and pressure–temperature conditions. *Geological*  
913 *Society of America Bulletin* 122, 772–793.

914

- 915 Lin, W., Ji, W., Faure, M., Wu, L., Li, Q., Shi, Y., Scharer, U., Wang, F., and Wang, Q.,  
916 2015. Early Cretaceous extensional reworking of the Triassic HP-UHP metamorphic orogen  
917 in Eastern China. *Tectonophysics*, 662, 256-270.
- 918
- 919 Liu, J.L, Tran, M., Tang, Y., Nguyen, Q.L., Tran, T.H., Wu, W., Chen, J., Zhang, Z., Zhao,  
920 Z., 2012a. Permo-Triassic granitoids in the northern part of the Truong Son belt, NW  
921 Vietnam: geochronology, geochemistry and tectonic implications. *Gondwana Research* 122,  
922 628–644.
- 923
- 924 Maluski, H., Lepvrier, C., Leyreloup, A., Vu, Van Tich, Truon Thi, Phan, 2005.  
925  $^{40}\text{Ar}$ - $^{39}\text{Ar}$  geochronology of the charnokites and granulites of the Kan Nack  
926 complex, Kon Tum Massif, Vietnam. *Journal of Asian Earth Science* 25, 653–677.
- 927
- 928 Martinez Catalan, J. R., Arenas, R. et al. , 2009. A rootless suture and the loss of the roots of a  
929 mountain chain: the Variscan belt of NW Iberia. *Comptes Rendus Géosciences*, 341, 114–  
930 126.
- 931
- 932 Mattauer, M., Matte, Malavieille, J., Tapponnier, P., Maluski, H., Xu, Z., Lu, Tang, Y., 1985.  
933 Tectonics of the Qinling Belt: buildup and evolution of eastern Asia, *Nature* 317, 496–500.
- 934
- 935 Mériaux, C. A., Mériaux, A. S., Schellart, W.P., Duarte, J. C., Duarte, S. S, Chen, Z., 2016.  
936 Mantle plumes in the vicinity of subduction zones. *Earth and Planetary Science Letters*, 454,  
937 166–177.
- 938

- 939 Metcalfe, I., 1996. Pre-Cretaceous evolution of SE Asian terranes. In: Hall, R. & Blundell, D.  
940 (Ed.), Tectonic Evolution of Southeast Asia. Geological Society Special Publication, vol. 106,  
941 pp. 97–122.
- 942
- 943 Metcalfe, I., 2013. Gondwana dispersion and Asian accretion: tectonic and palaeogeographic  
944 evolution of eastern Tethys. *Journal of Asian Earth Sciences* 66, 1–33.
- 945
- 946 Murphy, B., Oppliger, G., Brimhall, G., 1998. Plume-modified orogeny: An example from the  
947 western United States. *Geology*, 26, 731–734.
- 948
- 949 Nagy, E.A., Maluski, H., Lepvrier, C., Schärer, U., Truong Thi, Phan, Leyreloup, A., Vu, V.  
950 T., 2001. Geodynamic significance of the Kontum massif in central Vietnam: composite  
951  $^{40}\text{Ar}/^{39}\text{Ar}$  and U–Pb ages from Paleozoic to Triassic. *The Journal of Geology*, Chicago, 109,  
952 755–770.
- 953
- 954 Nakano, N., Osanai, Y., Owada, M., Tran, N.N., Tsunogae, T., Toyoshima, T., Pham B.,  
955 2004. Decompression process of mafic granulite from eclogite to granulite facies under  
956 ultrahigh-temperature condition in the Kontum massif, central Vietnam. *Journal of*  
957 *Mineralogical and Petrological Sciences* 99, 242–256.
- 958
- 959 Nakano, N., Osanai, Y., Owada, M., Ngoc N. T., Toyoshima, T., Binh, P., Tsunogae, T.,  
960 Kagami, H., 2007. Geologic and metamorphic evolution of the basement complexes in the  
961 Kontum Massif, central Vietnam. *Gondwana Research* 12, 438–453.
- 962



- 963 Nakano, N., Osanai, Y., Owada, M., Hayasaka, Y., Ngoc N., T., 2009. Permo-Triassic  
964 Barrovian-type metamorphism in the ultrahigh-temperature Kontum Massif, central Vietnam:  
965 constraints on continental collision tectonics in Southeast Asia. *The Island Arc* 18, 126–143.  
966
- 967 Nakano, N., Osanai, Y., Sajejev, K., Hayasaka, Y., Miyamoto, T., Nguyen Thi Minh, Owada,  
968 M., 2010. Triassic eclogite from northern Vietnam: inferences and geological significance.  
969 *Journal of Metamorphic Geology* 28, 59–76.  
970
- 971 Nakano, N., Osanai, Y., Owada, M., Nam, T.N., Charusiri, P., Khamphavong, K., 2013.  
972 Tectonic evolution of high-grade metamorphic terranes in central Vietnam: constraints  
973 from large-scale monazite geochronology. *Journal of Asian Earth Sci.* 73, 520–539.  
974
- 975 Nam, T.N., Sano, Y., Terada, K., Toriumi, M., Quynh, P.V., Dung, L.T., 2001. First SHRIMP  
976 U–Pb zircon dating of granulites from the Kontum massif (Vietnam) and tectono-thermal  
977 implications. *Journal of Asian Earth Sciences*, 19, 77–84.  
978
- 979 Nguyen, V.V., Hansen, B., Wemmer, K., Lepvrier, C., Vu, V.T., Ta, T.T., 2013. U/Pb and  
980 Sm/Nd dating on ophiolitic rocks of the Song Ma suture zone (northern Vietnam): evidence  
981 for Upper Paleozoic Paleotethyan lithospheric remnants. *J. of Geodynamics* 69, 140–147.  
982
- 983 Osanai, Y., Nakano, N., Owada, M., Nam, T.N., Toyoshima, T., Tsunogae, T., Pham, B.,  
984 2004. Permo-Triassic ultrahigh-temperature metamorphism in the Kontum massif, central  
985 Vietnam. *Journal of Mineralogical and Petrological Sciences*, 99, 224–241.  
986
- 987 Osanai, Y., Nakano, N., Owada, M., Ngoc Nam, Tran, Miyamoto, T., Thi Minh, Nguyen,

- 988 Van Nam, Nguyen, Van Tri, Tran, 2008. Collision zone metamorphism in Vietnam and  
989 adjacent Southeastern Asia: proposition for Trans-Vietnam Orogenic Belt. *Journal of*  
990 *Mineralogical and Petrological Sciences* 103, 226–241.
- 991
- 992 Otofujii, Y., Trung, V.D., Fujihara, M., Tanaka, M., Yokoyama, M., Kitada, K., Zaman, H.,  
993 2012. Tectonic deformation of the southeastern tip of the Indochina Peninsula during its  
994 southward displacement in the Cenozoic time. *Gondwana Research* 22, 615-627.
- 995
- 996 Owada, M., Osanai, Y., Hokada, T., Nakano, N., 2006. Timing of metamorphism and  
997 formation of garnet granite in the Kontum massif, central Vietnam: evidence from monazite  
998 EMP dating. *Journal of Mineralogy and Petrological Sciences* 101, 324–328.
- 999
- 1000 Owada, M., Osanai, Y., Nakano, N., Adachi, T., Kitano, I., Tran, V.T., Kagami, H., 2016.  
1001 Late Permian plume-related magmatism and tectonothermal events in the Kontum massif, in  
1002 central Vietnam. *J. Mineral. Petrol. Sciences*, 111, 181-195.
- 1003
- 1004 Pham, T.D., Tran, T.H., Ngo, T.P., Tran, T.A., Bui, A.N., 2006. Characteristics of mineral  
1005 composition (Olivine, Pyroxene, Chromian Spinel) of ultramafic intrusions in the edge of the  
1006 Kon Tum geoblock. *Journal of Geology, Hanoi, Series B*, 47-57.
- 1007
- 1008 Roger, F., Maluski, H., Leyreloup, A., Lepvrier, C., Truong Tri, Phan, 2007. U–Pb dating  
1009 of high temperature metamorphic episodes in the Kon Tum Massif (Vietnam). *Journal of*  
1010 *Asian Earth Sciences* 30, 565–572.
- 1011

- 1012 Roger, F., Jolivet, M., Malavieille, J., 2008. Tectonic evolution of the Triassic fold belts of  
1013 Tibet. *C. R. Geoscience* 340, 180–189.  
1014
- 1015 Roger, F., Maluski, H., Lepvrier, C., Vu Van, T., Paquette, J.-L., 2012. LA-ICPMS zircons  
1016 U/Pb dating of Permo-Triassic and Cretaceous magmatisms in Northern Vietnam-geodynamic  
1017 implications. *Journal of Asian Earth Sciences* 48, 72–82.  
1018
- 1019 Royden, L. H., B. C. Burchfiel, R. D. Van der Hilst, 2008. The geological evolution of the  
1020 Tibetan Plateau. *Science* 321, 1054–1058.  
1021
- 1022 Saurin, E., 1935. Etudes géologiques sur l'Indochine du Sud-Est (Sud Annam, Cochinchine,  
1023 Cambodge Oriental). *Bull. Serv. Geol. Indochine* 22, pp. 419.  
1024
- 1025 Schulmann, K., Lexa, O., Stipska, P., Racek, M., Tajcmanova, L., Konopasek, J., Edel, J.B.,  
1026 A. Peschler, A., Lehmann, J., 2008. Vertical extrusion and horizontal channel flow of  
1027 orogenic lower crust: key exhumation mechanisms in large hot orogens? *J. Metamorphic*  
1028 *Geol.*, 26, 273–297.  
1029
- 1030 Shellnutt, G., Usuki, T., Kennedy, A., Chiu, H. Y., 2015. A lower crust origin of some flood  
1031 basalts of the Emeishan large igneous province, SW China. *Journal of Asian Earth Sciences*  
1032 109, 74–85.  
1033
- 1034 Shi, M.F., Lin, F.C., Fan, W.Y., Deng, Q., Cong, F., Tran, M.D., Zhu, H.P., Wang, H., 2015.  
1035 Zircon U–Pb ages and geochemistry of granitoids in the Truong Son terrane, Vietnam:  
1036 Tectonic and metallogenic implications. *J. Asian Earth Sciences* 101, 101–120.

- 1037
- 1038 Shi, G., Wang, H., Huang, C., Yang, S., Song, G. 2016. Provenance and tectonic setting of  
1039 middle-upper Devonian sandstones in the Qinling Orogen (Shanyang area): New insights  
1040 from geochemistry, heavy minerals and tourmaline chemistry. *Tectonophysics* 688, 11-25.  
1041
- 1042 Tapponnier, P., Lacassin, R., Leloup, P.H., Schärer, U., Zhong, D.L., Liu, X.H., Ji, S.C.,  
1043 Zhang, L.S., Zhong, J.Y., 1990. The Ailao Shan/Red River metamorphic belt: Tertiary left  
1044 lateral shear between Indochina and South China. *Nature* 343, 431–437.  
1045
- 1046 Thanh, T-D., Janvier, P., Phuong, T.H., 1996. Fish suggest continental connections between  
1047 the Indochina and South China blocks in Middle Devonian time. *Geology* 24, 571-574.  
1048
- 1049 Thanh, T-D., Phuong, T.H., Janvier, P., Nguyen, H.H., Nguyen, T.T.C., Nguyen T.D., 2013.  
1050 Silurian and Devonian in Vietnam-Stratigraphy and facies. *J. of Geodynamics* 69, 165-185.  
1051
- 1052 Touret, J. HL.R., Santosh, M., Huizenga, J.M. 2016. High-temperature granulites and  
1053 supercontinents. *Geoscience Frontiers* 7, 101-113.  
1054
- 1055 Tran Ngoc Nam, 2004. Zircon U-Pb isotopic age at 436 Ma within the Song Re complex, Kon  
1056 Tum massif. *Journal of Geology, Serie A.*, 281, 3-4 (in Vietnamese).  
1057
- 1058 Tran Duan., Than Duc Duyen., Vu Nhu Hung., Trinh Van Long., Bui Minh Vo., 2006. Report  
1059 on mapping and mineral resources investigation of Kon Tum sheet group, scale 1:50.000.  
1060 General Department of Geology and Minerals of Vietnam (in Vietnamese).  
1061

1062

1063

1064 Tran, T.H., Tran, T.A., Ngo, T.P., Pham, T.D., Tran, V.A., Izokh, A.E., Borisenko, A.S., Lan,  
1065 C.Y., Chung, S.L., Lo, C.H., 2008. Permo-Triassic intermediate-felsic magmatism of the  
1066 Truong Son belt, eastern margin of Indochina. *Comptes Rendus Geoscience* 340, 112-126.

1067

1068 Tran, V.T., Vu, K., Vu (Eds.), 2011. *Geology and Earth Resources of Vietnam*, General Dept  
1069 of Geology, and Minerals of Vietnam, Hanoi, Publishing House for Science and Technology,  
1070 pp. 634.

1071

1072 Tran, H.T., Zaw, K., Halpin, J., Manaka, T., Meffre, S., Lai, C.K., Lee, Y., Le, H.V., Dinh, S.,  
1073 2014. The Tam Ky-Phuoc Son Shear Zone in central Vietnam: Tectonic and metallogenic  
1074 implications. *Gondwana Research*, 26, 144-164.

1075

1076 Usuki, T., Lan, C.-Y., Yui, T.-F., Iizuka, Y., Van Tich, Vu, Tuan Anh, Tran, Okamoto, K.,  
1077 Wooden, J.L., Liou, J.G., 2009. Early Paleozoic medium-pressure metamorphism in central  
1078 Vietnam: evidence from SHRIMP U–Pb zircon ages. *Geoscience Journal* 13, 245–256.

1079

1080 Vu V.T., Maluski, H., Nguyen V. V., 2007. Ar-Ar age of metamorphic and mylonitic rocks  
1081 in northern part of the Kon Tum massif: evidence for the Indosinian movement along shear  
1082 zones between Kon Tum massif and Truong Son belt. *VNU Journal of Science, Earth  
1083 Sciences* 23, 253-264.

1084

- 1085 Xu, Z., Dilek, Y., Cao, H., Yang, J., Robinson, P., Mae, C., Li, H., Jolivet, M., Roger, F.,  
1086 Chen, X., 2015. Paleo-Tethyan evolution of Tibet as recorded in the East Cimmerides and  
1087 West Cathaysides. *Journal of Asian Earth Sciences* 105, 320-337.  
1088
- 1089 Xu, Y. J., Cawood, P., Du, Y.S., 2016. Intraplate orogenesis in response to Gondwana  
1090 assembly: Kwangsi Orogeny, South China. *American Journal of Science* 316, 329–362.  
1091
- 1092 Yonemura, K., Osanai, Y., Nakano, N., Owada, M., Baba, S., 2013. Petrology, geochemistry,  
1093 and origin of metamorphosed mafic rocks of the Trans Vietnam Orogenic Belt, Southeast  
1094 Asia. *Journal of Mineralogical and Petrological Sciences* 108, 55–86.

## 1095 Figure Captions

1096

1097 Fig. 1. Tectonic sketch of SE Asia with emphasis of the Early-Middle Triassic (Indosinian),  
1098 and Late Triassic-Early Jurassic (Cimmerian) sutures. Modified from Sone and Metcalfe,  
1099 2008; and Lepvrier et al., 1997).

1100

1101 Fig 2. Simplified litho-structural map of the Kontum Massif and the northern area (South  
1102 Truong Son). The dashed line between Dak To and Tu Mo Rong is the Dak To Kan shear  
1103 zone proposed by Nakano et al. (2007, 2012); and Osanai et al. (2008).

1104

1105 Fig. 3. Field pictures of the Kham Duc complex. A: pillow basalts (GPS N15.60017;  
1106 E107.43735); B: partially melted metagabbro (GPS N15.44366; E107.84521); C: partly  
1107 melted mylonitic paragneiss, the white layers are Early Paleozoic quartz-feldspar aggregates  
1108 corresponding to leucosomes deformed by the Triassic D<sub>1</sub> deformation (GPS N15.51322;  
1109 E107.81195); D: isoclinally folded leucosomes formed during the D<sub>1</sub> or pre-D<sub>1</sub> event (GPS N  
1110 15.53013; E107.81676).

1111

1112 Fig. 4. Field pictures of the Ngoc Linh complex. A: Triassic metatexite derived from a  
1113 graywacke protoliths (W. of Quang Ngai; GPS N.14.99382; E108.78465); B: Garnet-biotite-  
1114 sillimanite-cordierite gneiss, the K-feldspar megacrysts suggest that it derived from an  
1115 orthogneiss (near Tu Mo Rong; GPS N14.99383; E107.89547); C: metatexite with mafic  
1116 xenolith (N. of Di Lang; GPS N14.92865; E108.64905); D: Early Paleozoic folded metatexite  
1117 block with a pre-D<sub>1</sub> foliation enclosed in the Triassic migmatite (S. of Di Lang; GPS  
1118 N15.00483; E108.50844).

1119

1120 Fig. 5. Structural elements of the Kon Tum massif. Stereoplots (Schmidt net, lower  
 1121 hemisphere) showing the statistical preferred orientation of the structural elements:  $S_1$ ,  $L_1$ ,  $S_2$ ,  
 1122  $L_2$ ,  $F_2$ . The stereogram with grey back compiles the total  $L_1$  measurements for the Ngoc Linh  
 1123 complex. The foliation trajectory outlines the domal shape of the Ngoc Linh complex, and the  
 1124 dominantly S-ward dip of the Kan Nack complex. Arrow indicates the top-to-the NW sense of  
 1125 shear related to the  $D_1$  event.

1126

1127 Fig. 6. Sketch of collapse fold, similar to the one shown in Fig. 7C, responsible for the  
 1128 reorientation of  $L_1$ . The NW-SE trend of  $L_1$  in the normal limb is changed to NE-SW in the  
 1129 inverted one. The top-to-the NW sense of shear becomes top-to-the-NE.

1130

1131 Fig. 7. Kinematic indicators related to the  $D_1$  and  $D_2$  events in the Ngoc Linh complex. A:  
 1132 Sigmoidal leucosome with top-to-the-NW sense of shear in a subhorizontal foliation (W. of  
 1133 Quang Ngai; GPS N14.92498; E108.67458); B: top-to-the-NW sigmoidal leucosome in a  
 1134 South dipping foliation (S. of Di Lang; GPS N14.82420; E108.54218); C: NW verging  
 1135 "collapse" fold (N. of Nam Tra My; GPS N15.23340; E108.09438); D:  $D_2$  upright fold  
 1136 developed in the Ngoc Linh migmatite (S. of Tra My; GPS N15.29946; E108.12762); E:  
 1137 sigmoidal leucocratic boudins in metatexite (W. of Quang Ngai; GPS N15.20117;  
 1138 E108.10903); F: pluri-meter sized retort-shaped leucosomic boudin indicating a top-to-the-  
 1139 NW sense of shear (N. of Nam Tra My; GPS N15.18497; E108.11713).

1140

1141 Fig. 8. Field-scale kinematic indicators in the Phuoc Son complex and Chu Lai migmatite. A:  
 1142 asymmetric amphibolite boudin indicating a top-to-the-NW sense of shear (Phuoc Son; GPS  
 1143 N15.51221; E107.81304); B: NW verging  $F_2$  folds refolding a  $D_1$  or pre- $D_1$  isoclinal fold  
 1144 (Phuoc Son; GPS N15.46803; E107.83970); C: Upright  $F_2$  fold developed in the Early



1145 Paleozoic Chu Lai migmatite (GPS N15.35611; E108.23472); D: Intrafolial pre-D<sub>1</sub> fold  
1146 coeval with the Early Paleozoic Chu Lai migmatite (GPS N15.35611; E108.23472).

1147

1148 Fig. 9. Thin section-scale kinematic indicators of the D<sub>1</sub> event showing a top-to-the NW sense  
1149 of shear in the Ngoc Linh and Phuoc Son complexes. A: D<sub>1</sub> sigmoidal biotite aggregate in  
1150 paragneiss (Phuoc Son; GPS N15.53013; E107.81676); B: pre-D<sub>1</sub> kyanite relict enclosed in a  
1151 sigmoidal plagioclase porphyroblast deformed during D<sub>1</sub> (GPS N15.49237; E107. 82388) C:  
1152 pre-D<sub>1</sub> kyanite relict enclosed in a sigmoidal plagioclase porphyroblast (Phuoc Son; GPS  
1153 N15.49237; E107.82388); D, E: sigmoidal muscovite in micaschist (GPS N14.92548;  
1154 E108.60463, N15.22334; E108.43497, respectively); F: quartz grain with an oblique shape  
1155 fabric (dashed line) due to synkinematic recrystallization (N15.20898; E108.41955).

1156

1157 Fig. 10. Geometric relationships between the Early Paleozoic Chu Lai migmatite, and the  
1158 Triassic Ngoc Linh migmatite. The S<sub>1</sub> foliation, and L<sub>1</sub> stretching lineation formed during the  
1159 D<sub>1</sub> event in the Ngoc Linh migmatite are folded by the D<sub>2</sub> event, coeval with the right lateral  
1160 strike-slip faulting along the Tra Bong fault. The Chu Lai migmatite, containing intrafolial  
1161 folds with high-angle dipping axes, is deformed by the post-folial D<sub>2</sub> upright folds (see Fig.  
1162 8C, D for a natural example).

1163

1164 Fig. 11. P-T-t paths for the Kan Nack (KN), Ngoc Linh (NL), and Kham Duc (KD)  
1165 metamorphic complexes (modified from Roger et al., 2007, and Nakano et al., 2013). A:  
1166 Early Paleozoic event. The rocks of the Kham Duc and Ngoc Linh complexes reached their  
1167 maximum pressure around 460 Ma. The retrograde paths were coeval with HT and crustal  
1168 melting around 450-420 Ma. B: Early to Middle Triassic event. The rocks of the Ngoc Linh  
1169 and Kan Nack complexes experienced a decompression melting event at UHT conditions.

1170

1171 Fig. 12. Crustal scale cross section of the S. Truong Son belt, and Kon Tum Massif. The Early  
1172 Paleozoic Tam Ky-Phuoc Son suture zone separates a northern block (N. Vietnam or Viet-  
1173 Lao block) from a southern block (S. Vietnam or Viet-Cambodia block). The weakly  
1174 deformed Paleozoic series of the Viet-Lao block argues for a north-directed subduction. The  
1175 Early to Middle Triassic Ngoc Linh metamorphic core complex (MCC) superimposes on the  
1176 Early Paleozoic belt. The northern border of the Ngoc Linh MCC is cut by E-W striking  
1177 dextral fault of the Tra Bong shear zone; the detachment fault is interpreted.

1178

1179 Fig. 13. Present tectonic map showing the different orogenic belts of S. China and Indochina  
1180 blocks: Neoproterozoic (Jiangnan), Early Paleozoic, and Triassic (simplified from Faure et  
1181 al., 2016a). JSJ: Jin Sha Jiang; AS: Ailaoshan, SM: Song Ma, SC: Song Chay; PKF: Po Ko  
1182 Fault, DBF: Bien Bien Fu Fault; RRF: Red River Fault, TS: Truong Son, XFS: Xuefengshan.

1183

1184 Fig. 14. Reconstruction of the Indochina-South China blocks tectonic framework in Early  
1185 Triassic. A: Palinspastic map after removing the Cenozoic sinistral strike-slip motion, and the  
1186 Triassic dextral strike-slip along the late Indosinian faults. In this reconstruction, the  
1187 Emeishan-Song Da LIP is about 1200 km from the Ngoc Linh MCC. B: schematic crustal  
1188 scale cross section from N. China to Kon Tum massif.

1189

1190 Fig. 15. Permian-Triassic geodynamic evolution of the SCB-Indochina lithosphere with  
1191 respect to the Emeishan plume fixed. In the Late Permian, a post-collisional slab break off is  
1192 inferred to allow heat transfer from the mantle plume to the Indochina lithosphere.

1193

1194 Fig. 16. Early Paleozoic to Late Carboniferous geodynamic evolution model from the North  
1195 China block (NCB) to South Vietnam. The Middle to Late Ordovician (around 460 Ma)  
1196 continental collision between the N. Vietnam (Viet-Lao) and S. Vietnam (Viet-Cambodia)  
1197 blocks is interpreted as the driving force for the intracontinental belt of S. China. The NCB-  
1198 SCB collision took place later, around 420 Ma. The Paleotethys ocean opened in Late  
1199 Carboniferous.





Fig. 1



*Sedimentary-volcanic formations*

- Quaternary (Q) and Neogene basalt (β)
- K2 Late Cretaceous red beds
- J Jurassic sandstone
- T3 Late Triassic sandstone
- unconformity*
- Permian-Early Triassic (T1)
- Middle Triassic (T2) terrigenous and felsic volcanics
- Devonian terrigenous rocks
- unconformity*
- Early Paleozoic rocks

*Metamorphic-migmatitic complexes in Kon Tum massif*

- Kham Duc Complex: HP/MT Early Paleozoic ophiolitic melange (σ: serpentinite, ω: gabbro) with a Triassic HT/LP metamorphic overprint
- Chu Lai migmatite and partly melted rocks
- Ngoc Linh complex: Early Triassic (ca 250 Ma) migmatite with Early Paleozoic restites
- Kan Nack complex: Early Triassic (ca 250 Ma) migmatite, granulite and charnockite with Early Paleozoic restites
- Foliated granitoid
- Isotropic granitoid
- Foliation and mineral lineation with dip

*Post-T1 Ngoc Linh granitoids*

- Two mica granite: Hai Van, Ba Na, Van Canh, and Cha Val, Phu Loc mafic plutons (ca 240-225 Ma)

*Pre-T1 Ngoc Linh granitoids*

- Ben Giang-Que Son diorite-gabbro-granodiorite (ca 260 Ma)
- Dai Loc porphyritic biotite-granite, two mica granite (ca 410 Ma)
- Dien Binh granodiorite (ca 450-444 Ma)
- Tra Bong granodiorite

Fig. 2





Fig. 3






Fig. 4




Fig. 5






Fig. 6




Fig. 7






Fig. 8






Fig. 9





Fig. 10

A: Early Paleozoic event

i) ca 460 Ma - HP/MT (Kham Duc) to UHP/MT (Ngoc Linh)

ii) ca 450-420 Ma - HT (Kham Duc, Ngoc Linh, Kannack) to UHT (Kannack)



B: Early - Middle Triassic event HT to UHT coeval with extensional tectonics




Fig. 11






Fig. 12




Fig. 13








Fig. 14






Fig. 15




Fig. 16