



HAL
open science

Formality of a higher-codimensional Swiss-Cheese operad

Najib Idrissi

► **To cite this version:**

Najib Idrissi. Formality of a higher-codimensional Swiss-Cheese operad. Algebraic and Geometric Topology, 2022, 22, 10.21136/HS.2021.09 . hal-01878406v3

HAL Id: hal-01878406

<https://hal.science/hal-01878406v3>

Submitted on 3 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Formality of a higher-codimensional Swiss-Cheese operad

Najib Idrissi*

November 3, 2020

We study bicolored configurations of points in the Euclidean n -space that are constrained to remain either inside or outside a fixed Euclidean m -subspace, with $n - m \geq 2$. We define a higher-codimensional variant of the Swiss-Cheese operad, called the complementarily constrained disks operad CD_{mn} , associated to such configurations. The operad CD_{mn} is weakly equivalent to the operad of locally constant factorization algebras on the stratified space $\{\mathbb{R}^m \subset \mathbb{R}^n\}$. We prove that this operad is formal over \mathbb{R} .

Contents

1. Prerequisites	4
2. Definition of CFM_{mn} and comparison	6
3. (Co)homology of CD_{mn}	9
4. Graph complexes	14
5. Proof of the formality	29
A. Relative cooperadic twisting	39
B. Compactifications and projections	41

Introduction

The little disks operads \mathcal{D}_n (for $n \geq 1$) represent operations acting on n -fold loop spaces. They have had many applications in homotopy theory over the years (see e.g. [17] for

*Université de Paris and Sorbonne Université, CNRS, IMJ-PRG, F-75006 Paris, France. Email: najib.idrissi-kaitoumi@imj-prg.fr

a survey). Elements of $D_n(k)$ consist of configurations of k disks inside the unit n -disk. Voronov’s Swiss-Cheese operads SC_n [50] (for $n \geq 2$) are relative versions of the little disks operads. They encode central actions of D_n -algebras on D_{n-1} -algebras. Elements of $SC_n(k, l)$ are given by configurations of k half-disks and l full disks in the unit upper half-disk.

In this article, we introduce higher-codimensional variants of the Swiss-Cheese operads CD_{mn} (for $n - 2 \geq m \geq 1$), called the “complementarily constrained (little) disks operads”.¹ These operads encode actions of D_n -algebras on a D_m -algebras by central derivations (see Section 3.2). Elements of $CD_{mn}(k, l)$ are given by configurations of k “terrestrial” disks centered on $D^m \subset D^n$ and l “aerial” entirely contained in $D^n \setminus D^m$.

A fundamental property of D_n is its formality [30, 47, 32, 38, 19]: the cohomology $H^*(D_n; \mathbb{Q})$ is quasi-isomorphic (as a cooperad in CDGAs) to the forms on D_n . Unlike the little disks operads, the Swiss-Cheese operads are not formal [33, 55]. In this paper, we establish the following result:

Theorem A (See Theorem 5.35). *For $n - 2 \geq m \geq 1$, the complementarily constrained disks operad CD_{mn} is formal over \mathbb{R} .*

The operad $CD_{(n-1)n}$ contains the Swiss-Cheese operad SC_n as a suboperad of connected components. It thus follows from arguments of Livernet [33] that $CD_{(n-1)n}$ is not formal (Remark 3.22). By [55, Section 5.1], nonformality of SC_n is equivalent to nonformality of the inclusion $D_{n-1} \hookrightarrow D_n$ (established in [48]). Since the inclusion $D_m \hookrightarrow D_n$ is formal for $n \geq m + 2$, Theorem A might not be a surprise. However, Willwacher studied another generalization of the Swiss-Cheese operad, the “extended Swiss-Cheese operad” ESC_{mn} [55]. He proved that its formality is equivalent to the formality of $D_m \subset D_n$ [55, Section 5.1]. The difference with CD_{mn} is that in CD_{mn} , the aerial disks are forbidden from touching the “ground” D^m , whereas this is allowed in ESC_{mn} (see Remark 2.11). The argument used for the formality of ESC_{mn} thus does not seem easily adaptable: CD_{mn} is obtained by removing a subspace from ESC_{mn} , an operation which is usually difficult to deal with in homotopy theory. It is not clear that formality of $D_m \subset D_n$ directly implies formality of CD_{mn} , or conversely.

Motivation The general motivation for this article comes from the study of configuration spaces started in previous works. Campos–Willwacher [8] and the author [27] provided combinatorial models for the real homotopy types of configuration spaces of simply connected closed smooth manifolds. Campos, Lambrechts, Willwacher, and the author [7] provided similar models for configuration spaces of compact, simply connected smooth manifolds of dimension ≥ 4 . Campos, Ducoulombier, Willwacher, and the author [6] studied framed configuration spaces of orientable closed manifolds, i.e. configurations of points equipped with trivializations of the tangent spaces.

In each of these articles, knowing models for the little disks operads or their variants was essential. Configuration spaces of \mathbb{R}^n are intimately linked to the little disks operad

¹In previous versions of this article, this operad was denoted VSC_{mn} and called the “variant Swiss-Cheese operad”.

D_n , and closed manifolds are locally homeomorphic to \mathbb{R}^n . The formality of D_n , and more precisely its proof by Kontsevich and Lambrechts–Volić, was thus essential in finding models for configuration spaces of closed manifolds in [8, 27]. Similarly, a manifold with boundary is locally homeomorphic to the upper half-space $\mathbb{H}^n \subset \mathbb{R}^n$, and configuration spaces of \mathbb{H}^n are linked to the Swiss-Cheese operad SC_n . While SC_n is not formal, Willwacher [53] defined a model for the real homotopy type of SC_n which was used extensively in [7]. For framed configuration spaces [6], we used the model for the framed little disks operad due to Khoroshkin–Willwacher [28].

Our goal is to study the configuration spaces of the complement $N \setminus M$, where N is a closed n -manifold and M is a closed sub- m -manifold of codimension ≥ 2 (e.g. the complement of a knot $S^3 \setminus K$). Such a pair (N, M) is locally homeomorphic to the stratified space $(\mathbb{R}^n, \mathbb{R}^m)$. Using the analogy above, configuration spaces of $(\mathbb{R}^n, \mathbb{R}^m)$ are linked to the operad CD_{mn} . Based on the previous works cited above, we are led to expect that models for CD_{mn} will produce models for $\text{Conf}_{N \setminus M}$ by adapting and generalizing the proof, just like models for D_n produced to models for configuration spaces of closed manifolds. In this article, we find that models for CD_{mn} are as simple as possible: the operad is formal, i.e. the cohomology $H^*(\text{CD}_{mn})$ is a model.

Proof strategy and outline The proof of our theorem is inspired by Kontsevich’s [30] proof of the formality of the little disks operad and its improvement by Lambrechts–Volić [32]. We first define the compactifications $\text{CFM}_{mn}(k, l)$ of $\text{Conf}_{\mathbb{R}^m(k)} \times \text{Conf}_{\mathbb{R}^n \setminus \mathbb{R}^m(l)}$, inspired by the Fulton–MacPherson compactification [3, 20, 45]. These compactifications form an operad with the homotopy type of CD_{mn} . We build an intermediate cooperad, cgraphs_{mn} , which serves as a bridge between the cohomology $H^*(\text{CFM}_{mn})$ and the piecewise semi-algebraic forms (see [24]) $\Omega_{\text{PA}}^*(\text{CFM}_{mn})$. The definition of cgraphs_{mn} is inspired by Willwacher’s model for the Swiss-Cheese operad [53] and by [28, Section 8]. The map $\text{cgraphs}_{mn} \rightarrow \Omega_{\text{PA}}^*(\text{CFM}_{mn})$ is defined by integrals. We cannot find a direct map $\text{cgraphs}_{mn} \rightarrow H^*(\text{CFM}_{mn})$, as the differential of cgraphs_{mn} depends on non-explicit integrals. Using vanishing results on the cohomology of some graph complex, we are able to simplify cgraphs_{mn} up to homotopy, and then map it to $H^*(\text{CFM}_{mn})$.

Section 1 contains background on operads, the little disks operads, and piecewise semi-algebraic forms. In Section 2, we define the compactifications CFM_{mn} and we compare them with CD_{mn} . We give examples of CD_{mn} -algebras based on relative iterated loop spaces. In Section 3, we compute the cohomology of CD_{mn} . We give a presentation of its homology $\text{cd}_{mn} = H_*(\text{CD}_{mn})$. In Section 4, we start by reviewing Kontsevich’s proof of the formality of the little disks operad, and we define the cooperad cgraphs_{mn} that will be used to adapt that proof to CFM_{mn} . We moreover construct the map from cgraphs_{mn} to $\Omega_{\text{PA}}^*(\text{CFM}_{mn})$ using integrals. In Section 5, we first show that certain integrals used in the definition of cgraphs_{mn} can be simplified. We construct the map into $H^*(\text{CFM}_{mn})$ and we show that all our maps are quasi-isomorphisms. In Appendix A, we define twisting of relative cooperads. In Appendix B, we sketch a proof that $\text{CFM}_{mn}(U, V)$ is an SA manifold and that canonical projections are SA bundles.

Acknowledgments The author would like to thank Benoit Fresse, Matteo Felder, Muriel Livernet, and Thomas Willwacher for fruitful discussions and helpful comments. The author would also like to thank the anonymous referees for detailed and thoughtful reports and many suggestions for improvements. The author acknowledges support from ERC StG 678156–GRAPHCPX.

1. Prerequisites

We work with cohomologically graded modules over the base field \mathbb{R} (except in Section 3.1 which is over \mathbb{Z}). The cohomology (resp. homology) of a space is concentrated in nonnegative (resp. nonpositive) degrees. For a graded vector space V , the free graded symmetric algebra $S(V)$ is $\mathbb{R}[V^{\text{even}}] \otimes \Lambda(V^{\text{odd}})$, i.e. the tensor product of the polynomial algebra on even elements by the exterior algebra on odd elements. Homogeneous elements x, y satisfy $yx = (-1)^{(\deg x)(\deg y)}xy$.

1.1. Operads

We work extensively with operads and cooperads and we assume basic proficiency with the theory. General references include [34] and [15, Part I(a)]. We will usually label inputs by elements of arbitrary finite sets rather than numbers. Briefly, let Bij be the category of finite sets and bijections. A symmetric collection is a functor $\text{Bij} \rightarrow \mathcal{C}$ where \mathcal{C} is a symmetric monoidal category. For $k \geq 0$, we let $\underline{k} = \{1, \dots, k\}$. A symmetric collection \mathbf{M} can equivalently be seen as a symmetric sequence $\{\mathbf{M}(n) := \mathbf{M}(\underline{n})\}_{n \geq 0}$ with actions of $\Sigma_n = \text{Aut}_{\text{Bij}}(\underline{n})$ on $\mathbf{M}(n)$.

For a pair of finite sets $W \subset U$, we define the quotient $U/W := (U \setminus W) \sqcup \{*\}$ (note that $U/\emptyset = U \sqcup \{*\}$). For $u \in U$, we let $[u] \in U/W$ be its class in the quotient. An operad \mathbf{P} is a symmetric collection equipped with composition maps $\circ_W : \mathbf{P}(U/W) \otimes \mathbf{P}(W) \rightarrow \mathbf{P}(U)$, for each pair $W \subset U$, satisfying the usual axioms. A cooperad \mathbf{C} is a symmetric collection equipped with cocomposition maps $\circ_W^\vee : \mathbf{C}(U) \rightarrow \mathbf{C}(U/W) \otimes \mathbf{C}(W)$. A Hopf cooperad [15] is a cooperad in the category of commutative differential-graded algebras (CDGAs).

We also deal with some special particular bicolored operads called “relative operads” [50] or “Swiss-Cheese type operads” [54]. Given an operad \mathbf{P} , a relative \mathbf{P} -operad is an operad in the category of right \mathbf{P} -modules. Equivalently, a relative \mathbf{P} -operad is a bisymmetric collection, i.e. a functor $\mathbf{Q} : \text{Bij} \times \text{Bij} \rightarrow \mathcal{C}$, equipped with composition maps:

$$\begin{aligned} \circ_T : \mathbf{Q}(U, V/T) \otimes \mathbf{P}(T) &\rightarrow \mathbf{Q}(U, V) && \text{for } V \subset T, \\ \circ_{W,T} : \mathbf{Q}(U/W, V) \otimes \mathbf{Q}(W, T) &\rightarrow \mathbf{Q}(U, V \sqcup T) && \text{for } W \subset U. \end{aligned}$$

We will often write “the operad \mathbf{Q} ” when \mathbf{P} is implicit from the context. Relative cooperad are defined dually. We also apply the adjective “Hopf” to refer to such cooperads in the category of CDGAs.

1.2. Little disks and variants

Fix some $n \geq 0$ and let $D^n = \{x \in \mathbb{R}^n \mid \|x\| \leq 1\}$ be the closed disk. The space $D_n(r)$ is the space of maps $c : (D^n)^{\sqcup r} \rightarrow D^n$ such that: (i) each $c_i : D^n \rightarrow D^n$ is an embedding given by the composition of a translation and a positive rescaling; (ii) the interiors of two different little disks are disjoint, i.e. $c_i(\overset{\circ}{D}^n) \cap c_j(\overset{\circ}{D}^n) = \emptyset$ for $i \neq j$. Using composition of embeddings, the collection $D_n = \{D_n(r)\}_{r \geq 0}$ forms a topological operad called the little n -disks operad.

Let us now fix notations for the (Axelrod–Singer–)Fulton–MacPherson operad [20, 3] (see also [45] and [32, Chapter 5]). Given some space X , and a finite set U , we define the configuration space:

$$\text{Conf}_X(U) := \{(x_u)_{u \in U} \in X^U \mid \forall u \neq v, x_u \neq x_v\}. \quad (1.1)$$

Consider the quotient $\underline{\text{Conf}}_n(U) := \text{Conf}_{\mathbb{R}^n}(U)/\mathbb{R}^n \rtimes \mathbb{R}_{>0}$ by the action of translations and positive rescalings. This space embeds in $(S^{n-1})^{\text{Conf}_U(2)} \times [0, +\infty]^{\text{Conf}_U(3)}$ using the maps θ_{ij} and δ_{ijk} from (2.2), (2.3). The Fulton–MacPherson compactification $\text{FM}_n(r)$ is the closure of the image of this embedding. The collection FM_n forms an operad [32, Section 5.2] with the same homotopy type as D_n , see [36] and [42, Proposition 4.3].

Remark 1.2. There is an operad $\text{Disk}_n^{\text{fr}}$ related to locally constant framed factorization algebras on $\overset{\circ}{D}^n$ and which has the same homotopy type as D_n , see [35, Definition 5.4.2.10] or [4, Notation 2.8]. This operad $\text{Disk}_n^{\text{fr}}$ is associated, in some sense, to the trivial stratification of \mathbb{R}^n . The present article is devoted to the operad $\text{Disk}_{m \subset n}^{\text{fr}}$ associated to the stratification $\{\mathbb{R}^m \subset \mathbb{R}^n\}$ [4, Section 4.3].

1.3. Semi-algebraic sets and PA forms

For technical reasons, we will use the technology of semi-algebraic (SA) sets and piecewise semi-algebraic (PA) forms. We use [24] as a general reference. Recall in particular that the CDGA $\Omega_{\text{PA}}^*(X)$ of all PA forms on a compact SA set X is a model for the real homotopy type of X [24, Theorem 6.1].

If \mathbb{P} is an operad in compact SA sets, then the symmetric sequence $\Omega_{\text{PA}}^*(\mathbb{P})$ is not a Hopf cooperad: the Künneth quasi-isomorphisms go in the wrong direction. However, for a Hopf cooperad \mathbb{C} , we can define a “morphism” $\mathbb{C} \rightarrow \Omega_{\text{PA}}^*(\mathbb{P})$ as a collection of maps $\mathbb{C}(U) \rightarrow \Omega_{\text{PA}}^*(\mathbb{P}(U))$ making the obvious diagrams commute [32, Chapter 3]. We call it a quasi-isomorphism if it is a quasi-isomorphism in each arity. For simplicity, we will treat $\Omega_{\text{PA}}^*(\mathbb{P})$ as if it were an actual Hopf cooperad; when we write a morphism into $\Omega_{\text{PA}}^*(\mathbb{P})$, we actually have a “morphism” as defined above. Results of Fresse [16, Discussion after Proposition 4.4] can be adapted to Ω_{PA}^* : if \mathbb{P} is cofibrant in the category of operads satisfying $\mathbb{P}(0) = \{*\}$ (e.g. $\mathbb{P} = \text{FM}_n$), then any Hopf cooperad quasi-isomorphic to $\Omega_{\text{PA}}^*(\mathbb{P})$ encodes the real homotopy type of \mathbb{P} . Briefly, there is an operadic upgrade of the functor Ω_{PA}^* which turns operads into Hopf cooperads, and this operadic upgrade is part of a Quillen adjunction. We will thus say that an operad \mathbb{P} in compact SA sets is formal if there exists a zigzag of quasi-isomorphisms of Hopf cooperads $H^*(\mathbb{P}; \mathbb{R}) \leftarrow \cdot \rightarrow \Omega_{\text{PA}}^*(\mathbb{P})$.

These constructions can be extended to relative operads over a given base, as they can be seen as operads in a given symmetric monoidal category (of right operadic modules).

2. Definition of CFM_{mn} and comparison

From now on and until the end of the article, we fix integers $n - 2 \geq m \geq 1$. (In some tangential remarks, we will consider $n = m + 1$.) We identify \mathbb{R}^m as the subspace of \mathbb{R}^n given by $\mathbb{R}^m \times \{0\}^{n-m}$.

2.1. The compactification and its boundary

Let U and V be finite sets. We define the colored configuration spaces by:

$$\text{Conf}_{mn}(U, V) := \text{Conf}_{\mathbb{R}^m}(U) \times \text{Conf}_{\mathbb{R}^n \setminus \mathbb{R}^m}(V) \subset \text{Conf}_{\mathbb{R}^n}(U \sqcup V). \quad (2.1)$$

Roughly speaking, $\text{Conf}_{mn}(U, V)$ is the set of configurations of bicolored points in \mathbb{R}^n : U “terrestrial” points in \mathbb{R}^m , and V “aerial” points in $\mathbb{R}^n \setminus \mathbb{R}^m$. We will reuse the terminology “aerial/terrestrial” throughout the document.

The group $\mathbb{R}^m \rtimes \mathbb{R}_{>0}$ of translations and positive rescalings acts on $\text{Conf}_{mn}(U, V)$. Let $\underline{\text{Conf}}_{mn}(U, V)$ be the quotient. Elements of $\underline{\text{Conf}}_{mn}(U, V)$ can be seen as configurations of radius 1 (i.e. $\max(\|x_i\|)_{i \in U \sqcup V} = 1$) with a barycenter in $\{0\}^m \times \mathbb{R}^{n-m} \subset \mathbb{R}^n$. Since $\mathbb{R}^m \rtimes \mathbb{R}_{>0}$ is contractible, the quotient map is a weak homotopy equivalence. If $\#U + 2\#V \geq 2$, then the action is free, smooth, and proper, thus $\underline{\text{Conf}}_{mn}(U, V)$ is a manifold of dimension $m\#U + n\#V - m - 1$. However, if $\#U \leq 1$ and $\#V = 0$, then $\underline{\text{Conf}}_{mn}(U, V)$ is merely a point. We embed $\underline{\text{Conf}}_{mn}(U, V)$ into $(S^{n-1})^{\text{Conf}_{U \sqcup V}(2)} \times [0, +\infty]^{\text{Conf}_{U \sqcup V}(3)} \times (S^{n-m-1})^V \times [0, +\infty]^{\text{Conf}_V(2)} \times [0, +\infty]^{\text{Conf}_{U \sqcup V}(2) \times V}$ using the maps:

$$\theta_{ij}(x) := (x_i - x_j) / \|x_i - x_j\|, \quad \text{for } i \neq j \in U \sqcup V; \quad (2.2)$$

$$\delta_{ijk}(x) := \|x_i - x_j\| / \|x_i - x_k\|, \quad \text{for } i \neq j \neq k \neq i \in U \sqcup V; \quad (2.3)$$

$$\alpha_v(x) := p_{(\mathbb{R}^m)^\perp}(x_v) / \|p_{(\mathbb{R}^m)^\perp}(x_v)\|, \quad \text{for } v \in V; \quad (2.4)$$

$$\rho_{vv'}(x) := \|p_{(\mathbb{R}^m)^\perp}(x_v)\| / \|p_{(\mathbb{R}^m)^\perp}(x_{v'})\|, \quad \text{for } v, v' \in V; \quad (2.5)$$

$$\sigma_{ijv}(x) := \|x_i - x_j\| / \|p_{(\mathbb{R}^m)^\perp}(x_v)\|, \quad \text{for } i, j \in U \sqcup V, v \in V; \quad (2.6)$$

where $p_{(\mathbb{R}^m)^\perp}$ is the orthogonal projection on $(\mathbb{R}^m)^\perp = \{0\}^m \times \mathbb{R}^{n-m} \subset \mathbb{R}^n$.

Definition 2.7. The space $\text{CFM}_{mn}(U, V)$ is the closure of the image of the embedding $(\theta_{ij}, \delta_{ijk}, \alpha_v, \rho_{vv'}, \sigma_{ijv})$.

Example 2.8. We have $\text{CFM}_{mn}(U, \emptyset) = \text{FM}_m(U)$ and $\text{CFM}_{mn}(0, 1) = S^{n-m-1}$.

Proposition 2.9 (Appendix B). *The space $\text{CFM}_{mn}(U, V)$ is a compact semi-algebraic manifold and a smooth manifold with corners. Its dimension is $m\#U + n\#V - m - 1$ if $\#U + 2\#V \geq 2$, and zero otherwise. The projections $p_{U, V} : \text{CFM}_{mn}(U \sqcup I, V \sqcup J) \rightarrow \text{CFM}_{mn}(U, V)$ are SA bundles.*

Proposition 2.10. *The collection CFM_{mn} forms a relative FM_n -operad.*

Proof. On the coordinates θ_{ij} and δ_{ijk} , the formulas are identical to FM_n , see [32, Section 5.2]. For $v \in V$, one simply define $\alpha_v(x \circ_T y) = \alpha_{[v]}(x)$; and $\alpha_v(x' \circ_{W,T} y') = \alpha_v(x')$ if $v \notin T$, or $\alpha_v(y')$ if $v \in T$. For $v \neq v' \in V$, then $\rho_{vv'}(x \circ_T y) = 1$ if $v, v' \in T$, or $\rho_{[v][v']}(x)$ otherwise; and $\rho_{vv'}(x' \circ_{W,T} y') = \rho_{vv'}(x')$ if $v, v' \notin T$, or $\rho_{vv'}(y')$ if $v, v' \in T$, or 0 if $v \in T, v' \notin T$, or $+\infty$ if $v' \in T, v \notin T$. Finally, $\sigma_{ijv}(x \circ_T y) = 0$ if $i, j \in T$, or $\sigma_{[i][j][v]}(x)$ otherwise; and $\sigma_{ijv}(x' \circ_{W,T} y') = \sigma_{ijv}(y')$ if $i, j, v \in W \cup T$, or 0 if $i, j \in W \cup T, v \notin T$, or $\sigma_{[i][j]v}(x')$ if $v \notin T$ and $\#\{i, j\} \cap (W \cup T) = 1$, or ∞ otherwise. \square

Remark 2.11. The operad CFM_{mn} is not homotopy equivalent to the operad ESC_{mn} considered by Willwacher [55]. Recall that $\text{ESC}_{mn}(U, V) := \text{D}_n(U \sqcup V) \times_{\text{D}_n(U)} \text{D}_m(U)$, where $\text{D}_n(U \sqcup V) \rightarrow \text{D}_n(U)$ is the projection that forgets disks and $\text{D}_m(U) \rightarrow \text{D}_n(U)$ is the usual embedding. The difference is that we do not allow ‘‘aerial’’ points to be on \mathbb{R}^m , so e.g. $\text{CFM}_{mn}(0, 1) = S^{n-m-1} \not\cong \text{ESC}_{mn}(0, 1) \simeq *$, and $\text{CFM}_{mn}(1, 1) \simeq S^{n-m-1} \not\cong \text{ESC}_{mn}(1, 1) \simeq S^{n-1}$.

Proposition 2.12. *We have a decomposition in terms of faces:*

$$\partial \text{CFM}_{mn}(U, V) = \bigcup_{T \in \mathcal{BF}'(V)} (\text{im} \circ_T) \cup \bigcup_{(W, T) \in \mathcal{BF}''(U; V)} (\text{im} \circ_{W, T}),$$

where the subsets $\mathcal{BF}'(V) \subset V$ and $\mathcal{BF}''(U; V) \subset U \times V$ are respectively defined by the conditions $\#T \geq 2$ and by $(W \cup T \subsetneq U \cup V$ and $2 \cdot \#T + \#W \geq 2)$. Each of these boundary faces is a compact SA subset of the boundary, and the intersection of two distinct faces is of positive codimension in $\partial \text{CFM}_{mn}(U, V)$.

Proof. Adapting the proofs of [32, Proposition 5.4.1] is straightforward. \square

If $p : E \rightarrow B$ is an SA bundle of rank k , then its fiberwise boundary $p^\partial : E^\partial \rightarrow B$ is an SA bundle of rank $k - 1$, where $E^\partial = \bigcup_{x \in B} \partial p^{-1}(x)$, see [24, Definition 8.1]. The fiberwise Stokes formula reads $d(p_* \alpha) = p_*(d\alpha) \pm p_*^\partial \alpha$ [24, Proposition 8.12], where $p_* : \Omega_{\min}^*(E) \rightarrow \Omega_{\text{PA}}^{*-k}(B)$ is integration along fibers.

Proposition 2.13. *The fiberwise boundary of the projection $p_{U, V} : \text{CFM}_{mn}(U \sqcup I, V \sqcup J) \rightarrow \text{CFM}_{mn}(U, V)$ is the subset of $\text{CFM}_{mn}(U \sqcup I, V \sqcup J)$ given by:*

$$\text{CFM}_{mn}^\partial(U \sqcup I, V \sqcup J) = \bigcup_{T \in \mathcal{BF}'(V, J)} (\text{im} \circ_T) \cup \bigcup_{(W, T) \in \mathcal{BF}''(U, I; V, J)} (\text{im} \circ_{W, T}),$$

where the subsets $\mathcal{BF}'(V, J) \subset \mathcal{BF}'(V \sqcup J)$ and $\mathcal{BF}''(U, I; V, J) \subset \mathcal{BF}''(U \sqcup I, V \sqcup J)$ are respectively defined by the conditions $\#(T \cap J) \leq 1$ and by $((U \subset W$ and $V \subset T)$ or $(V \cap T = \emptyset$ and $\#(U \cap W) \leq 1)$.

Proof. We can adapt the proof of [32, Proposition 5.7.1] immediately. We simply use the decomposition of Proposition 2.12 and we check easily that $\mathcal{BF}'(V, J)$ and $\mathcal{BF}''(U, I; V, J)$ index the faces which are sent to the interior of $\text{Conf}_{mn}(U, V)$ under the projection $p_{U, V}$. \square

2.2. Comparison with CD_{mn}

In this section, we compare CFM_{mn} with CD_{mn} , the complementarily constrained (little) disks operad. Let $D^m = D^n \cap \mathbb{R}^m$ for convenience.

Definition 2.14. The space $\text{CD}_{mn}(U, V)$ is the space of maps $c : (D^n)^{U \sqcup V} \hookrightarrow D^n$ satisfying: (i) for all i , $c_i : D^n \hookrightarrow D^n$ is an embedding obtained by composing a translation and a positive rescaling; (ii) for $i \neq j$, we have $c_i(\overset{\circ}{D}^n) \cap c_j(\overset{\circ}{D}^n) = \emptyset$; (iii) for $u \in U$, we have $c_u(D^m) \subset D^m$; (iv) for $v \in V$, we have $c_v(D^n) \cap D^m = \emptyset$. Using composition of embeddings, CD_{mn} is a relative D_n -operad, called the complementarily constrained disks operad.


Remark 2.15. The usual Swiss-Cheese operad SC_n is the suboperad of $\text{CD}_{(n-1)n}$ formed by the components where all the aerial disks are in the upper half-disk.

Proposition 2.16. *There exists a zigzag of weak homotopy equivalences of operads $(\text{CD}_{mn}, \text{D}_n) \simeq (\text{CFM}_{mn}, \text{FM}_n)$.*

Proof. We can adapt the proof of Salvatore [42, Proposition 3.9] directly. Briefly, we use the Boardman–Vogt resolution $W\text{CD}_{mn} \xrightarrow{\sim} \text{CD}_{mn}$. Elements of $W\text{CD}_{mn}(U, V)$ are rooted trees with (U, V) leaves, bicolored edges, internal vertices labeled by CD_{mn} , and internal edges labeled by time parameters $t \in [0, 1]$. Operadic composition is grafting (new edges are decorated by 1). If $t = 0$ then the corresponding edge is collapsed and the decorations are composed. The map $W\text{CD}_{mn} \rightarrow \text{CFM}_{mn}$ is defined on trees with edge decorations < 1 by rescaling the disks by one minus edge decoration, composing in CD_{mn} , and keeping the centers of the remaining disks. This extends continuously to composite trees and defines a weak equivalence of operads. \square

Let us also give examples of CD_{mn} -algebras. Recall first that for a pointed space $* \in X$, the iterated loop space $\Omega^n X$ is the space of maps $\gamma : D^n \rightarrow X$ such that $\gamma(\partial D^n) = *$. The space $\Omega^n X$ is an algebra over D_n . Conversely, the recognition principle states that any “group-like” D_n -algebra is weakly equivalent to an iterated loop space [5, 37].

For a pair of pointed topological spaces $* \in A \subset X$, the iterated loop space $\Omega^n(X, A)$ is $\text{hofib}(\Omega^{n-1} A \rightarrow \Omega^{n-1} X)$. Concretely, let $D_h^n := D^n \cap \mathbb{H}^n$ be the upper half-disk. Its boundary ∂D_h^n is the union of the disk $\partial_- D_h^n := D^n \cap \partial \mathbb{H}^n \cong D^{n-1}$ and the upper hemisphere $\partial_+ D_h^n := \partial D^n \cap \mathbb{H}^n \cong D^{n-1}$ along the equator $\partial D^n \cap \partial \mathbb{H}^n \cong S^{n-2}$. The relative iterated loop space $\Omega^n(X, A)$ is the space of maps $\gamma : D_h^n \rightarrow X$ such that $\gamma(\partial_- D_h^n) \subset A$ and $\gamma(\partial_+ D_h^n) = *$. For example, $\Omega^1(X, A) = \{\gamma : [0, 1] \rightarrow X \mid \gamma(0) \in A, \gamma(1) = *\}$. A sketch for $n = 2$ is on the side. The pair $(\Omega^n(X, A), \Omega^n X)$ is an algebra over the operad SC_n . By the relative recognition principle, any SC_n -algebra satisfying appropriate properties is weakly equivalent to such a pair [13, 41, 26, 49].



By analogy, we define the (n, m) -relative iterated loop space:

$$\Omega^{n,m}(X, A) := \{\gamma : D^n \rightarrow X \mid \gamma(D^m) \subset A \text{ and } \gamma(\partial D^n) = *\}. \quad (2.17)$$

The pair $(\Omega^{n,m}(X, A), \Omega^n X)$ is an algebra over the operad \mathbf{CD}_{mn} . We conjecture that an analogous relative recognition principle holds: any \mathbf{CD}_{mn} -algebra satisfying appropriate conditions should be weakly equivalent to such a pair.

3. (Co)homology of \mathbf{CD}_{mn}

In this section, we compute the integral cohomology of \mathbf{CD}_{mn} (Definition 2.14). We then give a presentation of the operad $H_*(\mathbf{CD}_{mn})$ by generators and relations. Unless specified, the ring of coefficients is \mathbb{Z} in this section.

3.1. The cohomology as a Hopf cooperad

We will first compute the cohomology of $\mathbf{Conf}_W(l)$ with $W := \mathbb{R}^n \setminus \mathbb{R}^m$ for $n - m \geq 2$. The computation is inspired by the methods of [46]. We prove that it is free as an abelian group, thus we will be able to apply Künneth's formula to get the cohomology of $\mathbf{CD}_{mn}(k, l) \simeq \mathbf{Conf}_{\mathbb{R}^m}(k) \times \mathbf{Conf}_W(l)$ as a tensor product. Then we study the maps induced on cohomology by the operad structure of \mathbf{CD}_{mn} .

Definition 3.1. The Poincaré polynomial of X is $\mathcal{P}(X) := \sum_{i \geq 0} (\text{rk } H^i(X)) \cdot t^i$. For $P, Q \in \mathbb{N}[[t]]$, we say that $P \preceq Q$ if the coefficients of $Q - P$ are nonnegative.

Proposition 3.2. *For $n - m \geq 2$, the Poincaré polynomial of $\mathbf{Conf}_W(l)$ satisfies:*

$$\mathcal{P}(\mathbf{Conf}_W(l)) \preceq \prod_{i=0}^{l-1} (1 + t^{n-m-1} + it^{n-1}). \quad (3.3)$$

Moreover, if the equality is reached and the homology of $\mathbf{Conf}_W(l-1)$ is free as a \mathbb{Z} -module, then the homology of the total space $\mathbf{Conf}_W(l)$ is free too.

Proof. We use the Serre spectral sequence of the Fadell–Neuwirth fibrations:

$$S^{n-m-1} \vee (S^{n-1})^{\vee(l-1)} \hookrightarrow \mathbf{Conf}_W(l) \xrightarrow{\pi} \mathbf{Conf}_W(l-1). \quad (3.4)$$

For $n - m = 2$, the base $\mathbf{Conf}_W(l-1)$ is not simply connected. However, we can adapt the arguments of [9, Lemma 6.3] to show that the coefficient system is trivial. Let $c_1, c_2 \in \mathbf{Conf}_W(l-1)$ be two configurations and $F_1, F_2 = \pi^{-1}(c_1), \pi^{-1}(c_2) \subset \mathbf{Conf}_W(l)$ the fibers ($\simeq S^1 \vee (S^{n-1})^{\vee(l-1)}$). Any path $\gamma \in \Omega_{c_1, c_2} \mathbf{Conf}_W(l-1)$ lifts to a path in $\mathbf{Conf}_W(l)$ by putting the l th point far from all the others (e.g. outside a ball B enclosing the compact subset $\text{im}(\gamma)$). Let us show that the induced isomorphism $h_\gamma : H_*(F_1) \rightarrow H_*(F_2)$ is the identity.

It is clear that h_γ does not affect the fundamental class of $S^{n-m-1} = S^1$, as we can choose a representative with the l th point rotating around the axis R^m outside the ball B . The class of the i th S^{n-1} in the fiber corresponds to the l th point rotating around the i th point. This can be represented by concatenating a path η_{ij} from l to i with a small sphere σ_i around i . Consider the path γ_i given by the i th coordinate of the path γ . Then $(h_\gamma)_*[S^{n-1}]$ can be represented by $\eta_{ij} \cdot \gamma_i \cdot \sigma_i$ (see Figure 3.1 for an example). But this $(n-1)$ st homology class is homologous to $\eta_{ij} \cdot \sigma_i$ for any path γ . \square




Figure 3.1: The effect of h_γ on the fundamental class (in blue) of S^{n-1}

Remark 3.5. If $n = m + 1$, then $W = \mathbb{R}^n \setminus \mathbb{R}^{n-1} \cong \mathbb{R}^n \sqcup \mathbb{R}^n$ is not even connected. However, we then have $\text{Conf}_W(l) = \bigsqcup_{l=l'+l''} \text{Conf}_{\mathbb{R}^n}(l') \times \text{Conf}_{\mathbb{R}^n}(l'') \times_{\Sigma_{l'} \times \Sigma_{l''}} \Sigma_l$. Its Poincaré polynomial is $\sum_{l=l'+l''} \frac{l!}{(l')!(l'')!} \prod_{i=0}^{l'-1} (1 + it^{n-1}) \prod_{j=0}^{l''-1} (1 + jt^{n-1})$, which is the RHS of (3.3) (here, $\prod_{i=0}^{l-1} (2 + t^{n-1})$) by induction.

The classical presentation of $e_n^\vee(l) := H^*(\text{Conf}_{\mathbb{R}^n}(l))$ [1, 9] is ($\deg \omega_{ij} = n - 1$):

$$e_n^\vee(l) = S(\omega_{ij})_{1 \leq i \neq j \leq l} / (\omega_{ji} - (-1)^n \omega_{ij}, \omega_{ij}^2, \omega_{ij}\omega_{jk} + \omega_{jk}\omega_{ki} + \omega_{ki}\omega_{ij}). \quad (3.6)$$

Definition 3.7. We define an algebra, with generators η_i of degree $n - m - 1$:

$$\text{cd}_{mn}^\vee(0, l) := e_n^\vee(l) \otimes S(\eta_i)_{1 \leq i \leq l} / (\eta_i^2, \eta_i \omega_{ij} - \eta_j \omega_{ij}). \quad (3.8)$$

Elements of $e_n^\vee(l)$ can be seen as linear combinations of simple oriented graphs on l vertices, modulo orientation signs, double edges, and a local three-term Arnold relation. In $\text{cd}_{mn}^\vee(0, l)$, each connected component is decorated by 1 or η (formally, vertices can be decorated and decorations can move along edges; two η classes on the same vertex give 0).

Remark 3.9. This algebra is very similar to the Lambrechts–Stanley model \mathbf{G}_A [27] applied to $A = H^*(D^n \setminus D^m)$ with vanishing diagonal class.

Remark 3.10. The space $\text{Conf}_W(l) = (\mathbb{R}^n)^l \setminus (\bigcup_{i \neq j} \{x_i = x_j\} \cup \bigcup_{i=1}^l p_i^{-1}(\mathbb{R}^m))$ is a real subspace arrangement. Its Betti numbers can be computed by [23, Theorem A]. For even n and m , this is also a complex arrangement and the ring structure of the cohomology can also be computed using [10, Section 5].

Lemma 3.11. *The Poincaré polynomial of $\text{cd}_{mn}^\vee(0, l)$ is the RHS of (3.3).*

Proof. Let V_d be the graded vector space with basis $\langle 1, \omega_{1d}, \dots, \omega_{(d-1)d}, \eta_d \rangle$. Its Poincaré polynomial is $1 + t^{n-m-1} + (d-1)t^{n-1}$. It thus suffices to show that $\text{cd}_{mn}^\vee(0, l) \cong V_1 \otimes \dots \otimes V_l$, i.e. that $\text{cd}_{mn}^\vee(0, l)$ has the basis $\omega_{i_1 j_1} \dots \omega_{i_r j_r} \eta_{k_1} \dots \eta_{k_s}$, where $j_1 < \dots < j_r$, $i_a < j_a$ and $k_b \neq j_a$ for all a, b (that is, graphs where edges increase and are ordered by their target, no two edges have the same target, and no target is decorated by η).

We use the theory of commutative PBW bases [40, Section 4.8]. Generators of $\text{cd}_{mn}^\vee(0, l)$ are ω_{ij} for $1 \leq i < j \leq l$ and η_i for $1 \leq i \leq l$. We order them by $\eta_1 > \dots > \eta_l$, $\omega_{ij} < \omega_{kl}$ if $i < k$ or $i = k \wedge j < l$, and e.g. $\eta_k < \omega_{ij}$ for all i, j, k . Monomials are ordered

lexicographically. Quadratic relations, written as “rewriting rules” (higher monomial = lower monomials), are $\omega_{jk}\omega_{ik} = (-1)^{n+1}\omega_{ij}\omega_{jk} - \omega_{ik}\omega_{ij}$ for all $i < j < k$, $\eta_j\omega_{ij} = \eta_i\omega_{ij}$ and $\omega_{ij}^2 = 0$ for all $i < j$, and $\eta_i^2 = 0$ for all i . The claimed basis is exactly the PBW generating set.

We need to check that overlapping rewriting rules are confluent, i.e. we get to the same element using only the rewriting rules. The critical monomials are $\omega_{lk}\omega_{jk}\omega_{ik}$, ω_{ij}^3 , η_i^3 , $\omega_{jk}^2\omega_{ik}$, $\eta_k\omega_{jk}\omega_{ik}$, $\eta_j\omega_{ij}^2$, and $\eta_j^2\omega_{ij}$ (where $i < j < k < l$). These are all easily seen to be confluent. For example, $\eta_k\omega_{jk}\omega_{ik}$ can be rewritten in two ways. The first way, $(\eta_k\omega_{jk})\omega_{ik} = \eta_j\omega_{jk}\omega_{ik} = \eta_j((-1)^{n+1}\omega_{ij}\omega_{jk} - \omega_{ik}\omega_{ij}) = (-1)^{n+1}\eta_j\omega_{ij}\omega_{jk} - \eta_i\omega_{ik}\omega_{ij}$. The second way, $\eta_k(\omega_{jk}\omega_{ik}) = \eta_k((-1)^{n+1}\omega_{ij}\omega_{jk} - \omega_{ik}\omega_{ij}) = (-1)^{n+1}\eta_j\omega_{ij}\omega_{jk} - \eta_i\omega_{ik}\omega_{ij} = (-1)^{n+1}\eta_i\omega_{ij}\omega_{jk} - \eta_i\omega_{ik}\omega_{ij}$. \square

Proposition 3.12. *For $n-2 \geq m \geq 1$, we have a well-defined algebra map $\text{cd}_{mn}^\vee(0, l) \rightarrow H^*(\text{Conf}_W(l))$ given by $\omega_{ij} \mapsto \theta_{ij}^*(\text{vol}_{n-1})$ and $\eta_i \mapsto \alpha_i^*(\text{vol}_{n-m-1})$.*

Proof. Clearly $\alpha_i^*(\text{vol}_{n-m-1})^2 = 0$, so we just need to check is that $\eta_i\omega_{ij} = \eta_j\omega_{ij}$. It is sufficient to check this on $\text{Conf}_W(2)$. The product $S^{n-1} \times S^{n-m-1}$ maps into $\text{Conf}_W(2)$: the S^{n-1} describes the rotation of point 2 around point 1, while the S^{n-m-1} describes the rotation of the pair around \mathbb{R}^m . By (3.3), $\dim H^{n-m-1+n-1}(\text{Conf}_2(W)) \leq 1$. Since $\eta_1\omega_{12}$ and $\eta_2\omega_{12}$ pair to 1 with the pushforward of the fundamental class of $S^{n-1} \times S^{n-m-1}$, we get $\eta_1\omega_{12} = \eta_2\omega_{12}$. \square

Proposition 3.13. *For $n-2 \geq m \geq 1$, the map $\text{cd}_{mn}^\vee(0, l) \rightarrow H^*(\text{Conf}_W(l))$ is an isomorphism.*

Proof. Our proof is inspired by the proof of [46, Theorem 4.9] (see e.g. [44] for other uses of the pairing defined below). Given Inequality (3.3), the universal coefficients theorem, and Proposition 3.12, it is sufficient to show that the map is injective over any field $\mathbb{k} \in \{\mathbb{Q}, \mathbb{F}_p\}$.

Let us define classes in $H_*(\text{Conf}_W(l); \mathbb{k})$ using “solar system” [46, Section 2]. With this point of view, any forest consisting of binary trees whose roots are possibly decorated by a loop induces a class in $H_*(\text{Conf}_W(l); \mathbb{k})$. The difference with classical solar systems is that orbital centers are chosen far enough from \mathbb{R}^m , and that if a root is decorated by a loop then the orbital center itself orbits around \mathbb{R}^m in the shape of S^{n-m-1} . For example, the embedding $S^{n-1} \times S^{n-m-1} \rightarrow \text{Conf}_W(2)$ of Proposition 3.12 corresponds to the decorated binary tree with two leaves.

The homology classes induced by undecorated trees satisfy the Jacobi relation, as these homology classes come from the subspace $\text{Conf}_{\mathbb{R}^n}(l)$ (with \mathbb{R}^n being e.g. the upper half-space). The homology/cohomology pairing is given by the pairing between graphs (using the description after Definition 3.7) and trees, which is nondegenerate by [46, Theorem 4.7], tensored with the pairing between the loops and the classes η (which is clearly nondegenerate). We therefore get that $\text{cd}_{mn}^\vee(l) \rightarrow H^*(\text{Conf}_W(l))$ is injective, establishing the result. \square

Definition 3.14. We define, for integers $k, l \geq 0$, $\text{cd}_{mn}^\vee(k, l) := e_m^\vee(k) \otimes \text{cd}_{mn}^\vee(0, l)$.

For cosmetic reasons, for $m \geq 2$ we will write $\tilde{\omega}_{ij}$ for the generators of $\mathbf{e}_m^\vee(k)$, to distinguish them from the generators of $\mathbf{cd}_{mn}^\vee(0, l)$. (Recall $\mathbf{e}_1^\vee(k)$ is simply the algebra of functions on Σ_k .) The CDGA $\mathbf{cd}_{mn}^\vee(k, l)$ is equipped with the obvious action of $\Sigma_k \times \Sigma_l$ and we can therefore view \mathbf{cd}_{mn}^\vee as a bisymmetric collection.

Proposition 3.15. *There is an isomorphism of algebras $\mathbf{cd}_{mn}^\vee(k, l) \cong H^*(\mathbf{CD}_{mn}(k, l))$.*

Proof. This follows directly from the Künneth formula and the homotopy equivalence $\mathbf{CD}_{mn}(k, l) \simeq \text{Conf}_{\mathbb{R}^m}(k) \times \text{Conf}_{\mathbb{R}^n \setminus \mathbb{R}^m}(l)$. \square

We now turn to the cooperad structure of \mathbf{cd}_{mn}^\vee . In \mathbf{e}_n^\vee , we have $\circ_W^\vee(\omega_{uv}) = 1 \otimes \omega_{uv}$ if $u, v \in W$, and $\circ_W^\vee(\omega_{uv}) = \omega_{[u][v]} \otimes 1$ otherwise. Moreover, $\mathbf{e}_1^\vee(U) = \mathbf{Ass}^\vee(U) = \mathbb{R}[\Sigma_U]^\vee$ is the algebra of functions on $\Sigma_U := \text{Bij}(U, \{1, \dots, \#U\})$, the set of linear orders on U . The cocomposition of \mathbf{e}_1^\vee is dual to block composition:

$$\Sigma_{U/W} \times \Sigma_W \rightarrow \Sigma_U, \quad (\sigma, \tau) \mapsto \sigma \circ_W \tau, \quad (3.16)$$

where $\sigma \circ_W \tau$ is the linear order on U obtained by inserting the order on W at the position $*$ in the order σ , e.g. $(a < * < b) \circ_{\{x, y\}} (x < y) = (a < x < y < b)$.

Proposition 3.17. *For $m \geq 2$, the isomorphism of Proposition 3.13 induces cooperadic structure maps $\circ_T^\vee : \mathbf{cd}_{mn}^\vee(U, V) \rightarrow \mathbf{cd}_{mn}^\vee(U, V/T) \otimes \mathbf{e}_n^\vee(T)$ and $\circ_{W, T}^\vee : \mathbf{cd}_{mn}^\vee(U, V \sqcup T) \rightarrow \mathbf{cd}_{mn}^\vee(U/W, V) \otimes \mathbf{cd}_{mn}^\vee(W, T)$ given by:*

$$\begin{aligned} \circ_T^\vee(\tilde{\omega}_{uu'}) &= \tilde{\omega}_{uu'} \otimes 1. & \circ_{W, T}^\vee(\tilde{\omega}_{uu'}) &= \begin{cases} 1 \otimes \tilde{\omega}_{uu'}, & \text{if } u, u' \in W; \\ \tilde{\omega}_{[u][u']} \otimes 1, & \text{otherwise.} \end{cases} \\ \circ_T^\vee(\omega_{vv'}) &= \begin{cases} 1 \otimes \omega_{vv'}, & \text{if } v, v' \in T; \\ \omega_{[v][v']} \otimes 1 & \text{otherwise.} \end{cases} & \circ_{W, T}^\vee(\omega_{vv'}) &= \begin{cases} 1 \otimes \omega_{vv'}, & \text{if } v, v' \in T; \\ \omega_{vv'} \otimes 1 & \text{if } v, v' \in V; \\ 0 & \text{otherwise.} \end{cases} \\ \circ_T^\vee(\eta_v) &= \eta_{[v]} \otimes 1. & \circ_{W, T}^\vee(\eta_v) &= \begin{cases} \eta_v \otimes 1 & \text{if } v \in V; \\ 1 \otimes \eta_v & \text{if } v \in T. \end{cases} \end{aligned}$$

For $m = 1$, the maps \circ_T^\vee and $\circ_{W, T}^\vee$ have the same behavior as above on the generators $\omega_{vv'}$ and η_v . On $\mathbf{e}_1^\vee(U) = \mathbf{Ass}^\vee(U) = \mathbb{R}[\Sigma_U]^\vee$, the map \circ_T^\vee is the identity, and $\circ_{W, T}^\vee$ is the dual of the block composition of Equation (3.16).

Proof. Our proof is similar to [46, Theorem 6.3]. Recall the maps $\theta_{uu'} : \mathbf{CD}_{mn}(U, V) \rightarrow S^{m-1}$, $\theta_{vv'} : \mathbf{CD}_{mn}(U, V) \rightarrow S^{n-1}$, and $\alpha_v : \mathbf{CD}_{mn}(U, V) \rightarrow S^{n-m-1}$ from Section 2. Let us first check that, when they are composed with the insertion maps $\circ_T : \mathbf{CD}_{mn}(U, V/T) \times \mathbf{D}_n(T) \rightarrow \mathbf{CD}_{mn}(U, V)$, we obtain the behavior indicated above on generators. For $\theta_{uu'}$ and $\theta_{vv'}$, this is identical to the proof that \mathbf{e}_n^\vee (resp. \mathbf{sc}_m^\vee) is the cohomology of \mathbf{D}_n (resp. \mathbf{SC}_m). Now, let us consider the composite $\alpha_v(- \circ_T -) : \mathbf{CD}_{mn}(U, V/T) \times \mathbf{D}_n(T) \rightarrow S^{n-m-1}$ for some $v \in V$. If $v \notin T$, then the composite map $\alpha_v(- \circ_T -)$ is equal to $\alpha_v \circ \text{proj}_1$, therefore:

$$\circ_T^\vee(\eta_v) = (\alpha_v(- \circ_T -))^*(\text{vol}_{n-m-1}) = \text{proj}_1^*(\alpha_v^*(\text{vol}_{n-m-1})) = \eta_v \otimes 1. \quad (3.18)$$

If $v \in T$, consider the homotopy which precomposes the embedding indexed by $[v] \in V/T$ with $x \mapsto tx$. At the limit $t = 0$, we find the map $\alpha_* \circ \text{proj}_1$. Since the homotopy class of the map is constant as t varies, we also get $\circ_T^\vee(\eta_v) = \eta_{[v]} \otimes 1$.

Now consider $\circ_{W,T} : \text{CD}_{mn}(U/W, V) \times \text{CD}_{mn}(W, T) \rightarrow \text{D}_{mn}(U, V \sqcup T)$ and let us check that we get the maps from the proposition. For $\theta_{uu'}$, this is again identical to the computation for e_n^\vee . For η_v , it is easy to see that $\eta_v(- \circ_{W,T} -)$ factors by the projection on one of the two factors in the product. Similarly, for $\theta_{vv'}$, if v, v' are both in V or both in T , then $\theta_{vv'}(- \circ_{W,T} -)$ also factors by one of the projections. Otherwise, WLOG assume that $v \in V$ and $v' \in T$. If we contract the appropriate disks by a homotopy which linearly decreases the radius, then the limit is a constant map, thus $\circ_{W,T}^\vee(\omega_{vv'}) = 0$. \square

Remark 3.19. We can compare cd_{mn}^\vee with the cohomology of ESC_{mn} (see Remark 2.11). We have $H^*(\text{ESC}_{mn}(U, V)) = e_m^\vee(U) \otimes_{e_n^\vee(U)} e_n^\vee(U \sqcup V)$, where $e_n^\vee(U) \rightarrow e_n^\vee(U \sqcup V)$ is the obvious inclusion and $e_n^\vee(U) \rightarrow e_m^\vee(U)$ sends all the generators $\omega_{uu'}$ to zero [55, Proposition 4.1]. Thus $H^*(\text{ESC}_{mn}(U, V)) = e_m^\vee(U) \otimes_{e_n^\vee(U \sqcup V)/e_n^\vee(U)}$. The cooperadic structure maps are defined by formulas similar to Proposition 3.17 (forgetting the η_v). The inclusion $\text{CD}_{mn} \subset \text{ESC}_{mn}$ induces on cohomology the composite $e_m^\vee(U) \otimes_{e_n^\vee(U \sqcup V)/e_n^\vee(U)} \rightarrow e_m^\vee(U) \otimes_{e_n^\vee(V)} \hookrightarrow \text{cd}_{mn}^\vee(U, V)$. An interesting question would be whether this inclusion is formal.

Remark 3.20. The equality $\text{CD}_{mn}(-, \emptyset) = \text{D}_m$ induces a left D_m -module structure on $\text{CD}_{mn}(\emptyset, -)$, using the operad structure of CD_{mn} . This structure dualizes to the map $\Delta : \text{cd}_{mn}^\vee(\emptyset, \bigsqcup_{u \in U} V_u) \rightarrow \text{cd}_{mn}^\vee(U, \emptyset) \otimes \bigotimes_{u \in U} \text{cd}_{mn}^\vee(\emptyset, V_u)$ given by $\Delta(\eta_v) = 1 \otimes \eta_v$ (put in the corresponding $\text{cd}_{mn}^\vee(\emptyset, V_u)$), $\Delta(\omega_{vv'}) = 1 \otimes \omega_{vv'}$ if v and v' are in the same V_u and $\Delta(\omega_{vv'}) = 0$ otherwise.

3.2. Generators and relations for the homology

We take the opportunity to describe a presentation of $\text{cd}_{mn} := H_*(\text{CFM}_{mn})$ by generators and relations. Let us first recall the presentation of $e_n := H_*(\text{FM}_n)$, due to Cohen [9]. An algebra over e_1 is a unital associative algebra. For $n \geq 2$, an algebra over e_n is a unital Poisson n -algebra, i.e. a unital commutative algebra equipped with a Lie bracket of (cohomological) degree $1 - n$ which is a biderivation for the product.

Proposition 3.21. *For $n - 2 \geq m \geq 1$, an algebra over cd_{mn} is the data (A, B, f, δ) consisting of an e_m -algebra A , an e_n -algebra B , a central morphism of algebras $f : B \rightarrow A$, and its central derivation $\delta : B[n - m - 1] \rightarrow A$.*

Central means that f and δ land in the center $Z(A) = \{a \in A \mid \forall b \in A, [a, b] = 0\}$, where the bracket is the graded commutator ($m = 1$) or the shifted Lie bracket ($m \geq 2$). The map δ is a derivation with respect to f : $\delta(xy) = \delta(x)f(y) \pm f(x)\delta(y)$.

Proof. An algebra is a pair (A, B) where B is an e_n -algebra and A is a cd_{mn} -algebra. Since $\text{cd}_{mn}(-, \emptyset) = e_m$, we know that A is an e_m -algebra. Given that $\text{cd}_{mn}(0, 1) = H_*(S^{n-m-1})$, we obtain two classes $f : B \rightarrow A$ of degree 0 and $\delta : B[n - m - 1] \rightarrow A$ of degree $1 + m - n$. The two possible compositions $\text{cd}_{mn}(2, 0) \times \text{cd}_{mn}(0, 1) \rightarrow \text{cd}_{mn}(1, 1)$ in either input of the

product are homotopic, so f and δ are central. Similarly, f is a morphism of algebras. The proof that δ is a derivation is the same as the proof that the Lie bracket in \mathfrak{e}_n is a biderivation.

We thus get a suboperad of \mathbf{cd}_{mn} that contains exactly the data of the proposition. Elements of this suboperad can be described as forests of rooted binary trees (like elements of \mathfrak{e}_n) with roots possibly decorated by a loop of degree $1 + m - n$ (the element δ). These are exactly the trees that appeared in the proof of Proposition 3.13 and we proved there that they generate the whole homology. \square

We can rephrase Proposition 3.21 more compactly. Let A be an \mathfrak{e}_m -algebra and consider $A[\varepsilon] := A \otimes \mathbb{R}[\varepsilon]/(\varepsilon^2)$ where $\deg \varepsilon = n - m - 1$. If $m \geq 2$, then there is a Poisson bracket on $A[\varepsilon]$ given by $[x + \varepsilon y, x' + \varepsilon y'] = [x, x'] + \varepsilon([x, y'] \pm [x', y])$. A \mathbf{cd}_{mn} -algebra is the data of an \mathfrak{e}_m -algebra A , an \mathfrak{e}_n -algebra B , and a central morphism $f + \varepsilon\delta : B \rightarrow A[\varepsilon]$.

Compare this result with the ∞ -categorical counterparts from [4, Section 4.3]. An algebra over the ∞ -categorical version $\mathcal{D}isk_{m \subset n}^{\text{fr}}$ consist of a $\mathcal{D}isk_m^{\text{fr}}$ -algebra A , a $\mathcal{D}isk_n^{\text{fr}}$ -algebra B , and a morphism of $\mathcal{D}isk_{m+1}^{\text{fr}}$ -algebras $\alpha : \int_{S^{n-m-1}} B \rightarrow \text{HC}_{\mathbb{D}_m}^*(A)$, where $\int_{S^{n-m-1}} B$ is the “factorization homology” of S^{n-m-1} with coefficients in B , and $\text{HC}_{\mathbb{D}_m}^*$ refers to Hochschild cochains. We view this as a “up to homotopy” version of an \mathbf{cd}_{mn} -algebra, the morphism $f + \varepsilon\delta$ being the part $\int_{S^{n-m-1}} B \rightarrow \text{HC}_{\mathbb{D}_m}^0(A)$ and the higher terms being homotopies. It would be interesting to make this observation precise.

Remark 3.22. An algebra over the homology $\mathbf{sc}_n := H_*(\mathbf{SC}_n)$ of the Swiss-Cheese operad is the data of (A, B, f) as in the proposition, see [33, Section 4.3]. However, our computation for $H^*(\mathbf{CD}_{(n-1)n})$ in Section 3.1 above does not apply (e.g. the class $\omega_{12} \in H^*(\mathbf{CD}_{(n-1)n}(0, 2))$ vanishes on some connected components). Instead, we get that $\mathbf{cd}_{(n-1)n}$ -algebras are given by quadruples (A, B, f, g) where (A, B, f) are as above and $g : B \rightarrow A$ is another central morphism (i.e. A is a unitary B -bimodule). There is an embedding $\mathbf{SC}_n \subset \mathbf{CD}_{(n-1)n}$. On homology, an $\mathbf{cd}_{(n-1)n}$ -algebra (A, B, f, g) viewed as an \mathbf{sc}_n -algebra is simply (A, B, f) , i.e. we forget the right action. Livernet [33, Theorem 4.3] proved that \mathbf{SC}_n is not formal by exhibiting a nontrivial operadic Massey product $\langle \mu_B, f, \lambda_A \rangle$, where μ_B represents the product of B and λ_A represents the Lie bracket of A . This shows that the operad of chains $C_*(\mathbf{CFM}_{(n-1)n}; \mathbb{Q})$ cannot be formal either, because we obtain a nontrivial Massey product there too.

4. Graph complexes

In this section, we define a bicolored Hopf cooperad, whose operations in the second color are given by Kontsevich’s cooperad \mathbf{graphs}_n [30], and whose operations in the first color will be called $\mathbf{cgraphs}_{mn}$. Our tool of choice to define $\mathbf{cgraphs}_{mn}$ will be “operadic twisting” [51, 11], just like in [53]. To give an idea of how $\mathbf{cgraphs}_{mn}$ is built, we first recall the steps in the definition of \mathbf{graphs}_n . We also refer to [27, Sections 1.5–1.6] for more details with matching notation.

Remark 4.1. We use the notation \mathbf{graphs}_n for the cooperad rather than its dual operad. Its linear dual \mathbf{graphs}_n^\vee is an operad which is quasi-isomorphic to $H_*(\mathbf{D}_n)$. We make this

choice because we never deal with \mathbf{graphs}_n^\vee . Kontsevich's graph complex GC_n^\vee where the differential splits vertices will also be denoted with a dual sign. See e.g. [27, 7] for matching notations.

4.1. Recollections: the cooperad \mathbf{graphs}_n

Untwisted The first step is to define an untwisted Hopf cooperad \mathbf{Gra}_n , given in each arity by the following CDGA, with generators e_{uv} of degree $n - 1$:

$$\mathbf{Gra}_n(U) := S(e_{uv})_{u,v \in U} / (e_{vu} = (-1)^n e_{uv}). \quad (4.2)$$

Graphically, $\mathbf{Gra}_n(U)$ is spanned by graphs on the set of vertices U . The monomial $e_{u_1 v_1} \dots e_{u_r v_r}$ corresponds to the graph with edges $\overrightarrow{u_1 v_1}, \dots, \overrightarrow{u_r v_r}$. The identification $e_{vu} = \pm e_{uv}$ allows us to view the graphs as undirected, although we need directions to define the signs precisely for odd n . If n is even then $\deg e_{uv}$ is odd, so we need to order edges to get precise signs. We explicitly allow tadpoles (e_{uu}) and double edges (e_{uv}^2). However, for even n , $e_{uv}^2 = 0$ because $\deg e_{uv}$ is odd; and for odd n , $e_{uu} = (-1)^n e_{uu} = -e_{uu}$ thus $e_{uu} = 0$. The product is given by gluing graphs along their vertices. The cooperadic structure is given by subgraph contraction. Explicitly, the map $\circ_W^\vee : \mathbf{Gra}_n(U) \rightarrow \mathbf{Gra}_n(U/W) \otimes \mathbf{Gra}_n(W)$ is given by $\circ_W^\vee(e_{uv}) = e_{**} \otimes 1 + 1 \otimes e_{uv}$ if $u, v \in W$, and by $\circ_W^\vee(e_{uv}) = e_{[u][v]} \otimes 1$ otherwise.

Remark 4.3. The quotient of \mathbf{Gra}_1 by disconnected graph forms an operad denoted by $\mathcal{G}r$ in [44]. It was used to present Harrison cohomology and find bases for cofree Lie coalgebras.

One may then produce a first zigzag of Hopf cooperads:

$$H^*(\mathbf{FM}_n) = \mathbf{e}_n^\vee \leftarrow \mathbf{Gra}_n \xrightarrow{\omega'} \Omega_{\mathrm{PA}}^*(\mathbf{FM}_n), \quad \omega_{uv} \leftarrow e_{uv} \mapsto p_{uv}^*(\varphi_n), \quad (4.4)$$

where $p_{uv} : \mathbf{FM}_n(U) \rightarrow \mathbf{FM}_n(2)$ is the projection and φ_n is the ‘‘propagator’’:

$$\varphi_n := \mathrm{vol}_{n-1} = \mathrm{cst} \cdot \sum_{i=1}^n \pm x_i dx_1 \wedge \dots \widehat{dx_i} \dots \wedge dx_n \in \Omega_{\mathrm{PA}}^{n-1}(\mathbf{FM}_n(2)) = \Omega_{\mathrm{PA}}^{n-1}(S^{n-1}). \quad (4.5)$$

Given a graph $\Gamma \in \mathbf{Gra}_n(U)$, one may define its *coefficient* $\mu(\Gamma)$ by $\mu(\Gamma) = 0$ if $\#U \leq 1$ and $\mu(\Gamma) = \int_{\mathbf{FM}_n(U)} \omega'(\Gamma)$ otherwise. This element μ has a simple description: it vanishes on all graphs, except for the one with exactly two vertices and an edge between them [32, Lemma 9.4.3]. In other words, in the dual basis:

$$\mu = \textcircled{1} \text{---} \textcircled{2} \in \mathbf{Gra}_n^\vee(2) \subset \prod_{i \geq 0} \mathbf{Gra}_n^\vee(i). \quad (4.6)$$

Twisting The second step is to twist the cooperad \mathbf{Gra}_n . Briefly, the idea of twisting is to turn an operad \mathbf{P} that encodes algebras that are also Lie algebras (i.e. there is a morphism $\mathrm{Lie} \rightarrow \mathbf{P}$) into an operad that encode \mathbf{P} -algebras twisted by a Maurer–Cartan element, see [51, Appendix I]. In the case of \mathbf{Gra}_n , this turns out to produce the Arnold

relations, which are themselves the algebraic shadow of the stratification of FM_n . See Appendix A.1 for concepts and notations.

The element μ of (4.6) defines a morphism from Lie_n to Gra_n^\vee : it sends the generating bracket $\lambda_2 \in \mathrm{Lie}_n(2)$ to the graph appearing in (4.6). By composing with the canonical resolution $\mathrm{hoLie}_n \rightarrow \mathrm{Lie}_n$, we thus obtain a morphism $\mathrm{hoLie}_n \rightarrow \mathrm{Gra}_n^\vee$, which sends the binary bracket to μ and the k -ary brackets to 0 for $k \geq 3$. One can check immediately that this graph satisfies the Jacobi relation. Alternatively, this can be deduced from the definition as an integral and the decomposition of the boundary of $\mathrm{FM}_n(U)$, cf. Proposition 4.32.

Cooperadic twisting produces a dg-cooperad $\mathrm{Tw Gra}_n$ from Gra_n and μ . Let us now describe it. As a graded module,

$$\mathrm{Tw Gra}_n(r) := \left(\bigoplus_{i \geq 0} (\mathrm{Gra}_n(r+i) \otimes \mathbb{k}[n]^{\otimes i})_{\Sigma_i}, d_\mu \right). \quad (4.7)$$

The CDGA structure is defined using the fact that $\mathrm{Gra}_n(\emptyset) = \mathbb{R}$ (see [27, Lemma 9]) and the fact that μ vanishes on disconnected graphs as well as graphs admitting a cut point (i.e. if a point that disconnects the graph when removed).

Let us now give a graphical interpretation of this definition. The CDGA $\mathrm{Tw Gra}_n(U)$ is spanned by graphs with two kinds of vertices: external vertices, which are in bijection with U , and internal vertices, which are indistinguishable (in pictures they will be drawn in black). Given a graph Γ , its differential $d\Gamma = \sum_e \pm \Gamma/e$ is obtained as a sum, over all the edges $e \in E_\Gamma$ connected to at least one internal vertex, of the graphs obtained by contracting these edges. Note that in this differential, edges connected to univalent vertices are not contracted, roughly speaking because the contraction appears twice in $d\Gamma$ and cancels out; but if both endpoints of the edge are univalent, then the edge is contracted with a minus sign, see [51, Appendix I.3] and Description 4.12. The product of two graphs is the graph obtained by gluing them along their external vertices. The cooperadic structure is given by subgraph contraction (summing over all choices of whether internal vertices and edges are in the subgraph).

One checks that the zigzag of Equation (4.4) extends to a zigzag:

$$e_n^\vee \leftarrow \mathrm{Tw Gra}_n \xrightarrow{\omega} \Omega_{\mathrm{PA}}^*(\mathrm{FM}_n). \quad (4.8)$$

This follows from the results of [32, Section 9–10] and [51, Section 3]. The left-pointing map sends all graphs with internal vertices to zero. The right-pointing map is given by the following integrals. Given a graph $\Gamma \in \mathrm{Gra}_n(U \sqcup I) \subset \mathrm{Tw Gra}_n(U)$, the form $\omega(\Gamma)$ is the integral of $\omega'(\Gamma)$ along the fiber of the PA bundle $p_U : \mathrm{FM}_n(U \sqcup I) \rightarrow \mathrm{FM}_n(U)$ which forgets points in the configuration:

$$\omega(\Gamma) := (p_U)_*(\omega'(\Gamma)) = \int_{\mathrm{FM}_n(U \sqcup I) \rightarrow \mathrm{FM}_n(U)} \omega'(\Gamma). \quad (4.9)$$

Graph complex fGC_n We record the following definition for future use.

Definition 4.10. The full graph complex fGC_n is defined to be $\mathrm{Tw}\mathrm{Gra}_n(\emptyset)[-n]$. It is spanned by graphs with only internal vertices. The degree of $\gamma \in \mathrm{fGC}_n$ is $\deg \gamma = (n-1)\#E_\gamma - n\#V_\gamma + n$. We describe its differential below.

Remark 4.11. This degree shift by n comes from the fact that we mod out by $\mathbb{R}^n \times \mathbb{R}_{>0}$ in the definition of FM_n . To get a Maurer–Cartan element (of degree 1) out of ω in the Lie algebra GC_n^\vee defined below, a shift by n is thus used.

The module fGC_n is a shifted algebra, with a product given by disjoint union of graphs. If $\mathrm{GC}_n \subset \mathrm{fGC}_n$ is the subcomplex of connected graphs, then there is an isomorphism of shifted algebras $\mathrm{fGC}_n = S(\mathrm{GC}_n[n])[-n]$. The module GC_n is a (pre)Lie coalgebra. Its cobracket Δ is given by subgraph contraction (i.e. it is inherited from the cooperad structure on Gra_n). Dually, GC_n^\vee is a (pre)Lie algebra, with a bracket given by graph insertion.

Description 4.12. Let us now describe the differential of GC_n . It is given by $(-\mu \otimes 1 + 1 \otimes \mu)\Delta$, where $\mu \in \mathrm{GC}_n^\vee$ is the Maurer–Cartan element defined in Equation (4.6). The summand $(1 \otimes \mu)\Delta$ is given by the sum of contractions of edges, while the summand $(\mu \otimes 1)\Delta$ is given by the sum of contractions of edges attached to a univalent vertex. The contraction of an edge attached to exactly one univalent vertex appears twice in the sum with opposite signs, which thus cancels out. However, if both endpoints of the edge are univalent, then the contractions of this edge appears once with a plus and twice with a minus, which leaves one term with a minus. (See [51, Appendix I.3] for more details)

Dually, the differential on GC_n^\vee is $[\mu, -]$. It is roughly speaking given by vertex splitting (with the caveat about univalent vertices explained above). The space $\mathrm{Tw}\mathrm{Gra}_n(U)$ is a module over the shifted CDGA fGC_n , by taking disjoint unions.

Reduction The next step is to mod out graphs with *internal components*, i.e. connected components with only internal vertices, to obtain a new Hopf cooperad Graphs_n . Formally, we consider the tensor product

$$\mathrm{Graphs}_n(U) := \mathrm{Tw}\mathrm{Gra}_n(U) \otimes_{\mathrm{fGC}_n} \mathbb{R}, \quad (4.13)$$

where fGC_n acts trivially on \mathbb{R} . In other words, in Graphs_n , a graph with a connected component consisting entirely of internal vertices is set equal to zero. The map ω of Equation (4.9) factors through the quotient defining Graphs_n [32, Lemma 9.3.7], and the quotient map $\mathrm{Tw}\mathrm{Gra}_n \rightarrow \mathrm{e}_n^\vee$ clearly does.

We can moreover reduce further the operad. The quotient graphs_n is given by modding out graphs containing: internal vertices that are univalent or bivalent; double edges; or tadpoles. It follows again from the lemmas of [32, Section 9.3] that ω factors through this quotient (and the map $\mathrm{Graphs}_n \rightarrow \mathrm{e}_n^\vee$ clearly does).

Recall that even though $\Omega_{\mathrm{PA}}^*(\mathrm{FM}_n)$ is not a Hopf cooperad, we take the convention that a morphism into it is a morphism in the sense defined in Section 1.3.

Theorem 4.14 ([30], [32, Chapter 10]). *This defines quasi-isomorphisms of Hopf cooperads $\mathrm{e}_n^\vee \xleftarrow{\sim} \mathrm{graphs}_n \xrightarrow{\sim} \Omega_{\mathrm{PA}}^*(\mathrm{FM}_n)$, thus FM_n is formal over \mathbb{R} .*

4.2. The cooperad \mathbf{CGr}_m

Let us now define \mathbf{CGr}_m , using the same methodology that was used to define \mathbf{Gr}_n . Note that we must take special care of the case $m = 1$. We define the further reduced cooperad \mathbf{cgr}_m in the next section.

Untwisted The first step is the definition of the untwisted graph cooperad.

Definition 4.15. For $n - 2 \geq m \geq 2$, the untwisted graph cooperad is a relative \mathbf{CGra}_m -cooperad given in each arity by $(\deg \tilde{e}_{uu'} = m - 1$ and $\deg e_{ij} = n - 1)$:

$$\mathbf{CGra}_m(U, V) := S(\tilde{e}_{uu'})_{u, u' \in U} \otimes S(e_{ij})_{i, j \in U \sqcup V} / (e_{ji} = (-1)^n e_{ij}, \tilde{e}_{uu'} = (-1)^m \tilde{e}_{u'u}).$$

Definition 4.16. For $n \geq 3$, we let (where $\Sigma_U = \text{Ass}(U) = \text{Bij}(U, \{1, \dots, \#U\})$):

$$\mathbf{CGra}_1(U, V) := S(e_{ij})_{i, j \in U \sqcup V} \otimes \mathbb{R}[\Sigma_U]^\vee / (e_{ji} = (-1)^n e_{ij}).$$

Let us now give a graphical interpretation of $\mathbf{CGra}_m(U, V)$. We will concentrate first on the case $m \geq 2$. As a vector space, it is spanned by graphs with two kinds of vertices: terrestrial (in bijection with U) and aerial (in bijection with V). We will draw the aerial vertices as circles, and the terrestrial vertices as semicircles, below the aerial ones. There are also two kind of edges: full edges (corresponding to the e_{ij}) between any two vertices, and dashed edges (corresponding to the $\tilde{e}_{uu'}$) between two terrestrial vertices. Note that we allow tadpoles and double edges. See Figure 4.1 for an example.




Figure 4.1: Example of a graph in \mathbf{CGra}_m (for $m \geq 2$)

If $m = 1$, the interpretation is similar, except that there are no dashed edges, and in addition the set of terrestrial vertices is ordered. Note that we are implicitly using the dual of the canonical basis of $\mathbb{R}[\Sigma_U]$ (i.e. we consider the basis elements of $\mathbb{R}[\Sigma_U]^\vee$ which vanish on all linear orders except one).

The product glues graphs along their vertices, and the differential is zero. These CDGAs assemble to form a relative Hopf \mathbf{Gra}_n -cooperad. The cooperadic structure maps act on the generators by the following formulas (compare with Proposition 3.17), with the convention $u, u' \in U, v, v' \in V$:

$$\circ_T^\vee(\tilde{e}_{uu'}) = \tilde{e}_{uu'} \otimes 1, \quad \circ_{W, T}^\vee(\tilde{e}_{uu'}) = \begin{cases} \tilde{e}_{**} \otimes 1 + 1 \otimes \tilde{e}_{uu'}, & \text{if } u, u' \in W; \\ \tilde{e}_{[u][u']} \otimes 1, & \text{otherwise.} \end{cases}$$

$$\circ_T^\vee(e_{vv'}) = \begin{cases} e_{**} \otimes 1 + 1 \otimes e_{vv'}, & \text{if } v, v' \in T; \\ \circ_T^\vee(e_{vv'}) = e_{[v][v']} \otimes 1, & \text{otherwise.} \end{cases} \quad \circ_{W,T}^\vee(e_{vv'}) = \begin{cases} 1 \otimes e_{vv'}, & \text{if } v, v' \in T; \\ e_{vv'} \otimes 1 & \text{if } v, v' \in V; \\ 0 & \text{otherwise.} \end{cases}$$

$$\begin{aligned} \circ_T^\vee(e_{uu'}) &= e_{uu'} \otimes 1, & \circ_{W,T}^\vee(e_{uu'}) &= \begin{cases} e_{**} \otimes 1 + 1 \otimes e_{uu'}, & \text{if } u, u' \in W, \\ e_{[u][u']} \otimes 1, & \text{otherwise,} \end{cases} \\ \circ_T^\vee(e_{uv}) &= e_{u[v]} \otimes 1, & \circ_{W,T}^\vee(e_{uv}) &= \begin{cases} e_{**} \otimes 1 + 1 \otimes e_{uv}, & \text{if } u \in W, v \in T \\ e_{u*} \otimes 1, & \text{if } u \notin W, v \in T, \\ e_{[u]v} \otimes 1 & \text{otherwise.} \end{cases} \end{aligned}$$

Moreover, if $m = 1$, then \circ_T^\vee is the identity on $\mathbb{R}[\Sigma_U]^\vee$, and $\circ_{W,T}^\vee$ is the dual of block composition from Equation (3.16). Graphically, the cooperadic structure map $\circ_{W,T}$ is given by subgraph contraction, which we now explain.

Definition 4.17 (Quotient graph, case $m \geq 2$). Let $\Gamma \in \text{CGra}_{mn}(U, V)$ be a graph and $\Gamma' \subset \Gamma$ be a subgraph, not necessarily full. We define the quotient graph Γ/Γ' as follows. The set of vertices of Γ/Γ' is the quotient set $V_\Gamma/V_{\Gamma'}$, identifying all the vertices of Γ' to produce a new vertex $[\Gamma']$, always terrestrial even if Γ' contains no terrestrial vertices. The set of edges of Γ/Γ' is the difference $E_\Gamma \setminus E_{\Gamma'}$ (we informally view the edges of Γ' as being “contracted”). If $e \in E_\Gamma \setminus E_{\Gamma'}$ has an endpoint in Γ' , then the corresponding endpoint of $e \in E_{\Gamma/\Gamma'}$ is the new terrestrial vertex $[\Gamma']$. In particular, if an edge is not in Γ' but both of its endpoints are, then this edge becomes a tadpole on the vertex $[\Gamma']$ in Γ/Γ' .

Example 4.18. The quotient Γ/\emptyset is Γ with a new isolated terrestrial vertex.

Definition 4.19 (Quotient graph, case $m = 1$). For $m = 1$, the definition of the quotient graph Γ/Γ' is slightly modified to deal with the linear order on terrestrial vertices. Suppose that $\Gamma' \subset \Gamma$ is a subgraph. The linear order of the terrestrial vertices of Γ/Γ' is dual to the block composition of linear orders from Equation (3.16). More concretely, Γ/Γ' is a linear combination of graphs:

- If Γ' has no terrestrial vertices, then Γ/Γ' is a sum of several terms. Each term is a quotient graph as in Definition 4.17. The sum ranges over all possible positions for the new vertex: if $f \in \mathbf{e}_1^\vee(U)$ is the decoration of Γ , then the decoration of Γ/Γ' is $(\sigma \in \Sigma_{U/\emptyset}) \mapsto f(\sigma|_U)$.
- If the terrestrial vertices of Γ' are consecutive, then Γ/Γ' is defined as in Definition 4.17, and the linear order on the terrestrial vertices of Γ/Γ' is inherited from Γ , with the new terrestrial vertex $[\Gamma']$ in the position of the vertices of Γ' .
- If the terrestrial vertices of Γ' are not consecutive, then $\Gamma/\Gamma' = 0$.

We can interpret $\circ_{W,T}(\Gamma)$ as the sum of the $\Gamma/\Gamma' \otimes \Gamma'$ for all (not necessarily full) subgraphs Γ' with set of vertices (W, T) .

Let us now define a zigzag of Hopf cooperad maps

$$H^*(\text{CFM}_{mn}) = \text{cd}_{mn}^\vee \leftarrow \text{CGra}_{mn} \rightarrow \Omega_{\text{PA}}^*(\text{CFM}_{mn}). \quad (4.20)$$

The left-pointing map is defined by $\tilde{e}_{uu} \mapsto 0$, $\tilde{e}_{uu'} \mapsto \tilde{\omega}_{uu'}$ for $u \neq u' \in U$, $e_{vv} \mapsto 0$, $e_{vv'} \mapsto \omega_{vv'}$ for $v \neq v' \in V$, and $e_{ij} \mapsto 0$ if $i \in U$ or $j \in U$. The right-pointing map is defined using the following three ‘‘propagators’’:

- Using the identification $\text{CFM}_{mn}(2, 0) = \text{FM}_m(2) \cong S^{m-1}$, we can use the propagator $\varphi_m := \text{vol}_{m-1} \in \Omega_{\text{PA}}^{m-1}(\text{FM}_m(2))$ of Equation (4.5).
- Recall the map $\theta_{12} : \text{CFM}_{mn}(1, 1) \rightarrow S^{n-1}$ which records the direction from the aerial point to the terrestrial point. The propagator ψ_{mn} is the pullback of the volume form of S^{n-1} along θ_{12} :

$$\psi_{mn}^\partial := \theta_{12}^*(\text{vol}_{n-1}) \in \Omega_{\text{PA}}^{n-1}(\text{CFM}_{mn}(1, 1)). \quad (4.21)$$

- Similarly, there is another map $\vartheta_{12} : \text{CFM}_{mn}(0, 2) \rightarrow S^{n-1}$ which records the direction from the second point to the first point. The propagator is:

$$\psi_{mn} := \vartheta_{12}^*(\text{vol}_{n-1}) \in \Omega_{\text{PA}}^{n-1}(\text{CFM}_{mn}(0, 2)). \quad (4.22)$$

Remark 4.23. By construction, these propagators are all minimal forms on CFM_{mn} , because the volume form on S^d is $\text{vol}_d = \text{cst} \cdot \sum_i (-1)^i x_i dx_1 \wedge \dots \wedge \widehat{dx}_i \wedge \dots \wedge dx_d$. Hence they can be pushed forward (once) along PA bundles [24].

Remark 4.24. The map $\text{CGra}_{mn} \rightarrow \text{cd}_{mn}^\vee$ is not surjective as its image does not contain the classes η_i . These classes are only reached once CGra_{mn} is twisted: it is the image of a graph with internal vertices (see Remark 4.52 and Section 5.2).

We may then define a morphism

$$\omega' : \text{CGra}_{mn}(U, V) \rightarrow \Omega_{\text{PA}}^*(\text{CFM}_{mn}(U, V)) \quad (4.25)$$

as follows, using the convention that $u, u' \in U$ and $v, v' \in V$:

$$\begin{aligned} \omega'(\tilde{e}_{uu'}) &:= p_{uu'}^*(\varphi_m); & \omega'(e_{vv'}) &:= p_{vv'}^*(\psi_{mn}); \\ \omega'(e_{uu'}) &:= 0; & \omega'(e_{vu}) &:= p_{vu}^*(\psi_{mn}^\partial). \end{aligned} \quad (4.26)$$

Here, $p_{ij}(x) = (p_i(x), p_j(x))$ (including if $i = j$). The following lemma is immediate:

Lemma 4.27. *This defines a zigzag $\text{cd}_{mn}^\vee \leftarrow \text{CGra}_{mn} \rightarrow \Omega_{\text{PA}}^*(\text{CFM}_{mn})$.* \square

Given a graph $\Gamma \in \text{CGra}_{mn}(U, V)$, we define $V_\Gamma = V_\Gamma^t \cup V_\Gamma^a = U \cup V$ to be its set of vertices, partitioned into terrestrial and aerial ones. Similarly, $E_\Gamma = E_\Gamma^f \cup E_\Gamma^d$ is its set of edges, split into full edges and dashed edges. The graph Γ induces:

$$\Phi_\Gamma : \text{CFM}_{mn}(U, V) \rightarrow (S^{m-1})^{E_\Gamma^d} \times (S^{n-1})^{E_\Gamma^f}, \quad (4.28)$$

obtained using the maps θ_{ij} from Section 2. We also define

$$\text{vol}_\Gamma \in \Omega_{\text{PA}}^{\text{deg } \Gamma}((S^{m-1})^{E_\Gamma^d} \times (S^{n-1})^{E_\Gamma^f}) \quad (4.29)$$

to be the product of the volume forms. Then by definition, $\omega'(\Gamma) = \Phi_\Gamma^*(\text{vol}_\Gamma)$.

Given a graph $\Gamma \in \text{CGra}_{mn}(U, V)$ with $\#U + 2\#V \geq 2$, we may define its *coefficient* $c(\Gamma)$ by

$$c(\Gamma) := \int_{\text{CFM}_{mn}(U, V)} \omega'(\Gamma). \quad (4.30)$$

If $\#U + 2\#V \leq 1$ then we just set $c(\Gamma) = 0$ (this is due to the special case in the definition of \mathcal{BF}'' , see Proposition 2.12).

Remark 4.31. In the case $(n, m) = (2, 1)$, these are analogous to the coefficients in Kontsevich’s universal formality morphism $T_{\text{poly}} \rightarrow D_{\text{poly}}$ from [31]. For $(n, m) = (3, 1)$, these are related to local versions of Kontsevich integrals [29], which are conjectured to equal Bott–Taubes integrals [39].

Twisting In this section, we are going to twist the cooperad CGra_{mn} using the theory developed in Appendix A.3. The general motivation of twisting is to encode Maurer–Cartan equations. Here, since we are going to twist with respect to the coefficients c , which come from the stratification of CFM_{mn} , we get a differential which matches the way the boundary facets of CFM_{mn} interact.

Recall the relative hoLie_n -operad hoLie_{mn} from Appendix A.3. Recall also the coefficient $\mu : \text{hoLie}_n \rightarrow \text{Gra}_n^\vee$ from Equation (4.6).

Proposition 4.32. *Together with μ , the collection of coefficients c defines a morphism of colored operads $(\text{hoLie}_n, \text{hoLie}_{mn}) \rightarrow (\text{Gra}_n^\vee, \text{CGra}_{mn}^\vee)$. The generating bracket $\lambda_{U, V} \in \text{hoLie}_{mn}(U, V)$ is sent to the element $c(\lambda_{U, V}) \in \text{CGra}_{mn}^\vee(U, V)$ given by $\Gamma \mapsto c(\Gamma)$, which we can write as the possibly infinite sum $c = \sum_{\Gamma} c(\Gamma)\Gamma$.*

Proof. The proof is similar to Kontsevich’s proof that μ defines a morphism $\text{hoLie}_n \rightarrow \text{Gra}_n^\vee$. Let C_*^{SA} be the functor of semi-algebraic chains [24, Section 3]. It is lax-monoidal [24, Proposition 3.8] so $C_*^{SA}(\text{CFM}_{mn})$ is a dg-operad. We can dualize Lemma 4.27 to get a morphism $I : C_*^{SA}(\text{CFM}_{mn}) \rightarrow \Omega_{\text{PA}}^*(\text{CFM}_{mn})^\vee \rightarrow \text{CGra}_{mn}^\vee, \sigma \mapsto \langle \omega'(-), \sigma \rangle$. The claimed morphism $c : \text{hoLie}_{mn} \rightarrow \text{CGra}_{mn}^\vee$ is the composition of I with:

$$\llbracket \text{CFM}_{mn} \rrbracket : \text{hoLie}_{mn} \rightarrow C_*^{SA}(\text{CFM}_{mn})$$

which maps the generating bracket $\lambda_{U, V} \in \text{hoLie}_{mn}(U, V)$ to the fundamental chain $\llbracket \text{CFM}_{mn}(U, V) \rrbracket$. So we just need to check that $\llbracket \text{CFM}_{mn} \rrbracket$ is a morphism.

Since hoLie_{mn} is quasi-free, we just need to check that our prescription on generators is compatible with the differential. Recall [24, Theorem 3.5] that for a compact oriented SA manifold M , $d\llbracket M \rrbracket = \llbracket \partial M \rrbracket$. If we use the description of $\partial \text{CFM}_{mn}(U, V)$ from Proposition 2.12, we find that the differential in $C_*^{SA}(\text{CFM}_{mn})$ matches the cobar differential. The faces of type \mathcal{BF}' correspond to cocompositions $\overrightarrow{\text{Com}}_{mn}^\vee(U, V) \rightarrow \overrightarrow{\text{Com}}_{mn}^\vee(U, V/T) \otimes \text{Com}_n^\vee(T)$, and the faces of type \mathcal{BF}'' correspond to cocompositions $\overrightarrow{\text{Com}}_{mn}^\vee(U, V) \rightarrow \overrightarrow{\text{Com}}_{mn}^\vee(U/W, V \setminus T) \otimes \overrightarrow{\text{Com}}_{mn}^\vee(W, T)$. The “missing” faces (when $\#T < 2$, $(W, T) = (U, V)$, or $2\#T + \#W < 2$) come from the fact that the cobar construction uses the coaugmentation quotient of the cooperad and from $\text{Com}_n^\vee(\emptyset) = \overrightarrow{\text{Com}}_{mn}^\vee(\emptyset, \emptyset) = 0$. \square

Definition 4.33. The twisted graph cooperad Tw CGra_{mn} is the relative (Tw Gra_n) -cooperad obtained by twisting CGra_{mn} with respect to μ and c .

This twisted cooperad has a graphical description. The CDGA $\text{Tw CGra}_{mn}(U, V)$ is spanned by graphs with four types of vertices: external terrestrial vertices, in bijection with U (drawn as semicircles); external aerial vertices, in bijection with V (drawn as circles); internal terrestrial vertices, indistinguishable among themselves, of degree $-m$ (drawn as black semicircles); internal aerial vertices, indistinguishable among themselves, of degree $-n$ (drawn as black circles).

There are two kinds of edges: full edges of degree $n - 1$, and dashed edges of degree $m - 1$. The product glues graphs along external vertices. If $m = 1$, then there are no dashed edges, and the whole set of terrestrial vertices is ordered (if we view $e_1^\vee(U \sqcup I)$ as spanned by the dual basis of $e_1(U \sqcup I)$). In this basis, the product of two graphs $\Gamma \cdot \Gamma'$ vanishes if the external vertices of Γ and Γ' are ordered differently. Otherwise $\Gamma \cdot \Gamma'$ is obtained by gluing the two graphs along external vertices, and summing all possible ways of ordering the union (along external vertices) of the terrestrial vertices of Γ and Γ' in a way that the new order restricts to the given orders in Γ and Γ' .

Description 4.34. The differential is given by the twisting procedure (Appendix A.3). Recall the subgraph contractions defined in Definitions 4.17, 4.19. Given a graph Γ , its differential $d\Gamma$ is given as a sum (with signs in Section A.3, the terms corresponding respectively to $(-) \cdot \mu$, $(-) \cdot (c - c_1)$, and $c_1 \cdot (-)$):

1. contractions of full edges between an aerial internal vertex and an aerial vertex of any kind, including edges connected to a univalent internal vertex (this uses the simple description of μ in Equation (4.6));
2. contractions of subgraphs Γ' containing at most one external (necessarily terrestrial) vertex, with result $c(\Gamma') \cdot \Gamma/\Gamma'$;
3. forgetting of all vertices outside a subgraph Γ'' which contains all the external vertices, with result $c(\Gamma/\Gamma'') \cdot \Gamma''$.

Note that the differential is not purely combinatorial: it depends on c , which is defined by integrals.

Since CGra_{mn} is a Hopf cooperad satisfying $\text{CGra}_{mn}(\emptyset, \emptyset) = \mathbb{k}$, and since c vanishes on disconnected graphs (see Lemma 4.37) and graphs with a terrestrial cut point (by a simple degree argument: $\dim \text{CFM}_{mn}(i+i'+1, j+j') > \dim \text{CFM}_{mn}(i+1, j) \times \text{CFM}_{mn}(i'+1, j')$), we find again that Tw CGra_{mn} inherits a Hopf cooperad structure, similarly to the uncolored case. For $m = 1$, this is also true: the differential preserves (up to quotient) the order of the terrestrial vertices, and if two consecutive terrestrial vertices in $\Gamma \cdot \Gamma'$ come respectively from Γ and Γ' (which could potentially produce an unwanted term in $d(\Gamma \cdot \Gamma')$ due to the exception in Lemma 4.37), then they also appear with the opposite order in the product and the two terms in the differential cancel out.

Lemma 4.35. For $n - 2 \geq m \geq 1$, the morphism ω' extends to a morphism of Hopf cooperads $\omega : \text{Tw CGra}_{mn} \rightarrow \Omega_{\text{PA}}^*(\text{CFM}_{mn})$ by setting, for $\Gamma \in \text{CGra}_{mn}(U \sqcup I, V \sqcup J) \subset$

Tw $\text{CGra}_{mn}(U, V)$ and $\#U + 2\#V \geq 2$:

$$\omega(\Gamma) := (p_{U,V})_*(\omega'(\Gamma)) = \int_{\text{CFM}_{mn}(U \sqcup I, V \sqcup J) \rightarrow \text{CFM}_{mn}(U, V)} \omega'(\Gamma).$$

If $\#U + 2\#V \leq 1$, then $\omega(\Gamma) = 1$ if Γ has no edges and no internal vertices, and $\omega(\Gamma) = 0$ otherwise.

Proof. We use the double pushforward formula [24, Proposition 8.13], Stokes' formula [24, Proposition 8.12] and the decomposition of the fiberwise boundary $\text{CFM}_{mn}(U, V)$ from Section 2.1 (compare with [32, Section 9.4]).

We deal separately with the case $\#U + 2\#V \leq 1$ (cf. [32, Section 9.1]). Indeed, the fiber of $p_{U,V} : \text{CFM}_{mn}(U \sqcup I, V \sqcup J) \rightarrow \text{CFM}_{mn}(U, V)$ has dimension $m\#I + n\#J$ in general, but $\dim \text{CFM}_{mn}(U, V) = 0$ for $\#U + 2\#V \leq 1$, so the dimension of the fiber of $p_{U,V}$ is either $m\#I + n\#J - 1$ or $m\#I + n\#J - m - 1$ and ω' would not preserve degrees. In these cases $\text{CFM}_{mn}(U, V)$ is merely a singleton, so we just have to check that any cocycle is mapped to zero. This is clear if $U = V = \emptyset$, as the differential of a nonempty graph is nonempty. For $\Gamma \in \text{Tw CGra}_{mn}(1, 0)$ with at least one edge, the graph with no edges appears exactly twice with opposite signs in $d\Gamma$ (see Description 4.34): for $\Gamma' = \Gamma$ in $-\Gamma \cdot c_1 = -c(\Gamma')\Gamma/\Gamma' + \dots$, and for Γ'' containing only the external vertex in $c_1 \cdot \Gamma = c(\Gamma/\Gamma'')\Gamma'' + \dots$. \square

Graph complex fCGC_{mn} We now mimic Definition 4.10:

Definition 4.36. For $n - 2 \geq m \geq 1$, we define $\text{fCGC}_{mn} := \text{Tw CGra}_{mn}(\emptyset, \emptyset)[-m]$.

This shifted CDGA is free and generated by its submodule of connected graphs. We still denote by $c \in \text{fCGC}_{mn}^\vee$ the restriction of the morphism c from (4.30).

Lemma 4.37. Let $n - 2 \geq m \geq 1$. Given a disconnected graph $\gamma \in \text{fCGC}_{mn}$, the coefficient $c(\gamma)$ vanishes, unless $m = 1$ and $\gamma = (\bullet \quad \bullet)$.

Proof. Let us first assume that γ has no isolated terrestrial vertices. Then γ factors as a product $\gamma = \gamma_1 \cdot \gamma_2$, with corresponding sets of vertices (I_1, J_1) and (I_2, J_2) both satisfying $\#I_\bullet + 2\#J_\bullet \geq 2$ for $\bullet \in \{1, 2\}$. We need $\deg \text{vol}_\gamma = \dim \text{CFM}_{mn}(I, J)$, otherwise $c(\gamma) = \int_{\text{CFM}_{mn}(I, J)} \Phi_\gamma^*(\text{vol}_\gamma)$ obviously vanishes. But thanks to the hypothesis Φ_γ factors through $\text{CFM}_{mn}(I_1, J_1) \times \text{CFM}_{mn}(I_2, J_2)$, which is of strictly lower dimension. Hence $\Phi_\gamma^*(\text{vol}_\gamma) = 0 \implies c(\gamma) = 0$.

Now let γ have an isolated terrestrial vertex $i \in I$. Then Φ_γ factors through $\text{CFM}_{mn}(I \setminus \{i\}, J)$. If $(\#I - 1) + 2\#J \geq 2$, then the codimension with $\text{CFM}_{mn}(I, J)$ is $m > 0$, and so similarly $c(\gamma) = 0$. However, if γ is the graph with two terrestrial vertices and no aerial ones (i.e. $(\#I, \#J) = (2, 0)$), then the codimension is $m - 1$, because $\dim \text{CFM}_{mn}(1, 0) = 0$. Thus we only get $c(\gamma) = 0$ if $m > 1$. \square

Remark 4.38. In the case $m = 1$, we have that $\text{CFM}_{1n}(2, 0) \cong \text{FM}_1(2) \cong S^0$. We find that for $\gamma = (\bullet \quad \bullet) \in \text{fCGC}_{1n}$, we have $c(\gamma) = 1$.

Definition 4.39. For $n - 2 \geq m \geq 2$, the connected complementarily constrained graph complex CGC_{mn} as the quotient of fCGC_{mn} by disconnected graphs.

The case $m = 1$ is special. Mimicking [53, Section 4], we define an alternate basis of $\text{Tw Gra}_{1n}(U, V)$. Recall that $\text{Gra}_{1n}(U, \emptyset)$ contains $\text{Ass}^\vee(U)$, where Ass encodes associative algebras. The symmetric sequence Ass is isomorphic to $\text{Pois} = \text{Com} \circ \text{Lie}$ [34, Section 13.3]. We can thus decree that terrestrial internal vertices are in the same ‘‘Lie component’’ if they are connected by brackets.

Definition 4.40 (Cf. [53, Section 4]). A Lie-decorated graph is a graph Γ defined similarly as an element of $\text{Tw Gra}_{1n}(U, V)$ (Definition 4.33), but instead of giving a function on linear orders of terrestrial vertices, we give a function on commutative products of Lie words of terrestrial vertices. More precisely, we use the dual of the PBW isomorphism to get an isomorphism and we obtain a new basis of $\text{Tw CGra}_{1n}(U, V)$ through the twisting procedure:

$$\text{CGra}_{1n}(U, V) \cong S(e_{ij})_{i,j \in U \sqcup V} \otimes \text{Pois}^\vee(U) / (e_{ji} = (-1)^n e_{ij}).$$

One possible way to describe a Lie-decorated graph $\Gamma \in \text{Tw CGra}_{1n}(U, V)$ (with internal vertices I) is to take a usual basis of $\text{Pois}(U \sqcup I) = (\text{Com} \circ \text{Lie})(U \sqcup I)$ and consider its dual basis in $\text{Pois}^\vee(U \sqcup I)$, i.e. functions on products of Lie words that vanish on all such elements except one. See Figure 4.2 for an example.




Figure 4.2: Lie-decorated graph; the depicted product of Lie words $[x_1, x_2] \wedge x_3 \wedge x_4$ is actually the element in the dual basis of $\text{Pois}^\vee(4)$

It is actually easier to describe the differential in of a Lie-decorated graph using coLie words. Recall that a coLie word U is an element in the cofree coalgebra $T^{\geq 1}(U)^\vee$ which vanishes on the image of $S^{\geq 2}(\text{Lie}(U))$ under the PBW isomorphism. These words are the image of the first Eulerian idempotent $e^{(1)} = \log_\star(\text{id})$, where \star is the convolution product of the bialgebra $T(U)^\vee$ (see e.g. [14, Appendix 4.2]). CoLie words themselves are described by graphs (see [44]), one can also thus think of Lie-decorated graphs as ‘‘graph-decorated graphs’’. With this basis, the differential merges coLie words (with a description similar to the one of Tw CGra_{mn} above).

Lemma 4.41. *The vector space spanned by Lie-decorated graphs with external vertices (U, V) is isomorphic to $\text{Tw Gra}_{1n}(U, V)$.*

Proof. We use the dual of the PBW isomorphism $\text{Pois}^\vee(U \sqcup I) \cong \text{Ass}^\vee(U \sqcup I)$ to identify linear combinations of Lie-decorated graphs with elements of $\text{Tw Gra}_{1n}(U, V)$. For the signs, terrestrial vertices have degree -1 . \square

Definition 4.42. Two vertices in a Lie-decorated graph are in the same Lie component if they are connected by full edges or if they are both terrestrial and in the same Lie word. A Lie-decorated graph is Lie-connected if all its vertices are in the same Lie component, and Lie-disconnected otherwise.

Lemma 4.43. *The coefficient c vanishes on the image of a Lie-disconnected graph under the isomorphism of Lemma 4.41.*

Proof. This follows from Lemma 4.37. □

Definition 4.44. For $n \geq 3$, the Lie-connected complementarily constrained graph complex CGC_{1n} as the quotient of fCGC_{1n} by the image of the Lie-disconnected graph under the isomorphism of Lemma 4.41.

Lemma 4.45. *For $n - 2 \geq m \geq 1$, the coefficient $c : \text{fCGC}_{mn} \rightarrow \mathbb{R}$ factors through the quotient defining CGC_{mn} . Abusing notation, we denote by c the induced map $\text{CGC}_{mn} \rightarrow \mathbb{R}$.*

Proof. This also follows from Lemma 4.37, in both cases $m = 1$ and $m \geq 2$. □

Remark 4.46. The dual CGC_{mn}^\vee is a (pre-)Lie algebra and the Lie algebra GC_n^\vee acts on it by derivations, in both cases using insertion of graphs (respectively at terrestrial and aerial vertices). The differential of CGC_{mn}^\vee is given by $[\mu + c, -]$ where c represents the coefficients from Equation (4.30). The elements μ and c satisfy the Maurer–Cartan equation, i.e. $[\mu, \mu] = 0$ and $[\mu, c] + \frac{1}{2}[c, c] = 0$.

Reduction We now mod out internal components.

Definition 4.47. For $n - 2 \geq m \geq 2$, we define the graph cooperad CGraphs_{mn} to be the relative Graphs_n -cooperad given by quotient of Tw CGra_{mn} by the Hopf cooperadic ideal of graphs with internal components. For $n - 2 \geq m = 1$, we define CGraphs_{1n} to be the quotient of Tw CGra_{1n} by the Hopf cooperadic ideal of Lie-disconnected graphs (Definition 4.42).

Remark 4.48. Compare this with Willwacher’s model SGraphs_n for the Swiss-Cheese operad [53]. In SGraphs_n , there are no dashed edges, the full edges are oriented, and their source is always aerial.

Proposition 4.49. *The morphism ω factors through the quotient and defines a morphism $\omega : \text{CGraphs}_{mn} \rightarrow \Omega_{\text{PA}}^*(\text{CFM}_{mn})$.*

Proof. The proof is identical to the proof of [32, Lemma 9.3.7]. Briefly, let $\gamma \in \text{Tw CGra}_{mn}(U, V)$ be a graph whose edges all are between internal vertices. Let (I, J) be the sets of internal vertices of γ and assume $\#U + 2\#V \geq 2$ otherwise the claim

is obvious (see Lemma 4.35). We can fill the diagram, where ρ is the product of two projections:

$$\begin{array}{ccc}
\mathrm{CFM}_{mn}(U \sqcup I, V \sqcup J) & \xrightarrow{\Phi_\gamma} & (S^{m-1})^{E_\Gamma^t} \times (S^{n-1})^{E_\Gamma^a} \\
\downarrow p_{U,V} & \searrow \rho & \nearrow \exists \Phi'_\gamma \\
\mathrm{CFM}_{mn}(U, V) & \xleftarrow{\mathrm{pr}_1} \mathrm{CFM}_{mn}(U, V) \times \mathrm{CFM}_{mn}(I, J) &
\end{array}$$

We therefore find that $\omega(\Gamma) = (p_{U,V})_*(\omega'(\Gamma)) = (p_{U,V})_*(\rho^*(\Phi'_\gamma \mathrm{vol}_\Gamma))$. The dimension of the fibers of $p_{U,V}$ and pr_1 are respectively $m\#I + n\#J > m\#I + n\#J - m$. Therefore $\omega(\Gamma) = 0$ by [24, Proposition 5.1.2]. For $m = 1$, we note that a Lie-disconnected graph is in particular disconnected. \square

4.3. Vanishing lemmas and $\mathrm{cgraphs}_{mn}$

We now prove some vanishing lemmas about ω and c , which allows us to define the further reduced cooperad $\mathrm{cgraphs}_{mn}$. This will be useful to prove that the quotient map $\mathrm{CGra}_{mn} \rightarrow \mathrm{cd}_{mn}^\vee$ extends to $\mathrm{cgraphs}_{mn}$ in Section 5.

Lemma 4.50. *The morphism ω' vanishes on graphs with loops or double edges.*

Proof. This follows from simple dimension arguments, cf. [32, Section 9.3]. \square

Lemma 4.51 (Cf. [32, Lemma 9.3.9]). *The morphism ω vanishes on graphs containing univalent aerial internal vertices, or univalent terrestrial internal vertices connected to another terrestrial vertex.*

Proof. The lemma is trivial if $\#U + 2\#V \leq 1$ (see Lemma 4.35), so let us assume that we are not in this case.

Let us first deal with univalent aerial internal vertices connected to another aerial vertex. The graph $\Gamma_{\mathrm{uni}} = \textcircled{1} \text{---} \bullet$ is of negative degree, thus $\omega(\Gamma_{\mathrm{uni}}) = 0$. For a general graph Γ containing Γ_{uni} , the argument is the same as [27, Corollary 44] or [32, Lemma 9.3.9]. The map Φ_Γ of (4.28) factors through $\mathrm{CFM}_{mn}(\emptyset, \{v, j\}) \times \mathrm{CFM}_{mn}(U \sqcup I, V \sqcup J \setminus \{j\})$, and so does the canonical projection. The two factorizations make the obvious diagram (like in Proposition 4.49) commute. Using [24, Propositions 8.10, 8.13] and $\omega(\Gamma_{\mathrm{uni}}) = 0$, we get $\omega(\Gamma) = 0$.

Let us now assume that Γ is a graph with a univalent internal vertex connected i to a terrestrial vertex u . We can assume that i is aerial or that the incident edge is dashed, as otherwise $\omega(\Gamma) = 0$ by definition (4.26). Let $(U \sqcup I, V \sqcup J)$ be the sets of vertices of Γ . Then both Φ_Γ and $p_{U,V}$ factor through either $X = \mathrm{CFM}_{mn}(\{u\}, \{i\}) \times \mathrm{CFM}_{mn}(U \sqcup I, V \sqcup J \setminus \{i\})$ (if i is aerial) or $X = \mathrm{CFM}_{mn}(\{u, i\}, \emptyset) \times \mathrm{CFM}_{mn}(U \sqcup I \setminus \{i\}, V \sqcup J)$ if i is terrestrial, and the factorizations make the obvious diagram commute. The dimension of the fiber of $p_{U,V}$ is $m\#I + n\#J$, while the dimension of the fiber of the projection $X \rightarrow \mathrm{CFM}_{mn}(U, V)$ is $m\#I + n\#J - 1$ in both cases. Hence, by [24, Proposition 8.14], $\omega(\Gamma) = 0$. \square

Remark 4.52. The argument fails for graphs containing a univalent internal terrestrial vertex connected to an aerial vertex. If we consider the following graph, then we find that $\omega(\Gamma) \in \Omega_{\text{PA}}^{n-m-1}(\text{CFM}_{mn}(0, 1))$ represents η from Section 3.1:

$$\Gamma = \textcircled{1} \text{---} \bullet \in \text{cgraphs}_{mn}(0, 1), \quad (4.53)$$

Indeed, by the definition of ψ_{mn}^∂ in (4.21) and by [24, Proposition 8.10], we find that $\int_{\text{CFM}_{mn}(1,1)} \psi_{mn}^\partial = \int_{S^{n-1}} \text{vol}_{n-1} = 1$. Therefore $\int_{\text{CFM}_{mn}(0,1)} \omega(\Gamma) = 1$ by [24, Proposition 8.13]. This also implies that if γ is obtained from the Γ above by making the external vertex internal, then we get $c(\gamma) = 1$.

Lemma 4.54. *The coefficient c vanishes on graphs with more than two vertices and which either contain a univalent aerial vertex, or which contain a univalent terrestrial vertex connected to another terrestrial vertex. It also vanishes on the graph with exactly two vertices, both aerial, and a (full) edge between the two.*

Proof. We can reuse the proof of Lemma 4.51 almost verbatim. There is one caveat: the graph γ with the univalent vertex removed must still satisfy the hypothesis $\#I + 2\#J \geq 2$. Otherwise, a degree shift occurs, to deal with the fact that $\dim \text{CFM}_{mn}(I, J) = 0$ and not -1 or $-m - 1$ as the general formula would give. This is the case unless γ has exactly two vertices with at most one aerial. If γ has exactly two aerial vertices and one edge, then $c(\gamma)$ vanishes for degree reasons ($\dim \text{CFM}_{mn}(0, 2) = 2n - m - 1 > n - 1$). \square

Remark 4.55. The restriction about the number of vertices is necessary. Indeed, for $\gamma = \bullet \text{----} \bullet \in \text{CGC}_{mn}$. we find $c(\gamma) = \int_{\text{CFM}_{mn}(2,0)} \varphi_m = \int_{S^{m-1}} \text{vol}_{S^{m-1}} = 1$.

Lemma 4.56. *The morphism ω and the coefficient c vanish on graphs with bivalent internal terrestrial vertices connected to two terrestrial vertices.*

Proof. Let us first consider the case of the graph

$$\Gamma_{\text{biv}} = \textcircled{u} \text{----} \bullet \text{----} \textcircled{v}. \quad (4.57)$$

Using $\text{CFM}_{mn}(U, \emptyset) \cong \text{FM}_m(U)$ and the fact that the terrestrial propagator is the same as the one used in the definition of the map $\text{Graphs}_m \rightarrow \Omega_{\text{PA}}^*(\text{FM}_m)$, we deduce that $\omega(\Gamma_{\text{biv}}) = 0$ from [32, Lemma 9.3.9] (see also [29, Lemma 2.1] for the origin of this result: there is an involution on $\text{FM}_m(U)$ which leaves $\omega'(\Gamma_{\text{biv}})$ invariant but reverses the orientation, thus $\omega(\Gamma_{\text{biv}}) = -\omega(\Gamma_{\text{biv}})$). If Γ contains Γ_{biv} as a subgraph, then we use the same reasoning as Lemma 4.51 to show that $\omega(\Gamma) = 0$. Finally, for the other cases, simply note that ω and c vanish by definitions on graphs with full edges between terrestrial vertices. \square

Definition 4.58. Let $I(U, V) \subset \text{CGraphs}_{mn}(U, V)$ be the vector space spanned by graphs containing loops, double edges (of any type), univalent aerial internal vertices, univalent internal terrestrial vertices connected to another terrestrial vertex, or bivalent internal terrestrial vertices connected to two terrestrial vertices by dashed edges. For $m = 1$, we take graphs containing loops, double edges, and univalent aerial internal vertices.

Lemma 4.59. *The subspace $I(U, V)$ define a CDGA ideal in $\mathbf{CGraphs}_{mn}(U, V)$, and the subcollection $I \subset \mathbf{CGraphs}_{mn}$ define a cooperadic coideal.*

Proof. It is clear that $I(U, V)$ is an algebra ideal. Let us show that it is a differential ideal. The fact that I defines a cooperadic coideal can also be checked easily case-by-case. We deal with $m \geq 2$, and the case $m = 1$ is also checked by a similar case-by-case argument. Let $\Gamma \in I(U, V)$ be a graph. Let us show that $d\Gamma \in I(U, V)$. The three summands in Description 4.34 are called $d_1 = (-) \cdot (\mu - \mu_1)$, $d_2 = (-) \cdot (c - c_1)$, and $d_3 = (c_1) \cdot (-)$, where the coefficients c are defined analytically by integrals.

- If Γ contains a loop, then all the summands in $d\Gamma$ contain a loop, as both μ and c vanish on loops.
- Suppose Γ contains a univalent aerial internal vertex i , with incident edge e . In $d_2\Gamma = \sum_{\Gamma'} c(\Gamma')\Gamma/\Gamma'$, if Γ' contains e then $c(\Gamma')$ vanishes by Lemma 4.54, and otherwise Γ/Γ' contains either a loop (if $i \in \Gamma'$) or a univalent aerial internal vertex (if $i \notin \Gamma'$). Similarly, all terms in $d_3\Gamma = \sum_{\Gamma''} c(\Gamma/\Gamma'')\Gamma''$ either vanish, contain a loop, or contain a univalent aerial internal vertex. The only problem in $d_3\Gamma$ is for $\Gamma'' = \Gamma \setminus e$: then $c(\Gamma/\Gamma'') = 1$ (see Equation (5.2)) and Γ'' does not have a univalent vertex anymore. But this term cancels with the contraction of e in $d_1\Gamma$, and all the other summands of $d_1\Gamma$ contain a univalent aerial internal vertex.
- If Γ contains a univalent terrestrial vertex connected to another terrestrial vertex, then Lemma 4.54 and an argument similar to the previous one show that all summands of $d\Gamma$ either vanish, contain a loop, or contain a univalent terrestrial vertex connected to another terrestrial vertex.
- Suppose that Γ contains a bivalent internal terrestrial vertices connected to two terrestrial vertices by dashed edges. Let i be the internal terrestrial vertex, and e, e' its incident edges. All terms in $d_1\Gamma$ contain a similar subgraph. In $d_2\Gamma = \sum_{\Gamma'} c(\Gamma')\Gamma/\Gamma'$:
 - If $i \notin \Gamma'$, then Γ/Γ' still contains a bivalent internal terrestrial vertices connected to two terrestrial vertices.
 - If i is in Γ' but e, e' are not, then $c(\Gamma') = 0$ by Lemma 4.54.
 - The term with $\Gamma' = e$ is cancelled with the terms with $\Gamma' = e'$.
 - If $e \in \Gamma'$ but $e' \notin \Gamma'$ and $\Gamma' \neq e$, then $c(\Gamma') = 0$ by Lemma 4.54. The case $e' \in \Gamma'$ and $e \notin \Gamma'$ is symmetric.
 - Finally if both $e, e' \in \Gamma'$, then $c(\Gamma') = 0$ by Lemma 4.56.

The fact that $d_3\Gamma \in I(U, V)$ follows by similar arguments. □

Definition 4.60. The reduced graph cooperad $\mathbf{cgraphs}_{mn}$ is the relative \mathbf{graphs}_n -cooperad given in each arity by the quotient of $\mathbf{CGraphs}_{mn}(U, V)$ by $I(U, V)$.

Proposition 4.61. *The CDGA $\mathbf{cgraphs}_{mn}(U, V)$ is well-defined, and ω factors through the quotient to define $\mathbf{cgraphs}_{mn}(U, V) \xrightarrow{\omega} \Omega_{\text{PA}}^*(\text{CFM}_{mn}(U, V))$.*

Proof. This follows from Lemmas 4.50, 4.51, 4.54, and 4.56. □

5. Proof of the formality

In this section we complete the proof of the formality of the operad CFM_{mn} . We first show that, up to homotopy, the differential of cgraphs_{mn} can be simplified: the Maurer–Cartan element c is gauge equivalent to a much simpler one c_0 . To prove this, we compute a terrestrially-bound version of the twisted graph complex $(\text{CGC}_{mn}^{\vee}, d + [c_0, -])$ (denoted with a \downarrow superscript). The difference $c - c_0$ belongs to this terrestrially-bound complex. We prove that $\text{CGC}_{mn}^{\vee, c_0, \downarrow}$ quasi-isomorphic to the classical hairy graph complex HGC_{mn}^{\vee} , whose cohomology is known; in particular, it vanishes in the right degree. We can then apply classical deformation-theoretical theorems to prove that $c \simeq c_0$. This enables us to define a simpler version cgraphs_{mn}^0 (replacing c with c_0 in the definition) which is quasi-isomorphic to cgraphs_{mn} . Then we construct a map from cgraphs_{mn}^0 to $H^*(\text{CFM}_{mn})$ by explicit formulas. We prove by combinatorial arguments that this map is a quasi-isomorphism. Since ω is clearly surjective in cohomology, the theorem will follow.

5.1. Change of Maurer–Cartan element and cgraphs_{mn}^0

We would like to define a morphism $\text{cgraphs}_{mn} \rightarrow \text{cd}_{mn}^{\vee}$. However, cgraphs_{mn} depends on the Maurer–Cartan element $c \in \text{CGC}_{mn}^{\vee}$ from Equation (4.30), and we do not know the precise form of c . We just know its leading terms:

Proposition 5.1. *For $n - 2 \geq m \geq 1$, we have $c = c_0 + (\dots)$, where (\dots) denotes terms where $\#\{\text{terr. vert.}\} + 2\#\{\text{aer. vert.}\} > 3$, and:*

$$c_0 := \begin{cases} \begin{array}{c} \bullet \text{---} \bullet + \bullet \text{---} \bullet \\ \bullet \quad \quad \bullet + \bullet \text{---} \bullet \end{array} \in \text{CGC}_{mn}^{\vee}, & \text{if } m \geq 2; \\ \begin{array}{c} \bullet \text{---} \bullet + \bullet \text{---} \bullet \\ \bullet \quad \quad \bullet + \bullet \text{---} \bullet \end{array} \in \text{CGC}_{1n}^{\vee}, & \text{if } m = 1. \end{cases} \quad (5.2)$$

Proof. This follows from Remark 4.55, Remark 4.52, and Remark 4.38 for $m = 1$. Note that c vanishes by definition on the graph with just one internal terrestrial vertex (see after Equation (4.30)). There are no other (Lie-)connected graphs that satisfy $\#\{\text{terr. vert.}\} + 2\#\{\text{aer. vert.}\} \leq 3$, with no loops, double edges, full edges between terrestrial vertices, or bivalent terrestrial vertices connected by dashed edges to other terrestrial vertices (see Lemmas 4.50 and 4.56). \square

If we knew that $c = c_0$, then we would be able to build a map $\text{cgraphs}_{mn} \rightarrow \text{cd}_{mn}^{\vee}$ easily (see Section 5.2). In this section, we show that c is gauge equivalent to c_0 . For this we show that the cohomology of CGC_{mn}^{\vee} twisted by c_0 vanishes in the right degree. Obstruction theory then shows that c is gauge equivalent to c_0 . As we will see, the cohomology of the graph complex

$$\text{CGC}_{mn}^{\vee, c_0} := (\text{CGC}_{mn}^{\vee}, d + [c_0, -]) \quad (5.3)$$

is related to the cohomology of GC_n^{\vee} and the cohomology of the “hairy graph complex” HGC_{mn}^{\vee} that will be defined below (see e.g. [19, Section 2.2.6] or [2]).

Proposition 5.4 ([19, Proposition 2.2.3]). *The cohomology of GC_n^\vee splits as $H^*(\mathrm{GC}_n^{\vee, \geq 3}) \oplus \bigoplus_{l \equiv 2n+1 \pmod{4}} \mathbb{k}\gamma_l$, where $\deg \gamma_l = l - n$ and $\mathrm{GC}_n^{\vee, \geq 3}$ is the Lie subalgebra of graphs whose vertices are all at least trivalent. The class γ_l is represented by the loop with l vertices. Moreover $H^{>-n}(\mathrm{GC}_n^{\vee, \geq 3}) = 0$ for $n \geq 3$.*

Definition 5.5. For $k \geq 0$, let $\mathrm{Graphs}'_n(k)$ be the quotient of $\mathrm{Graphs}_n(k)$ by the ideal spanned by graphs which are disconnected or whose external vertices are not univalent. The hairy graph complex is (with differential from Graphs'_n):

$$\mathrm{HGC}_{mn}^\vee := \prod_{k \geq 1} (\mathrm{Graphs}'_n(k)^\vee \otimes (\mathbb{R}[m])^{\otimes k})^{\Sigma_k}[-m]. \quad (5.6)$$

The full hairy graph complex is the dual of the shifted CDGA $S(\mathrm{HGC}_{mn}[m])[-m]$. The complex HGC_{mn}^\vee is spanned by (infinite sums of) graphs whose external vertices are exactly univalent and indistinguishable. The differential is given by vertex splitting. Each external vertex, together with its only incident edge, can be seen as a “hair”, which justifies the terminology. This hairy graph complex is of great topological interest, as it can e.g. be used to compute spaces of higher-codimensional long knots [2] or the mapping space $\mathrm{Map}(\mathbb{D}_m, \mathbb{D}_n)$ [18].

Proposition 5.7 ([19, Proposition 2.2.7]). *When $n - m \geq 2$, the cohomology of the hairy graph complex HGC_{mn}^\vee vanishes in degrees > -1 .*

Proof. Note that our definition of the hairy graph complex (denoted by HGC_{mn} without the dual in [19]) is slightly different, as we allow bivalent and univalent internal vertices. However, we can reuse their arguments to show that the inclusion of their complex into ours is a quasi-isomorphism (see also [51, Proposition 3.4] for a similar argument). Briefly, we can filter both complexes by the number of internal vertices of valence ≥ 3 . Both spectral sequences collapse starting on page E^2 , and the inclusion induces an isomorphism on this page. We can then use [19, Proposition 2.2.7] to show the vanishing of the homology in degrees > -1 (note that in the reference, homologically graded complexes are used, so we just use the natural correspondence that reverse degrees). \square

For $m \geq 2$, there is a natural preLie product on HGC_{mn}^\vee , induced by the operad structure of Graphs_n^\vee . Roughly speaking, $\Gamma \circ \Gamma'$ is obtained by inserting Γ' in an external vertex of Γ and reconnecting the incident edge to a (non-hair) vertex of Γ' , in all possible ways. Moreover, there is a natural action of the Lie algebra GC_n^\vee (see Definition 4.10) on HGC_{mn}^\vee . Given $\Gamma \in \mathrm{HGC}_{mn}^\vee$ and $\gamma \in \mathrm{GC}_n^\vee$, the action $\Gamma \cdot \gamma$ is given by inserting γ at a vertex of Γ in all possible ways.

When $m = 1$, this simple Lie algebra structure is not right. There is an L_∞ -structure on HGC_{1n}^\vee , called the Shoikhet structure [52]. It is defined by a certain Maurer–Cartan element m_{trans} [43] (in an oriented version of the graph complex GC_n^\vee). The hairy graph complex with this L_∞ -structure is denoted by HGC'_{1n} and encodes the deformation complex of the map $\mathbf{e}_1 = \mathrm{Ass} \rightarrow \mathbf{e}_n$ rather than $\mathrm{Pois} \rightarrow \mathbf{e}_n$ (we refer to [18, Theorem 7.16] where this connection is spelled out in detail). See [52, Section 7] for examples of this L_∞ structure.

Definition 5.8. Let $\text{fCGC}_{mn}^{\vee, c_0, \downarrow}$ be the submodule of $\text{fCGC}_{mn}^{\vee, c_0}$ spanned by graphs whose connected components all have at least one terrestrial vertex and at least one full edge. Let $\text{CGC}_{mn}^{\vee, c_0, \downarrow} \subset \text{fCGC}_{mn}^{\vee, c_0, \downarrow}$ be the submodule of connected graphs. Then clearly:

Lemma 5.9. *The submodule $\text{CGC}_{mn}^{\vee, c_0, \downarrow} \subset \text{CGC}_{mn}^{\vee, c_0}$ is a dg-Lie subalgebra and a Lie GC_n^{\vee} -submodule.* \square

Lemma 5.10. *There is an inclusion of dg-modules $\text{fHGC}_{mn}^{\vee} \subset \text{fCGC}_{mn}^{\vee, c_0, \downarrow}$ obtained by considering all external vertices as terrestrial, with no dashed edges. On the connected parts, for $m \geq 2$ this inclusion is compatible with the Lie structure and the action of the Lie algebra GC_n^{\vee} on both sides. For $m = 1$, it can be extended to an L_{∞} -morphism whose linear part is the inclusion.*

Proof. Simple inspection shows that the inclusion is well-defined, and that it is compatible with the differential. The compatibility with the Lie bracket is clear for $m \geq 2$. The case $m = 1$ follows by adapting the proof of [52, Proposition 5.1], replacing the Hochschild complex of Graphs_n^{\vee} with $\text{CGC}_{1n}^{\vee, c_0, \downarrow}$. \square

Proposition 5.11. *The inclusion $\text{fHGC}_{mn}^{\vee} \subset \text{fCGC}_{mn}^{\vee, c_0, \downarrow}$ is a quasi-isomorphism.*

Proof. The proof is similar to [51, Lemma 4.4]. Indeed, $\text{fCGC}_{mn}^{\vee, c_0, \downarrow}$ is very close to the deformation complex of the morphism $\text{Graphs}_m^{\vee} \rightarrow \text{Graphs}_n^{\vee}$ (denoted by $\text{Def}(\text{hoe}_m \rightarrow \text{Graphs}_n)$ in [51]).

We first filter both complexes by the number of full edges, which is the only kind of edges in fHGC_{mn}^{\vee} . The differential of fHGC_{mn}^{\vee} always increases this number strictly by 1. Let us write $c_0 = c'_0 + c''_0$, where c'_0 is the part with two terrestrial vertices, and c''_0 with one vertex of each kind (see Equation (5.2)). The differential of $\text{fCGC}_{mn}^{\vee, c_0, \downarrow}$ increases the filtration number by 1 (for the action of $\mu + c''_0$) or keeps it constant (for the action of c'_0). Hence on the associated spectral sequences, the differential of $E^0 \text{fHGC}_{mn}^{\vee}$ vanishes, while the differential of $E^0 \text{fCGC}_{mn}^{\vee, c_0, \downarrow}$ is just the bracket $[c'_0, -]$.

We now check that the inclusion induces a quasi-isomorphism on these E^0 pages, from which the proposition follows. Let us first assume that $m \geq 2$. Given $\Gamma \in \text{fCGC}_{mn}^{\vee, c_0, \downarrow}$, define its character $[\Gamma] \in \text{fHGC}_{mn}^{\vee}$ as follow: remove all terrestrial vertices and dashed edges, and call the full edges that used to be connected to terrestrial vertices “dangling”, then make the dangling edges into hairs (see [51, Lemma 4.4] for an analogous definition). The differential $[c'_0, -]$ does not change the character of a graph. Hence $E^0 \text{fCGC}_{mn}^{\vee, c_0, \downarrow}$ splits:

$$E^0 \text{fCGC}_{mn}^{\vee, c_0, \downarrow} = \prod_{\gamma \in \text{fHGC}_{mn}^{\vee}} \underbrace{\{\Gamma \in E^0 \text{fCGC}_{mn}^{\vee, c_0, \downarrow} \mid [\Gamma] = \gamma\}}_{=: C_{\gamma}}. \quad (5.12)$$

Let $\gamma \in \text{fHGC}_{mn}^{\vee}$ be a graph with hairs $\{h_1, \dots, h_k\}$. Let G be the group of permutations of hairs. Then C_{γ} is isomorphic to $C'_{\gamma} = (\text{Graphs}_m^{\vee} \circ \bar{S}^c(H_1, \dots, H_k)^{(1, \dots, 1)})^G$, where $\bar{S}^c(H_1, \dots, H_k)$ is the (non counital) cofree cocommutative coalgebra on variables H_i of degree $-m$, $\text{Graphs}_m^{\vee} \circ -$ is the free Graphs_m^{\vee} -algebra functor, and $(-)^{(1, \dots, 1)}$ is the subcomplex where each H_i appears exactly once. Indeed, we can view $\Xi \in C'_{\gamma}$ as a linear combination of graphs from $\text{Graphs}_m^{\vee}(r)$ with each external vertex decorated by one or

more H_i , with each H_i appearing once. We can identify Ξ with an element of C_γ by making its edges dashed, its vertices terrestrial, and we glue γ to the graph obtained, connecting the hair h_i to the vertex decorated by H_i . The hairs are indistinguishable, but Ξ is invariant under G so this is well-defined. This is illustrated by (with Ξ at the bottom):

$$\begin{array}{c} \begin{array}{c} \boxed{\gamma} \\ \swarrow h_2 \quad \downarrow h_1 \quad \searrow h_3 \\ H_2 \quad H_1, H_3 \end{array} \\ \text{---} \text{---} \text{---} \end{array} \mapsto \begin{array}{c} \boxed{\gamma} \\ \swarrow \quad \downarrow \quad \searrow \\ \text{---} \text{---} \text{---} \end{array} \in C_\gamma. \quad (5.13)$$

For example, full edges between terrestrial vertices may be obtained when γ contains a copy of the “line graph”, i.e. the only connected hairy graph with no internal vertices. The differential $[c'_0, -]$ replicates the differential of $\text{Graphs}_m^\vee(k)$ (i.e. vertex splitting), thus this is an isomorphism of dg-modules.

The homology of Graphs_m^\vee is the m -Poisson operad (Theorem 4.14). Checking the degrees and the induced differential $[\mu + c''_0, -]$, we can identify the page $E^1 \text{fCGC}_{mn}^{\vee, c_0, \downarrow}$ with (a shift of) the deformation complex $\text{Def}(\text{hoe}_m \xrightarrow{*} \text{Graphs}_n^\vee)$ considered in [51]. Note that there, the case $n = m$ is considered and so the map $\text{hoe}_n \rightarrow \text{Graphs}_n^\vee$ sends the Lie bracket to a nonzero element; however, in [51], the part of the differential induced by this element is discarded, so the complex considered is $\text{Def}(\text{hoe}_m \xrightarrow{*} \text{Graphs}_n^\vee)$ up to shifts. Compare also with the results of [2, Section 5], where the full hairy graph complex is called $HH^{m,n}$.

The differential of fHGC_{mn}^\vee raises the number of edges by 1, so the page $E^1 \text{fHGC}_{mn}^\vee$ is just fHGC_{mn}^\vee . Thanks to [51, Lemma 4.4], the inclusion $\text{fHGC}_{mn}^\vee \rightarrow \text{Def}(\text{hoe}_n \xrightarrow{*} \text{Graphs}_n^\vee)$ is a quasi-isomorphism, thus our morphism induces an isomorphism on the E^2 page of the spectral sequence and so it is a quasi-isomorphism itself.

For $m = 1$, the proof is similar, but Lie clusters replace dashed edges. We get that $E^1 \text{fCGC}_{1n}^{\vee, c_0, \downarrow}$ is the (chains) deformation complex of $\text{hoe}_1 \rightarrow \text{Graphs}_n^\vee(k)$, whose homology is the full hairy graph complex fHGC_{1n} . The induced morphism on the E^2 pages is the identity, from which the result follows. \square

Corollary 5.14. *The inclusion $\text{HGC}_{mn}^\vee \subset \text{CGC}_{mn}^{\vee, c_0, \downarrow}$ is a quasi-isomorphism.*

Proof. Both CDGAs fHGC_{mn}^\vee and $\text{fCGC}_{mn}^{\vee, c_0, \downarrow}$ are free as CDGAs, so they are in particular cofibrant. The functor of indecomposables is a left Quillen adjoint [34, Section 12.1.3]. It thus preserves quasi-isomorphisms between cofibrant objects. Since the indecomposables of the two CDGAs mentioned above are respectively HGC_{mn}^\vee and $\text{CGC}_{mn}^{\vee, c_0, \downarrow}$, we conclude by Lemma 5.10 and Proposition 5.11. \square

Corollary 5.15. *The Maurer–Cartan element $c - c_0 \in \text{CGC}_{mn}^{\vee, c_0}$ is gauge equivalent to zero; equivalently, c and c_0 are gauge equivalent.*

Proof. Let $C \subset \text{CGC}_{mn}^{\vee, c_0}$ be the subalgebra spanned by graphs which are not the loops γ_l from Proposition 5.4. Similarly let $C' \subset \text{GC}_n^\vee$ be the subalgebra spanned by graphs which are not the loops. We have a short exact sequence $0 \rightarrow \text{CGC}_{mn}^{\vee, c_0, \downarrow} \rightarrow C \rightarrow C' \rightarrow 0$.

The coefficient $c - c_0$ belongs to the subalgebra C . Indeed, c vanishes on the loops γ_l by degree reasons (and so does c_0). Moreover, we can compute c on purely terrestrial/dashed graphs and show that it agree with c . If $m = 1$, then this follows by immediate degree reasons. For $m \geq 2$, the restriction of c to purely terrestrial/dashed graphs is equal to Kontsevich's coefficient $\mu \in \text{GC}_m^\vee$ ($\text{CFM}_{mn}(U, \emptyset) = \text{FM}_m(U)$ and the integral is identical). So the fact that c and c_0 agree on such graphs follows from the explicit description of μ in Equation (4.6).

We can then combine Propositions 5.4, 5.7 and 5.11 to get that the homology of C vanishes in degrees > -1 . We conclude by applying the Goldman–Millson theorem [22] (see [21, 12] for modern accounts that explicitly deal with MC elements) to the inclusion of the truncation $\tau_{<0} \text{CGC}_{mn}^{\vee, c_0} \subset \text{CGC}_{mn}^{\vee, c_0}$. \square

Definition 5.16. Let CGraphs_{mn}^0 be the variant of CGraphs_{mn} where we use c_0 instead of c to twist the Hopf cooperad CGra_{mn} in the step of Definition 4.33.

Corollary 5.17. *The Hopf cooperads CGraphs_{mn} and CGraphs_{mn}^0 are quasi-isomorphic.*

Proof. This follows from the same general arguments of [7, Section 5.4]. Let us briefly describe them. Let $S(t, dt)$ be the algebra of polynomial forms on the interval $[0, 1]$, with $\deg t = 0$ and $\deg dt = 1$. Let $\text{CGC}_{mn}^{\vee, \sim}$ be the Lie algebra with differential $[\mu, -]$, i.e. we are only allowed to split aerial vertices. Both c and c_0 are Maurer–Cartan elements, i.e. they satisfy $[\mu, c] + \frac{1}{2}[c, c] = [\mu, c_0] + \frac{1}{2}[c_0, c_0] = 0$. The Lie algebra CGC_{mn}^\vee is the twist of $\text{CGC}_{mn}^{\vee, \sim}$ with respect to c .

The gauge equivalence between c and c_0 is a Maurer Cartan element $c_t \in \text{CGC}_{mn}^{\vee, \sim} \otimes S(t, dt)$ whose restriction at $t = 1$ (resp. $t = 0$) is c (resp. c_0). This element c_t produces a differential on $\text{CGraphs}_{mn} \otimes S(t, dt)$ such that restriction at $t = 1$ (resp. $t = 0$) gives CGraphs_{mn} (resp. CGraphs_{mn}^0). We thus have a zigzag:

$$\text{CGraphs}_{mn} \xleftarrow{\text{ev}_{t=1}} \text{CGraphs}_{mn} \otimes S(t, dt) \xrightarrow{\text{ev}_{t=0}} \text{CGraphs}_{mn}^0. \quad (5.18)$$

The evaluation maps $\text{ev}_{t=0}, \text{ev}_{t=1} : S(t, dt) \rightarrow \mathbb{R}$ are quasi-isomorphisms of CDGAs. This implies that the two maps above are quasi-isomorphisms. \square

Definition 5.19. Let cgraphs_{mn}^0 be the quotient of CGraphs_{mn}^0 defined similarly to how cgraphs_{mn} is a quotient of CGraphs_{mn} (see Definition 4.60).

Lemma 5.20. *The quotient cgraphs_{mn}^0 is a relative Hopf graphs $_n$ -cooperad.*

Proof. See Proposition 4.61: c_0 satisfies the same vanishing lemmas as c . \square

Proposition 5.21. *The quotient map $\text{CGraphs}_{mn}^0 \rightarrow \text{cgraphs}_{mn}^0$ is a quasi-isomorphism.*

Proof. This follows from the same arguments as in the proof of [51, Proposition 3.8]. One can set up spectral sequences (counting bivalent vertices of the appropriate type) to see that univalent vertices and bivalent terrestrial vertices with dashed incident edges are killed up to homotopy. Similarly another spectral sequence shows that loops (called tadpoles in [51]) are killed up to homotopy. \square

5.2. Connecting the graphs to the cohomology

The goal of this section is to describe a quasi-isomorphism of Hopf cooperads $\pi : \mathbf{cgraphs}_{mn}^0 \rightarrow \mathbf{cd}_{mn}^\vee$, where $\mathbf{cd}_{mn}^\vee = H^*(\mathbf{CFM}_{mn})$ was obtained in Section 3.1. We will describe this map on generators. The CDGA $\mathbf{cgraphs}_{mn}^0(U, V)$ is free as an algebra for $m \geq 2$. Its generators are the ‘‘internally connected’’ graphs, i.e. the graphs which stay connected when all the external vertices are removed. For example, if Γ has no internal vertices, then it is internally connected iff it has exactly one edge (an empty graph is not connected).

Definition 5.22. If Γ is an internally connected graph, then $\pi(\Gamma) \in \mathbf{cd}_{mn}^\vee(U, V)$ is given by:

- If $\Gamma = e_{vv'}$ has no internal vertices and one full edge between $v \neq v' \in V$, then $\pi(\Gamma) = \omega_{vv'}$.
- If $\Gamma = \tilde{e}_{uu'}$ (for $m \geq 2$) has no internal vertices and one dashed edge between $u \neq u' \in U$, then $\pi(\Gamma) = \tilde{\omega}_{uu'}$.
- If Γ is the graph of (4.53), then $\pi(\Gamma) = \eta_u$.
- In all other cases, $\pi(\Gamma) = 0$.

This is extended to the whole algebra. For $m = 1$, a graph $\Gamma \in \mathbf{cgraphs}_{mn}^0(U, V)$ is additionally decorated (in the dual basis) with an order on $U \sqcup I$ where I is the set of terrestrial internal vertices. The element $\pi(\Gamma) \in \mathbf{cd}_{1n}^\vee(U, V)$ is defined as above. It is decorated with the order on U given by the restriction of the order on $U \sqcup I$, multiplied by the number of ways the internal vertices can be reordered while still keeping the same graph. (This normalization is due to the canonical isomorphism between invariants and coinvariants.)

Proposition 5.23. *The map π defined above is a quasi-isomorphism of Hopf cooperads $\mathbf{cgraphs}_{mn}^0 \rightarrow \mathbf{cd}_{mn}^\vee$.*


The proof of this proposition is split in a series of lemmas, which occupies the rest of this section (until the conclusion, Theorem 5.35).

Lemma 5.24. *The map π is a well-defined algebra map and is equivariant with the symmetric group actions.*

Proof. For $m \geq 2$, we defined π on the generators of a free algebra (forgetting about the differential), so it is well-defined. It is moreover clearly equivariant. For $m = 1$, we need to check the compatibility with the order on the terrestrial vertices. One can directly compute that the coefficients match. Let us first illustrate with an example:

$$\left(\begin{array}{c} \textcircled{1} \quad \textcircled{2} \quad \textcircled{3} \quad \textcircled{4} \quad \textcircled{5} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \text{---} \end{array} \right) \cdot \left(\begin{array}{c} \textcircled{1} \quad \textcircled{2} \quad \textcircled{3} \quad \textcircled{4} \quad \textcircled{5} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \text{---} \end{array} \right) \mapsto \left(\frac{1}{2} \eta_1 \eta_3 \eta_4 \right) \cdot \eta_2 \eta_5,$$

The product on the LHS is given by:



The normalization factor in the formula for π is $\frac{1}{2!3!} = \frac{1}{12}$ which cancels with the 6 terms to give the $\frac{1}{2}$ in the RHS. More generally, to multiply $\Gamma, \Gamma' \in \mathbf{cgraphs}_{mn}^0(U, V)$, if the restriction of the orders on U differ then the result is zero. Otherwise let I, I' be the respective sets of internal vertices and let $k = \#U$. The linear orders on $U \sqcup I$ and $U \sqcup I'$ split I and I' in $(k+1)$ blocks consecutive vertices, of respective sizes i_0, \dots, i_k , and i'_0, \dots, i'_k . The normalization factor in $\pi(\Gamma)$ (resp. $\pi(\Gamma')$) is then $\prod_j (i_j)!^{-1}$ (resp. $\prod_j (i'_j)!^{-1}$). In $\Gamma \cdot \Gamma'$, for each $0 \leq j \leq k$, the i_j vertices of block j in I are shuffled with the i'_j vertices of block j in I' , yielding in total $(i_j + i'_j)! (i_j)!^{-1} (i'_j)!^{-1}$ shuffles. The normalization factor in $\pi(\Gamma \cdot \Gamma')$ is $\prod_j (i_j + i'_j)!^{-1}$ which is equal to the product of the number of shuffles and the normalization factors of Γ and Γ' . \square

Lemma 5.25. *The map π commutes with the differentials, i.e. $\pi d = 0$.*

Proof. Since π is an algebra map and the differential is a derivation, it is sufficient to check this on generators. Let Γ be an internally connected graph. If Γ has no internal vertices, then $d\Gamma = 0$ thus $\pi d\Gamma = 0$. If Γ has one internal vertex, then $\pi d\Gamma = 0$ follows from the Arnold relations and the fact that full edges incident to terrestrial vertices are mapped to zero.

Assume that Γ has at least two internal vertices. If a summand in $d\Gamma$ is nonzero, then after contracting one edge, all remaining edges are between external vertices or between an external aerial vertex and a univalent terrestrial internal one. There can thus be at most one aerial internal vertex. Since contracting an edge cannot reduce the valence of the remaining vertices (contracting dead ends is forbidden), there can only be one internal vertex of valence greater than one, necessarily aerial. Using the internal connectedness of Γ , this special vertex must be connected to all the univalent terrestrial vertices by a full edge. In other words, the graph Γ must be of this type (plus disconnected external vertices):



The Arnold relations in \mathbf{e}_n^\vee , the symmetry relation $\eta_v \omega_{vv'} = \eta_{v'} \omega_{vv'}$, and $\eta_v^2 = 0$ (if there is more than one terrestrial vertex) show that $\pi d\Gamma = 0$. The case $m = 1$ is identical except that everything is multiplied by the number of ways of reordering the internal vertices. \square

Lemma 5.27. *The map π commutes with the cooperad structure maps.*

Proof. It is sufficient to check this on generators, i.e. internally connected graphs, which is completely straightforward but tedious. \square

5.3. Proof that π is a quasi-isomorphism

The last step for Proposition 5.23 is proving that π is a quasi-isomorphism. We split this proof in several sub-lemmas.

- W : the last external vertex is at least bivalent, or univalent and connected by a full edge to an internal vertex.

Let $\mathcal{Q} = \mathbf{cgraphs}_{mn}^0(k+1, 0)_{\text{cn}}/U \cong (V \oplus W, d)$. We filter V by the number of edges, and W by the number of edges minus 1. In the 0th page of the spectral sequence $E^0\mathcal{Q}$, the differential maps V isomorphically onto W , so \mathcal{Q} is acyclic and $U \simeq \mathbf{cgraphs}_{mn}^0(k+1, 0)_{\text{cn}}$. For $k = 0$, $U = \mathbb{R} = \mathbf{e}_m^\vee(1)$ as expected. For $k > 0$, U is isomorphic to $\bigoplus_{i=0}^k \mathbf{cgraphs}_{mn}^0(k, 0)_{\text{cn}}[1-m]$ (by removing the vertex $k+1$ and its incident edge). Hence $\beta^j(k+1, 0) = k \cdot \beta^{j-m+1}(k, 0)$ as expected too. \square

Let us now turn to the second step of the proof. We prove that we can, in some sense, “split” our graph complex in two: external aerial and external terrestrial.

Lemma 5.32. *Let $k, l \geq 1$ and let $I_{k,l} \subset \mathbf{cgraphs}_{mn}^0(k, l)$ be the module spanned by graphs where one of the connected components contains an external aerial vertex and an external terrestrial vertex. Then I is an acyclic dg-ideal.*

Proof. The subspace $I_{k,l}$ is a dg-ideal: contracting edges does not affect connected components, and gluing along external vertices can merge connected components but never split them. Let us prove that it is acyclic. We only deal with connected graphs (the general case follows by the Künneth formula). The proof is similar to Lemma 5.30: we fix $l \geq 1$ and work by induction on $k \geq 1$.

For $k = 1$ we check acyclicity directly. We split $I_{1,l}$ in two submodules (either the external vertex is univalent with a dashed edge, or not) and we filter like in Lemma 5.30 to get a trivial E^1 page. Let us now assume that the claim is true for a given $k \geq 1$. Just like in the proof of Lemma 5.30, we can split $I_{k+1,l}$ in three summands, depending on whether the last external terrestrial vertex is: univalent, connected by a dashed edge to an external vertex; univalent, connected by a dashed edge to an internal vertex; all other cases. The first summand is isomorphic to $\bigoplus_{i=1}^k I_{k,l} \simeq 0$. The quotient by this summand can be filtered like in Lemma 5.30 and is thus also acyclic. \square

Lemma 5.33. *The map $\pi : \mathbf{cgraphs}_{mn}^0(0, l) \rightarrow \mathbf{cd}_{mn}^\vee(0, l)$ is a quasi-isomorphism.*

Proof. This final lemma is also proved by induction. Once again π is clearly surjective on cohomology, so it suffices to prove that both complexes have the same Betti numbers. Using the results of Section 3.1, the Poincaré polynomial of $\mathbf{CFM}_{mn}(0, l) \simeq \mathbf{Conf}_{\mathbb{R}^n \setminus \mathbb{R}^m}(l)$ is $\mathcal{P}(\mathbf{Conf}_{\mathbb{R}^n \setminus \mathbb{R}^m}(l)) = \prod_{i=0}^{l-1} (1 + t^{n-m-1} + it^{n-1})$.

We can again work with the connected part of the graph complex $\mathbf{cgraphs}_{mn}^0(0, l)_{\text{cn}}$. Note that the case $l = 0$ is covered by Lemma 5.30. The base case that we need to prove is $\beta^0(0, 1) = \beta^{n-m-1}(0, 1) = 1$, and $\beta^j(0, 1) = 0$ for other j . The recurrence relation is $\beta^j(0, l+1) = l \cdot \beta^{j-n+1}(0, l)$ for all j and all $l \geq 1$.

For $l = 1$, we split $\mathbf{cgraphs}_{mn}^0(0, 1)_{\text{cn}} = \mathbf{cgraphs}_{mn}^0(0, 1)$ according to the valence of the only external vertex:

- U : the external vertex is zero-valent (i.e. $\Gamma = 1$) or univalent, connected to a univalent internal terrestrial vertex (i.e. $\Gamma = \textcircled{1} \text{---} \bullet$).
- V : the external vertex is at least bivalent.

- V' : the external vertex is univalent, connected to an aerial internal vertex.
- W : the external vertex is univalent, connected to a terrestrial internal vertex; this vertex is itself bivalent and its other incident edge is dashed.
- W' : the external vertex is univalent, connected to a terrestrial internal vertex; this vertex is itself either at least trivalent, or bivalent and both incident edges are full.

Let $\mathcal{Q} = \text{cgraphs}_{mn}^0(0, 1)/U$. We can set up a spectral sequence just like in Lemma 5.30 so that $E^0\mathcal{Q} = \left(V' \xrightarrow[\cong]{d} V \oplus W' \xrightarrow[\cong]{d} W \right)$. Thus \mathcal{Q} is acyclic and $U \simeq \text{cgraphs}_{mn}^0(0, 1)$ is thus a quasi-isomorphism, as we wanted.

For the induction step, we split $\text{cgraphs}_{mn}^0(0, l+1)_{\text{cn}}$ as above. We just replace U by $U = \bigoplus_{i=1}^l U_i$ where U_i is spanned by graphs where the external vertex $(l+1)$ is univalent, connected to the external vertex i . The others are similar but the valence conditions are on the external vertex $(l+1)$ instead. The same argument shows that $U \subset \text{cgraphs}_{mn}^0(0, l+1)_{\text{cn}}$ is a quasi-isomorphism. We have an isomorphism $U_i \cong \text{cgraphs}_{mn}^0(0, l)[1-m]$ given by removing the last external vertex and its incident edge. The Betti numbers thus satisfy the expected recurrence relation: $\beta^j(0, l+1) = l \cdot \beta^{j-n+1}(0, l)$. \square

5.3.2. Case $m = 1$

We deal separately with the case $m = 1$, because e_1 is the associative operad and not the Poisson operad. To summarize the differences, recall that: the graphs do not have dashed edges, and the terrestrial vertices are ordered (Definition 4.16); the notion of “disconnected” is replaced by “Lie-disconnected” (Definition 4.42); the differential $[c_0, -]$ merges coLie clusters (5.2).

Proposition 5.34. *The map $\pi : \text{cgraphs}_{1n}^0(k, l) \rightarrow \text{cd}_{1n}^\vee(k, l)$ is a quasi-isomorphism.*

Proof. As in the case $m = 2$, the map π is clearly surjective on cohomology, so we just need to check that $\text{cgraphs}_{1n}^0(k, l)$ has the correct Betti numbers. The proofs of Lemmas 5.30, 5.32, and 5.33 can be adapted in a straightforward manner. We can follow the same proofs, replacing m with 1. The crucial difference will be in the splitting of the complex $\text{cgraphs}_{1n}^0(k, l)_{\text{cn}}$ or of $I_{k,l}$.

- In $\text{cgraphs}_{1n}^0(k, 0)_{\text{cn}}$ (for Lemma 5.30), we set U to be the submodule where the $(k+1)$ th external vertex is isolated but not adjacent to a terrestrial internal vertex (terrestrial vertices are ordered for $m = 1$), V the submodule where the $(k+1)$ th external vertex is isolated and adjacent to a terrestrial internal vertex, and W all other kinds of graphs.
- In $I_{k,l}$ (for Lemma 5.32), we use the same splitting as for $\text{cgraphs}_{1n}^0(k, 0)_{\text{cn}}$.
- In $\text{cgraphs}_{1n}^0(0, l)$ (for Lemma 5.33), we keep the same U , V , and V' as in the proof of Lemma 5.33. We change the submodules W and W' : in W , we require the last external vertex to be connected to a univalent internal terrestrial vertex, while in W' we put all other graphs.

With this, we obtain the correct recurrence relations on the Betti numbers. \square

5.3.3. Conclusion

Theorem 5.35. *The operad CFM_{mn} is formal over \mathbb{R} for $n - 2 \geq m \geq 1$.*

Proof. We have a zigzag, where cd_{mn}^\vee is defined in Section 3.1, CGraphs_{mn} in Definition 4.47, CGraphs_{mn}^0 in Definition 5.16, $\text{CGraphs}_{mn} \otimes S(t, dt)$ in Corollary 5.17, and cgraphs_{mn}^0 in Definition 5.19, the map π is defined at the beginning of Section 5.2, and the map ω is defined in Proposition 4.61:

$$\begin{aligned} \text{cd}_{mn}^\vee \xleftarrow{\pi} \text{cgraphs}_{mn}^0 \leftarrow \text{CGraphs}_{mn}^0 \leftarrow \\ \leftarrow \text{CGraphs}_{mn} \otimes S(t, dt) \rightarrow \text{CGraphs}_{mn} \xrightarrow{\omega} \Omega_{\text{PA}}^*(\text{CFM}_{mn}), \end{aligned}$$

We proved in Proposition 3.17 that $\text{cd}_{mn}^\vee \cong H^*(\text{CFM}_{mn})$ as Hopf cooperads. We moreover proved in Corollary 5.17 that the two maps involving the three variants of CGraphs_{mn} were quasi-isomorphisms of Hopf cooperads. We also proved that the quotient $\text{CGraphs}_{mn}^0 \rightarrow \text{cgraphs}_{mn}^0$ is a quasi-isomorphism in Proposition 5.21. In addition, we proved that π was a quasi-isomorphism of Hopf cooperads in Proposition 5.23 (for $m \geq 2$) and Proposition 5.34 (for $m = 1$). Therefore it just remains to check that ω is a quasi-isomorphism of Hopf cooperads to conclude.

We already know that CGraphs_{mn} and $\Omega_{\text{PA}}^*(\text{CFM}_{mn})$ have the same cohomology cd_{mn}^\vee . Thus we only need ω to be surjective on cohomology, which is clear (η_v is obtained by graphs of the type seen in Remark 4.52). \square

A. Relative cooperadic twisting

Operadic twisting is a tool originally introduced in [51, Appendix I], studied in further detail in [11], and generalized to certain types of colored operads in [54, Appendix C]. In this appendix, we quickly recall operadic twisting for cooperads and right comodules, and we combine both to deal with relative cooperads.

A.1. Twisting cooperads

General references for twisting of plain operads are [51, Appendix I] and [11]. The dual notion of cooperadic twisting is spelled out in [27, Section 1.5]. Let $\text{Lie}_n = \text{Lie}\{n - 1\}$ be the operad governing Lie algebras with a bracket of homological degree $n - 1$ (so $\text{Lie}_n \subset \mathbf{e}_n$). Let $\text{holie}_n = \Omega(\text{Lie}_n^\dagger) = \Omega(\text{Com}^\vee\{n\})$ be its Koszul resolution. Suppose that \mathbf{C} is a cooperad with finite-dimensional components (so that \mathbf{C}^\vee is an operad) equipped with a morphism $\mu : \text{holie}_n \rightarrow \mathbf{C}^\vee$. We consider the following convolution Lie algebra:

$$\mathfrak{g}_{\mathbf{C}} := \text{hom}_\Sigma(\text{Com}^\vee\{n\}, \mathbf{C}^\vee) = \left(\prod_{i \geq 0} (\mathbf{C}^\vee(i) \otimes \mathbb{R}[-n]^{\otimes i})^{\Sigma_i} [n], d, [-, -] \right).$$

The differential is induced from \mathbf{C} . The Lie bracket of $f, g \in \mathfrak{g}_{\mathbf{C}}$ is $[f, g] = f \star g \mp g \star f$, where \star is the convolution product. Thanks to [34, Theorem 6.5.7], the morphism $\mu : \text{holie}_n \rightarrow \mathbf{C}^\vee$ can equivalently be seen as a Maurer–Cartan element $\mu \in \mathfrak{g}_{\mathbf{C}}$.

The twist of \mathbf{C} with respect to μ is the dg-cooperad given in each arity by:

$$\mathrm{Tw} \mathbf{C}(U) := \bigoplus_{i \geq 0} (\mathbf{C}(U \sqcup \{1, \dots, i\}) \otimes \mathbb{R}[n]^{\otimes i})_{\Sigma_i}.$$

The entries labeled by U are called “external”, whereas the entries that were labeled by $\{1, \dots, i\}$ before taking coinvariants are called “internal”. The cooperadic structure is inherited from \mathbf{C} . Let $\mu_1 \in \prod_{j \geq 0} \mathbf{C}^\vee(\{1, \dots, j, *\})^{\Sigma_j} = \mathrm{Tw} \mathbf{C}^\vee(*)$ (up to shifts and signs) be the element obtained from μ by summing over all possible ways of distinguishing one of the inputs. The differential of $x \in \mathrm{Tw} \mathbf{C}$ is $dx = d_{\mathbf{C}}x + x \cdot \mu - x \cdot \mu_1 - \mu_1 \cdot x$, i.e. the sum of the internal differential of \mathbf{C} with a threefold action of μ that we now describe term by term (see [51] for details): (i) co-insertion of μ in an internal input of x in all possible ways; (ii) co-insertion of $-\mu_1$ in an external input of x in all possible ways; (iii) co-insertion of x in the external input of $-\mu_1$. One checks that $\mu - \mu_1$ defines a Maurer–Cartan element in $\mathfrak{g}_{\mathbf{C}} \times \mathrm{Tw} \mathbf{C}^\vee(*)$, so this differential squares to zero. The compatibility with the cooperad structure is immediate by coassociativity.

A.2. Twisting right comodules

We now recall twisting of right comodules (see [54, Appendix C.1] for the dual case of right modules). Fix $\mu : \mathrm{hoLie}_n \rightarrow \mathbf{C}^\vee$ as in Section A.1. Suppose that \mathbf{M} is a right \mathbf{C} -comodule. Then as a graded module,

$$\mathrm{Tw} \mathbf{M}(U) := \bigoplus_{i \geq 0} (\mathbf{M}(U \sqcup \{1, \dots, i\}) \otimes \mathbb{R}[n]^{\otimes i})_{\Sigma_i}.$$

This inherits a right $(\mathrm{Tw} \mathbf{C})$ -comodule structure from the \mathbf{C} -comodule structure of \mathbf{M} . The differential of $x \in \mathrm{Tw} \mathbf{M}(U)$ is given by $dx = d_{\mathbf{M}}x + x \cdot \mu - x \cdot \mu_1$ (where one uses the comodule structure instead of the cooperad structure). Note that since \mathbf{M} is only a right module, there can be no term of the type $\mu_1 \cdot x$.

A.3. Twisting relative cooperads

Let us finally deal with relative cooperads (see Section 1.1 for the definition). The definition is inspired by the case of “moperads” (i.e. relative operads which can only admit operations with zero or one terrestrial input) [54, Appendix C.3].

Let $\overrightarrow{\mathrm{Lie}}_{mn}$ be the relative Lie_n -operad governing triples $(\mathfrak{g}, \mathfrak{h}, f)$ where \mathfrak{g} is a Lie_m -algebra, \mathfrak{h} is a Lie_n -algebra, and $f : \mathfrak{h}[m-n] \rightarrow \mathfrak{g}$ is a morphism of shifted Lie algebras. We define below an operad hoLie_{mn} over $\overrightarrow{\mathrm{Lie}}_{mn}$. We will not prove that $\overrightarrow{\mathrm{Lie}}_{mn}$ is Koszul, although this seems doable using techniques similar to [25].

Let $\overrightarrow{\mathrm{Com}}_{mn}$ be the operad governing triples (A, B, α) where A is a $\mathrm{Com}\{n\}$ -algebra, B is a $\mathrm{Com}\{m\}$ -algebra, and $\alpha : A \rightarrow B[n-m]$ is a morphism of shifted commutative algebras. In particular, $\overrightarrow{\mathrm{Com}}_{mn}(U, V) \cong \mathbb{R}[-m]^{\otimes U} \otimes \mathbb{R}[-n]^{\otimes V} \otimes \mathbb{R}[m]$ is one-dimensional for all pairs $(U, V) \neq (\emptyset, \emptyset)$ (and $\overrightarrow{\mathrm{Com}}_{mn}(\emptyset, \emptyset) = 0$). By definition, $\overrightarrow{\mathrm{Com}}_{mn}$ is a relative $\mathrm{Com}\{n\}$ -operad. Thus the cobar construction $\mathrm{hoLie}_{mn} := \Omega(\overrightarrow{\mathrm{Com}}_{mn}^\vee)$ is a relative hoLie_n -operad (since $\mathrm{hoLie}_n = \Omega(\mathrm{Com}_n^\vee)$).

Let D be a relative C -cooperad equipped with a map $(c, \mu) : (\overrightarrow{\text{hoLie}}_{mn}, \text{hoLie}_n) \rightarrow (D, C)$. This map can equivalently be seen as a Maurer–Cartan element in $\mathfrak{g}_C \oplus \mathfrak{g}_{C,D}$ (i.e. $d\mu + \frac{1}{2}[\mu, \mu] = dc + [\mu, c] + \frac{1}{2}[c, c] = 0$), where $\mathfrak{g}_{C,D} = \text{hom}_\Sigma(\overrightarrow{\text{Com}}_{mn}^V, D)$.

Let us define the twisted relative (Tw C)-cooperad $\text{Tw } D$. As a graded module,

$$\text{Tw } D(U, V) := \bigoplus_{i, j \geq 0} (D(U \sqcup \{1, \dots, i\}, V \sqcup \{1, \dots, j\}) \otimes \mathbb{R}[m]^{\otimes i} \otimes \mathbb{R}[n]^{\otimes j})_{\Sigma_i \times \Sigma_j}.$$

The relative (Tw C)-cooperad structure is inherited from D . Let c_1 be the element obtained from c by summing over all possible ways of distinguishing one of the terrestrial inputs of c (similarly to how μ_1 is defined from μ). Then the differential of $x \in \text{Tw } D(U, V)$ is given by $d_D x + x \cdot \mu + x \cdot (c - c_1) - c_1 \cdot x$.


Proposition A.1. *The collection $\text{Tw } D$ defines a relative (Tw C)-cooperad.*

Proof. Generalizing the proofs of [51, Appendix I] and [54, Appendix C.3] to this setting is straightforward. One checks that $\mu + c + c_1$ defines a Maurer–Cartan element in $\mathfrak{g}_C \rtimes \mathfrak{g}_{C,D} \rtimes \text{Tw } D^V(*, \emptyset)$ – which acts by cooperadic coderivations on $\text{Tw } D$ – so the differential above squares to zero. Compatibility with the cooperad structure follows from the coassociativity of the cooperad structure. \square

B. Compactifications and projections

In this appendix, we sketch a proof of Proposition 2.9: $\text{CFM}_{mn}(U, V)$ is a compact SA manifold and a smooth manifold with corners, and the canonical projection maps are SA bundles. Our proofs are heavily inspired by [32, Section 5.9].

Let (U, V) be a pair of finite sets. A relative (rooted) tree \mathcal{T} with leaves (U, V) is a rooted tree with dashed and full edges. We require that the leaves with incident full (resp. dashed) edge are in bijection with U (resp. V), that if a vertex has only one incoming edge then this edge is full, and that if an edge is full then all the edges above it are full. An example is on the side.



For a relative tree \mathcal{T} , we let $V_{\mathcal{T}}$ be the set of all its vertices, $V_{\mathcal{T}}^0 = V_{\mathcal{T}} \setminus \text{root}$, and $V_{\mathcal{T}}^* = V_{\mathcal{T}} \setminus (U \cup V)$. The set $V_{\mathcal{T}}$ is partially ordered by considering that a vertex is smaller than any vertex above it. For $i \in V_{\mathcal{T}}$, we let $\text{in}(i) = \text{in}_t(i) \cup \text{in}_a(i) = \{\text{incoming dashed edges}\} \cup \{\text{incoming full edges}\}$ and $\text{par}(i) \in V_{\mathcal{T}}$ be the immediate predecessor of i . Finally, we let:

$$\underline{\text{Conf}}_{mn}^{\mathcal{T}} := \prod_{i \in V_{\mathcal{T}}^*} \underline{\text{Conf}}_{mn}(\text{in}_t(i), \text{in}_a(i)).$$

The spaces $\underline{\text{Conf}}_{mn}^{\mathcal{T}}$ will be used to give a decomposition of $\text{CFM}_{mn}(U, V)$ as in [32, Section 5.9.1]. Let $\xi = (\xi_i)_{i \in V_{\mathcal{T}}^*} \in \underline{\text{Conf}}_{mn}^{\mathcal{T}}$. We can represent ξ_i by a configuration $\bar{\xi}_i \in \text{Conf}_{mn}(\text{in}_t(v), \text{in}_a(i))$ of radius 1 and whose barycenter is in $\{0\}^m \times \mathbb{R}^{n-m}$. For $i \in V_{\mathcal{T}}^0$, we let $\xi(i) := \bar{\xi}_{\text{par}(i)}(i)$. We then define, for $r > 0$ and $i \in V_{\mathcal{T}}$:

$$x(\xi, r, i) := \sum_{j \in V_{\mathcal{T}}^0, j \leq i} \xi(j) \cdot r^{\text{height}(j)} \in \mathbb{R}^n.$$

Then $(x(\xi, r, i))_{i \in U \sqcup V}$ is a configuration for r small enough. Let us define $h_{\mathcal{T}} : \underline{\text{Conf}}_{mn}^{\mathcal{T}} \hookrightarrow \text{CFM}_{mn}(U, V)$ by $h_{\mathcal{T}}(\xi) = \lim_{r \rightarrow 0} (x(\xi, r, i))_{i \in U \sqcup V}$. The map $h_{\mathcal{T}}$ is a homeomorphism onto its image, $\{\text{im}(h_{\mathcal{T}})\}_{\mathcal{T}}$ covers $\text{CFM}_{mn}(U, V)$, and the interior of $\text{CFM}_{mn}(U, V)$ is the stratum corresponding to a corolla.

Now let $x = h_{\mathcal{T}}(\xi) \in \text{CFM}_{mn}(U, V)$. We want to build an SA chart around x . Let

$$r_1 := \frac{1}{4} \min\{\|\bar{\xi}_i(a) - \bar{\xi}_i(b)\| \mid i \in V_{\mathcal{T}}^0, a \neq b \in \text{in}(i)\} \cup \{d(\bar{\xi}_i(v), \mathbb{R}^m) \mid i \in V_{\mathcal{T}}^0, v \in \text{in}_a(i)\}.$$

Note that $r_1 \leq \frac{1}{2}$, because $\bar{\xi}_i$ has radius 1. Define a neighborhood of ξ by $W = \{\zeta \in \underline{\text{Conf}}_{mn}^{\mathcal{T}} \mid \forall i, \|\xi(i) - \zeta(i)\| \leq r_1^{\#U + \#V + 1}\}$ (thanks to the distance condition, distinct points stay distinct and aerial points stay aerial). For $\tau \in [0, r_1]^{V_{\mathcal{T}}^*}$ with $\tau_{\text{root}} = 0$ and $0 \leq r \leq r_1$, we let $y(\zeta, \tau, r, \text{root}) := 0$ and

$$y(\zeta, \tau, r, i) := y(\zeta, \tau, r, \text{par}(i)) + \xi(i) \cdot \prod_{j < i} \max(r, \tau_j).$$

We can then define $\Phi : W \times [0, r_1]^{V_{\mathcal{T}}^* \setminus \{\text{root}\}} \rightarrow \text{CFM}_{mn}(U, V)$ by $\Phi(\zeta, \tau) := \lim_{r \rightarrow 0} (y(\zeta, \tau, r, i))_{i \in U \sqcup V}$. We also let V be the image of Φ . The proof that Φ is an SA chart onto a compact neighborhood of x is identical to [32, Lemma 5.9.3]. This proves the first part of Proposition 2.9.

We would now like to prove that $p_{U, V} : \text{CFM}_{mn}(U \sqcup I, V \sqcup J) \rightarrow \text{CFM}_{mn}(U, V)$ is an SA bundle. Since the composite of two SA bundles is an SA bundle [24, Proposition 8.5], it is sufficient to check that the following are SA bundles:

$$p : \text{CFM}_{mn}(U \sqcup *, V) \rightarrow \text{CFM}_{mn}(U, V), \quad q : \text{CFM}_{mn}(U, V \sqcup *) \rightarrow \text{CFM}_{mn}(U, V).$$

We will describe the fibers explicitly as complements of open balls. The fiber of p will be almost identical to the one in [32, Section 5.9.4]. However the fiber of q is slightly different, because the new aerial point cannot touch the ground.

Let $x = h_{\mathcal{T}}(\xi)$, $r_1 > 0$, and W as before. For $\zeta \in W$ and $i \in V_{\mathcal{T}}$, define $x_1(\zeta, i) := x(\zeta, r_1, i)$ and $\varepsilon(i) := 4r_1^{\text{height}(i)+1}$. Let $B_i(\zeta) := B(x_1(\zeta, i), \varepsilon(i))$ be the closed ball. Then $B_i(\zeta) \subset 1/3B_j(\zeta)$ if $i < j$ and $B_i(\zeta) \cap B_j(\zeta) = \emptyset$ otherwise.

Recall $\phi : \mathbb{R}^n \times [0, 1] \times [0, 2] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$, $(c, r, \varepsilon, x) \mapsto \phi_r^{c, \varepsilon}(x)$ from [32, Lemma 5.9.5]. It is such that $\phi_r^{c, \varepsilon}$ is radial, the identity outside $B(c, \varepsilon)$, and shrinks $B(c, \varepsilon/3)$ by a factor r . Moreover for a configuration $x \in \text{Conf}_{B(c, \varepsilon/3)}(k)$, $\phi_r^{c, \varepsilon}(x)$ is a configuration in $\underline{\text{Conf}}_n(k)$ that does not depend on r , and ϕ behaves well with respect to other points $z \in B(c, \varepsilon)$ (see the reference for details). We note in addition that, thanks to the properties of ϕ , if $c \in \mathbb{R}^m$ then $\phi_r^{c, \varepsilon}(\mathbb{R}^m) \subset \mathbb{R}^m$, and if $c \notin \mathbb{R}^m$, then $\phi_r^{c, \varepsilon}(\mathbb{R}^n \setminus B(\mathbb{R}^m, \varepsilon)) \subset \mathbb{R}^n \setminus B(\mathbb{R}^m, \varepsilon)$ (where $B(\mathbb{R}^m, \varepsilon) = \bigcup_{z \in \mathbb{R}^m} B(z, \varepsilon)$).

Now, fix $\zeta \in W$ and $\tau \in [0, r_1]^{V_{\mathcal{T}}}$ s.t. $\tau_{\text{root}} = 0$ and $\tau_i = 0$ for $i \in U \sqcup V$ a leaf. Then for $i \in V_{\mathcal{T}}$ and $0 < r \leq r_1$, let $\phi_r^i = \phi_{\max(r, \tau(i))/r_1}^{x_1(\zeta, i), \varepsilon(i)}$. Moreover, let ϕ_r be the composition (in any order thanks to the disjointness of the balls) of the ϕ_r^i for $i \in U \sqcup V$. (Despite the notation, ϕ_r depends on x, ζ and τ .) We then check, like in [32, Lemma 5.9.6], that $\phi_r(x_1(\zeta, i)) = x(\zeta, \tau, r, i)$ for $r > 0$ and $i \in U \sqcup V$.

We can now check the local trivialities of p and q . Let us first deal with $p : \mathbf{CFM}_{mn}(U \sqcup *, V) \rightarrow \mathbf{CFM}_{mn}(U, V)$. We define $F_\zeta := B(x_1(\zeta, \text{root}), \varepsilon(\text{root})/2) \setminus \bigcup_{u \in U} B(x_1(\zeta, u), \varepsilon(u)/2) \cap \mathbb{R}^m$ for $\zeta \in W$, which will be the fiber of p over $h_{\mathcal{T}}(\zeta)$. We also set $F := B(0, \#U + 1) \setminus \bigcup_{i=1}^{\#U} \mathring{B}((i, 0, \dots, 0), 1/4) \cap \mathbb{R}^m$. Note that F is a compact SA manifold (a closed m -ball with U open balls removed). There is an SA-homeomorphism $\Theta_\zeta : F \cong F_\zeta$ since W is small enough. Let

$$\begin{aligned} \widehat{\Phi} : W \times [0, 1]^{V_{\mathcal{T}}^* \setminus \text{root}} \times F &\rightarrow \mathbf{CFM}_{mn}(U \sqcup *, V), \\ (\zeta, \tau, z) &\mapsto \lim_{r \rightarrow 0} ((\phi_r(x_1(\zeta, i)))_{i \in U \sqcup V}, \phi_r(\Theta_\zeta(z))). \end{aligned}$$

Then $\widehat{\Phi}$ covers Φ . The proof that $\widehat{\Phi}(\zeta, \tau, -)$ maps F_ζ homeomorphically onto $p^{-1}(\Phi(\zeta, \tau))$ is identical to the proof in [32, Section 5.9.4].

Now let us deal with $q : \mathbf{CFM}_{mn}(U, V \sqcup *) \rightarrow \mathbf{CFM}_{mn}(U, V)$. Its fiber over $h_{\mathcal{T}}(\zeta)$ will be $G_\zeta := B(x_1(\zeta, \text{root}), \varepsilon(\text{root})/2) \setminus \bigcup_{v \in V} \mathring{B}(x_1(\zeta, v), \varepsilon(v)/2) \setminus \mathring{B}(\mathbb{R}^m, r_1)$. We also have $G := B(0, \#V + 1) \setminus \mathring{B}(\mathbb{R}^m, 1/4) \setminus \bigcup_{i=1}^{\#V} \mathring{B}((i, 0, \dots, 0, 1, 0, \dots, 0), 1/4)$ (where the 1 in the open ball is in position $m + 1$). This is a compact SA manifold (a closed n -ball with an open tubular neighborhood of \mathbb{R}^m and V open n -balls removed) and we have an SA-homeomorphism $\Theta_\zeta : G \cong G_\zeta$ since, again, W is small enough. We can then define a chart $\widehat{\Phi} : W \times [0, 1]^{V_{\mathcal{T}}^* \setminus \text{root}} \times G \rightarrow \mathbf{CFM}_{mn}(U \sqcup *, V)$ with a formula similar to the one above. This map covers Φ , and showing that $\widehat{\Phi}(\zeta, \tau, -)$ maps G_ζ SA-homeomorphically to $q^{-1}(\Phi(\zeta, \tau))$ is a straightforward adaption of the arguments in [32, Section 5.9.4]. This completes the proof of the second part of Proposition 2.9.

References

- [1] Vladimir I. Arnold. “The cohomology ring of the colored braid group”. In: *Math. Notes* 5.2 (1969), pp. 138–140. ISSN: 0025-567X. DOI: [10.1007/BF01098313](https://doi.org/10.1007/BF01098313).
- [2] Gregory Arone and Victor Turchin. “Graph-complexes computing the rational homotopy of high dimensional analogues of spaces of long knots”. In: *Ann. Inst. Fourier* 65.1 (2015), pp. 1–62. DOI: [10.5802/aif.2924](https://doi.org/10.5802/aif.2924). arXiv: [1108.1001](https://arxiv.org/abs/1108.1001).
- [3] Scott Axelrod and I. M. Singer. “Chern–Simons perturbation theory. II”. In: *J. Differential Geom.* 39.1 (1994), pp. 173–213. DOI: [10.4310/jdg/1214454681](https://doi.org/10.4310/jdg/1214454681). arXiv: [hep-th/9304087](https://arxiv.org/abs/hep-th/9304087).
- [4] David Ayala, John Francis, and Hiro Lee Tanaka. “Factorization homology of stratified spaces”. In: *Selecta Math. New ser.* 23.1 (2017), pp. 293–362. ISSN: 1022-1824. DOI: [10.1007/s00029-016-0242-1](https://doi.org/10.1007/s00029-016-0242-1). arXiv: [1409.0848v3](https://arxiv.org/abs/1409.0848v3).
- [5] J. Michael Boardman and Rainer M. Vogt. “Homotopy-everything H -spaces”. In: *Bull. Amer. Math. Soc.* 74 (1968), pp. 1117–1122. DOI: [10.1090/S0002-9904-1968-12070-1](https://doi.org/10.1090/S0002-9904-1968-12070-1).
- [6] Ricardo Campos, Julien Ducoulombier, Najib Idrissi, and Thomas Willwacher. *A model for framed configuration spaces of points*. 2018. arXiv: [1807.08319](https://arxiv.org/abs/1807.08319). Submitted.

- [7] Ricardo Campos, Najib Idrissi, Pascal Lambrechts, and Thomas Willwacher. *Configuration Spaces of Manifolds with Boundary*. 2018. arXiv: [1802.00716](#).
- [8] Ricardo Campos and Thomas Willwacher. *A model for configuration spaces of points*. 2016. arXiv: [1604.02043](#).
- [9] Frederick R. Cohen. “The homology of \mathcal{C}_{n+1} spaces, $n \geq 0$ ”. In: Frederick R. Cohen, Thomas J. Lada, and J. Peter May. *The homology of iterated loop spaces*. Lecture Notes in Mathematics 533. Berlin Heidelberg: Springer, 1976. Chap. 3, pp. 207–351. ISBN: 978-3-540-07984-2. DOI: [10.1007/BFb0080467](#).
- [10] Corrado De Concini and Claudio Procesi. “Wonderful models of subspace arrangements”. In: *Selecta Math. New ser.* 1.3 (1995), pp. 459–494. ISSN: 1022-1824. DOI: [10.1007/BF01589496](#).
- [11] Vasily Dolgushev and Thomas Willwacher. “Operadic twisting. With an application to Deligne’s conjecture”. In: *J. Pure Appl. Algebra* 219.5 (2015), pp. 1349–1428. DOI: [10.1016/j.jpaa.2014.06.010](#).
- [12] Vasily A. Dolgushev and Christopher L. Rogers. “A version of the Goldman–Millson theorem for filtered L_∞ -algebras”. In: *J. Algebra* 430 (2015), pp. 260–302. ISSN: 0021-8693. DOI: [10.1016/j.jalgebra.2015.01.032](#).
- [13] Julien Ducoulombier. “Swiss-cheese action on the totalization of operads under the monoid actions actions operad”. In: *Algebr. Geom. Topol.* 16.3 (2016), pp. 1683–1726. DOI: [10.2140/agt.2016.16.1683](#). arXiv: [1410.3236](#).
- [14] Benoit Fresse. “Théorie des opérades de Koszul et homologie des algèbres de Poisson”. In: *Ann. Math. Blaise Pascal* 13.2 (2006), pp. 237–312. DOI: [10.5802/ambp.219](#).
- [15] Benoit Fresse. *Homotopy of Operads and Grothendieck–Teichmüller Groups*. Mathematical Surveys and Monographs 217. Providence, RI: Amer. Math. Soc., 2017, pp. xlvii+532. ISBN: 978-1-4704-3481-6.
- [16] Benoit Fresse. “The extended rational homotopy theory of operads”. In: *Georgian Math. J.* 25.4 (2018), pp. 493–512. DOI: [10.1515/gmj-2018-0061](#). arXiv: [1805.00530](#).
- [17] Benoit Fresse. “Little discs operads, graph complexes and Grothendieck–Teichmüller groups”. In: *Handbook of Homotopy Theory*. Ed. by Haynes R. Miller. CRC Press/Chapman Hall Handb. Math. Ser. Boca Raton, FL: Chapman and Hall/CRC, 2019. Chap. 11. ISBN: 978-0-8153-6970-7. arXiv: [1811.12536](#).
- [18] Benoit Fresse, Victor Turchin, and Thomas Willwacher. *The rational homotopy of mapping spaces of E_n operads*. 2017. arXiv: [1703.06123](#).
- [19] Benoit Fresse and Thomas Willwacher. “The intrinsic formality of E_n -operads”. In: *J. Eur. Math. Soc.* 22.7 (2020), pp. 2047–2133. DOI: [10.4171/JEMS/961](#). arXiv: [1503.08699](#).
- [20] William Fulton and Robert MacPherson. “A compactification of configuration spaces”. In: *Ann. of Math. (2)* 139.1 (1994), pp. 183–225. DOI: [10.2307/2946631](#).

- [21] Ezra Getzler. “Lie theory for nilpotent L_∞ -algebras”. In: *Ann. of Math.* 2nd ser. 170.1 (2009), pp. 271–301. ISSN: 0003-486X. DOI: [10.4007/annals.2009.170.271](https://doi.org/10.4007/annals.2009.170.271).
- [22] William M. Goldman and John J. Millson. “The deformation theory of representations of fundamental groups of compact Kähler manifolds”. In: *Publ. Math. Inst. Hautes Études Sci.* 67 (1988), pp. 43–96. ISSN: 0073-8301. URL: http://www.numdam.org/item?id=PMIHES_1988__67__43_0.
- [23] Mark Goresky and Robert MacPherson. *Stratified Morse theory*. Ergebnisse der Mathematik und ihrer Grenzgebiete (3) 14. Berlin: Springer, 1988. 272 pp. ISBN: 3-540-17300-5. DOI: [10.1007/978-3-642-71714-7](https://doi.org/10.1007/978-3-642-71714-7).
- [24] Robert Hardt, Pascal Lambrechts, Victor Turchin, and Ismar Volić. “Real homotopy theory of semi-algebraic sets”. In: *Algebr. Geom. Topol.* 11.5 (2011), pp. 2477–2545. DOI: [10.2140/agt.2011.11.2477](https://doi.org/10.2140/agt.2011.11.2477). arXiv: [0806.0476](https://arxiv.org/abs/0806.0476).
- [25] Eduardo Hoefel and Muriel Livernet. “On the spectral sequence of the Swiss-cheese operad”. In: *Algebr. Geom. Topol.* 13.4 (2013), pp. 2039–2060. DOI: [10.2140/agt.2013.13.2039](https://doi.org/10.2140/agt.2013.13.2039). arXiv: [1209.6122](https://arxiv.org/abs/1209.6122).
- [26] Eduardo Hoefel, Muriel Livernet, and Jim Stasheff. “ A_∞ -actions and recognition of relative loop spaces”. In: *Topology Appl.* 206 (2016), pp. 126–147. DOI: [10.1016/j.topol.2016.03.023](https://doi.org/10.1016/j.topol.2016.03.023). arXiv: [1312.7155](https://arxiv.org/abs/1312.7155).
- [27] Najib Idrissi. “The Lambrechts–Stanley Model of Configuration Spaces”. In: *Invent. Math.* 216.1 (2019), pp. 1–68. DOI: [10.1007/s00222-018-0842-9](https://doi.org/10.1007/s00222-018-0842-9). arXiv: [1608.08054](https://arxiv.org/abs/1608.08054).
- [28] Anton Khoroshkin and Thomas Willwacher. *Real models for the framed little n -disks operads*. 2017. arXiv: [1705.08108](https://arxiv.org/abs/1705.08108).
- [29] Maxim Kontsevich. “Feynman diagrams and low-dimensional topology”. In: *First European Congress of Mathematics*. Vol. 2: *Invited lectures* (Paris, July 6–10, 1992). Progr. Math. 120. Basel: Birkhäuser, 1994, pp. 97–121. DOI: [10.1007/978-3-0348-9112-7_5](https://doi.org/10.1007/978-3-0348-9112-7_5).
- [30] Maxim Kontsevich. “Operads and motives in deformation quantization”. In: *Lett. Math. Phys.* 48.1 (1999), pp. 35–72. DOI: [10.1023/A:1007555725247](https://doi.org/10.1023/A:1007555725247). arXiv: [math/9904055](https://arxiv.org/abs/math/9904055).
- [31] Maxim Kontsevich. “Deformation quantization of Poisson manifolds”. In: *Lett. Math. Phys.* 66.3 (2003), pp. 157–216. DOI: [10.1023/B:MATH.0000027508.00421.bf](https://doi.org/10.1023/B:MATH.0000027508.00421.bf). arXiv: [q-alg/9709040](https://arxiv.org/abs/q-alg/9709040).
- [32] Pascal Lambrechts and Ismar Volić. “Formality of the little N -disks operad”. In: *Mem. Amer. Math. Soc.* 230.1079 (2014), pp. viii+116. DOI: [10.1090/memo/1079](https://doi.org/10.1090/memo/1079). arXiv: [0808.0457](https://arxiv.org/abs/0808.0457).
- [33] Muriel Livernet. “Non-formality of the Swiss-cheese operad”. In: *J. Topol.* 8.4 (2015), pp. 1156–1166. DOI: [10.1112/jtopol/jtv018](https://doi.org/10.1112/jtopol/jtv018). arXiv: [1404.2484](https://arxiv.org/abs/1404.2484).

- [34] Jean-Louis Loday and Bruno Vallette. *Algebraic operads*. Grundlehren der mathematischen Wissenschaften 346. Berlin-Heidelberg: Springer, 2012. 634 pp. ISBN: 978-3-642-30361-6. DOI: [10.1007/978-3-642-30362-3](https://doi.org/10.1007/978-3-642-30362-3).
- [35] Jacob Lurie. *Higher Algebra*. Sept. 2017. URL: <http://www.math.harvard.edu/~lurie/papers/HA.pdf>.
- [36] Martin Markl. “A compactification of the real configuration space as an operadic completion”. In: *J. Algebra* 215.1 (1999), pp. 185–204. DOI: [10.1006/jabr.1998.7709](https://doi.org/10.1006/jabr.1998.7709). arXiv: [hep-th/9608067](https://arxiv.org/abs/hep-th/9608067).
- [37] J. Peter May. *The geometry of iterated loop spaces*. Lectures Notes in Mathematics 271. Berlin: Springer-Verlag, 1972, pp. viii+175. DOI: [10.1007/BFb0067491](https://doi.org/10.1007/BFb0067491).
- [38] Dan Petersen. “Minimal models, GT-action and formality of the little disk operad”. In: *Selecta Math.* New ser. 20.3 (2014), pp. 817–822. ISSN: 1022-1824. DOI: [10.1007/s00029-013-0135-5](https://doi.org/10.1007/s00029-013-0135-5). arXiv: [1303.1448](https://arxiv.org/abs/1303.1448).
- [39] Sylvain Poirier. “The configuration space integral for links in \mathbb{R}^3 ”. In: *Algebr. Geom. Topol.* 2 (2002), pp. 1001–1050. ISSN: 1472-2747. DOI: [10.2140/agt.2002.2.1001](https://doi.org/10.2140/agt.2002.2.1001). arXiv: [math/0005085](https://arxiv.org/abs/math/0005085).
- [40] Alexander Polishchuk and Leonid Positsel’ski. *Quadratic algebras*. University Lecture Series 37. Providence, RI: Amer. Math. Soc., 2005. 159 pp. ISBN: 0-8218-3834-2.
- [41] Alexandre Quesney. *Swiss Cheese type operads and models for relative loop spaces*. 2015. arXiv: [1511.05826](https://arxiv.org/abs/1511.05826).
- [42] Paolo Salvatore. “Configuration spaces with summable labels”. In: *Cohomological methods in homotopy theory*. Barcelona conference on algebraic topology (Bellaterra, Spain, June 5–10, 1998). Ed. by Jaume Aguadé, Carles Broto, and Carles Casacuberta. Progr. Math. 196. Basel: Birkhäuser, 2001, pp. 375–395. ISBN: 978-3-0348-9513-2. DOI: [10.1007/978-3-0348-8312-2_23](https://doi.org/10.1007/978-3-0348-8312-2_23). arXiv: [math/9907073](https://arxiv.org/abs/math/9907073).
- [43] Boris Shoikhet. “An L_∞ algebra structure on polyvector fields”. In: *Selecta Math.* New ser. 24.2 (2018), pp. 1691–1728. ISSN: 1022-1824. DOI: [10.1007/s00029-017-0382-y](https://doi.org/10.1007/s00029-017-0382-y). arXiv: [0805.3363](https://arxiv.org/abs/0805.3363).
- [44] Dev Sinha and Benjamin Walter. “Lie coalgebras and rational homotopy theory I. Graph coalgebras”. In: *Homology Homotopy Appl.* 13.2 (2011), pp. 263–292. ISSN: 1532-0073. DOI: [10.4310/HHA.2011.v13.n2.a16](https://doi.org/10.4310/HHA.2011.v13.n2.a16). arXiv: [math/0610437](https://arxiv.org/abs/math/0610437).
- [45] Dev P. Sinha. “Manifold-theoretic compactifications of configuration spaces”. In: *Selecta Math.* New ser. 10.3 (2004), pp. 391–428. DOI: [10.1007/s00029-004-0381-7](https://doi.org/10.1007/s00029-004-0381-7). arXiv: [math/0306385](https://arxiv.org/abs/math/0306385).
- [46] Dev P. Sinha. “The (non-equivariant) homology of the little disks operad”. In: *Proceedings of the school and conference. Operads 2009* (Luminy, Apr. 20–30, 2009). Ed. by Jean-Louis Loday and Bruno Vallette. Sémin. Congr. 26. Paris: Soc. Math. France, 2013, pp. 253–279. ISBN: 978-2-85629-363-8. arXiv: [math/0610236](https://arxiv.org/abs/math/0610236).
- [47] Dmitry E. Tamarkin. “Formality of chain operad of little discs”. In: *Lett. Math. Phys.* 66.1-2 (2003), pp. 65–72. DOI: [10.1023/B:MATH.0000017651.12703.a1](https://doi.org/10.1023/B:MATH.0000017651.12703.a1).

- [48] Victor Turchin and Thomas Willwacher. “Relative (non-)formality of the little cubes operads and the algebraic Cerf Lemma”. In: *Am. J. Math.* 140.2 (2018), pp. 277–316. DOI: [10.1353/ajm.2018.0006](https://doi.org/10.1353/ajm.2018.0006). arXiv: [1409.0163](https://arxiv.org/abs/1409.0163).
- [49] Renato Vasconcello Vieira. “Relative Recognition Principle”. In: *Algebr. Geom. Topol.* 20.3 (2020), pp. 1431–1486. DOI: [10.2140/agt.2020.20.1431](https://doi.org/10.2140/agt.2020.20.1431). arXiv: [1802.01530](https://arxiv.org/abs/1802.01530).
- [50] Alexander A. Voronov. “The Swiss-cheese operad”. In: *Homotopy invariant algebraic structures* (Baltimore, MD, 1998). Contemp. Math. 239. Providence, RI: American Mathematical Society, 1999, pp. 365–373. DOI: [10.1090/conm/239/03610](https://doi.org/10.1090/conm/239/03610). arXiv: [math/9807037](https://arxiv.org/abs/math/9807037).
- [51] Thomas Willwacher. “M. Kontsevich’s graph complex and the Grothendieck–Teichmüller Lie algebra”. In: *Invent. Math.* 200.3 (2014), pp. 671–760. DOI: [10.1007/s00222-014-0528-x](https://doi.org/10.1007/s00222-014-0528-x). arXiv: [1009.1654](https://arxiv.org/abs/1009.1654).
- [52] Thomas Willwacher. *Deformation quantization and the Gerstenhaber structure on the homology of knot spaces*. 2015. arXiv: [1506.07078](https://arxiv.org/abs/1506.07078).
- [53] Thomas Willwacher. *Models for the n-Swiss Cheese operads*. 2015. arXiv: [1506.07021](https://arxiv.org/abs/1506.07021).
- [54] Thomas Willwacher. “The Homotopy Braces Formality Morphism”. In: *Duke Math. J.* 165.10 (2016), pp. 1815–1964. DOI: [10.1215/00127094-3450644](https://doi.org/10.1215/00127094-3450644). arXiv: [1109.3520](https://arxiv.org/abs/1109.3520).
- [55] Thomas Willwacher. *(Non-)formality of the extended Swiss Cheese operads*. June 2017. arXiv: [1706.02945](https://arxiv.org/abs/1706.02945).