



HAL
open science

Optimal investment with possibly non-concave utilities and no-arbitrage: a measure theoretical approach

Romain Blanchard, Laurence Carassus, Miklos Rasonyi

► To cite this version:

Romain Blanchard, Laurence Carassus, Miklos Rasonyi. Optimal investment with possibly non-concave utilities and no-arbitrage: a measure theoretical approach. *Mathematical Methods of Operations Research*, 2018, 88, pp.241-281. 10.1007/s00186-018-0635-3 . hal-01883419

HAL Id: hal-01883419

<https://hal.science/hal-01883419v1>

Submitted on 28 Sep 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Optimal investment with possibly non-concave utilities and no-arbitrage: a measure theoretical approach

Romain Blanchard

LMR, Université Reims Champagne-Ardenne (URCA), France
romain.blanchard@etudiant.univ-reims.fr

Laurence Carassus

Research Center, Pôle Universitaire de Vinci and LMR, URCA, France
laurence.carassus@univ-reims.fr

Miklós Rásonyi

MTA Alfréd Rényi Institute of Mathematics, Hungary
rasonyi@renyi.mta.hu

January 20, 2018

Abstract

We consider a discrete-time financial market model with finite time horizon and investors with utility functions defined on the non-negative half-line. We allow these functions to be random, non-concave and non-smooth. We use a dynamic programming framework together with measurable selection arguments to establish both the characterization of the no-arbitrage property for such markets and the existence of an optimal portfolio strategy for such investors.

Key words: no-arbitrage condition ; non-concave utility functions; optimal investment

AMS 2000 subject classification: Primary 93E20, 91B70, 91B16 ; secondary 91G10, 28B20

1 Introduction

We consider investors trading in a multi-asset and discrete-time financial market. We revisit two classical problems: the characterization of no arbitrage and the maximisation of expected utility from the terminal wealth of an investor. There are strong connections between them: on one hand, no-arbitrage is a necessary condition for the existence of optimal strategies, see e.g. Pliska [1997], on the other hand, utility maximization is a means to construct dual objects (e.g. equivalent martingale measures), see Rogers [1994] in the frictionless case and Sass and Smaga [2014] for the case under transaction costs.

We consider a possibly non-concave and non-smooth *random* utility function U , defined on the non-negative half-line. For instance, U may be “ S -shaped” (i.e. convex up to a certain level of wealth and concave beyond that) but our results apply to a broader class of utility functions, e.g. to piecewise concave ones. We provide sufficient conditions which guarantee the existence of an optimal strategy. Such problems lie outside mainstream optimization theory and constitute an area of intensive study in recent years, see e.g. Bensoussan et al. [2015], He and Zhou [2011], Jin and Zhou [2008], Carlier and Dana [2011].

We are working in the setting of Carassus et al. [2015] and extend its results in several directions. First, we remove a restrictive integrability hypothesis [Carassus et al., 2015, Assumption 2.7]: $EU^-(\cdot, 0) < \infty$. Recall that the non-random version of this assumption: $U(0) < -\infty$ does not hold

true for the most widely used (concave) utility functions (logarithmic, $U(x) = -x^\alpha$ for $\alpha < 0$). Hence Carassus et al. [2015] excluded some of the most important cases. Note also that $U(0) = -\infty$ has the rather natural interpretation of an investor being infinitely averse of bankruptcy. We propose here an alternative, weaker integrability condition.

Second, we introduce a weaker version of the asymptotic elasticity assumption (see Assumption 4.10 below). In particular, Assumption 4.10 holds true for concave functions (see Remark 4.15) and therefore our result extends the one obtained in Rásonyi and Stettner [2006] to random utility functions.

Next, instead of using some Carathéodory utility function U as in Carassus et al. [2015] (*i.e.* U measurable in ω and continuous in x), we consider U which is measurable in ω and only upper semi-continuous (usc in the rest of the paper) in x . Allowing non-continuous U is unusual in the financial mathematics literature (see, however, Sass [2005]) but it is common in optimization theory. We highlight that this generalisation has a potential to model the investor's behaviour which can change suddenly after reaching a desired wealth level. Such a change can be expressed by a jump of U .

Next, we do not require that the value function is finite for all initial wealth as it is usually postulated in the maximisation of expected utility theory; instead we only assume the less restrictive and more tractable Assumption 4.7.

Finally, in contrast to the preceding literature, we do not need complete sigma-algebras for our arguments to work. This is an interesting technical aspect that can be useful in robust utility maximization where multiple probabilities are considered which lead to varying families of null sets.

We use methods that are different from the ones in Rásonyi and Stettner [2005], Rásonyi and Stettner [2006], Carassus and Rásonyi [2016] and Carassus et al. [2015], where similar multistep problems were treated. To solve our optimisation problem, we use dynamic programming as in those papers but here we propose a different approach which provides simpler proofs. As in Nutz [2016], we consider first a one period case where strategies are (deterministic) elements of \mathbb{R}^d . Then we use dynamic programming and measurable selection arguments, namely the Aumann Theorem (see, for example, Corollary 1 in Sainte-Beuve [1974]) to solve the multi-period problem. We will also often rely on the Projection Theorem (see for example Theorem 3.23 in Castaing and Valadier [1977]). Our modelisation of $(\Omega, \mathcal{F}, \mathfrak{F}, P)$ is more general than in Nutz [2016] since, as opposed to Nutz [2016], we consider only one probability measure so we don't have to postulate Borel spaces or analytic sets. The same methodology is used to reprove classical results on the characterization of no-arbitrage (see Rásonyi and Stettner [2005] and Jacod and Shiryaev [1998]) in our context of possibly incomplete sigma-algebras, we present this in Section 3 below.

Our paper can also be regarded as a review of problems in non-concave optimisation which arise in finance so detailed proofs are given. The paper is organized as follows: in section 2 we introduce our setup; section 3 contains the results on no-arbitrage; section 4 presents the main theorem on expected utility maximisation from terminal wealth; section 5 establishes the existence of an optimal strategy for the one period case; we prove our main theorem on utility maximisation in section 6. Finally, section 7 collects some technical results and proofs as well as elements about the measurability of random sets.

2 Set-up

Fix a time horizon $T \in \mathbb{N}$ and let $(\Omega_t)_{1 \leq t \leq T}$ be a sequence of non-empty sets equipped with the respective sigma-algebras $(\mathcal{G}_t)_{1 \leq t \leq T}$. For $t = 1, \dots, T$, we denote by Ω^t the t -fold Cartesian product $\Omega^t = \Omega_1 \times \dots \times \Omega_t$. An element of Ω^t will be denoted by $\omega^t = (\omega_1, \dots, \omega_t)$ for $(\omega_1, \dots, \omega_t) \in \Omega_1 \times \dots \times \Omega_t$. We also denote by \mathcal{F}_t the product sigma-algebra on Ω^t , $\mathcal{F}_t = \mathcal{G}_1 \otimes \dots \otimes \mathcal{G}_t$. For the sake of simplicity, we set $\Omega^0 := \{\omega_0\}$ and $\mathcal{F}_0 = \mathcal{G}_0 = \{\emptyset, \Omega^0\}$. We will omit the dependency in ω_0 in the rest of the paper. We denote by \mathfrak{F} the filtration $(\mathcal{F}_t)_{0 \leq t \leq T}$.

Let P_1 be a probability measure on \mathcal{F}_1 and q_{t+1} be a stochastic kernel on $\mathcal{G}_{t+1} \times \Omega^t$ for $t = 1, \dots, T-1$.

Namely we assume that for all $\omega^t \in \Omega^t$, $B \in \mathcal{G}_{t+1} \rightarrow q_{t+1}(B|\omega^t)$ is a probability on \mathcal{G}_{t+1} and for all $B \in \mathcal{G}_{t+1}$, $\omega^t \in \Omega^t \rightarrow q_{t+1}(B|\omega^t)$ is \mathcal{F}_t -measurable. Here we DO NOT assume that \mathcal{G}_1 contains the null sets of P_1 or that \mathcal{G}_{t+1} contains the null sets of $q_{t+1}(\cdot|\omega^t)$ for all $\omega^t \in \Omega^t$. Then we define for $A \in \mathcal{F}_t$ the probability P_t by Fubini's Theorem for stochastic kernels (see Lemma 7.1) via

$$P_t(A) = \int_{\Omega_1} \int_{\Omega_2} \cdots \int_{\Omega_t} 1_A(\omega_1, \dots, \omega_t) q_t(d\omega_t|\omega^{t-1}) \cdots q_2(d\omega_2|\omega^1) P_1(d\omega_1). \quad (1)$$

Finally, $(\Omega, \mathcal{F}, \mathfrak{F}, P) := (\Omega^T, \mathcal{F}_T, \mathfrak{F}, P_T)$ will be our basic filtered probability space. The expectation under P_t will be denoted by E_{P_t} ; when $t = T$, we simply write E .

We introduce \mathcal{F}_t^T , the filtration on Ω^T associated to \mathcal{F}_t , defined by

$$\mathcal{F}_t^T = \mathcal{G}_1 \otimes \dots \otimes \mathcal{G}_t \otimes \{\emptyset, \Omega_{t+1}\} \dots \otimes \{\emptyset, \Omega_T\}.$$

Let Ξ_t^T be the set of \mathcal{F}_t^T -measurable random variables from Ω^T to \mathbb{R}^d . Let $X_t: \Omega^T \rightarrow \Omega_t$, $X_t(\omega_1, \dots, \omega_T) = \omega_t$ be the coordinate mapping corresponding to t . Then $\mathcal{F}_t^T = \sigma(X_1, \dots, X_t)$. So $h \in \Xi_t^T$ if and only if there exists some $g \in \Xi_t$ such that $h = g(X_1, \dots, X_t)$. This implies that $h(\omega^T) = g(\omega^t)$. For ease of notation we will identify h and g and also $\mathcal{F}_t, \mathcal{F}_t^T, \Xi_t$ and Ξ_t^T .

Remark 2.1 If we choose Polish spaces for Ω_t , $t = 1, \dots, T$ then any probability measure P on Ω can be decomposed in the form of (1) (see Dellacherie and Meyer [1979] III.70-7).

The positive (resp. negative) part of a number or a random variable X is denoted by X^+ (resp. X^-). We will also write $f^\pm(X)$ for $(f(X))^\pm$ for any random variable X and (possibly random) function f .

In the rest of the paper we will use generalised integral: for some $f_t: \Omega^t \rightarrow \mathbb{R} \cup \{\pm\infty\}$, \mathcal{F}_t -measurable, such that $\int_{\Omega^t} f_t^+(\omega^t) P_t(d\omega^t) < \infty$ or $\int_{\Omega^t} f_t^-(\omega^t) P_t(d\omega^t) < \infty$, we define

$$\int_{\Omega^t} f_t(\omega^t) P_t(d\omega^t) := \int_{\Omega^t} f_t^+(\omega^t) P_t(d\omega^t) - \int_{\Omega^t} f_t^-(\omega^t) P_t(d\omega^t) \in \mathbb{R} \cup \{-\infty, \infty\}.$$

We refer to Lemma 7.1, Definition 7.2 and Proposition 7.3 of the Appendix for more details and properties. In particular, if f_t is non-negative or if f_t is such that $\int_{\Omega^t} f_t^+(\omega^t) P_t(d\omega^t) < \infty$ (this will be the two cases of interest in the paper) we can apply Fubini's Theorem¹ to get

$$\int_{\Omega^t} f_t(\omega^t) P_t(d\omega^t) = \int_{\Omega_1} \int_{\Omega_2} \cdots \int_{\Omega_t} f_t(\omega_1, \dots, \omega_t) q_t(d\omega_t|\omega^{t-1}) \cdots q_2(d\omega_2|\omega^1) P_1(d\omega_1).$$

We will denote by \mathcal{N}_{P_t} the set of P_t negligible sets of Ω^t i.e $\mathcal{N}_{P_t} = \{N \subset \Omega^t, \exists M \in \mathcal{F}_t, N \subset M \text{ and } P_t(M) = 0\}$. Let $\overline{\mathcal{F}}_t = \{A \cup N, A \in \mathcal{F}_t, N \in \mathcal{N}_{P_t}\}$ and $\overline{P}_t(A \cup N) = P_t(A)$ for $A \cup N \in \overline{\mathcal{F}}_t$. Then it is well known that \overline{P}_t is a measure on $\overline{\mathcal{F}}_t$ which coincides with P_t on \mathcal{F}_t , $(\Omega^t, \overline{\mathcal{F}}_t, \overline{P}_t)$ is a complete probability space and \overline{P}_t restricted to \mathcal{N}_{P_t} is zero. For $t = 0, \dots, T-1$, let Ξ_t be the set of \mathcal{F}_t -measurable random variables mapping Ω^t to \mathbb{R}^d .

The following lemma makes the link between conditional expectation and kernel.

Lemma 2.2 *Let $0 \leq s \leq t \leq T$. Let $h \in \Xi_t$ such that $\int_{\Omega^t} h^+ dP_t < \infty$ then*

$$\begin{aligned} E(h|\mathcal{F}_s) &= \varphi(X_1, \dots, X_s) P_s\text{-a.s., where} \\ \varphi(\omega_1, \dots, \omega_s) &= \int_{\Omega_{s+1} \times \dots \times \Omega_t} h(\omega_1, \dots, \omega_s, \omega_{s+1}, \dots, \omega_t) q_t(\omega_t|\omega^{t-1}) \cdots q_{s+1}(\omega_{s+1}|\omega^s). \end{aligned}$$

¹From now, we call Fubini's theorem the Fubini theorem for stochastic kernel (see eg Lemma 7.1, Proposition 7.3).

Proof. For the sake of completeness, the proof is reported in Section 7.3 of the Appendix. \square

Let $\{S_t, 0 \leq t \leq T\}$ be a d -dimensional \mathcal{F}_t -adapted process representing the price of d risky securities in the financial market in consideration. There exists also a riskless asset for which we assume a constant price equal to 1. The notation $\Delta S_t := S_t - S_{t-1}$ will often be used. If $x, y \in \mathbb{R}^d$ then the concatenation xy stands for their scalar product. The symbol $|\cdot|$ denotes the Euclidean norm on \mathbb{R}^d (or on \mathbb{R}).

Trading strategies are represented by d -dimensional predictable processes $(\phi_t)_{1 \leq t \leq T}$, where ϕ_t^i denotes the investor's holdings in asset i at time t ; predictability means that $\phi_t \in \Xi_{t-1}$. The family of all predictable trading strategies is denoted by Φ .

We assume that trading is self-financing. As the riskless asset's price is constant 1, the value at time t of a portfolio ϕ starting from initial capital $x \in \mathbb{R}$ is given by

$$V_t^{x,\phi} = x + \sum_{i=1}^t \phi_i \Delta S_i.$$

Example 2.3 For a better understanding of the present, abstract setting one may consider a simple trinomial tree model where $\Omega_t = \{-1, 0, 1\}$, $\mathcal{G}_t := 2^{\Omega_t}$ for each $t = 1, \dots, T$. Let $S_t := \sum_{j=1}^t X_j$, $t = 1, \dots, T$ where the X_j are the respective coordinate mappings. The kernels $q_{t+1}(\cdot|\omega^t)$ then assign probabilities to the 3 possible outcomes $\{-1, 0, 1\}$ conditionally to the past path $\omega^t = (\omega_1, \dots, \omega_t)$. P_1 is just a probability on $\{-1, 0, 1\}$. Strategies can be identified with functions $\phi_t : \Omega^{t-1} \rightarrow \mathbb{R}$ expressing the holding in the risky asset in the interval $(t-1, t]$.

3 No-arbitrage condition

The following absence of arbitrage condition or NA condition is standard, it is equivalent to the existence of a risk-neutral measure in discrete-time markets with finite horizon, see e.g. Dalang et al. [1990].

(NA) If $V_T^{0,\phi} \geq 0$ P-a.s. for some $\phi \in \Phi$ then $V_T^{0,\phi} = 0$ P-a.s.

Definition 3.1 Let (Ω, \mathcal{F}) be a measurable space and (T, \mathcal{T}) a topological space. A random set R is a set valued function that assigns to each $\omega \in \Omega$ a subset $R(\omega)$ of T . We write $R : \Omega \rightarrow T$. We say that R is measurable if for any open set $O \in T$ $\{\omega \in \Omega, R(\omega) \cap O \neq \emptyset\} \in \mathcal{F}$. The graph of R is defined as $Gr(R) := \{(\omega, t) \in \Omega \times T : t \in R(\omega)\}$.

We now provide classical tools and results about the (NA) condition and its ‘‘concrete’’ local characterization, see Proposition 3.6, that we will use in the rest of the paper. We start with the set D^{t+1} (see Definition 3.1) where $D^{t+1}(\omega^t)$ is the smallest affine subspace of \mathbb{R}^d containing the support of the distribution of $\Delta S_{t+1}(\omega^t, \cdot)$ under $q_{t+1}(\cdot|\omega^t)$. If $D^{t+1}(\omega^t) = \mathbb{R}^d$ then, intuitively, there are no redundant assets. Otherwise, for $\phi_{t+1} \in \Xi_t$, one may always replace $\phi_{t+1}(\omega^t, \cdot)$ by its orthogonal projection $\phi_{t+1}^\perp(\omega^t, \cdot)$ on $D^{t+1}(\omega^t)$ without changing the portfolio value since $\phi_{t+1}(\omega^t) \Delta S_{t+1}(\omega^t, \cdot) = \phi_{t+1}^\perp(\omega^t) \Delta S_{t+1}(\omega^t, \cdot)$, $q_{t+1}(\cdot|\omega^t)$ a.s., see Remark 5.2 and Lemma 7.16 below as well as Remark 9.1 of Föllmer and Schied [2002].

Definition 3.2 Let $0 \leq t \leq T$ be fixed. We define the random set $\tilde{D}^{t+1} : \Omega^t \rightarrow \mathbb{R}^d$ by

$$\tilde{D}^{t+1}(\omega^t) := \bigcap \left\{ A \subset \mathbb{R}^d, \text{ closed, } q_{t+1}(\Delta S_{t+1}(\omega^t, \cdot) \in A | \omega^t) = 1 \right\}. \quad (2)$$

For $\omega^t \in \Omega^t$, $\tilde{D}^{t+1}(\omega^t) \subset \mathbb{R}^d$ is the support of the distribution of $\Delta S_{t+1}(\omega^t, \cdot)$ under $q_{t+1}(\cdot|\omega^t)$. We also define the random set $D^{t+1} : \Omega^t \rightarrow \mathbb{R}^d$ by $D^{t+1}(\omega^t) := \text{Aff}(\tilde{D}^{t+1}(\omega^t))$, where Aff denotes the affine hull of a set.

Lemma 3.3 *Let $0 \leq t \leq T$ be fixed. Then \tilde{D}^{t+1} and D^{t+1} are both non-empty, closed-valued and \mathcal{F}_t -measurable random sets (see Definition 3.1). In particular, $\text{Graph}(D^{t+1}) \in \mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$.*

Proof. The proof is reported in Section 7.3 of the Appendix. \square

In Lemma 3.4, which is used in the proof of Lemma 3.5 for projection purposes, we obtain a well-known result : for $\omega^t \in \Omega^t$ fixed and under a local version of (NA), $D^{t+1}(\omega^t)$ is a vector subspace of \mathbb{R}^d (see for instance Theorem 1.48 of Föllmer and Schied [2002]). Then in Lemma 3.5 we prove that under the (NA) assumption, for P_t almost all ω^t , $D^{t+1}(\omega^t)$ is a vector subspace of \mathbb{R}^d . We also provide a local version of the (NA) condition (see (4)).

Lemma 3.4 *Let $0 \leq t \leq T$ and $\omega^t \in \Omega^t$ be fixed. Assume that for all $h \in D^{t+1}(\omega^t) \setminus \{0\}$*

$$q_{t+1}(h\Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) < 1.$$

Then $0 \in D^{t+1}(\omega^t)$ and the set $D^{t+1}(\omega^t)$ is actually a vector subspace of \mathbb{R}^d .

Proof. The proof is reported in Section 7.3 of the Appendix. \square

Lemma 3.5 *Assume that the (NA) condition holds true. Then for all $0 \leq t \leq T - 1$, there exists a full measure set Ω_{NA1}^t such that for all $\omega^t \in \Omega_{NA1}^t$, $0 \in D^{t+1}(\omega^t)$, i.e $D^{t+1}(\omega^t)$ is a vector space of \mathbb{R}^d . Moreover, for all $\omega^t \in \Omega_{NA1}^t$ and all $h \in \mathbb{R}^d$ we get that*

$$q_{t+1}(h\Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) = 1 \Rightarrow q_{t+1}(h\Delta S_{t+1}(\omega^t, \cdot) = 0 | \omega^t) = 1. \quad (3)$$

In particular, if $\omega^t \in \Omega_{NA1}^t$ and $h \in D^{t+1}(\omega^t)$ we obtain that

$$q_{t+1}(h\Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) = 1 \Rightarrow h = 0. \quad (4)$$

Proof. Let $0 \leq t \leq T$ be fixed. We introduce the following random set:

$$\Pi^t := \{ \omega^t \in \Omega^t, \exists h \in D^{t+1}(\omega^t), h \neq 0, q_{t+1}(h\Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) = 1 \}.$$

Assume for a moment that $\Pi^t \in \overline{\mathcal{F}}_t$ and that $\overline{P}_t(\Pi^t) = 0$. Let $\omega^t \in \Omega^t \setminus \Pi^t$. The fact that $0 \in D^{t+1}(\omega^t)$ directly follows from the definition of Π^t and from Lemma 3.4. We now prove (3). Let $h \in \mathbb{R}^d$ be fixed such that $q_{t+1}(h\Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) = 1$. If $h = 0$ there is nothing to prove. If $h \in D^{t+1}(\omega^t) \setminus \{0\}$, $\omega^t \in \Pi^t$ which is impossible. Now if $h \notin D^{t+1}(\omega^t)$ and $h \neq 0$, let h' be the orthogonal projection of h on $D^{t+1}(\omega^t)$. We first show that $q_{t+1}(h'\Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) = 1$. Indeed, if it were not the case the set $B := \{h'\Delta S_{t+1}(\omega^t, \cdot) < 0\}$ would verify $q_{t+1}(B | \omega^t) > 0$. Set $L^{t+1}(\omega^t) := (D^{t+1}(\omega^t))^\perp$. Using Lemma 7.16, since $(h - h') \in L^{t+1}(\omega^t)$, $q_{t+1}(A | \omega^t) = 1$ where $A := \{(h - h')\Delta S_{t+1}(\omega^t, \cdot) = 0\}$. We would therefore obtain that $q_{t+1}(A \cap B | \omega^t) > 0$ which implies that $q_{t+1}(h\Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) < 1$, a contradiction. Thus $q_{t+1}(h'\Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) = 1$. If $h' \neq 0$, as $h' \in D^{t+1}(\omega^t)$, $\omega^t \in \Pi^t$ which is again a contradiction. Thus $h' = 0$ and as $A \cap \{h'\Delta S_{t+1}(\omega^t, \cdot) = 0\} \subset \{h\Delta S_{t+1}(\omega^t, \cdot) = 0\}$, $q_{t+1}(h\Delta S_{t+1}(\omega^t, \cdot) = 0 | \omega^t) = 1$.

As $\Omega^t \setminus \Pi^t \in \overline{\mathcal{F}}_t$ there exists $\Omega_{NA1}^t \in \mathcal{F}_t$ and $N^t \in \mathcal{N}_{P_t}$ such that $\Omega^t \setminus \Pi^t = \Omega_{NA1}^t \cup N^t$ and $P_t(\Omega_{NA1}^t) = \overline{P}_t(\Omega^t \setminus \Pi^t) = 1$. Since $\Omega_{NA1}^t \subset \Omega^t \setminus \Pi^t$, it follows that for all $\omega^t \in \Omega_{NA1}^t$, $0 \in D^{t+1}(\omega^t)$ and for all $h \in \mathbb{R}^d$, (3) holds true.

We prove (4). Assume now that $\omega^t \in \Omega_{NA1}^t$ and $h \in D^{t+1}(\omega^t)$ are such that $q_{t+1}(h\Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) = 1$. Using (3) and Lemma 7.16 we get that $h \in L^{t+1}(\omega^t)$. So $h \in D^{t+1}(\omega^t) \cap L^{t+1}(\omega^t) = \{0\}$ and (4) holds true.

It remains to prove that $\Pi^t \in \overline{\mathcal{F}}_t$ and $\overline{P}_t(\Pi^t) = 0$. We introduce the random set

$$H^t(\omega^t) := \{ h \in D^{t+1}(\omega^t), h \neq 0, q_{t+1}(h\Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) = 1 \}.$$

Then $\Pi^t = \{\omega^t \in \Omega^t, H^t(\omega^t) \neq \emptyset\} = \text{proj}_{|\Omega^t} \text{Graph}(H^t)$.

We prove now that $\text{Graph}(H^t) \in \mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$. Indeed, we can rewrite that

$$\text{Graph}(H^t) = \text{Graph}(D^{t+1}) \cap \left\{ (\omega^t, h) \in \Omega^t \times \mathbb{R}^d, q_{t+1}(h \Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) = 1 \right\} \cap \left(\Omega^t \times \mathbb{R}^d \setminus \{0\} \right).$$

As from Lemma 7.7, $\{(\omega^t, h) \in \Omega^t \times \mathbb{R}^d, q_{t+1}(h \Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) = 1\} \in \mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$ and from Lemma 3.3, $\text{Graph}(D^{t+1}) \in \mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$, we obtain that $\text{Graph}(H^t) \in \mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$. The Projection Theorem applies and $\Pi^t = \{H^t \neq \emptyset\} = \text{proj}_{|\Omega^t} \text{Graph}(H^t) \in \overline{\mathcal{F}}_t$. From the Aumann Theorem there exists a $\overline{\mathcal{F}}_t$ -measurable selector $\bar{h}_{t+1} : \Pi^t \rightarrow \mathbb{R}^d$ such that $\bar{h}_{t+1}(\omega^t) \in H^t(\omega^t)$ for every $\omega^t \in \Pi^t$. We now extend \bar{h}_{t+1} on Ω^t by setting $\bar{h}_{t+1}(\omega^t) = 0$ for $\omega^t \in \Omega^t \setminus \Pi^t$. It is clear that \bar{h}_{t+1} remains $\overline{\mathcal{F}}_t$ -measurable. Applying Lemma 7.8, there exists $h_{t+1} : \Omega^t \rightarrow \mathbb{R}^d$ which is \mathcal{F}_t -measurable and satisfies $h_{t+1} = \bar{h}_{t+1}$ P_t -almost surely. Then if we set

$$\begin{aligned} \varphi(\omega^t) &= q_{t+1}(h_{t+1}(\omega^t) \Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t), \\ \bar{\varphi}(\omega^t) &= q_{t+1}(\bar{h}_{t+1}(\omega^t) \Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t), \end{aligned}$$

we get from Proposition 7.7 that φ is \mathcal{F}_t -measurable and from Proposition 7.4 *iii*) that $\bar{\varphi}$ is $\overline{\mathcal{F}}_t$ -measurable. Furthermore as $\{\omega^t \in \Omega^t, \varphi(\omega^t) \neq \bar{\varphi}(\omega^t)\} \subset \{\omega^t \in \Omega^t, h_t(\omega^t) \neq \bar{h}_{t+1}(\omega^t)\}$, $\varphi = \bar{\varphi}$ P_t -almost surely. This implies that $\int_{\Omega^t} \bar{\varphi} d\bar{P}_t = \int_{\Omega^t} \varphi dP_t$. Now we define the predictable process $(\phi_i)_{1 \leq i \leq T}$ by $\phi_{t+1} = h_{t+1}$ and $\phi_i = 0$ for $i \neq t+1$. Then

$$\begin{aligned} P(V_T^{0,\phi} \geq 0) &= P(h_{t+1} \Delta S_{t+1} \geq 0) = P_{t+1}(h_{t+1} \Delta S_{t+1} \geq 0) \\ &= \int_{\Omega^t} \varphi(\omega^t) P_t(d\omega^t) = \int_{\Omega^t} \bar{\varphi}(\omega^t) \bar{P}_t(d\omega^t) \\ &= \int_{\Pi^t} q_{t+1}(\bar{h}_t(\omega^t) \Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) \bar{P}_t(d\omega^t) + \\ &\quad \int_{\Omega^t \setminus \Pi^t} q_{t+1}(0 \times \Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) \bar{P}_t(d\omega^t) \\ &= \bar{P}_t(\Pi^t) + \bar{P}_t(\Omega^t \setminus \Pi^t) = 1, \end{aligned}$$

where we have used that if $\omega^t \in \Pi^t$, $\bar{h}_{t+1}(\omega^t) \in H^t(\omega^t)$ and otherwise $\bar{h}_{t+1}(\omega^t) = 0$. With the same arguments we obtain that

$$\begin{aligned} P(V_T^{0,\phi} > 0) &= P_t(h_{t+1} \Delta S_{t+1} > 0) \\ &= \int_{\Pi^t} q_{t+1}(\bar{h}_{t+1}(\omega^t) \Delta S_{t+1}(\omega^t, \cdot) > 0 | \omega^t) \bar{P}_t(d\omega^t) + \int_{\Omega^t \setminus \Pi^t} q_{t+1}(0 > 0 | \omega^t) \bar{P}_t(d\omega^t) \\ &= \int_{\Pi^t} q_{t+1}(\bar{h}_{t+1}(\omega^t) \Delta S_{t+1}(\omega^t, \cdot) > 0 | \omega^t) \bar{P}_t(d\omega^t). \end{aligned}$$

Let $\omega^t \in \Pi^t$ then $q_{t+1}(\bar{h}_{t+1}(\omega^t) \Delta S_{t+1}(\omega^t, \cdot) > 0 | \omega^t) > 0$. Indeed, if it is not the case then $q_{t+1}(\bar{h}_{t+1}(\omega^t) \Delta S_{t+1}(\omega^t, \cdot) \leq 0 | \omega^t) = 1$. As $\omega^t \in \Pi^t$, $\bar{h}_{t+1}(\omega^t) \in D^{t+1}(\omega^t)$ and $q_{t+1}(\bar{h}_{t+1}(\omega^t) \Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) = 1$, Lemma 7.16 applies and $\bar{h}_{t+1}(\omega^t) \in L^{t+1}(\omega^t)$. Thus we get that $\bar{h}_{t+1}(\omega^t) \in L^{t+1}(\omega^t) \cap D^{t+1}(\omega^t) = \{0\}$, a contradiction. So if $\bar{P}_t(\Pi^t) > 0$ we obtain that $P(V_T^{0,\phi} > 0) > 0$. This contradicts the (NA) condition and we obtain $\bar{P}_t(\Pi^t) = 0$, the required result. \square

Similarly as in Rásonyi and Stettner [2005] and Jacod and Shiryaev [1998], we prove a ‘‘quantitative’’ characterization of (NA).

Proposition 3.6 *Assume that the (NA) condition holds true and let $0 \leq t \leq T$. Then there exists $\Omega_{NA}^t \in \mathcal{F}_t$ with $P_t(\Omega_{NA}^t) = 1$ and $\Omega_{NA}^t \subset \Omega_{NA1}^t$ (see Lemma 3.5 for the definition of Ω_{NA1}^t) such that for all $\omega^t \in \Omega_{NA}^t$, there exists a \mathcal{F}_t -measurable $\omega^t \rightarrow \alpha_t(\omega^t) \in (0, 1]$ such that for all $h \in D^{t+1}(\omega^t)$*

$$q_{t+1}(h \Delta S_{t+1}(\omega^t, \cdot) \leq -\alpha_t(\omega^t) | h | \omega^t) \geq \alpha_t(\omega^t). \quad (5)$$

Proof. Let $\omega^t \in \Omega_{NA1}^t$ be fixed (Ω_{NA1}^t is defined in Lemma 3.5).

Step 1 : Proof of (5). Introduce the following set for $n \geq 1$

$$A_n(\omega^t) := \left\{ h \in D^{t+1}(\omega^t), |h| = 1, q_{t+1} \left(h \Delta S_{t+1}(\omega^t, \cdot) \leq -\frac{1}{n} |\omega^t \right) < \frac{1}{n} \right\}. \quad (6)$$

Let $\bar{n}_0(\omega^t) := \inf\{n \geq 1, A_n(\omega^t) = \emptyset\}$. Note that if $D^{t+1}(\omega^t) = \{0\}$, then $\bar{n}_0(\omega^t) = 1 < \infty$. We assume now that $D^{t+1}(\omega^t) \neq \{0\}$ and we prove by contradiction that $\bar{n}_0(\omega^t) < \infty$. Assume that $\bar{n}_0(\omega^t) = \infty$ i.e for all $n \geq 1$, $A_n(\omega^t) \neq \emptyset$. We thus get $h_n(\omega^t) \in D^{t+1}(\omega^t)$ with $|h_n(\omega^t)| = 1$ and such that $q_{t+1}(h_n(\omega^t) \Delta S_{t+1}(\omega^t, \cdot)) \leq -\frac{1}{n} |\omega^t| < \frac{1}{n}$. By passing to a sub-sequence we can assume that $h_n(\omega^t)$ tends to some $h^*(\omega^t) \in D^{t+1}(\omega^t)$ (recall that the set $D^{t+1}(\omega^t)$ is closed by definition) with $|h^*(\omega^t)| = 1$. Then $\{h^*(\omega^t) \Delta S_{t+1}(\omega^t, \cdot) < 0\} \subset \liminf_n B_n(\omega^t)$, where $B_n(\omega^t) := \{h_n(\omega^t) \Delta S_{t+1}(\omega^t, \cdot) \leq -1/n\}$. Furthermore as $1_{\liminf_n B_n(\omega^t)} = \liminf_n 1_{B_n(\omega^t)}$, Fatou's Lemma implies that

$$\begin{aligned} q_{t+1}(h^*(\omega^t) \Delta S_{t+1}(\omega^t, \cdot) < 0 | \omega^t) &\leq \int_{\Omega_{t+1}} 1_{\liminf_n B_n(\omega^t)}(\omega_{t+1}) q_{t+1}(\omega_{t+1} | \omega^t) \\ &\leq \liminf_n \int_{\Omega_{t+1}} 1_{B_n(\omega^t)}(\omega_{t+1}) q_{t+1}(\omega_{t+1} | \omega^t) = 0. \end{aligned}$$

This implies that $q_{t+1}(h^*(\omega^t) \Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) = 1$, and thus from (4) in Lemma 3.5 we get that $h^*(\omega^t) = 0$ which contradicts $|h^*(\omega^t)| = 1$. Thus $\bar{n}_0(\omega^t) < \infty$ and we can set for $\omega^t \in \Omega_{NA1}^t$, $\bar{\alpha}_t(\omega^t) = \frac{1}{\bar{n}_0(\omega^t)}$. It is clear that $\bar{\alpha}_t \in (0, 1]$. Then for all $\omega^t \in \Omega_{NA1}^t$, for all $h \in D^{t+1}(\omega^t)$ with $|h| = 1$, by definition of $A_{\bar{n}_0(\omega^t)}(\omega^t)$ we obtain

$$q_{t+1}(h \Delta S_{t+1}(\omega^t, \cdot) \leq -\bar{\alpha}_t(\omega^t) |\omega^t|) \geq \bar{\alpha}_t(\omega^t). \quad (7)$$

Step 2 : Measurability.

We now construct a function α_t which is \mathcal{F}_t -measurable and satisfies (5) as well. To do that we use the Aumann Theorem again as in the proof of Lemma 3.5 but this time applied to the random set $A_n : \Omega^t \rightarrow \mathbb{R}^d$ where $A_n(\omega^t)$ is defined in (6) if $\omega^t \in \Omega_{NA1}^t$ and $A_n(\omega^t) = \emptyset$ otherwise.

We prove that $\text{graph}(A_n) \in \mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$. From Lemma 7.7, the function $(\omega^t, h) \rightarrow q_{t+1}(h \Delta S_{t+1}(\omega^t, \cdot) \leq -\frac{1}{n} |\omega^t|)$ is $\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable. From Lemma 3.3, $\text{Graph}(D^{t+1}) \in \mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$ and the result follows from

$$\begin{aligned} \text{Graph}(A_n) &= \text{Graph}(D^{t+1}) \cap \left(\Omega_{NA1}^t \times \{h \in \mathbb{R}^d, |h| = 1\} \right) \\ &\cap \left\{ (\omega^t, h) \in \Omega^t \times \mathbb{R}^d, q_{t+1} \left(h \Delta S_{t+1}(\omega^t, \cdot) \leq -\frac{1}{n} |\omega^t \right) < \frac{1}{n} \right\}. \end{aligned}$$

Using the Projection Theorem, we get that $\{\omega^t \in \Omega^t, A_n(\omega^t) \neq \emptyset\} \in \bar{\mathcal{F}}_t$. We now extend \bar{n}_0 to Ω^t by setting $\bar{n}_0(\omega^t) = 1$ if $\omega^t \notin \Omega_{NA1}^t$. Then $\{\bar{n}_0 \geq 1\} = \Omega^t \in \mathcal{F}_t \subset \bar{\mathcal{F}}_t$ and for $k > 1$

$$\{\bar{n}_0 \geq k\} = \Omega_{NA1}^t \cap \bigcap_{1 \leq n \leq k-1} \{A_n \neq \emptyset\} \in \bar{\mathcal{F}}_t,$$

this implies that \bar{n}_0 and thus $\bar{\alpha}_t$ is $\bar{\mathcal{F}}_t$ -measurable. Using Lemma 7.8, we get some \mathcal{F}_t -measurable function α_t such that $\alpha_t = \bar{\alpha}_t P_t$ almost surely, i.e there exists $M^t \in \mathcal{F}_t$ such that $P_t(M^t) = 0$ and $\{\alpha_t \neq \bar{\alpha}_t\} \subset M^t$. We set $\Omega_{NA}^t := \Omega_{NA1}^t \cap (\Omega^t \setminus M^t)$. Then $P_t(\Omega_{NA}^t) = 1$ and as α_t is \mathcal{F}_t -measurable it remains to check that (5) holds true.

For $\omega^t \in \Omega_{NA}^t$, $\alpha_t(\omega^t) = \bar{\alpha}_t(\omega^t)$ (recall that $\omega^t \in \Omega^t \setminus M^t$) and since $\omega^t \in \Omega_{NA1}^t$, (7) holds true and consequently (5) as well. It is also clear that $\alpha_t(\omega^t) \in (0, 1]$ and the proof is completed. \square

Remark 3.7 The characterization of (NA) given by (5) works only for $h \in D^{t+1}(\omega^t)$. This is the reason why we will have to project the strategy $\phi_{t+1} \in \Xi_t$ onto $D^{t+1}(\omega^t)$ in our proofs.

Example 3.8 Getting back to Example 2.3, (NA) holds iff, for all $t = 1, \dots, T-1$ and $\omega^t \in \Omega^t$, either both $q_{t+1}(\{1\} | \omega^t) > 0$ and $q_{t+1}(\{-1\} | \omega^t) > 0$ or $q_{t+1}(\{0\} | \omega^t) = 1$ hold, and either both $P_1(\{1\}) > 0$ and $P_1(\{-1\}) > 0$ or $P_1(\{0\}) = 1$ hold.

4 Utility problem and main result

We now describe the investor's risk preferences by a possibly non-concave, random utility function.

Definition 4.1 A *random utility* is any function $U : \Omega \times \mathbb{R} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ satisfying the following conditions

- for every $x \in \mathbb{R}$, the function $U(\cdot, x) : \Omega \rightarrow \mathbb{R} \cup \{\pm\infty\}$ is \mathcal{F} -measurable,
- for all $\omega \in \Omega$, the function $U(\omega, \cdot) : \mathbb{R} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ is non-decreasing and usc on \mathbb{R} ,
- $U(\cdot, x) = -\infty$, for all $x < 0$.

We introduce the following notations.

Definition 4.2 For all $x \geq 0$, we denote by $\Phi(x)$ the set of all strategies $\phi \in \Phi$ such that $P_T(V_T^{x,\phi}(\cdot) \geq 0) = 1$ and by $\Phi(U, x)$ the set of all strategies $\phi \in \Phi(x)$ such that $EU(\cdot, V_T^{x,\phi})$ exists in a generalised sense, i.e. either $EU^+(\cdot, V_T^{x,\phi}(\cdot)) < \infty$ or $EU^-(\cdot, V_T^{x,\phi}(\cdot)) < \infty$.

Remark 4.3 Under (NA), if $\phi \in \Phi(x)$ then we have that $P_t(V_t^{x,\phi}(\cdot) \geq 0) = 1$ for all $1 \leq t \leq T$ see Lemma 7.17.

We now formulate the problem which is our main concern in the sequel.

Definition 4.4 Let $x \geq 0$. The *portfolio optimization problem* on a finite horizon T with initial wealth x is

$$u(x) := \sup_{\phi \in \Phi(U, x)} EU(\cdot, V_T^{x,\phi}(\cdot)). \quad (8)$$

Remark 4.5 Assume that there exists some P -full measure set $\tilde{\Omega} \in \mathcal{F}$ such that for all $\omega \in \tilde{\Omega}$, $x \rightarrow U(\omega, x)$ is non-decreasing and usc on $[0, +\infty)$ We set $\bar{U} : \Omega \times \mathbb{R} \rightarrow \mathbb{R} \cup \{\pm\infty\}$

$$\bar{U}(\omega, x) := U(\omega, x)1_{\tilde{\Omega} \times [0, +\infty)}(\omega, x) + (-\infty)1_{\Omega \times (-\infty, 0)}(\omega, x).$$

Then \bar{U} satisfies Definition 4.1, see Lemma 7.9 for the second item. Moreover, the value function does not change $u(x) = \sup_{\phi \in \Phi(U, x)} E\bar{U}(\cdot, V_T^{x,\phi}(\cdot))$, and if there exists some $\phi^* \in \Phi(U, x)$ such that $u(x) = E\bar{U}(\cdot, V_T^{x,\phi^*}(\cdot))$, then ϕ^* is an optimal solution for (8).

Remark 4.6 Let U be a utility function defined only on $(0, \infty)$ and verifying for every $x \in (0, \infty)$, $U(\cdot, x) : \Omega \rightarrow \mathbb{R} \cup \{\pm\infty\}$ is \mathcal{F} -measurable and for all $\omega \in \Omega$, $U(\omega, \cdot) : (0, \infty) \rightarrow \mathbb{R} \cup \{\pm\infty\}$ is non-decreasing and usc on $(0, \infty)$. We may extend U on \mathbb{R} by setting, for all $\omega \in \Omega$, $\bar{U}(\omega, 0) = \lim_{x \rightarrow 0} U(\omega, x)$ and for $x < 0$, $\bar{U}(\omega, x) = -\infty$. Then, as before, \bar{U} verifies Definition 4.1 and the value function has not changed.

We now present conditions on U which allow to assert that if $\phi \in \Phi(x)$ then $EU(\cdot, V_T^{x,\phi}(\cdot))$ is well-defined and that there exists some optimal solution for (8).

Assumption 4.7 For all $\phi \in \Phi(U, 1)$, $EU^+(\cdot, V_T^{1,\phi}(\cdot)) < \infty$.

Assumption 4.8 $\Phi(U, 1) = \Phi(1)$.

Remark 4.9 Assumptions 4.7 and 4.8 are connected but play a different role. Assumption 4.8 guarantees that $EU\left(\cdot, V_T^{1,\phi}(\cdot)\right)$ is well-defined for all $\Phi \in \Phi(1)$ and allows us to relax Assumption 2.7 of Carassus et al. [2015] on the behavior of U around 0, namely that $EU^-(\cdot, 0) < \infty$. Then Assumption 4.7 (together with Assumption 4.10) is used to show that $u(x) < \infty$ for all $x > 0$.

In Proposition 6.1, we will show that under Assumptions 4.7, 4.8 and 4.10, $EU^+\left(\cdot, V_T^{x,\phi}(\cdot)\right) < \infty$ for all $x \geq 0$ and $\phi \in \Phi(x)$. Thus $\Phi(U, x) = \Phi(x)$. Note that if there exists some $\Phi \in \Phi(U, x)$ such that $EU^+\left(\cdot, V_T^{x,\phi}(\cdot)\right) = \infty$ and $EU^-\left(\cdot, V_T^{x,\phi}(\cdot)\right) < \infty$ then $u(x) = \infty$ and the problem is ill-posed.

We propose some examples where Assumptions 4.7 or 4.8 hold true. Example *ii*) illustrates the distinction between Assumptions 4.7 and 4.8 and justifies why we do not merge both assumptions and postulate that $EU^+\left(\cdot, V_T^{1,\phi}(\cdot)\right) < \infty$, for all $\phi \in \Phi(1)$.

i) If U is bounded above then both Assumptions are trivially true. We get directly that $\Phi(U, x) = \Phi(x)$ for all $x \geq 0$.

ii) Assume that $EU^-(\cdot, 0) < \infty$ holds true. Let $x \geq 0$ and $\phi \in \Phi(x)$ be fixed. Using that U^- is non-increasing for all $\omega \in \Omega$ we get that

$$EU^-(\cdot, V_T^{x,\phi}(\cdot)) \leq EU^-(\cdot, 0) < +\infty,$$

Thus $EU(\cdot, V_T^{x,\phi}(\cdot))$ is well-defined, $\Phi(U, x) = \Phi(x)$ and Assumption 4.8 holds true.

iii) Assume that there exists some $\hat{x} \geq 1$ such that $U(\cdot, \hat{x} - 1) \geq 0$ P -almost surely and

$$\hat{u}(\hat{x}) := \sup_{\phi \in \Phi(\hat{x})} EU(\cdot, V_T^{\hat{x},\phi}(\cdot)) < \infty,$$

where we set for $\phi \in \Phi(\hat{x}) \setminus \Phi(U, \hat{x})$, $EU(\cdot, V_T^{\hat{x},\phi}(\cdot)) = -\infty$. Let $\phi \in \Phi(1)$ be fixed. Then using that U is non-decreasing for all $\omega \in \Omega$, we have that P -almost surely

$$U(\cdot, V_T^{1,\phi}(\cdot) + \hat{x} - 1) \geq U(\cdot, \hat{x} - 1) \geq 0.$$

Therefore $U(\cdot, V_T^{1,\phi}(\cdot) + \hat{x} - 1) = U^+(\cdot, V_T^{1,\phi}(\cdot) + \hat{x} - 1)$ P -almost surely. Now using that U^+ is non-decreasing for all $\omega \in \Omega$ we get that for all $\phi \in \Phi(1)$

$$EU^+(\cdot, V_T^{1,\phi}(\cdot)) \leq EU^+(\cdot, V_T^{1,\phi}(\cdot) + \hat{x} - 1) = EU(\cdot, V_T^{1,\phi}(\cdot) + \hat{x} - 1) \leq \hat{u}(\hat{x}) < +\infty$$

and Assumptions 4.7 and 4.8 are satisfied. Instead of stipulating that $\hat{u}(\hat{x}) < \infty$ it is enough to assume that $EU(\cdot, V_T^{\hat{x},\phi}(\cdot)) < \infty$ for all $\phi \in \Phi(\hat{x})$.

iv) We will prove in Theorem 4.17 that under the (NA) condition and Assumption 4.10, Assumptions 4.7 and 4.8 hold true if $EU^+(\cdot, 1) < +\infty$ and if for all $0 \leq t \leq T$ $|\Delta S_t|, \frac{1}{\alpha_t} \in \mathcal{W}_t$ (see (14) for the definition of \mathcal{W}_t).

Assumption 4.10 We assume that there exist some constants $\bar{\gamma} \geq 0$, $K > 0$, as well as a random variable C satisfying $C(\omega) \geq 0$ for all $\omega \in \Omega$ and $E(C) < \infty$ such that for all $\omega \in \Omega$, $\lambda \geq 1$ and $x \in \mathbb{R}$, we have

$$U(\omega, \lambda x) \leq K\lambda^{\bar{\gamma}} \left(U\left(\omega, x + \frac{1}{2}\right) + C(\omega) \right). \quad (9)$$

Remark 4.11 First note that the constant $\frac{1}{2}$ in (9) has been chosen arbitrarily. Indeed, assume there exists some constant $\bar{x} \geq 0$ such that for all $\omega \in \Omega$, $\lambda \geq 1$ and $x \in \mathbb{R}$

$$U(\omega, \lambda x) \leq K\lambda^{\bar{\gamma}} (U(\omega, x + \bar{x}) + C(\omega)). \quad (10)$$

Using the monotonicity of U , we can always assume $\bar{x} > 0$. Set for all $\omega \in \Omega$ and $x \in \mathbb{R}$, $\bar{U}(\omega, x) = U(\omega, 2\bar{x}x)$. Then for all $\omega \in \Omega$, $\lambda \geq 1$ and $x \in \mathbb{R}$, we have that

$$\bar{U}(\omega, \lambda x) = U(\omega, 2\lambda\bar{x}x) \leq K\lambda^{\bar{\gamma}} (U(\omega, 2\bar{x}x + \bar{x}) + C(\omega)) = K\lambda^{\bar{\gamma}} \left(\bar{U} \left(\omega, x + \frac{1}{2} \right) + C(\omega) \right),$$

and \bar{U} satisfies (9). It is clear that if ϕ^* is an optimal solution for the problem

$\bar{u}(x) := \sup_{\phi \in \Phi(\bar{U}, \frac{x}{2\bar{x}})} E\bar{U}(\cdot, V_T^{\frac{x}{2\bar{x}}, \phi}(\cdot))$ then $2\bar{x}\phi^*$ is an optimal solution for (8). Note as well that, since $K > 0$ and $C \geq 0$, it is immediate to see that for all $\omega \in \Omega$, $\lambda \geq 1$ and $x \in \mathbb{R}$

$$U^+(\omega, \lambda x) \leq K\lambda^{\bar{\gamma}} \left(U^+ \left(\omega, x + \frac{1}{2} \right) + C(\omega) \right). \quad (11)$$

Remark 4.12 We now provide some insight on Assumption 4.10. As the inequality (9) is used to control the behaviour of $U^+(\cdot, x)$ for large values of x , the usual assumption in the non-concave case (see Assumption 2.10 in Carassus et al. [2015]) is that there exists some $\hat{x} \geq 0$ such that $EU^+(\cdot, \hat{x}) < \infty$ as well as a random variable C_1 satisfying $E(C_1) < \infty$ and $C_1(\omega) \geq 0$ for all ω ² such that for all $x \geq \hat{x}$, $\lambda \geq 1$ and $\omega \in \Omega$

$$U(\omega, \lambda x) \leq \lambda^{\bar{\gamma}} (U(\omega, x) + C_1(\omega)). \quad (12)$$

We prove now that if (12) holds true then (10) is verified with $\bar{x} = \hat{x}$, $K = 1$ and $C = C_1$. Indeed, for $x \geq 0$, using the monotonicity of U , we have for all $\omega \in \Omega$ and $\lambda \geq 1$ that

$$U(\omega, \lambda x) \leq U(\omega, \lambda(x + \hat{x})) \leq \lambda^{\bar{\gamma}} (U(\omega, x + \hat{x}) + C_1(\omega)).$$

Therefore (10) is a weaker assumption than (12). Note as well that if we assume that (12) holds true for all $x > 0$, then if $0 < x < 1$ and $\omega \in \Omega$ we have

$$U(\omega, 1) \leq \left(\frac{1}{x} \right)^{\bar{\gamma}} (U(\omega, x) + C_1(\omega)),$$

and $U(\omega, 0) := \lim_{x \rightarrow 0, x > 0} U(\omega, x) \geq -C_1(\omega)$. This excludes for instance the case where U is the logarithm. Furthermore, this also implies that $EU^-(\cdot, 0) \leq EC_1 < \infty$ and we are back to Assumption 2.7 of Carassus et al. [2015]

Alternatively, recalling the way the concave case is handled (see Lemma 2 in Rásonyi and Stettner [2005]), we could have introduced that there exists a random variable C_2 satisfying $E(C_2) < \infty$ and $C_2 \geq 0$ such that for all $x \in \mathbb{R}$, $\omega \in \Omega$

$$U^+(\omega, \lambda x) \leq \lambda^{\bar{\gamma}} (U^+(\omega, x) + C_2(\omega)). \quad (13)$$

We have not done so as it is difficult to prove that this inequality is preserved through the dynamic programming procedure when considering non-concave functions unless we assume that $EU^-(\cdot, 0) < \infty$ as in Carassus et al. [2015].

Remark 4.13 If there exists some set $\Omega_{AE} \in \mathcal{F}$ with $P(\Omega_{AE}) = 1$ such that (9) holds true only for $\omega \in \Omega_{AE}$, then setting as in Remark 4.5, $\bar{U}(\omega, x) := U(\omega, x)1_{\Omega_{AE} \times \mathbb{R}}(\omega, x)$, \bar{U} satisfies (9) and the value function in (8) does not change. We also assume without loss of generality that $C(\omega) \geq 0$ for all ω in (9). Indeed, if $C \geq 0$ P -a.s, we could consider $\tilde{C} := C\mathbb{1}_{C \geq 0}$. Then Assumption 4.10 would hold true with \tilde{C} instead of C .

²In the cited paper $C_1 \geq 0$ a.s but this is not an issue, see Remark 4.13 below

Remark 4.14 In the case where (12) holds true, we refer to remark 2.5 of Carassus and Rásonyi [2016] and remark 2.10 of Carassus et al. [2015] for the interpretation of $\bar{\gamma}$: for $C_1 = 0$, it can be seen as a generalization of the “asymptotic elasticity” of U at $+\infty$ (see Kramkov and Schachermayer [1999]). So (12) requires that the (generalized) asymptotic elasticity at $+\infty$ is finite. In this case and if U is differentiable there is a nice economic interpretation of the “asymptotic elasticity” as the ratio of “marginal utility”: $U'(x)$ and the “average utility”: $\frac{U(x)}{x}$, see again Section 6 of Kramkov and Schachermayer [1999] for further discussions. The case $C_1 > 0$ allows bounded utilities. In Carassus et al. [2015] it is proved that unlike in the concave case, the fact that U is bounded from above (and therefore satisfies (10)) does not imply that the asymptotic elasticity is bounded.

We propose now an example of an unbounded utility function satisfying (10) and such that $\limsup_{x \rightarrow \infty} \frac{xU'(x)}{U(x)} = +\infty$. This shows (as the counterexample of Carassus et al. [2015]), that Assumption 4.10 is less strong than the usual “asymptotic elasticity”. Let $U : \mathbb{R} \rightarrow \mathbb{R}$ be defined by

$$U(x) = -\infty 1_{(-\infty, 0)}(x) + \sum_{p \geq 0} p 1_{[p, p+1 - \frac{1}{2^{p+1}})}(x) + f_p(x) 1_{[p+1 - \frac{1}{2^{p+1}}, p+1)}(x)$$

where $f_p(x) = 2^{p+1}x + (p+1)(1 - 2^{p+1})$ for $p \in \mathbb{N}$. Then U satisfies Definition 4.1 and we have

$$U'(x) = \sum_{p \geq 0} 2^{p+1} 1_{[p+1 - \frac{1}{2^{p+1}}, p+1)}(x).$$

We prove that (10) holds true. Note that for all $x \geq 0$ we have $x - 1 \leq U(x) \leq x + 1$. Let $x \geq 0$ and $\lambda \geq 1$ be fixed. Then we get that

$$U(\lambda x) \leq \lambda x + 1 \leq \lambda(U(x+1) + 1) + 1 \leq \lambda(U(x+1) + 2),$$

and (10) is true with $K = \bar{x} = 1$ and $C = 2$. Now for $k \geq 0$, let $x_k = k + 1 - \frac{1}{2^{k+2}}$. We have $U(x_k) = f_k(x_k) = k + \frac{1}{2}$ and

$$\frac{x_k U'(x_k)}{U(x_k)} = 2^{k+1} \frac{(k+1 - \frac{1}{2^{k+2}})}{k + \frac{1}{2}} \rightarrow_{k \rightarrow \infty} +\infty.$$

Remark 4.15 We propose further examples where Assumption 4.10 holds true.

- i) Assume that U is bounded from above by some integrable random constant $C_1 \geq 0$ and that $EU^-(\cdot, \frac{1}{2}) < \infty$. Then for all $x \geq 0$, $\lambda \geq 1$, $\omega \in \Omega$ we have

$$\begin{aligned} U(\omega, \lambda x) &\leq C_1(\omega) \leq \lambda U\left(\omega, x + \frac{1}{2}\right) + \lambda \left(C_1(\omega) - U\left(\omega, x + \frac{1}{2}\right)\right) \\ &\leq \lambda U\left(\omega, x + \frac{1}{2}\right) + \lambda \left(C_1(\omega) + U^-\left(\omega, \frac{1}{2}\right)\right), \end{aligned}$$

and (9) holds true for $x \geq 0$ with $K = 1$, $\bar{\gamma} = 1$ and $C(\cdot) = C_1(\cdot) + U^-(\cdot, \frac{1}{2})$. As $U(\cdot, x) = -\infty$ for $x < 0$, (9) is true for all $x \in \mathbb{R}$.

- ii) Assume that U satisfies Definition 4.1 and that the restriction of U to $[0, \infty)$ is concave and non-decreasing and that $EU^-(\cdot, 1) < \infty$. We use similar arguments as in Lemma 2 in Rásonyi and Stettner [2006]. Indeed, let $x \geq 2$, $\lambda \geq 1$ be fixed we have

$$\begin{aligned} U(\omega, \lambda x) &\leq U(\omega, x) + U'(\omega, x)(\lambda x - x) \leq U(\omega, x) + \frac{U(\omega, x) - U(\omega, 1)}{x - 1}(\lambda - 1)x \\ &\leq U(\omega, x) + 2(\lambda - 1)(U(\omega, x) - U(\omega, 1)) \\ &\leq U(\omega, x) + 3\left(\lambda - \frac{1}{3}\right)(U(\omega, x) - U(\omega, 1)) \\ &\leq 3\lambda(U(\omega, x) + U^-(\omega, 1)), \end{aligned}$$

where we have used the concavity of U for the first two inequalities and the fact that $x \geq 2$ and U is non-decreasing for the other ones. Thus from the proof that (12) implies (10), we obtain that (10) holds true with $K = 3$, $\bar{\gamma} = 1$, $\bar{x} = 2$ and $C(\cdot) = U^-(\cdot, 1)$.

We can now state our main result.

Theorem 4.16 *Assume the (NA) condition and that Assumptions 4.7, 4.8 and 4.10 hold true. Let $x \geq 0$. Then, $u(x) < \infty$ and there exists some optimal strategy $\phi^* \in \Phi(U, x)$ such that*

$$u(x) = EU(\cdot, V_T^{x, \phi^*}(\cdot)).$$

Moreover $\phi_t^*(\cdot) \in D^t(\cdot)$ a.s. for all $0 \leq t \leq T$.

We will use dynamic programming in order to prove our main result. We will combine the approach of Rásonyi and Stettner [2005], Rásonyi and Stettner [2006], Carassus and Rásonyi [2016], Carassus et al. [2015] and Nutz [2016]. As in Nutz [2016], we will consider a one period case where the initial filtration is trivial (so that strategies are in \mathbb{R}^d) and thus the proofs are much simpler than in the other cited papers. The price to pay is that in the multi-period case where we use intensively measurable selection arguments (as in Nutz [2016]) in order to obtain Theorem 4.16. In our model, there is only one probability measure, so we don't have to introduce Borel spaces and analytic sets. Thus our modelisation of $(\Omega, \mathcal{F}, \mathfrak{F}, P)$ is more general than the one of Nutz [2016] restricted to one probability measure. As we are in a non concave setting we use similar ideas to those of Carassus and Rásonyi [2016] and Carassus et al. [2015].

Finally, as in Rásonyi and Stettner [2005], Rásonyi and Stettner [2006], Carassus and Rásonyi [2016] and Carassus et al. [2015], we propose the following result as a simpler but still general setting where Theorem 4.16 applies. We introduce for all $0 \leq t \leq T$

$$\mathcal{W}_t := \{X : \Omega^t \rightarrow \mathbb{R} \cup \{\pm\infty\}, \mathcal{F}_t\text{-measurable, } E|X|^p < \infty \text{ for all } p > 0\} \quad (14)$$

Theorem 4.17 *Assume the (NA) condition and that Assumption 4.10 hold true. Assume furthermore that $EU^+(\cdot, 1) < +\infty$ and that for all $0 \leq t \leq T$ $|\Delta S_t|, \frac{1}{\alpha_t} \in \mathcal{W}_t$. Let $x \geq 0$. Then, for all $\phi \in \Phi(x)$ and all $0 \leq t \leq T$, $V_t^{x, \phi} \in \mathcal{W}_t$. Moreover, there exists some optimal strategy $\phi^* \in \Phi(U, x)$ such that*

$$u(x) = EU(\cdot, V_T^{x, \phi^*}(\cdot)) < \infty$$

Proof of the main theorems will appear in Section 6.

Remark 4.18 The assumptions of Theorem 4.17 clearly hold for the model of Example 2.3 above when U is deterministic satisfying Assumption 4.10. However, they also hold for many models with an infinite Ω .

5 One period case

Let $(\bar{\Omega}, \mathcal{H}, Q)$ be a probability space (we denote by E the expectation under Q) and $Y(\cdot)$ an \mathcal{H} -measurable \mathbb{R}^d -valued random variable. Let $D \subset \mathbb{R}^d$ be the smallest affine subspace of \mathbb{R}^d containing the support of the distribution of $Y(\cdot)$. We assume that D contains 0. The condition corresponding to (NA) in the present setting is

Assumption 5.1 There exists some constant $0 < \alpha \leq 1$ such that for all $h \in D$

$$Q(hY(\cdot) \leq -\alpha|h|) \geq \alpha. \quad (15)$$

Remark 5.2 below is exactly Remark 8 of Carassus and Rásonyi [2016] (see also Lemma 2.6 of Nutz [2016]).

Remark 5.2 Let $h \in \mathbb{R}^d$ and let $h' \in \mathbb{R}^d$ be the orthogonal projection of h on D . Then $h - h' \perp D$ hence $\{Y(\cdot) \in D\} \subset \{(h - h')Y(\cdot) = 0\}$. It follows that

$$Q(hY(\cdot) = h'Y(\cdot)) = Q((h - h')Y(\cdot) = 0) \geq Q(Y(\cdot) \in D) = 1$$

by the definition of D . Hence $Q(hY(\cdot) = h'Y(\cdot)) = 1$.

Assumption 5.3 We consider a *random utility* $V : \bar{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$ satisfying the following conditions

- for every $x \in \mathbb{R}$, the function $V(\cdot, x) : \bar{\Omega} \rightarrow \mathbb{R}$ is \mathcal{H} -measurable,
- for every $\omega \in \bar{\Omega}$, the function $V(\omega, \cdot) : \mathbb{R} \rightarrow \mathbb{R}$ is non-decreasing and usc on \mathbb{R} ,
- $V(\cdot, x) = -\infty$, for all $x < 0$.

Let $x \geq 0$ be fixed. We define

$$\mathcal{H}_x := \left\{ h \in \mathbb{R}^d, Q(x + hY(\cdot) \geq 0) = 1 \right\}, \quad (16)$$

$$D_x := \mathcal{H}_x \cap D. \quad (17)$$

It is clear that \mathcal{H}_x and D_x are closed subsets of \mathbb{R}^d . We define

$$v(x) = (-\infty)1_{(-\infty, 0)}(x) + 1_{[0, +\infty)}(x) \sup_{h \in \mathcal{H}_x} EV(\cdot, x + hY(\cdot)). \quad (18)$$

From Remark 5.2,

$$v(x) = (-\infty)1_{(-\infty, 0)}(x) + 1_{[0, +\infty)}(x) \sup_{h \in D_x} EV(\cdot, x + hY(\cdot)). \quad (19)$$

Remark 5.4 It will be shown in Lemma 5.9 that under Assumptions 5.1, 5.3, 5.5 and 5.7, for all $h \in \mathcal{H}_x$, $E(V(\cdot, x + hY(\cdot)))$ is well-defined and more precisely that $EV^+(\cdot, x + hY(\cdot)) < +\infty$. So, under this set of assumptions, $\Phi(V, x)$, the set of $h \in \mathcal{H}_x$ such that $EV(\cdot, x + hY(\cdot))$ is well-defined, equals \mathcal{H}_x .

We present now the assumptions which allow to assert that there exists some optimal solution for (18). First we introduce an ‘‘asymptotic elasticity’’ assumption.

Assumption 5.5 There exist some constants $\bar{\gamma} \geq 0$, $K > 0$, as well as some \mathcal{H} -measurable C with $C(\omega) \geq 0$ for all $\omega \in \bar{\Omega}$ and $E(C) < \infty$, such that for all $\omega \in \bar{\Omega}$, for all $\lambda \geq 1$, $x \in \mathbb{R}$ we have

$$V(\omega, \lambda x) \leq K\lambda^{\bar{\gamma}} \left(V\left(\omega, x + \frac{1}{2}\right) + C(\omega) \right). \quad (20)$$

Remark 5.6 The same comments as in Remark 4.13 apply. Furthermore, note that since $K > 0$ and $C \geq 0$ we also have that for all $\omega \in \bar{\Omega}$, all $\lambda \geq 1$ and $x \in \mathbb{R}$

$$V^+(\omega, \lambda x) \leq K\lambda^{\bar{\gamma}} \left(V^+\left(\omega, x + \frac{1}{2}\right) + C(\omega) \right). \quad (21)$$

We introduce now some integrability assumption on V^+ .

Assumption 5.7 For every $h \in \mathcal{H}_1$,

$$EV^+(\cdot, 1 + hY(\cdot)) < \infty. \quad (22)$$

The following lemma corresponds to Lemma 2.1 of Rásonyi and Stettner [2006] in the deterministic case.

Lemma 5.8 Assume that Assumption 5.1 holds true. Then $D_x \subset B(0, \frac{x}{\alpha})$ where $B(0, \frac{x}{\alpha}) = \{h \in \mathbb{R}^d, |h| \leq \frac{x}{\alpha}\}$ and D_x is a convex, compact subspace of \mathbb{R}^d .

Proof. Let $h \in D_x$. Assume that $|h| > \frac{x}{\alpha}$ and let $\omega \in \{hY(\cdot) \leq -\alpha|h|\}$. Then $x + hY(\omega) < x - \alpha|h| < 0$ and from Assumption 5.1 $Q(x + hY(\cdot) < 0) \geq Q(hY(\cdot) \leq -\alpha|h|) \geq \alpha > 0$, a contradiction. The convexity and the closedness of D_x are clear and compactness follows from the boundness property. \square

This lemma corresponds in the deterministic case to Lemma 4.8 of Carassus et al. [2015] (see also Lemma 2.3 of Rásonyi and Stettner [2006] and Lemma 2.8 of Nutz [2016]).

Lemma 5.9 Assume that Assumptions 5.1, 5.3, 5.5 and 5.7 hold true. Then there exists a \mathcal{H} -measurable $L \geq 0$ satisfying $E(L) < \infty$ and such that for all $x \geq 0$ and $h \in \mathcal{H}_x$

$$V^+(\cdot, x + hY(\cdot)) \leq ((2x)^{\bar{\gamma}}K + 1)L(\cdot)Q - a.s. \quad (23)$$

Proof. The proof is reported in Section 7.3 of the Appendix \square

Lemma 5.10 Assume that Assumptions 5.1, 5.3, 5.5 and 5.7 hold true. Let \mathcal{D} be the set valued function that assigns to each $x \geq 0$ the set D_x . Then $Graph(\mathcal{D}) := \{(x, h) \in [0, +\infty) \times \mathbb{R}^d, h \in D_x\}$ is a closed subset of $\mathbb{R} \times \mathbb{R}^d$. Let $\psi : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R} \cup \{\pm\infty\}$ be defined by

$$\psi(x, h) := \begin{cases} EV(\cdot, x + hY(\cdot)), & \text{if } (x, h) \in Graph(\mathcal{D}) \\ -\infty, & \text{otherwise.} \end{cases} \quad (24)$$

Then ψ is usc on $\mathbb{R} \times \mathbb{R}^d$ and $\psi < +\infty$ on $Graph(\mathcal{D})$.

Proof. Let $(x_n, h_n)_{n \geq 1} \in Graph(\mathcal{D})$ be a sequence converging to some $(x^*, h^*) \in \mathbb{R} \times \mathbb{R}^d$. We prove first that $(x^*, h^*) \in Graph(\mathcal{D})$. It is clear that $x^* \geq 0$. Set for $n \geq 1$ $E_n := \{\omega \in \bar{\Omega}, x_n + h_n Y(\omega) \geq 0\}$ and $E^* := \{\omega \in \bar{\Omega}, x^* + h^* Y(\omega) \geq 0\}$. Since $\limsup_n E_n \subset E^*$, Fatou's lemma gives

$$Q(x^* + h^* Y(\cdot) \geq 0) = E1_{E^*}(\cdot) \geq E \limsup_n 1_{E_n}(\cdot) \geq \limsup_n E1_{E_n}(\cdot) = 1,$$

so $h^* \in \mathcal{H}_{x^*}$. Since D is closed by definition we have $h^* \in D_{x^*}$ and $(x^*, h^*) \in Graph(\mathcal{D})$.

We prove now that ψ is usc on $Graph(\mathcal{D})$. The upper semicontinuity on $\mathbb{R} \times \mathbb{R}^d$ will follow immediately from Lemma 7.9. By Assumption 5.3 $x \in \mathbb{R} \rightarrow V(x, \omega)$ is usc on \mathbb{R} for all $\omega \in \bar{\Omega}$ and thus

$$\limsup_n V(\omega, x_n + h_n Y(\omega)) \leq V(\omega, x^* + h^* Y(\omega)).$$

By Lemma 5.9 for all $\omega \in \bar{\Omega}$

$$V(\omega, x_n + h_n Y(\cdot)) \leq V^+(\omega, x_n + h_n Y(\cdot)) \leq (|2x_n|^{\bar{\gamma}}K + 1)L(\omega) \leq (|2x^*|^{\bar{\gamma}}K + 2)L(\omega)$$

for n big enough. We can apply Fatou's Lemma (the *limsup* version) and ψ is usc on $Graph(\mathcal{D})$. From Lemma 5.9 it is also clear that $\psi < +\infty$ on $Graph(\mathcal{D})$. \square

We are now able to state our main result of this section.

Theorem 5.11 Assume that Assumptions 5.1, 5.3, 5.5 and 5.7 hold true. Then for all $x \geq 0$, $v(x) < \infty$ and there exists some optimal strategy $\hat{h} \in D_x$ such that

$$v(x) = E(V(\cdot, x + \hat{h}Y(\cdot))).$$

Moreover, $v : \mathbb{R} \rightarrow [-\infty, \infty)$ is non-decreasing and usc on \mathbb{R} .

Proof. Fix $x \geq 0$. We show first that $v(x) < \infty$. Indeed, using Lemma 5.9, for all $h \in D_x$,

$$E(V(\cdot, x + hY(\cdot))) \leq E(V^+(\cdot, x + hY(\cdot))) \leq ((2x)^{\bar{\gamma}}K + 1)EL(\cdot) < \infty.$$

From Lemma 5.10, $h \in \mathbb{R}^d \rightarrow E(V(\cdot, x + hY(\cdot)))$ is usc on \mathbb{R}^d and thus on D_x (recall that D_x is closed and see Lemma 7.9). Since by (19), $v(x) = \sup_{h \in D_x} E(\cdot, V(x + hY(\cdot)))$ and D_x is compact (see Lemma 5.8), applying Theorem 2.43 of Aliprantis and Border [2006] there exists some $\hat{h} \in D_x$ such that

$$v(x) = E(V(\cdot, x + \hat{h}Y(\cdot))). \quad (25)$$

We show that v is usc on $[0, +\infty)$. As previously, the upper semicontinuity on \mathbb{R} will follow immediately from Lemma 7.9. Let $(x_n)_{n \geq 0}$ be a sequence of non-negative numbers converging to some $x^* \in [0, +\infty)$. Let $\hat{h}_n \in D_{x_n}$ be the associated optimal strategies to x_n in (25). Let $(n_k)_{k \geq 1}$ be a subsequence such that $\limsup_n v(x_n) = \lim_k v(x_{n_k})$. By Lemma 5.8 $|\hat{h}_{n_k}| \leq x_{n_k}/\beta \leq (x^* + 1)/\beta$ for k big enough. So we can extract a subsequence (that we still denote by $(n_k)_{k \geq 1}$) such that there exists some \underline{h}^* with $\hat{h}_{n_k} \rightarrow \underline{h}^*$. As the sequence $(x_{n_k}, \hat{h}_{n_k})_{k \geq 1} \in \text{Graph}(\mathcal{D})$ converges to (x^*, \underline{h}^*) and $\text{Graph}(\mathcal{D})$ is closed (see Lemma 5.10), we get that $\underline{h}^* \in D_{x^*}$. Using Lemma 5.10

$$\limsup_n v(x_n) = \lim_k v(x_{n_k}) = \lim_k EV(\cdot, x_{n_k} + \hat{h}_{n_k}Y(\cdot)) \leq EV(\cdot, x^* + \underline{h}^*Y(\cdot)) \leq v(x^*),$$

where the last inequality holds true because $\underline{h}^* \in D_{x^*}$ and therefore v is usc on $[0, +\infty)$ and it is trivially non-decreasing. \square

6 Multi-period case

Proposition 6.1 *Let Assumptions 4.7, 4.8 and 4.10 hold true. Then $EU^+(\cdot, V_T^{x,\phi}(\cdot)) < \infty$ for all $x \geq 0$ and $\phi \in \Phi(x)$. This implies that $\Phi(U, x) = \Phi(x)$.*

Proof. Fix $0 \leq x \leq 1$ and let $\phi \in \Phi(x)$. Then $V_T^{x,\phi} \leq V_T^{1,\phi}$, $\phi \in \Phi(1) = \Phi(1, U)$ and $EU^+(\cdot, V_T^{x,\phi}(\cdot)) \leq EU^+(\cdot, V_T^{1,\phi}(\cdot)) < \infty$ by Assumption 4.7. Now, if $x \geq 1$ and $\phi \in \Phi(x)$ then Assumption 4.10 implies that for all $\omega \in \Omega$

$$U(\omega, V_T^{x,\phi}(\omega)) = U\left(\omega, 2x \left(\frac{1}{2} + \sum_{t=1}^T \frac{\phi_t(\omega^{t-1})}{2x} \Delta S_t(\omega^t)\right)\right) \leq (2x)^{\bar{\gamma}}K \left(U(\omega, V_T^{1, \frac{\phi}{2x}}(\omega)) + C(\omega)\right).$$

By Assumption 4.8, $\frac{\phi}{2x} \in \Phi(\frac{1}{2}) \subset \Phi(1) = \Phi(1, U)$. Thus

$$EU^+(\cdot, V_T^{x,\phi}(\cdot)) \leq (2x)^{\bar{\gamma}}K \left(EU^+(\cdot, V_T^{1, \frac{\phi}{2x}}(\cdot)) + E(C)\right) < \infty$$

using Assumption 4.7 and Assumption 4.10). We conclude that $\Phi(x) = \Phi(U, x)$. \square

We introduce now the dynamic programming procedure. First we set for all $t \in \{0, \dots, T-1\}$, $\omega^t \in \Omega^t$ and $x \geq 0$

$$\mathcal{H}_x^{t+1}(\omega^t) := \left\{ h \in \mathbb{R}^d, q_{t+1}(x + h\Delta S_{t+1}(\omega^t, \cdot)) \geq 0 | \omega^t \right\}, \quad (26)$$

$$\mathcal{D}_x^{t+1}(\omega^t) := \mathcal{H}_x^{t+1}(\omega^t) \cap D^{t+1}(\omega^t). \quad (27)$$

For $x < 0$ we set $\mathcal{H}_x^{t+1}(\omega^t) = \emptyset$.

We define for all $t \in \{0, \dots, T\}$ the following functions U_t from $\Omega^t \times \mathbb{R} \rightarrow \mathbb{R}$. Starting with $t = T$, we set for all $x \in \mathbb{R}$, all $\omega^T \in \Omega$

$$U_T(\omega^T) := U(\omega^T). \quad (28)$$

Using for $t \geq 1$ the full-measure set $\tilde{\Omega}^t \in \mathcal{F}_t$ that will be defined by induction in Propositions 6.7 and 6.8, we set for all $x \in \mathbb{R}$ and $\omega^t \in \Omega^t$

$$U_t(\omega^t, x) := (-\infty)1_{(-\infty, 0)}(x) + 1_{\tilde{\Omega}^t \times [0, +\infty)}(\omega^t, x) \sup_{h \in \mathcal{H}_x^{t+1}(\omega^t)} \int_{\Omega_{t+1}} U_{t+1}(\omega^t, \omega_{t+1}, x + h\Delta S_{t+1}(\omega^t, \omega_{t+1})) q_{t+1}(d\omega_{t+1} | \omega^t). \quad (29)$$

Finally, for $t = 0$,

$$U_0(x) := (-\infty)1_{(-\infty, 0)}(x) + 1_{[0, +\infty)}(x) \sup_{h \in \mathcal{H}_x^1} \int_{\Omega_1} U_1(\omega_1, x + h\Delta S_1(\omega_1)) P_1(d\omega_1). \quad (30)$$

Remark 6.2 We will prove by induction that U_t is well-defined (see (32)), i.e the integrals in (29) and (30) are well-defined in the generalised sense.

The P_t full measure set $\tilde{\Omega}^t$ in (29) is related to measurability issues that will be tackled in Proposition 6.9.

Lemma 6.3 *Let $0 \leq t \leq T - 1$ and H be a fixed \mathbb{R} -valued and \mathcal{F}_t -measurable random variable. Then the random sets \mathcal{H}_H^{t+1} , \mathcal{D}_H^{t+1} are closed-valued and with graph in $\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$.*

Proof. First it is clear that \mathcal{H}_H^{t+1} is closed-valued. As \mathcal{D}^{t+1} is closed-valued (see Lemma 3.3) it follows that \mathcal{D}_H^{t+1} is closed-valued as well. The fact that $\text{Graph}(\mathcal{H}_H^{t+1}) \in \mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$ follows immediately from

$$\text{Graph}(\mathcal{H}_H^{t+1}) = \left\{ (\omega^t, h) \in \Omega^t \times \mathbb{R}^d, H(\omega^t) \geq 0, q_{t+1}(\{H(\omega^t) + h\Delta S_{t+1}(\omega^t, \cdot) \geq 0\}) = 1 | \omega^t \right\},$$

and Lemma 7.7. We know from Lemma 3.3 that $\text{Graph}(\mathcal{D}^{t+1}) \in \mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$ and it follows that $\text{Graph}(\mathcal{D}_H^{t+1}) = \text{Graph}(\mathcal{D}^{t+1}) \cap \text{Graph}(\mathcal{H}_H^{t+1}) \in \mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$, too. \square

Finally we introduce

$$\begin{aligned} C_T(\omega^T) &:= C(\omega^T), \text{ for } \omega^T \in \Omega^T, \text{ where } C \text{ is defined in Assumption 4.10} \\ C_t(\omega^t) &:= \int_{\Omega_{t+1}} C_{t+1}(\omega^t, \omega_{t+1}) q_{t+1}(d\omega_{t+1} | \omega^t) \text{ for } t \in \{0, \dots, T-1\}, \omega^t \in \Omega^t. \end{aligned} \quad (31)$$

Lemma 6.4 *The functions $\omega^t \in \Omega^t \rightarrow C_t(\omega^t)$ are well-defined, non-negative (for all ω^t), \mathcal{F}_t -measurable and satisfy $E(C_t) = E(C_T) < \infty$. Furthermore, for all $t \in \{1, \dots, T\}$, there exists $\Omega_C^t \in \mathcal{F}_t$ and with $P_t(\Omega_C^t) = 1$ and such that $C_t < \infty$ on Ω_C^t . For $t = 0$ we have $C_0 < \infty$.*

Proof. We proceed by induction. Assumption 4.10 implies the statements for $t = T$. Assume now that C_{t+1} is \mathcal{F}_{t+1} -measurable, $C_{t+1} \geq 0$ and $E(C_{t+1}) = E(C_T) < \infty$. From Proposition 7.4 *i)* applied to $f = C_{t+1}$ we get that $\omega^t \rightarrow C_t(\omega^t)$ is \mathcal{F}_t -measurable and clearly $C_t(\omega^t) \geq 0$ for all ω^t . From the Fubini theorem

$$\begin{aligned} E(C_t) &= \int_{\Omega^t} \int_{\Omega_{t+1}} C_{t+1}(\omega^t, \omega_{t+1}) q_{t+1}(d\omega_{t+1} | \omega^t) P_t(d\omega^t) \\ &= \int_{\Omega^{t+1}} C_{t+1}(\omega^{t+1}) P_{t+1}(d\omega^{t+1}) = E(C_{t+1}) = E(C_T) < \infty. \end{aligned}$$

and the induction step is complete. For the second part of the lemma, we apply Lemma 7.5 to $f = C_{t+1}$ and we obtain that $\Omega_C^t := \{\omega^t \in \Omega^t, C_t(\omega^t) < \infty\} \in \mathcal{F}_t$ and $P_t(\Omega_C^t) = 1$. \square

Propositions 6.6 to 6.9 below carry out a dynamic programming procedure. They hold true under the following set of conditions. Let $1 \leq t \leq T$ be fixed.

$$U_t(\omega^t, \cdot) : \mathbb{R} \rightarrow \mathbb{R} \text{ is well-defined, non-decreasing and usc on } \mathbb{R} \text{ for all } \omega^t \in \Omega^t, \quad (32)$$

$$U_t(\cdot, \cdot) : \Omega^t \times \mathbb{R} \rightarrow \mathbb{R}\{\pm\infty\} \text{ is } \mathcal{F}_t \otimes \mathcal{B}(\mathbb{R})\text{-measurable,} \quad (33)$$

$$\int_{\Omega^t} U_t^+(\omega^t, H(\omega^{t-1}) + \xi(\omega^{t-1})\Delta S_t(\omega^t)) P_t(d\omega^t) < \infty, \quad (34)$$

for all $\xi \in \Xi_{t-1}$ and $H = x + \sum_{s=1}^{t-1} \phi_s \Delta S_s$ where $x \geq 0$, $\phi_1 \in \Xi_0, \dots, \phi_{t-1} \in \Xi_{t-2}$ and $P_t(H(\cdot) + \xi(\cdot)\Delta S_t(\cdot) \geq 0) = 1$,

$$U_t(\omega^t, \lambda x) \leq \lambda^{\bar{\gamma}} K \left(U_t \left(\omega^t, x + \frac{1}{2} \right) + C_t(\omega^t) \right), \text{ for all } \omega^t \in \Omega^t, \lambda \geq 1, x \in \mathbb{R}. \quad (35)$$

Remark 6.5 Note that from (32) and (33) we have that $-U_t$ is a $\bar{\mathcal{F}}_t$ -normal integrand (see Definition 14.27 in Rockafellar and Wets [1998] and Corollary 14.34 of Rockafellar and Wets [1998]). However to prove that this property is preserved in the dynamic programming procedure we need to show separately that (32) and (33) are true.

The next proposition is a first step in the construction of $\tilde{\Omega}^t$.

Proposition 6.6 *Let $0 \leq t \leq T - 1$ be fixed. Assume that (NA) condition holds true and that (32), (33), (34) and (35) hold true at stage $t + 1$. Then there exists $\tilde{\Omega}_1^t \in \mathcal{F}_t$ such that $P_t(\tilde{\Omega}_1^t) = 1$ and such that for all $\omega^t \in \tilde{\Omega}_1^t$ the function $(\omega_{t+1}, x) \rightarrow U_{t+1}(\omega^t, \omega_{t+1}, x)$ satisfies the assumptions of Theorem 5.11 with $\bar{\Omega} = \Omega_{t+1}$, $\mathcal{H} = \mathcal{G}_{t+1}$, $Q(\cdot) = q_{t+1}(\cdot|\omega^t)$, $Y(\cdot) = \Delta S_{t+1}(\omega^t, \cdot)$, $V(\cdot, y) = U_{t+1}(\omega^t, \cdot, y)$.*

Proof. We will review one by one the assumptions needed to apply Theorem 5.11 in the context $\bar{\Omega} = \Omega_{t+1}$, $\mathcal{H} = \mathcal{G}_{t+1}$, $Q(\cdot) = q_{t+1}(\cdot|\omega^t)$, $Y(\cdot) = \Delta S_{t+1}(\omega^t, \cdot)$, $V(\cdot, y) = U_{t+1}(\omega^t, \cdot, y)$. In the sequel we shortly call this the context $t + 1$.

From (32) at $t + 1$ for all $\omega^t \in \Omega^t$ and $\omega_{t+1} \in \Omega_{t+1}$, the function $x \in \mathbb{R} \rightarrow U_{t+1}(\omega^t, \omega_{t+1}, x)$ is non-decreasing and usc on \mathbb{R} . From (33) at $t + 1$ for all fixed $\omega^t \in \Omega^t$ and $x \in \mathbb{R}$, the function $\omega_{t+1} \in \Omega_{t+1} \rightarrow U_{t+1}(\omega^t, \omega_{t+1}, x)$ is \mathcal{G}_{t+1} -measurable and thus Assumption 5.3 is satisfied in the context $t + 1$. We move now to the assumptions that are verified for ω^t chosen in some specific P_t -full measure set. First from Lemma 3.5 for all $\omega^t \in \Omega_{NA1}^t$ we have $0 \in D^{t+1}(\omega^t)$ (recall that in Section 5 we have assume that D contains 0). From Proposition 3.6, Assumption 5.1 holds true for all $\omega^t \in \Omega_{NA}^t$ in the context $t + 1$.

We handle now Assumption 5.5 in context $t + 1$. Let $\omega^t \in \Omega_C^t$ be fixed where Ω_C^t is defined in Lemma 6.4. From (35) at $t + 1$ we have that for all $\omega_{t+1} \in \Omega_{t+1}$, $\lambda \geq 1$ and $x \in \mathbb{R}$

$$U_{t+1}(\omega^t, \omega_{t+1}, \lambda x) \leq \lambda^{\bar{\gamma}} K \left(U_{t+1} \left(\omega^t, \omega_{t+1}, x + \frac{1}{2} \right) + C_{t+1}(\omega^t, \omega_{t+1}) \right).$$

Now from Lemma 6.4 since $\omega^t \in \Omega_C^t$, we get that $\int_{\Omega_{t+1}} C_{t+1}(\omega^t, \omega_{t+1}) q_{t+1}(\omega_{t+1}|d\omega^t) = C_t(\omega^t) < \infty$ and thus Assumption 5.5 in context $t + 1$ is verified for all $\omega^t \in \Omega_C^t$. We want to show that for ω^t in some P_t full measure set to be determined and for all $h \in \mathcal{H}_1^{t+1}(\omega^t)$ we have that

$$\int_{\Omega_{t+1}} U_{t+1}^+(\omega^t, \omega_{t+1}, 1 + h\Delta S_{t+1}(\omega^t, \omega_{t+1})) q_{t+1}(d\omega_{t+1}|\omega^t) < \infty.$$

We introduce the following random set $I_1 : \Omega^t \rightarrow \mathbb{R}^d$

$$I_1(\omega^t) := \left\{ h \in \mathcal{H}_1^{t+1}(\omega^t), \int_{\Omega_{t+1}} U_{t+1}^+(\omega^t, \omega_{t+1}, 1 + h\Delta S_{t+1}(\omega^t, \omega_{t+1})) q_{t+1}(d\omega_{t+1}|\omega^t) = \infty \right\}. \quad (36)$$

Arguing by contradiction and using measurable selection arguments we will prove that $I_1(\omega^t) = \emptyset$ for P_t -almost all $\omega^t \in \Omega^t$. We show first that $\text{Graph}(I_1) \in \mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$. It is clear from (33) at $t + 1$

that $(\omega^t, \omega_{t+1}, h) \rightarrow U_{t+1}^+(\omega^t, \omega_{t+1}, 1 + h\Delta S_{t+1}(\omega^t, \omega_{t+1}))$ is $\mathcal{F}_t \otimes \mathcal{G}_{t+1} \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable. Using Proposition 7.4 ii) we get that $(\omega^t, h) \rightarrow \int_{\Omega_{t+1}} U_{t+1}^+(\omega^t, \omega_{t+1}, 1 + h\Delta S_{t+1}(\omega^t, \omega_{t+1}))q_{t+1}(d\omega_{t+1}|\omega^t)$ is $\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable (taking potentially the value $+\infty$). From Lemma 6.3, we obtain $Graph(\mathcal{H}_1^{t+1}) \in \mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$ and $Graph(I_1) \in \mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$ follows.

Applying the Projection Theorem we obtain that $\{I_1 \neq \emptyset\} \in \overline{\mathcal{F}}_t$ and using the Aumann Theorem there exists some $\overline{\mathcal{F}}_t$ -measurable $\bar{h}_1 : \{I_1 \neq \emptyset\} \rightarrow \mathbb{R}^d$ such that for all $\omega^t \in \{I_1 \neq \emptyset\}$, $\bar{h}_1(\omega^t) \in I_1(\omega^t)$. We extend \bar{h}_1 on all Ω^t by setting $\bar{h}_1(\omega^t) = 0$ on $\Omega^t \setminus \{I_1 \neq \emptyset\}$. Using Lemma 7.8 we get some \mathcal{F}_t -measurable $h_1 : \Omega^t \rightarrow \mathbb{R}^d$ and $\Omega_{I_1}^t \in \mathcal{F}_t$ such that $P_t(\Omega_{I_1}^t) = 1$ and $\Omega_{I_1}^t \subset \{\omega^t \in \Omega^t, h_1(\omega^t) = \bar{h}_1(\omega^t)\}$. Arguing as in the proof of Lemma 3.5 and using the Fubini Theorem (see Lemma 7.1) we get that

$$\begin{aligned} P_{t+1}(1 + h_1(\cdot)\Delta S_{t+1}(\cdot) \geq 0) &= \int_{\Omega^t} q_{t+1}(1 + h_1(\omega^t)\Delta S_{t+1}(\omega^t, \cdot) \geq 0|\omega^t)P_t(d\omega^t) \\ &= \int_{\Omega^t} q_{t+1}(1 + \bar{h}_1(\omega^t)\Delta S_{t+1}(\omega^t, \cdot) \geq 0|\omega^t)\bar{P}_t(d\omega^t) = 1. \end{aligned}$$

Now assume that $\bar{P}_t(\{I_1 \neq \emptyset\}) > 0$. Since $h_1 \in \Xi_t$ and $P_{t+1}(1 + h_1(\cdot)\Delta S_{t+1}(\cdot) \geq 0) = 1$ from (34) at $t + 1$ applied to $H = 1$

$$\int_{\Omega_{t+1}} U_{t+1}^+(\omega^{t+1}, 1 + h_1(\omega^t)\Delta S_{t+1}(\omega^{t+1}))P_{t+1}(d\omega^{t+1}) < \infty.$$

We argue as in Lemma 3.5 again. Let

$$\begin{aligned} \varphi_1(\omega^t) &= \int_{\Omega_{t+1}} U_{t+1}^+(\omega^t, \omega_{t+1}, 1 + h_1(\omega^t)\Delta S_{t+1}(\omega^t, \omega_{t+1}))q_{t+1}(d\omega_{t+1}|\omega^t), \\ \bar{\varphi}_1(\omega^t) &= \int_{\Omega_{t+1}} U_{t+1}^+(\omega^t, \omega_{t+1}, 1 + \bar{h}_1(\omega^t)\Delta S_{t+1}(\omega^t, \omega_{t+1}))q_{t+1}(d\omega_{t+1}|\omega^t). \end{aligned}$$

It is clear that φ_1 is \mathcal{F}_t -measurable and that $\bar{\varphi}_1$ is $\overline{\mathcal{F}}_t$ -measurable. Furthermore as $\{\omega^t \in \Omega^t, \varphi_1(\omega^t) \neq \bar{\varphi}_1(\omega^t)\} \subset \{\omega^t \in \Omega^t, h_1(\omega^t) \neq \bar{h}_1(\omega^t)\}$, $\varphi_1 = \bar{\varphi}_1$ P_t -almost surely. This implies that $\int_{\Omega^t} \bar{\varphi}_1 d\bar{P}_t = \int_{\Omega^t} \varphi_1 dP_t$ and using again the Fubini Theorem (see Lemma 7.1) we get that

$$\begin{aligned} &\int_{\Omega_{t+1}} U_{t+1}^+(\omega^{t+1}, x + h_1(\omega^t)\Delta S_{t+1}(\omega^{t+1}))P_{t+1}(d\omega^{t+1}) \\ &= \int_{\Omega^t} \varphi_1(\omega^t)P_t(d\omega^t) = \int_{\Omega^t} \bar{\varphi}_1(\omega^t)\bar{P}_t(d\omega^t) \\ &\geq \int_{\{I_1 \neq \emptyset\}} (+\infty)\bar{P}_t(d\omega^t) = +\infty. \end{aligned}$$

Therefore we must have $\bar{P}_t(\{I_1 \neq \emptyset\}) = 0$ i.e. $\bar{P}_t(\{I_1 = \emptyset\}) = 1$. Now since $\{I_1 = \emptyset\} \in \overline{\mathcal{F}}_t$ there exists $\Omega_{int}^t \subset \{I_1 = \emptyset\}$ such that $\Omega_{int}^t \in \mathcal{F}_t$ and $P_t(\Omega_{int}^t) = \bar{P}_t(\{I_1 = \emptyset\}) = 1$. For all $\omega^t \in \Omega_{int}^t$, Assumption 5.7 in the context $t + 1$ is true and we can now define $\tilde{\Omega}_1^t \subset \Omega^t$

$$\tilde{\Omega}_1^t := \Omega_{NA}^t \cap \Omega_{int}^t \cap \Omega_C^t. \quad (37)$$

It is clear that $\tilde{\Omega}_1^t \in \mathcal{F}_t$, $P_t(\tilde{\Omega}_1^t) = 1$ and the proof is complete. \square

The next proposition enables us to initialize the induction argument that will be carried on in Proposition 6.9.

Proposition 6.7 *Assume that the (NA) condition and Assumptions 4.7, 4.8 and 4.10 hold true. Then U_T satisfies (32), (33), (34) and (35) for $t = T$. We set $\tilde{\Omega}^T = \Omega$.*

Proof. As $U_T = U$, (32) for $t = T$ is true. We show that for all $\omega^T \in \Omega^T$, $x \in \mathbb{R} \rightarrow U_T(\omega^T, x)$ is right-continuous and for all $x \in \mathbb{R}$, $\omega^T \in \Omega^T \rightarrow U_T(x, \omega^T)$ is \mathcal{F}_T -measurable (this is just the second point of Definition 4.1) so that we can use Lemma 7.14 and establish (33) for $t = T$. Let $\omega^T \in \Omega^T$ be fixed. From (32) at T , $x \in \mathbb{R} \rightarrow U_T(\omega^T, x)$ is non-decreasing and usc on \mathbb{R} , thus applying Lemma 7.10 we get that $x \in \mathbb{R} \rightarrow U_T(\omega^T, x)$ is right-continuous on \mathbb{R} .

Let $\xi \in \Xi_{T-1}$ and $H = x + \sum_{t=1}^{T-1} \phi_t \Delta S_t$ where $x \geq 0$, $\phi_1 \in \Xi_0, \dots, \phi_{T-1} \in \Xi_{T-2}$ and $P_T(H(\cdot) + \xi(\cdot) \Delta S_T(\cdot) \geq 0) = 1$. Let $(\phi_i^\xi)_{1 \leq i \leq T} \in \Phi$ be defined by $\phi_T^\xi = \xi$ and $\phi_i^\xi = \phi_i$ for $1 \leq i \leq T-1$ then $V_T^{x, \phi^\xi} = H + \xi \Delta S_T$ and thus $\phi^\xi \in \Phi(x)$. Using Proposition 6.1 we get that $EU^+(\cdot, V_T^{x, \phi^\xi}(\cdot)) = EU_T^+(\cdot, H(\cdot) + \xi(\cdot) \Delta S_T(\cdot)) < \infty$ (recall that $U = U_T$). Therefore (34) is verified for $t = T$. Finally, from Assumption 4.10, (35) for $t = T$ is true. \square

The next proposition proves that if (32), (33), (34) and (35) hold true at $t+1$ then they are also true at U_t for some well chosen $\tilde{\Omega}^t$.

Proposition 6.8 *Let $0 \leq t \leq T-1$ be fixed. Assume that the (NA) condition holds true and that (32), (33), (34) and (35) are true at $t+1$ (where U_{t+1} is defined from a given $\tilde{\Omega}^{t+1}$ see (29)). Then there exists some $\tilde{\Omega}^t \in \mathcal{F}_t$ with $P_t(\tilde{\Omega}^t) = 1$ such that (32), (33), (34) and (35) are true for t .*

Moreover for all $H = x + \sum_{s=1}^t \phi_s \Delta S_s$, with $x \geq 0$ and $\phi_1 \in \Xi_0, \dots, \phi_t \in \Xi_{t-1}$, such that $P_t(H \geq 0) = 1$ there exists some $\tilde{\Omega}_H^t \in \mathcal{F}_t$ such that $P(\tilde{\Omega}_H^t) = 1$, $\tilde{\Omega}_H^t \subset \tilde{\Omega}^t$ and some $\hat{h}_{t+1}^H \in \Xi_t$ such that for all $\omega^t \in \tilde{\Omega}_H^t$, $\hat{h}_{t+1}^H(\omega^t) \in \mathcal{D}_{H(\omega^t)}^{t+1}(\omega^t)$ and

$$U_t(\omega^t, H(\omega^t)) = \int_{\Omega_{t+1}} U_{t+1}(\omega^t, \omega_{t+1}, H(\omega^t) + \hat{h}_{t+1}^H(\omega^t) \Delta S_{t+1}(\omega^t, \omega_{t+1})) q_{t+1}(d\omega_{t+1} | \omega^t). \quad (38)$$

Proof. First we define $\tilde{\Omega}^t$ and prove that (32) and (33) are true for U_t . Applying Proposition 6.6, we get that for all $\omega^t \in \tilde{\Omega}_1^t$, the function $(\omega_{t+1}, x) \rightarrow U_{t+1}(\omega^t, \omega_{t+1}, x)$ satisfies the assumptions of Lemma 5.9 and Theorem 5.11 with $\bar{\Omega} = \Omega_{t+1}$, $\mathcal{H} = \mathcal{G}_{t+1}$, $Q = q_{t+1}(\cdot | \omega^t)$, $Y(\cdot) = \Delta S_{t+1}(\omega^t, \cdot)$, $V(\cdot, y) = U_{t+1}(\omega^t, \cdot, y)$ where V is defined on $\Omega_{t+1} \times \mathbb{R}$. In particular, for $\omega^t \in \tilde{\Omega}_1^t$ and all $h \in \mathcal{H}_x^{t+1}(\omega^t)$, recalling (23) we have

$$\int_{\Omega_{t+1}} U_{t+1}^+(\omega^t, \omega_{t+1}, x + h \Delta S_{t+1}(\omega^t, \omega_{t+1})) q_{t+1}(d\omega_{t+1} | \omega^t) < \infty. \quad (39)$$

Now, we introduce $\bar{U}_t : \Omega^t \times \mathbb{R}$ defined by

$$\bar{U}_t(\omega^t, x) := (-\infty) 1_{(-\infty, 0)}(x) + 1_{[0, \infty)}(x) 1_{\tilde{\Omega}_1^t}(\omega^t) \sup_{h \in \mathcal{D}_x^{t+1}(\omega^t)} \int_{\Omega_{t+1}} U_{t+1}(\omega^t, \omega_{t+1}, x + h \Delta S_{t+1}(\omega^t, \omega_{t+1})) q_{t+1}(d\omega_{t+1} | \omega^t).$$

From (39), \bar{U}_t is well-defined (in the generalised sense). First, we prove that \bar{U}_t is $\bar{\mathcal{F}}_t \otimes \mathbb{R}$ -measurable and then we will show that this implies that U_t is $\mathcal{F}_t \otimes \mathbb{R}$ -measurable for a well chosen $\tilde{\Omega}^t$. To show that \bar{U}_t is $\bar{\mathcal{F}}_t \otimes \mathcal{B}(\mathbb{R})$ -measurable, we use Lemma 7.14 (and Remark 7.15) after having proved that it is an extended Carathéodory function (see Definition 7.13). Applying Theorem 5.11, we get that for all $\omega^t \in \tilde{\Omega}_1^t$, the function $x \in \mathbb{R} \rightarrow \bar{U}_t(\omega^t, x)$ is non-decreasing and usc on \mathbb{R} . Actually, this is true for all $\omega^t \in \Omega^t$ since outside $\tilde{\Omega}_1^t$, $x \in \mathbb{R} \rightarrow U_t(\omega^t, x)$ is constant equal to zero on $[0, \infty)$ and to $-\infty$ on $(-\infty, 0)$. Let now $\omega^t \in \Omega^t$ be fixed. As $x \in \mathbb{R} \rightarrow \bar{U}_t(\omega^t, x)$ is non-decreasing and usc on \mathbb{R} we can apply Lemma 7.10 and we get that $x \in \mathbb{R} \rightarrow \bar{U}_t(\omega^t, x)$ is right-continuous on \mathbb{R} . For $x \geq 0$ fixed, applying Lemma 6.9 with $H = x$ (here $\Omega_H^t = \tilde{\Omega}_1^t$) we obtain that $\omega^t \in \Omega^t \rightarrow \sup_{h \in \mathbb{R}^d} u_x(\omega^t, h)$ is $\bar{\mathcal{F}}_t$ -measurable. Finally, from the definitions of \bar{U}_t and u_x , we get that

$$\bar{U}_t(\omega^t, x) = (-\infty) 1_{(-\infty, 0)}(x) + 1_{[0, \infty)}(x) 1_{\tilde{\Omega}_1^t}(\omega^t) \sup_{h \in \mathbb{R}^d} u_x(\omega^t, h),$$

and this implies that $\omega^t \in \Omega^t \rightarrow \bar{U}_t(\omega^t, x)$ is $\bar{\mathcal{F}}_t$ -measurable for all $x \in \mathbb{R}$ and thus that \bar{U}_t is an extended Carathéodory function as claimed

Finally, we prove the $\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R})$ -measurability of U_t . To do that we apply Lemma 7.11 and we obtain some $\Omega_{mes}^t \in \mathcal{F}_t$ such that $P_t(\Omega_{mes}^t) = 1$ and some $\mathcal{F}_t \otimes \mathbb{R}$ -measurable $\bar{U}_t : \Omega^t \times \mathbb{R} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ such that for all $x \in \mathbb{R}$, $\{\omega^t \in \Omega^t, \bar{U}_t(\omega^t, x) \neq \tilde{U}_t(\omega^t, x)\} \subset \Omega^t \setminus \Omega_{mes}^t$. We are now in a position to define $\tilde{\Omega}^t$ and set

$$\tilde{\Omega}^t := \tilde{\Omega}_1^t \cap \Omega_{mes}^t. \quad (40)$$

We have that for all $x \in \mathbb{R}$, $\omega^t \in \Omega^t$,

$$\begin{aligned} U_t(\omega^t, x) &= (-\infty)1_{(-\infty, 0)}(x) + 1_{[0, \infty)}(x)1_{\Omega_{mes}^t}(\omega^t)1_{\tilde{\Omega}_1^t}(\omega^t) \sup_{h \in \mathcal{H}_x^{t+1}(\omega^t)} \int_{\Omega_{t+1}} U_{t+1}(\omega^t, \omega_{t+1}, x + h\Delta S_{t+1}(\omega^t, \omega_{t+1}))q_{t+1}(d\omega_{t+1}|\omega^t) \\ &= (-\infty)1_{(-\infty, 0)}(x) + 1_{[0, \infty)}(x)1_{\Omega_{mes}^t}(\omega^t)1_{\tilde{\Omega}_1^t}(\omega^t) \sup_{h \in \mathcal{D}_x^{t+1}(\omega^t)} \int_{\Omega_{t+1}} U_{t+1}(\omega^t, \omega_{t+1}, x + h\Delta S_{t+1}(\omega^t, \omega_{t+1}))q_{t+1}(d\omega_{t+1}|\omega^t) \\ &= 1_{\Omega_{mes}^t}(\omega^t)\bar{U}_t(\omega^t, x) + (-\infty)1_{\Omega^t \setminus \Omega_{mes}^t}(\omega^t)1_{(-\infty, 0)}(x) \\ &= 1_{\Omega_{mes}^t}(\omega^t)\tilde{U}_t(\omega^t, x) + (-\infty)1_{\Omega^t \setminus \Omega_{mes}^t}(\omega^t)1_{(-\infty, 0)}(x), \end{aligned}$$

and the $\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R})$ -measurability of U_t follows immediately, *i.e.* (33) is true at t . From the third equality (32) is true for t since for all $\omega^t \in \Omega^t$, $x \in \mathbb{R} \rightarrow \bar{U}_t(\omega^t, x)$ is well-defined, non-decreasing and usc on \mathbb{R} . We turn now to the assumption on asymptotic elasticity *i.e.* (35) for t . If $\omega^t \notin \tilde{\Omega}^t$, then (35) is true since $C_t(\omega^t) \geq 0$ for all ω^t . Let $\omega^t \in \tilde{\Omega}^t$ be fixed. Let $x \geq 0$, $\lambda \geq 1$, $h \in \mathbb{R}^d$ such that $q_{t+1}(\lambda x + h\Delta S_{t+1}(\omega^t, \cdot) \geq 0|\omega^t) = 1$ be fixed. By (35) for $t+1$ for all $\omega_{t+1} \in \Omega_{t+1}$, we have that

$$U_{t+1}(\omega^t, \omega_{t+1}, \lambda x + h\Delta S_{t+1}(\omega^t, \omega_{t+1})) \leq \lambda^{\bar{\gamma}} K U_{t+1}\left(\omega^t, \omega_{t+1}, x + \frac{1}{2} + \frac{h}{\lambda}\Delta S_{t+1}(\omega^t, \omega_{t+1})\right) + \lambda^{\bar{\gamma}} C_{t+1}(\omega^t, \omega_{t+1}).$$

By integrating both sides (recall (39)) we get that

$$\begin{aligned} &\int_{\Omega_{t+1}} U_{t+1}(\omega^t, \omega_{t+1}, \lambda x + h\Delta S_{t+1}(\omega^t, \omega_{t+1})) q_{t+1}(d\omega_{t+1}|\omega^t) \leq \\ &\lambda^{\bar{\gamma}} K \int_{\Omega_{t+1}} U_{t+1}\left(\omega^t, \omega_{t+1}, x + \frac{1}{2} + \frac{h}{\lambda}\Delta S_{t+1}(\omega^t, \omega_{t+1})\right) q_{t+1}(d\omega_{t+1}|\omega^t) + \lambda^{\bar{\gamma}} K \int_{\Omega_{t+1}} C_{t+1}(\omega^t, \omega_{t+1}) q_{t+1}(d\omega_{t+1}|\omega^t). \end{aligned}$$

Since $C_t(\omega^t) = \int_{\Omega_{t+1}} C_{t+1}(\omega^t, \omega_{t+1})q_{t+1}(d\omega_{t+1}|\omega^t)$ (see Lemma 6.4) and $h \in \mathcal{H}_{\lambda x}^{t+1}(\omega^t)$ implies that $\frac{h}{\lambda} \in \mathcal{H}_x^{t+1}(\omega^t) \subset \mathcal{H}_{x+\frac{1}{2}}^{t+1}(\omega^t)$, we obtain by definition of U_t (see (29)) that

$$\int_{\Omega_{t+1}} U_{t+1}(\omega^t, \omega_{t+1}, \lambda x + h\Delta S_{t+1}(\omega^t, \omega_{t+1})) q_{t+1}(d\omega_{t+1}|\omega^t) \leq \lambda^{\bar{\gamma}} K U_t\left(\omega^t, x + \frac{1}{2}\right) + \lambda^{\bar{\gamma}} K C_t(\omega^t).$$

Taking the supremum over all $h \in \mathcal{H}_{\lambda x}^{t+1}(\omega^t)$ we conclude that (35) is true for t for $x \geq 0$. If $x < 0$, then (35) is true by definition of U_t .

We now prove (38) for U_t . First, from Proposition 6.6 and Theorem 5.11 and since $\tilde{\Omega}^t \subset \tilde{\Omega}_1^t$, we have for all $\omega^t \in \tilde{\Omega}^t$ and $x \geq 0$ that there exists some $\xi^* \in \mathcal{D}_x^{t+1}(\omega^t)$ such that

$$U_t(\omega^t, x) = \int_{\Omega_{t+1}} U_{t+1}(\omega^t, \omega_{t+1}, x + \xi^* \Delta S_{t+1}(\omega^t, \omega_{t+1}))q_{t+1}(d\omega_{t+1}|\omega^t). \quad (41)$$

Let $H = x + \sum_{s=1}^{t-1} \phi_s \Delta S_s$, with $x \geq 0$ and $\phi_s \in \Xi_s$ for $s \in \{1, \dots, t-1\}$, be fixed such that $P(H \geq 0) = 1$. Let $\tilde{\Omega}_H^t := \tilde{\Omega}^t \cap \{\omega^t \in \Omega^t, H(\omega) \geq 0\}$. Then $\tilde{\Omega}_H^t \in \mathcal{F}_t$ and $P(\tilde{\Omega}_H^t) = 1$. We introduce

$$\psi_H(\omega^t) := \left\{ h \in \mathcal{D}_H^{t+1}(\omega^t) : U_t(\omega^t, H(\omega^t)) = \int_{\Omega_{t+1}} U_{t+1}(\omega^t, \omega_{t+1}, H(\omega^t) + h\Delta S_{t+1}(\omega^t, \omega_{t+1})) q_{t+1}(d\omega_{t+1}|\omega^t) \right\},$$

for $\omega^t \in \tilde{\Omega}_H^t$ and $\psi_H(\omega^t) = \emptyset$ otherwise. To prove (38) it is enough to find a \mathcal{F}_t -measurable selector for ψ_H . From the definitions of ψ_H and u_H (see (43)) we obtain that (recall that $\tilde{\Omega}_H^t \subset \tilde{\Omega}^t$ and $\tilde{\Omega}_H^t \subset \Omega_H^t$, see (40) and the definition of Ω_H^t in Lemma 6.9).

$$Graph(\psi_H) = \left\{ (\omega^t, h) \in \left(\tilde{\Omega}_H^t \times \mathbb{R}^d \right) \cap Graph(\mathcal{D}_H^{t+1}), U_t(\omega^t, H(\omega^t)) = u_H(\omega^t, h) \right\}.$$

From Lemma 6.3 we have that $\text{Graph}(\mathcal{D}_H^{t+1}) \in \mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$. We have already proved that $(\omega^t, y) \rightarrow U_t(\omega^t, y)$ is $\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R})$ -measurable and, as H is \mathcal{F}_t -measurable, we obtain that $\omega^t \rightarrow U_t(\omega^t, H(\omega^t))$ is \mathcal{F}_t -measurable. Now applying Lemma 6.9 we obtain that u_H is $\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable. The fact that $\text{Graph}(\psi_H) \in \mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$ follows immediately.

So we can apply the Projection Theorem and get that $\{\psi_H \neq \emptyset\} \in \overline{\mathcal{F}}_t$ and, using the Aumann Theorem, there exists some $\overline{\mathcal{F}}_t$ -measurable $\overline{h}_{t+1}^H : \{\psi_H \neq \emptyset\} \rightarrow \mathbb{R}^d$ such that for all $\omega^t \in \{\psi_H \neq \emptyset\}$, $\overline{h}_{t+1}^H(\omega^t) \in \psi_H(\omega^t)$. Then we extend \overline{h}_{t+1}^H on all Ω^t by setting $\overline{h}_{t+1}^H = 0$ on $\Omega^t \setminus \{\psi_H \neq \emptyset\}$. Now applying Lemma 7.8 we get some \mathcal{F}_t -measurable $\widehat{h}_{t+1}^H : \Omega^t \rightarrow \mathbb{R}^d$ and some $\overline{\Omega}_H^t \in \mathcal{F}_t$ such that $P(\overline{\Omega}_H^t) = 1$ and $\overline{\Omega}_H^t \subset \{\overline{h}_{t+1}^H = \widehat{h}_{t+1}^H\}$. We prove now that the set $\{\psi_H \neq \emptyset\}$ is of full measure. Indeed, let $\omega^t \in \widetilde{\Omega}_H^t$ be fixed. Using (41) for $x = H(\omega^t) \geq 0$, there exists $h^*(\omega^t) \in \psi_H(\omega^t)$. Therefore $\widetilde{\Omega}_H^t \subset \{\psi_H \neq \emptyset\}$ and $\overline{P}_t(\{\psi_H \neq \emptyset\}) = 1$. So for all $\omega^t \in \overline{\Omega}_H^t \cap \widetilde{\Omega}_H^t$ we have

$$\begin{aligned} U_t(\omega^t, H(\omega^t)) &= \int_{\Omega_{t+1}} U_{t+1}(\omega^t, \omega_{t+1}, H(\omega^t) + \overline{h}_{t+1}^H(\omega^t) \Delta S_{t+1}(\omega^t, \omega_{t+1})) q_{t+1}(d\omega_{t+1} | \omega^t) \\ &= \int_{\Omega_{t+1}} U_{t+1}(\omega^t, \omega_{t+1}, H(\omega^t) + \widehat{h}_{t+1}^H(\omega^t) \Delta S_{t+1}(\omega^t, \omega_{t+1})) q_{t+1}(d\omega_{t+1} | \omega^t). \end{aligned}$$

So setting

$$\widetilde{\Omega}_H^t = \widetilde{\Omega}_H^t \cap \overline{\Omega}_H^t \subset \widetilde{\Omega}^t \quad (42)$$

(38) is proved for t .

We are now left with the proof of (34) for U_t . Let $\xi \in \Xi_{t-1}$ and $H = x + \sum_{s=1}^{t-1} \phi_s \Delta S_s$ where $x \geq 0$ and $\phi_1 \in \Xi_0, \dots, \phi_{t-1} \in \Xi_{t-2}$ and such that $P_t(H(\cdot) + \xi(\cdot) \Delta S_t(\cdot) \geq 0) = 1$. We fix some $\omega^t \in \widetilde{\Omega}^t$. Let $X(\omega^t) = H(\omega^{t-1}) + \xi(\omega^{t-1}) \Delta S_t(\omega^t)$ then X is \mathcal{F}_t -measurable. We apply (38) to $X(\omega^t)$ (and $\mathcal{D}_{X(\omega^t)}^{t+1}(\omega^t)$), and we get some $\omega^t \in \Omega^t \rightarrow \widehat{h}_{t+1}(\omega^t)$ which is \mathcal{F}_t -measurable and $\widetilde{\Omega}_X^t \in \mathcal{F}_t$ such that $P_t(\widetilde{\Omega}_X^t) = 1$ and such that for all $\omega^t \in \widetilde{\Omega}_X^t$, $q_{t+1}(X(\omega^t) + \widehat{h}_{t+1}(\omega^t) \Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) = 1$ and

$$U_t(\omega^t, X(\omega^t)) = \int_{\Omega_{t+1}} U_{t+1}(\omega^t, \omega_{t+1}, X(\omega^t) + \widehat{h}_{t+1}(\omega^t) \Delta S_{t+1}(\omega^t, \omega_{t+1})) q_{t+1}(d\omega_{t+1} | \omega^t).$$

Using Jensen's Inequality

$$U_t^+(\omega^t, X(\omega^t)) \leq \int_{\Omega_{t+1}} U_{t+1}^+(\omega^t, \omega_{t+1}, X(\omega^t) + \widehat{h}_{t+1}(\omega^t) \Delta S_{t+1}(\omega^t, \omega_{t+1})) q_{t+1}(d\omega_{t+1} | \omega^t).$$

Thus as $P_t(\widetilde{\Omega}_X^t) = 1$

$$\begin{aligned} \int_{\widetilde{\Omega}_X^t} U_t^+(\omega^t, X(\omega^t)) P_t(d\omega^t) &= \int_{\Omega^t} U_t^+(\omega^t, X(\omega^t)) P_t(d\omega^t) \\ &\leq \int_{\Omega^{t+1}} U_{t+1}^+(\omega^{t+1}, X(\omega^t) + \widehat{h}_{t+1}(\omega^t) \Delta S_{t+1}(\omega^{t+1})) P_{t+1}(d\omega^{t+1}) < \infty, \end{aligned}$$

because of (34) for $t+1$ which applies since $X = x + \sum_{s=1}^{t-1} \phi_s \Delta S_s + \xi \Delta S_t$ where $x \geq 0$, $\phi_1 \in \Xi_1, \dots, \phi_{t-1} \in \Xi_{t-2}$, $\xi \in \Xi_{t-1}$ and $\widehat{h}_{t+1} \in \Xi_t$: (34) for t is proved. \square

Lemma 6.9 Fix some $0 \leq t \leq T-1$ and $x \geq 0$. Let $H := x + \sum_{s=1}^{t-1} \phi_s \Delta S_s$, where $\phi_1 \in \Xi_0, \dots, \phi_{t-1} \in \Xi_{t-2}$ and $P_t(H \geq 0) = 1$. Assume that the (NA) condition holds true and that (32), (33), (34) and (35) are

true at $t + 1$. Let $u_H : \Omega^t \times \mathbb{R}^d \rightarrow \mathbb{R} \cup \{\pm\infty\}$ be defined by

$$u_H(\omega^t, h) := \begin{cases} \int_{\Omega_{t+1}} U_{t+1}(\omega^t, \omega_{t+1}, H(\omega^t) + h\Delta S_{t+1}(\omega^t, \omega_{t+1}))q_{t+1}(d\omega_{t+1}|\omega^t), \\ \quad \text{if } (\omega^t, h) \in (\Omega_H^t \times \mathbb{R}^d) \cap \text{Graph}(\mathcal{D}_H^{t+1}), \\ -\infty \text{ if } (\omega^t, h) \notin \text{Graph}(\mathcal{D}_H^{t+1}), \\ 0 \text{ otherwise.} \end{cases} \quad (43)$$

where $\Omega_H^t := \tilde{\Omega}_1^t \cap \{\omega^t \in \Omega^t, H(\omega^t) \geq 0\}$. Then u_H is well-defined, $\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable and for all $\omega^t \in \Omega^t$, $u_H(\omega^t, \cdot)$ is usc. Moreover, $\omega^t \in \Omega^t \rightarrow \sup_{h \in \mathbb{R}^d} u_H(\omega^t, h)$ is \mathcal{F}_t -measurable.

Remark 6.10 In the proof below we will show that for $(\omega^t, h) \in (\Omega_H^t \times \mathbb{R}^d) \cap \text{Graph}(\mathcal{D}_H^{t+1})$ the integral in (43) is well-defined. Note that this is not necessarily the case for all $(\omega^t, h) \in \Omega^t \times \mathbb{R}^d$.

Proof. From (33) at $t + 1$, U_{t+1} is $\mathcal{F}_t \otimes \mathcal{G}_{t+1} \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable and since H and ΔS_{t+1} are respectively \mathcal{F}_t and \mathcal{F}_{t+1} -measurable, we obtain that $(\omega^t, \omega_{t+1}, h) \in \Omega^t \times \Omega_{t+1} \times \mathbb{R}^d \rightarrow U_{t+1}(\omega^t, \omega_{t+1}, H(\omega^t) + h\Delta S_{t+1}(\omega^t, \omega_{t+1}))$ is $\mathcal{F}_t \otimes \mathcal{G}_{t+1} \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable. We introduce

$$\tilde{u}_H : (\omega^t, h) \in (\Omega_H^t \times \mathbb{R}^d) \cap \text{Graph}(\mathcal{D}_H^{t+1}) \rightarrow \int_{\Omega_{t+1}} U_{t+1}(\omega^t, \omega_{t+1}, H(\omega^t) + h\Delta S_{t+1}(\omega^t, \omega_{t+1}))q_{t+1}(d\omega_{t+1}|\omega^t).$$

First we show that \tilde{u}_H is well-defined in the generalised sense. Indeed, let $(\omega^t, h) \in (\Omega_H^t \times \mathbb{R}^d) \cap \text{Graph}(\mathcal{D}_H^{t+1})$ be fixed. We can show as in Proposition 6.8 that (39) holds true and thus

$$\int_{\Omega_{t+1}} U_{t+1}^+(\omega^t, \omega_{t+1}, H(\omega^t) + h\Delta S_{t+1}(\omega^t, \omega_{t+1}))q_{t+1}(d\omega_{t+1}|\omega^t) < \infty,$$

So \tilde{u}_H is well-defined (but may be infinite-valued).

We now prove that u_H is $\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable. We can apply Proposition 7.4 *iv*) to $\mathcal{S} = (\Omega_H^t \times \mathbb{R}^d) \cap \text{Graph}(\mathcal{D}_H^{t+1})$, with $f(\omega^t, h, \omega_{t+1})$ equal to both $U_{t+1}^\pm(\omega^t, \omega_{t+1}, H(\omega^t) + h\Delta S_{t+1}(\omega^t, \omega_{t+1}))$, since $(\Omega_H^t \times \mathbb{R}^d) \cap \text{Graph}(\mathcal{D}_H^{t+1}) \in \mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$ (see Lemma 6.3), and both $(\omega^t, h, \omega_{t+1}) \in \Omega^t \times \mathbb{R}^d \times \Omega_{t+1} \rightarrow U_{t+1}^\pm(\omega^t, \omega_{t+1}, H(\omega^t) + h\Delta S_{t+1}(\omega^t, \omega_{t+1}))$ are $\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d) \otimes \mathcal{G}_{t+1}$ -measurable. So we obtain that \tilde{u}_H is $[\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)]_{\mathcal{S}}$ -measurable, where $[\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)]_{\mathcal{S}}$ denotes the trace sigma algebra of $\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$ on \mathcal{S} . Now we extend \tilde{u}_H to $\Omega^t \times \mathbb{R}^d$ by setting $\tilde{u}_H(\omega^t, h) = -\infty$ if $(\omega^t, h) \notin \text{Graph}(\mathcal{D}_H^{t+1})$ and $\tilde{u}_H(\omega^t, h) = 0$ if $(\omega^t, h) \in \text{Graph}(\mathcal{D}_H^{t+1})$ and $\omega^t \notin \Omega_H^t$. Since $[\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)]_{\mathcal{S}} \subset \mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$, $\Omega_H^t \in \mathcal{F}_t$ and $\text{Graph}(\mathcal{D}_H^{t+1}) \in \mathcal{F}_t \times \mathcal{B}(\mathbb{R}^d)$, this extension of \tilde{u}_H is again $\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable. As it is clear that this extension of \tilde{u}_H and u_H coincide, the measurability of u_H is proved.

We turn now to the usc property. Let $\omega^t \in \Omega_H^t \subset \tilde{\Omega}_1^t$ be fixed. Applying Proposition 6.6 to U_{t+1} and we get U_{t+1} satisfies the assumptions of Lemma 5.10 with $\bar{\Omega} = \Omega_{t+1}$, $\mathcal{H} = \mathcal{G}_{t+1}$, $Q = q_{t+1}(\cdot|\omega^t)$, $Y(\cdot) = \Delta S_{t+1}(\omega^t, \cdot)$, $V(\cdot, y) = U_{t+1}(\omega^t, \cdot, y)$ where V is defined on $\Omega_{t+1} \times \mathbb{R}$. Therefore the function $\phi_{\omega^t}(\cdot, \cdot)$ defined on $\mathbb{R} \times \mathbb{R}^d$ by

$$\phi_{\omega^t}(x, h) = \begin{cases} \int_{\Omega_{t+1}} U_{t+1}(\omega^t, \omega_{t+1}, x + h\Delta S_{t+1}(\omega^t, \omega_{t+1}))q_{t+1}(d\omega_{t+1}|\omega^t) \text{ if } x \geq 0 \text{ and } h \in D_x^{t+1}(\omega^t) \\ -\infty \text{ otherwise} \end{cases}$$

is usc on $\mathbb{R} \times \mathbb{R}^d$ (see (24)). In particular, for $x = H(\omega^t) \geq 0$ fixed, the function $h \in \mathbb{R}^d \rightarrow u_H(\omega^t, h) = \phi_{\omega^t}(H(\omega^t), h)$ is usc on \mathbb{R}^d . Now for $\omega^t \notin \Omega_H^t$, as u_H is equal to 0 if $h \in D_{H(\omega^t)}^{t+1}(\omega^t)$ and to $-\infty$ otherwise, Lemma 7.9 applies and $h \in \mathbb{R}^d \rightarrow u_H(\omega^t, h)$ is usc on all \mathbb{R}^d .

Finally, we apply Corollary 14.34 in Rockafellar and Wets [1998] and find that $-u_H$ is a $\overline{\mathcal{F}}_t$ -normal integrand³. Now from Theorem 14.37 of Rockafellar and Wets [1998], we obtain that $\omega^t \in \Omega^t \rightarrow$

³Corollary 14.34 of Rockafellar and Wets [1998] holds true only for complete σ -algebra. That is the reason why $-u_H$ is a $\overline{\mathcal{F}}_t$ -normal integrand and not a \mathcal{F}_t -normal integrand.

$\sup_{h \in \mathbb{R}^d} u_H(\omega^t, h)$ is $\overline{\mathcal{F}}_t$ -measurable and this concludes the proof. \square

Proof. of Theorem 4.16. Integrability issues. We fix some $\phi \in \Phi(x) = \Phi(U, x)$ (recall Proposition 6.1). Since Proposition 6.7 holds true, we can apply Proposition 6.8 for $t = T - 1$, and by backward induction, we can therefore apply Proposition 6.8 for all $t = T - 2, \dots, 0$. In particular, we get that (34) holds true for all $0 \leq t \leq T$. So choosing $H = V_{t-1}^{x, \phi}$ and $\xi = \phi_t$ we get that (recall Remark 4.3, from $\phi \in \Phi(x)$ we get that $P_t(V_t^{x, \phi}(\cdot) \geq 0) = 1$)

$$\int_{\Omega^t} U_t^+ \left(\omega^t, V_t^{x, \phi}(\omega^t) \right) P_t(d\omega^t) < \infty. \quad (44)$$

This implies that $\int_{\Omega^t} U_t \left(\omega^t, V_t^{x, \phi}(\omega^t) \right) P_t(d\omega^t)$ is defined in the generalised sense and that we can apply the Fubini Theorem for generalised integral (see Proposition 7.3)

$$\int_{\Omega^t} U_t \left(\omega^t, V_t^{x, \phi}(\omega^t) \right) P_t(d\omega^t) = \int_{\Omega^{t-1}} \int_{\Omega_t} U_t \left(\omega^{t-1}, \omega_t, V_t^{x, \phi}(\omega^{t-1}, \omega_t) \right) q_{t-1}(d\omega_t | \omega^{t-1}) P_{t-1}(d\omega^{t-1}). \quad (45)$$

Construction of ϕ^*

We fix some $x \geq 0$ and build our candidate for the optimal strategy by induction. We start at $t = 0$ and use (38) in Proposition 6.8 with $H = x \geq 0$. We set $\phi_1^* := \widehat{h}_1^x$ and we obtain that (recall that $\mathcal{F}_0 = \{\emptyset, \Omega^0\}$), $P_1(x + \phi_1^* \Delta S_1(\cdot) \geq 0) = 1$ and $U_0(x) = \int_{\Omega_1} U_1(\omega_1, x + \phi_1^* \Delta S_1(\omega_1)) P_1(d\omega_1)$. Assume that until some $t \geq 1$ we have found some $\phi_1^* \in \Xi_0, \dots, \phi_t^* \in \Xi_{t-1}$ and some $\overline{\Omega}^1 \in \mathcal{F}_1, \dots, \overline{\Omega}^{t-1} \in \mathcal{F}_{t-1}$ such that for all $i = 1, \dots, t-1$, $\overline{\Omega}^i \subset \widetilde{\Omega}^i$, $P_i(\overline{\Omega}^i) = 1$, for all $i = 0, \dots, t-1$, $\phi_{i+1}^*(\omega^i) \in D^{i+1}(\omega^i)$ and

$$P_t(x + \phi_1^* \Delta S_1(\omega_1) + \dots + \phi_t^*(\omega^{t-1}) \Delta S_t(\omega^{t-1}, \omega_t) \geq 0) = 1,$$

and finally, for all $\omega^t \in \overline{\Omega}^t$

$$U_{t-1} \left(\omega^{t-1}, V_{t-1}^{x, \phi^*}(\omega^{t-1}) \right) = \int_{\Omega_t} U_t \left(\omega^{t-1}, \omega_t, V_t^{x, \phi^*}(\omega^{t-1}, \omega_t) + \phi_t^*(\omega^{t-1}) \Delta S_t(\omega^{t-1}, \omega_t) \right) q_t(d\omega_t | \omega^{t-1}).$$

We apply Proposition 6.8 with $H(\cdot) = V_t^{x, \phi^*}(\cdot) = V_{t-1}^{x, \phi^*}(\cdot) + \phi_t^*(\cdot) \Delta S_t(\cdot)$ (recall that $P_t(V_t^{x, \phi^*} \geq 0) = 1$) and there exists $\overline{\Omega}^t := \widetilde{\Omega}_{V_t^{x, \phi^*}}^t \in \mathcal{F}_t$ such that $\overline{\Omega}^t \subset \widetilde{\Omega}^t$, $P_t(\overline{\Omega}^t) = 1$ and some \mathcal{F}_t -measurable $\omega^t \rightarrow \phi_{t+1}^*(\omega^t) := \widehat{h}_{t+1}^{V_t^{x, \phi^*}}(\omega^t)$ such that for all $\omega^t \in \overline{\Omega}^t$, $\phi_{t+1}^*(\omega^t) \in D^{t+1}(\omega^t)$

$$q_{t+1}(V_t^{x, \phi^*}(\omega^t) + \phi_{t+1}^*(\omega^t) \Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) = 1,$$

$$U_t \left(\omega^t, V_t^{x, \phi^*}(\omega^t) \right) = \int_{\Omega_{t+1}} U_{t+1} \left(\omega^t, \omega_{t+1}, V_{t+1}^{x, \phi^*}(\omega^t, \omega_{t+1}) + \phi_{t+1}^*(\omega^t) \Delta S_{t+1}(\omega^t, \omega_{t+1}) \right) q_{t+1}(d\omega_{t+1} | \omega^t). \quad (46)$$

Now since $P_t(\overline{\Omega}^t) = 1$, we obtain by the Fubini Theorem that

$$P_{t+1}(V_{t+1}^{x, \phi^*} \geq 0) = \int_{\Omega^t} q_{t+1}(V_t^{x, \phi^*}(\omega^t) + \phi_{t+1}^*(\omega^t) \Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) P_t(d\omega^t) = 1$$

and we can continue the recursion.

Thus, we have found $\phi^* = (\phi_t^*)_{1 \leq t \leq T}$ such that for all $t = 0, \dots, T$, $P_t(V_t^{x, \phi^*} \geq 0) = 1$, i.e $\phi^* \in \Phi(x)$. We have also found some $\overline{\Omega}^t \in \mathcal{F}_t$, such that $\overline{\Omega}^t \subset \widetilde{\Omega}^t$, $P_t(\overline{\Omega}^t) = 1$ and for all $\omega^t \in \overline{\Omega}^t$, (46) holds true for all $t = 0, \dots, T-1$. Moreover, from Proposition 6.1, $\phi^* \in \Phi(U, x)$ and we have that $E(U(V_T^{x, \phi^*})) < \infty$.

Optimality of ϕ^*

Step 1: Using (45) with $\phi = \phi^*$ and the fact that $P_{T-1}(\overline{\Omega}^{T-1}) = 1$, we get that

$$E(U(V_T^{x, \phi^*})) = \int_{\overline{\Omega}^{T-1}} \int_{\Omega_T} U_T \left(\omega^{T-1}, \omega_T, V_{T-1}^{x, \phi^*}(\omega^{T-1}, \omega_T) + \phi_T^*(\omega^{T-1}) \Delta S_T(\omega^{T-1}, \omega_T) \right) q_T(d\omega_T | \omega^{T-1}) P_{T-1}(d\omega^{T-1}).$$

Using (46) for $t = T - 1$ and again the fact that $P_{T-1}(\bar{\Omega}^{T-1}) = 1$, we have that

$$E(U(V_T^{x,\phi^*})) = \int_{\Omega^{T-1}} U_{T-1}(\omega^{T-1}, V_{T-1}^{x,\phi^*}(\omega^{T-1})) P_{T-1}(d\omega^{T-1}).$$

By backward induction, we therefore obtain that $E(U(V_T^{x,\phi^*})) = U_0(x)$. As $\phi^* \in \Phi(U, x)$, we get that $U_0(x) \leq u(x)$. So ϕ^* will be optimal if $U_0(x) \geq u(x)$.

Step 2: We fix again some $\phi \in \Phi(U, x)$ (recall Proposition 6.1). We get that $V_t^{x,\phi} \geq 0$ P_t -a.s. for all $t = 1, \dots, T$ (recall Remark 4.3). As $\phi_1 \in \mathcal{H}_x^1$ we obtain that

$$U_0(x) \geq \int_{\Omega_1} U_1(\omega_1, x + \phi_1 \Delta S_1(\omega_1)) P_1(d\omega_1).$$

As $P_2(V_1^{x,\phi} + \phi_2 \Delta S_2 \geq 0) = 1$, there exists some P_1 -full measure set $\widehat{\Omega}^1 \in \mathcal{F}_1$ such that for all $\omega_1 \in \widehat{\Omega}^1$, $q_2(V_1^{x,\phi}(\omega_1) + \phi_2(\omega_1) \Delta S_2(\omega_1, \cdot)) \geq 0 | \omega_1) = 1$ i.e $q_2(\phi_2(\omega_1) \in \mathcal{H}_{V_1^{x,\phi}(\omega_1)}^2(\omega_1) | \omega_1) = 1$ (see Lemma 7.7).

So for $\omega_1 \in \widehat{\Omega}^1$, we have that

$$U_1(\omega_1, V_1^{x,\phi}(\omega_1)) \geq \int_{\Omega_2} U_2(\omega_1, \omega_2, V_1^{x,\phi}(\omega_1) + \phi_2(\omega_1) \Delta S_1(\omega_1, \omega_2)) q_2(d\omega_2 | \omega_1). \quad (47)$$

From (44), $\int_{\Omega^2} U_2^+(\omega^2, V_2^{x,\phi}(\omega^2)) P_2(d\omega^2) < \infty$ and we can apply the Fubini Theorem (see (45)) and

$$\int_{\Omega^2} U_2(\omega^2, V_2^{x,\phi}(\omega^2)) P_2(d\omega^2) = \int_{\widehat{\Omega}^1} \int_{\Omega_2} U_2(\omega_1, \omega_2, V_1^{x,\phi}(\omega_1) + \phi_2 \Delta S_1(\omega_1, \omega_2)) q_2(d\omega_2 | \omega_1) P_1(d\omega_1).$$

Using again (44), $\int_{\Omega^1} U_1^+(\omega^1, V_1^{x,\phi}(\omega^1)) P_1(d\omega^1) < \infty$ and integrating (in the generalised sense) both side of (47) we obtain

$$\int_{\Omega^1} U_1(\omega_1, V_1^{x,\phi}(\omega_1)) P_1(d\omega_1) \geq \int_{\Omega^2} U_2(\omega^2, V_2^{x,\phi}(\omega^2)) P_2(d\omega^2).$$

Therefore $U_0(x) \geq \int_{\Omega^2} U_2(\omega^2, V_2^{x,\phi}(\omega^2)) P_2(d\omega^2)$. We can go forward since for P_2 -almost all ω^2 we have that $q_3(\phi_3(\omega^2) \in \mathcal{H}_{V_2^{x,\phi}(\omega^2)}^3(\omega^2) | \omega^2) = 1, \dots$, for P_{T-1} almost all ω^{T-1} we have that

$$q_T(\phi_T(\omega^{T-1}) \in \mathcal{H}_{V_{T-1}^{x,\phi}(\omega^{T-1})}^T(\omega^{T-1}) | \omega^{T-1}) = 1,$$

we obtain using again (44) and the Fubini Theorem (see (45)) that

$$U_0(x) \geq \int_{\Omega_1} \int_{\Omega_2} \dots \int_{\Omega_T} U(\omega^T, V_T^{x,\phi}(\omega^T)) q_T(d\omega_T | \omega^{T-1}) \dots q_2(d\omega_2 | \omega^1) P_1(d\omega_1). \quad (48)$$

So we have that $U_0(x) \geq E(U(\cdot, V_T^{x,\phi}(\cdot)))$ for any $\phi \in \Phi(U, x)$ and the proof is complete since $u(x) = E(U(\cdot, V_T^{x,\phi^*}(\cdot))) < \infty$. \square

Proof. of Theorem 4.17. To prove Theorem 4.17, we want to apply Theorem 4.16 and thus we need to establish that Assumptions 4.7 and 4.8 hold true. To do so we will prove (51) below. First we show that for all $x \geq 0$, $\phi \in \Phi(x)$ and $0 \leq t \leq T$, we have for P_t -almost all $\omega^t \in \Omega^t$

$$|V_t^{x,\phi}(\omega^t)| \leq x \prod_{s=1}^t \left(1 + \frac{|\Delta S_s(\omega^s)|}{\alpha_{s-1}(\omega^{s-1})} \right). \quad (49)$$

To do so we first fix $x \geq 0$, some $\phi = (\phi_t)_{t=1, \dots, T} \in \Phi(x)$ and $1 \leq t \leq T$. For $\omega^{t-1} \in \Omega^{t-1}$ fixed, we denote by $\phi_t^\perp(\omega^{t-1})$ the orthogonal projection of $\phi_t(\omega^{t-1})$ on $D^t(\omega^t)$. Recalling Remark 5.2 we have

$$q_t \left(\phi_t^\perp(\omega^{t-1}) \Delta S_t(\omega^{t-1}, \cdot) = \phi_t(\omega^{t-1}) \Delta S_t(\omega^{t-1}, \cdot) | \omega^{t-1} \right) = 1,$$

and thus $\phi_t^\perp(\omega^{t-1}) \in \mathcal{D}_{V_{t-1}^{x, \phi}(\omega^{t-1})}^t(\omega^{t-1})$ (see (27) for the definition of \mathcal{D}_x^t). As the NA condition holds true, Lemma 3.5 applies and $0 \in D^t(\omega^{t+1})$. We can then apply Lemma 5.8 and we obtain that

$$|\phi_t^\perp(\omega^{t-1})| \leq \frac{V_{t-1}^{x, \phi}(\omega^{t-1})}{\alpha_{t-1}(\omega^{t-1})}. \quad (50)$$

Furthermore, as it is well-known that $\omega^{t-1} \in \Omega^{t-1} \rightarrow \phi_t^\perp(\omega^{t-1})$ is \mathcal{F}_{t-1} -measurable we obtain, applying the Fubini Theorem (see Lemma 7.1), that $P_t(\phi_t^\perp \Delta S_t = \phi_t \Delta S_t) = 1$ and we denote by Ω_{EQ}^t the P_t -full measure set on which this equality is verified. We need to slightly modify the set Ω_{EQ}^t to use it for different periods. We proceed by induction. We start at $t = 1$ (recall that $\Omega^0 := \{\omega_0\}$) with Ω_{EQ}^1 . For $t = 2$ we reset, with an abuse of notation, $\Omega_{EQ}^2 = \Omega_{EQ}^2 \cap (\Omega_{EQ}^1 \times \Omega_2)$ and we reiterate the process until T . To prove (49) we proceed by induction. It is clear at $t = 0$. Fix some $t \geq 0$ and assume that (49) holds true at t . Let $\omega^{t+1} \in \Omega_{EQ}^{t+1}$, using (49) at t and (50) we get that

$$\begin{aligned} |V_{t+1}^{x, \phi}(\omega^{t+1})| &= \left| V_t^{x, \phi}(\omega^t) + \phi_{t+1}(\omega^t) \Delta S_{t+1}(\omega^{t+1}) \right| = \left| V_t^{x, \phi}(\omega^t) + \phi_{t+1}^\perp(\omega^t) \Delta S_{t+1}(\omega^{t+1}) \right| \\ &\leq \left| V_t^{x, \phi}(\omega^t) \right| \left(1 + \frac{|\Delta S_{t+1}(\omega^{t+1})|}{\alpha_t(\omega^t)} \right) \leq x \prod_{s=1}^{t+1} \left(1 + \frac{|\Delta S_s(\omega^s)|}{\alpha_{s-1}(\omega^{s-1})} \right) \end{aligned}$$

and (49) is proven for $t + 1$. It follows since for all $0 \leq s \leq t$, $|\Delta S_s| \in \mathcal{W}_s$ and $\frac{1}{\alpha_s} \in \mathcal{W}_s$ that $V_t^{x, \phi} \in \mathcal{W}_t$. We will prove that for all $\Phi \in \Phi(x)$ and ω^T in a full measure set

$$U^+(\omega^T, V_T^{x, \phi}(\omega^T)) \leq 2^{\bar{\gamma}} K \max(x, 1)^{\bar{\gamma}} \left(\prod_{s=1}^T \left(1 + \frac{|\Delta S_s(\omega^s)|}{\alpha_{s-1}(\omega^{s-1})} \right) \right)^{\bar{\gamma}} (U^+(\omega^T, 1) + C_T(\omega^T)). \quad (51)$$

Since by assumptions $EU^+(\cdot, 1) < \infty$, $EC_T < \infty$ and since for all $0 \leq t \leq T$, $|\Delta S_t| \in \mathcal{W}_t$ and $\frac{1}{\alpha_t} \in \mathcal{W}_t$, we get that $EU^+(\cdot, V_T^{x, \phi}(\cdot)) < \infty$ for all $\Phi \in \Phi(x)$ and both Assumptions 4.7 and 4.8 hold true. We prove now (51). We fix some $x \geq 0$ and $\phi \in \Phi(x)$. Then from the monotonicity of U^+ , (49), Assumption 4.10, the fact that $\prod_{s=1}^T \left(1 + \frac{|\Delta S_s(\omega^s)|}{\alpha_{s-1}(\omega^{s-1})} \right) \geq 1$, we have for all $\omega^T \in \Omega_{EQ}^T \cap \tilde{\Omega}_T$ that

$$\begin{aligned} U^+(\omega^T, V_T^{x, \phi}(\omega^T)) &\leq U^+ \left(\omega^T, \max(x, 1) \prod_{s=1}^T \left(1 + \frac{|\Delta S_s(\omega^s)|}{\alpha_{s-1}(\omega^{s-1})} \right) \right) \\ &\leq K \left(2 \max(x, 1) \prod_{s=1}^T \left(1 + \frac{|\Delta S_s(\omega^s)|}{\alpha_{s-1}(\omega^{s-1})} \right) \right)^{\bar{\gamma}} (U^+(\omega^T, 1) + C_T(\omega^T)). \end{aligned}$$

□

7 Appendix

7.1 Generalised integral and Fubini's Theorem

For ease of the reader we provide some well-known results on measure theory, stochastic kernels and integrals. The first lemma provides a version of the Fubini Theorem for non-negative functions (see

for instance to Theorem 10.7.2 in Bogachev [2007]). We then present our definition of generalised integral and provide another version of the Fubini Theorem for generalised integral (see Proposition 7.3), which is essential throughout the paper.

Let (H, \mathcal{H}) and (K, \mathcal{K}) be two measurable spaces, p be a probability measure on (H, \mathcal{H}) and q a stochastic kernel on (K, \mathcal{K}) given (H, \mathcal{H}) , i.e such that for any $h \in H$, $C \in \mathcal{K} \rightarrow q(C|h)$ is a probability measure on (K, \mathcal{K}) and for any $C \in \mathcal{K}$, $h \in H \rightarrow q(C|h)$ is \mathcal{H} -measurable. Furthermore, for any $A \in \mathcal{H} \otimes \mathcal{K}$ and any $h \in H$, the section of A along h is defined by

$$(A)_h := \{k \in K, (h, k) \in A\}. \quad (52)$$

Lemma 7.1 *Let $A \in \mathcal{H} \otimes \mathcal{K}$ be fixed. For any $h \in H$ we have $(A)_h \in \mathcal{K}$ and we define P by*

$$P(A) := \int_H \int_K 1_A(h, k) q(dk|h) p(dh) = \int_H q((A)_h | h) p(dh). \quad (53)$$

Then P is a probability measure on $(H \times K, \mathcal{H} \otimes \mathcal{K})$.

Furthermore, if $f : H \times K \rightarrow \mathbb{R}_+ \cup \{+\infty\}$ is non-negative and $\mathcal{H} \otimes \mathcal{K}$ -measurable then $h \in H \rightarrow \int_K f(h, k) q(dk|h)$ is \mathcal{H} -measurable with value in $\mathbb{R}_+ \cup \{\infty\}$ and we have

$$\int_{H \times K} f dP := \int_{H \times K} f(h, k) P(dh, dk) = \int_H \int_K f(h, k) q(dk|h) p(dh). \quad (54)$$

Proof. Let $h \in H$ be fixed. Let $\mathcal{T} = \{A \in \mathcal{H} \otimes \mathcal{K} \mid (A)_h \in \mathcal{K}\}$. It is easy to see that \mathcal{T} is a sigma algebra on $H \times K$ and is included in $\mathcal{H} \otimes \mathcal{K}$. Let $A = B \times C \in \mathcal{H} \times \mathcal{K}$ then $(A)_h = \emptyset$ if $h \notin B$ and $(A)_h = C$ if $h \in B$. Thus $(A)_h \in \mathcal{K}$ and $\mathcal{H} \times \mathcal{K} \subset \mathcal{T}$. As \mathcal{T} is a sigma-algebra, $\mathcal{H} \otimes \mathcal{K} \subset \mathcal{T}$ and $\mathcal{T} = \mathcal{H} \otimes \mathcal{K}$ follows.

We show now that

$$h \rightarrow \int_K 1_A(h, k) q(dk|h) = \int_K 1_{(A)_h}(k) q(dk|h) = q((A)_h | h)$$

is \mathcal{H} -measurable for any $A \in \mathcal{H} \otimes \mathcal{K}$.

Let $\mathcal{E} = \{A \in \mathcal{H} \otimes \mathcal{K} \mid h \in H \rightarrow q((A)_h | h) \text{ is } \mathcal{H}\text{-measurable}\}$. It is easy to see that \mathcal{E} is a sigma algebra on $H \times K$ and is included in $\mathcal{H} \otimes \mathcal{K}$. Let $A = B \times C \in \mathcal{H} \times \mathcal{K}$ then $q((A)_h | h)$ equals to 0 if $h \notin B$ and to $q(C|h)$ if $h \in B$. So by definition of $q(\cdot|h)$, $\mathcal{H} \times \mathcal{K} \subset \mathcal{E}$. As \mathcal{E} is a sigma-algebra, $\mathcal{H} \otimes \mathcal{K} \subset \mathcal{E}$ and $\mathcal{E} = \mathcal{H} \otimes \mathcal{K}$ follows. Thus the last integral in (53) is well-defined. We verify that P defines a probability measure on $(H \times K, \mathcal{H} \otimes \mathcal{K})$. It is clear that $P(\emptyset) = 0$ and $P(H \times K) = 1$. The sigma-additivity property follows from the monotone convergence theorem.

(54) is clear when $f = 1_A$ with $A \in \mathcal{H} \otimes \mathcal{K}$ and then it for linear combinations and finally for arbitrary non-negative measurable f in a standard way. \square

Definition 7.2 *Let $f : H \times K \rightarrow \mathbb{R} \cup \{\pm\infty\}$ be a $\mathcal{H} \otimes \mathcal{K}$ -measurable function. If $\int_{H \times K} f^+ dP < \infty$ or $\int_{H \times K} f^- dP < \infty$, we define the generalised integral of f by*

$$\int_{H \times K} f dP := \int_{H \times K} f^+ dP - \int_{H \times K} f^- dP.$$

Proposition 7.3 *Let $f : H \times K \rightarrow \mathbb{R} \cup \{\pm\infty\}$ be a $\mathcal{H} \otimes \mathcal{K}$ -measurable function such that $\int_{H \times K} f^+ dP < \infty$. Then, we have*

$$\int_{H \times K} f dP = \int_H \int_K f(h, k) q(dk|h) p(dh). \quad (55)$$

Proof. Using Definition 7.2 and applying Lemma 7.1 to f^+ and f^- we obtain that

$$\begin{aligned} \int_{H \times K} f dP &= \int_{H \times K} f^+ dP - \int_{H \times K} f^- dP \\ &= \int_H \int_K f^+ q(dk|h) p(dh) + \int_H \int_K f^- q(dk|h) p(dh). \end{aligned}$$

To establish (55), assume for a moment that the following linearity result have been proved: let $g_i : H \times K \rightarrow \mathbb{R} \cup \{\pm\infty\}$ be some $\mathcal{H} \otimes \mathcal{K}$ -measurable functions such that $\int_{H \times K} g_i^+ dP < \infty$ for $i = 1, 2$. Then

$$\int_H (g_1 + g_2) dp = \int_H g_1 dp + \int_H g_2 dp. \quad (56)$$

We apply (56) with $g_1(h) = \int_K f^+(h, k)q(dh|k)$ and $g_2 = -\int_K f^-(h, k)q(dh|k)$ since by Lemma 7.1,

$$\begin{aligned} \int_H g_1^+ dp &= \int_H \left(\int_K f^+(h, k)q(dh|k) \right) p(dh) \\ &= \int_{H \times K} f^+(h, k)q(dh|k)p(dh) = \int_{H \times K} f^+ dP < \infty \end{aligned}$$

and clearly $\int_H g_2^+ dp = 0 < \infty$. So we obtain that

$$\begin{aligned} &\int_H \int_K f^+(h, k)q(dk|h)p(dh) - \int_H \int_K f^-(h, k)q(dk|h)p(dh) \\ &= \int_H \left(\int_K f^+(h, k)q(dk|h) - \int_K f^-(h, k)q(dk|h) \right) p(dh) \\ &= \int_H \int_K f(h, k)q(dk|h)p(dh), \end{aligned}$$

where the second equality comes from the definition of the generalised integral of $f(h, \cdot)$ with respect to $q(\cdot|h)$ and (55) is proven.

We prove now (56). If $\int_H g_i^- dp < \infty$ for $i = 1, 2$ this is trivial. From $\int_H g_i^+ dp < \infty$ we get that $g_i^+ < \infty$ p -almost surely for $i = 1, 2$, so the sum $g_1 + g_2$ is p -almost surely well-defined, taking its value in $[-\infty, \infty)$. As $(g_1 + g_2)^+ \leq g_1^+ + g_2^+$, using the linearity of the integral for non-negative functions we get that

$$\int_H (g_1 + g_2)^+(h)p(dh) \leq \int_H g_1^+ dp + \int_H g_2^+ dp < \infty.$$

Now from

$$g_1^+ + g_2^+ - g_1^- - g_2^- = g_1 + g_2 = (g_1 + g_2)^+ - (g_1 + g_2)^-,$$

using again the linearity of the integral for non-negative functions we get that

$$\int_H (g_1 + g_2)^+ dp + \int_H g_1^- dp + \int_H g_2^- dp = \int_H (g_1 + g_2)^- dp + \int_H g_1^+ dp + \int_H g_2^+ dp.$$

Checking the different cases, i.e. $\int_H g_1^- dp = \infty$ and $\int_H g_2^- dp < \infty$ (and the opposite case) as well as $\int_H g_i^- dp = \infty$ for $i = 1, 2$ we get that (56) is true. \square

7.2 Further measure theory issues

We present now specific applications or results that are used throughout the paper. We start with four extensions of the Fubini results presented previously. As noted in Remark 6.10, the introduction of the trace sigma-algebra is the price to pay in order to avoid using the convention $\infty - \infty = -\infty$.

Proposition 7.4 Fix some $t \in \{1, \dots, T\}$.

- i) Let $f : \Omega^t \rightarrow \mathbb{R}_+ \cup \{+\infty\}$ be a non-negative \mathcal{F}_t -measurable function. Then $\omega^{t-1} \in \Omega^{t-1} \rightarrow \int_{\Omega_t} f(\omega^{t-1}, \omega_t)q_t(d\omega_t|\omega^{t-1})$ is \mathcal{F}_{t-1} -measurable with values in $\mathbb{R}_+ \cup \{+\infty\}$.
- ii) Let $f : \Omega^t \times \mathbb{R}^d \rightarrow \mathbb{R}_+ \cup \{+\infty\}$ be a non-negative $\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable function. Then $(\omega^{t-1}, h) \in \Omega^{t-1} \times \mathbb{R}^d \rightarrow \int_{\Omega_t} f(\omega^{t-1}, \omega_t, h)q_t(d\omega_t|\omega^{t-1})$ is $\mathcal{F}_{t-1} \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable with values in $\mathbb{R}_+ \cup \{+\infty\}$

- iii) Let $f : \Omega^t \rightarrow \mathbb{R}_+ \cup \{+\infty\}$ be a non-negative $\overline{\mathcal{F}}_{t-1} \otimes \mathcal{G}_t$ -measurable function. Then $\omega^{t-1} \in \Omega^{t-1} \rightarrow \int_{\Omega_t} f(\omega^{t-1}, \omega_t) q_t(d\omega_t | \omega^{t-1})$ is $\overline{\mathcal{F}}_{t-1}$ -measurable with values in $\mathbb{R}_+ \cup \{+\infty\}$.
- iv) Let $\mathcal{S} \in \mathcal{F}_{t-1} \otimes \mathcal{B}(\mathbb{R}^d)$. Introduce $[\mathcal{F}_{t-1} \otimes \mathcal{B}(\mathbb{R}^d)]_{\mathcal{S}} := \{A \cap \mathcal{S}, A \in \mathcal{F}_{t-1} \otimes \mathcal{B}(\mathbb{R}^d)\}$ the trace sigma-algebra of $\mathcal{F}_{t-1} \otimes \mathcal{B}(\mathbb{R}^d)$ on \mathcal{S} . Let $f : \Omega^{t-1} \times \mathbb{R}^d \times \Omega_t \rightarrow \mathbb{R}_+ \cup \{+\infty\}$ be a non-negative $\mathcal{F}_{t-1} \otimes \mathcal{B}(\mathbb{R}^d) \otimes \mathcal{G}_t$ -measurable function. Then $(\omega^{t-1}, h) \in \mathcal{S} \rightarrow \int_{\Omega_t} f(\omega^{t-1}, h, \omega_t) q_t(d\omega_t | \omega^{t-1})$ is $[\mathcal{F}_{t-1} \otimes \mathcal{B}(\mathbb{R}^d)]_{\mathcal{S}}$ -measurable with values in $\mathbb{R}_+ \cup \{+\infty\}$.

Proof. Statement *i*) is a direct application of Lemma 7.1 for $H = \Omega^{t-1}$, $\mathcal{H} = \mathcal{F}_{t-1}$, $K = \Omega_t$, $\mathcal{K} = \mathcal{G}_t$ and $q(\cdot|\cdot) = q_t(\cdot|\cdot)$. To prove statement *ii*), let \bar{q}_t be defined by

$$\bar{q}_t : (G, \omega^{t-1}, h) \in \mathcal{G}_t \times \Omega^{t-1} \times \mathbb{R}^d \rightarrow \bar{q}_t(G | \omega^{t-1}, h) := q_t(G | \omega^{t-1}). \quad (57)$$

It is easy to see that \bar{q}_t is a stochastic kernel on \mathcal{G}_t given $\Omega^{t-1} \times \mathbb{R}^d$ where measurability is with respect to $\mathcal{F}_{t-1} \otimes \mathcal{B}(\mathbb{R}^d)$. Statement *ii*) follows by an application of Lemma 7.1 for $H = \Omega^{t-1} \times \mathbb{R}^d$, $\mathcal{H} = \mathcal{F}_{t-1} \otimes \mathcal{B}(\mathbb{R}^d)$, $K = \Omega_t$, $\mathcal{K} = \mathcal{G}_t$ and $q(\cdot|\cdot) = \bar{q}_t(\cdot|\cdot)$. To prove statement *iii*) note that since $\mathcal{F}_{t-1} \subset \overline{\mathcal{F}}_{t-1}$ it is clear that q_t is a stochastic kernel on $(\Omega_t, \mathcal{G}_t)$ given $(\Omega^{t-1}, \overline{\mathcal{F}}_{t-1})$ (*i.e.* measurability is with respect to $\overline{\mathcal{F}}_{t-1}$). And statement *iii*) follows immediately from an application of Lemma 7.1 for $H = \Omega^{t-1}$, $\mathcal{H} = \overline{\mathcal{F}}_{t-1}$, $K = \Omega_t$, $\mathcal{K} = \mathcal{G}_t$ and $q(\cdot|\cdot) = q_t(\cdot|\cdot)$. We prove now the last statement. Let \tilde{q}_t be defined by

$$\tilde{q}_t : (G, \omega^{t-1}, h) \in \mathcal{G}_t \times \mathcal{S} \rightarrow \tilde{q}_t(G | \omega^{t-1}, h) := q_t(G | \omega^{t-1}). \quad (58)$$

\tilde{q}_t is clearly a stochastic kernel on $(\Omega_t, \mathcal{G}_t)$ given $(\mathcal{S}, [\mathcal{F}_{t-1} \otimes \mathcal{B}(\mathbb{R}^d)]_{\mathcal{S}})$. Now let f_S be the restriction of f to $\mathcal{S} \times \Omega_t$. Using similar arguments and the fact that

$$[\mathcal{F}_{t-1} \otimes \mathcal{B}(\mathbb{R}^d) \otimes \mathcal{G}_t]_{\mathcal{S} \times \Omega_t} = [\mathcal{F}_{t-1} \otimes \mathcal{B}(\mathbb{R}^d)]_{\mathcal{S}} \otimes \mathcal{G}_t, \quad (59)$$

we obtain that f_S is $[\mathcal{F}_{t-1} \otimes \mathcal{B}(\mathbb{R}^d)]_{\mathcal{S}} \otimes \mathcal{G}_t$ -measurable. Finally, statement *iv*) follows from another application of Lemma 7.1 for $H = \mathcal{S}$, $\mathcal{H} = [\mathcal{F}_{t-1} \otimes \mathcal{B}(\mathbb{R}^d)]_{\mathcal{S}}$, $K = \Omega_t$, $\mathcal{K} = \mathcal{G}_t$ and $q(\cdot|\cdot) = \tilde{q}_t(\cdot|\cdot)$. \square

Lemma 7.5 Let $f : \Omega^{t+1} \rightarrow \mathbb{R}_+ \cup \{\infty\}$ be \mathcal{F}_{t+1} -measurable, non-negative and such that $\int_{\Omega_{t+1}} f(\omega^{t+1}) P_{t+1}(d\omega^{t+1}) < \infty$. Then $\omega^t \in \Omega^t \rightarrow \int_{\Omega_{t+1}} f(\omega^t, \omega_{t+1}) q_{t+1}(d\omega_{t+1} | \omega^t)$ is \mathcal{F}_t -measurable. Furthermore, let $N^t := \{\omega^t \in \Omega^t, \int_{\Omega_{t+1}} f(\omega^t, \omega_{t+1}) q_{t+1}(d\omega_{t+1} | \omega^t) = \infty\}$. Then $N^t \in \mathcal{F}_t$ and $P_t(N^t) = 0$

Proof. The first assertion of the lemma is a direct application of *i*) of Proposition 7.4. So it is clear that $N^t \in \mathcal{F}_t$. Furthermore, applying the Fubini Theorem (see Lemma 7.1) we get that

$$\int_{\Omega^t} \int_{\Omega_{t+1}} f(\omega^t, \omega_{t+1}) q_{t+1}(d\omega_{t+1} | \omega^t) P_t(d\omega^t) = \int_{\Omega^{t+1}} f(\omega^{t+1}) P_{t+1}(d\omega^{t+1}) < \infty$$

so necessarily $P(N^t) = 0$. \square

The next lemma, loosely speaking, allows to obtain “nice” sections (*i.e.* set of full measure for a certain probability measure). We use it in the proofs of Theorem 4.17 and Lemma 7.7.

Lemma 7.6 Fix some $t \in \{1, \dots, T\}$. Let $\tilde{\Omega}^t \in \mathcal{F}_t$ such that $P_t(\tilde{\Omega}^t) = 1$ and $\tilde{\Omega}^{t-1} \in \mathcal{F}_{t-1}$ such that $P_{t-1}(\tilde{\Omega}^{t-1}) = 1$ and set

$$\overline{\Omega}^{t-1} := \left\{ \omega^{t-1} \in \tilde{\Omega}^{t-1}, q_t \left(\left(\tilde{\Omega}^t \right)_{\omega^{t-1}} | \omega^{t-1} \right) = 1 \right\}$$

see Lemma 7.1 for the definition of $\left(\tilde{\Omega}^t \right)_{\omega^{t-1}}$. Then $\overline{\Omega}^{t-1} \in \mathcal{F}_{t-1}$ and $P_t(\overline{\Omega}^{t-1}) = 1$.

Proof. From Lemma 7.1 we know $\omega^{t-1} \rightarrow q_t \left(\left(\tilde{\Omega}^t \right)_{\omega^{t-1}} | \omega^{t-1} \right)$ is \mathcal{F}_{t-1} -measurable and the fact that $\bar{\Omega}^{t-1} \in \mathcal{F}_{t-1}$ follows immediately.

Furthermore, using the Fubini Theorem (see Lemma 7.1) we have that

$$\begin{aligned} 1 &= P_t(\tilde{\Omega}^t) = \int_{\Omega^{t-1}} \int_{\Omega_t} 1_{\tilde{\Omega}^t}(\omega^{t-1}, \omega_t) q_t(d\omega_t | \omega^{t-1}) P_{t-1}(d\omega^{t-1}) \\ &= \int_{\tilde{\Omega}^{t-1}} \int_{\Omega_t} 1_{(\tilde{\Omega}^t)_{\omega^{t-1}}}(\omega_t) q_t(d\omega_t | \omega^{t-1}) P_{t-1}(d\omega^{t-1}) \\ &= \int_{\tilde{\Omega}^{t-1}} 1 \times P_{t-1}(d\omega^{t-1}) + \int_{\tilde{\Omega}^{t-1} \setminus \bar{\Omega}^{t-1}} q_t \left(\left(\tilde{\Omega}^t \right)_{\omega^{t-1}} | \omega^{t-1} \right) P_{t-1}(d\omega^{t-1}), \end{aligned}$$

where we have used for the second line the fact that $P(\tilde{\Omega}^{t-1}) = 1$.

But if $P(\tilde{\Omega}^{t-1} \setminus \bar{\Omega}^{t-1}) > 0$ then we have that, by definition of $\bar{\Omega}^{t-1}$, that

$$\int_{\tilde{\Omega}^{t-1} \setminus \bar{\Omega}^{t-1}} q_t \left(\left(\tilde{\Omega}^t \right)_{\omega^{t-1}} | \omega^{t-1} \right) P_{t-1}(d\omega^{t-1}) < P_{t-1}(\tilde{\Omega}^{t-1} \setminus \bar{\Omega}^{t-1}),$$

and thus $1 < P_{t-1}(\bar{\Omega}^{t-1}) + P_{t-1}(\tilde{\Omega}^{t-1} \setminus \bar{\Omega}^{t-1}) = 1$, which is absurd and thus $P_{t-1}(\tilde{\Omega}^{t-1} \setminus \bar{\Omega}^{t-1}) = 0$. We conclude using again that $P_{t-1}(\tilde{\Omega}^{t-1}) = 1$. \square

The following lemma is used throughout the paper. In particular, the last statement is used in the proof of the main theorem.

Lemma 7.7 *Let $0 \leq t \leq T-1$, $B \in \mathcal{B}(\mathbb{R})$, $H : \Omega^t \rightarrow \mathbb{R}$ and $h_t : \Omega^t \rightarrow \mathbb{R}^d$ be \mathcal{F}_t -measurable. Then*

$$(\omega^t, h) \in \Omega^t \times \mathbb{R}^d \rightarrow q_{t+1}(H(\omega^t) + h \Delta S_{t+1}(\omega^t, \cdot) \in B | \omega^t), \quad (60)$$

$$\omega^t \in \Omega^t \rightarrow q_{t+1}(H(\omega^t) + h_t(\omega^t) \Delta S_{t+1}(\omega^t, \cdot) \in B | \omega^t), \quad (61)$$

are respectively $\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable and \mathcal{F}_t -measurable. Furthermore, assume that

$P_{t+1}(H(\cdot) + h_t(\cdot) \Delta S_{t+1}(\cdot) \in B) = 1$, then there exists some P_t -full measure set $\bar{\Omega}^t$ such that for all $\omega^t \in \bar{\Omega}^t$, $q_{t+1}(H(\omega^t) + h_t(\omega^t) \Delta S_{t+1}(\omega^t, \cdot) \in B | \omega^t) = 1$.

Proof. As $h \in \mathbb{R}^d \rightarrow h \Delta S_{t+1}(\omega^t, \omega_{t+1})$ is continuous for all $(\omega^t, \omega_{t+1}) \in \Omega^t \times \Omega_{t+1}$ and $(\omega^t, \omega_{t+1}) \in \Omega^t \times \Omega_{t+1} \rightarrow h \Delta S_{t+1}(\omega^t, \omega_{t+1})$ is $\mathcal{F}_{t+1} = \mathcal{F}_t \otimes \mathcal{G}_{t+1}$ -measurable for all $h \in \mathbb{R}^d$ (recall that S_t and S_{t+1} are respectively \mathcal{F}_t and \mathcal{F}_{t+1} measurable by assumption), $(\omega^t, \omega_{t+1}, h) \in \Omega^t \times \Omega_{t+1} \times \mathbb{R}^d \rightarrow h \Delta S_{t+1}(\omega^t, \omega_{t+1})$ is $\mathcal{F}_t \otimes \mathcal{G}_{t+1} \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable as a Carathéodory function. As H is \mathcal{F}_t -measurable we obtain that $\psi : (\omega^t, \omega_{t+1}, h) \in \Omega^t \times \Omega_{t+1} \times \mathbb{R}^d \rightarrow H(\omega^t) + h \Delta S_{t+1}(\omega^t, \omega_{t+1})$ is also $\mathcal{F}_t \otimes \mathcal{G}_{t+1} \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable. Therefore, for any $B \in \mathcal{B}(\mathbb{R})$, $f_B : (\omega^t, \omega_{t+1}, h) \in \Omega^t \times \Omega_{t+1} \times \mathbb{R}^d \rightarrow 1_{\psi(\cdot, \cdot, \cdot) \in B}(\omega^t, \omega_{t+1}, h)$ is $\mathcal{F}_t \otimes \mathcal{G}_{t+1} \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable. We conclude using statement *i*) of Proposition 7.4 applied to f_B and (60) is proved. We prove (61) using similar arguments. Since h_t is \mathcal{F}_t -measurable, it is clear that $\psi_{h_t} : (\omega^t, \omega_{t+1}) \in \Omega^t \times \Omega_{t+1} \rightarrow H(\omega^t) + h_t(\omega^t) \Delta S_{t+1}(\omega^t, \omega_{t+1})$ is $\mathcal{F}_t \otimes \mathcal{G}_{t+1}$ -measurable. Therefore, for any $B \in \mathcal{B}(\mathbb{R})$, $f_{B, h_t} : (\omega^t, \omega_{t+1}) \in \Omega^t \times \Omega_{t+1} \rightarrow 1_{\psi_{h_t}(\cdot, \cdot) \in B}(\omega^t, \omega_{t+1})$ is $\mathcal{F}_t \otimes \mathcal{G}_{t+1}$ -measurable. We conclude applying *i*) of Proposition 7.4 to f_{B, h_t} .

For the last statement, we set

$$\tilde{\Omega}^{t+1} := \{ \omega^{t+1} = (\omega^t, \omega_{t+1}) \in \Omega^t \times \Omega_{t+1}, H(\omega^t) + h_t(\omega^t) \Delta S_{t+1}(\omega^t, \omega_{t+1}) \in B \}.$$

It is clear that $\tilde{\Omega}^{t+1} \in \mathcal{F}_{t+1}$ and that $P_{t+1}(\tilde{\Omega}^{t+1}) = 1$. We can then apply Lemma 7.6 and we obtain some P_t -full measure set $\bar{\Omega}^t$ such that for all $\omega^t \in \bar{\Omega}^t$, $q_{t+1}(H(\omega^t) + h_t(\omega^t) \Delta S_{t+1}(\omega^t, \cdot) \in B | \omega^t) = 1$. \square

Lemma 7.8 is often used in conjunction with the Aumann Theorem.

Lemma 7.8 Let $f : \Omega^t \rightarrow \mathbb{R}$ be $\overline{\mathcal{F}}_t$ -measurable. Then there exists $g : \Omega^t \rightarrow \mathbb{R}$ that is \mathcal{F}_t -measurable and such that $f = g$ P_t -almost surely, i.e there exists $\Omega_{fg}^t \in \mathcal{F}_t$ with $P_t(\Omega_{fg}^t) = 1$ and $\Omega_{fg}^t \subset \{f = g\}$.

Proof. Let $f = 1_B$ with $B \in \overline{\mathcal{F}}_t$ then $B = A \cup N$, with $A \in \mathcal{F}_t$ and $N \in \mathcal{N}_{P_t}$. Let $g = 1_A$. Then g is \mathcal{F}_t -measurable. Clearly, $\{f \neq g\} = N \in \mathcal{N}_{P_t}$, thus $f = g$ P_t a.s. By taking linear combinations, the lemma is proven for step functions using the same argument for each indicator function. Then it is always possible to approximate some $\overline{\mathcal{F}}_t$ -measurable function f by a sequence of step function $(f_n)_{n \geq 1}$. From the preceding step for all $n \geq 1$, we get some \mathcal{F}_t -measurable step functions g_n such that $f_n = g_n$ P_t -almost surely. Let $g = \limsup g_n$, g is \mathcal{F}_t -measurable and we conclude since $\{f \neq g\} \subset \cup_{n \geq 1} \{f_n \neq g_n\}$ which is again in \mathcal{N}_{P_t} . \square

Next we provide some simple but useful results on usc functions.

Lemma 7.9 Let C be a closed subset of \mathbb{R}^m for some $m \geq 1$. Let $g : \mathbb{R}^m \rightarrow \mathbb{R} \cup \{\pm\infty\}$ be such that $g = -\infty$ on $\mathbb{R}^m \setminus C$. Then g is usc on \mathbb{R}^m if and only if g is usc on C .

Proof. We prove that if g is usc on C then it is usc on \mathbb{R}^m as the reverse implication is trivial. Let $\alpha \in \mathbb{R}$ be fixed. We prove that $S_\alpha := \{x \in \mathbb{R}^m, g(x) \geq \alpha\}$ is closed in \mathbb{R}^m . Let $(x_n)_{n \geq 1} \subset S_\alpha$ converge to $x \in \mathbb{R}^m$. Then $x_n \in C$ for all $n \geq 1$ and as C is a closed set, $x \in C$. As g is usc on C , (i.e the set $\{x \in C, g(x) \geq \alpha\}$ is closed for the induced topology of \mathbb{R}^m on C) we get that $g(x) \geq \alpha$, i.e $x \in S_\alpha$ and g is usc on \mathbb{R}^m . \square

Lemma 7.10 Let $S \subset \mathbb{R}$ be a closed subset of \mathbb{R} . Let $f : \mathbb{R} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ be such that f is usc and non-decreasing on S . Then f is right-continuous on S .

Proof. Let $(x_n)_{n \geq 1} \subset S$ be a sequence converging to some x^* from above. Then $x^* \in S$ since S is closed. As $x \in S \rightarrow f(x)$ is non-decreasing, for all $n \geq 1$ we have that $f(x_n) \geq f(x^*)$ and thus $\liminf_n f(x_n) \geq f(x^*)$. Now as f is usc on S , we get that $\limsup_n f(x_n) \leq f(x^*)$. The right continuity of f on S follows immediately. \square

We now establish a useful extension of Lemma 7.8.

Lemma 7.11 Let $f : \Omega^t \times \mathbb{R} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ be an $\overline{\mathcal{F}}_t \otimes \mathcal{B}(\mathbb{R})$ -measurable function such that for all $\omega^t \in \Omega^t$, $x \in \mathbb{R} \rightarrow f(\omega^t, x)$ is usc and non-decreasing. Then, there exists some $\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R})$ -measurable function g from $\Omega^t \times \mathbb{R}$ to $\mathbb{R} \cup \{\pm\infty\}$ and some $\Omega_{mes}^t \in \mathcal{F}_t$ such that $P_t(\Omega_{mes}^t) = 1$ and $f(\omega^t, x) = g(\omega^t, x)$ for all $(\omega^t, x) \in \Omega_{mes}^t \times \mathbb{R}$.

Remark 7.12 In particular, for all $\omega^t \in \Omega_{mes}^t$, $x \in \mathbb{R} \rightarrow g(\omega^t, x)$ is usc and non-decreasing.

Proof. Let $n \geq 1$ and $k \in \mathbb{Z}$ be fixed. We apply Lemma 7.8 to $f(\cdot) = f(\cdot, \frac{k}{2^n})$ that is $\overline{\mathcal{F}}_t$ -measurable by assumption and we get some \mathcal{F}_t -measurable $g_{n,k} : \Omega^t \rightarrow \mathbb{R} \cup \{\pm\infty\}$ and some $\Omega_{n,k}^t \in \mathcal{F}_t$ such that $P_t(\Omega_{n,k}^t) = 1$ and $\Omega_{n,k}^t \subset \{\omega^t \in \Omega^t, f(\omega^t, \frac{k}{2^n}) = g_{n,k}(\omega^t)\}$. We set $\Omega_{mes}^t := \bigcap_{n \geq 1, k \in \mathbb{Z}} \Omega_{n,k}^t$. It is clear that $\Omega_{mes}^t \in \mathcal{F}_t$ and that $P_t(\Omega_{mes}^t) = 1$. Now, we define for all $n \geq 1$, $g_n : \Omega^t \times \mathbb{R} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ by

$$g_n(\omega^t, x) := \sum_{k \in \mathbb{Z}} 1_{\left(\frac{k-1}{2^n}, \frac{k}{2^n}\right]}(x) g_{n,k}(\omega^t).$$

It is clear that g_n is $\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R})$ -measurable for all $n \geq 1$. Finally, we define $g : \Omega^t \times \mathbb{R} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ by $g(\omega^t, x) := \lim_n g_n(\omega^t, x)$. Then g is again $\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R})$ -measurable and it remains to prove that $f(\omega^t, x) = g(\omega^t, x)$ for all $(\omega^t, x) \in \Omega_{mes}^t \times \mathbb{R}$. Let $(\omega^t, x) \in \Omega_{mes}^t \times \mathbb{R}$ be fixed. For all $n \geq 1$, there exists $k_n \in \mathbb{Z}$ such that $\frac{k_n-1}{2^n} < x \leq \frac{k_n}{2^n}$ and such that $g_n(\omega^t, x) = g_{n,k_n}(\omega^t) = f(\omega^t, \frac{k_n}{2^n})$. Applying Lemma 7.10 to $f(\cdot) = f(\omega^t, \cdot)$ (and $S = \mathbb{R}$), we get that $x \in \mathbb{R} \rightarrow f(\omega^t, x)$ is right-continuous on \mathbb{R} . As $(\frac{k_n}{2^n})_{n \geq 1}$ converges to x from above, it follows that $g(\omega^t, x) = \lim_n f(\omega^t, \frac{k_n}{2^n}) = f(\omega^t, x)$ and this concludes the proof. \square

Finally, we introduce the following definition.

Definition 7.13 Let S be a closed interval of \mathbb{R} . A function $f : \Omega^t \times S \rightarrow \mathbb{R}$ is an extended Carathéodory function if

- i) for all $\omega^t \in \Omega^t$, $x \in S \rightarrow f(\omega^t, x)$ is right-continuous,
- ii) for all $x \in S$, $\omega^t \in \Omega^t \rightarrow f(\omega^t, x)$ is \mathcal{F}_t -measurable.

And we prove the following lemma that is an extension of a well-known result on Carathéodory functions (see for example 4.10 in Aliprantis and Border [2006])

Lemma 7.14 Let $S \subset \mathbb{R}$ be a closed interval of \mathbb{R} and $f : \Omega^t \times S \rightarrow \mathbb{R}$ be an extended Carathéodory function. Then f is $\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R})$ -measurable.

Proof. We define for all $n \geq 1$, $f_n : \Omega^t \times \mathbb{R} \rightarrow \mathbb{R}$ by

$$f_n(\omega^t, x) := \sum_{k \in \mathbb{Z}} 1_{\left(\frac{k-1}{2^n}, \frac{k}{2^n}\right]}(x) 1_S\left(\frac{k}{2^n}\right) f(\omega^t, \frac{k}{2^n}).$$

It is clear that f_n is $\mathcal{F}_t \otimes \mathcal{B}(\mathbb{R})$ -measurable. From the right continuity of f , we can show as in the proof of Lemma 7.11 that $f(\omega^t, x) = \lim_n f_n(\omega^t, x)$ for all $(\omega^t, x) \in \Omega^t \times S$ and the proof is complete (recall that $\Omega \times S \in \mathcal{F}_t \otimes \mathcal{B}(\mathbb{R})$ as S is a closed subset of \mathbb{R}). \square

Remark 7.15 Note that we have the same result if we replace \mathcal{F}_t with $\overline{\mathcal{F}_t}$.

7.3 Proof of technical results

Finally, we provide the missing results and proofs of the paper. We start with the following results from Section 2.

Proof of Lemma 2.2. We refer to Section 6.1 of Carassus and Rásonyi [2016] for the definition and various properties of generalized conditional expectations. In particular since $E(h^+) = \int_{\Omega^t} h^+ dP_t < \infty$, $E(h|\mathcal{F}_s)$ is well-defined (in the generalised sense) for all $0 \leq s \leq t$ (see Lemma 6.2 of Carassus and Rásonyi [2016]). Similarly, from Proposition 7.3 we have that $\varphi : \Omega^s \rightarrow \mathbb{R} \cup \{\pm\infty\}$ is well-defined (in the generalised sense) and \mathcal{F}_s -measurable.

As $\varphi(X_1, \dots, X_s)$ is \mathcal{F}_s -measurable, it remains to prove that $E(gh) = E(g\varphi(X_1, \dots, X_s))$ for all $g : \Omega^s \rightarrow \mathbb{R}_+$ non-negative, \mathcal{F}_s -measurable and such that $E(gh)$ is well-defined in the generalised sense, i.e such that $E(gh)^+ < \infty$ or $E(gh)^- < \infty$. Recalling the notations of the beginning of Section 2 and using the Fubini Theorem for the third and fourth equality (see Proposition 7.3) we get that

$$\begin{aligned} E(gh) &= E(g(X_1, \dots, X_s)h(X_1, \dots, X_t)) = \int_{\Omega^T} g(\omega_1, \dots, \omega_s)h(\omega_1, \dots, \omega_t)P(d\omega^T) \\ &= \int_{\Omega^t} g(\omega_1, \dots, \omega_s)h(\omega_1, \dots, \omega_t)q_t(\omega_t|\omega^{t-1}) \dots q_{s+1}(\omega_{s+1}|\omega^s)P_s(d\omega^s) \\ &= \int_{\Omega^s} g(\omega_1, \dots, \omega_s) \left(\int_{\Omega_{s+1} \times \dots \times \Omega_t} h(\omega_1, \dots, \omega_s, \omega_{s+1}, \dots, \omega_t)q_t(\omega_t|\omega^{t-1}) \dots q_{s+1}(\omega_{s+1}|\omega^s) \right) P_s(d\omega^s) \\ &= \int_{\Omega^s} g(\omega_1, \dots, \omega_s)\varphi(\omega_1, \dots, \omega_s)P_s(d\omega^s) \\ &= E(g(X_1, \dots, X_s)\varphi(X_1, \dots, X_t)), \end{aligned}$$

which concludes the proof. \square

We give now the proof of results of Section 3.

Proof of Lemma 3.3. Clearly, for all $\omega^t \in \Omega^t$, $\tilde{D}^{t+1}(\omega^t)$ is a non-empty and closed subset of \mathbb{R}^d . We now

show that \tilde{D}^{t+1} is measurable. Let O be a fixed open set in \mathbb{R}^d and introduce

$$\mu_O : \omega^t \in \Omega^t \rightarrow \mu_O(\omega^t) := q_{t+1}(\Delta S_{t+1}(\omega^t, \cdot) \in O | \omega^t) = \int_{\Omega_{t+1}} 1_{\Delta S_{t+1}(\cdot, \cdot) \in O}(\omega^t, \omega_{t+1}) q_{t+1}(d\omega_{t+1} | \omega^t).$$

We prove that μ_O is \mathcal{F}_t -measurable. As $(\omega^t, \omega_{t+1}) \in \Omega^t \times \Omega_{t+1} \rightarrow \Delta S_{t+1}(\omega^t, \omega_{t+1})$ is $\mathcal{F}_t \otimes \mathcal{G}_{t+1}$ -measurable and $O \in \mathcal{B}(\mathbb{R}^d)$, $(\omega^t, \omega_{t+1}) \rightarrow 1_{\Delta S_{t+1}(\cdot, \cdot) \in O}(\omega^t, \omega_{t+1})$ is $\mathcal{F}_t \otimes \mathcal{G}_{t+1}$ -measurable and the result follows from Proposition 7.7.

By definition of $\tilde{D}^{t+1}(\omega^t)$ we get that

$$\{\omega^t \in \Omega^t, \tilde{D}^{t+1}(\omega^t) \cap O \neq \emptyset\} = \{\omega^t \in \Omega^t, \mu_O(\omega^t) > 0\} \in \mathcal{F}_t.$$

Clearly, D^{t+1} is a non-empty and closed-valued random set. We prove that D^{t+1} is \mathcal{F}_t -measurable. As \tilde{D}^{t+1} is \mathcal{F}_t -measurable, applying the Castaing representation (see Theorem 14.5 of Rockafellar and Wets [1998]), we obtain a countable family of \mathcal{F}_t -measurable functions $(f_n)_{n \geq 1} : \Omega^t \rightarrow \mathbb{R}^d$ such that for all $\omega^t \in \Omega^t$, $\tilde{D}^{t+1}(\omega^t) = \overline{\{f_n(\omega^t), n \geq 1\}}$. Let $\omega^t \in \Omega^t$ be fixed. It can be easily shown that

$$D^{t+1}(\omega^t) = \text{Aff}(\tilde{D}^{t+1}(\omega^t)) = \overline{\left\{ f_1(\omega^t) + \sum_{i=2}^p \lambda_i (f_i(\omega^t) - f_1(\omega^t)), (\lambda_2, \dots, \lambda_p) \in \mathbb{Q}^{p-1}, p \geq 2 \right\}}. \quad (62)$$

So, using again the Castaing representation we obtain that $D^{t+1}(\omega^t)$ is \mathcal{F}_t -measurable. From Theorem 14.8 of Rockafellar and Wets [1998], $\text{Graph}(D^{t+1}) \in \mathcal{F}_t \otimes \mathcal{B}(\mathbb{R}^d)$ (recall that D^{t+1} is closed-valued). \square

Proof of Lemma 3.4. Introduce $C^{t+1}(\omega^t) := \overline{\text{Conv}(\tilde{D}^{t+1}(\omega^t))}$ the closed convex hull generated by $\tilde{D}^{t+1}(\omega^t)$. We will prove that $0 \in C^{t+1}(\omega^t)$. Since $C^{t+1}(\omega^t) \subset D^{t+1}(\omega^t)$ by assumption, for all $h \in C^{t+1}(\omega^t) \setminus \{0\}$

$$q_{t+1}(h \Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) < 1. \quad (63)$$

Thus if we find some $h_0 \in C^{t+1}(\omega^t)$ such that $q_{t+1}(h_0 \Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) = 1$ then $h_0 = 0$. We distinguish two cases. First assume that for all $h \in \mathbb{R}^d$, $h \neq 0$, $q_{t+1}(h \Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) < 1$. Then the polar cone of $C^{t+1}(\omega^t)$, i.e the set

$$(C^{t+1}(\omega^t))^\circ := \{y \in \mathbb{R}^d, yx \leq 0, \forall x \in C^{t+1}(\omega^t)\}$$

is reduced to $\{0\}$. Indeed if this is not the case there exists $y_0 \in \mathbb{R}^d$ such that $-y_0 x \geq 0$ for all $x \in C^{t+1}(\omega^t)$. As $A := \{\omega_{t+1} \in \Omega_{t+1}, \Delta S_{t+1}(\omega^t, \omega_{t+1}) \in \tilde{D}^{t+1}(\omega^t)\} \subset \{\omega_{t+1} \in \Omega_{t+1}, -y_0 \Delta S_{t+1}(\omega^t, \omega_{t+1}) \geq 0\}$ and $q_{t+1}(A | \omega^t) = 1$ we obtain that $q_{t+1}(-y_0 \Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) = 1$ a contradiction. As $((C^{t+1}(\omega^t))^\circ)^\circ = \text{cone}(C^{t+1}(\omega^t))$ where $\text{cone}(C^{t+1}(\omega^t))$ denote the cone generated by $C^{t+1}(\omega^t)$ we get that $\text{cone}(C^{t+1}(\omega^t)) = \mathbb{R}^d$. Let $u \neq 0 \in \text{cone}(C^{t+1}(\omega^t))$ then $-u \in \text{cone}(C^{t+1}(\omega^t))$ and there exist $\lambda_1 > 0$, $\lambda_2 > 0$ and $v_1, v_2 \in C^{t+1}(\omega^t)$ such that $u = \lambda_1 v_1$ and $-u = \lambda_2 v_2$. Thus $0 = \frac{\lambda_1}{\lambda_1 + \lambda_2} v_1 + \frac{\lambda_2}{\lambda_1 + \lambda_2} v_2 \in C^{t+1}(\omega^t)$ by convexity of $C^{t+1}(\omega^t)$.

Now we assume that there exists some $h_0 \in \mathbb{R}^d$, $h_0 \neq 0$ such that $q_{t+1}(h_0 \Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) = 1$. Note that since $h_0 \in \mathbb{R}^d$ we cannot use (63). Introduce the orthogonal projection on $C^{t+1}(\omega^t)$ (recall that $C^{t+1}(\omega^t)$ is a closed convex subset of \mathbb{R}^d)

$$p : h \in \mathbb{R}^d \rightarrow p(h) \in C^{t+1}(\omega^t).$$

Then p is continuous and we have $(h - p(h))(x - p(h)) \leq 0$ for all $x \in C^{t+1}(\omega^t)$. Fix $\omega_{t+1} \in \{\omega_{t+1} \in \Omega_{t+1}, \Delta S_{t+1}(\omega^t, \omega_{t+1}) \in \tilde{D}^{t+1}(\omega^t)\} \cap \{\omega_{t+1} \in \Omega_{t+1}, h_0 \Delta S_{t+1}(\omega^t, \omega_{t+1}) \geq 0\}$ and $\lambda \geq 0$. Let $h = \lambda h_0$ and $x = \Delta S_{t+1}(\omega^t, \omega_{t+1}) \in C^{t+1}(\omega^t)$ in the previous equation, we obtain (recall that $\tilde{D}^{t+1}(\omega^t) \subset C^{t+1}(\omega^t)$)

$$\begin{aligned} 0 &\leq \lambda h_0 \Delta S_{t+1}(\omega^t, \omega_{t+1}) = (\lambda h_0 - p(\lambda h_0)) \Delta S_{t+1}(\omega^t, \omega_{t+1}) + p(\lambda h_0) \Delta S_{t+1}(\omega^t, \omega_{t+1}) \\ &\leq (\lambda h_0 - p(\lambda h_0)) p(\lambda h_0) + p(\lambda h_0) \Delta S_{t+1}(\omega^t, \omega_{t+1}). \end{aligned}$$

As this is true for all $\lambda \geq 0$ we may take the limit when λ goes to zero and use the continuity of p

$$p(0)\Delta S_{t+1}(\omega^t, \omega_{t+1}) \geq |p(0)|^2 \geq 0$$

As $q_{t+1} \left(\left\{ \omega_{t+1} \in \Omega_{t+1}, \Delta S_{t+1}(\omega^t, \omega_{t+1}) \in \tilde{D}^{t+1}(\omega^t) \right\} | \omega^t \right) = 1$ by definition of $\tilde{D}^{t+1}(\omega^t)$ and as $q_{t+1}(h_0 \Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) = 1$ as well we have obtained that

$$q_{t+1}(p(0)\Delta S_{t+1}(\omega^t, \cdot) \geq 0 | \omega^t) = 1.$$

The fact that $p(0) \in C^{t+1}(\omega^t)$ together with (63) implies that $p(0) = 0$ and $0 \in C^{t+1}(\omega^t)$ follows.

□

The following lemma has been used in the proof of Lemma 3.5. It corresponds to Lemma 2.5 of Nutz [2016]

Lemma 7.16 *Let $\omega^t \in \Omega^t$ be fixed. Recall that $L^{t+1}(\omega^t) := (D^{t+1}(\omega^t))^\perp$ is the orthogonal space of $D^{t+1}(\omega^t)$. Then for $h \in \mathbb{R}^d$ we have that*

$$q_{t+1}(h\Delta S_{t+1}(\omega^t, \cdot) = 0 | \omega^t) = 1 \iff h \in L^{t+1}(\omega^t).$$

Proof. Assume that $h \in L^{t+1}(\omega^t)$. Then $\{\omega \in \Omega_t, \Delta S_{t+1}(\omega^t, \omega) \in D^{t+1}(\omega^t)\} \subset \{\omega \in \Omega_t, h\Delta S_{t+1}(\omega^t, \omega) = 0\}$. As $q_{t+1}(\Delta S_{t+1}(\omega^t, \cdot) \in D^{t+1}(\omega^t) | \omega^t) = 1$, we conclude that $q_{t+1}(h\Delta S_{t+1}(\omega^t, \cdot) = 0 | \omega^t) = 1$. Conversely, assume that $h \notin L^{t+1}(\omega^t)$. We first show that there exists $v \in \tilde{D}^{t+1}(\omega^t)$ such that $hv \neq 0$. If not, for all $v \in \tilde{D}^{t+1}(\omega^t)$, $hv = 0$ and for any $w \in D^{t+1}(\omega^t)$ with $w = \sum_{i=1}^m \lambda_i v_i$ where $\lambda_i \in \mathbb{R}$, $\sum_{i=1}^m \lambda_i = 1$ and $v_i \in \tilde{D}^{t+1}(\omega^t)$, we get that $hw = 0$, a contradiction. Furthermore there exists an open ball centered in v with radius $\varepsilon > 0$, $B(v, \varepsilon)$, such that $hv' \neq 0$ for all $v' \in B(v, \varepsilon)$. Assume that $q_{t+1}(\Delta S_{t+1}(\omega^t, \cdot) \in B(v, \varepsilon) | \omega^t) = 0$ or equivalently that $q_{t+1}(\Delta S_{t+1}(\omega^t, \cdot) \in \mathbb{R}^d \setminus B(v, \varepsilon) | \omega^t) = 1$. By definition of the support, $\tilde{D}^{t+1}(\omega^t) \subset \mathbb{R}^d \setminus B(v, \varepsilon)$: this contradicts $v \in \tilde{D}^{t+1}(\omega^t)$. Therefore $q_{t+1}(\Delta S_{t+1}(\omega^t, \cdot) \in B(v, \varepsilon) | \omega^t) > 0$. Let $\omega \in \{\Delta S_{t+1}(\omega^t, \cdot) \in B(v, \varepsilon)\}$, then $h\Delta S_{t+1}(\omega^t, \omega) \neq 0$ i.e. $q_{t+1}(h\Delta S_{t+1}(\omega^t, \cdot) = 0 | \omega^t) < 1$. □

We prove now the following result of Section 5.

Proof of Proposition 5.9. We start with the proof of (23) when $h \in D_x$. Since D is a vectorial subspace of \mathbb{R}^d and $0 \in \mathcal{H}_x$, the affine hull of D_x is also a vector space that we denote by $\text{Aff}(D_x)$. If $x \leq 1$ we have by Assumption 5.3 that for all $\omega \in \bar{\Omega}$, $h \in D_x$,

$$V^+(\omega, x + hY(\omega)) \leq V^+(\omega, 1 + hY(\omega)). \quad (64)$$

If $x > 1$ using Assumption 5.5 (see (21) in Remark 5.6) we get that for all $\omega \in \bar{\Omega}$, $h \in D_x$

$$V^+(\omega, x + hY(\omega)) = V^+\left(2x \left(\frac{1}{2} + \frac{h}{2x}Y(\omega)\right)\right) \leq (2x)^{\bar{\gamma}}K \left(V^+\left(\omega, 1 + \frac{h}{2x}Y(\omega)\right) + C(\omega)\right). \quad (65)$$

When $\text{Dim}(\text{Aff}(D_x)) = 0$, using (64) and (65), we obtain

$$V^+(\omega, x + hY(\omega)) \leq V^+(\omega, 1) + (2x)^{\bar{\gamma}}K (V^+(\omega, 1) + C(\omega)) \leq ((2x)^{\bar{\gamma}}K + 1)(V^+(\omega, 1) + C(\omega)). \quad (66)$$

We assume now that $\text{Dim}(\text{Aff}(D_x)) > 0$. If $x = 0$ then $Y = 0$ Q -a.s. as easily seen, and by Assumption 5.3 we get that for all $\omega \in \bar{\Omega}$, $h \in D_0$,

$$V^+(\omega, 0 + hY(\omega)) \leq V^+(\omega, 1).$$

From now we assume that $x > 0$. Then as $g \in D_x$ if and only if $\frac{g}{x} \in D_1$, we have that $\text{Aff}(D_x) = \text{Aff}(D_1)$. We set $d' := \text{Dim}(\text{Aff}(D_1))$. Let $(e_1, \dots, e_{d'})$ be an orthonormal basis of $\text{Aff}(D_1)$ (which is a

sub-vector space of \mathbb{R}^d) and $\varphi : (\lambda_1, \dots, \lambda_{d'}) \in \mathbb{R}^{d'} \rightarrow \sum_{i=1}^{d'} \lambda_i e_i \in \text{Aff}(D_1)$. Then φ is an isomorphism and it is also a homeomorphism between $\mathbb{R}^{d'}$ and $\text{Aff}(D_1)$. Since D_1 is compact by Lemma 5.8, $\varphi^{-1}(D_1)$ is a compact subset of $\mathbb{R}^{d'}$. So there exists some $c \geq 0$ such that for all $h = \sum_{i=1}^{d'} \lambda_i e_i \in D_1$, $|\lambda_i| \leq c$ for all $i = 1, \dots, d'$. We complete the family of vector $(e_1, \dots, e_{d'})$ in order to obtain an orthonormal basis of \mathbb{R}^d , denoted by $(e_1, \dots, e_{d'}, e_{d'+1}, \dots, e_d)$. For all $\omega \in \Omega$, let $(y_i(\omega))_{i=1, \dots, d}$ be the coordinate of $Y(\omega)$ in this basis.

Now let $h \in D_x$ be fixed. Then $\frac{h}{2x} \in D_{\frac{1}{2}} \subset D_1$ and $\frac{h}{2x} = \sum_{i=1}^{d'} \lambda_i e_i$ for some $(\lambda_1, \dots, \lambda_{d'}) \in \mathbb{R}^{d'}$ with $|\lambda_i| \leq c$ for all $i = 1, \dots, d'$. Note that as $\frac{h}{2x} \in D_1$, $\lambda_i = 0$ for $i \geq d' + 1$. Then as (e_1, \dots, e_d) is an orthonormal basis of \mathbb{R}^d , we obtain for all $\omega \in \Omega$

$$1 + \frac{h}{2x} Y(\omega) = 1 + \sum_{i=1}^{d'} \lambda_i y_i(\omega) \leq 1 + \sum_{i=1}^{d'} |\lambda_i| |y_i(\omega)| \leq 1 + c \sum_{i=1}^{d'} |y_i(\omega)|.$$

Thus from Assumption 5.3 for all $\omega \in \bar{\Omega}$ we get that

$$V^+ \left(\omega, 1 + \frac{h}{2x} Y(\omega) \right) \leq V^+ \left(\omega, 1 + c \sum_{i=1}^{d'} |y_i(\omega)| \right).$$

We set

$$L(\cdot) := V^+ \left(\omega, 1 + c \sum_{i=1}^{d'} |y_i(\omega)| \right) 1_{d' > 0} + V^+(\cdot, 1) + C(\cdot).$$

As $d' = \text{Dim}(\text{Aff}(D_1))$ it is clear that L does not depend on x . It is also clear that L is \mathcal{H} -measurable. Then using (64), (65) and (66) we obtain that for all $\omega \in \bar{\Omega}$

$$V^+(\omega, x + hY(\omega)) \leq ((2x)^{\bar{\gamma}} K + 1)L(\omega).$$

Note that the first term in L is used in the above inequality if $x \neq 0$ and $\text{Dim}(\text{Aff}(D_x)) > 0$. The second and the third one are there for both the case of $\text{Dim}(\text{Aff}(D_x)) = 0$ and the case of $x = 0$ and $\text{Dim}(\text{Aff}(D_x)) > 0$. As by Assumptions 5.5 and 5.7, $E(V^+(\cdot, 1) + C(\cdot)) < \infty$, it remains to prove that $d' > 0$ implies $E \left(V^+ \left(\cdot, 1 + c \sum_{i=1}^{d'} |y_i(\cdot)| \right) \right) < \infty$.

Introduce W , the finite set of \mathbb{R}^d whose coordinates on $(e_1, \dots, e_{d'})$ are 1 or -1 and 0 on $(e_{d'+1}, \dots, e_d)$. Then $W \subset \text{Aff}(D_1)$ and the vectors of W will be denoted by θ^j for $j \in \{1, \dots, 2^{d'}\}$. Let θ^ω be the vector whose coordinates on $(e_1, \dots, e_{d'})$ are $(\text{sign}(y_i(\omega)))_{i=1 \dots d'}$ and 0 on $(e_{d'+1}, \dots, e_d)$. Then $\theta^\omega \in W$ and we get that

$$V^+ \left(\omega, 1 + c \sum_{i=1}^{d'} |y_i(\omega)| \right) = V^+(\omega, 1 + c \theta^\omega Y(\omega)) \leq \sum_{j=1}^{2^{d'}} V^+(\omega, 1 + c \theta^j Y(\omega)).$$

So to prove that $EL < \infty$ it is sufficient to prove that if $d' > 0$ for all $1 \leq j \leq 2^{d'}$, $EV^+(\cdot, 1 + c \theta^j Y(\cdot)) < \infty$. Recall that $\theta^j \in \text{Aff}(D_1)$. Let $ri(D_1) = \{y \in D_1, \exists \alpha > 0 \text{ s.t } \text{Aff}(D_1) \cap B(y, \alpha) \subset D_1\}$ ⁴ denote the relative interior of D_1 . As D_1 is convex and non-empty (recall $d' > 0$), $ri(D_1)$ is also non-empty and convex and we fix some $e^* \in ri(D_1)$. We prove that $\frac{e^*}{2} \in ri(D_1)$. Let $\alpha > 0$ be such that $\text{Aff}(D_1) \cap B(e^*, \alpha) \subset D_1$ and $g \in \text{Aff}(D_1) \cap B(\frac{e^*}{2}, \frac{\alpha}{2})$. Then $2g \in \text{Aff}(D_1) \cap B(e^*, \alpha)$ and thus $2g \in D_1$. As D_1 is convex and $0 \in D_1$, we get that $g \in D_1$ and $\text{Aff}(D_1) \cap B(\frac{e^*}{2}, \frac{\alpha}{2}) \subset D_1$ which proves that $\frac{e^*}{2} \in ri(D_1)$. Now let ε_j be such that $\varepsilon_j(\frac{c}{2}\theta^j - \frac{e^*}{2}) \in B(0, \frac{\alpha}{2})$. It is easy to see that one can chose $\varepsilon_j \in (0, 1)$. Then as $\bar{e}^j := \frac{e^*}{2} + \frac{\varepsilon_j}{2}(c\theta^j - e^*) \in \text{Aff}(D_1) \cap B(\frac{e^*}{2}, \frac{\alpha}{2})$ we deduce that $\bar{e}^j \in D_1$. Using (21) we obtain that for

⁴Here $B(y, \alpha)$ is the ball of \mathbb{R}^d centered at y and with radius α .

Q -almost all ω

$$\begin{aligned}
V^+(\omega, 1 + c\theta^j Y(\omega)) &= V^+(\omega, 1 + e^* Y(\omega) + (c\theta^j - e^*) Y(\omega)) \\
&\leq \left(\frac{2}{\varepsilon_j}\right)^{\bar{\gamma}} K \left[V^+ \left(\omega, \frac{\varepsilon_j}{2} (1 + e^* Y(\omega)) + \frac{\varepsilon_j}{2} (c\theta^j - e^*) Y(\omega) + \frac{1}{2} \right) + C(\omega) \right] \\
&\leq \left(\frac{2}{\varepsilon_j}\right)^{\bar{\gamma}} K \left[V^+ \left(\omega, \frac{1}{2} + \frac{e^*}{2} Y(\omega) + \frac{\varepsilon_j}{2} (c\theta^j - e^*) Y(\omega) + \frac{1}{2} \right) + C(\omega) \right] \\
&\leq \left(\frac{2}{\varepsilon_j}\right)^{\bar{\gamma}} K [V^+(\omega, 1 + \bar{e}^j Y(\omega)) + C(\omega)],
\end{aligned}$$

where the second inequality follows from the fact that $1 + e^* Y(\cdot) \geq 0$ Q -a.s and the monotonicity property of V . Note that the above inequalities are true even if $1 + c\theta^j Y(\omega) < 0$ since (21) and the monotonicity property of V hold true for all $x \in \mathbb{R}$.

From Assumption 5.7 we get that $EV^+(\cdot, 1 + \bar{e}^j Y(\cdot)) < \infty$ and Assumption 5.5 implies $EC < \infty$, therefore $EV^+(\cdot, 1 + c\theta^j Y(\cdot)) < \infty$ and (23) is proven for $h \in D_x$. Now let $h \in \mathcal{H}_x$ and h' its orthogonal projection on D , then $hY(\cdot) = h'Y(\cdot)$ Q -a.s (see Remark 5.2). It is clear that $h' \in D_x$ thus $V^+(\cdot, x + hY(\cdot)) = V^+(\cdot, x + h'Y(\cdot))$ Q -a.s and (23) is true also for $h \in \mathcal{H}_x$. \square

To conclude, the following lemma was used in the proof of Theorem 4.16.

Lemma 7.17 *Assume that (NA) holds true. Let $\phi \in \Phi$ such that $V_T^{x,\phi} \geq 0$ P -a.s, then $V_t^{x,\phi} \geq 0$ P_t -a.s.*

Proof. Assume that there is some t such that $P_t(V_t^{x,\phi} \geq 0) < 1$ or equivalently $P_t(V_t^{x,\phi} < 0) > 0$ and let $n = \sup\{t | P_t(V_t^{x,\phi} < 0) > 0\}$. Then $P_n(V_n^{x,\phi} < 0) > 0$ and for all $s \geq n + 1$, $P_s(V_s^{x,\phi} \geq 0) = 1$. Let $\Psi_s(\omega) = 0$ if $s \leq n$ and $\Psi_s(\omega) = 1_A \phi_s(\omega)$ if $s \geq n + 1$ with $A = \{V_n^\Phi < 0\}$. Then

$$V_s^{0,\Psi} = \sum_{k=1}^s \Psi_k \Delta S_k = \sum_{k=n+1}^s \Psi_k \Delta S_k = 1_A (V_s^{x,\phi} - V_n^{x,\phi})$$

If $s \geq n + 1$ $P_s(V_s^{x,\phi} \geq 0) = 1$ and on A , $-V_n^\Phi > 0$ thus $P_T(V_T^{0,\Psi} \geq 0) = 1$ and $V_T^{0,\Psi} > 0$ on A . As by the (usual) Fubini Theorem $P_T(A) = P_n(V_n^{x,\phi} < 0) > 0$, we get an arbitrage opportunity, a contradiction. \square

Acknowledgments.

L. Carassus thanks LPMA (UMR 7599) for support. M. Rásonyi was supported by the ‘‘Lendület’’ grant LP2015-6 of the Hungarian Academy of Sciences and by the NKFIH (National Research, Development and Innovation Office, Hungary) grant KH 126505.

References

- C. D. Aliprantis and K. C. Border. *Infinite Dimensional Analysis : A Hitchhiker’s Guide*. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 3rd edition, 2006.
- A. Bensoussan, A. Cadenillas, and H. K. Koo. Entrepreneurial decisions on effort and project with a nonconcave objective function. *Math. Oper. Res.*, 40(4):902–914, 2015.
- V.I. Bogachev. *Measure Theory, vol 2*. Springer-Verlag, Berlin, 2007.

- L. Carassus and M. Rásonyi. Maximization of nonconcave utility functions in discrete-time financial market models. *Math. Oper. Res.*, 41(1):146–173, 2016.
- L. Carassus, Rásonyi M., and A. M. Rodrigues. Non-concave utility maximisation on the positive real axis in discrete time. *Mathematics and Financial Economics*, 9(4):325–349, 2015.
- G. Carlier and R.-A. Dana. Optimal demand for contingent claims when agents have law invariant utilities. *Math. Finance*, 21:169–201, 2011.
- C. Castaing and M. Valadier. *Convex Analysis and Measurable Multifunctions*, volume 580. Springer, Berlin, 1977.
- R.C. Dalang, A. Morton, and W. Willinger. Equivalent martingale measures and no-arbitrage in stochastic securities market models. *Stochastics Stochastics Rep.*, 29:185–201, 1990.
- C. Dellacherie and P.-A. Meyer. *Probability and potential*. North-Holland, Amsterdam, 1979.
- H. Föllmer and A. Schied. *Stochastic Finance: An Introduction in Discrete Time*. Walter de Gruyter & Co., Berlin, 2002.
- X. He and X.Y. Zhou. Portfolio choice under cumulative prospect theory: An analytical treatment. *Management Science*, 57:315–331, 2011.
- J. Jacod and A. N. Shiryaev. Local martingales and the fundamental asset pricing theorems in the discrete-time case. *Finance Stoch.*, 2:259–273, 1998.
- H. Jin and X.Y. Zhou. Behavioural portfolio selection in continuous time. *Math. Finance*, 18:385–426, 2008.
- D. O. Kramkov and W. Schachermayer. The asymptotic elasticity of utility functions and optimal investment in incomplete markets. *Ann. Appl. Probab.*, 9:904–950, 1999.
- M. Nutz. Utility maximization under model uncertainty in discrete time. *Math. Finance*, 26(2):252–268, 2016.
- S. R. Pliska. *Introduction to mathematical finance: discrete time models*. Wiley, 1997.
- M. Rásonyi and L. Stettner. On the utility maximization problem in discrete-time financial market models. *Ann. Appl. Probab.*, 15:1367–1395, 2005.
- M. Rásonyi and L. Stettner. On the existence of optimal portfolios for the utility maximization problem in discrete time financial models. In: *Kabanov, Y.; Lipster, R.; Stoyanov, J. (Eds), From Stochastic Calculus to Mathematical Finance*, Springer., pages 589–608, 2006.
- R. T. Rockafellar and R. J.-B. Wets. *Variational analysis*, volume 317 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1998. ISBN 3-540-62772-3.
- L. C. G. Rogers. Equivalent martingale measures and no-arbitrage. *Stochastics Stochastics Rep.*, 51(1-2):41–49, 1994.
- M.-F. Sainte-Beuve. On the extension of von Neumann-Aumann’s theorem. *J. Functional Analysis*, 17(1):112–129, 1974.
- J. Sass. Portfolio optimization under transaction costs in the CRR model. *Math. Methods Oper. Res.*, 61(2):239–259, 2005.
- J. Sass and M. Smaga. FTAP in finite discrete time with transaction costs by utility maximization. *Finance Stoch.*, 18(4):805–823, 2014.