



HAL
open science

Collet, Eckmann and the bifurcation measure

Matthieu Astorg, Thomas Gauthier, Nicolae Mihalache, Gabriel Vigny

► **To cite this version:**

Matthieu Astorg, Thomas Gauthier, Nicolae Mihalache, Gabriel Vigny. Collet, Eckmann and the bifurcation measure. *Inventiones Mathematicae*, 2019, 217 (3), pp.749-797. 10.1007/s00222-019-00874-5 . hal-01888305

HAL Id: hal-01888305

<https://hal.science/hal-01888305v1>

Submitted on 4 Oct 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

COLLET, ECKMANN AND THE BIFURCATION MEASURE

by

Matthieu Astorg, Thomas Gauthier, Nicolae Mihalache & Gabriel Vigny

Abstract. — The moduli space \mathcal{M}_d of degree $d \geq 2$ rational maps can naturally be endowed with a measure μ_{bif} detecting maximal bifurcations, called the bifurcation measure. We prove that the support of the bifurcation measure μ_{bif} has positive Lebesgue measure. To do so, we establish a general sufficient condition for the conjugacy class of a rational map to belong to the support of μ_{bif} and we exhibit a large set of Collet-Eckmann rational maps which satisfy this condition. As a consequence, we get a set of Collet-Eckmann rational maps of positive Lebesgue measure which are approximated by hyperbolic rational maps.

Contents

1. Introduction.....	1
2. Basics on bifurcation currents.....	5
3. Generalized large scale condition and the bifurcation currents.....	8
4. Misiurewicz maps and the generalized large scale condition.....	13
5. The large scale condition and good Collet-Eckmann maps.....	17
6. The proofs of Theorem A and Corollary 1.....	33
References.....	34

1. Introduction

For an integer $d \geq 2$, we let Rat_d be the space of rational maps of degree d of the Riemann sphere \mathbb{P}^1 . The space Rat_d of degree d rational maps is a quasi-projective subvariety of dimension $2d + 1$. More precisely, there exists an (irreducible) variety V such that $\text{Rat}_d \simeq \mathbb{P}^{2d+1} \setminus V$. The *J-stability locus* in Rat_d is defined as the set of maps f which are structurally stable on their Julia set \mathcal{J}_f . By the seminal work [MSS] of Mañé, Sad and Sullivan, it is an open and dense subset of Rat_d . The *bifurcation locus* in Rat_d is its complement.

One can give a measurable description of this bifurcation as follows: any rational map $f \in \text{Rat}_d$ has a unique measure μ_f of maximal entropy $\log d$. The *Lyapunov exponent* of f with respect to the measure μ_f can be defined as $L(f) := \int_{\mathbb{P}^1} \log |f'| \mu_f$, where $|\cdot|$ is any hermitian metric on \mathbb{P}^1 . DeMarco [De] proved that the function $L : f \in \text{Rat}_d \mapsto L(f) \in \mathbb{R}_+$ is plurisubharmonic (*p.s.h* for short) and continuous and that the bifurcation locus is the support of the positive closed $(1, 1)$ -current $T_{\text{bif}} := dd^c L$.

Moreover, Möbius transformations act by conjugacy on Rat_d and the quotient space is an orbifold affine variety of dimension $2d - 2$ which is known as the *moduli space* \mathcal{M}_d of degree d rational maps. As the function L is invariant by conjugacy, the function L induces a *p.s.h* and continuous function $\mathcal{L} : \mathcal{M}_d \rightarrow \mathbb{R}$. The support of $dd^c \mathcal{L}$ coincides with the image of the bifurcation locus under the quotient map $\Pi : \text{Rat}_d \rightarrow \mathcal{M}_d$.

Definition 1.1. — *The bifurcation measure of the space \mathcal{M}_d is the positive measure*

$$\mu_{\text{bif}} := (dd^c \mathcal{L})^{2d-2}.$$

Notice that the support of μ_{bif} is strictly contained in the bifurcation locus. This measure has been introduced by Bassanelli and Berteloot in [BB]. Its support is exactly the accumulation set of the hyperbolic (conjugacy class of) rational maps (see [BB] and also Buff and Epstein [BE] and the second author [G2] for other dynamical characterizations). Recall that a rational map is *hyperbolic* if it is uniformly expanding on its Julia set \mathcal{J}_f , i.e. there exist $C > 0$ and $\lambda > 1$ such that for any $n \geq 1$ and any $z \in \mathcal{J}_f$, $|(f^n)'(z)| \geq C \cdot \lambda^n$. Another way to define μ_{bif} is as follows: it is the only finite measure on \mathcal{M}_d such that $\Pi^*(\mu_{\text{bif}}) = T_{\text{bif}}^{2d-2}$ as closed positive $(2d - 2, 2d - 2)$ -currents on Rat_d (see [BB, §6]).

In the family $\{z^2 + c\}_{c \in \mathbb{C}}$ of quadratic polynomials, this measure is exactly the harmonic measure of the Mandelbrot set and its support is exactly the boundary of the Mandelbrot set. One of the important questions in complex dynamics which is still open is the following.

Question. — *Does the boundary of the Mandelbrot set have positive area?*

It is natural to ask similar questions in the moduli space \mathcal{M}_d of degree d rational maps. For that, let us *naturally* endow \mathcal{M}_d with a volume form. the moduli space \mathcal{M}_d is a affine variety in some \mathbb{P}^N . Let $\omega_{\mathbb{P}^N}$ be the Fubini-Study form on \mathbb{P}^N . Its restriction defines a $(1, 1)$ form ω_d on \mathcal{M}_d and we consider the volume form on \mathcal{M}_d

$$\text{Vol}_{\mathcal{M}_d} := \omega_d^{2d-2}.$$

This is (up to renormalization) a probability measure on \mathcal{M}_d . Moreover, in any (smooth) local chart of \mathcal{M}_d it is smoothly equivalent to the Lebesgue measure, hence it is a non-degenerate volume form in \mathcal{M}_d . For the bifurcation locus, Rees showed that it has positive $\text{Vol}_{\mathcal{M}_d}$ -measure [R]. This leaves the question of the volume of the support of the measure μ_{bif} in \mathcal{M}_d .

Another major problem concerning the moduli space \mathcal{M}_d is the *Hyperbolicity Conjecture*:

Conjecture (Hyperbolicity). — *Hyperbolic rational maps are dense in Rat_d .*

In the present paper, we are interested in the following simpler but related problem: is any Collet-Eckmann rational map approximated by hyperbolic rational maps? Refining Rees' results, Aspenberg [Asp2] showed that there is a set of positive $\text{Vol}_{\mathcal{M}_d}$ -measure of (suitable) Collet-Eckmann rational map (such maps are in the bifurcation locus). A rational map $f \in \text{Rat}_d$ is *Collet-Eckmann* if the critical set $\mathcal{C}(f)$ of f is contained in \mathcal{J}_f and if there exist $\gamma, \gamma_0 > 0$ such that

$$(\text{CE}(\gamma, \gamma_0)) \quad |(f^n)'(f(c))| \geq e^{n\gamma - \gamma_0},$$

for any $c \in \mathcal{C}(f)$ and any $n \geq 0$.

Our main result is the following

Theorem A. — *Pick an integer $d \geq 2$ and let $[f] \in \text{supp}(\mu_{\text{bif}})$. Then for any open neighborhood $\Omega \subset \mathcal{M}_d$ of $[f]$, there is a set $CE_\Omega \subset \Omega \cap \text{supp}(\mu_{\text{bif}})$ of Collet-Eckmann maps with:*

$$\text{Vol}_{\mathcal{M}_d}(CE_\Omega) > 0.$$

This implies the following result related to the above conjecture.

Corollary 1. — *There exists a set of positive $\text{Vol}_{\text{Rat}_d}$ -measure of Collet-Eckmann rational maps f of Rat_d with $[f] \in \text{supp}(\mu_{\text{bif}})$ such that f is approximated by hyperbolic rational maps.*

We rely on the following strategy: we first give a very general sufficient condition for a conjugacy class of rational maps to belong to the support of the bifurcation measure (see Theorem B). Then we exhibit a large set of rational maps fulfilling this condition. Theorem A becomes a corollary of Theorem B, Theorem C and Theorem D below and of the Main Theorem of [BE].

When a rational map has simple critical points, one can follow them holomorphically as maps $c_1, \dots, c_{2d-2} : V \rightarrow \mathbb{P}^1$ where V is an open neighborhood of f in Rat_d . For any $(2d-2)$ -tuple of positive integers $\underline{n} := (n_1, \dots, n_{2d-2})$, one also can define a holomorphic map $\mathfrak{V}_{\underline{n}} : V \rightarrow (\mathbb{P}^1)^{2d-2}$ by letting

$$\mathfrak{V}_{\underline{n}}(g) := (g^{n_1}(c_1(g)), \dots, g^{n_{2d-2}}(c_{2d-2}(g))).$$

This map detects, in a certain sense, the collective stability/instability of the critical points c_1, \dots, c_{2d-2} . In the sequel, we will use the following definition in a crucial way.

Definition 1.2. — *Let $f \in \text{Rat}_d$ with simple critical values and let $\Gamma_{\underline{n}}$ denote the graph of $\mathfrak{V}_{\underline{n}}$. We say that f satisfies the large scale condition if there exist a (local) complex submanifold $\Lambda_f \subset \text{Rat}_d$ of dimension $2d-2$, a sequence $\underline{n}_k = (n_{k,1}, \dots, n_{k,2d-2})$ of $(2d-2)$ -tuples with $n_{k,j} \rightarrow +\infty$ for all j , a basis of neighborhoods $\{\Omega_k\}_{k \geq 1}$ of f in Λ_f and $\delta > 0$ such that for any k , the connected component of $\Gamma_{\underline{n}_k} \cap \Omega_k \times (\mathbb{D}(f^{n_{k,1}}(c_1(f)), \delta) \times \dots \times \mathbb{D}(f^{n_{k,2d-2}}(c_{2d-2}(f)), \delta))$ containing $(f, \mathfrak{V}_{\underline{n}_k}(f))$ is contained in $\Omega'_k \times (\mathbb{D}(f^{n_{k,1}}(c_1(f)), \delta) \times \dots \times \mathbb{D}(f^{n_{k,2d-2}}(c_{2d-2}(f)), \delta))$ for some $\Omega'_k \Subset \Omega_k$.*

This means that a rational map f satisfies the large scale condition if, infinitely many times, the map $\mathfrak{V}_{\underline{n}}$ sends an arbitrarily small neighborhood of $[f]$ in the moduli space \mathcal{M}_d to a polydisk of fixed size in $(\mathbb{P}^1)^{2d-2}$ and its graph is vertical-like near $[f]$. The first step of our proof consists in proving that the large scale condition is actually a sufficient condition for maximal bifurcations to occur at f . More precisely, we prove the following.

Theorem B. — *Pick $f \in \text{Rat}_d$. Assume that $\omega(c) \subset \mathcal{J}_f$ for all $c \in \mathcal{C}(f)$, that f has simple critical points and that f satisfies the large scale condition. Then $[f] \in \text{supp}(\mu_{\text{bif}})$.*

This condition extracts the core of the ones previously used, such as having a uniform expansion along the postcritical set, the famous Misiurewicz condition. To illustrate the interest of this condition, we show in §4 that this condition is satisfied by Misiurewicz maps. The proof drastically simplifies the proof of [G1, Theorem 1.4]. In particular, we now are able to avoid any linearization process along the critical orbits which was a crucial step in the proof given by the second author (see [G1, §5]). Indeed, we only use here the transversality and the holomorphic motion of the hyperbolic set containing the post-critical set. The proof of Theorem B is based on a phase-parameter transfer (giving a measurable version of Tan Lei's work [Ta]). For clarity, we give the proof in the easier case of one critical point in Section 3.1.

A rational map is *strongly Misiurewicz* if all its critical points are preperiodic to repelling cycles. Reformulating the main result of Aspenberg [**Asp2**] in terms of the conditions $CE(\gamma, \gamma_0)$, $CE2(\mu, \mu_0)$, $BA(\alpha)$ and $FA(\eta, \iota)$ (defined at the beginning of Section 5) gives the following.

Theorem C. — *Assume that $f \in \text{Rat}_d$ is strongly Misiurewicz, has simple critical points and is not a flexible Lattès map. Then, there exist $\mu, \mu_0, \gamma, \gamma_0 > 0$ and $\hat{\alpha} > 0$ such that for all $\alpha < \min(\frac{\gamma}{200}, \hat{\alpha})$, there exist $\hat{\eta} > 0$ and $\hat{\iota} > 0$ such that for all $\eta < \hat{\eta}$ and for all $\iota < \hat{\iota}$, the map f is a Lebesgue density point of rational maps satisfying $CE(\gamma, \gamma_0)$, $CE2(\mu, \mu_0)$, $BA(\alpha)$ and $FA(\eta, \iota)$.*

The last key ingredient to prove Theorem A can be formulated as follows (conditions (K1-6) are defined in Section 5.1).

Theorem D. — *Let $\gamma, \gamma_0, \mu, \mu_0, \eta, \kappa > 0$ and $\alpha < \gamma/200$. There exists $\iota > 0$ such that any $f \in \text{Rat}_d$ with simple critical points and satisfying $CE(\gamma, \gamma_0)$, $CE2(\mu, \mu_0)$, $BA(\alpha)$, $FA(\eta, \iota)$ and (K1-6) satisfies the large scale condition.*

To prove Theorem D, we follow Tsujii's generalization [**Ts**] of Benedicks and Carleson construction [**BC**] that we need to adapt to the complex setting. The strategy of the proof is summarized below.

- Take a Collet-Eckmann map f and a small ball in the dynamical plane centered at a critical value. That ball will go to the large scale with exponential growth and good distortion estimates under f^n by (CE) as long as its orbit stays far away from the critical set $\mathcal{C}(f)$. For each n we will choose such a starting ball B_n .
- Passages of the critical orbit near $\mathcal{C}(f)$ impose upper bounds of the size of B_n . The assertion (BA) gives lower bounds for the approach rate to $\mathcal{C}(f)$. After a close visit near $\mathcal{C}(f)$, the image is even closer to the visited critical value. Lemma 5.4 shows that the sequel of the critical orbit copies the good properties of a long prefix of the orbit of the visited critical value.
- Lemma 5.10 guarantees that just before going (again) near a critical point, we gain expansivity by (CE2), so we have restored the exponential growth and the bound for the distortion on B_n . We use a Lemma à la Măné (Proposition 5.11) for the suffix of a (finite) critical orbit that does not visit a neighborhood of $\mathcal{C}(f)$.
- Lemma 5.13 gives a large density of times n for which B_n goes to the large scale ((FA) tells that the critical orbit does not go too often near the critical set). We intersect over all the critical values and still have positive density of times for which the starting balls go to the large scale for all the critical values.
- Using Lemma 5.15 allows to bound parametric distortion on complex lines passing through f thanks to the transversality of critical relations (using again (CE) and (BA)).
- Finally, we use a result of Sibony and Wong [**SW**] and the transversality to extend the distortion to a ball on a neighborhood of f to get the large scale condition (Theorem 5.16).

Structure of the paper. — In Section 2, we recall facts on bifurcation currents. In Section 3, we define a (generalized) large scale condition and show that if a parameter satisfies it for some m then it is in the support of T_{bif}^m . In Section 4, we apply this result to k -Misiurewicz maps. In Section 5, we prove Theorems C and D. In Section 6, we prove Theorem A and a strengthened version of Corollary 1.

2. Basics on bifurcation currents

This section is devoted to the bifurcation currents. We begin with giving a description of the bifurcation currents. We then give a new formula for the higher bifurcation currents.

2.1. The bifurcation currents of critical points

Let $(f_\lambda)_{\lambda \in \Lambda}$ be a holomorphic family of rational maps equipped with $2d - 2$ marked critical points $c_1, \dots, c_{2d-2} : \Lambda \rightarrow \mathbb{P}^1$.

Definition 2.1. — *A critical point c is said to be marked if there exists a holomorphic function $c : \Lambda \rightarrow \mathbb{P}^1$ satisfying $f'_\lambda(c(\lambda)) = 0$ for every $\lambda \in \Lambda$.*

We say that the critical point c is active at $\lambda_0 \in \Lambda$ if $(f_\lambda^n(c(\lambda)))_{n \geq 0}$ is not a normal family in any neighborhood of λ_0 . Otherwise we say that c is passive at λ_0 . The activity locus of c is the set of parameters $\lambda \in \Lambda$ at which c is active.

Let us construct the bifurcation current of the critical point c_i . One can define a fibered dynamical system \hat{f} acting on $\Lambda \times \mathbb{P}^1$

$$\begin{aligned} \hat{f} : \Lambda \times \mathbb{P}^1 &\longrightarrow \Lambda \times \mathbb{P}^1 \\ (\lambda, z) &\longmapsto (\lambda, f_\lambda(z)) . \end{aligned}$$

We denote by $p_\Lambda : \Lambda \times \mathbb{P}^1 \rightarrow \Lambda$ and $p_{\mathbb{P}^1} : \Lambda \times \mathbb{P}^1 \rightarrow \mathbb{P}^1$ the respective natural projections and by $\hat{\omega} := (p_{\mathbb{P}^1})^* \omega_{\text{FS}}$, where ω_{FS} is the Fubini-Study form on \mathbb{P}^1 normalized so that $\int_{\mathbb{P}^1} \omega_{\text{FS}} = 1$. We say that a function ψ is $\hat{\omega}$ -psh if it can be locally written as the sum of a smooth function and a plurisubharmonic function (psh for short) and $dd^c \psi + \hat{\omega} \geq 0$ in the sense of currents. Then, there exists a $\hat{\omega}$ -psh function g such that

$$(1) \quad (\hat{f})^* (\hat{\omega} + dd^c g) = d \cdot (\hat{\omega} + dd^c g) .$$

Indeed, since $d^{-1}(\hat{f})^* \hat{\omega}$ and $\hat{\omega}$ are in the same cohomology class, there exists a smooth $\hat{\omega}$ -psh function u such that $d^{-1}(\hat{f})^* \hat{\omega} = \hat{\omega} + dd^c u$. Taking

$$u_n := \sum_{j=0}^{n-1} \frac{u \circ (\hat{f})^j}{d^j} ,$$

we defined $g := \lim_n u_n$. The function g is continuous and $\hat{\omega}$ -plurisubharmonic on $\Lambda \times \mathbb{P}^1$, since $\|g - u_n\|_\infty = O(d^{-n})$. The function g is the *Green function* of \hat{f} and is unique up to an additive constant. We shall use the following notation in the sequel

$$\hat{T} := \hat{\omega} + dd^c g .$$

One can give the following definition.

Definition 2.2. — *The bifurcation current of the critical point c_i in $(f_\lambda)_{\lambda \in \Lambda}$ is*

$$T_i := (p_\Lambda)_* \left((\hat{\omega} + dd^c g) \wedge [\hat{V}_i] \right) ,$$

where $\hat{V}_i = \{(\lambda, v_i(\lambda)) : \lambda \in \Lambda\}$ is the graph of the map $v_i(\lambda) := f_\lambda(c_i(\lambda))$.

The holomorphic family $(f_\lambda)_{\lambda \in \Lambda}$ admits local lifts, i.e. for any small enough $V \subset \Lambda$, there exists a holomorphic family $(F_\lambda)_{\lambda \in V}$ of non-degenerate homogeneous degree d polynomial endomorphisms of \mathbb{C}^2 , and the *Green function* of the lift is then

$$G(\lambda, x, y) := \lim_{n \rightarrow \infty} d^{-n} \log \|F_\lambda^n(x, y)\| , \quad (x, y) \in \mathbb{C}^2 \setminus \{0\} .$$

It actually is a continuous and psh function on $V \times (\mathbb{C}^2 \setminus \{0\})$. According to [B, Section 3.2.2], one can prove that for any local holomorphic section $\sigma : U \subset \mathbb{P}^1 \rightarrow \mathbb{C}^2 \setminus \{0\}$ of the canonical projection $\mathbb{C}^2 \setminus \{0\} \rightarrow \mathbb{P}^1$, then

$$\hat{\omega} + dd^c g = dd^c G(\lambda, \sigma(z)) \text{ , on } V \times U.$$

We thus can locally write $T_i = dd^c G(\lambda, \sigma \circ v_i(\lambda)) = d \cdot dd^c G(\lambda, \sigma \circ c_i(\lambda))$ on U .

Remark. — The definition we give here is not the classical one. The usual definition of the bifurcation currents is to take locally $\tilde{T}_i := dd^c G(\lambda, \sigma \circ c_i(\lambda)) = d^{-1} \cdot T_i$, which does not change the support of the current.

The important information concerning the current T_i is the following (see [DF]).

Proposition 2.3 (Dujardin-Favre). — *The support of T_i is the activity locus of c_i .*

Another way to characterize the bifurcation current of c_i is the following. Let $\xi_n^i : \Lambda \rightarrow \mathbb{P}^1$ be the map given by $\xi_n^i(\lambda) := f_\lambda^n(v_i(\lambda))$, for $n \geq 0$ and $1 \leq i \leq 2d - 2$. The sequence of forms $d^{-n}(\xi_n^i)^* \omega_{\text{FS}}$ converge to the current T_i in the sense of currents (see e.g. [Du]).

2.2. Bifurcation currents of a holomorphic family

Recall that $f \in \text{Rat}_d$ admits a unique maximal entropy measure μ_f . The *Lyapounov exponent* of f with respect to the measure μ_f is the real number $L(f) := \int_{\mathbb{P}^1} \log |f'| \mu_f$. For a holomorphic family $(f_\lambda)_{\lambda \in \Lambda}$ of degree d rational maps, we denote by $L(\lambda) := L(f_\lambda)$. Then, the function $\lambda \mapsto L(\lambda)$ is called the *Lyapounov function* of the family $(f_\lambda)_{\lambda \in \Lambda}$. It is a psh and continuous function on Λ (see [BB] Corollary 3.4). The Margulis-Ruelle inequality implies that $L(f) \geq \frac{\log d}{2}$.

When $(f_\lambda)_{\lambda \in \Lambda}$ is with $2d - 2$ distinct marked critical points c_1, \dots, c_{2d-2} , the bifurcation locus in the sense of Mañé-Sad-Sullivan and Lyubich (see [Ly, MSS]) coincides with the union of the activity loci of the c_i 's. According to DeMarco [De], we have the following.

Theorem 2.4 (DeMarco). — *Let $(f_\lambda)_{\lambda \in \Lambda}$ be a holomorphic family of degree d rational maps with $2d - 2$ distinct marked critical points. Then the current $T_{\text{bif}} := dd^c L$ is exactly supported by the bifurcation locus. Moreover, $T_{\text{bif}} = d^{-1} \sum_{i=1}^{2d-2} T_i$.*

The current T_{bif} is the *bifurcation current* of the family $(f_\lambda)_{\lambda \in \Lambda}$. The self-intersections of the current T_{bif} have been first studied by Bassanelli and Berteloot [BB].

Definition 2.5. — *We define the m^{th} -bifurcation current of the family $(f_\lambda)_{\lambda \in \Lambda}$ by setting*

$$T_{\text{bif}}^m := \bigwedge_{i=1}^m T_{\text{bif}} \text{ .}$$

It is known that for all $1 \leq i \leq 2d - 2$, we have $T_i \wedge T_i = 0$ (see [G1, Theorem 6.1]) and one can show that

$$(2) \quad T_{\text{bif}}^m = m! \sum_{i_1 < \dots < i_m} T_{i_1} \wedge \dots \wedge T_{i_m} \text{ .}$$

2.3. A formula for higher bifurcation currents

We want here to give a similar expression as the one given in Definition 2.2 for the higher bifurcation current associated to a m -tuple of critical points. Let us introduce some notations. Let $(f_\lambda)_{\lambda \in \Lambda}$ be a holomorphic family of degree d rational maps with m marked critical points, $c_1, \dots, c_m : \Lambda \rightarrow \mathbb{P}^1$, with $1 \leq m \leq \min(2d - 2, \dim \Lambda)$. As above, we define $v_j : \Lambda \rightarrow \mathbb{P}^1$ for $\lambda \in \Lambda$ by $v_j(\lambda) := f_\lambda(c_j(\lambda))$. This time, we let

$$\mathcal{V}_j := \{(\lambda, z) \in \Lambda \times (\mathbb{P}^1)^m : z_j = v_j(\lambda)\}$$

We finally let $\pi_\Lambda : \Lambda \times (\mathbb{P}^1)^m \rightarrow \Lambda$ and, for $1 \leq j \leq m$, we let

$$\begin{aligned} \pi_j : \Lambda \times (\mathbb{P}^1)^m &\longrightarrow \Lambda \times \mathbb{P}^1 \\ (\lambda, z) &\longmapsto (\lambda, z_j) \end{aligned}$$

be the respective natural projection. Let T_i be the bifurcation current of c_i in $(f_\lambda)_{\lambda \in \Lambda}$.

Lemma 2.6. — *With the above notations, we have*

$$\bigwedge_{j=1}^m T_j = (\pi_\Lambda)_* \left(\bigwedge_{j=1}^m (\pi_j)^* (\widehat{T}) \wedge [\mathcal{V}_j] \right).$$

Proof. — It is a local problem, so we can assume that $\Lambda = \mathbb{B}$ is a ball of \mathbb{C}^N for some $N \geq 1$. Recall that, up to reducing the ball, one can also write $\widehat{T} = dd_{\lambda, z}^c G_\lambda(\sigma(z))$, where G_λ is the Green function of a holomorphic family of non-degenerate homogeneous polynomial lift F_λ of f_λ , i.e.

$$G_\lambda(z_1, z_2) := \lim_{n \rightarrow \infty} d^{-n} \log \|F_\lambda^n(z_1, z_2)\|, \quad (z_1, z_2) \in \mathbb{C}^2 \setminus \{0\},$$

and σ is any local section of the natural projection $\mathbb{C}^2 \setminus \{0\} \rightarrow \mathbb{P}^1$. Let σ_j be such a section which up to reducing the ball contains the image of the map $v_j : \mathbb{B} \rightarrow \mathbb{P}^1$, then $dd^c G_\lambda(\sigma_j \circ v_j(\lambda)) = T_j$ (see e.g. [B, Section 3.2.2]).

Let now ϕ be a smooth test $(N - m, N - m)$ -form on \mathbb{B} and let $\mathcal{V} := \bigcap_j \mathcal{V}_j$. Let $p(\lambda) := (\lambda, v_1(\lambda), \dots, v_m(\lambda))$, so that $\pi_\Lambda \circ p = \text{id}_{\mathbb{B}}$ and $p(\mathbb{B}) = \mathcal{V}$. Then

$$\begin{aligned}
\left\langle (\pi_\Lambda)_* \left(\bigwedge_{j=1}^m ((\pi_j)^* \widehat{T}) \wedge [\mathcal{V}_j] \right), \phi \right\rangle &= \int \bigwedge_{j=1}^m (\pi_j)^* (\widehat{T}) \wedge [\mathcal{V}_j] \wedge (\pi_\Lambda)^* \phi \\
&= \int_{\mathcal{V}} \bigwedge_{j=1}^m (\pi_j)^* (\widehat{T}) \wedge (\pi_\Lambda)^* \phi \\
&= \int_{\mathcal{V}} \bigwedge_{j=1}^m (\pi_j)^* dd_{\lambda, z}^c (G_\lambda \circ \sigma_j) \wedge (\pi_\Lambda)^* \phi \\
&= \int_{p(\mathbb{B})} \bigwedge_{j=1}^m dd_{\lambda, z}^c G_\lambda(\sigma_j(z_j)) \wedge (\pi_\Lambda)^* \phi \\
&= \int_{\mathbb{B}} \bigwedge_{j=1}^m dd^c G_\lambda(\sigma_j \circ v_j(\lambda)) \wedge ((p^*(\pi_\Lambda)^* \phi)) \\
&= \left\langle \bigwedge_{j=1}^m T_j, \phi \right\rangle.
\end{aligned}$$

□

3. Generalized large scale condition and the bifurcation currents

For the whole section, we let $(f_\lambda)_{\lambda \in \Lambda}$ be a holomorphic family of rational maps with $m \geq 1$ marked critical points $c_1, \dots, c_m : \Lambda \rightarrow \mathbb{P}^1$. As above, we use the notation $v_j(\lambda) := f_\lambda(c_j(\lambda))$. We set $\mathbf{c} := (c_1, \dots, c_m)$ and we also use the following notation: For any m -tuple of positive integers $\underline{n} = (n_1, \dots, n_m)$, any $1 \leq j \leq m$ and any $\lambda \in \Lambda$, we let

$$\xi_{n_j}^j(\lambda) := f_\lambda^{n_j}(v_j(\lambda)), \text{ and } \mathfrak{V}_{\underline{n}}^{\mathbf{c}}(\lambda) := (\xi_{n_1}^1(\lambda), \dots, \xi_{n_m}^m(\lambda)).$$

This way, we define a holomorphic map $\mathfrak{V}_{\underline{n}}^{\mathbf{c}} : \Lambda \rightarrow (\mathbb{P}^1)^m$. We denote by $\mathcal{V}_{\underline{n}}$ the graph of $\mathfrak{V}_{\underline{n}}^{\mathbf{c}}$.

We now can define the (generalized) large scale condition.

Definition 3.1 (Generalized large scale condition). — *We say that a parameter $\lambda_0 \in \Lambda$ satisfies the generalized large scale condition at f_{λ_0} for the m -tuple (c_1, \dots, c_m) in Λ if there exist a sequence $\underline{n}_k = (n_{k,1}, \dots, n_{k,m})$ of m -tuples with $n_{k,j} \rightarrow +\infty$ and a basis of neighborhood $\{\Omega_k\}_{k \geq 1}$ of f in Λ and $\delta > 0$ such that for any k , the connected component of $\mathcal{V}_{\underline{n}_k} \cap \Omega_k \times \left(\mathbb{D}(\xi_{n_{k,1}}^1(\lambda_0), \delta) \times \dots \times \mathbb{D}(\xi_{n_{k,m}}^m(\lambda_0), \delta) \right)$ containing $(\lambda_0, \mathfrak{V}_{\underline{n}_k}^{\mathbf{c}}(\lambda_0))$ is contained in $\Omega'_k \times \left(\mathbb{D}(\xi_{n_{k,1}}^1(\lambda_0), \delta) \times \dots \times \mathbb{D}(\xi_{n_{k,m}}^m(\lambda_0), \delta) \right)$ for some $\Omega'_k \Subset \Omega_k$.*

We prove in this section the following result which we view as a general sufficient condition for a parameter to belong to the support of a (higher) bifurcation current.

Theorem 3.2. — *Let $1 \leq m \leq \min(2d - 2, \dim \Lambda)$ be an integer and let $\lambda_0 \in \Lambda$. Assume that λ_0 satisfies the generalized large scale condition for $\mathbf{c} := (c_1, \dots, c_m)$ in a local submanifold $S \ni \lambda_0$ of Λ with $\dim S = m$. Assume in addition that $\omega(c_i(\lambda_0)) \subset \mathcal{I}_{\lambda_0}$ for all $1 \leq i \leq m$. Then $\lambda_0 \in \text{supp}(T_1 \wedge \dots \wedge T_m)$.*

Proof of Theorem B. — By definition of μ_{bif} , we have $[f] \in \text{supp}(\mu_{\text{bif}})$ if and only if $f \in \text{supp}(T_{\text{bif}}^{2d-2})$. On the other hand, since f has simple critical points, there exist a neighborhood $U \subset \text{Rat}_d$ of f and holomorphic maps $c_1, \dots, c_{2d-2} : U \rightarrow \mathbb{P}^1$ with $\mathcal{C}(g) = \{c_1(g), \dots, c_{2d-2}(g)\}$ for all $g \in U$. We then can apply the above Theorem 3.2 to $\mathbf{c} = (c_1, \dots, c_{2d-2})$. Since, by (2), $T_{\text{bif}}^{2d-2} = (2d-2)!T_1 \wedge \dots \wedge T_{2d-2}$ on U , the result follows. \square

Remark. — Note that the assumption $\omega(c_j(\lambda_0)) \subset \mathcal{J}_{\lambda_0}$ is satisfied not only when $c_j(\lambda_0) \in \mathcal{J}_{\lambda_0}$, but also when $c_j(\lambda_0)$ belongs to a parabolic basin.

First, note that it is sufficient to treat the case when $\dim \Lambda = m$ and λ_0 satisfies the large scale condition for $\mathbf{c} := (c_1, \dots, c_m)$ in a Λ . Indeed, by [G1, Lemma 6.3], if $\lambda_0 \in \text{supp}((T_1 \wedge \dots \wedge T_m)|_S)$, then $\lambda_0 \in \text{supp}(T_1 \wedge \dots \wedge T_m)$. We hence may assume $S = \Lambda$ has dimension m and let

$$\mu := T_1 \wedge \dots \wedge T_m .$$

It defines a positive measure on Λ .

Let $(\lambda_1, \dots, \lambda_m)$ be a local system of holomorphic coordinates centered at λ_0 . We let \mathbb{D}_δ^m be the polydisk of radius $\delta > 0$ of Λ centered at λ_0 in those coordinates.

3.1. The case $m = 1$: a toy-model for the general case

We give here the proof of Theorem 3.2 in the case $m = 1$. We let $c : \Lambda \rightarrow \mathbb{P}^1$ be the marked critical point satisfying the large scale condition at λ_0 . As above, for $n \geq 0$ and $\lambda \in \Lambda$, write

$$\xi_n(\lambda) := f_\lambda^{n+1}(c(\lambda)) = f_\lambda^n(\xi_0(\lambda)) .$$

Of course, in that case, it is easy to see that the large scale condition implies the non-normality of the family $(f_\lambda^n(c(\lambda)))_n$ but we provide here a proof that can be adapted to work with higher degree bifurcation currents. For that, recall that \mathcal{V}_n is the graph of $\xi_n : \Lambda \rightarrow \mathbb{P}^1$. Recall that $d^{-1}(\hat{f})^*\hat{T} = \hat{T}$. Choose any $\epsilon > 0$. We shall prove that $\mu(\mathbb{D}(\lambda_0, \epsilon)) > 0$ for all $\epsilon > 0$ small enough. For any $n \geq 1$

$$\begin{aligned} \mu(\mathbb{D}(\lambda_0, \epsilon)) &= \int_{\mathbb{D}(\lambda_0, \epsilon) \times \mathbb{P}^1} \hat{T} \wedge [\mathcal{V}_0] \\ &= d^{-n} \int_{\mathbb{D}(\lambda_0, \epsilon) \times \mathbb{P}^1} (\hat{f}^n)^* (\hat{T}) \wedge [\mathcal{V}_0] . \end{aligned}$$

As a consequence,

$$\begin{aligned} I_n &:= d^n \mu(\mathbb{D}(\lambda_0, \epsilon)) = \int_{\mathbb{D}(\lambda_0, \epsilon) \times \mathbb{P}^1} (\hat{f}^n)^* (\hat{T}) \wedge [\mathcal{V}_0] \\ &= \int_{\mathbb{D}(\lambda_0, \epsilon) \times \mathbb{P}^1} \hat{T} \wedge (\hat{f}^n)_* [\mathcal{V}_0] = \int_{\mathbb{D}(\lambda_0, \epsilon) \times \mathbb{P}^1} \hat{T} \wedge [\mathcal{V}_n], \end{aligned}$$

since on one hand $\hat{f}^{-1}(\mathbb{D}(\lambda_0, \epsilon) \times \mathbb{P}^1) = \mathbb{D}(\lambda_0, \epsilon) \times \mathbb{P}^1 = \hat{f}(\mathbb{D}(\lambda_0, \epsilon) \times \mathbb{P}^1)$ and, on the other hand, $(\hat{f}^n)_* [\mathcal{V}_0] = [\mathcal{V}_n]$. Let now $\delta > 0$, (n_k) and (Ω_k) be given by the large scale condition. Up to extraction, we can assume $\xi_{n_k} := \xi_{n_k}(\lambda_0)$ converges to some point $x \in \mathcal{J}_{\lambda_0}$. As a consequence, there exists $k_0 \geq 1$ such that $\Omega_k \subset \mathbb{D}(\lambda_0, \epsilon)$ and

$$\xi_{n_k}(\Omega_k) \supset \mathbb{D}(x, \delta/2) ,$$

for any $k \geq k_0$. We now let S_k be the connected component of $\mathcal{V}_{n_k} \cap \Omega_k \times \mathbb{D}(x, \delta/2)$ containing $(0, \xi_{n_k})$. Then $S_k \subset \Omega_k \times \mathbb{D}(x, \delta/2)$, $[S_k]/\|[S_k]\|$ has mass 1 and

$$[\mathcal{V}_{n_k}] \geq \mathbf{1}_{\mathbb{D}(\lambda_0, \epsilon) \times \mathbb{D}(x, \delta/2)} [\Gamma_{n_k}] \geq [S_k] .$$

for all $k \geq k_0$. Let now S be any weak limit of the sequence $[S_k]/\|[S_k]\|$. Then $\text{supp}(S) = \{\lambda_0\} \times \mathbb{D}(x, \delta/2)$ and S has mass 1 by the large scale condition. By extremality of the current $[\{\lambda_0\} \times \mathbb{D}(x, \delta/2)]$, we deduce $S = M \cdot [\{\lambda_0\} \times \mathbb{D}(x, \delta/2)]$, where $M^{-1} > 0$ is the Fubini-Study area of $\mathbb{D}(x, \delta/2)$. As \widehat{T} has continuous potential, $\widehat{T} \wedge [S_k]/\|[S_k]\| \rightarrow \widehat{T} \wedge S$ as $k \rightarrow \infty$ in the sense of measures so:

$$\begin{aligned} \liminf_{k \rightarrow \infty} \|[S_k]\|^{-1} \cdot I_{n_k} &\geq \liminf_{k \rightarrow \infty} \int \widehat{T} \wedge \frac{[S_k]}{\|[S_k]\|} \geq \int \widehat{T} \wedge S = M \cdot \int \widehat{T} \wedge [\{\lambda_0\} \times \mathbb{D}(x, \delta/2)] \\ &= M \cdot \mu_{\lambda_0}(\mathbb{D}(x, \delta/2)) \end{aligned}$$

as $\widehat{T}|_{\lambda=0} = \mu_{\lambda_0}$. Since $x \in \mathcal{J}_f = \text{supp}(\mu_{\lambda_0})$ then $\mu_{\lambda_0}(\mathbb{D}(x, \delta/2)) > 0$, we get

$$\liminf_{k \rightarrow \infty} \|[S_k]\|^{-1} \cdot I_{n_k} > 0 ,$$

which means that $I_{n_k} > 0$ for k large enough so $\mu(\mathbb{D}(\lambda_0, \epsilon)) > 0$. Since this holds for all $\epsilon > 0$, this ends the proof.

3.2. First step: pulling-back by a fibered dynamical system

Let us define a family of fibered dynamical systems acting on $\Lambda \times (\mathbb{P}^1)^m$ as follows: for any m -tuple $\underline{n} := (n_1, \dots, n_m) \in (\mathbb{N}^*)^m$, we let

$$\begin{aligned} F_{\underline{n}} : \Lambda \times (\mathbb{P}^1)^m &\longrightarrow \Lambda \times (\mathbb{P}^1)^m \\ (\lambda, z_1, \dots, z_m) &\longmapsto (\lambda, f_{\lambda}^{n_1}(z_1), \dots, f_{\lambda}^{n_m}(z_m)) . \end{aligned}$$

For a m -tuple $\underline{n} = (n_1, \dots, n_m)$ of positive integers, we also set

$$|\underline{n}| := n_1 + \dots + n_m .$$

Let us first partially rewrite the mass of μ on any open set in terms of iterated pull-back by one the $F_{\underline{n}}$'s.

Lemma 3.3. — *For any m -tuple $\underline{n} = (n_1, \dots, n_m)$ of positive integers, we let $\mathcal{V}_{\underline{n}}$ be the graph in $\Lambda \times (\mathbb{P}^1)^m$ of $\mathfrak{V}_{\underline{n}}$. Then, for any Borel set $\Omega \subset \Lambda$, we have*

$$\mu(\Omega) = d^{-|\underline{n}|} \int_{\Omega \times (\mathbb{P}^1)^m} \left(\bigwedge_{j=1}^m (\pi_j)^* (\widehat{T}) \right) \wedge [\mathcal{V}_{\underline{n}}] .$$

Proof. — Recall that we defined in Section 2.1 a dynamical system \widehat{f} acting on $\Lambda \times \mathbb{P}^1$ by

$$\begin{aligned} \widehat{f} : \Lambda \times \mathbb{P}^1 &\longrightarrow \Lambda \times \mathbb{P}^1 \\ (\lambda, z) &\longmapsto (\lambda, f_{\lambda}(z)) . \end{aligned}$$

By definition of $F_{\underline{n}}$ and \widehat{f} , for all j , the following diagram commutes

$$\begin{array}{ccc} \Lambda \times (\mathbb{P}^1)^m & \xrightarrow{F_{\underline{n}}} & \Lambda \times (\mathbb{P}^1)^m \\ \pi_j \downarrow & & \downarrow \pi_j \\ \Lambda \times \mathbb{P}^1 & \xrightarrow{\widehat{f}^{n_j}} & \Lambda \times \mathbb{P}^1 . \end{array}$$

In particular, by (1), we get

$$\begin{aligned} d^{-n_j}(F_{\underline{n}})^* \left((\pi_j)^* \left(\widehat{T} \right) \right) &= d^{-n_j}(\pi_j \circ F_{\underline{n}})^* \left(\widehat{T} \right) = d^{-n_j}(\widehat{f}^{n_j} \circ \pi_j)^* \left(\widehat{T} \right) \\ &= d^{-n_j}(\pi_j)^* \left((\widehat{f}^{n_j})^* \left(\widehat{T} \right) \right) = (\pi_j)^* \left(\widehat{T} \right). \end{aligned}$$

According to Lemma 2.6, the change of variable formula gives

$$\begin{aligned} \mu(\Omega) &= \int_{\Omega} (\pi_{\Lambda})_* \left(\bigwedge_{j=1}^m (\pi_j)^* \left(\widehat{T} \right) \wedge [\mathcal{V}_j] \right) \\ &= \int_{\pi_{\Lambda}^{-1}(\Omega)} \bigwedge_{j=1}^m (\pi_j)^* \left(\widehat{T} \right) \wedge [\mathcal{V}_j] \\ &= d^{-|\underline{n}|} \int_{\Omega \times (\mathbb{P}^1)^m} (F_{\underline{n}})^* \left(\bigwedge_{j=1}^m (\pi_j)^* \left(\widehat{T} \right) \right) \wedge \left[\bigcap_{j=1}^m \mathcal{V}_j \right] \end{aligned}$$

where the last equality comes from $\pi_{\Lambda}^{-1}(\Omega) = \Omega \times (\mathbb{P}^1)^m$. Whence

$$\mu(\Omega) = d^{-|\underline{n}|} \int_{\Omega \times (\mathbb{P}^1)^m} \left(\bigwedge_{j=1}^m (\pi_j)^* \left(\widehat{T} \right) \right) \wedge (F_{\underline{n}})_* \left[\bigcap_{j=1}^m \mathcal{V}_j \right]$$

where we used $(F_{\underline{n}})_* \bigwedge_j [\mathcal{V}_j] = [\mathcal{V}_{\underline{n}}]$. \square

3.3. Second step: a phase-parameter transfer phenomenon

For the sake of simplicity, we let in the sequel $\mathfrak{V}_k := \mathfrak{V}_{\underline{n}_k}^c$, where \underline{n}_k is given by the large scale condition. Up to extracting a subsequence, we may assume that $f_{\lambda_0}^{n_k}(c_j(\lambda_0)) \rightarrow x_j \in \mathcal{J}_{\lambda_0}$ for all $1 \leq j \leq m$. We also let $x := (x_1, \dots, x_m) \in (\mathbb{P}^1)^m$.

We want to reduce the problem to a purely dynamical datum of f . Building on the large scale condition, one actually gets the following.

Proposition 3.4. — *There exist $k_0 \geq 1$ and $\delta, \alpha > 0$ such that for any $k \geq k_0$, we have*

$$\int_{\Omega_k \times (\mathbb{P}^1)^m} \left(\bigwedge_{j=1}^m (\pi_j)^* \left(\widehat{T} \right) \right) \wedge [\mathcal{V}_{\underline{n}_k}] > 0.$$

Proof of Proposition 3.4. — Set

$$I_k := \int_{\Omega_k \times (\mathbb{P}^1)^m} \left(\bigwedge_{j=1}^m (\pi_j)^* \left(\widehat{T} \right) \right) \wedge [\mathcal{V}_{\underline{n}_k}],$$

and let δ be given by the large scale condition. Let S_k be the connected component of $\mathcal{V}_{\underline{n}_k} \cap \Omega_k \times \mathbb{D}_{\delta}^m(x)$ containing $(0, \mathfrak{V}_{\underline{n}_k}(0))$. Up to replacing δ with $\delta/2$, for any $k \geq k_1$, the current $[S_k]/\|[S_k]\|$ is of a vertical current of mass 1 in $\Lambda \times \mathbb{D}_{\delta}^m(v_{\infty})$ and

$$\text{supp}([S_k]) = S_k \subset \Omega_k \times \mathbb{D}_{\delta}^m(x).$$

As in the case $m = 1$, let S be any weak limit of the sequence $[S_k]/\|[S_k]\|$. Then S is a closed positive (m, m) -current of mass 1 in $\mathbb{B}_{\epsilon} \times \mathbb{D}_{\delta}^m(x)$ with $\text{supp}(S) = \{\lambda_0\} \times \mathbb{D}_{\delta}^m(x)$ by the large scale condition. Hence, by extremality of $[\{\lambda_0\} \times \mathbb{D}_{\delta}^m(x)]$, we have that $S = M \cdot [\{\lambda_0\} \times \mathbb{D}_{\delta}^m(x)]$,

where $M^{-1} > 0$ is the volume of $\mathbb{D}_\delta^m(x)$ for the volume form $\bigwedge_j \omega_j$, where $\omega_j = (p_j)^* \omega_{\text{FS}}$ and $p_j : (\mathbb{P}^1)^m \rightarrow \mathbb{P}^1$ is the projection on the j -th coordinate.

As a consequence, $[S_k]/\|[S_k]\|$ weakly converges to S as $k \rightarrow \infty$. Since $\bigwedge_{j=1}^m (\pi_j)^* (\widehat{T})$ is the wedge product of $(1, 1)$ current with continuous potentials, we have

$$\bigwedge_{j=1}^m (\pi_j)^* (\widehat{T}) \wedge \frac{[S_k]}{\|[S_k]\|} \rightarrow \bigwedge_{j=1}^m (\pi_j)^* (\widehat{T}) \wedge S$$

so

$$\liminf_{k \rightarrow \infty} (\|[S_k]\|^{-1} \cdot I_k) \geq \liminf_{k \rightarrow \infty} \int \bigwedge_{j=1}^m (\pi_j)^* (\widehat{T}) \wedge \frac{[S_k]}{\|[S_k]\|} \geq \int \bigwedge_{j=1}^m (\pi_j)^* (\widehat{T}) \wedge S.$$

By the above, this gives

$$\begin{aligned} \liminf_{k \rightarrow \infty} (\|[S_k]\|^{-1} \cdot I_k) &\geq M \cdot \int_{\{\lambda_0\} \times \mathbb{D}_\delta^m(x)} \bigwedge_{j=1}^m (\pi_j)^* (\widehat{T}) \\ &\geq \int \bigwedge_{j=1}^m (\pi_j)^* (\widehat{T}) \wedge [\{\lambda_0\} \times \mathbb{D}_\delta^m(x)]. \end{aligned}$$

The proof of the proposition directly follows from the following lemma. \square

Lemma 3.5. — *For any $\delta > 0$, and any $x = (x_1, \dots, x_m) \in (\mathcal{J}_f)^m$, we have*

$$\int \bigwedge_{j=1}^m (\pi_j)^* (\widehat{T}) \wedge [\{\lambda_0\} \times \mathbb{D}_\delta^m(x)] = \prod_{j=1}^m \mu_{\lambda_0}(\mathbb{D}(x_j, \delta)) > 0.$$

Proof. — Let us set $\omega_j := ((\pi_j)^* \widehat{\omega})|_{\lambda=\lambda_0}$. First, we can remark that $g|_{\lambda=\lambda_0} = g_{\lambda_0}$ is the Green function of the rational map f_{λ_0} . We denote by $p_j : (\mathbb{P}^1)^k \rightarrow \mathbb{P}^1$ the canonical projection onto the j^{th} coordinate. A classical slicing argument gives

$$((\pi_j)^* dd^c g)|_{\lambda=\lambda_0} = dd_z^c(g_{\lambda_0} \circ p_j) = (p_j)^* dd^c g_{\lambda_0}.$$

In particular, since $\omega_j = (p_j)^* \omega_{\text{FS}}$, we have

$$\left((\pi_j)^* (\widehat{T}) \right) \Big|_{\lambda=\lambda_0} = ((\pi_j)^* (\omega_{\text{FS}} + dd^c g)) \Big|_{\lambda=\lambda_0} = (p_j)^* (\mu_{\lambda_0}),$$

where μ_{λ_0} is the maximal entropy measure of f_{λ_0} , hence

$$\begin{aligned} I &:= \int \bigwedge_{j=1}^m (\pi_j)^* (\widehat{T}) \wedge [\{\lambda_0\} \times \mathbb{D}_\delta^m(x)] = \int_{\{\lambda_0\} \times \mathbb{D}_\delta^m(x)} \bigwedge_{j=1}^m (\pi_j)^* (\widehat{T}) \\ &= \int_{\mathbb{D}_\delta^m(x)} \bigwedge_{j=1}^m \left((\pi_j)^* (\widehat{T}) \right) \Big|_{\lambda=\lambda_0} \\ &= \int_{\mathbb{D}_\delta^m(x)} \bigwedge_{j=1}^m (p_j)^* (\mu_{\lambda_0}). \end{aligned}$$

Since $\text{supp}(\mu_{\lambda_0}) = \mathcal{J}_{\lambda_0}$ and $x_1, \dots, x_m \in \mathcal{J}_{\lambda_0}$, Fubini Theorem yields

$$I = \prod_{j=1}^m \left(\int_{\mathbb{D}(x_j, \delta)} \mu_{\lambda_0} \right) = \prod_{j=1}^m \mu_{\lambda_0}(\mathbb{D}(x_j, \delta)) > 0,$$

which ends the proof of Lemma 3.5. \square

The proof of Theorem 3.2 directly follows from Lemma 3.3 and Proposition 3.4.

4. Misiurewicz maps and the generalized large scale condition

Fix $d \geq 2$ and pick $1 \leq k \leq 2d - 2$. A rational map $f \in \text{Rat}_d$ is *k-Misiurewicz* if the following properties hold:

- f has no parabolic periodic points,
- f has k critical points in its Julia set, counted with multiplicity,
- for any $c \in \mathcal{C}(f) \cap \mathcal{J}_f$, we have $\omega(c) \cap \mathcal{C}(f) = \emptyset$.

In this section, we want to emphasize that the large scale condition is the good condition for proving that specific parameters lie in the support of the bifurcation measure. Our motivation here is also to provide a simpler and more intrinsic proof of Theorem 1.4 of [G1]. Recall that for a critical point c_j , we denote by v_j the critical value $f(c_j)$. According to Theorem 3.2, it is sufficient to prove the existence of a (local) submanifold in which f satisfies the (generalized) large scale condition. More precisely, we prove the following.

Theorem 4.1. — *Let $f \in \text{Rat}_d$ and $1 \leq k \leq 2d - 2$. Assume that f is k -Misiurewicz and that $\mathcal{C}(f) \cap \mathcal{J}_f = \{c_1, \dots, c_k\}$. Assume that for all $n \in \mathbb{N}$ and all $i \neq j \leq k$, $f^n(v_j) \neq v_i$. If f is not a flexible Lattès map, then there exists a k -dimensional local submanifold $\Lambda_f \subset \text{Rat}_d$ such that f satisfies the generalized large scale condition for (c_1, \dots, c_k) in Λ_f .*

This approach allows us to exhibit the key expansion and distortion arguments, without using more elaborate tools that will necessarily be missing in a more general situation, such as linearizing coordinates along repelling orbits.

Observe also that the condition "for all $n \in \mathbb{N}$ and all $i \neq j \leq k$, $f^n(v_j) \neq v_i$ " is not an issue: any k -Misiurewicz map that does not satisfy this condition can be approximated by k -Misiurewicz maps that do satisfy it using Montel theorem. In particular, it is in the support of the bifurcation current $T_1 \wedge \dots \wedge T_k$.

4.1. Hyperbolic sets and holomorphic motions

We recall some classical definitions and facts. Let $f \in \text{Rat}_d$ and $E \subset \mathbb{P}^1$ be a non-empty compact f -invariant set, i.e. such that $f(E) \subset E$. We say that E is *f-hyperbolic* if one of the following equivalent conditions is satisfied:

1. there exist $C > 0$ and $\alpha > 1$ such that $|(f^n)'(z)| \geq C\alpha^n$ for all $z \in E$ and all $n \geq 0$,
2. for some appropriate metric on \mathbb{P}^1 , there exists $K > 1$ such that $|f'(z)| \geq K$ for all $z \in E$.

Such a constant K is called a *hyperbolicity constant* for f .

Recall also that a *holomorphic motion* of a set $X \subset \mathbb{P}^1$ over a complex manifold Λ centered at $\lambda_0 \in \Lambda$ is a map $h : \Lambda \times X \rightarrow \mathbb{P}^1$ such that:

- $h_{\lambda_0} := h(\lambda_0, \cdot) : X \rightarrow \mathbb{P}^1$ is the identity map,
- $h_\lambda := h(\lambda, \cdot) : X \rightarrow \mathbb{P}^1$ is injective for all $\lambda \in \Lambda$,
- $\lambda \mapsto h(\lambda, x)$ is holomorphic on Λ for all $x \in X$.

First, notice that the classical λ -lemma of Mañé-Sad-Sullivan [MSS] says that any holomorphic motion of X extends continuously to a holomorphic motion of the closure of X . It is known that a hyperbolic set E admits a natural holomorphic motion (see, e.g., [S, Property (1.2) page 229] or [J, Theorem C]).

Theorem 4.2. — *Let $(f_\lambda)_{\lambda \in \mathbb{B}(0,r)}$ be a holomorphic family of degree d rational maps parametrized by a ball $\mathbb{B}(0,r) \subset \mathbb{C}^m$. Let $E_0 \subset \mathbb{P}^1$ be a compact f_0 -hyperbolic set. Then there exist $0 < \rho \leq r$ and a unique holomorphic motion*

$$\begin{aligned} h : \mathbb{B}(0,\rho) \times E_0 &\longrightarrow \mathbb{P}^1 \\ (\lambda, z) &\longmapsto h_\lambda(z) \end{aligned}$$

centered at 0 and such that $f_\lambda \circ h_\lambda(z) = h_\lambda \circ f_0(z)$ for all $z \in E_0$.

The proof of Theorem 4.2 relies on the compactness of E_0 and on the next lemma, which is an immediate corollary of the Implicit Function Theorem. Let $K > 1$ be a hyperbolicity constant for E_0 for a suitable metric α . In what follows, we denote by $|\cdot|_\alpha, \mathbb{D}_\alpha$ the distance and disk with respect to that metric. Up to reducing K and r , we can find a δ -neighborhood \mathcal{N}_δ of E_0 in \mathbb{P}^1 such that

$$(3) \quad |f'_\lambda(z)|_\alpha \geq K > 1 \text{ for all } (z, \lambda) \in \mathcal{N}_\delta \times \mathbb{B}(0,r).$$

Lemma 4.3. — *Under the assumption of Theorem 4.2, there exist $\varepsilon > 0$ and $0 < \rho \leq r$, such that for all $z_0 \in E_0$, there exists a map $f_{z_0,\lambda}^{-1}(w)$ which depends holomorphically on $(\lambda, w) \in \mathbb{B}(0,\rho) \times \mathbb{D}_\alpha(f_0(z_0), \varepsilon)$ and taking values in $\mathbb{D}_\alpha(z_0, \varepsilon)$ which satisfies*

1. $f_{z_0,0}^{-1}(f_0(z_0)) = z_0$,
2. $f_\lambda(f_{z_0,\lambda}^{-1}(w)) = w$ for all $(\lambda, w) \in \mathbb{B}(0,\rho) \times \mathbb{D}_\alpha(f_0(z_0), \varepsilon)$, and
3. $\left| (f_{z_0,\lambda}^{-1})'(w) \right|_\alpha \leq \frac{1}{K}$ for all $(\lambda, w) \in \mathbb{B}(0,\rho) \times \mathbb{D}_\alpha(f_0(z_0), \varepsilon)$.

4.2. Transversality for Misiurewicz maps

It is well known that if f is a k -Misiurewicz degree d rational map but not a flexible Lattès map, then all periodic Fatou components of f are attracting basins and f does not carry any invariant line field on its Julia set. Moreover, there exists a positive integer $n_0 \geq 1$ such that the set

$$E_f := \overline{\{f^n(c); n \geq n_0 \text{ and } c \in \mathcal{C}(f) \cap \mathcal{J}_f\}}$$

is a compact hyperbolic f -invariant set (see e.g. [Asp1, G1]).

If $f \in \text{Rat}_d$ is k -Misiurewicz, $\mathcal{C}(f) \cap \mathcal{J}_f = \{c_1, \dots, c_k\}$ and, for all $n \in \mathbb{N}$ and all $i \neq j \leq k$, $f^n(v_j) \neq v_i$, where $v_i = f(c_i)$, we let $h : \mathbb{B}(f,r) \times E_f \rightarrow \mathbb{P}^1$ be the dynamical holomorphic motion of E_f . Since all the critical points in the Julia set of f are simple ($v_i \neq v_j$), we may follow them holomorphically in the neighborhood of f , as well as the critical values: denote by $v_i(\lambda)$, $1 \leq i \leq k$, the corresponding critical values of f_λ .

Given a holomorphic curve $\lambda \mapsto f_\lambda \in \text{Rat}_d$ passing through $f = f_{\lambda_0}$, it will be convenient to denote by η the meromorphic vector field on \mathbb{P}^1 defined by

$$\eta(z) := \frac{\frac{d}{d\lambda}|_{\lambda=\lambda_0} f_\lambda(z)}{f'(z)}.$$

Recall that the pullback of η by f^n is given in coordinates by $(f^n)^*\eta(z) = \frac{\eta \circ f^n(z)}{(f^n)'(z)}$. As before, let $\xi_n^i(\lambda) := f_\lambda^n(v_i(\lambda))$.

Lemma 4.4. — *Let $(f_\lambda)_{\lambda \in \Lambda}$ be a holomorphic curve of rational maps with a marked critical value $v_i(\lambda)$. Assume that the orbit of $v_i := v_i(\lambda_0)$ does not meet the critical set of $f := f_{\lambda_0}$. We*

have, for all $n \geq 1$: :

$$\frac{1}{(f^n)'(v_i)} \frac{d}{d\lambda}|_{\lambda=\lambda_0} \xi_n^i(\lambda) = \frac{d}{d\lambda}|_{\lambda=\lambda_0} v_i(\lambda) + \sum_{k=0}^{n-1} (f^k)^* \eta(v_i).$$

Moreover, if v_i lies in a hyperbolic set E_f and $h_\lambda(v_i)$ denote its holomorphic motion, then

$$\frac{d}{d\lambda}|_{\lambda=\lambda_0} h_\lambda(v_i) = - \sum_{n=0}^{+\infty} (f^n)^* \eta(v_i).$$

Proof. — To lighten the notations, we will denote by \dot{v}_i , $\dot{f}(z)$, etc. the derivatives $\frac{d}{d\lambda}|_{\lambda=\lambda_0} v_i(\lambda)$, $\frac{d}{d\lambda}|_{\lambda=\lambda_0} f_\lambda(z)$, etc.

We have: $\xi_n^i(\lambda) = f_\lambda^n(v_i(\lambda)) = f_\lambda \circ f_\lambda^{n-1}(v_i(\lambda))$, so

$$\frac{d}{d\lambda}|_{\lambda=\lambda_0} \xi_n^i(\lambda) = \dot{f} \circ f^{n-1}(v_i) + f' \circ f^{n-1}(v_i) \cdot \frac{d}{d\lambda}|_{\lambda=\lambda_0} \xi_{n-1}^i(\lambda).$$

Therefore

$$\frac{\frac{d}{d\lambda}|_{\lambda=\lambda_0} \xi_n^i(\lambda)}{(f^n)'(v_i)} = \frac{\eta \circ f^{n-1}(v_i)}{(f^{n-1})'(v_i)} + \frac{\frac{d}{d\lambda}|_{\lambda=\lambda_0} \xi_{n-1}^i(\lambda)}{(f^{n-1})'(v_i)} = (f^{n-1})^* \eta(v_i) + \frac{\frac{d}{d\lambda}|_{\lambda=\lambda_0} \xi_{n-1}^i(\lambda)}{(f^{n-1})'(v_i)}.$$

The first statement then follows by induction on n .

Let us now prove the second statement. By Theorem 4.2, there exists a holomorphic family of homeomorphisms $h_\lambda : E_f \rightarrow \mathbb{P}^1$, with $h_{\lambda_0} = \text{Id}$ and such that

$$h_\lambda \circ f = f_\lambda \circ h_\lambda.$$

Differentiating with respect to λ at $\lambda = \lambda_0$, we get:

$$\dot{h} \circ f = \dot{f} + f' \cdot \dot{h},$$

which we may rewrite as $f^* \dot{h} - \dot{h} = \eta$. Since E_f is hyperbolic, the operator f^* is strictly contracting on the space of vector fields on E_f , and therefore

$$\dot{h}(v_i) = - \sum_{n=0}^{+\infty} (f^n)^* \eta(v_i),$$

ending the proof. □

We can deduce the following from Theorem B of [Ast] (see the discussion after Theorem B in [Ast]):

Proposition 4.5. — *Let $f \in \text{Rat}_d$ be k -Misiurewicz, $\mathcal{C}(f) \cap \mathcal{J}_f = \{c_1, \dots, c_k\}$ and, for all $n \in \mathbb{N}$ and all $i \neq j \leq k$, $f^n(v_j) \neq v_i$. Then, there exists a local complex submanifold $\Lambda_f \subset \text{Rat}_d$ which contains f , such that the holomorphic map*

$$\lambda \in \Lambda_f \mapsto (v_1(\lambda) - h_\lambda(v_1), \dots, v_k(\lambda) - h_\lambda(v_k)) \in \mathbb{C}^k$$

defines local coordinates at $f_0 = f$ in Λ_f . Furthermore, the limits

$$\tau_i := \lim_{n \rightarrow \infty} \frac{D\xi_n^i(0)}{(f^n)'(v_i)}$$

exist as linear maps on $T_0\Lambda_f$, for $1 \leq i \leq k$, and τ_1, \dots, τ_k are linearly independent.

4.3. The generalized large scale condition for k -Misiurewicz maps

We now turn to the proof of Theorem 4.1.

Proof of Theorem 4.1. — Consider the set

$$\Omega_n := \left\{ \lambda \in \Lambda_f, \forall i \leq k, |h_\lambda(v_i) - v_i(\lambda)| \leq \frac{1}{|(f^n)'(v_i)|} \right\},$$

then, by Proposition 4.5, it can be written as $L_n^{-1}(\Omega)$ where

$$L_n(t_1, \dots, t_k) = \left(\frac{t_1}{|(f^n)'(v_1)|}, \dots, \frac{t_k}{|(f^n)'(v_k)|} \right)$$

and Ω is some polydisk around f in Λ_f . For all $1 \leq i \leq k$, let $\phi_n^i : \Omega \rightarrow \mathbb{C}$ be defined by:

$$\phi_n^i(\lambda) := \xi_n^i \left(\frac{\lambda}{|(f^n)'(v_i)|} \right) - \xi_n^i(0).$$

Let $\phi_n : \Omega \rightarrow \mathbb{C}^k$ be defined by $\phi_n := (\phi_n^i)_{1 \leq i \leq k}$.

We start by proving that the sequence $(\phi_n)_{n \in \mathbb{N}}$ is bounded, hence normal, on some small polydisk $\tilde{\rho} \cdot \Omega$. Let $i \leq k$, let us bound $|\xi_n^i(\lambda) - \xi_n^i(0)|$. We have:

$$(4) \quad \begin{aligned} |\xi_n^i(\lambda) - \xi_n^i(0)| &= |f_\lambda^n(v_i(\lambda)) - f^n(v_i)| \\ &\leq |f_\lambda^n(v_i(\lambda)) - f_\lambda^n(h_\lambda(v_i))| + |f_\lambda^n(h_\lambda(v_i)) - f^n(v_i)|. \end{aligned}$$

By chaining Lemma 4.3, there exists $\epsilon > 0$ such that for all λ small enough, for all $n \in \mathbb{N}$, there is an inverse branch $g_{n,\lambda}$ of f_λ^n that is well-defined on $\mathbb{D}_\alpha(f_\lambda^n(h_\lambda(v_i)), \epsilon)$, and that maps $f_\lambda^n(h_\lambda(v_i))$ to $h_\lambda(v_i)$. As the standard metric and the metric α are equivalent, we have that the map $g_{n,\lambda}$ is univalent on some disk $\mathbb{D}(f_\lambda^n(h_\lambda(v_i)), C \cdot \epsilon)$ for some constant C that does not depend on n nor λ . By Koebe's distortion theorem, the distortion of $g_{n,\lambda}$ is uniformly bounded on $\mathbb{D}(f_\lambda^n(h_\lambda(v_i)), C \cdot \epsilon/2)$.

We prove by induction on n that there is a constant $0 < \tilde{\rho} < 1$ depending only on f such that if $|h_\lambda(v_i) - v_i(\lambda)| \leq \frac{\tilde{\rho}}{|(f^n)'(v_i)|}$, then $|f_\lambda^n(v_i(\lambda)) - f_\lambda^n(h_\lambda(v_i))| \leq \frac{C\epsilon}{4}$. Assume it is true for $n-1$. Observe first that by Lemma 6.6 of [G1], there exists $C_1 > 0$ depending only on f and on the metric, such that we have:

$$(5) \quad \max \left(\frac{|(f_\lambda^{n-1})'(h_\lambda(v_i))|}{|(f^{n-1})'(v_i)|}, \frac{|(f^{n-1})'(v_i)|}{|(f_\lambda^{n-1})'(h_\lambda(v_i))|} \right) \leq e^{(n-1)C_1\|\lambda\|} \leq 2,$$

by our choice of λ . We can thus apply the induction hypothesis and the bound on the distortion of $g_{n-1,\lambda}$, and then (5) to find:

$$\begin{aligned} |f_\lambda^{n-1}(v_i(\lambda)) - f_\lambda^{n-1}(h_\lambda(v_i))| &\leq 2|(f_\lambda^{n-1})'(h_\lambda(v_i))| \cdot |v_i(\lambda) - h_\lambda(v_i)| \\ &\leq 4|(f^{n-1})'(v_i)| \cdot |v_i(\lambda) - h_\lambda(v_i)|. \end{aligned}$$

Let $M := \sup_{\lambda \in \mathbb{B}(0, \rho)} \|f'_\lambda\|_\infty$ and $m := \|(f')^{-1}\|_{E_f, \infty} < +\infty$ we deduce:

$$\begin{aligned} |f_\lambda^n(v_i(\lambda)) - f_\lambda^n(h_\lambda(v_i))| &\leq M |f_\lambda^{n-1}(v_i(\lambda)) - f_\lambda^{n-1}(h_\lambda(v_i))| \\ &\leq 4M |(f^{n-1})'(v_i)| \cdot |v_i(\lambda) - h_\lambda(v_i)| \\ &\leq 4Mm |(f^n)'(v_i)| \cdot |v_i(\lambda) - h_\lambda(v_i)| \\ &\leq 4Mm |(f^n)'(v_i)| \cdot \frac{\tilde{\rho}}{|(f^n)'(v_i)|} \\ &\leq \frac{C\varepsilon}{4} \end{aligned}$$

for $\tilde{\rho}$ small enough, independent of n . In particular,

$$|f_\lambda^n(v_i(\lambda)) - f_\lambda^n(h_\lambda(v_i))| = O(1).$$

It remains to bound the second term of (4). To do that, just note that $|f_\lambda^n(h_\lambda(v_i)) - f^n(v_i)| = |h_\lambda(f^n(v_i)) - f^n(v_i)|$, and that h_λ extends continuously to a holomorphic motion of \mathbb{P}^1 by the λ -lemma, hence in particular h is uniformly continuous on $\mathbb{B}(f, r) \times E_f$.

Therefore (ϕ_n) is bounded, hence normal on $\tilde{\rho} \cdot \Omega$. If ϕ is a limit of a subsequence of (ϕ_n) , then Proposition 4.5 implies that $D\phi(0)$ is invertible, hence that ϕ is open near 0. So there is some polydisk $\mathbb{D}(0, 2\delta)^k$ such that $\mathbb{D}(0, 2\delta)^k \subset \phi(\tilde{\rho} \cdot \Omega)$ and therefore, for arbitrarily large n , $\mathbb{D}(0, \delta)^k \subset \phi_n(\tilde{\rho} \cdot \Omega)$. Again, as $D\phi(0)$ is invertible, its graph is vertical near the origin (restricting δ if necessary); by normality, this is also the case for ϕ_n for n large enough which is exactly what we want. \square

5. The large scale condition and good Collet-Eckmann maps

In the present section, we prove Theorem D and Theorem C. We start by recalling the conditions appearing in Theorem C, which correspond to "good" Collet-Eckmann maps for which we will be able to prove that they also satisfy the large scale condition.

Let f be a rational map, and denote by $\mathcal{C}(f)$ its critical set. Recall the Collet-Eckmann condition:

Collet-Eckmann (CE) : there exist $\gamma, \gamma_0 > 0$ such that for all $v \in f(\mathcal{C}(f))$ and $n \geq 0$

$$|(f^n)'(v)| \geq e^{n\gamma - \gamma_0}$$

and the so-called *backward Collet-Eckmann condition*:

Backward Collet-Eckmann (CE2) : for all $n \geq 0$ and $x \in f^{-n}(\mathcal{C}(f))$

$$|(f^n)'(x)| > e^{n\mu - \mu_0}.$$

Additionally, recall the following conditions appearing in the papers [Ts] and [Asp2]:

Basic Assumption (BA) : there exists $\alpha > 0$ such that for all $v \in f(\mathcal{C}(f))$ and $n \geq 0$

$$\ln |f'(f^n(v))| > -n\alpha.$$

Free Assumption (FA) : there exist $\eta, \iota > 0$ such that for all $v \in f(\mathcal{C}(f))$ and $n > 0$

$$\sum_{\substack{j=0 \\ d(f^j(v), \mathcal{C}(f)) \leq \eta}}^{n-1} \ln |f'(f^j(v))| > -n\iota.$$

Aspenberg's Basic Assumption (BA') : there exists $\alpha > 0$ such that for all $v \in f(\mathcal{C}(f))$, $c \in \mathcal{C}(f)$ and $n \geq 0$

$$|f^n(v) - c| > e^{-n\alpha}$$

We will now describe a somewhat more technical key condition from [Asp2] (definition 1.15), which we will soon prove implies the condition (FA) with appropriate constants. Let $\gamma, \gamma_0 > 0$ and assume that f satisfies CE with constants γ, γ_0 . Fix $0 < \beta \ll \gamma$ and $0 < \delta \ll 1$. For some $v \in f(\mathcal{C}(f))$, we say that f makes a *deep return* at time ν if

$$d(f^\nu(v), \mathcal{C}(f)) < \delta^2.$$

The corresponding *bound period* p is defined by

$$p := \min\{k \geq 0 : |f^{k+1}(f^\nu(v)) - f^{k+1}(c)| \geq e^{-\beta k}\},$$

where c is the closest critical point to $f^\nu(v)$. Only deep returns outside bound periods are considered. Thus, for a given critical value v we can enumerate them

$$\nu_0(v) < \nu_0(v) + p_0(v) < \nu_1(v) < \dots$$

The length of the *free orbit* between two consecutive deep returns is in this case

$$\mu_i(v) := \nu_i(v) - (\nu_{i-1}(v) + p_{i-1}(v)) \text{ and } \mu_0(v) := \nu_0(v).$$

We omit the definition of *shallow returns* in [Asp2], as they are not needed here.

Aspenberg's Free Assumption (FA') : there exist $\delta, \beta > 0$ and $\tau \in (0, 1)$ such that for all $v \in f(\mathcal{C}(f))$ and $s \geq 0$

$$\sum_{i=0}^s \mu_i(v) > (1 - \tau)(\nu_s(v) + p_s(v)).$$

We will use the notation $\text{BA}(\alpha)$, $\text{CE}(\gamma, \gamma_0)$, $\text{FA}(\eta, \iota)$, etc. to refer to those conditions with the specified constants $\alpha, \gamma, \eta, \iota$, etc.

5.1. Getting a positive measure set of good parameters

This subsection is devoted to the proof of Theorem C. In order to find a set of positive measure of good Collet-Eckmann rational maps, our strategy is as follows: first, we deduce from the main result of [Asp2] the existence of a positive measure set of Collet-Eckmann rational maps satisfying all of the previous conditions with appropriate constants. Then, we follow Tsujii's presentation and improvement [Ts] of the Benedicks-Carleson theory for interval maps [BC]. This will enable us to obtain precise distortion estimates for those good parameters, from which we will deduce that the large scale condition is satisfied.

From now on and through all of this section, f_0 is a fixed Collet-Eckmann rational map such that each critical point of f_0 is simple. We can parametrize analytically a neighborhood O of f_0 in parameter space by means of an analytic map $F : O \times \mathbb{P}^1 \rightarrow \mathbb{P}^1$, and up to restricting O we may assume that each of the following conditions hold for a certain constant $\kappa > 0$:

- (K1) : $\|F\|_{\mathcal{C}^2} < \kappa$.
- (K2) : $\frac{1}{\kappa} < \frac{|f'_\lambda(z)|}{d(z, \mathcal{C}(f_\lambda))} < \kappa$, if $z \notin \mathcal{C}(f_\lambda)$.
- (K3) : $|\partial_z \log f'_\lambda(z)| < \frac{\kappa}{d(z, \mathcal{C}(f_\lambda))}$, if $z \notin \mathcal{C}(f_\lambda)$.
- (K4) : $d(c_i(\lambda), c_j(\lambda')) > \frac{5}{\kappa}$, for all $\lambda' \in O$ and $i \neq j$.
- (K5) : $\|c_i\|_{\mathcal{C}^1} < \kappa$, for all i .

(K6) : if $0 < r < \kappa^{-1}$ and $c \in \mathcal{C}(f_\lambda)$, then

$$\mathbb{D}(c, \kappa^{-1}\sqrt{r}) \subseteq f_\lambda^{-1}(\mathbb{D}(f_\lambda(c), r)) \subseteq \mathbb{D}(c, \kappa\sqrt{r}).$$

We start by recalling the definition of the distortion of an analytic map.

Definition 5.1. — *Let $\psi : \Omega \rightarrow \mathbb{C}$ be holomorphic. We define its complex distortion by*

$$\text{Dist}(\psi, \Omega) := \sup_{x, y \in \Omega} |\log \psi'(x) - \log \psi'(y)| \in [0, \infty].$$

We are only interested in small values of the distortion, and therefore \log (referring to the principal branch of the logarithm) will always be well-defined on $\psi'(\Omega)$. Let $g_i : U_i \rightarrow U_{i+1}$ be holomorphic maps for $i = 1, \dots, n$. Observe that

$$(6) \quad \text{Dist}(g_n \circ \dots \circ g_1, U_1) \leq \sum_{i=1}^n \text{Dist}(g_i, U_i).$$

Assume that $\psi : \mathbb{D}(a, r) \rightarrow \mathbb{C}$ has $\text{Dist}(\psi, \mathbb{D}(a, r)) \leq 50^{-1}$. It is relatively easy to check that

$$(7) \quad \mathbb{D}\left(\psi(a), \frac{9r}{10}|\psi'(a)|\right) \subseteq \psi(\mathbb{D}(a, r)) \subseteq \mathbb{D}\left(\psi(a), \frac{11r}{10}|\psi'(a)|\right).$$

Definition 5.2. — *Let $U \subseteq \mathbb{C}$ be a connected open set. We define its inner diameter as follows*

$$\delta(U) := \sup_{x, y \in U} \inf_{\Gamma \subseteq U} l(\Gamma),$$

where the \inf is taken over all curves $\Gamma \subseteq U$ which connect x to y .

Combining with the definition of the distortion, we obtain the following upper bound

$$(8) \quad \text{Dist}(\psi, U) \leq \delta(U) \sup_{z \in U} |\partial_z \log \psi'(z)|.$$

Another natural property of the inner diameter is

$$(9) \quad \delta(\psi(U)) \leq \delta(U) \sup_{z \in U} |\psi'(z)|.$$

For any $\lambda \in O$, $n \in \mathbb{N}^*$ and $z \in \mathbb{P}^1 \setminus \bigcup_{j=0}^{n-1} f_\lambda^{-j}(\mathcal{C}(f_\lambda))$, we define

$$(10) \quad a^+ = a^+(z, n, f_\lambda) := \left(400e\kappa^2 \sum_{j=0}^{n-1} \frac{|(f_\lambda^j)'(z)|}{|f_\lambda'(f_\lambda^j(z))|} \right)^{-1}.$$

As an immediate consequence of the previous definition, we observe

$$(11) \quad a^+(z, n; f_\lambda) \cdot |(f_\lambda^n)'(z)| = \left(400e\kappa^2 \sum_{j=0}^{n-1} \frac{1}{|(f_\lambda^{n-j})'(f_\lambda^j(z))| \cdot |f_\lambda'(f_\lambda^j(z))|} \right)^{-1} < \frac{1}{400e\kappa}.$$

Lemma 5.3. — *Let f be a rational map satisfying (K2), (K3), (K6). Let $\mathbb{D}(y, r) \subseteq \mathbb{D}(z, a^+)$. Then for all $0 \leq j < m \leq n$*

$$\text{Dist}\left(f_\lambda^{m-j}, f_\lambda^j(\mathbb{D}(y, r))\right) < \frac{r}{100a^+}.$$

Proof. — We show by induction that for all $j = 0, \dots, n-1$

$$(12) \quad \text{Dist} \left(f_\lambda, f_\lambda^j(\mathbb{D}(y, r)) \right) < 4e\kappa^2 r \frac{|(f_\lambda^j)'(z)|}{|f_\lambda'(f_\lambda^j(z))|},$$

and conclude by the definition (10) of a^+ and the bound (6). Let $0 \leq j < n$ and assume (12) holds for all $0 \leq i < j$ and all $\mathbb{D}(y, r) \subseteq \mathbb{D}(z, a^+)$. Using (6),

$$\text{Dist}(f_\lambda^j, \mathbb{D}(z, a^+)) \leq \sum_{i=0}^{j-1} \text{Dist}(f_\lambda, f_\lambda^i(\mathbb{D}(z, a^+))) < 10^{-2},$$

thus, by (K2)

$$\frac{\text{diam}(f_\lambda^j(\mathbb{D}(z, a^+)))}{d(f_\lambda^j(z), C(f_\lambda))} < 2e^{0.01} \kappa a^+ \frac{|(f_\lambda^j)'(z)|}{|f_\lambda'(f_\lambda^j(z))|} < 10^{-2}.$$

As a consequence of the above inequalities, using bounds (8), (K3), (K2) and (9), we obtain

$$\begin{aligned} \text{Dist}(f_\lambda, f_\lambda^j(\mathbb{D}(y, r))) &< \kappa \frac{\delta(f_\lambda^j(\mathbb{D}(y, r)))}{d(f_\lambda^j(\mathbb{D}(y, r)), C(f_\lambda))} \leq \kappa \frac{\delta(f_\lambda^j(\mathbb{D}(y, r)))}{d(f_\lambda^j(\mathbb{D}(z, a^+)), C(f_\lambda))} \\ &< 1.01\kappa \frac{\delta(f_\lambda^j(\mathbb{D}(y, r)))}{d(f_\lambda^j(z), C(f_\lambda))} < 1.01e^{0.01} \kappa^2 \frac{|(f_\lambda^j)'(y)| \delta(\mathbb{D}(y, r))}{|f_\lambda'(f_\lambda^j(z))|} \\ &< e^{0.03} \kappa^2 \frac{2r |(f_\lambda^j)'(z)|}{|f_\lambda'(f_\lambda^j(z))|}, \end{aligned}$$

which completes the inductive proof. \square

For any set $A \subseteq \mathbb{P}^1$ and any $r > 0$, we denote the r -neighborhood of A by

$$B(A, r) := \bigcup_{z \in A} \mathbb{D}(z, r)$$

and its punctured neighborhood by

$$B(A, r)^* := B(A, r) \setminus A.$$

Let $C(f, \delta) := B(C(f), \delta)$ and $C(f, \delta)^* := B(C(f), \delta)^*$.

For $z \in C(f, \kappa^{-1})^*$, we define

$$(13) \quad k(z) := \min\{k \geq 1; \log |(f^k)'(f(c))| > 1 - 1.9 \log |f'(z)|\},$$

where c is the closest critical point of f to z .

Lemma 5.4. — *Let $\alpha, \gamma, \gamma_0, \mu, \mu_0 > 0$, and let f be a rational map satisfying $BA(\alpha)$, $CE(\gamma, \gamma_0)$ and $CE2(\mu, \mu_0)$. If $\alpha < \frac{\gamma}{200}$, there exist $\delta_0 > 0$ such that for all $0 < \delta < \delta_0$ and $z \in C(f, \delta)^*$*

- (1) $-\frac{\log |f'(z)|}{\log \kappa} < k(z) + 1 < -2 \frac{\log |f'(z)|}{\gamma}$,
- (2) $\log |(f^{k(z)+1})'(z)| > -0.9 \log |f'(z)| > \frac{\gamma}{3}(k(z) + 1) + \gamma_0 + \mu_0 + 1$,
- (3) $\log |f'(f^j(z))| > 0.1 \log |f'(z)|$ for all $0 < j \leq k(z)$,
- (4) $\left| \log \left| \frac{f^j(f(z)) - f^j(f(c))}{f(z) - f(c)} \right| - \log |(f^j)'(f(c))| \right| \leq 1$, for all $0 \leq j \leq k(z)$.

Proof. — Let c be the critical point which is closest to z and $v := f(c)$. Claim (1) follows from (K1) and (CE). From (BA) we deduce that

$$\log |f'(f^i(v))| > -k(z) \frac{\gamma}{200}, \text{ for all } 0 \leq i \leq k(z),$$

provided δ is sufficiently small and thus $k(z)$ is large. Combined with claim (1), we obtain

$$\log |f'(f^i(v))| > 10^{-2} \log |f'(z)|, \text{ for all } 0 \leq i \leq k(z).$$

By (K2)

$$(14) \quad d(f(z), v) < \kappa^3 |f'(z)|^2.$$

By the definition (10), we immediately obtain

$$(15) \quad a^+(z, n; f) \geq \left(400e\kappa^2 n \max_{0 \leq j < n} \frac{|(f^j)'(z)|}{|f'(f^j(z))|} \right)^{-1}.$$

Combining the previous three inequalities with the definition (13) of $k(z)$, we obtain

$$(16) \quad d(f(z), v) < a^+(v, k(z); f).$$

All other claims follow from Lemma 5.3 in v . \square

Remark. — An inspection of the proof of Lemma 5.4 shows that we did not in fact use the property CE2. However, in further applications of this lemma, we will apply it to rational maps that do satisfy CE2, and we will need the estimate from item (2) in which μ_0 appears.

We are now able to reconcile the difference in the definition of the conditions FA and FA':

Lemma 5.5. — *Let $\gamma, \gamma_0, \mu, \mu_0 > 0$, let $\alpha < \gamma/200$ and let $f_\lambda \in \mathcal{O}$ satisfying $CE(\gamma, \gamma_0)$, $CE2(\mu, \mu_0)$, $BA'(\alpha)$ and (K1-6). Let $\tau > 0$. There exist $\hat{\delta} > 0$ and $\hat{\beta} > 0$ such that if f_λ satisfies $FA'(\delta, \beta, \tau)$ with $\delta < \hat{\delta}$ and $\beta < \hat{\beta}$, then f_λ satisfies $FA(\eta, \iota)$ with $\eta := \delta^2$ and $\iota := \tau \log \kappa \cdot \max \left(1, 3^{\frac{\beta + \log \kappa}{\gamma + \beta}} - 1 \right)$.*

Proof. — Let $\hat{\delta}^2 := \delta_0$ from Lemma 5.4, and let $\hat{\beta} := 0.05\gamma$. Let $f := f_\lambda$ be a rational map satisfying all the above conditions and $FA'(\beta, \delta, \tau)$ with $\delta < \hat{\delta}$ and $\beta < \hat{\beta}$. Let ν be a deep return for a critical value v' , i.e. $d(f^\nu(v'), c) \leq \delta^2$ for some critical point c . Let $v = f(c)$.

Let $k_0 := k(f^\nu(v'))$ and $z_0 = f^{\nu+1}(v')$. Then, by (16), $z_0 \in D := \mathbb{D}(v, a^+(v, k(f^\nu(v'))))$. We claim that

$$(17) \quad k_0 + 1 < p$$

where p is the length of the bound period. In order to prove the claim, we need to prove that for all $j \leq k_0$, $|f^{j+\nu}(v') - f^j(c)| < e^{-\beta j}$ (by definition of p). Let $z := f^\nu(v)$.

By Lemma 5.4(4), we have that for all $j \leq k_0$,

$$|f^{j+\nu}(v') - f^j(c)| \leq e |(f^j)'(f(c))| \cdot |f(z) - f(c)|$$

and by (K6), we have $d(f(z), f(c)) \leq \kappa d(z, c)^2 \leq \kappa^3 |f'(z)|^2$. By definition of k_0 , we also have $|(f^j)'(f(c))| \leq e \cdot |f'(z)|^{-1.9}$. Therefore we have

$$|f^{j+\nu}(v') - f^j(c)| \leq \kappa^3 |f'(z)|^{0.1}.$$

On the other hand, by Lemma 5.4(1),

$$e^{-\beta j} \geq e^{-\beta k_0} \geq |f'(z)|^{2\beta/\gamma}.$$

By the choice of $\hat{\beta}$, we have $\frac{2\hat{\beta}}{\gamma} \leq 0.1$, and therefore, up to taking an even smaller $\hat{\delta}$, we have proved the claim (17).

We consider the following sets:

$$\begin{aligned} V_k &:= D \cap f^{-k}(\mathbb{D}(f^k(v), 3e^{-k\beta})), & V^k &:= \bigcap_{j=k_0}^k V_j, & V &:= V^p \\ U_k &:= D \cap f^{-k}\left(\mathbb{D}\left(v, \frac{2e^{-k\beta}}{|(f^k)'(v)|}\right)\right), & U^k &:= \bigcap_{j=k_0}^k U_j, & U &:= U^p \\ W_k &:= D \cap f^{-k}(\mathbb{D}(f^k(v), e^{-k\beta})), & W^k &:= \bigcap_{j=k_0}^k W_j, & W &:= W^p \end{aligned}$$

Let us show that for all $k_0 \leq k \leq p$

$$(18) \quad \text{Dist}(f^k, V^k) < \frac{1}{50}.$$

By inequality (6):

$$\begin{aligned} \text{Dist}(f^k, V^k) &\leq \text{Dist}(f^{k_0}, V^k) + \sum_{j=k_0}^{k-1} \text{Dist}(f, f^j(V^k)) \\ &\leq \text{Dist}(f^{k_0}, D) + \sum_{j=k_0}^{k-1} \text{Dist}(f, \mathbb{D}(f^j(v), 3e^{-j\beta})). \end{aligned}$$

By Lemma 5.3, the first term is $< 10^{-2}$. By Lemma 2.1 in [Asp2], it is enough to show that for all $y \in \mathbb{D}(f^j(v), 3e^{-j\beta})$

$$\sum_{j=k_0}^{k-1} \left| \frac{f'(y)}{f'(f^j(v))} - 1 \right| < 10^{-3}.$$

By (BA) and (K1), for $y \in \mathbb{D}(f^j(v), 3e^{-j\beta})$, one has:

$$\left| \frac{f'(y)}{f'(f^j(v))} - 1 \right| < \kappa 3e^{-j(\beta-\alpha)}.$$

By Lemma 5.4(1) and (K2), $k_0 > -\frac{\log |f'(f^{k_0}(v))|}{\log \kappa} - 1 > -\frac{2 \log \delta}{\log \kappa} - 2$. So, by Lemma 2.1 in [Asp2]:

$$\begin{aligned} \sum_{j=k_0}^{k-1} \left| \frac{f'(y)}{f'(f^j(v))} - 1 \right| &< \sum_{j=k_0}^{k-1} \kappa e^{-j(\beta-\alpha)} \\ &< \frac{3\kappa e^{-k_0(\beta-\alpha)}}{1 - e^{-(\beta-\alpha)}} < \frac{\kappa e^{2\frac{(\beta-\alpha)}{\log \kappa}}}{1 - e^{-(\beta-\alpha)}} \end{aligned}$$

which completes our argument, as δ can be taken arbitrarily small.

We show that $U^k \subset V^k$ for all $k_0 \leq k \leq p$. If not, let $k_0 < k$ such that there is $x \in U^k \setminus V^k$. Taking k and x minimal, we assume that $[x, v] \in V^{k-1}$, $[v, x] \subset V^k$ and $x \in \partial V_k$. In particular,

$$\text{Dist}(f^k, [v, x]) \leq \frac{1}{50}$$

hence, as $x \in U_k$:

$$3e^{-k\beta} = |f^k(x) - f^k(v)| \leq 1.03|x - v| \cdot |(f^k)'(v)| \leq 2.06e^{-k\beta},$$

a contradiction.

We show now that $W^k \subset U^k$ for all $k_0 \leq k \leq p$. If not, let $k_0 < k$ and $x \in W^k$ minimal, such that $[x, v] \in U^k \subset V^k$, $[v, x] \subset V^k$ and $x \in \partial U_k$. In particular,

$$\text{Dist}(f^k, [v, x]) \leq \frac{1}{50}$$

hence, as $x \in \partial U_k$:

$$e^{-k\beta} > |f^k(x) - f^k(v)| \leq 0.97|x - v| \cdot |(f^k)'(v)| = 1.94e^{-k\beta},$$

a contradiction.

In particular, we have $W \subset U \subset V$ therefore $[v, z_0] \subset V$ which implies that $\forall k_0 \leq k \leq p$:

$$(19) \quad 0.97 \cdot |z_0 - v| \cdot |(f^k)'(v)| < |f^k(z_0) - f^k(v)| < 1.03 \cdot |z_0 - v| \cdot |(f^k)'(z_0)|$$

We have seen that $p > k_0 + 1 > -\frac{\log |f'(f^{\nu}(v))|}{\log \kappa}$. Thus, by (FA'):

$$(20) \quad \sum_{i=0}^s -\log |f'(f^{\nu_i}(v))| < \log \kappa \sum_{i=0}^s p_i(v) < \tau \log \kappa (\nu_s(v) + p_s(v)).$$

By inequalities (K2) and (19) applied for $j = p - 1$, (CE) and the definition of p ,

$$\kappa^{-2} |f'(z)|^2 < |f(z) - f(c)| < e^{\gamma + \gamma_0 - p(\gamma + \beta) + 1},$$

thus

$$p < -\frac{3}{\gamma + \beta} \log |f'(z)|.$$

Let Σ_1 and Σ_2 be the sums of $\log |f'(f^j(z))|$ respectively over the sets

$$\{0 < j \leq p : f^j(z) \in C(f, \delta^2)\} \text{ and } \{0 < j \leq p : f^j(z) \notin C(f, \delta^2)\}.$$

Observe that $\Sigma := \Sigma_1 + \Sigma_2 = \log |(f^p)'(z_0)|$. By (K2), inequalities (19) and (7) and the definition of p , we have

$$\Sigma + 2 \log |f'(z)| > \Sigma + \log |f(z) - f(c)| - 2 \log \kappa \geq -p\beta - 3 \log \kappa,$$

thus by (K1)

$$\begin{aligned} -\Sigma_1 &< \Sigma_2 + 2 \log |f'(z)| + p\beta + 3 \log \kappa < \log |f'(z)| + p(\beta + \log \kappa) \\ &< -\log |f'(z)| \left(3 \frac{\beta + \log \kappa}{\gamma + \beta} - 1 \right). \end{aligned}$$

Combined with inequality (20) it shows that Aspenberg's (FA') implies (FA) with constants $\eta := \delta^2$ and $\iota := \tau \log \kappa \cdot \max \left(1, 3 \frac{\beta + \log \kappa}{\gamma + \beta} - 1 \right)$. \square

By the main result of [GS], Collet-Eckmann rational maps satisfy the backward Collet-Eckmann condition at critical points of maximal multiplicity. In our setting, all critical points of f_0 are simple. However, we need uniform constants for the backward contraction in a neighborhood of O of f_0 .

Let $\gamma, \gamma_0 > 0$ such that f_0 satisfies $CE(\gamma, \gamma_0)$. A careful inspection of the proof of Proposition 1 in [GS] reveals the following facts. Dynamically defined constants M, R' and R can be chosen to be uniform in a neighborhood of f_0 . They also depend on λ_1 and C_1 , which in our setting are expressed by constants γ, γ_0 . A \mathcal{C}^1 continuity argument, similar to the proof of the above proposition, shows that constants C_{2t} and L can be chosen uniformly for maps f_λ satisfying $CE(\gamma, \gamma_0)$ in a neighborhood of f_0 . Observe that the constant $K_{3.1}$ only depends on R' .

This observation proves the following:

Proposition 5.6. — *There exist O , a neighborhood of f_0 in parameter space, and constants $\mu, \mu_0 > 0$ such that every f_λ with $\lambda \in O$ satisfying $CE(\gamma, \gamma_0)$ also satisfies $CE2(\mu, \mu_0)$.*

We now restate for the reader's convenience the main result from [Asp2]:

Theorem 5.7 (Aspenberg, [Asp2]). — *Let f_0 be a strongly Misiurewicz rational map with simple critical points. If f_0 is not a flexible Lattès map, there exist $\gamma, \gamma_0 > 0$ and $\hat{\alpha}, \hat{\delta}, \hat{\tau} > 0$ such that for all $\alpha < \hat{\alpha}, \delta < \hat{\delta}, \tau < \hat{\tau}$, the map f_0 is a Lebesgue point of density of rational maps satisfying $CE(\gamma, \gamma_0)$, $BA'(\alpha)$, and $FA'(\beta, \delta, \tau)$ with $\beta := 10\alpha$.*

Putting together those results, we are now ready to prove Theorem C:

Proof of Theorem C. — Let $\gamma, \gamma_0 > 0$ be the constants given by Theorem 5.7. Let $\mu, \mu_0 > 0$ be the constants given by Proposition 5.6. Let $\hat{\eta} := \hat{\delta}^2$ and

$$\hat{\iota} := \sup_{0 < \beta \leq 10\hat{\alpha}} \hat{\tau} \log \kappa \cdot \max \left(1, 3 \frac{\beta + \log \kappa}{\gamma + \beta} - 1 \right).$$

Let $\alpha < \min(\hat{\alpha}, \gamma/200)$, $\eta < \hat{\eta}$ and $\iota < \hat{\iota}$.

Let E be the set of rational maps satisfying $CE(\gamma, \gamma_0)$, $BA'(\alpha/\kappa)$, and $FA'(\beta, \delta, \tau)$ with $\beta := 10\alpha$, $\delta^2 := \eta$, and $\tau > 0$ small enough to ensure that $\tau \max \left(1, 3 \frac{\beta + \log \kappa}{\gamma + \beta} - 1 \right) \leq \iota$.

By Lemma 5.5, each $f \in E$ satisfies $FA(\eta, \iota)$. By Proposition 5.6, each $f \in E$ satisfies $CE2(\mu, \mu_0)$. By (K2), each $f \in E$ satisfies $BA(\alpha)$. By Theorem 5.7, f_0 is a Lebesgue density point of E . \square

5.2. Proving the large scale condition for good parameters

In the present section, we prove Theorem D. Let \mathcal{F} be the space of complex lines passing through 0 in \mathbb{C}^N and let \mathcal{L} be the unique volume form on \mathcal{F} which is invariant under the action of the unitary group $U(N)$ and of total mass 1. We shall rely on the following result of Sibony and Wong [SW].

Theorem 5.8 (Sibony-Wong). — *Let $m > 0$ be a positive constant. Let $\mathcal{F}' \subset \mathcal{F}$ be such that $\mathcal{L}(\mathcal{F}') \geq m$ and let Σ denote the intersection of the family \mathcal{F}' with the ball $\mathbb{B}(0, r)$ of \mathbb{C}^N . Then any holomorphic function h on a neighborhood of Σ can be extended to a holomorphic function on $\mathbb{B}(0, \tilde{\rho}r)$ where $0 < \tilde{\rho} \leq 1$ is a constant depending on m and N but independent of \mathcal{F}' and r . Moreover, we have*

$$\sup_{\mathbb{B}(0, \tilde{\rho}r)} |h| \leq \sup_{\Sigma} |h|.$$

Another key ingredient will be a transversality property of the differentials of the ξ_n^i . This property first appeared in the work of Tsujii in the real dynamics case. It was proved for some class of rational maps (containing notably Collet-Eckmann polynomials) by Levin in [Le]. In [Ast], this property was extended to rational maps having no invariant line fields, and so (see Lemma 5.14) this property is satisfied by all Collet-Eckmann rational maps.

From now on, we will not rely on the result of Aspenberg ([Asp2]). We will also not make any more use of the conditions FA' or BA' . Instead, we fix in this whole subsection some rational map f satisfying $CE(\gamma, \gamma_0)$, $CE2(\mu, \mu_0)$, $BA(\alpha)$ and $FA(\eta, \iota)$. We denote by \mathbb{S}_{2d-2} the unit sphere in \mathbb{C}^{2d-2} .

The main goal of this section is to prove the following distortion estimate for rational maps f satisfying all the previous conditions:

Proposition 5.9. — *Let $\gamma, \gamma_0, \mu, \mu_0 > 0$ and $\alpha < \gamma/200$. Let $\eta > 0$. There exists $\iota > 0$ such that if f satisfy $CE(\gamma, \gamma_0)$, $CE2(\mu, \mu_0)$, $BA(\alpha)$ and $FA(\eta, \iota)$, then there exist $C > 0$, a set*

$E \subset \mathbb{S}_{2d-2}$ such that $\lambda(E) > 0$ and an infinite set $G \subseteq \mathbb{N}$ of natural numbers with the following property: for every $n \in G$ and all $c_i \in \mathcal{C}(f)$ there exists $r_{n,i} > 0$ such that for every $u \in E$

$$\text{Dist}(\xi_n^i, \mathbb{D}(0, r_{n,i}) \cdot u) < 10^{-2}, \text{ and}$$

$$\frac{1}{C} < r_{n,i} |(f^n)'(f(c_i))| < C.$$

5.3. Distortion estimates for a single CE map

Lemma 5.10. — Assume $\alpha < \gamma/200$. There exist $\delta_1 > 0$ such that for all $0 < \delta < \delta_1$, if $n > 0$ and $z \in \mathbb{P}^1$ satisfy

$$(21) \quad f^j(z) \notin C(f, \delta) \text{ for all } 0 \leq j < n \text{ and } f^n(z) \in C(f, \delta),$$

we have

$$|(f^n)'(z)| > e^{n\mu - \mu_0 - 1}.$$

Proof. — For $\delta > 0$ and $n \in \mathbb{N}$ denote

$$A(n, \delta) := f^{-n}(\mathcal{C}(f)) \setminus \bigcup_{j=0}^{n-1} f^{-j}(C(f, \delta^2)).$$

Let us prove that if W is a connected component of $f^{-n}(C(f, \delta))$ which intersects $A(n, \delta)$ then

$$(22) \quad \text{Dist}(f^n, W) < 10^{-2}.$$

Let $z \in W \cap A(n, \delta)$, $l := -2\mu^{-1} \log \delta$ and $\Sigma_1, \Sigma_2, \Sigma_3$ the sums of

$$|(f^{n-j})'(f^j(z)) \cdot f'(f^j(z))|^{-1}$$

on the following sets of $j \in \mathbb{N}$, respectively:

$$\{n-l \leq j < n : f^j(z) \in C(f, \sqrt{\delta})\},$$

$$\{n-l \leq j < n : f^j(z) \notin C(f, \sqrt{\delta})\}, \{0 \leq j < n-l\}.$$

If δ is small and $y \in C(f, \sqrt{\delta})$ then Lemma 5.4, claims (3) and (4), provides $k > 0$ such that $\log |(f^{k+1})'(y)| > -0.9 \log |f'(y)|$ and for all $0 < j \leq k$, $f^j(y) \notin C(f, \delta)$. Combined with (K2) and (CE2) at $f^n(z) \in \mathcal{C}(f)$, we obtain

$$\Sigma_1 < l \cdot \left(\frac{\delta^2}{\kappa}\right)^{-0.1} e^{\mu_0}, \quad \Sigma_2 < l \cdot \kappa \delta^{-0.5} e^{\mu_0}, \quad \Sigma_3 < \sum_{j>l} \kappa \delta^{-2} e^{-j\mu + \mu_0}.$$

By inequality (11)

$$a^+(z, n, f) \cdot |(f^n)'(z)| = (400e\kappa^2(\Sigma_1 + \Sigma_2 + \Sigma_3))^{-1} > 2\delta.$$

Lemma 5.3 and observation (7) complete the proof of claim (22).

Let $z \in \mathbb{P}^1$ satisfying (21) for some $n > 0$. Let $0 \leq k_0 \leq n$ be the smallest integer such that $f^{k_0}(z) \in W$, a connected component of $f^{-n+k_0}(C(f, \delta))$ which contains $z' \in A(n - k_0, \delta)$. We prove that $k_0 = 0$ and use inequality (22) and (CE2) to complete the proof.

Suppose $k_0 > 0$. Let J be the connected component of $f^{-1}(W)$ containing $y := f^{k_0-1}(z)$. There are two possibilities.

Case 1. There exists $c \in \mathcal{C}(f) \cap J$. Then by (21), $y \notin \mathbb{D}(c, \delta)$ and by (K6) $\text{diam}(W) \geq d(f(y), f(c)) > \kappa^{-2}\delta^2$. Then by (22) and (7)

$$|(f^{n-k_0})'(f(c))| < 2 \frac{2\delta}{\text{diam}(W)} < 4\kappa^2\delta^{-1}.$$

Then by (CE)

$$\gamma(n - k_0) < \log |(f^{n-k_0})'(f(c))| + \gamma_0 < \log(\delta^{-1}) + \log(4\kappa^2) + \gamma_0.$$

As $f^{n-k_0+1}(c) \in C(f, \delta)$, by (BA) and (K2) we obtain

$$-(n - k_0)\alpha < \log |f'(f^{n-k_0+1}(c))| < \log \delta + \log \kappa.$$

As $\alpha < \gamma/200$, taking δ sufficiently small, we obtain a contradiction.

Case 2. $J \cap \mathcal{C}(f) = \emptyset$. Let $z'' := (f|_J)^{-1}(z') \notin A(n - k_0 + 1, \delta)$. There exists $c \in \mathcal{C}(f)$ with $d(z'', c) < \delta^2$. As $c \notin J$, we may choose $x \in [z'', c] \cap \partial J$. By (K6), $d(z', f(x)) < \kappa^2 \delta^4$. As $f(x) \in \partial W$, by (22) and (7) applied to $f^{-(n-k_0)}$

$$W \subseteq \mathbb{D}(z', 2\kappa^2 \delta^4) \text{ and hence } J \subseteq \mathbb{D}(z'', \sqrt{2}\kappa^2 \delta^2).$$

But this implies $d(y, c) \leq \text{diam}(J) + d(z'', c) < \delta$, which contradicts (21). \square

Let $\delta_2 := \min(\delta_0, \delta_1)$ such that lemmas 5.4 and 5.10 apply for all $\delta < \delta_2$.

We will use the following variant of a well known result due to Mañé [Ma]. We provide a short proof using the ExpShrink property of (CE) maps [PRLS, Main Theorem]: there exist $r, \gamma' > 0$ such that for all $z \in \mathcal{J}$ (the Julia set of f), every $n > 0$ and every connected component W of $f^{-n}(\mathbb{D}(z, r))$

$$\text{diam}(W) < e^{-n\gamma'}.$$

Proposition 5.11. — *Let $K := \mathbb{P}^1 \setminus C(f, \delta_2)$ be a compact set. There exist $\gamma', \gamma'_0 > 0$ such that for all $n \in \mathbb{N}$ and $z \in \cap_{i=0}^{n-1} f^{-i}(K)$*

$$|(f^n)'(z)| > e^{n\gamma' - \gamma'_0}.$$

Proof. — As all critical orbits of f satisfy the Collet-Eckmann condition, by the classification of Fatou domains, $\mathcal{J} = \mathbb{P}^1$. Using ExpShrink, by continuity and by eventually shrinking r , we may assume that for all $i < n$

$$\text{diam}(f^i(W)) < \frac{d(K, \mathcal{C}(f))}{2},$$

where W is an arbitrary connected component of $f^{-n}(\mathbb{D}(z, r))$. Then f^n is univalent on W and we conclude by the Koebe lemma applied to f^{-n} on $\mathbb{D}(z, r/2)$ that if n is large enough and $f^i(z) \in K$ for $i = 0, \dots, n-1$ then

$$|(f^n)'(z)| > 3.$$

which completes the proof. \square

Let γ' and γ'_0 be provided by Proposition 5.11. Let

$$(23) \quad \sigma := \min(\gamma/3, \mu, \gamma') \text{ and } \sigma_0 := \log \kappa + \gamma' + \mu_0 + \gamma'_0 + 1.$$

Lemma 5.12. — *Let $z \in \mathbb{P}^1$ and $n > 0$ such that for all $0 \leq i < n$, $f^i(z) \notin \mathcal{C}(f)$. Let $I := \{0 \leq i < n : f^i(z) \in C(f, \delta_2) \text{ and } i + k(f^i(z)) \geq n\}$. Let $\beta := 1$ if $I = \emptyset$, otherwise $\beta := \min\{d(f^i(z), \mathcal{C}(f)) : i \in I\}$. Then*

$$\log |(f^n)'(z)| > \log \beta + n\sigma - \sigma_0.$$

Proof. — If for all $0 \leq i < n$, $f^i(z) \notin C(f, \delta_2)$, $I = \emptyset$ and we apply Proposition 5.11 to complete the proof.

Let $i_1 := \min\{0 \leq i < n : f^i(z) \in C(f, \delta_2)\}$ and for each $p \geq 1$, $i_{p+1} := \min\{i_p + k(f^{i_p}(z)) < i < n : f^i(z) \in C(f, \delta_2)\}$. Let s be the largest index such that $i_s < \infty$.

We use Lemma 5.10 at z and obtain

$$\log |(f^{i_1})'(z)| > i_1\mu - \mu_0 - 1 \geq i_1\sigma - \mu_0 - 1.$$

Let $k_p := k(f^{i_p}(z))$ for all $1 \leq p \leq s$. By Lemma 5.4(2), we get

$$\log |(f^{k_1+1})'(f^{i_1}(z))| > (k_1 + 1)\frac{\gamma}{3} + \mu_0 + 1 \geq (k_1 + 1)\sigma + \mu_0 + 1.$$

We iterate this chaining of lemmas 5.10 and 5.4 from $i_{p-1} + k_{p-1} + 1$ to i_p and respectively from i_p to $i_p + k_p + 1$.

The only exception occurs for $p = s$ if $i_s \in I$. In this case, we stop at i_s and estimate directly by (K2)

$$\log |f'(f^{i_s}(z))| > \log d(f^{i_s}(z), \mathcal{C}(f)) - \log \kappa \geq \log \beta - \log \kappa.$$

We apply Proposition 5.11 from $i_s + 1$ to n and obtain

$$\log |(f^{n-i_s-1})'(f^{i_s+1}(z))| > (n - i_s - 1)\gamma' - \gamma'_0 \geq (n - i_s)\sigma - \gamma' - \gamma'_0.$$

If $i_s \notin I$, then we apply Proposition 5.11 from $i_s + k_s + 1$ to n . We conclude by summing up all above estimates. \square

Recall that η, ι characterize the (FA) condition on f . Shrinking η only improves condition (FA), so we may assume $\eta \leq \delta_2$.

Lemma 5.13. — *There exists $\rho > 0$ such that for all $v \in f(\mathcal{C}(f))$ and $m > 0$, there exists $G \subseteq \mathbb{N} \cap (m, 2m]$ with*

$$(24) \quad \frac{|G|}{m} > 1 - \frac{4\iota}{\gamma} \left(1 + 1.2 \frac{\log \kappa}{\sigma} \right),$$

such that for all $n \in G$

$$(25) \quad |(f^n)'(v)| \cdot a^+(v, n; f) > \rho.$$

Proof. — Fix $v \in f(\mathcal{C}(f))$ and let $v_j := f^j(v)$ for $j \geq 0$. For all $n > 0$ let $R(n) := \{0 \leq j < n : v_j \in \mathcal{C}(f, \eta)\} = \{j_1 < j_2 < \dots < j_s\}$. Note that s depends on n . Note that by (FA), for all $n > 0$

$$(26) \quad \sum_{j \in R(n)} \ln |f'(v_j)| > -n\iota.$$

For $i = 1, \dots, s$, let $k_i := k(v_{j_i})$, $J'_i := \mathbb{N} \cap (j_i, j_i + k_i + 1]$ and $J'(n) := \cup_{i=1}^s J'_i$. For any $m \geq 0$ we define

$$G' := \mathbb{N} \cap (m, 2m] \setminus J'(2m).$$

Let $\theta > 1$, $J_i := \mathbb{N} \cap (j_i, j_i + \theta(k_i + 1)]$ and $J(n) := \cup_{i=1}^s J_i$. As before,

$$G := \mathbb{N} \cap (m, 2m] \setminus J(2m).$$

Let us denote

$$d_j^n := \log |(f^{n-j})'(v_j)|, \quad d_j := d_j^{j+1} \quad \text{and} \quad d^n := d_0^n.$$

By Lemma 5.4(1), $k_i + 1 < -2\gamma^{-1}d_{j_i}$, thus by (26)

$$|J'(2m)| < 4\frac{m\iota}{\gamma} \quad \text{and} \quad |J(2m)| < 4\frac{m\iota\theta}{\gamma}.$$

It suffices to choose

$$\theta := 1 + 1.2 \frac{\log \kappa}{\sigma}$$

and show inequality (25) for all $n \in G \subseteq G'$.

We employ Lemma 5.12 to estimate d_j^n for all $j < n$. First, observe that as $n \in G'$, $\beta \geq \eta$ for all $0 \leq j < n$. Therefore, for all $j < n$ we have

$$(27) \quad d_j^n > \log \eta + (n - j)\sigma - \sigma_0.$$

By Lemma 5.4(2) and (27) for $j = j_i + k_i + 1$

$$d_{j_i} + d_{j_i}^n > 0.1d_{j_i} + d_{j_i+k_i+1}^n > 0.1d_{j_i} + (n - j_i - k_i - 1)\sigma - \sigma_0.$$

If $n \in G$, then by Lemma 5.4(1) $n - j_i - k_i - 1 > (\theta - 1)(k_i + 1) > -d_{j_i} \frac{\theta - 1}{\log \kappa}$. Combine the previous two inequalities with the choice of θ to obtain

$$(28) \quad d_{j_i} + d_{j_i}^n > -d_{j_i} + (n - j_i - \theta(k_i + 1))\sigma > -d_{j_i},$$

as $-d_{j_i} > -\log \eta - \log \kappa$.

To complete the proof, using inequality (11), we need to bound

$$S(n) := \sum_{j=0}^{n-1} \frac{1}{|(f^{n-j})'(v_j)| \cdot |f'(v_j)|} = \sum_{j=0}^{n-1} \exp(-d_j - d_j^n).$$

Let $S_0(n) := \sum_{j \in R(n)} \exp(-d_j - d_j^n)$ and $S_1(n) := S(n) - S_0(n)$. By (27) and (K2),

$$S_1(n) < \kappa \eta^{-2} e^{\sigma_0} \sum_{j=0}^{n-1} e^{-j\sigma}.$$

Let $Q(n) := \{j \in R(n) : -d_j > (n - j)\sigma/2\}$. Then by inequalities (28) and (27) we can estimate

$$\begin{aligned} S_0(n) &= \sum_{j \in Q(n)} \exp(-d_j - d_j^n) + \sum_{j \in R(n) \setminus Q(n)} \exp(-d_j - d_j^n) \\ &< \sum_{j \in Q(n)} e^{-\frac{(n-j)\sigma}{2}} + \sum_{j \in R(n) \setminus Q(n)} e^{-\frac{(n-j)\sigma}{2} - \log \eta + \sigma_0} < \eta^{-1} e^{\sigma_0} \sum_{j=0}^{n-1} e^{-j\frac{\sigma}{2}}. \end{aligned}$$

This ends the proof. □

Remark. — The scale ρ only depends on σ, σ_0 and η .

5.4. Using the transversality property

From now on, we consider a local parametrization of the moduli space near f , given by a holomorphic injection $F : O \times \mathbb{P}^1 \rightarrow \mathbb{P}^1$ where O is a small neighborhood of the origin in \mathbb{C}^{2d-2} , and $F(0, \cdot) = f$. For all $\lambda \in O$, we denote $f_\lambda := F(\lambda, \cdot)$.

In [Ast], the first author proves a transversality property under the assumption that f has no invariant line fields. We show that it is the case for Collet-Eckmann rational maps which is not a flexible Lattès map. Though it is known by the experts, we did not find a complete proof in the literature, so we provide it here.

Lemma 5.14. — *Let $f \in \text{Rat}_d$ be Collet-Eckmann. If f carries an invariant line field on its Julia set, then f is a flexible Lattès map.*

Proof. — Recall that a point $z_0 \in \mathcal{J}_f$ is said to be a *conical point* of f if there exist $z_j \rightarrow z_0$, integers $n_j \rightarrow +\infty$ and positive numbers $\rho_j \rightarrow 0$ such that $f^{n_j}(z_j + \rho_j \cdot z)$ converges uniformly on \mathbb{D} to some non-constant holomorphic function (see, e.g., [BM] page 109). Denote by Λ_f the

set of conical points of f . Przytycki proved (see [P] Proposition A.3.4) that for any $f \in \text{Rat}_d$ which is a Collet-Eckmann rational map, we have $\Lambda_f = \mathcal{J}_f$. In particular $\text{Leb}(\Lambda_f) > 0$.

As a consequence, if f is Collet-Eckmann and carries an invariant line field on its Julia set, by Theorem VII.22 of [BM], this implies that f is a Lattès map. Finally, Corollary 3.18 of [Mc] asserts that if f is a Lattès map and carries an invariant line field on its Julia set, then f is a flexible Lattès map. \square

Therefore, by Theorem B of [Ast] and Lemma 4.4, Collet-Eckmann rational maps satisfy the following property:

Transversality (T) : The following limits exist (as linear forms on $T_0O \simeq \mathbb{C}^{2d-2}$), and they are linearly independent:

$$\tau_i := u \mapsto \lim_{n \rightarrow \infty} \frac{\frac{d}{dt}|_{t=0} \xi_n^i(tu)}{(f^n)'(f(c_i))} = \sum_{j=0}^{\infty} \frac{\frac{d}{dt}|_{t=0} F(tu, f^j(c_i))}{(f^j)'(f(c_i))}$$

where $\xi_n^i(\lambda) := f_\lambda^{n+1}(c_i(\lambda))$.

Note that the equality is just the first part of Lemma 4.4 with different notations. Indeed, observe that $\dot{v}_i = \frac{d}{dt}|_{t=0} F(tu, c_i)$.

Since the (τ_i) are linearly independent, we can choose $2d-2$ unit vectors $u_1, \dots, u_{2d-2} \in \mathbb{C}^{2d-2}$ such that $\tau_i(u_j) \neq 0$ if and only if $i = j$. Moreover, there is an open subset E of \mathbb{S}_{2d-2} such that for all $u \in E$ and all i ,

$$(29) \quad \|\tau_i(u)\| \geq \frac{1}{2d} \min_j \|\tau_j(u_j)\| > 0$$

(we can take E to be a small enough neighborhood of $u = \|\sum_{j=1}^{2d-2} u_j\|^{-1} \sum_{j=1}^{2d-2} u_j$).

We fix some critical point c_i . From the transversality condition (T) and the definition (29) of the set of directions E in parameter space, we get $m_0 > 0$ such that for all $m \geq m_0$ and all $u \in E$, $\partial_u \xi_m^i(0) \neq 0$. For some $r \geq 0$, let us define the corresponding parameter neighborhood of 0 in direction u

$$(30) \quad V(u, r) := (\mathbb{D}(0, r) \cdot u) \cap O.$$

For any $m \geq m_0$ and each i , we say that $\Gamma^u(m) \geq 0$ satisfies (T1), (T2) and (T3) if:

(T1) : if $V := V(u, \Gamma^u(m))$, then for all $m_0 \leq j \leq m$

$$\text{Dist}(\xi_j^i, V) \leq 10^{-2},$$

$$(T2) : \Sigma(u, m, V) := \sum_{j=0}^{m-1} \max_{\alpha, \beta \in V} \left| \log \frac{f'_\alpha(\xi_{j+1}^i(\alpha))}{f'_\beta(\xi_{j+1}^i(\beta))} \right| \leq 10^{-2},$$

(T3) : $\text{diam}(\xi_j^i(V)) \leq \frac{1}{e^\kappa}$ for all $0 \leq j \leq m$.

Combining (T1) and (T3) with (7) we observe that

$$(31) \quad \Gamma^u(m) < |\partial_u \xi_m^i(0)|^{-1}.$$

Recall that $\tau_i(u) \neq 0$ is a condition of the transversality hypothesis (T) and the definition (29) of E . Denote by $v_i = v_i(0) := f_0(c_i(0))$.

Lemma 5.15. — *There exist $l_1 > 1$ and $m_1 > m_0$ such that for all $u \in E$ and $m \geq m_1$, then*

$$(32) \quad \Gamma^u(m) = l_1^{-1} \cdot a^+(v_i, m; f_0),$$

satisfies (T1), (T2) and (T3).

Proof. — By (CE) and (T), there exists $m' > m_0$ such that for all $u \in E$:

$$(33) \quad \kappa M \sum_{j=m'}^{\infty} e^{-j\gamma+\gamma_0+3} < 10^{-3},$$

$$(34) \quad \left| \sum_{j=0}^{m'-1} \frac{\frac{d}{dt}|_{t=0} F(tu, f_0^j(c_i))}{\tau_i(u)(f_0^m)'(v_i)} - 1 \right| < 10^{-3},$$

where $M := 1 + \max\{|\tau_i(u)|, |\tau_i(u)|^{-1} : u \in E\}$. Furthermore, for $k \geq m'$, one has by (T) and (CE) that

$$(35) \quad \left| \log \frac{\partial_u \xi_k^i(0)}{\tau_i(u)(f_0^k)'(v_i)} \right| < 10^{-2} \text{ and } |\partial_u \xi_k^i(0)| > 2e\kappa.$$

We choose $0 < \varepsilon_1$ small enough such that

$$\text{diam}(\xi_j^i(V(u, 2\varepsilon_1)) < e^{-1}\kappa^{-1} \text{ for any } u \in E \text{ and all } 0 \leq j \leq m',$$

$$\text{Dist}(\xi_j^i, V(u, 2\varepsilon_1)) < 10^{-3} \text{ for any } u \in E \text{ and all } m_0 \leq j \leq m' \text{ and}$$

$$\Sigma(u, m', V(u, 2\varepsilon_1)) < 10^{-3},$$

recalling that V was defined by (30) and Σ in condition (T2). Put

$$l_1 := 11e\kappa M$$

and choose $m_1 > m'$ such that

$$e^{m_1\gamma_1-\gamma_0} > \max(4e\kappa M, \varepsilon_1^{-1}).$$

We prove the lemma for these l_1 and m_1 .

Take any $m \geq m_1$, $u \in E$. Let

$$\Lambda := V(u, l_1^{-1}a^+(v_i, m; f_0)),$$

$$W := \mathbb{D}(v_i, 10^{-1}a^+(v_i, m; f_0)),$$

and prove by induction the following claim for all $m_0 \leq j \leq m$

$$(*)_j \text{ Dist}(\xi_j^i, \Lambda) < 10^{-2} \text{ and } \Sigma(u, j, \Lambda) < 10^{-2}.$$

Since $a^+(v_i, m; f) < |(f_0^m)'(v_i)|^{-1} < \varepsilon_1$ by the choice of m_1 , we can get $(*)_j$ for $m_0 \leq j \leq m'$ from the choice of ε_1 .

Assume that for some $m' < j \leq m$, $(*)_k$ holds for all $m_0 \leq k < j$. Observe that $\xi_k^i(\Lambda) \subseteq f_0^k(W)$ for all $m' \leq k < j$. On one hand, we use (35), $\text{Dist}(\xi_k^i, \Lambda) < 10^{-2}$, (7) and the choice of l_1 . On the other, we use Lemma 5.3 applied to f_0^m on W and (7). In particular, by Lemma 5.3, for all $\alpha \in \Lambda$ and $m' \leq k < j$

$$(36) \quad \left| \log \frac{f_0'(\xi_k^i(\alpha))}{f_0'(\xi_k^i(0))} \right| < 10^{-3}.$$

Summing (12) for $k = m'$ to $j - 1$ implies:

$$\sum_{k=m'}^{j-1} \text{Dist}(f_0, f_0^k(W)) < 10^{-3}a^+(v_i, m; f)400e\kappa^2 \sum_{k=m'}^{j-1} \frac{|(f_0^k)'(v_i)|}{|f_0'(f_0^k(v_i))|},$$

hence, by definition of $a^+(v_i, m; f)$:

$$(37) \quad \sum_{k=m'}^{j-1} \text{Dist}(f_0, f_0^k(W)) < 10^{-3}.$$

Also, for all $\alpha \in \Lambda$ and $m' \leq k < j$, by (K1), $(*)_k$, (35) and direct calculation, we get

$$\begin{aligned} |f'_\alpha(\xi_k^i(\alpha)) - f'_0(\xi_k^i(\alpha))| &< \kappa|\alpha| < \kappa l_1^{-1} a^+(v_i, m; f) < \kappa |(f_0^k)'(v_i)|^{-1}, \\ \left| \frac{\partial_u \xi_{k+1}^i(\alpha)}{\partial_u \xi_k^i(\alpha)} - f'_\alpha(\xi_k^i(\alpha)) \right| &= \frac{|\frac{d}{dt}|_{t=0} F(\alpha + tu, \xi_k^i(\alpha))|}{|\partial_u \xi_k^i(\alpha)|} < \frac{\kappa e^{0.02}}{|\tau_i(u)(f_0^k)'(v_i)|}. \end{aligned}$$

Divide the last two inequalities by $|f'_0(\xi_k^i(\alpha))|$, use $\text{Dist}(f_0, \xi_k^i(\Lambda)) < 10^{-3}$, the triangle inequality, (CE) and (33) to obtain

$$\left| \frac{\partial_u \xi_{k+1}^i(\alpha)}{\partial_u \xi_k^i(\alpha) f'_0(\xi_k^i(\alpha))} - 1 \right| < \frac{\kappa M e}{|(f_0^{k+1})'(v_i)|} < \kappa M e^{-(k+1)\gamma_1 + \gamma_0 + 2} < 10^{-3},$$

which immediately implies

$$\left| \log \frac{\partial_u \xi_{k+1}^i(\alpha)}{\partial_u \xi_k^i(\alpha)} - \log f'_0(\xi_k^i(\alpha)) \right| < \kappa M e^{-(k+1)\gamma_1 + \gamma_0 + 3}.$$

Observe that

$$\text{Dist}(\xi_{k+1}^i, \Lambda) \leq \text{Dist}(\xi_k^i, \Lambda) + \sup_{\alpha, \beta \in \Lambda} \left| \log \frac{\partial_u \xi_{k+1}^i(\alpha)}{\partial_u \xi_k^i(\alpha)} - \log \frac{\partial_u \xi_{k+1}^i(\beta)}{\partial_u \xi_k^i(\beta)} \right|.$$

The last two inequalities with the choice of ε_1 , inequality (37), the choice of W and bound (33) imply:

$$\begin{aligned} \text{Dist}(\xi_j^i, \Lambda) &\leq \text{Dist}(\xi_{m'}^i, \Lambda) + \sum_{k=m'}^{j-1} \text{Dist}(f_0, f_0^k(W)) \\ &\quad + 2 \sum_{k=m'}^{j-1} \kappa M e^{-(k+1)\gamma_1 + \gamma_0 + 3} < 4 \cdot 10^{-3}. \end{aligned}$$

Similarly, by the choice of ε_1 , inequality (37), the bound for $|f'_\alpha(\xi_k^i(\alpha)) - f'_0(\xi_k^i(\alpha))|$ above, inequality (12), the choice of W and bound (33) imply

$$\Sigma(u, j, \Lambda) < \Sigma(u, m', \Lambda) + \sum_{k=m'}^{j-1} \left(\text{Dist}(f_0, f_0^k(W)) + 2\kappa e^{-k\gamma_1 + \gamma_0 + 1} \right) < 3 \cdot 10^{-3}.$$

Therefore $(*)_j$ holds for all $m_0 \leq j \leq m$. Thus by inequality (35), the definition of Λ and inequality (11)

$$\text{diam}(\xi_j^i(\Lambda)) < eM |(f_0^j)'(v_i)| \text{diam}(\Lambda) < e^{-1} \kappa^{-1}.$$

Conditions (T1-3) are satisfied by $\Gamma^u(m) := l_1^{-1} \cdot a^+(v_i, m; f)$, which completes the proof. \square

5.5. Proof of Proposition 5.9

Proof of Proposition 5.9. — We assume that we have a rational map $f = f_0$ that satisfies all the assumptions of Proposition 5.9 and such that

$$\theta := 1 - \frac{4\iota}{\gamma} \left(1 + 1.2 \frac{\log \kappa}{\sigma} \right) > 1 - \frac{1}{2d-2},$$

where σ is defined by (23).

We apply Lemma 5.15. Fix $\rho > 0$ provided by Lemma 5.13, applied for each $i = 1, \dots, 2d-2$. The intersection G of sets G_1, \dots, G_{2d-2} of densities at least θ in $\mathbb{N} \cap (m, 2m]$ has positive density, thus the union of such sets G is infinite.

By inequalities (32), (25) and (35), for each $n \in G$ and $i = 1, \dots, 2d-2$ we compute, for $u \in E$:

$$\Gamma^i(n) = l_1^{-1} \cdot a^+(v_i, n; f) > \frac{l_1^{-1} \rho}{|(f^n)'(v_i)|} > \frac{l_1^{-1} \rho M}{e |\partial_u \xi_n^i(0)|},$$

where $M := 1 + \max\{|\tau_i(u)|, |\tau_i(u)|^{-1} : u \in E\}$. The bounds for $r_{n,i} := \Gamma^i(n)$ compared to $C := e^2 l_1 M \rho^{-1}$ follow from the bounds (35), (7) and (T1) in the definition of $\Gamma^i(n)$. This gives the lower bound for $r_{n,i} |(f^n)'(v_i)|$. On the other hand, (31) gives

$$r_{n,i} |(f^n)'(v_i)| < \frac{|(f^n)'(v_i)|}{|\partial_u \xi_n^i(0)|} \leq 2M$$

for all $n \in G$ large enough and the upper bound follows. This completes the proof. \square

Theorem 5.16. — *Let f be a rational map satisfying the conditions of Proposition 5.9. Then f satisfies the large scale condition. Furthermore, it is possible to choose the sequence $(\underline{n}_k)_k$ in Definition 1.2 of the form $\underline{n}_k = (n_k, \dots, n_k)$ with $n_k \rightarrow +\infty$.*

Proof. — Let $f = f_0$ be such a rational map, and let $(f_\lambda)_{\lambda \in O}$ be a local holomorphic parametrization of the moduli space, where O is some neighborhood of the origin in \mathbb{C}^{2d-2} . Recall that by the transversality property (T), there are unit vectors $(u_j)_{1 \leq j \leq 2d-2}$ in \mathbb{C}^{2d-2} such that $\tau_i(u_j) \neq 0$ if and only if $i = j$. Let $r_{n,i}$, $G \subset \mathbb{N}$ and $E \subset \mathbb{S}_{2d-2}$ be provided by Proposition 5.9. Recall that E is an open neighborhood of $\|\sum_{j=1}^{2d-2} u_j\|^{-1} \sum_{j=1}^{2d-2} u_j$ in the unit sphere \mathbb{S}_{2d-2} of \mathbb{C}^{2d-2} , such that for all $1 \leq j \leq 2d-2$ we have $\tau_j(u) \neq 0$. We now fix an affine chart on \mathbb{P}^1 for the rest of the proof.

For all $n \in G$, let

$$\phi_n(\lambda) := (\xi_n^i(r_{n,i}\lambda) - \xi_n^i(0))_{1 \leq i \leq 2d-2}$$

As $r_{n,i} \asymp |(f^n)'(v_i)|^{-1} \rightarrow 0$ when $n \rightarrow \infty$, for all i by transversality property (T), for all $n \in G$ large enough ϕ_n is defined in the neighborhood of a ball of fixed size in \mathbb{C}^{2d-2} . Up to rescaling, we may assume it is the unit ball $\mathbb{B} \subset \mathbb{C}^{2d-2}$. Observe that for all $1 \leq i \leq 2d-2$, we have

$$\frac{d}{dt} \Big|_{t=0} \phi_n^i(tu) = r_{n,i} D\xi_n^i(0) \cdot u$$

so that by the definitions of τ_i we have

$$\frac{d}{dt} \Big|_{t=0} \phi_n^i(tu) = r_{n,i} \tau_i(u) (f^n)'(v_i) + o(1), \text{ as } n \rightarrow \infty.$$

Using again Proposition 5.9 and inequality (29), we find that there is a constant $C' > 0$ depending only on f such that for all $n \in G$ and all $u \in E$,

$$\frac{1}{C'} \leq \left| \frac{d}{dt} \Big|_{t=0} \phi_n^i(tu) \right| \leq C'.$$

From the distortion control of Proposition 5.9 and equation (7), we therefore have that for all $u \in E$, for all $t \in \mathbb{D}$, and for all $1 \leq i \leq 2d - 2$,

$$|\phi_n^i(tu)| \leq \frac{11}{10}C'.$$

Now applying Sibony-Wong's theorem (Theorem 5.8) to ϕ_n on E , we see that there is a constant $0 < \tilde{\rho} < 1$ independent of $n \in G$ such that for all $z \in \mathbb{B}(0, \tilde{\rho})$,

$$\|\phi_n(z)\| \leq (2d - 2)\frac{11}{10}C'.$$

Therefore the sequence $(\phi_n)_{n \in G}$ is normal near the origin, hence up to extraction converges to some holomorphic map $\phi : \mathbb{B}(0, \tilde{\rho}) \rightarrow \mathbb{C}^{2d-2}$. By the transversality property (T) and the fact that $r_{n,i}|(f^n)'(v_i)| \geq \frac{1}{C}$, we have that $D\phi(0)$ is invertible. Therefore $\phi(\mathbb{B}(0, \tilde{\rho}))$ contains some ball $\mathbb{B}(0, 2r)$ for some $r > 0$, and so for all large enough $n \in G$, $\phi_n(\mathbb{B}(0, \tilde{\rho}))$ contains the ball $\mathbb{B}(0, r)$. Recalling the definition of ϕ_n , we claim that this implies that f satisfies the large scale condition for all $n \in G$ with

$$\Omega_n := \mathbb{B}\left(0, \max_{1 \leq i \leq 2d-2} r_{n,i}\right).$$

Indeed, since $D\phi(0)$ is invertible, in particular, the graph of ϕ is vertical-like in a neighborhood of the origin, restricting r if necessary; by normality, this is also the case for ϕ_n for n large enough which is exactly what we want. \square

6. The proofs of Theorem A and Corollary 1

A map $f \in \text{Rat}_d$ is a *flexible Lattès map* if there exist an elliptic curve E , an affine map $\ell : E \rightarrow E$ with integral linear part and finite branched cover $\pi : E \rightarrow \mathbb{P}^1$ such that the following diagram commutes

$$\begin{array}{ccc} E & \xrightarrow{\ell} & E \\ \downarrow \pi & & \downarrow \pi \\ \mathbb{P}^1 & \xrightarrow{f} & \mathbb{P}^1 \end{array}$$

It follows from the definition that any flexible Lattès map is strongly Misiurewicz, that $\mathcal{P}(f) \cap \mathcal{C}(f) = \emptyset$ and that $\text{Card}(\mathcal{P}(f)) \leq 4$ and that its degree d is the square of the linear part of ℓ (see e.g. [Mi]). In particular, if f is a flexible Lattès map, then $d \geq 4$.

Proof of Theorem A. — Recall that a rational map is strongly Misiurewicz if all its critical points are preperiodic to repelling cycles. Let us denote by $\mathcal{X} \subset \mathcal{M}_d$ the set of all conjugacy classes of strongly Misiurewicz degree d rational maps and by $\mathcal{X}^* \subset \mathcal{X}$ those classes of rational maps with $\mathcal{P}(f) \cap \mathcal{C}(f) = \emptyset$ with simple critical values. Choose $[f] \in \mathcal{X}^*$. If $d \leq 3$, then f can not be a flexible Lattès map. If $d = \deg(f) \geq 4$, since f has simple critical values, $\text{Card}(\mathcal{P}(f)) \geq \text{Card}(f(\mathcal{C}(f))) = 2d - 2 > 4$ hence \mathcal{X}^* does not contain flexible Lattès maps. Observe that \mathcal{X}^* is countable.

Pick any $[f] \in \text{supp}(\mu_{\text{bif}})$ and $\Omega \subset \mathcal{M}_d$ an open neighborhood of $[f]$. According to [BE, Main Theorem], the set \mathcal{X}^* is dense in $\text{supp}(\mu_{\text{bif}})$, hence in particular $\mathcal{X}^* \cap \Omega \neq \emptyset$. Choose now such $[f_0] \in \mathcal{X}^* \cap \Omega$. By the above argument, we apply Theorem C: there exist $\mu(f_0), \mu_0(f_0), \gamma(f_0), \gamma_0(f_0) > 0$ and $\hat{\alpha}(f_0) > 0$ such that for all $\alpha < \min(\frac{\gamma(f_0)}{200}, \hat{\alpha}(f_0))$, there exist $\hat{\eta}(f_0) > 0$ and $\hat{i}(f_0) > 0$ such that for all $\eta < \hat{\eta}(f_0)$ and for all $\iota < \hat{i}(f_0)$, the map f_0 is

a Lebesgue point of density of rational maps satisfying $CE(\gamma(f_0), \gamma_0(f_0))$, $CE2(\mu(f_0), \mu_0(f_0))$, $BA(\alpha)$ and $FA(\eta, \iota)$.

We let $CE_{[f_0]}$ be those conjugacy classes $[g] \in \Omega$ of maps g satisfying the properties $CE(\gamma(f_0), \gamma_0(f_0))$, $CE2(\mu(f_0), \mu_0(f_0))$, $BA(\alpha)$ and $FA(\eta, \iota)$ with $\eta < \hat{\eta}(f_0)$ and $\iota < \hat{\iota}(f_0)$. Up to reducing $\hat{\iota}(f_0)$, we may assume that such $[g]$ satisfies the Large scale condition by Theorem D. The set $CE_{[f_0]}$ is contained in $\Omega \cap \text{supp}(\mu_{\text{bif}})$ by Theorem B and has positive Lebesgue volume by Theorem C. This concludes the proof of Theorem A. \square

Combining Theorem D with Theorem B, we also get the following result. In the light of the proof above, it is an enhanced version of Corollary 1.

Theorem 6.1. — *Let $\gamma, \gamma_0, \mu, \mu_0 > 0$, let $\alpha < \gamma/200$ and let $\eta > 0$. Let $\iota > 0$ be given by Theorem D. Assume that $f \in \text{Rat}_d$ has simple critical points and satisfies $CE(\gamma, \gamma_0)$, $CE2(\mu, \mu_0)$, $BA(\alpha)$ and $FA(\eta, \iota)$, then $[f] \in \text{supp}(\mu_{\text{bif}})$. In particular,*

1. *f is approximated by strongly Misiurewicz rational maps;*
2. *for any given $\theta_1, \dots, \theta_{2d-2} \in \mathbb{R} \setminus 2\pi\mathbb{Z}$, f is approximated by rational maps having $2d - 2$ distinct neutral cycles with respective multipliers $e^{i\theta_1}, \dots, e^{i\theta_{2d-2}}$;*
3. *for any given $w_1, \dots, w_{2d-2} \in \mathbb{D}$, f is approximated by hyperbolic rational maps with $2d - 2$ distinct attracting cycles with respective multipliers w_1, \dots, w_{2d-2} .*

Proof. — Combining Theorem D with Theorem B, we see that for any such Collet-Eckmann rational map f , the class $[f]$ belongs to $\text{supp}(\mu_{\text{bif}})$, or equivalently that $f \in \text{supp}(T_{\text{bif}}^{2d-2})$. The three points respectively follow from [BE, Theorem 1], from [G2, Theorem 1.1] and from Theorem 134 of [B]. \square

The support of the measure μ_{bif} has been proven to be the closure of classes of rational maps having a maximal number of neutral cycles with given multipliers. The following is a direct consequence of Theorem A and [G2, Theorem 1.1].

Corollary 6.2. — *Fix any $\theta_1, \dots, \theta_{2d-2} \in \mathbb{R} \setminus 2\pi\mathbb{Z}$, and let $\mathcal{N}(\theta_1, \dots, \theta_{2d-2})$ be the set of degree d rational maps having $(2d - 2)$ distinct neutral cycles of respective multipliers $\exp(i\theta_1), \dots, \exp(i\theta_{2d-2})$. Then*

$$\text{Vol}_{\mathcal{M}_d} \left(\overline{\{[f] \in \mathcal{M}_d; f \in \mathcal{N}(\theta_1, \dots, \theta_{2d-2})\}} \right) > 0.$$

It is actually difficult to exhibit one example of such a rational map. Here, we prove that they are dense in a set which is *not* negligible.

References

- [Asp1] Magnus Aspenberg. Rational Misiurewicz maps are rare. *Comm. Math. Phys.*, 291(3):645–658, 2009.
- [Asp2] Magnus Aspenberg. The Collet-Eckmann condition for rational functions on the Riemann sphere. *Math. Z.*, 273(3-4):935–980, 2013.
- [Ast] Matthieu Astorg. Summability Condition and Rigidity for Finite Type Maps. *ArXiv e-prints*, February 2016.
- [B] Francois Berteloot. Bifurcation currents in holomorphic families of rational maps. In *Pluripotential theory*, volume 2075 of *Lecture Notes in Math.*, pages 1–93. Springer, Heidelberg, 2013.
- [BB] Giovanni Bassanelli and Francois Berteloot. Bifurcation currents in holomorphic dynamics on \mathbb{P}^k . *J. Reine Angew. Math.*, 608:201–235, 2007.

- [BC] Michael Benedicks and Lennart Carleson. On iterations of $1 - ax^2$ on $(-1, 1)$. *Ann. of Math. (2)*, 122(1):1–25, 1985.
- [BE] Xavier Buff and Adam L. Epstein. Bifurcation measure and postcritically finite rational maps. In *Complex dynamics : families and friends / edited by Dierk Schleicher*, pages 491–512. A K Peters, Ltd., Wellesley, Massachusetts, 2009.
- [BM] Francois Berteloot and Volker Mayer. *Rudiments de dynamique holomorphe*, volume 7 of *Cours Spécialisés*. Société Mathématique de France, Paris, 2001.
- [De] Laura DeMarco. Dynamics of rational maps: a current on the bifurcation locus. *Math. Res. Lett.*, 8(1-2):57–66, 2001.
- [Du] Romain Dujardin. The supports of higher bifurcation currents. *Ann. Fac. Sci. Toulouse Math. (6)*, 22(3):445–464, 2013.
- [DF] Romain Dujardin and Charles Favre. Distribution of rational maps with a preperiodic critical point. *Amer. J. Math.*, 130(4):979–1032, 2008.
- [G1] Thomas Gauthier. Strong bifurcation loci of full Hausdorff dimension. *Ann. Sci. Éc. Norm. Sup. (4)*, 45(6):947–984, 2012.
- [G2] Thomas Gauthier. Higher bifurcation currents, neutral cycles, and the Mandelbrot set. *Indiana Univ. Math. J.*, 63(4):917–937, 2014.
- [GS] Jacek Graczyk and Stanislas Smirnov. Collet, Eckmann and Hölder. *Invent. Math.*, 133(1):69–96, 1998.
- [J] Mattias Jonsson. Holomorphic motions of hyperbolic sets. *Michigan Math. J.*, 45(2):409–415, 1998.
- [Le] Genadi Levin. Perturbations of weakly expanding critical orbits. *Frontiers in Complex Dynamics: In Celebration of John Milnor's 80th Birthday*, pages 163–196, 2014.
- [Ly] Mikhail Yu. Lyubich. Investigation of the stability of the dynamics of rational functions. *Teor. Funktsii Funktsional. Anal. i Prilozhen.*, (42):72–91, 1984. Translated in *Selecta Math. Soviet.* **9** (1990), no. 1, 69–90.
- [Ma] Ricardo Mañé. On the uniqueness of the maximizing measure for rational maps. *Bol. Soc. Brasil. Mat.*, 14(1):27–43, 1983.
- [Mc] Curtis T. McMullen. *Complex dynamics and renormalization*, volume 135 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1994.
- [Mi] John Milnor. On Lattès maps. In *Dynamics on the Riemann sphere*, pages 9–43. Eur. Math. Soc., Zürich, 2006.
- [MSS] Ricardo Mañé, Paulo Sad, and Dennis Sullivan. On the dynamics of rational maps. *Ann. Sci. École Norm. Sup. (4)*, 16(2):193–217, 1983.
- [P] Feliks Przytycki. Conical limit set and Poincaré exponent for iterations of rational functions. *Trans. Amer. Math. Soc.*, 351(5):2081–2099, 1999.
- [PRLS] Feliks Przytycki, Juan Rivera-Letelier, and Stanislav Smirnov. Equivalence and topological invariance of conditions for non-uniform hyperbolicity in the iteration of rational maps. *Invent. Math.*, 151(1):29–63, 2003.
- [R] Mary Rees. Positive measure sets of ergodic rational maps. *Ann. Sci. École Norm. Sup. (4)*, 19(3):383–407, 1986.
- [S] Mitsuhiro Shishikura. The Hausdorff dimension of the boundary of the Mandelbrot set and Julia sets. *Ann. of Math. (2)*, 147(2):225–267, 1998.
- [SW] Nessim Sibony and Pit Mann Wong. Some results on global analytic sets. In *Séminaire Pierre Lelong-Henri Skoda (Analyse). Années 1978/79 (French)*, volume 822 of *Lecture Notes in Math.*, pages 221–237. Springer, Berlin, 1980.
- [Ta] Lei Tan. Similarity between the Mandelbrot set and Julia sets. *Comm. Math. Phys.*, 134(3):587–617, 1990.
- [Ts] Masato Tsujii. Positive Lyapunov exponents in families of one-dimensional dynamical systems. *Invent. Math.*, 111(1):113–137, 1993.

MATTHIEU ASTORG, MAPMO, Université d'Orléans, Route de Chartres B.P. 6759 - 45067 ORLÉANS Cedex 2,
FRANCE • *E-mail* : `matthieu.astorg@univ-orleans.fr`

THOMAS GAUTHIER, LAMFA, UPJV, 33 rue Saint-Leu, 80039 AMIENS Cedex 1, FRANCE
CMLS, École polytechnique, CNRS, Université Paris-Saclay, 91128 PALAISEAU Cedex, FRANCE
E-mail : `thomas.gauthier@u-picardie.fr`

NICOLAE MIHALACHE, LAMA, UPEC, 61 avenue du Général de Gaulle, 94010 CRÉTEIL Cedex, FRANCE
E-mail : `nicolae.mihalache@u-pec.fr`

GABRIEL VIGNY, LAMFA, UPJV, 33 rue Saint-Leu, 80039 AMIENS Cedex 1, FRANCE
E-mail : `gabriel.vigny@u-picardie.fr`