



HAL
open science

A computable analysis of variable words theorems

Lu Liu, Benoit Monin, Ludovic Patey

► **To cite this version:**

Lu Liu, Benoit Monin, Ludovic Patey. A computable analysis of variable words theorems. Proceedings of the American Mathematical Society, 2019, 147 (2), pp.823–834. 10.1090/proc/14269 . hal-01888789

HAL Id: hal-01888789

<https://hal.science/hal-01888789v1>

Submitted on 9 Oct 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A COMPUTABLE ANALYSIS OF VARIABLE WORDS THEOREMS

LU LIU, BENOIT MONIN, AND LUDOVIC PATEY

ABSTRACT. The Carlson-Simpson lemma is a combinatorial statement occurring in the proof of the Dual Ramsey theorem. Formulated in terms of variable words, it informally asserts that given any finite coloring of the strings, there is an infinite sequence with infinitely many variables such that for every valuation, some specific set of initial segments is homogeneous. Friedman, Simpson, and Montalban asked about its reverse mathematical strength. We study the computability-theoretic properties and the reverse mathematics of this statement, and relate it to the finite union theorem. In particular, we prove the Ordered Variable word for binary strings in ACA_0 .

1. INTRODUCTION

Let $(\mathbb{N})^k$ and $(\mathbb{N})^\infty$ denote the set of partitions of \mathbb{N} into exactly k and infinitely many non-empty pieces, respectively. For $X \in (\mathbb{N})^\infty$, $(X)^k$ is the set of all $Y \in (\mathbb{N})^k$ which are coarser than X .

Statement 1.1 (Dual Ramsey theorem). DRT^k is the statement “If $(\mathbb{N})^k$ is colored with finitely many Borel colors, then there is some $X \in (\mathbb{N})^\infty$ such that $(X)^k$ is monochromatic”.

The Dual Ramsey theorem was proven by Carlson and Simpson [1], and studied from a reverse mathematical viewpoint by Slaman [9], Miller and Solomon [5] and Dzhafarov et al. [3]. In this paper, we shall focus on a combinatorial lemma used by Carlson and Simpson to prove the Dual Ramsey theorem. This lemma can be formulated in terms of *variable words*.

Definition 1.2 (Variable word). An *infinite variable word* on a finite alphabet A is an ω -sequence W of elements of $A \cup \{x_i : i \in \mathbb{N}\}$ in which all variables occur at least once, and finitely often. Moreover, the first occurrence of x_i comes before the first occurrence of x_{i+1} . A *finite variable word* is an initial segment of an infinite variable word. A finite or infinite variable word is *ordered* if moreover all occurrences of x_i come before any occurrence of x_{i+1} . Given $\bar{a} = a_0 a_1 \dots a_{k-1} \in A^{<\omega}$, we let $W(\bar{a})$ denote the finite A -string obtained by replacing x_i with a_i in W and then truncating the result just before the first occurrence of x_k .

Statement 1.3 (Variable word theorem). $\text{VW}(n, r)$ is the statement “If $A^{<\omega}$ is colored with r colors for some alphabet A of cardinality n , there exists an infinite variable word W such that $\{W(\bar{a}) : \bar{a} \in A^{<\omega}\}$ is monochromatic. $\text{OVW}(n, r)$ is the same statement as $\text{VW}(n, r)$ but for ordered variable words.

In this paper, we study the computability-theoretic properties of the variable word theorems using the framework of reverse mathematics.¹

1.1. Reverse mathematics. Reverse mathematics is a vast foundational program aiming to determine the optimal axioms to prove ordinary theorems. It uses the framework of second-order arithmetic, with a base theory RCA_0 consisting of the axioms of Robinson arithmetic, the Σ_1^0 induction scheme and the Δ_1^0 comprehension scheme. The system RCA_0 arguably captures *computable mathematics*. Starting from a proof-theoretic perspective, modern reverse mathematics tends to be seen as a framework to analyse the computability-theoretic features of theorems. Among the distinguished statements, let us mention weak König’s lemma (WKL), asserting that every infinite binary tree has an infinite path, the arithmetic comprehension axiom (ACA), and the Π_1^1 comprehension axiom ($\Pi_1^1\text{CA}$), consisting of the comprehension scheme restricted to arithmetic and Π_1^1 formulas, respectively. See Simpson [8] for reference book on classical reverse mathematics.

¹The authors thank Damir Dzhafarov, Stephen Flood, Reed Solomon and Linda Brown Westrick for bringing the attention of the authors to the Carlson-Simpson lemma, and for numerous discussions. The authors are also thankful to Denis Hirschfeldt and Barbara Csima for showing them how to use Lovasz Local Lemma to prove lower bounds to combinatorial theorems.

The statements studied within this framework are mainly of the form $(\forall X)[\Phi(X) \rightarrow (\exists Y)\Psi(X, Y)]$, where Φ and Ψ are arithmetic formulas with set parameters, and can be considered as *problems*. Given a statement P of this form, a set X such that $\Phi(X)$ holds is an *instance* of P , and a set Y such that $\Psi(X, Y)$ holds is a *solution* to the P -instance X . In this paper, we shall consider exclusively statements of this kind.

Friedman and Simpson [4], and later Montalban [6], asked about the reverse mathematical strength of the ordered variable word. The statement $\text{OVW}(k, \ell)$ is known to be provable in $\text{RCA}_0 + \Pi_1^1\text{CA}$. Our main result is a direct combinatorial proof of $\text{OVW}(2, \ell)$ in $\text{RCA}_0 + \text{ACA}$.

Theorem 1.4. *For every $\ell \geq 2$, $\text{RCA}_0 + \text{ACA} \vdash \text{OVW}(2, \ell)$.*

On the lower bound hand, Miller and Solomon [5] constructed a computable instance c of $\text{OVW}(2, 2)$ with no Δ_2^0 solution, and deduced that $\text{RCA}_0 + \text{WKL}$ does not prove $\text{VW}(2, 2)$. Indeed, seeing the instance c of $\text{OVW}(2, 2)$ as an instance of $\text{VW}(2, 2)$, and noticing that the jump of a solution to $\text{VW}(2, 2)$ gives a solution to $\text{OVW}(2, 2)$, one can deduce that c has no low $\text{VW}(2, 2)$ -solution. In this paper, we improve their lower bound by constructing a computable instance of $\text{OVW}(2, 2)$ whose solutions are of DNC degree relative to \emptyset' .

1.2. Organization of the paper. In section 2, we shall give a simple proof of the ordered variable word for binary strings ($\text{OVW}(2, \ell)$) using the finite union theorem. Then, in section 3, we provide a direct combinatorial proof of the same statement over $\text{RCA}_0 + \text{ACA}$. Finally, in section 4, we give a new lower bound on the strength of $\text{OVW}(2, \ell)$ using a computable version of Lovasz Local Lemma.

1.3. Notation. Given two sets A and B , we write $A < B$ for the formula $(\forall x \in A)(\forall y \in B)x < y$. Given a set A , we write $A^{<\omega}$ for the set of finite A -valued strings. In particular, $2^{<\omega}$ is the set of binary strings. We denote by $\mathcal{P}_{fin}(\mathbb{N})$ the collection of finite *non-empty* subsets of \mathbb{N} . Given two strings $\sigma, \tau \in A^{<\omega}$, $\sigma * \tau$ denotes their concatenation. We may also write $\sigma\tau$ when there is no ambiguity. Given a string or a sequence X and some $n \in \omega$, we write $X \upharpoonright n$ for the initial segment of X of length n . In particular, $X \upharpoonright 0$ is the empty string, written ε .

2. A SIMPLE PROOF OF THE ORDERED VARIABLE WORD THEOREM FROM THE FINITE UNION THEOREM

Simpson first noted a relation between Hindman's theorem and the Carlson-Simpson lemma [1]. In this section, we give a formal counterpart to his observation by giving a simple proof of $\text{OVW}(2, \ell)$ using the Finite Union Theorem, a statement known to be equivalent to Hindman's theorem. A variation of the proof below was used by Dzhafarov et al. [3] to give an upper bound to the Open Dual Ramsey's theorem. A direct combinatorial proof of $\text{OVW}(2, \ell)$ in $\text{RCA}_0 + \text{ACA}$ will be given in the next section.

Definition 2.1. An *IP collection* is an infinite collection of finite sets $\mathcal{I} \subseteq \mathcal{P}_{fin}(\mathbb{N})$ which is closed under *non-empty* finite unions and contains an infinite subcollection of pairwise disjoint sets.

Note that any IP collection \mathcal{I} necessarily contains an infinite \mathcal{I} -computable sequence $S_0 < S_1 < \dots$.

Statement 2.2 (Finite union theorem). For every $\ell \in \mathbb{N}$, FUT_ℓ is the statement “For every coloring $c : \mathcal{P}_{fin}(\mathbb{N}) \rightarrow \ell$, there is a monochromatic IP collection”. wFUT_ℓ^2 is the statement “For every coloring $c : \mathcal{P}_{fin}(\mathbb{N}) \times \mathbb{N} \rightarrow \ell$, there is an IP collection \mathcal{I} and a color $i < \ell$ such that $c(S, \min T) = i$ for every $S < T \in \mathcal{I}$.”

Theorem 2.3. $\text{RCA}_0 \vdash \forall \ell (\text{FUT}_\ell \rightarrow \text{wFUT}_\ell^2)$.

Proof. Assume $\ell \geq 2$, the other cases being trivial. Let $f : \mathcal{P}_{fin}(\mathbb{N}) \times \mathbb{N} \rightarrow \ell$ be an instance of wFUT_ℓ^2 . Note that over RCA_0 , $\text{FUT}_\ell \rightarrow \text{ACA}$ and $\text{ACA} \rightarrow \text{COH}$. Let \vec{R} be a sequence of set defined for every $S \in \mathcal{P}_{fin}(\mathbb{N})$ and $i < \ell$ by $R_{S,i} = \{n \in \mathbb{N} : f(S, n) = i\}$. Apply COH to \vec{R} to obtain an infinite \vec{R} -cohesive set C . In particular, for every $S \in \mathcal{P}_{fin}(\mathbb{N})$, $\lim_{n \in C} f(S, n)$ exists.

Let $h : \omega \rightarrow C$ be a computable bijection. Let $\tilde{f} : \mathcal{P}_{fin}(\mathbb{N}) \rightarrow \ell$ be defined by $\tilde{f}(S) = \lim_{n \in C} f(h[S], n)$. \tilde{f} is a $\Delta_2^{0, f \oplus C}$ instance of FUT_ℓ , so by the finite union theorem, there is an IP collection $\mathcal{I} \subseteq \mathcal{P}_{fin}(\mathbb{N})$. and a color $i < \ell$ such that for every $S \in \mathcal{I}$, $\tilde{f}(S) = \lim_{n \in C} f(h[S], n) = i$. Note that for every $S \in \mathcal{I}$, $h[S] \in C$. Therefore, by f -computably thinning-out the set \mathcal{I} , we obtain an IP collection $\mathcal{J} \subseteq \mathcal{I}$ such that for every $S < T \in \mathcal{J}$, $f(h[S], \min h[T]) = i$. The set $\{h[S] : S \in \mathcal{J}\}$ is a solution to f . \square

Theorem 2.4. $\text{RCA}_0 \vdash \forall \ell (\text{wFUT}_\ell^2 \rightarrow \text{OVW}(2, \ell))$.

Proof. Let $f : 2^{<\omega} \rightarrow \ell$ be an instance of $\text{OVW}(2, \ell)$. Define an instance $g : \mathcal{P}_{fin}(\mathbb{N}) \times \mathbb{N} \rightarrow \ell$ of wFUT_ℓ^2 as follows: Given some $S \in \mathcal{P}_{fin}(\mathbb{N})$ and $n \in \mathbb{N}$, if $\max S < n$, then set $g(S, n) = f(\sigma)$, where σ is the binary string of length n defined by $\sigma(i) = 1$ iff $i \in S$. If $n \leq \max S$, set $g(S, n) = 0$. By wFUT_ℓ^2 , there is an IP collection \mathcal{I} and a color $i < \ell$ such that $g(S, \min T) = i$ for every $S < T \in \mathcal{I}$. Compute from \mathcal{I} an infinite increasing sequence of pairwise disjoint finite sets $F_0 < F_1 < \dots$. Let W be the infinite variable word defined by

$$W(n) = \begin{cases} 1 & \text{if } n \in F_0 \\ x_i & \text{if } n \in F_i \text{ for some } i \geq 1 \\ 0 & \text{otherwise} \end{cases}$$

The variable word W and the sequence of the F 's is a solution to the instance f of $\text{OVW}(2, \ell)$. \square

Corollary 2.5. $\text{RCA}_0 \vdash \text{ACA}^+ \rightarrow \forall \ell \text{OVW}(2, \ell)$.

Proof. Immediate since $\text{ACA}^+ \rightarrow \forall \ell \text{FUT}_\ell \rightarrow \forall \ell \text{wFUT}_\ell^2 \rightarrow \forall \ell \text{OVW}(2, \ell)$ over RCA_0 . \square

3. A PROOF OF THE ORDERED VARIABLE WORD THEOREM IN ACA

The proof of the previous section gave a very coarse computability-theoretic upper bound of the Ordered Variable Word theorem in terms of ω -jumps. In this section, we give a direct combinatorial proof of $\text{OVW}(2, \ell)$ in $\text{RCA}_0 + \text{ACA}$. Actually, every PA degree relative to \emptyset' is sufficient to compute a solution of a computable instance of $\text{OVW}(2, \ell)$. We thereby answer a question of Miller and Solomon [5].

Theorem 3.1. *For every $\ell \in \omega$, every computable instance c of $\text{OVW}(2, \ell)$, every PA degree over \emptyset' computes a solution to c .*

A formalization of Theorem 3.1 yields a proof of Theorem 1.4.

Proof of Theorem 1.4. The proof of Theorem 3.1 can be formalized within $\text{RCA}_0 + \text{ACA}$. Indeed, the arguments require only arithmetical induction to be carried out, and every model of $\text{RCA}_0 + \text{ACA}$ is a model of the statement “For every set X , there is a set of PA degree over the jump of X .” \square

Let us first introduce some notation. For a finite set F and a string $\sigma \in 2^{<\omega}$ let σ_F be the binary string of length $|\sigma|$ defined by $\sigma_F(i) = \sigma(i)$ if $i \notin F$, and $\sigma_F(i) = 1 - \sigma(i)$ otherwise. Let \leq_{lex} denote the shortlex order on $\omega^{<\omega}$, that is, the order with the shortest length first, and with the strings of same length sorted lexicographically.

In what follows, fix a coloring $\tilde{c} : 2^{<\omega} \rightarrow \ell$, and a string $\rho \in 2^{<\omega}$.

The main combinatorial lemma we use is Lemma 3.4. As a warm up, we first prove the following lemma 3.2, which is a consequence of Lemma 3.4 and the proof is somehow similar but much simpler. In the following lemma, one may think of $\rho_{P'}$ as a finite variable word, where the positions at \tilde{P} are replaced by a same variable kind.

Lemma 3.2. *For any $P \subseteq \{0, \dots, |\rho| - 1\}$ with $(\forall n \in P)[\rho(n) = 0] \wedge |P| \geq \ell$, there exist two subsets $P' < \tilde{P}$ of P with $\tilde{P} \neq \emptyset$ such that $\tilde{c}(\alpha) = \tilde{c}(\alpha_{\tilde{P}})$ where $\alpha = \rho_{P'}$.*

Proof. Suppose $P = \{p_0 < \dots < p_{m-1}\}$. Let ℓ_0, \dots, ℓ_m be defined by $\ell_i = \tilde{c}(\rho_{\{p_0, \dots, p_{i-1}\}})$. In particular, $\ell_0 = \tilde{c}(\rho)$. Since $|P| = m \geq \ell$, so among ℓ_0, \dots, ℓ_m , there must exist $i < j$ such that $\ell_i = \ell_j$. Let $P' = \{p_0, \dots, p_{i-1}\}$ (if $i = 0$ then $P' = \emptyset$), and $\tilde{P} = \{p_i, \dots, p_{j-1}\}$, let $\alpha = \rho_{P'}$. Clearly $P' < \tilde{P}$ and $\tilde{P} \neq \emptyset$. It is also easy to see that $\tilde{c}(\alpha) = \ell_i = \ell_j = \tilde{c}(\alpha_{\tilde{P}})$. \square

We now prove a technical lemma used in the proof of our main combinatorial lemma (Lemma 3.4). The sequence in the following lemma is obtained by a simple greedy algorithm, with finitely many resets.

Lemma 3.3. *There exists a nonempty set of colors $L \subseteq \{0, 1, \dots, \ell - 1\}$, $|L| + 1$ many sets of binary strings $\Gamma_0 = \{\tau^\eta\}_{\eta \in L}, \Gamma_1 = \{\tau^\eta\}_{\eta \in L^2}, \dots, \Gamma_{|L|} = \{\tau^\eta\}_{\eta \in L^{|L|+1}}$, such that, letting*

$$\tilde{\eta} = \underbrace{\max L * \max L * \dots * \max L}_{|L|+1 \text{ many}}$$

and letting $\tilde{\rho} = \tau^{\tilde{\eta}} * 0$, the following holds:

- (1) $\rho \prec \Gamma_0$ and $\tau^\eta \prec \tau^\beta \Leftrightarrow \eta \prec_{lex} \beta$;
- (2) $\tilde{\rho}(|\tau|) = 0$ for all $\tau \in \Gamma_i, i \leq |L|$;
- (3) for all $i \leq |L|$, $\eta \in L^{i+1}$, let $\eta_0 \prec \eta_1 \prec \dots \prec \eta_{i-1}$ denote all nonempty predecessors of η , let $Q = \{|\tau^{\eta_0}|, |\tau^{\eta_1}|, \dots, |\tau^{\eta_{i-1}}|\}$ (if $i = 0$ then $Q = \emptyset$), then $\tilde{c}(\tau_Q^\eta) = \eta(i)$;
- (4) let $P = \{|\tau^\eta|\}_{\eta \in L \leq |L|+1}$, for all subset Q of P , all $\tau \succeq \tilde{\rho}$, $\tilde{c}(\tau_Q) \in L$.

Moreover, $\Gamma_i, i \leq |L|$ is computable in the jump of \tilde{c} , uniformly in ρ .

Proof. We firstly show how to find Γ_0 . Start with $L = \{0, 1, \dots, \ell - 1\}$. At step 1, try to find a string $\tau \in 2^{<\omega}$ such that $\tilde{c}(\rho\tau) = 0$ and let $\tau^0 = \rho\tau$. Then try to find a τ such that $\tilde{c}(\tau^0 0\tau) = 1$ and let $\tau^1 = \tau^0 0\tau$. Generally, after τ^j is found, try to find τ such that $\tilde{c}(\tau^j 0\tau) = j + 1$ and let $\tau^{j+1} = \tau^j 0\tau$ if τ is found. If during the above process, after τ^{j-1} is defined ($\tau^{-1} = \rho$), there is no τ such that $\tilde{c}(\tau^{j-1} 0\tau) = j$, then we start all over again with ρ replaced by $\rho_1 = \tau^j 0$ and with L replaced by $L \setminus \{j\}$.

Generally, given a set of colors L and after τ^β is found, let η be the immediate successor (with respect to \leq_{lex} order restricted to L -strings) of β , let $\eta_0 \prec \eta_1 \prec \dots \prec \eta_{i-1}$ denote all nonempty predecessors of η , let $Q = \{|\tau^{\eta_0}|, |\tau^{\eta_1}|, \dots, |\tau^{\eta_{i-1}}|\}$ (if $i = 0$ then $Q = \emptyset$), we try to find τ such that $\tilde{c}((\tau^\beta 0\tau)_Q) = \eta(|\eta| - 1)$. If such a string τ does not exist then we start all over again with ρ replaced by $\tau^\beta 0_Q$ and L replaced by $L \setminus \{\eta(|\eta| - 1)\}$. If such τ exists then let $\tau^\eta = \tau^\beta 0\tau$.

Note that we have to start over for at most $\ell - 1$ times before we ultimately succeed since there are ℓ colors in total. It is plain to check all the four items. Also note that the sequence $\Gamma_0, \dots, \Gamma_{|L|}$ is \mathcal{C}' -computable since we only need to use the jump of \tilde{c} to know whether the next τ^η can be found. \square

Lemma 3.4. *There exists a string $\tilde{\rho} \succ \rho$ and a finite set $P \subseteq \{|\rho|, \dots, |\tilde{\rho}| - 1\}$ with $(\forall i \in P)[\tilde{\rho}(i) = 0]$ such that for all $\sigma \succeq \tilde{\rho}$ there exists two subsets $P' < \tilde{P}$ of P with $\tilde{P} \neq \emptyset$ such that, letting $\alpha = \sigma_{P'}$, $\tilde{c}(\alpha) = \tilde{c}(\alpha_{\tilde{P}}) = \tilde{c}(\alpha \upharpoonright \min \tilde{P})$. Moreover, $|P| < \ell^{\ell+2}$, and $\tilde{\rho}, P$, are computable in the jump of \tilde{c} , uniformly in ρ .*

Proof. Let L and $\tilde{\rho}$ satisfy Lemma 3.3. We claim that $\tilde{\rho}$ and $P = \{|\tau^\eta|\}_{\eta \in L \leq |L|+1}$ satisfy the current lemma. It is clear by item 1 of Lemma 3.3 that $\tilde{\rho} \succ \rho$ and by item 2 of Lemma 3.3 that $(\forall i \in P)[\tilde{\rho}(i) = 0]$.

Fix an arbitrary $\sigma \succeq \tilde{\rho}$. We now describe how to construct P' and \tilde{P} . Define $\ell_0, \dots, \ell_{|L|}$ and $p_0, \dots, p_{|L|}$ inductively by $\ell_0 = \tilde{c}(\sigma)$, $\ell_{i+1} = \tilde{c}(\sigma_{\{p_0, p_1, \dots, p_i\}})$, and $p_i = |\tau^{\ell_0 \dots \ell_i}|$ (where $\tau^{\ell_0 \dots \ell_i} \in \Gamma_i$). Since $\ell_0, \dots, \ell_{|L|} \in L$ (by item 4 of Lemma 3.3), there is some $i < j \leq |L|$ such that $\ell_i = \ell_j$. Let $P' = \{p_0, \dots, p_{i-1}\}$ (if $i = 0$ then $P' = \emptyset$), $\tilde{P} = \{p_i, \dots, p_{j-1}\}$, and let $\alpha = \sigma_{P'}$. We claim that $\tilde{c}(\alpha) = \tilde{c}(\alpha_{\tilde{P}}) = \tilde{c}(\alpha \upharpoonright \min \tilde{P})$. Note that $\min \tilde{P} = p_i = |\tau^{\ell_0 \dots \ell_i}|$. Therefore $\alpha \upharpoonright \min \tilde{P} = \tau_{P'}^{\ell_0 \dots \ell_i}$. By item 3 of Lemma 3.3, we have $\tilde{c}(\tau_{P'}^{\ell_0 \dots \ell_i}) = \ell_i$. Meanwhile, by definition of ℓ_i , $\tilde{c}(\sigma_{P'}) = \tilde{c}(\alpha) = \ell_i$. By definition of ℓ_j , $\tilde{c}(\sigma_{P' \cup \tilde{P}}) = \tilde{c}(\alpha_{\tilde{P}}) = \ell_j$. Thus, $\tilde{c}(\alpha) = \tilde{c}(\alpha_{\tilde{P}}) = \tilde{c}(\alpha \upharpoonright \min \tilde{P})$. \square

We say that $(\tilde{\rho}, P)$ is \tilde{c} -valid if P and $\tilde{\rho}$ satisfy Lemma 3.4. We say that (P', \tilde{P}) witnesses \tilde{c} -validity of $(\tilde{\rho}, P)$ for $\sigma \succeq \tilde{\rho}$ if $P' < \tilde{P} \subseteq P$, and letting $\alpha = \sigma_{P'}$, $\tilde{c}(\alpha) = \tilde{c}(\alpha_{\tilde{P}}) = \tilde{c}(\alpha \upharpoonright \min \tilde{P})$. Before proving Theorem 3.1, we start with the following simpler version.

Theorem 3.5. *For every $\ell \in \omega$, every computable instance $c : 2^{<\omega} \rightarrow \ell$ of OVW(2, ℓ), every PA degree over \emptyset'' computes a solution to c .*

Proof. It suffices to compute, given a PA degree relative to \emptyset'' , an infinite binary sequence $Y \in 2^\omega$ together with a sequence of finite sets $\tilde{P}_0 < \tilde{P}_1 < \dots$ with $(\forall i \in \omega)(\forall n \in \tilde{P}_i)[Y(n) = 0]$ such that the following holds:

Let $Position = \{\min \tilde{P}_i : i \geq 1\}$. There is some $\tilde{\ell} < \ell$ such that for all subset J of ω , letting $\tilde{P}_J = \bigcup_{i \in J} \tilde{P}_i$, then we have, $(\forall p \in Position)[c(Y_{\tilde{P}_J} \upharpoonright p) = \tilde{\ell}]$.

Using Lemma 3.4, we first construct a \emptyset' -computable sequence of strings $\tilde{\rho}_0 \prec \tilde{\rho}_1 \prec \dots$, a sequence of finite sets $P_i \subseteq \{|\tilde{\rho}_{i-1}|, \dots, |\tilde{\rho}_i| - 1\}$ and a sequence of colorings $c_i : [\tilde{\rho}_i]^\preceq \rightarrow L_i$ inductively as follows. $\tilde{\rho}_0 = \varepsilon$ and $c_0 = c$. Given $\tilde{\rho}_i$ and $c_i : [\tilde{\rho}_i]^\preceq \rightarrow L_i$, let $\tilde{\rho}_{i+1} \succeq \tilde{\rho}_i$ and $P_i \subseteq \{|\tilde{\rho}_i|, \dots, |\tilde{\rho}_{i+1}| - 1\}$ be such that $(\tilde{\rho}_{i+1}, P_i)$ - is c_i -valid, and let c_{i+1} be the coloring of $[\tilde{\rho}_{i+1}]^\preceq$ which on $\sigma \succeq \tilde{\rho}_{i+1}$ associates $\langle P', \tilde{P}, j \rangle$ such that (P', \tilde{P}) witnesses c_i -validity of $(\tilde{\rho}_{i+1}, P_i)$ for σ , and $c_i(\sigma_{P'}) = j$. If there are multiple such tuples, take the least one, in some arbitrary order. Note that the range of c_i is some finite set L_i .

We now analyze for $\sigma \succeq \tilde{\rho}_i$ what $c_i(\sigma) = \langle P', \tilde{P}, j \rangle$ means. Note that elements of $L_i, i \in \omega$ admit a natural partial order \triangleleft as follows: for $\langle P'_0, \tilde{P}_0, j_0 \rangle \in L_i, \langle P'_1, \tilde{P}_1, j_1 \rangle \in L_{i+1}, \langle P'_1, \tilde{P}_1, j_1 \rangle$ is an immediate successor of $\langle P'_0, \tilde{P}_0, j_0 \rangle$ if and only if $j_1 = \langle P'_0, \tilde{P}_0, j_0 \rangle$. Clearly every $j \in L_i$ admit a unique immediate predecessor.

Claim 3.6. *Fix some $n \geq 1$, and let $\tilde{\ell} \triangleleft \langle P'_0, \tilde{P}_0, j_0 \rangle \triangleleft \dots \triangleleft \langle P'_{n-1}, \tilde{P}_{n-1}, j_{n-1} \rangle = c_n(\sigma)$, Let $P' = \bigcup_{i \leq n-1} P'_i$ and $\alpha = \sigma_{P'}$. Then for any subset J of $\{0, \dots, n-1\}$,*

$$(\forall p \in \{ \min \tilde{P}_j : 1 \leq j \leq n-1 \} \cup \{|\alpha|\}) [c(\alpha_{\tilde{P}_J} \upharpoonright p) = \tilde{\ell}].$$

Proof. First we prove the claim for $n = 1$. By definition of $c_1(\sigma) = \langle P'_0, \tilde{P}_0, j_0 \rangle$, letting $\beta = \sigma_{P'_0}, c_0(\beta) = c_0(\beta_{\tilde{P}_0}) = j_0 = \tilde{\ell}$. In other words, for any subset $J \subseteq \{0\}$,

$$(\forall p \in \{ \min \{ \tilde{P}_j : 1 \leq j \leq 0 \} \} \cup \{|\beta|\}) [c(\beta_{\tilde{P}_J} \upharpoonright p) = \tilde{\ell}].$$

So the claim holds for $n = 1$. Suppose now the claim holds for $n-1$.

Suppose $c_n(\sigma) = \langle P'_{n-1}, \tilde{P}_{n-1}, j_{n-1} \rangle$. Let $\beta = \sigma_{P'_{n-1}}$. We have $c_{n-1}(\beta) = c_{n-1}(\beta_{\tilde{P}_{n-1}}) = c_{n-1}(\beta \upharpoonright \min \tilde{P}_{n-1}) = j_{n-1} = \langle P'_{n-2}, \tilde{P}_{n-2}, j_{n-2} \rangle$. As $c_{n-1}(\beta) = \langle P'_{n-2}, \tilde{P}_{n-2}, j_{n-2} \rangle$ and as $\tilde{\ell} \triangleleft \langle P'_{n-2}, \tilde{P}_{n-2}, j_{n-2} \rangle$, by induction hypothesis, for any subset J of $\{0, \dots, n-2\}$ we have:

$$(3.1) \quad c(\beta_{(\bigcup_{i \leq n-2} P'_i) \cup \tilde{P}_J}) = \tilde{\ell}.$$

Let $\beta' = \beta_{\tilde{P}_{n-1}}$. As $c_{n-1}(\beta') = \langle P'_{n-2}, \tilde{P}_{n-2}, j_{n-2} \rangle$ and as $\tilde{\ell} \triangleleft \langle P'_{n-2}, \tilde{P}_{n-2}, j_{n-2} \rangle$, by induction hypothesis, for any subset J of $\{0, \dots, n-2\}$ we have:

$$(3.2) \quad c(\beta'_{(\bigcup_{i \leq n-2} P'_i) \cup \tilde{P}_J}) = \tilde{\ell}.$$

As $c_{n-1}(\beta \upharpoonright \min \tilde{P}_{n-1}) = \langle P'_{n-2}, \tilde{P}_{n-2}, j_{n-2} \rangle$ and as $\tilde{\ell} \triangleleft \langle P'_{n-2}, \tilde{P}_{n-2}, j_{n-2} \rangle$, by induction hypothesis, for any subset J of $\{0, \dots, n-2\}$ we have:

$$(3.3) \quad (\forall p \in \{ \min \tilde{P}_j : 1 \leq j \leq n-2 \} \cup \{|\beta \upharpoonright \min \tilde{P}_{n-1}|\}) [c(\beta_{(\bigcup_{i \leq n-2} P'_i) \cup \tilde{P}_J} \upharpoonright p) = \tilde{\ell}].$$

But $|\beta \upharpoonright \min \tilde{P}_{n-1}| = \min \tilde{P}_{n-1}$. So (3.3) means for any subset J of $\{0, \dots, n-2\}$ we have:

$$(\forall p \in \{ \min \tilde{P}_j : 1 \leq j \leq n-1 \}) [c(\beta_{(\bigcup_{i \leq n-2} P'_i) \cup \tilde{P}_J} \upharpoonright p) = \tilde{\ell}].$$

Or equivalently, for any subset J of $\{0, \dots, n-1\}$ we have:

$$(3.4) \quad (\forall p \in \{ \min \tilde{P}_j : 1 \leq j \leq n-1 \}) [c(\beta_{(\bigcup_{i \leq n-2} P'_i) \cup \tilde{P}_J} \upharpoonright p) = \tilde{\ell}].$$

Now from 3.1, 3.2 and 3.4 we deduce that for any subset J of $\{0, \dots, n-1\}$ we have:

$$(\forall p \in \{ \min \tilde{P}_j : 1 \leq j \leq n-1 \} \cup \{|\beta|\}) [c(\beta_{(\bigcup_{i \leq n-2} P'_i) \cup \tilde{P}_J} \upharpoonright p) = \tilde{\ell}]$$

which completes the proof of the claim since $\beta_{\bigcup_{i \leq n-2} P'_i} = \alpha$. □

Let \mathcal{T}_0 be the \emptyset' -computable set of all γ such that $(\forall i \leq |\gamma|)[\gamma(i) \in L_i, \gamma(i) \triangleleft \gamma(i+1)$ and $\gamma(|\gamma|-1) = c_{|\gamma|-1}(\tilde{\rho}_{|\gamma|})$. Then, let \mathcal{T} be the downward closure of the set \mathcal{T}_0 by the prefix relation. The tree \mathcal{T} is infinite by construction of the strings $\tilde{\rho}_i$, the colors c_i and the sets P_i : a witness for the c_i -validity of $(\tilde{\rho}_{i+1}, P_{i+1})$ for ρ_{i+1} yields a node of \mathcal{T}_0 of length $i+2$. The tree \mathcal{T} is also \emptyset' -computably bounded, and \emptyset'' -computable. Let $j_0 * \langle P'_0, \tilde{P}_0, j_0 \rangle * \langle P'_1, \tilde{P}_1, j_1 \rangle * \dots$ be an infinite path through \mathcal{T} computed by any PA degree over \emptyset'' . By construction, $\langle P'_i, \tilde{P}_i, j_i \rangle \triangleleft \langle P'_{i+1}, \tilde{P}_{i+1}, j_{i+1} \rangle$. Let $X = \bigcup_{i \in \omega} \tilde{\rho}_i, P' = \bigcup_{i \in \omega} P'_i$ and let $Y = X_{P'}$. Clearly $(\forall i \forall n \in \tilde{P}_i)[Y(n) = 0]$ and Y is computable in the given PA degree relative to \emptyset'' . Therefore, letting $Position = \{ \min \tilde{P}_i : i \geq 1 \}$, it suffices to show that for all subsets J of ω ,

$$(\forall p \in Position) [c(Y_{\tilde{P}_J} \upharpoonright p) = j_0].$$

Without loss of generality, suppose $p = \min \tilde{P}_n$ and $J \subseteq \{0, \dots, n-1\}$. Since $j_0 * \langle P'_0, \tilde{P}_0, j_0 \rangle * \langle P'_1, \tilde{P}_1, j_1 \rangle * \dots * \langle P'_n, \tilde{P}_n, j_n \rangle$ is an initial segment of some element in \mathcal{T}_0 , there must exist some $N > n$ such that $c_N(\tilde{\rho}_{N+1}) = \langle P'_{N-1}, \tilde{P}_{N-1}, j_{N-1} \rangle$. Let $\sigma = \tilde{\rho}_{N+1}, \alpha = \sigma_{P'}$. Clearly $\alpha \prec Y \wedge |\alpha| > p$. Moreover, by Claim 3.6, $c(\alpha_{\tilde{P}_J} \upharpoonright p) = j_0$. Thus $c(Y_{\tilde{P}_J} \upharpoonright p) = j_0$. □

Finally, we slightly modify the proof of Theorem 3.5 to derive Theorem 3.1.

Proof of Theorem 3.1. The main point is to make the tree \mathcal{T} \emptyset' -computable. To ensure this, after we obtain $\tilde{\rho}_i, c_i$, we do not directly go to $\tilde{\rho}_{i+1}$. Instead, we \emptyset' -compute $\tilde{\rho}_i^0 \prec \tilde{\rho}_i^1 \prec \dots \prec \tilde{\rho}_i^{r_i}$ such that $\tilde{\rho}_i^0 \succ \tilde{\rho}_i$ and $c_i(\{\tau : \tau \succeq \tilde{\rho}_i^{r_i}\}) \subseteq c_i(\{\tilde{\rho}_i^0, \dots, \tilde{\rho}_i^{r_i}\})$. Then we \emptyset' -compute $\tilde{\rho}_{i+1} \succ \tilde{\rho}_i^{r_i}$ as in the proof of Theorem 3.5. Note that this indeed can be achieved using \emptyset' since c_i is computable. Define \mathcal{T} to be the set of all γ such that $(\forall i \leq |\gamma|)[\gamma(i) \in L_i, \gamma(i) \triangleleft \gamma(i+1)]$, and either $|\gamma| = 1 \wedge \gamma \in L_0$ or there exists $\tilde{\rho}_{|\gamma|-1}^u$ with $c_{|\gamma|-1}(\tilde{\rho}_{|\gamma|-1}^u) = \gamma(|\gamma| - 1)$. It is easy to see that \mathcal{T} is \emptyset' -computable since c_i is computable for all i and the sequences $\langle c_i : i \in \omega \rangle$ and $\langle \tilde{\rho}_i^v : i \in \omega, v \leq r_i \rangle$ are \emptyset' -computable.

Now we show that \mathcal{T} is a tree. Suppose $\gamma \in \mathcal{T}$, $|\gamma| = n+1$ with $n \geq 1$, and $c_n(\tilde{\rho}_n^u) = \gamma(n) = \langle P', \tilde{P}, j \rangle \in L_n$. We claim that $\gamma \upharpoonright n \in \mathcal{T}$. If $n = 1$, then $\gamma \upharpoonright 1 \in L_0 \subseteq \mathcal{T}$. Otherwise, let $\langle Q', \tilde{Q}, k \rangle \in L_{n-1}$ be the predecessor of $\langle P', \tilde{P}, j \rangle$. We need to show that there exists $\tilde{\rho}_{n-1}^v$ such that $c_{n-1}(\tilde{\rho}_{n-1}^v) = \langle Q', \tilde{Q}, k \rangle$. $c_n(\sigma) = \langle P', \tilde{P}, j \rangle$ implies that, letting $\alpha = \sigma_{P'}$, $c_{n-1}(\alpha) = c_{n-1}(\alpha \upharpoonright \min \tilde{P}) = j = \langle Q', \tilde{Q}, k \rangle$. Note that $\alpha \succeq \tilde{\rho}_{n-1}^{r_{n-1}}$ since $P' > |\tilde{\rho}_{n-1}^{r_{n-1}}|$. But $c_{n-1}(\{\tau : \tau \succeq \tilde{\rho}_{n-1}^{r_{n-1}}\}) \subseteq c_{n-1}(\{\tilde{\rho}_{n-1}^0, \dots, \tilde{\rho}_{n-1}^{r_{n-1}}\})$. Therefore there exists $\tilde{\rho}_{n-1}^v$ such that $c_{n-1}(\tilde{\rho}_{n-1}^v) = \langle Q', \tilde{Q}, k \rangle$. It follows that $\gamma \upharpoonright n \in \mathcal{T}$ and that \mathcal{T} is a tree. Any PA degree relative to \emptyset' computes an infinite path through \mathcal{T} . The rest of the proof goes exactly the same as Theorem 3.5. \square

We now give an alternative proof of Theorem 3.1 based on the definitional complexity of the solutions of c .

Second proof of Theorem 3.1. Let P_0, P_1, \dots be the \emptyset' -computable sequence defined in the proof of Theorem 3.5. We have seen that there exists an infinite ordered variable word such that the n th variable kind appears before the position $\max P_n$. Let \mathcal{T} be the tree of all finite ordered variable words which are finite solutions to c and such that the n th variable appears before the position $\max P_n$. By the previous observation, the tree is infinite, \emptyset' -computable, and \emptyset' -computably bounded. Any PA degree relative to \emptyset' computes an infinite variable word which, by construction of \mathcal{T} , is a solution to c . This completes the proof of Theorem 3.1. \square

Note that the above proof can be slightly modified to obtain a proof of a sequential version of the ordered variable word.

Statement 3.7. SeqOVW(n, ℓ) is the statement “If c_0, c_1, \dots is a sequence of ℓ -colorings of a fixed alphabet A of cardinality n , there exists a variable word W such that for every $i \in \omega$ and every $\bar{b} \in A^i$, $\{W(\bar{b}\bar{a}) : \bar{a} \in A^{<\infty}\}$ is monochromatic for c_i .”

Theorem 3.8. For every computable instance c_0, c_1, \dots of SeqOVW($2, \ell$), every PA degree relative to \emptyset' computes a solution to \bar{c} .

Proof. The proof is similar to Theorem 3.1. Using Lemma 3.4, we first construct a \emptyset' -computable sequence of strings $\tilde{\rho}_0 \prec \tilde{\rho}_1 \prec \dots$, a sequence of finite sets $P_i \subseteq \{|\tilde{\rho}_{i-1}|, \dots, |\tilde{\rho}_i| - 1\}$ and a sequence of colorings $d_i : [\tilde{\rho}_i]^\triangleleft \rightarrow L_i$ inductively as follows. $\tilde{\rho}_0 = \varepsilon$ and $d_0 = c_0$. Given $\tilde{\rho}_i$ and $d_i : [\tilde{\rho}_i]^\triangleleft \rightarrow L_i$, let $\tilde{\rho}_{i+1} \succeq \tilde{\rho}_i$ and $P_i \subseteq \{|\tilde{\rho}_i|, \dots, |\tilde{\rho}_{i+1}| - 1\}$ be such that $(\tilde{\rho}_{i+1}, P_i)$ - is d_i -valid, and let d_{i+1} be the coloring of $[\tilde{\rho}_{i+1}]^\triangleleft$ which on $\sigma \succeq \tilde{\rho}_{i+1}$ associates $\langle P', \tilde{P}, j, k \rangle$ such that (P', \tilde{P}) witnesses d_i -validity of $(\tilde{\rho}_{i+1}, P_i)$ for σ , $d_i(\sigma_{P'}) = j$ and $c_{i+1}(\sigma_{P'}) = k$. Note that the main difference with the previous construction is that we handle more and more colorings among c_0, c_1, \dots at each level. The remainder of the proof is the same as in Theorem 3.1. \square

The theorem above is optimal, in that we can obtain the following reversal.

Theorem 3.9. There is a computable instance c_0, c_1, \dots of SeqOVW($2, 2$), such that every solution is of PA degree relative to \emptyset' .

Proof. Let R_0, R_1, \dots be a uniformly computable sequence of sets such that for every e , if $\Phi_e^{\emptyset'}(e) \downarrow = 0$ then R_e is finite, and if $\Phi_e^{\emptyset'}(e) \downarrow = 1$ then R_e is cofinite. In particular, any function $f : \omega \rightarrow 2$ such that $f(e)$ gives a side of R_e which is infinite, is DNC₂ relative to \emptyset' , hence of PA degree relative to \emptyset' . Let $c_i : 2^{<\infty} \rightarrow 2$ be defined by $c_i(\sigma) = 1$ iff $|\sigma| \in R_i$, and let W be a solution to \bar{c} , that is, a variable word W such that for every $i \in \omega$ and every $\bar{b} \in A^i$, $\{W(\bar{b}\bar{a}) : \bar{a} \in A^{<\infty}\}$ is monochromatic for c_i . We claim that W computes such a function f . Given e , let $f(e) = c_e(W(\bar{b}))$, where $\bar{b} \in 2^e$ is arbitrary (this is well-defined, since $c_e(\bar{b})$ depends only on the length of \bar{b}). By definition of W , $\{W(\bar{b}\bar{a}) : \bar{a} \in A^{<\infty}\}$ is monochromatic for c_e , the color

of $c_e(W(\bar{b}))$ appears infinitely often in R_e . Therefore, W is of PA degree relative to \emptyset' . This completes the proof. \square

4. A DIFFICULT INSTANCE OF THE ORDERED VARIABLE WORD THEOREM

Miller and Solomon [5] constructed a computable instance of $\text{OVW}(2,2)$ with no Δ_2^0 solution. In this section, we strengthen their proof by constructing a computable instance of $\text{OVW}(2,2)$ such that every solution is of DNC degree relative to \emptyset' , using a significantly simpler argument.

The proof makes an essential use of a computable version of Lovasz Local Lemma proven by Remyantsev and Shen [7]. The idea of using Lovasz Local Lemma to analyse the computability-theoretic strength of problems in reverse mathematics comes from Csima and Dzhafarov, Hirschfeldt, Jockusch, Solomon and Westrick [2], who proved that a version of Hindman's theorem for subtractions is not computably true.

Definition 4.1. Fix a countable set of variables x_0, x_1, \dots . A (*disjunctive*) *clause* C is a tuple of the form $(x_{n_1} = i_1 \vee \dots \vee x_{n_k} = i_k)$, with $i_1, \dots, i_k < 2$. The *length* of C is the integer k . An *infinite CNF formula* is an infinite conjunction of disjunctive clauses. An infinite CNF formula $\bigwedge_n C_n$ is *computable* if the function which given n outputs a code for C_n is computable, and the set of n such that C_n contains the variable x_j is uniformly computable in j .

Theorem 4.2 (Remyantsev and Shen [7]). *For every $\alpha \in (0, 1)$, there exists some $N \in \omega$ such that every computable infinite CNF where each variable appears in at most $2^{\alpha n}$ clauses of size n (for every n) and all clauses have size at least N , has a computable satisfying assignment.*

Theorem 4.3. *There is a computable instance c of $\text{OVW}(2,2)$ and a computable function $h : \omega \rightarrow \omega$ such that if $\Phi_e^{\emptyset'}$ outputs a finite variable word in which the first $h(e)$ variable kinds occur, then $\Phi_e^{\emptyset'}$ is not extendible into an infinite solution to c .*

Proof. Fix $\alpha = 0.5$, and let N be the threshold of Theorem 4.2. For every index e and stage s , we interpret $\Phi_e^{\emptyset'}[s]$ as a finite variable word $W_{e,s}$ with exactly $N + e$ variable kinds, and where a new variable occurs right after $W_{e,s}$. Such a variable word induces a binary tree $T_{e,s}$ with 2^{N+e} leaves. Let $L_{e,s}$ be the set of leaves of $T_{e,s}$, that is, the set of all instantiations of the variable word $W_{e,s}$. Moreover, all the leaves of $T_{e,s}$ have the same length $n_{e,s}$.

The idea is the following: since the variable word is ordered and a new variable kind occurs right after $W_{e,s}$, no variable among the first $N + e$ variables can occur after $W_{e,s}$. If W is a solution to c with initial segment $W_e = \lim_s W_{e,s}$ for some color i , then W must be homogeneous for c for every instance of the variables, so in particular when setting all the variables after the $N + e$ first ones to 0. Hence, there must be infinitely many strings τ such that for every $\sigma \in \lim_s L_{e,s}$, $c(\sigma\tau) = i$. By ensuring that for cofinitely many τ , there is some $\sigma \in L_{e,|\tau|}$ such that $c(\sigma\tau) \neq i$, we force W_e not to be a solution to c for color i .

Fix a countable collection of variables $(x_\rho : \rho \in 2^{<\omega})$. Each variable x_ρ corresponds to the color of the string ρ . Given some $s \in \omega, \tau \in 2^{<\omega}$ and some $i < 2$, if $n_{e,s} + |\tau| = s$, then let $C_{e,s,\tau,i}$ be the disjunctive 2^{N+e} -clause

$$\bigvee \{x_{\sigma\tau} = i : \sigma \in L_{e,s}\}.$$

And let C be the conjunction

$$\bigwedge_{n_{e,s} + |\tau| = s} \{C_{e,s,\tau,i} : e \in \omega, \tau \in 2^{<\omega}, i < 2\}.$$

This infinite CNF formula is clearly computable. Clearly $C_{e,s,\tau,i}$ has length 2^{N+e} . Note that for every ρ, e , there exists at most one τ such that $(\exists \sigma \in L_{e,|\rho|})[\sigma\tau = \rho]$. Therefore, each variable x_ρ appears in at most 2 clauses of length 2^{N+e} , namely, $C_{e,|\rho|,\tau,0}$ and $C_{e,|\rho|,\tau,1}$, where τ is such that $(\exists \sigma \in L_{e,|\rho|})[\sigma\tau = \rho]$. Therefore, this formula satisfies the conditions of Theorem 4.2, and has a computable assignment $c : 2^{<\omega} \rightarrow 2$. By construction, letting $h(e) = N + e + 1$, the formula ensures that if $\Phi_e^{\emptyset'}$ outputs a finite variable word in which the first $h(e)$ variables kinds occur, then $\Phi_e^{\emptyset'}$ is not extendible into an infinite solution to c . \square

Definition 4.4. A function $f : \omega \rightarrow \omega$ is *diagonally non-computable relative to X* (or X -dnc) if for every e , $f(e) \neq \Phi_e^X(e)$.

Corollary 4.5. *There is a computable instance c of $\text{OVW}(2,2)$ such that every solution is of \emptyset' -dnc degree.*

Proof. Let c and h be as in Theorem 4.3. For every e , let α_e be a computable bijection from the finite variable words in which the first $h(e)$ variable kinds occur, to the set of the integers. By Kleene's fixpoint theorem, there is a computable function $g : \omega \rightarrow \omega$ such that for every e , $\Phi_{g(e)}^{\emptyset'} = \alpha_{g(e)}^{-1}(\Phi_e^{\emptyset'}(e))$.

Let W be a solution to c , that is, an infinite variable word. Let f be the W -computable function defined by $f(e) = \alpha_{g(e)}(w_e)$, where w_e is the first initial segment of W in which the first $h(g(e))$ variable kinds occur. We claim that f is \emptyset' -dnc. Indeed, given $e \in \omega$, $w_e \neq \Phi_{g(e)}^{\emptyset'}$, so

$$f(e) = \alpha_{g(e)}(w_e) \neq \alpha_{g(e)}(\Phi_{g(e)}^{\emptyset'}) = \Phi_e^{\emptyset'}(e)$$

This completes our proof. \square

We conclude this section with a small computational observation about $\text{VW}(2, 2)$ based on the syntactical form of the statement.

Definition 4.6. A function $g : \omega \rightarrow \omega$ *dominates* $f : \omega \rightarrow \omega$ if $(\forall x)f(x) < g(x)$. A function $f : \omega \rightarrow \omega$ is *hyperimmune* if it is not dominated by any computable function. A Turing degree is *hyperimmune-free* if it does not contain any hyperimmune function.

Lemma 4.7 (Folklore). *Let P be a statement of the form $(\forall X)[\Phi(X) \rightarrow (\exists Y)\Psi(X, Y)]$ where Φ is an arbitrary predicate, and Ψ is a Π_2^0 predicate. For every computable instance I of P , if I has a solution of hyperimmune-free degree, then every PA degree computes a solution to I .*

Proof. Say $\Psi(X, Y) \equiv (\forall x)(\exists y)\Theta(X \upharpoonright y, Y \upharpoonright y, x, y)$, where Θ is a decidable predicate. Let I be a computable P -instance with a solution S of hyperimmune-free degree. Let $h : \omega \rightarrow \omega$ be the S -computable function such that for every x , $\Theta(I, S, x, h(x))$ holds. In particular, there is a computable function $g : \omega \rightarrow \omega$ such that $(\forall x) \max(h(x), S(x)) < g(x)$. Let $T \subseteq \omega^{<\omega}$ be the computably bounded tree defined by

$$T = \left\{ \sigma \in \omega^{<\omega} : \begin{array}{l} (\forall x < |\sigma|)\sigma(x) < g(x) \wedge \\ (\forall x < |\sigma|)[g(x) < |\sigma| \rightarrow (\exists y < |\sigma|)\Theta(I \upharpoonright y, \sigma \upharpoonright y, x, y)] \end{array} \right\}$$

In particular, $S \in [T]$, so the tree is infinite. Moreover, any $R \in [T]$ is a solution to I , and any PA degree computes a member of $[T]$. This completes the proof. \square

Corollary 4.8. *There is a computable instance of $\text{VW}(2, 2)$ such that every solution is of hyperimmune degree.*

Proof. First, note that the statement $\text{VW}(2, 2)$ is of the form of Lemma 4.7. Let $c : 2^{<\omega} \rightarrow 2$ be the computable instance of $\text{VW}(2, 2)$ with no low solution constructed by Miller and Solomon [5] or by Theorem 4.3. Letting \mathbf{d} be a low PA degree, \mathbf{d} computes no solution to c , hence by Lemma 4.7, every solution to c is of hyperimmune degree. \square

It is still unknown whether there is a computable instance of $\text{OVW}(2, 2)$ such that every solution is PA over \emptyset' , or even just computes \emptyset' . In particular the following questions remain open:

Question 4.9. Does $\text{VW}(2, 2)$ or $\text{OVW}(2, 2)$ imply ACA over RCA_0 ?

Question 4.10. Is there a computable instance of $\text{VW}(2, 2)$ or $\text{OVW}(2, 2)$ such that the measure of oracles computing a solution to it is null?

REFERENCES

- [1] Timothy J. Carlson and Stephen G. Simpson. A dual form of Ramsey's theorem. *Adv. in Math.*, 53(3):265–290, 1984.
- [2] Barbara F Csima, Damir D Dzhabarov, Denis R Hirschfeldt, Carl G Jockusch Jr, Reed Solomon, and Linda Brown Westrick. The reverse mathematics of hindman's theorem for sums of exactly two elements. *arXiv preprint arXiv:1804.09809*, 2018.
- [3] Reed Solomon Linda Brown Westrick Damir D. Dzhabarov, Stephen Flood. Effectiveness for the dual ramsey theorem. To appear. Available at <http://www.math.uconn.edu/~damir/papers/dualRT.pdf>, 2017.
- [4] Harvey Friedman and Stephen G. Simpson. Issues and problems in reverse mathematics. In *Computability theory and its applications (Boulder, CO, 1999)*, volume 257 of *Contemp. Math.*, pages 127–144. Amer. Math. Soc., Providence, RI, 2000.
- [5] Joseph S. Miller and Reed Solomon. Effectiveness for infinite variable words and the dual Ramsey theorem. *Arch. Math. Logic*, 43(4):543–555, 2004.
- [6] Antonio Montalbán. Open questions in reverse mathematics. *Bulletin of Symbolic Logic*, 17(03):431–454, 2011.
- [7] Andrei Rumyantsev and Alexander Shen. Probabilistic constructions of computable objects and a computable version of lovász local lemma. *Fundamenta Informaticae*, 132(1):1–14, 2014.

- [8] Stephen G. Simpson. *Subsystems of Second Order Arithmetic*. Cambridge University Press, 2009.
- [9] Theodore A. Slaman. A note on dual ramsey theorem. Unpublished, January 1997.

DEPARTMENT OF MATHEMATICS, CENTRAL SOUTH UNIVERSITY, CHANGSHA 410083, PEOPLE'S REPUBLIC OF CHINA
E-mail address: `g.jiayi.liu@gmail.com`

LACL, DÉPARTEMENT D'INFORMATIQUE, FACULTÉ DES SCIENCES ET TECHNOLOGIE, 61 AVENUE DU GÉNÉRAL DE GAULLE,
94010 CRÉTEIL CEDEX
E-mail address: `benoit.monin@computability.fr`

INSTITUT CAMILLE JORDAN, UNIVERSITÉ CLAUDE BERNARD LYON 1, 43 BOULEVARD DU 11 NOVEMBRE 1918, F-69622
VILLEURBANNE CEDEX
E-mail address: `ludovic.patey@computability.fr`