



HAL
open science

Comparison of European eel sperm cryopreservation protocols with standardization as a target

J.G. Herranz-Jusado, V. Gallego, M. Morini, C. Rozenfeld, L. Pérez, E. Kása, T. Kollár, Alexandra Depince, Catherine Labbé, J.F. Asturiano

► **To cite this version:**

J.G. Herranz-Jusado, V. Gallego, M. Morini, C. Rozenfeld, L. Pérez, et al.. Comparison of European eel sperm cryopreservation protocols with standardization as a target. *Aquaculture*, 2019, 498, pp.539-544. 10.1016/j.aquaculture.2018.09.006 . hal-01892956

HAL Id: hal-01892956

<https://hal.science/hal-01892956>

Submitted on 23 Nov 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Comparison of European eel sperm cryopreservation protocols with standardization as a target

J.G. Herranz-Jusdado¹, V. Gallego¹, M. Morini¹, C. Rozenfeld¹, L. Pérez¹, E. Kása², T. Kollár², A. Depincé³, C. Labbé³, Á. Horváth², J.F. Asturiano^{1*}

¹*Grupo de Acuicultura y Biodiversidad. Instituto de Ciencia y Tecnología Animal. Universitat Politècnica de València, Edificio 7G, Camino de Vera s/n. 46022 Valencia (Spain).*

²*Department of Aquaculture, Szent István University, Páter Károly u. 1., H-2100 Gödöllő, Hungary*

³*INRA, Fish Physiology and Genomics UR 1037, Campus de Beaulieu, Rennes France.*

*Corresponding author:

Dr Juan F. Asturiano.

Email: jfastu@dca.upv.es

Phone: (+34) 96 387 93 85

Abstract

The critical situation of the European eel (*Anguilla anguilla*) has urged the development of sperm cryopreservation protocols for reproduction in captivity and cryobanking. In the last years, two research groups have developed their own protocols in Spain and Hungary with positive results, but difficult to compare.

Here, a series of experiments were conducted to test the quality of thawed sperm after using both protocols, determining which of them produce the best results and aiming for standardization. The quality of thawed sperm was assessed by studying the motility and kinetic values of thawed sperm from both cryopreservation protocols using a computer-assisted sperm analysis (CASA-Mot) system. In addition, a viability analysis was performed using flow cytometry to test if the cryoprotectants or the freezing-thawing process led to a reduction in spermatozoa survival. Furthermore, since during cryopreservation the sperm was treated with methylated cryoprotectants (DMSO or methanol) that may induce epigenetic changes in the sperm DNA (cytosine methylation) and could affect the offspring, we conducted a luminometric methylation assay (LUMA) to study the DNA methylation levels induced by both protocols.

In this work, all the above-mentioned parameters were analyzed in fresh and frozen-thawed sperm samples. Our results showed that thawed sperm samples from both protocols presented lower sperm motility and velocity, and lower percentage of live cells than those shown in fresh sperm samples. Furthermore, sperm samples from the methanol based protocol showed significantly higher motility, velocity and percentage of live spermatozoa than the same sperm samples treated with the DMSO based protocol. In addition, the DMSO based protocol induced a hypomethylation of sperm DNA compared to fresh samples whereas the methanol based protocol did not alter sperm DNA methylation level. Our results indicate that the methanol based protocol is a more suitable protocol that preserves better the motility and genetic qualities of the European eel sperm.

Keywords: *Anguilla anguilla*; Methanol; DMSO; DNA methylation; Epigenetics

Abbreviations: DMSO, dimethyl sulfoxide; ROS, reactive oxygen species; rhCG, recombinant human chorionic gonadotropin; CASA-Mot, computer assisted sperm analysis motility module; FBS, foetal bovine serum; HBSS, Hank's balanced salt solution.

1. Introduction

During the last years, a drastic decrease has been observed in the number of European eels (*Anguilla anguilla*) returning from Europe and North Africa to the spawning sites in the Atlantic Ocean (Dekker 2000; Jacoby & Gollock, 2014). Several impacts such as water pollution, overfishing or habitat fragmentation, have led the European eel to be included on the IUCN red list as critically endangered (Jacoby & Gollock, 2014). Consequently, the development of techniques and protocols for reproduction in captivity are necessary to reverse this situation.

The maturation of the European eel in captivity is only achieved by costly and long hormonal treatments (Asturiano et al., 2006; Gallego et al., 2012; Pérez et al., 2000), and still the production of gametes in both sexes can be unsynchronized (Asturiano et al., 2016). During the last years, several researchers have worked in the development of new maturation protocols such as alternative hormonal treatments with recombinant hormones (Peñaranda et al., 2018) or androgen implants (Di Biase et al., 2017; Mordenti et al., 2018), but the timing of final maturation in females is still highly variable and difficult to control (Mylonas et al., 2017). Therefore, the development of cryopreservation protocols for European eel sperm has been considered important for reproduction management, by guaranteeing the availability of both types of gametes when female spawns (Asturiano et al., 2017), besides its application for cryobanking and future broodstock management.

Cryopreservation of European eel sperm has been faced by different groups since early 2000s. Mainly two groups of research established successfully their own cryopreservation protocols in Spain (Asturiano et al., 2003; 2004; Peñaranda et al., 2009) and Hungary (Müller et al., 2004; Szabó et al., 2005). These protocols differ greatly in many aspects such as the composition of the extenders, the cryoprotectants used, the volume of the straws or the cooling rates within others, evidencing the need for standardization (Asturiano et al., 2017; Rosenthal et al., 2010).

The success of a sperm cryopreservation protocol is commonly assessed using parameters such as sperm viability and motility, fertilizing capacity and the quality of the offspring

(Cabrita et al., 2010). However, in the case of the eel both protocols have yielded high post-thaw sperm viability (58 to 63%) and motility values ranging between 18 and 38% (Asturiano et al., 2017). Furthermore, the fertilizing capacity of the Spanish protocol (from now on referred as DMSO protocol) was successfully tested by producing European eel larvae after fertilization with thawed sperm (Asturiano et al., 2016), and following the Hungarian protocol (from now on referred as methanol protocol), hybrid larvae were successfully produced using thawed sperm from European eel and eggs from Japanese eel (*Anguilla japonica*) (Müller et al., 2012; 2018). In this last study, Müller and collaborators showed that the malformation rate of larvae was higher when using cryopreserved sperm than in the control groups using fresh sperm, suggesting that the cryopreservation methodology needs further refinement. Additionally, a growing concern is that the epigenetic effects of cryopreservation on the sperm DNA might be altered by the freezing, cryobanking and thawing process (Labbé et al., 2017; Pérez-Cerezales et al., 2010). The use of methylated cryoprotectants such as methanol or dimethyl sulfoxide (DMSO) has been proven to produce reactive oxygen species (ROS) that can induce cytosine methylation in fish sperm (De Mello et al., 2017; Kawai et al., 2010). Methylation of cytosine residues in DNA is considered to be one of the major epigenetic mechanisms stabilizing gene silencing (Schaefer et al., 2007). Furthermore, cytosine methylation can be altered by cryopreservation, inducing hypo- and hypermethylation profiles in sperm DNA (Labbé et al., 2017). Therefore, the study of epigenetic effects of cryopreservation may be a good indicator of the success of a cryopreservation protocol, since damaged DNA or abnormal DNA regulation have been observed to have a negative effect on the generated embryos (Herráez et al., 2017).

The main objective of this work was to compare the main protocols previously developed for European eel sperm cryopreservation, aiming for standardization. The comparison was made considering sperm quality after thawing, when sperm motilities, sperm velocities, and sperm viability were analyzed. Furthermore, epigenetic effects of sperm cryopreservation were studied by analyzing whether DNA methylation patterns were affected by the different cryopreservation protocols.

2. Material and methods

2.1. Ethics statement

The protocol was approved by the Experimental Animal Ethics Committee from the Universitat Politècnica de València (UPV) and final permission was given by the local government (Generalitat Valenciana, Permit Number: 2015/VSC/PEA/00064).

2.2. Fish handling

For this experiment, 28 immature male European eels from the fish farm Valenciana de Acuicultura S.A. (Puzol, Valencia) were brought to our facilities in the Universitat Politècnica de València. Fish were distributed in two 200 L aquaria with recirculation systems, and thermostats and coolers to maintain water temperature at 20 °C. They were gradually acclimated to seawater (salinity 37 ± 0.2 g/L) increasing the salinity 10 ppt each 2 days for 8 days, and 2 days more of resting at 37 ppt. The aquaria were covered to maintain a constant shadow and reduce fish stress.

After 10 days of acclimation, male fish anesthetized with benzocaine (60 ppm) were weekly treated with injections of recombinant human chorionic gonadotropin (rhCG; Ovitrelle, Madrid, Spain, 1.5 IU/g fish) to induce maturation and spermiation (Gallego et al., 2012; Pérez et al., 2000). From the sixth week of hormonal treatment, sperm samples were collected weekly, 24h after the hormone injections.

For sperm collection, fish were anesthetized with benzocaine. Thereafter, the genital area was carefully cleaned with distilled water and thoroughly dried with paper to avoid contamination with feces, urine or seawater to avoid accidental sperm activation. Then, sperm was collected by applying a ventral massage from the pectoral fins to the genital opening and collected in graduated Falcon tubes using a vacuum pump.

Sperm samples were collected after 11-14 weeks of hormonal treatment. The samples were diluted 1:9 (sperm:extender) in P1 medium (in mM: NaCl 125, NaHCO₃ 20, MgCl₂ 2.5, CaCl₂

1, KCl 30; pH adjusted to 8.5, described by Peñaranda *et al.*, 2010) , kept in plastic tubes at 4 °C and evaluated for motility.

2.3. Evaluation of sperm motility

In a maximum of 2 h after the sperm extraction, sperm samples were evaluated in triplicates following the method described by Gallego *et al.* (2013). Briefly, each sperm sample was activated by mixing 0.5 µL of P1-diluted sperm sample with 4.5 µL of artificial seawater (in mM: NaCl 354.7, MgCl₂ 52.4, CaCl₂ 9.9, Na₂SO₄ 28.2, KCl 9.4, in distilled water) with 2% (w:v) bovine serum albumin (BSA), pH adjusted to 8.2 and osmolality of 1100 mOsm/kg. The activation was performed in an ISAS Spermtrack 10 counting chamber (Proiser R+D, S.L., Spain) on a microscope in negative phase with a 10X magnification (Nikon Eclipse 80i) connected to a computer with an ISAS 782M camera (Proiser R+D, S.L., Spain), recording 60 frames per second (fps). All samples were analyzed 15 s after activation, using the CASA module ISAS v1 software (Proiser R+D, S.L., Spain). Several kinetic parameters such as percentage of motile spermatozoa (MOT, %), progressive motility (pMOT, %), curvilinear velocity (VCL, µm/s), straight-line velocity (VSL, µm/s), and average path velocity (VAP, µm/s), as well as percentage of slow (average path velocity (VAP) = 10-50 µm/s), medium (VAP = 50-100 µm/s) and fast (VAP >100 µm/s) spermatozoa were recorded for further analysis (Gallego and Asturiano 2018a for details). Samples with motility values higher than 65% were selected for cryopreservation.

2.4. Experimental design

A total number of 18 sperm samples were selected for cryopreservation. Each sample was first evaluated for motility and then frozen and thawed following both protocols. In addition, before freezing, each sample was evaluated for motility approximately 10 minutes after diluted with the freezing media corresponding to each protocol. Then, four straws (IMV Technologies, l'Aigle, France) of 250 µL for the DMSO protocol and four straws of 500 µL for the methanol protocol were frozen. Therefrom, three straws per protocol were thawed and

immediately analyzed with CASA-Mot for sperm quality. Moreover, 50 μ L of fresh and thawed sperm from each sample were used for the viability analysis using the flow cytometer (see down). The left straw per protocol was maintained frozen in liquid nitrogen and was sent to INRA's lab in Rennes (France) for sperm epigenetic analysis, by studying the DNA methylation level. In addition, 100 μ L of fresh sperm from each sample were frozen as well by directly throwing the tube with the sperm into the liquid nitrogen and then storing it at -80 °C for DNA methylation analysis of the sperm control. We demonstrated previously that such snap freezing allows that the DNA methylation level of the fresh sperm is preserved (unpublished data).

2.5. Cryopreservation protocols

Every selected sample was frozen and thawed following both protocols. For the DMSO protocol, a freezing medium was prepared in advance by mixing a modified P1 extender solution (in mM: NaCl 50, NaHCO₃ 100, MgCl₂ 2.5, CaCl₂ 1, KCl 30; described by Peñaranda *et al.*, 2009; and named M5 in that paper), 25% (v/v) of fetal bovine serum (FBS) and 10% (v/v) of DMSO. The freezing medium was adjusted to a pH of 6.5, an osmolality of 330 mOsm/kg and maintained at 4 °C. Thereafter, a dilution 1:2 of sperm: freezing medium, was prepared and immediately packed in 250 μ L straws, sealed with modeling clay and frozen for 5 min in liquid nitrogen vapor 1 cm above the surface using a floating structure. Following, the straws were thrown into the liquid nitrogen where the sperm was preserved as long as needed. The thawing consisted in a water bath at 30 °C for 8 s.

For the methanol protocol, modified Tanaka's extender (in mM: NaCl 137, NaHCO₃ 76.2) was prepared in advance and maintained at 4 °C. Then, a dilution consisting in sperm:Tanaka's extender:methanol (1:8:1) was prepared and packed in 500 μ L straws, and frozen for 3 min in liquid nitrogen vapor 3 cm above the liquid nitrogen before throwing the straws into the liquid nitrogen. For thawing, the straws were immersed in a water bath at 40 °C for 13 s.

2.6. Thawed sperm evaluation

The quality of thawed sperm samples was assessed by analyzing several sperm motility parameters with CASA-Mot, sperm viability (membrane integrity) with a flow cytometer and epigenetic effects with an analysis of sperm DNA methylation pattern.

The motility analysis was performed using CASA-Mot as explained above. In addition, a viability analysis was conducted with flow cytometry using a fluorescence kit (LIVE/DEAD Sperm Viability Kit, Thermo Fisher Scientific, MA, USA) containing the membrane-permeating dye SYBR 14, that stains the nuclei of membrane-intact cells fluorescent green and the non-permeating propidium iodide (PI), that counterstains the nuclei of cells with a damaged membrane fluorescent red. Here, 0.5 μ L of SYBR 14 (100 μ M) and 2 μ L of PI (2.4 mM) were added to 50 μ L of fresh or thawed sperm samples and incubated at room temperature in the dark for 10 min. Thereafter, samples were diluted in 500 μ L of extender solution (P1 medium for the Spanish protocol or Tanaka's medium for the Hungarian protocol) and were analyzed with a flow cytometer (Beckman Coulter FC500). The analyses were performed using the voltages: SS= 199, FS= 199, FL1= 377 and FL2= 372; for a maximum number of 5,000 events or 15 s at low flow.

Finally, a study of DNA methylation level was conducted in fresh and thawed sperm. Sperm DNA was extracted using the phenol/chloroform method: about 20×10^6 spermatozoa in 10 μ L Hank's balanced salt solution (HBSS) 300 were digested overnight at 42 °C under agitation in 1 mL of TNES buffer (125 mM NaCl, 10 mM EDTA, 17 mM SDS, 4 M urea, 10 mM Tris-HCl, pH 8) with 75 μ g of proteinase K (Sigma Aldrich, P6556). One mL phenol-chloroform-isoamyl alcohol (25:24:1) was added and vigorously mixed. After centrifugation for 15 min at 8,000 g at 4 °C, the upper phase (800 μ L) was mixed with 200 μ L NaCl 5 M and 2 mL of cold (-20 °C) 100% ethanol. After centrifugation, the dried DNA pellet was mixed with 100 μ g/mL RNase in water (Promega, A7973) and incubated 1 h at 37 °C. Whole DNA methylation level was estimated using LUMA (luminometric methylation assay) (Karimi et al., 2006). Genomic DNA from each sperm (0.5-1 μ g) was digested 4 h at 37 °C with 7.5 units of either HpaII and EcoRI (NEB R3101) or MspI and EcoRI in a total volume of 30 μ L in duplicate. For

pyrosequencing of the digested samples, 20 μ L of digested DNA were mixed with 20 μ L of annealing buffer (Qiagen, 979009) and samples were placed in a Qiagen PyroMark Q96 ID. The instrument was programmed to add dNTPs in the following order: A, C+G, T, C+G, water, A, T. Peak heights (PH) were analyzed using the PyroMark Q96 software. A and T peaks refers to the amount of DNA cleaved by EcoRI (DNA content controls) whereas C + G peaks show the amount of DNA cleaved by MspI and HpaII. The percentage of methylation was calculated as $100 \times (1 - (\text{PH HpaII} / \text{PH MspI}))$. The PH HpaII/PH MspI ratio was calculated by doing $(\text{PH HpaII} / \text{PH EcoRI}) / (\text{PH MspI} / \text{PH EcoRI})$.

2.7. Statistical analysis

Sperm viability and motility parameters were subjected to analysis of variance (General Linear Model, GLM). As fixed effect was chosen fresh or thawed sperm from both protocols, i.e. “fresh sperm”, “thawed DMSO” and “thawed methanol”. For all models, an examination of the residual plots verified that no systematic patterns occurred in the errors. Model results of p -values < 0.05 were considered significant.

For the statistical analysis of DNA levels, a non-parametric test (paired Wilcoxon test) was performed. Differences were considered as significant if $p < 0.05$.

All analyses were conducted in the R-environment (R_Development_Core_Team, 2010).

3. Results

Results from this comparison experiment showed that all samples, independently of the protocol used, decreased their percentage of motile cells and cell velocity after cryopreservation (Figure 1). In addition, the motility results from thawed samples of sperm cryopreserved with the methanol protocol showed higher motility ($32.4 \pm 1.8\%$) than those from the DMSO protocol ($10.8 \pm 0.9\%$) (Figure 1). All the sperm kinetic parameters analyzed showed the same pattern, with higher motility and faster velocities in samples preserved with the methanol protocol than those preserved with the DMSO one (Figure 1). Furthermore, the proportion of fast cells (faster than $100 \mu\text{m/s}$) was also significantly reduced after

cryopreservation (Figure 2). Nevertheless, thawed samples of sperm cryopreserved with the methanol protocol presented a higher percentage ($47.9 \pm 1.5\%$) of fast cells than using the DMSO one ($29.6 \pm 2.1\%$). Note that the sperm was instantly activated when diluted in the freezing medium of the DMSO protocol before freezing, clearly affecting the motility after thawing (Figure 3), whereas samples diluted in the freezing medium containing methanol were not activated (no differences with fresh samples) and did not affect the sperm motility prior to freezing (Figure 3).

Cell viability results (Figure 4) showed that there were more live spermatozoa in thawed sperm samples from the methanol protocol than from the DMSO one, and although survival in both cases was high ($>75\%$), it was still lower than viability measured in fresh sperm samples.

The analysis of cysteine methylation in fresh and thawed sperm (Figure 5) showed that sperm samples treated with the DMSO protocol had lower DNA methylation than fresh samples and samples treated with the methanol protocol, whereas these two showed no differences between each other.

4. Discussion

In this work, we described and compared the two main protocols available for European eel sperm cryopreservation. Our results indicated that in every case, the sperm motility of thawed sperm was lower than in fresh sperm. The reduction in post-thawing sperm quality compared to fresh sperm is consistent with the available bibliography, although there is a great variation between fish species (Asturiano et al., 2017; Horváth et al., 2015). For instance, Dziewulska et al. (2011) used several cryoprotectants (DMSO and methanol as in the present study) to freeze fresh sperm samples of Atlantic salmon (*Salmo salar*) with a motility of 70-95%. The study showed that the sperm motility after thawing was significantly lower than in fresh samples, with post-thawing motility values in the best protocol of 8.2%, using DMSO as cryoprotectant. Oppositely, a different study with cryopreserved sperm from brown trout (*Salmo trutta*) using methanol as cryoprotectant, obtained motilities of thawed

sperm higher than 60%, which represented a reduction of only 20% of motility compared to fresh samples (Horváth et al., 2015).

In the present study, the data of sperm quality from thawed samples showed that cryopreservation using the methanol protocol, caused higher motility values than the DMSO protocol. Although the values obtained with the methanol protocol were consistent with the bibliography (Müller et al., 2004; Szabó et al., 2005), the motility results from the DMSO protocol were lower than previously reported (Asturiano et al., 2003, 2004; Peñaranda et al., 2009). Although the samples were frozen immediately after the addition of the freezing media containing DMSO to the sperm, it has been proved that the presence of DMSO in the freezing media activates the European eel sperm (Peñaranda et al., 2009), and lead to a reduced post-thawed sperm motility. Even though the DMSO protocol was improved to reduce activation by increasing the concentration of NaHCO_3 , decreasing the pH of the media (Peñaranda et al., 2009; Vílchez et al., 2017), fast manipulation was still required. In this study, we show that the sperm was activated after diluting in the freezing media (before freezing) containing DMSO. This pre-freezing activation naturally affects the final sperm motility of thawed sperm samples.

Although DMSO is the most used cryoprotectant for fish sperm (Gallego & Asturiano, 2018b; Martínez-Páramo et al., 2017), methanol has also been widely used in freshwater species such as sturgeons, salmonids, tench or Eurasian perch within others (reviewed by Asturiano *et al.*, 2017). Furthermore, it has been recently used in cryopreservation protocols for Japanese eel sperm (Koh et al., 2017; Müller et al., 2017, 2018). As cryoprotectant, methanol has been reported to penetrate more rapidly the cells and being less toxic than DMSO (Horváth et al., 2015). In addition, methanol is osmotically inert and therefore does not activate sperm by osmotic shock (De Baulny et al., 1997; Horváth et al., 2005). In our study, the methanol was apparently less toxic than the DMSO, because thawed samples from the methanol protocol presented higher survival than samples from the DMSO protocol. Furthermore, we confirmed that since methanol is osmotically inert, it did not activate the sperm, oppositely to the DMSO that activated the sperm due to the increase of osmolality.

This difference could partially explain the higher motility and velocity of thawed samples treated with the methanol protocol. Furthermore, both protocols differ in other aspects such as extender composition, dilution rate, volume and freezing rate that could also affect the thawed sperm motility.

Sperm from the DMSO protocol presented a loss of methylation compared to fresh sperm, whereas sperm from the methanol protocol remained similar to the fresh control. Changes in cytosine methylation levels after cryopreservation have been little explored in fish. Primarily, the concern arose for the use of methylating cryoprotectants that in the presence of ROS may led to cytosine methylation (Kawai et al., 2010). Indeed, Riesco and Robles (2013) observed in zebrafish that some promoter regions were hypermethylated after genital ridge cryopreservation in DMSO. However, in tambaqui (*Colossoma macropomum*) sperm, cryopreservation with either DMSO or methanol induced in both cases a sperm DNA hypomethylation (De Mello et al., 2017), contrarily to what could have been expected from the model study from Kawai et al. (2010). It is therefore not clear if the cryoprotectant molecule is the main parameter affecting DNA methylation. It was reported that cryopreservation-induced changes in DNA methylation could be species dependent (Labbé et al., 2017), and that cryopreservation with methods which are not optimal for a given species would induce more epigenetic effect (Labbé et al., 2014). In our case, the fact that the methanol protocol did not change the overall DNA methylation level would indicate that the epigenetic risk is reduced with this method.

5. Conclusions

In conclusion, this study show that the methanol cryopreservation protocol, is nowadays the most suitable protocol for European eel sperm cryopreservation, giving the best sperm motility, sperm velocity and cell survival values. Furthermore, the methylation level of sperm DNA from thawed samples with this method are the same as in fresh sperm, indicating that there are not drastic epigenetic changes when sperm is cryopreserved in this way.

Acknowledgements

Funded by the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement N° 642893 (IMPRESS), including the JGHJ and CR predoctoral contracts, the French CRB Anim project «Investissements d'avenir», ANR-11-INBS-0003, as well as the Hungarian EFOP 3.6.3.-VEKOP 16.-2017-00008 project, and by the Higher Education Institutional Excellence Program (1783-3/2018/FEKUTSRAT) awarded by the Ministry of Human Capacities within the framework of water related researches of Szent István University. MM, EK, TK and ÁH were granted with Short Term Scientific Missions by the COST Office (COST Action FA1205: AQUAGAMETE).

Declaration of interest

The authors declare no conflict of interests

References

- Asturiano, J.F., Pérez, L., Marco-Jiménez, F., Olivares, L., Vicente, J.S., Jover, M., 2003. Media and methods for the cryopreservation of European eel (*Anguilla anguilla*) sperm. *Fish Physiology and Biochemistry*. 28, 501-502. <https://doi.org/10.1023/B:FISH.0000030640.36044.c8>
- Asturiano, J.F., Pérez, L., Garzón, D.L., Marco-Jiménez, F., Peñaranda, D.S., Vicente, J.S., Jover, M., 2004. Physio-chemical characteristics of seminal plasma and development of media and methods for the cryopreservation of European eel sperm. *Fish Physiology and Biochemistry*. 30, 283-293. <https://doi.org/10.1007/s10695-005-1553-x>.
- Asturiano, J.F., Marco-Jiménez, F., Pérez, L., Balasch, S., Garzón, D.L., Peñaranda, D.S., Vicente, J.S., Viudes-de-Castro, M.P., Jover, M., 2006. Effects of hCG as spermiation inducer on European eel semen quality. *Theriogenology*. 66, 1012-1020. <https://doi.org/10.1016/j.theriogenology.2006.02.041>.
- Asturiano, J.F., Sørensen, S.R., Pérez, L., Lauesen, P., Tomkiewicz, J., 2016. First production of larvae using cryopreserved sperm: effects of preservation temperature and cryopreservation on European eel sperm fertilization capacity. *Reproduction in Domestic Animals*. 51, 485-491. <https://doi.org/10.1111/rda.12706>.
- Asturiano, J.F., Cabrita, E., Horváth, Á., 2017. Progress, challenges and perspectives on fish gamete cryopreservation: a mini-review. *General and Comparative Endocrinology*. 245, 69-76. <https://doi.org/10.1016/j.ygcen.2016.06.019>.
- Cabrita, E., Sarasquete, C., Martínez-Páramo, S., Robles, V., Beirao, J., Pérez-Cerezales, S., Herráez, M., 2010. Cryopreservation of fish sperm: applications and perspectives. *Journal of Applied Ichthyology*. 26, 623-635. <https://doi.org/10.1111/j.1439-0426.2010.01556.x>.
- De Baulny, B.O., Le Vern, Y., Kerboeuf, D., Maise, G., 1997. Flow cytometric evaluation of mitochondrial activity and membrane integrity in fresh and cryopreserved rainbow trout (*Oncorhynchus mykiss*) spermatozoa. *Cryobiology*. 34, 141-149. <https://doi.org/10.1006/cryo.1996.1992>.

- De Mello, F., Garcia, J.S., Godoy, L.C., Depincé, A., Labbé, C., Streit Jr, D.P., 2017. The effect of cryoprotectant agents on DNA methylation patterns and progeny development in the spermatozoa of *Colossoma macropomum*. *General and Comparative Endocrinology*. 245, 94-101. <https://doi.org/10.1016/j.ygcen.2016.06.003>
- Dekker, W., 2000. The fractal geometry of the European eel stock. *ICES Journal of Marine Science: Journal du Conseil*. 57, 109-121. <https://doi.org/10.1006/jmsc.1999.0562>.
- Di Biase, A., Lokman, P.M., Govoni, N., Casalini, A., Emmanuele, P., Parmeggiani, A., Mordenti, O., 2017. Co-treatment with androgens during artificial induction of maturation in female eel, *Anguilla anguilla*: Effects on egg production and early development. *Aquaculture*. 479, 508-515.
<https://doi.org/10.1016/j.aquaculture.2017.06.030>.
- Dziewulska, K., Rzemieniecki, A., Czerniawski, R., Domagała, J., 2011. Post-thawed motility and fertility from Atlantic salmon (*Salmo salar* L.) sperm frozen with four cryodiluents in straws or pellets. *Theriogenology*. 76, 300-311.
<https://doi.org/10.1016/j.theriogenology.2011.02.007>.
- Gallego, V., Mazzeo, I., Vílchez, M.C., Peñaranda, D.S., Carneiro, P.C.F., Pérez, L., Asturiano, J.F., 2012. Study of the effects of thermal regime and alternative hormonal treatments on the reproductive performance of European eel males (*Anguilla anguilla*) during induced sexual maturation. *Aquaculture*. 354, 7-16.
<https://doi.org/10.1016/j.aquaculture.2012.04.041>.
- Gallego, V., Carneiro, P.C.F., Mazzeo, I., Vílchez, M.C., Peñaranda, D.S., Soler, C., Pérez, L., Asturiano, J.F., 2013. Standardization of European eel (*Anguilla anguilla*) sperm motility evaluation by CASA software. *Theriogenology*. 79, 1034-1040.
<https://doi.org/10.1016/j.theriogenology.2013.01.019>.
- Gallego, V., Asturiano, J.F., 2018a. Sperm motility in fish: technical applications and perspectives through CASA-Mot systems. *Reproduction, Fertility and Development*. 30, 820-832. <https://doi.org/10.1071/RD17460>.

- Gallego, V., Asturiano, J.F., 2018b. Fish sperm motility assessment as a tool for aquaculture research: a historical approach. *Reviews in Aquaculture*, 1-28.
<https://doi.org/10.1111/raq.12253>.
- Herráez, M.P., Ausió, J., Devaux, A., González-Rojo, S., Fernández-Díez, C., Bony, S., Saperas, N., Robles, V., 2017. Paternal contribution to development: Sperm genetic damage and repair in fish. *Aquaculture*. 472, 45-59.
<https://doi.org/10.1016/j.aquaculture.2016.03.007>.
- Horváth, Á., Wayman, W.R., Urbányi, B., Ware, K.M., Dean, J.C., Tiersch, T.R., 2005. The relationship of the cryoprotectants methanol and dimethyl sulfoxide and hyperosmotic extenders on sperm cryopreservation of two North-American sturgeon species. *Aquaculture*. 247, 243-251. <https://doi.org/10.1016/j.aquaculture.2005.02.007>.
- Horváth, Á., Labbé, C., Jesenšek, D., Hoitsy, G., Bernáth, G., Kaczkó, D., Bokor, Z., Urbányi, B., 2015. Post-thaw storage of sperm from various salmonid species. *Journal of Applied Ichthyology*. 31, 119-124. <https://doi.org/10.1111/jai.12741>
- Jacoby, D., Gollock, M., 2014. *Anguilla anguilla*. The IUCN red list of threatened species. Version 2014.2.
- Karimi, M., Johansson, S., Stach, D., Corcoran, M., Grandér, D., Schalling, M., Bakalkin, G., Lyko, F., Larsson, C., Ekström, T.J., 2006. LUMA (LUminometric Methylation Assay)—a high throughput method to the analysis of genomic DNA methylation. *Experimental cell research*. 312, 1989-1995. <https://doi.org/10.1016/j.yexcr.2006.03.006>.
- Kawai, K., Li, Y.S., Song, M.F., Kasai, H., 2010. DNA methylation by dimethyl sulfoxide and methionine sulfoxide triggered by hydroxyl radical and implications for epigenetic modifications. *Bioorganic and Medicinal Chemistry Letters*. 20, 260-265.
<https://doi.org/10.1016/j.bmcl.2009.10.124>.
- Koh, I.C.C., Hamada, D., Tsuji, Y.-A., Okuda, D., Nomura, K., Tanaka, H., Ohta, H., 2017. Sperm cryopreservation of Japanese eel, *Anguilla japonica*. *Aquaculture*. 473, 487-492. <https://doi.org/10.1016/j.aquaculture.2017.03.011>.

- Labbé, C., Depincé, A., Milon, P., Morini, M., Riesco, M., Robles, V., Asturiano, J.F., Horváth, Á., Herráez, M.P., 2014. DNA methylation of fish germ cells and the risk of alteration after cryopreservation, 10th International Symposium on Reproductive Physiology of Fish. Olhão (Portugal). Book of abstracts, pp. 48.
- Labbé, C., Robles, V., Herráez, M.P., 2017. Epigenetics in fish gametes and early embryo. *Aquaculture*. 472, 93-106. <https://doi.org/10.1016/j.aquaculture.2016.07.026>.
- Martínez-Páramo, S., Horváth, Á., Labbé, C., Zhang, T., Robles, V., Herráez, P., Suquet, M., Adams, S., Viveiros, A., Tiersch, T.R., 2017. Cryobanking of aquatic species. *Aquaculture*. 472, 156-177. <https://doi.org/10.1016/j.aquaculture.2016.05.042>.
- Mordenti, O., Emmanuele, P., Casalini, A., Lokman, P.M., Zaccaroni, A., Di Biase, A., Parmeggiani, A., 2018. Effect of aromatizable androgen (17-methyltestosterone) on induced maturation of silver European eels (*Anguilla anguilla*): Oocyte performance and synchronization. *Aquaculture Research*. 49, 442-448. <https://doi.org/10.1111/are.13475>
- Müller, T., Urbányi, B., Váradi, B., Binder, T., Horn, P., Bercsényi, M., Horváth, Á., 2004. Cryopreservation of sperm of farmed European eel *Anguilla anguilla*. *Journal of the World Aquaculture Society*. 35, 225-231. <https://doi.org/10.1111/j.1749-7345.2004.tb01078.x>
- Müller, T., Horváth, Á., Takahashi, E., Kolics, B., Bakos, K., Decsi, K., Kovács, B., Taller, J., Urbányi, B., Bercsényi, M., 2012. Artificial hybridization of Japanese and European eel (*Anguilla japonica* × *A. anguilla*) by using cryopreserved sperm from freshwater reared males. *Aquaculture*. 350, 130-133. <https://doi.org/10.1016/j.aquaculture.2012.04.007>.
- Müller, T., Matsubara, H., Kubara, Y., Horváth, Á., Asturiano, J.F., Urbányi, B., 2017. Japanese eel (*Anguilla japonica* Temminck & Schlegel, 1846) propagation using cryopreserved sperm samples. *Journal of Applied Ichthyology*. 33, 550-552. <https://doi.org/10.1111/jai.13316>
- Müller, T., Matsubara, H., Kubara, Y., Horváth, Á., Kolics, B., Taller, J., Stéger, V., Kovács, B., Horváth, L., Asturiano, J.F., Peñaranda, D.S., Urbányi, B., 2018. Testing

- cryopreserved European eel sperm for hybridization (*A. japonica* × *A. anguilla*).
Theriogenology. 113, 153-158. <https://doi.org/10.1016/j.theriogenology.2018.02.021>.
- Mylonas, C.C., Duncan, N.J., Asturiano, J.F., 2017. Hormonal manipulations for the enhancement of sperm production in cultured fish and evaluation of sperm quality. Aquaculture. 472, 21-44. <https://doi.org/10.1016/j.aquaculture.2016.04.021>.
- Peñaranda, D.S., Pérez, L., Gallego, V., Jover, M., Asturiano, J.F., 2009. Improvement of European eel sperm cryopreservation method by preventing spermatozoa movement activation caused by cryoprotectants. Cryobiology. 59, 119-126.
<https://doi.org/10.1016/j.cryobiol.2009.06.001>.
- Peñaranda, D.S., Pérez, L., Gallego, V., Barrera, R., Jover, M., Asturiano, J.F., 2010. European eel sperm diluent for short-term storage. Reproduction in Domestic Animals. 45, 407-415. <https://doi.org/10.1111/j.1439-0531.2008.01206.x>.
- Peñaranda, D.S., Gallego, V., Rozenfeld, C., Herranz-Jusado, J.G., Pérez, L., Gómez, A., Giménez, I., Asturiano, J.F., 2018. Using specific recombinant gonadotropins to induce spermatogenesis and spermiation in the European eel (*Anguilla anguilla*).
Theriogenology. 107, 6-20. <https://doi.org/10.1016/j.theriogenology.2017.11.002>.
- Pérez-Cerezales, S., Martínez-Páramo, S., Beirão, J., Herráez, M.P., 2010. Evaluation of DNA damage as a quality marker for rainbow trout sperm cryopreservation and use of LDL as cryoprotectant. Theriogenology. 74, 282-289.
<https://doi.org/10.1016/j.theriogenology.2010.02.012>.
- Pérez, L., Asturiano, J.F., Tomás, A., Zegrari, S., Barrera, R., Espinós, F.J., Navarro, J.C., Jover, M., 2000. Induction of maturation and spermiation in the male European eel: assessment of sperm quality throughout treatment. Journal of Fish Biology. 57, 1488-1504. <https://doi.org/10.1111/j.1095-8649.2000.tb02227.x>
- R_Development_Core_Team, 2010. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.

Riesco, M.F., Robles, V., 2013. Cryopreservation causes genetic and epigenetic changes in zebrafish genital ridges. *PLoS One*. 8, e67614.

<https://doi.org/10.1371/journal.pone.0067614>

Rosenthal, H., Asturiano, J.F., Linhart, O., Horváth, A., 2010. On the biology of fish gametes: summary and recommendations of the Second International Workshop, Valencia, Spain, 2009. *Journal of Applied Ichthyology*. 26, 621-622.

Schaefer, C.B., Ooi, S.K.T., Bestor, T.H., Bourc'his, D., 2007. Epigenetic decisions in mammalian germ cells. *Science*. 316, 398-399.

<https://doi.org/10.1126/science.1137544>

Szabó, G., Müller, T., Bercsényi, M., Urbányi, B., Kucska, B., Horváth, A., 2005.

Cryopreservation of European eel (*Anguilla anguilla*) sperm using different extenders and cryoprotectants. *Acta Biologica Hungarica*. 56, 173-175.

<https://doi.org/10.1556/ABiol.56.2005.1-2.18>

Vílchez, M.C., Morini, M., Peñaranda, D.S., Gallego, V., Asturiano, J.F., Pérez, L., 2017.

Role of potassium and pH on the initiation of sperm motility in the European eel.

Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology.

203, 210-219. <https://doi.org/10.1016/j.cbpa.2016.09.024>.

Figure captions

Figure 1.

Sperm kinetic results from fresh sperm, thawed sperm from methanol cryopreservation protocol (Hungarian protocol) and thawed sperm from DMSO protocol (Spanish protocol). The motility analyses show MOT (motility) and PM (progressive motility). The velocity results presented here are VCL (curvilinear velocity), VSL (straight-line velocity) and VAP (average path velocity). Boxplots with different letters are significantly different ($p < 0.05$; $n = 16-18$).

Figure 2.

Comparison of the percentage of different velocity groups [slow (VAP = 10-50 $\mu\text{m/s}$), medium (VAP = 50-100 $\mu\text{m/s}$) and fast (VAP > 100 $\mu\text{m/s}$)] of thawed sperm samples from the DMSO and methanol protocols, and from fresh sperm. Different letters indicate significant differences between percentages of fast cells ($p < 0.05$; $n = 16-18$).

Figure 3.

Effect of freezing medium dilution on sperm motility. Percentage of motile cells after activation with artificial sea water. "Fresh" column shows motility from fresh samples. "Pre-cryo" columns shows the sperm motility of sea water-activated samples after being diluted with freezing medium containing DMSO or methanol before cryopreservation, and "Thawed" columns shows the sperm motility of thawed and sea water-activated samples from the DMSO or methanol protocol. Values are means \pm SEM of sperm from 16 samples. Means with different letters are significantly different ($p < 0.05$).

Figure 4.

Comparative viability data from flow cytometry of fresh sperm and thawed sperm from methanol and DMSO cryopreservation protocols. Values represent means \pm SEM ($n = 12$). Different letters indicate significant differences ($p < 0.05$) between means.

Figure 5

Global DNA methylation of eel sperm. Average percentage \pm SD (n=9) of 5-methylcytosine on fresh and thawed samples. Different letters indicate significant differences ($p < 0.05$).




Figure 1.




Figure 2.




Figure 3.




Figure 4.



Figure 5.

Highlights

- Two previously validated protocols for the cryopreservation of European eel were compared.
- This work represent an important step in standardization of the sperm cryopreservation protocol of a farmed and endangered species such as the European eel.
- The study have shown effects on sperm motility and epigenetic changes caused by cryopreservation in eel thawed sperm.