



HAL
open science

A_∞ weights and compactness of conformal metrics under $L^{n/2}$ curvature bounds

Clara L. Aldana, Gilles Carron, Samuel Tapie

► To cite this version:

Clara L. Aldana, Gilles Carron, Samuel Tapie. A_∞ weights and compactness of conformal metrics under $L^{n/2}$ curvature bounds. *Analysis & PDE*, 2021, 14 (7), pp.2163-2205. 10.2140/apde.2021.14.2163 . hal-01893599

HAL Id: hal-01893599

<https://hal.science/hal-01893599>

Submitted on 11 Oct 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A_∞ WEIGHTS AND COMPACTNESS OF CONFORMAL METRICS UNDER $L^{n/2}$ CURVATURE BOUNDS

CLARA L. ALDANA
UNIVERSITY OF LUXEMBOURG, GILLES CARRON
UNIVERSITÉ DE NANTES, AND SAMUEL TAPIE
UNIVERSITÉ DE NANTES

ABSTRACT. We study sequences of conformal deformations of a smooth closed Riemannian manifold of dimension n , assuming uniform volume bounds and $L^{n/2}$ bounds on their scalar curvatures. Singularities may appear in the limit. Nevertheless, we show that under such bounds the underlying metric spaces are pre-compact in the Gromov-Hausdorff topology. Our study is based on the use of A_∞ -weights from harmonic analysis, and provides geometric controls on the limit spaces thus obtained. Our techniques also show that any conformal deformation of the Euclidean metric on \mathbb{R}^n with infinite volume and finite $L^{n/2}$ norm of the scalar curvature satisfies the Euclidean isoperimetric inequality.

1. INTRODUCTION

This paper is devoted to the question of sequential compactness of Riemannian metrics inside a conformal class given uniform bounds on their volume and integral bounds on their curvatures. It was shown by Gursky in [23] that (unless g_0 is conformal to the round sphere), if the Riemann curvature tensor is bounded in L^p for some $p > \frac{n}{2}$ and if the volume is uniformly controlled, then the sequence of metric is pre-compact in the C^α topology for some $\alpha \in (0, 1)$.

It was later shown that no such compactness can hold when the curvature tensor is bounded in $L^{n/2}$. Indeed, in [16] Chang, Gursky and Wolff constructed two examples of families of sequences of conformal metrics with uniform volume and $L^{n/2}$ curvature bounds which have no convergent sub-sequences. Let us briefly recall these examples.

Example 1: spherical blowup. Let (\mathbb{S}^n, g_S) be the standard sphere, $N \in \mathbb{S}^n$ be some fixed point and S be its antipodal point. Let $\sigma : \mathbb{S}^n \setminus \{N\} \rightarrow \mathbb{R}^n$ be the stereographic projection with pole N , and let eucl denote the Euclidean metric on \mathbb{R}^n . For all $\lambda > 0$, let $g_\lambda = \sigma^*(\lambda \text{eucl})$: it is well known that g_λ extends to a smooth metric on \mathbb{S}^n , conformally equivalent to g_S .

Note that since g_λ is merely a pullback of g_S by some conformal diffeomorphisms, the volume of $(\mathbb{S}^n, g_\lambda)$ and all L^p norms of the curvature tensor are preserved. Moreover, as $\lambda \rightarrow +\infty$, the Riemannian measures $d\mu_{g_\lambda}$ converge to a Dirac mass at S .

It should also be noticed that for any $R > 0$, the g_S -ball $\mathbb{B}(S, R)$ of radius R contains asymptotically all the curvature of g_λ . In particular, we have

$$\lim_{\lambda \rightarrow +\infty} \left(\int_{\mathbb{B}(S, R)} (\text{Scal}_{g_\lambda})^{\frac{n}{2}} d\mu_{g_\lambda} \right)^{\frac{2}{n}} = \alpha(n, 2),$$

where $\alpha(n, 2) = \sigma_n^{\frac{2}{n}} n(n-1)$ is the $L^{n/2}$ -total scalar curvature of the standard round unit sphere, and $\sigma_n = \frac{2\pi^{n/2}}{\Gamma(n/2)}$ is its n -th volume.

It was shown in [16] that this family of conformal deformations of the sphere can be glued on any compact Riemannian manifold (M, g_0) at any point x_0 , giving rise to a sequence of conformal deformations $(g_\lambda = e^{2f_\lambda} g_0)_{\lambda>0}$ such that:

- the volume and diameter of the complement of any open ball centered at x_0 go to 0 as $\lambda \rightarrow +\infty$;
- the $L^{n/2}$ norm of the Riemann curvature tensor are uniformly bounded;
- the sequence of Riemannian measures $d\mu_{g_\lambda}$ develop a Dirac mass at x_0 and the volume stays in some fixed interval $[v, V]$.

In particular, this implies that the associated sequence of distances $(d_{g_\lambda})_{\lambda>0}$ is not precompact for the uniform C^0 topology. Once again, in this example, for any fixed $R > 0$, the g_0 -ball $\mathbb{B}(x_0, R)$ of radius R contains asymptotically a large part of the scalar curvature of g_λ . In particular it can be shown that, in this case, we have

$$\liminf_{\lambda \rightarrow +\infty} \left(\int_{\mathbb{B}(x_0, R)} (\text{Scal}_{g_\lambda}^+)^{\frac{2}{n}} d\mu_{g_\lambda} \right)^{\frac{n}{2}} \geq \alpha(n, 2).$$

We will see below that if we consider sequences of conformal measures whose volume and $L^{n/2}$ -total scalar curvature are uniformly bounded, and the scalar curvature concentration (in $L^{n/2}$ norm) in a g_0 -ball is *strictly less* than $\alpha(n, 2)$, then this kind of spherical blowup cannot occur.

Example 2: Schrödinger type blowup. The second construction in [16] exploits the Schrödinger-type structure of the conformal Laplacian, where the potential is only in $L^{n/2}$. We present a simple example inspired by their construction.

Let (M, g_0) be a compact smooth Riemannian manifold, let $x_0 \in M$ and R_0 be such that $\text{inj}_{x_0}(M, g_0) \geq 2R_0$, where $\text{inj}_{x_0}(M, g_0)$ denotes the injectivity radius of g_0 at x_0 . For $x \in M$, define

$$(1) \quad f_\infty(x) = \begin{cases} \sqrt{\left| \ln \left(\frac{d(x_0, x)}{R_0} \right) \right|} & \text{if } 0 < d(x_0, x) \leq R_0 \\ 0 & \text{if } d(x_0, x) \geq 2R_0 \end{cases}$$

with uniformly bounded second derivatives when $R_0 \leq d(x_0, x) \leq 2R_0$. Let $(f_k : M \rightarrow \mathbb{R})_{k \in \mathbb{N}}$ be a sequence of smooth maps which converges to f_∞ in $W^{2, \frac{n}{2}}$. We consider for all $k \geq 0$ the conformal metric $g_k = e^{2f_k} g_0$. The volume of (M, g_k) is uniformly bounded, and it follows from Theorem 1.159 in [4, p 58] that the $L^{\frac{n}{2}}$ norm of the Riemannian curvature of (M, g_k) are uniformly bounded. Note that eventhough the sequence of Riemannian metrics (g_k) will blow up at x_0 , the diameter of (M, g_k) remains bounded and the associated volume measures will not develop a Dirac mass. Using the sequence of metrics $(\tilde{g}_k = e^{-2f_k} g_0)_{k \in \mathbb{N}}$, we get an analogous example where the metric vanishes at x_0 as $k \rightarrow +\infty$.

Analogously, one can construct examples of sequences of conformal deformations whose volumes and $L^{\frac{n}{2}}$ norms of the scalar curvatures are uniformly bounded, for which the limit metric blows up (or goes to 0) at any countable (even dense) set of points.

The reader may easily show that in this example, eventhough the sequence of conformal factors is unbounded, the sequence of associated distances $(d_{g_k})_{k \in \mathbb{N}}$ converges for the C^0 -uniform topology to the Riemannian distance associated to the (singular) metric $g_\infty = e^{2f_\infty} g_0$ on M .

Our first theorem is a precompactness result which follows from an elementary application of Sobolev embeddings.

Theorem 1.1. *Let (M, g_0) be a compact smooth Riemannian manifold and let \mathcal{M}_V^δ be the space of conformal smooth metrics on M , $g_f = e^{2f} g_0$ with associated Riemannian measure $d\mu_f = e^{nf} d\text{vol}_{g_0}$, such that there exists $V > 0$, $\delta > 0$ such that:*

- (1) $\mu_f(M) \leq V$, and
- (2) $\|(\text{Scal}_{g_f})_+\|_{L^{\frac{n}{2}}(M, d\mu_f)} \leq \alpha(n, 2) - \delta$.

Then for some $\alpha \in (0, 1)$, the set of distances $\{d_{g_f} ; g_f \in \mathcal{M}_V^\delta\}$ is precompact for the C^α topology on $M \times M$. In particular, the set of metric spaces $\{(M, d_{g_f}), g_f \in \mathcal{M}_V^\delta\}$ is precompact for the Gromov-Hausdorff topology.

It follows from the Example 1 presented above that the upper bound

$$\|(\text{Scal}_{g_f})_+\|_{L^{\frac{n}{2}}(M, d\mu_f)} \leq \alpha(n, 2) - \delta = \sigma_n^{\frac{2}{n}} n(n-1) - \delta$$

is optimal to get C^α precompactness of the sequence of distances.

Theorem 1.1 does not exclude the possible collapsing of some part of the manifold in the limit of such conformal deformations. We will show in Corollary 2.4 that adding to the previous hypotheses a lower bound on the volume and a L^p bound on the *scalar* curvature for some $p > \frac{n}{2}$ is enough to recover C^α precompactness of the sequence of Riemannian metrics as in [23].

The rest of our paper is devoted to the exploration of conformal deformations for which only volume bounds and $L^{\frac{n}{2}}$ bounds on the scalar curvature are imposed. We will see in particular that, no collapsing can occur if the $L^{n/2}$ norm of the scalar curvature is small at small scales.

Our ideas come from a paper by Yi Wang [42], who was interested in conformal deformations with integral bounds on the Q -curvature related to the so-called Muckenhoupt weights, or A_∞ -weights. If (M, g) is a closed manifold, a non-negative L^1_{loc} function w is called an A_∞ -weight with respect to g if there exist $C > 0$, $q > 1$ such that for all g -geodesic balls $\mathbb{B} \subset M$,

$$\left(\int_{\mathbb{B}} w^q d\mu_g \right)^{\frac{1}{q}} \leq C \int_{\mathbb{B}} w d\mu_g,$$

where μ_g is the volume measure associated to g . Several equivalent characterizations of these A_∞ -weights will be presented in Section 3.

Theorem 1.2 below establishes that, under $L^{n/2}$ pinching conditions on the scalar curvature closely related to those in Theorem 1.1, the volume densities e^{nf} are *uniformly* A_∞ weights with respect to g_0 .

Theorem 1.2. *Let (M^n, g_0) be a closed Riemannian manifold, $R_0 \in (0, \text{diam}(M, g_0)]$ and $\delta, \Lambda > 0$. Let $f : M \rightarrow \mathbb{R}$ be a smooth function such that the metric $g_f = e^{2f} g_0$ and its associated Riemannian measure $d\mu_f = e^{nf} d\text{vol}_{g_0}$ satisfy*

$$i) \int_M |\text{Scal}_{g_f}|^{\frac{n}{2}} d\mu_f \leq \Lambda.$$

$$\text{ii) For any } x \in M, \left(\int_{\mathbb{B}(x, R_0)} (\text{Scal}_{g_f})_+^{\frac{n}{2}} d\mu_f \right)^{\frac{2}{n}} \leq \alpha(n, 2) - \delta.$$

Then e^{nf} is an A_∞ -weight with respect to g_0 , with constants only depending on n, g_0, R_0, δ and Λ .

The hypotheses as well as the conclusion of Theorem 1.2 are scale invariant. We will see in Section 3 that if we fix upper and lower volume bounds, having A_∞ control for e^{nf} implies C^α bounds for the distance associated to the metric $e^{2f}g$, for some $\alpha > 1$. This implies as well uniform control on certain analytic quantities associated to the conformal metric. Denoting $g_f = e^{2f}g_0$, the fact that e^{nf} is an A_∞ weight implies that for any g_0 -geodesic ball, its g_f -diameter is bounded from above in terms of its g_f -volume and that the measure $d\mu_f = e^{nf}d\mu_{g_0}$ is doubling, with constants only depending on δ, Λ and R_0 .

Notice that in the second family of examples presented above, for which Theorem 1.2 applies, the Riemannian volumes of the members of the family satisfy a *uniform Strong A_∞ bound* in the sense of [17], see Definition 4.1 in Section 4 below. This implies that for a sequence of metrics coming from these examples, the Gromov-Hausdorff limit (M_∞, d_∞) is homeomorphic to M and endows M with a distance d_∞ that is uniformly bi-Hölder to (M, d_{g_0}) . In addition, the metrics in this family of examples have uniform Sobolev, Poincaré and isoperimetric inequalities. Our main result shows that this is always the case, provided the $L^{n/2}$ -pinching of the scalar curvature is small enough at small scales.

Theorem 1.3. *Let (M^n, g_0) be a closed Riemannian manifold. Then there exists a constant $\Lambda_0 = \Lambda_0(g_0)$ with the following property: Let g_f be given by $e^{2f}g_0$ with $f \in C^2(M)$. If there is a $R_0 \in (0, \text{diam}(M, g_0)]$ such that*

$$(2) \quad \forall x \in M, \int_{\mathbb{B}(x, R_0)} |\text{Scal}_{g_f}|^{\frac{n}{2}} d\mu_f \leq \Lambda_0$$

then e^{nf} is a strong A_∞ weight with respect to g_0 , with constants only depending on g_0, R_0 and Λ_0 . Moreover, for all $\alpha \in (0, 1)$, the distances d_f and d_0 are uniformly α -bi-Hölder with constants depending only on g_0, R_0, Λ_0 and α .

Before we continue, let us introduce some notation. Let (M^n, g_0) be a closed Riemannian manifold, for $v, V, \delta, \Lambda > 0$ and $R_0 \in (0, \text{diam}(M, g_0)]$, define $\mathcal{M}_{v, V, R_0}^\Lambda$ as the set of smooth functions f on M such that the associated metric $g_f = e^{2f}g_0$ satisfies

- i) $v \leq \text{Vol}(M, g_f) \leq V$
- ii) for any $x \in M$: $\left(\int_{\mathbb{B}(x, R_0)} |\text{Scal}_{g_f}|^{\frac{n}{2}} d\mu_f \right)^{\frac{2}{n}} \leq \Lambda$.

As mentioned previously, strong A_∞ -weight control coupled with volume bounds has many interesting geometric consequences. Some of them are listed in the following corollary.

Corollary 1.4. *Let (M^n, g_0) be a closed Riemannian manifold. Then there exists $\Lambda_0 = \Lambda_0(g_0)$ such that for all $R_0 \in (0, \text{diam}(M, g_0)]$ and any v, V with $0 < v \leq V$, the set of metric spaces*

$$\{(M, d_{g_f}) ; f \in \mathcal{M}_{v, V, R_0}^{\Lambda_0}\}$$

is precompact for the Gromov-Hausdorff topology. In addition, any sequence in this set has a subsequence whose limit (M, d_∞) is α -bi-Hölder to (M, d_{g_0}) for all $\alpha \in (0, 1)$.

It is worth noticing that the conclusion of Theorem 1.2 is weaker than the conclusion of Corollary 1.4 but holds under an explicit (and optimal, due to examples of [16]) $L^{n/2}$ -pinching of the scalar curvature. The constant $\Lambda(g_0, R_0)$ can be explicitated in terms of the Ricci curvature of g_0 and its diameter. We could not produce a sequence of examples satisfying the hypotheses of Theorem 1.2 which would not be uniformly Strong A_∞ .

Remark 1.5. *Our proof of Theorem 1.3 shows that, under the same hypotheses, e^{nf} is actually a stronger A_∞ weight in the sense of [33]. This implies by Theorem 5.2 of [33] that there exists $N > 0$ such that any such $(M, g_f = e^{2f}g_0)$ can be embedded in \mathbb{R}^N via a bi-Lipschitz embedding, with uniform bi-Lipschitz constant. We will not further discuss this fact here.*

Eventhough our initial objective was to study conformal deformations of closed manifolds, our techniques also provide an interesting criterion to get strong A_∞ on $(\mathbb{R}^n, \text{eucl})$.

Theorem 1.6. *Let $g = e^{2f} \text{eucl}$ be a conformal deformation of the Euclidean metric on \mathbb{R}^n such that:*

- $\text{Vol}(\mathbb{R}^n, g) = +\infty,$
- $\int_{\mathbb{R}^n} |\text{Scal}_g|^{n/2} d\mu_g < +\infty.$

Then e^{nf} is a strong A_∞ weight on \mathbb{R}^n with respect to eucl .

As mentionned above, being a strong A_∞ weight has many implications in terms of geometric control. In particular, this theorem has the following striking geometric corollary, which seems to have remain unknown.

Corollary 1.7. *Let $g = e^{2f} \text{eucl}$ be a conformal deformation of the Euclidean metric on \mathbb{R}^n such that:*

- $\text{Vol}(\mathbb{R}^n, g) = +\infty,$
- $\int_{\mathbb{R}^n} |\text{Scal}_g|^{n/2} d\mu_g < +\infty.$

Then there are positive constants θ, γ such that any g -geodesic ball $B_g(x, r)$ satisfies

$$\theta^{-1} r^n \leq \mu_g(B_g(x, r)) \leq \theta r^n.$$

Moreover, (\mathbb{R}^n, g) satisfies the Euclidean isoperimetric inequality: for any bounded smooth open subset $\Omega \subset \mathbb{R}^n$ we have:

$$\gamma \mu_g(\Omega)^{1-\frac{1}{n}} \leq \mu_g(\partial\Omega).$$

Let us conclude by an intricate question. Let $(g_k = e^{2f_k}g_0)_{k \in \mathbb{N}}$ be a sequence of conformal deformations of g_0 with $f_k \in \mathcal{M}_{v, V, R_0}^{\Lambda_0}$, where Λ_0 satisfies the conclusion of Corollary 1.4. Then, up to extracting a subsequence, the sequence of associated metric spaces (M, d_{g_k}) converges in Gromov-Hausdorff topology to some distance d_∞ on M , which is bi-Hölder equivalent to d_0 . Moreover, a subsequence of the Riemannian volume measures $\mu_k = d\mu_{g_k} = e^{nf_k} d\mu_{g_0}$ converges in the weak- $*$ topology to a positive measure μ_∞ on M . Since all the μ_k are uniformly strong A_∞ -weights with respects to $d\text{vol}_{g_0}$, so is μ_∞ : there exists $e^{nf_\infty} \in L^1(M)$ such that $\mu_\infty = e^{nf_\infty} d\text{vol}_{g_0}$ and μ_∞ is still a strong A_∞ -weight with respect to $d\text{vol}_{g_0}$. Hence the weight e^{nf_∞} induces another distance d_{f_∞} on M , analogous to a conformal Riemannian distance (see Section 7 below or [33] for a precise definition).

A natural question is: *Under which assumption do the distances d_∞ and d_{f_∞} coincide?* We could not give a full answer to this question. We know that they do coincide in Example 2 presented above. On the other hand, in the last section of this paper we will describe another example which shows that uniform strong A_∞ -weight control is not enough to establish this equality in general. Nevertheless, volume bounds and a $L^{n/2}$ bound on the scalar curvature give a much stronger control on the geometry than just strong A_∞ -weight controls. Thus, the distances could still coincide under the integral pinching of the scalar curvature under consideration.

From now on, given a background Riemannian manifold (M, g_0) , for any conformal deformation $g_f = e^{2f} g_0$, we will denote by $d_f = d_{g_f}$ the associated Riemannian distance and by $d\mu_f = \text{vol}_{g_f}$ the associated Riemannian volume measure. In particular, d_0 and μ_0 are the distances and volume measure associated to g_0 , respectively.

Acknowledgements: The authors are grateful for the support of the ANR grants ACG (ANR-10-BLAN 0105) and CCEM (ANR-17-CE40-0034) which made this collaboration possible. C.L. Aldana was also supported by the Fonds National de la Recherche, Luxembourg 7926179.

2. INTEGRAL BOUNDS FOR THE SCALAR CURVATURE AND CONVERGENCE OF DISTANCES

This first section is devoted to the proof of Theorem 1.1. This theorem is an immediate consequence of Propositions 2.2 and 2.3 given below. The proof will show that for any sequence of metrics $(g_n)_{n \in \mathbb{N}}$ in \mathcal{M}_V^δ , the sequence of distances $(d_{g_n})_{n \in \mathbb{N}}$ converges in some Hölder topology up to extracting a subsequence. This implies the precompactness of the sequence of metric spaces $(M, d_{g_n})_{n \in \mathbb{N}}$ for the Gromov-Hausdorff topology, as shown by the following lemma which will be used several times in the sequel.

Lemma 2.1. *Let (X, d_0) be a compact metric space, $\alpha \in (0, 1)$, $C > 0$, and $(d_n)_{n \in \mathbb{N}}$ be a sequence of distances on M which is bounded in the α -Hölder topology with respect to d_0 , i.e. such that for all $x, x', y, y' \in M$ and all $n \in \mathbb{N}$,*

$$|d_n(x, y) - d_n(x', y')| \leq C (d_0(x, x')^\alpha + d_0(y, y')^\alpha).$$

Then up to extracting a subsequence, d_n is converging uniformly and in the α' -Hölder topology for all $\alpha' \in (0, \alpha)$ to a map $d_\infty : M \times M \rightarrow [0, +\infty)$. Moreover, the sequence of metric spaces (M, d_n) converges in the Gromov-Hausdorff topology to the metric space (M_∞, d_∞) where $M_\infty = M / \sim$, with $x \sim y$ if and only if $d_\infty(x, y) = 0$.

Proof. Let $(d_n)_{n \in \mathbb{N}}$ be a sequence of distances on M satisfying the above hypotheses. By Arzela-Ascoli Theorem, it has a subsequence which converges in the \mathcal{C}^0 -uniform topology on $M \times M$ to some non-negative map $d_\infty : M \times M \rightarrow [0, +\infty)$, satisfying the triangular inequality. By construction, d_∞ induces a distance on $M_\infty = M / \sim$, with $x \sim y$ if and only if $d_\infty(x, y) = 0$. The \mathcal{C}^0 -uniform convergence of the d_k implies that the sequence of metric spaces (M, d_n) converges in the Gromov-Hausdorff topology to the metric space (M_∞, d_∞) , see for instance [9] p.260, Example 7.4.4.

An elementary computation shows that the convergence of (d_n) is actually uniform in all $\mathcal{C}^{\alpha'}$ Hölder spaces for all $0 < \alpha' < \alpha$, which concludes the proof of our lemma. \square

Proposition 2.2. *Let (M, g_0) be a compact Riemannian manifold and let $(g_k = e^{2f_k} g_0)_{k \geq 0}$ be a sequence of smooth conformal metrics. Assume that there exist $p > n$ and $A > 0$*

such that for all $k \in \mathbb{N}$,

$$\int_M e^{pf_k} d\mu_0 \leq A.$$

Then the sequence of distances $(d_{f_k} : M \times M \rightarrow [0, +\infty))_{k \in \mathbb{N}}$ is precompact in the α -Hölder topology for all $\alpha \in (0, 1 - \frac{n}{p})$. In particular, the sequence of metric spaces $(M, d_{f_k})_{k \in \mathbb{N}}$ is precompact for the Gromov-Hausdorff topology.

Proof. Let $x \in M$ be fixed. Recall that for any smooth Riemannian metric g on M , $\|\nabla^g d_g(x, \cdot)\|_g = 1$ almost everywhere, see e.g. [31, Prop. 4.8]. Let $g_f = e^{2f} g_0$ be conformal metric, we have hence for almost all $y \neq x$,

$$\|\nabla^{g_0} d_f(x, \cdot)\|_{g_0}(y) = e^{f(y)}.$$

Therefore, there exists $C > 0$ such that

$$(3) \quad \int_M e^{pf} d\mu_0 \leq A \quad \Rightarrow \quad \|d_f(x, \cdot)\|_{W^{1,p}(M, g_0)} \leq CA,$$

where $W^{1,p}(M, g_0)$ is the Sobolev space of functions on M with distributional derivative in $L^p(g_0)$.

Now, let $(g_k = e^{2f_k} g_0)_{k \geq 0}$ be a sequence of smooth conformal metrics. Assume that there exist $p > 1$ and $A > 0$ such that for all $k \in \mathbb{N}$,

$$\int_M e^{pf_k} d\mu_0 \leq A.$$

It follows from (3) that the sequence of distances $(d_{f_k}(x, \cdot))_{k \in \mathbb{N}}$ is bounded in $W^{1,p}(M, g_0)$. Therefore, since for all $k \in \mathbb{N}$, $d_{f_k}(x, \cdot)$ vanishes at x , by Sobolev embeddings the sequence is bounded in C^α with $\alpha = 1 - \frac{n}{p}$ by some constant C' independent of x .

We have hence, for all $x, x', y, y' \in M$ and all $n \in \mathbb{N}$,

$$\begin{aligned} |d_{f_k}(x, y) - d_{f_k}(x', y')| &= |d_{f_k}(x, y) - d_{f_k}(x', y) + d_{f_k}(x', y) - d_{f_k}(x', y')| \\ &\leq C' (d_0(x, x')^\alpha + d_0(y, y')^\alpha). \end{aligned}$$

By Lemma 2.1, this implies Proposition 2.2. □

We now show that any conformal deformation of g_0 whose scalar curvature has a positive part with $L^{n/2}$ norm strictly less than the round sphere has a conformal factor that satisfies the integrability condition required in Proposition 2.2.

Proposition 2.3. *Let (M^n, g_0) be a closed Riemannian manifold. Let V, δ be positive real numbers. There exists $A = A(V, \delta) > 0$ such that for all $\varepsilon > 0$ small enough, if*

- i) $\text{Vol}(M, g_f) \leq V$,
- ii) $\|(\text{Scal}_{g_f})_+\|_{L^{\frac{n}{2}}(M, d\mu_f)} \leq \alpha(n, 2) - \delta$

then

$$(4) \quad \int_M e^{n(1+\varepsilon)f} d\mu_{g_0} \leq A.$$

Proof. We are going to use the following optimal Sobolev inequality due to E. Hebey and M. Vaugon ([24]) (this inequality had been conjectured by T. Aubin [2]): if we write again $\alpha(n, 2) = \sigma_n^{\frac{2}{n}} n(n-1)$, then there is a constant B such that for all $\varphi \in C^\infty(M)$,

$$(5) \quad \alpha(n, 2) \left(\int_M \varphi^{\frac{2n}{n-2}} d\mu_0 \right)^{1-\frac{2}{n}} \leq \frac{4(n-1)}{n-2} \int_M |d\varphi|_{g_0}^2 d\mu_{g_0} + \int_M B\varphi^2 dv_{g_0}.$$

If $g_f = u^{\frac{4}{n-2}} g_0$ if follows from [43] that

$$(6) \quad \text{Scal}_{g_f} u^{\frac{4}{n-2}} u = \frac{4(n-1)}{n-2} \Delta_{g_0} u + \text{Scal}_{g_0} u.$$

In particular for any $\epsilon > 0$ we get

$$\begin{aligned} \frac{4(n-1)}{n-2} \Delta_{g_0} u^{1+\epsilon} + (1+\epsilon) \text{Scal}_{g_0} u^{1+\epsilon} + (n-1) \frac{4\epsilon(1+\epsilon)}{n-2} u^{\epsilon-1} |du|_{g_0}^2 \\ = (1+\epsilon) \text{Scal}_{g_f} u^{\frac{4}{n-2}} u^{(1+\epsilon)}. \end{aligned}$$

Using the Hebey and Vaugon Sobolev inequality for $\varphi = u^{(1+\epsilon)}$, we get

$$\begin{aligned} \alpha(n, 2) \left(\int_M u^{(1+\epsilon)\frac{2n}{n-2}} d\mu_{g_0} \right)^{1-\frac{2}{n}} \\ \leq \frac{4(n-1)}{n-2} \int_M u^{(1+\epsilon)} \Delta_{g_0} u^{1+\epsilon} d\mu_{g_0} + B \int_M u^{2(1+\epsilon)} d\mu_{g_0} \\ \leq (1+\epsilon) \int_M (\text{Scal}_{g_f})_+ u^{\frac{4}{n-2}} u^{2(1+\epsilon)} d\mu_{g_0} \\ + ((1+\epsilon)S_0 + B) \int_M u^{2(1+\epsilon)} d\mu_{g_0}, \end{aligned}$$

where $S_0 = \|\text{Scal}_{g_0}\|_{L^\infty}$. When $\epsilon \leq 2/(n-2)$, using the Hölder inequality we get

$$\begin{aligned} \left((\alpha(n, 2) - (1+\epsilon)) \|(\text{Scal}_{g_f})_+\|_{L^{\frac{n}{2}}(M, d\mu_f)} \right) \left(\int_M u^{(1+\epsilon)\frac{2n}{n-2}} d\mu_{g_0} \right)^{1-\frac{2}{n}} \leq \\ ((1+\epsilon)S_0 + B) (\text{Vol}_{g_0}(M))^{\frac{2-(n-2)\epsilon}{n}} (\text{Vol}_{g_f}(M))^{(1+\epsilon)\frac{n-2}{n}}, \end{aligned}$$

which gives with $e^{2f} = u^{\frac{4}{n-2}}$,

$$\int_M e^{n(1+\epsilon)f} d\mu_{g_0} \leq \frac{((1+\epsilon)S_0 + B) (\text{Vol}_{g_0}(M))^{\frac{2-(n-2)\epsilon}{n}} V^{(1+\epsilon)\frac{n-2}{2}}}{\delta - \epsilon(\alpha(n, 2) - \delta)},$$

which is precisely (4) when ϵ is small enough, say $\epsilon < \frac{\delta}{\alpha(n, 2) - \delta}$. \square

Notice that under our very weak hypotheses, the limit space (M_∞, d_∞) obtained as the Gromov-Hausdorff limit of the sequence $(M, d_{g_k})_{k \in \mathbb{N}}$ may be very singular; the singularities depending on the set where d_∞ vanishes. Even if the volume of (M, g_k) remains constant (recall that the $L^{n/2}$ bound of the scalar curvature is invariant under rescaling), we cannot a priori prevent collapsing of some open part of the manifold. In the sequel, we look for stronger conditions on the scalar curvatures and the volumes so that we get a better geometric picture of the limit.

Let us first note that if we add to the hypotheses in Theorem 1.1 a lower bound on the volume and a uniform L^p bound on the *scalar curvature* (and not the full Riemann tensor) with $p > \frac{n}{2}$, then we get C^α -precompactness for the conformal factors:

Corollary 2.4. *Let (M^n, g_0) be a closed Riemannian manifold. Let v, V, Λ, δ be positive real numbers and let $p > n/2$. Then the set $\mathcal{M}_{v,V}^{p,\Lambda,\delta}$ of smooth functions $f \in C^\infty(M)$ such that $g_f = e^{2f}g_0$ satisfies*

- i) $v \leq \text{Vol}(M, g_f) \leq V$,
 - ii) $\int_M |\text{Scal}_{g_f}|^p d\mu_f \leq \Lambda$
 - iii) $\|(\text{Scal}_{g_f})_+\|_{L^{\frac{n}{2}}(M, d\mu_f)} \leq \sigma_n^{\frac{2}{n}} n(n-2) - \delta$
- is precompact in $C^\alpha(M)$ for $0 < \alpha < \frac{2p-n}{p}$.

Proof. It follows from Proposition 2.3 that for all metrics $g_f = e^{2f}g_0$ in $\mathcal{M}_{v,V}^{p,\Lambda,\delta}$, the volume density e^{nf} satisfies a $L^{1+\varepsilon}$ uniform bound given by (4).

Combining this uniform stronger integrability of e^f together with L^p bounds for the scalar curvature and volume bounds, a Harnack inequality due to Trudinger in [41] together with elliptic regularity will give C^α -precompactness for the space of metrics for $0 < \alpha < \frac{2p-n}{p}$. This is explained with details in Section 2.2 of [29]. \square

In particular, any sequence of metrics satisfying some L^p bound on the scalar curvature, $p > n/2$, and the above $L^{n/2}$ bounds on its positive part, converges (up to extraction of a subsequence) in some C^α -topology to a Riemannian metric $g_\infty = e^{2f_\infty}g_0$, where $f_\infty \in C^\alpha(M)$. This slightly improves the well known result by M. Gursky given in [23].

The rest of our paper is devoted to the study of sequences of conformal deformations with fixed volume bounds and a mere $L^{n/2}$ bound on the scalar curvature. It follows from [16] that in this setting, one cannot get C^0 precompactness for the sequence of conformal factors (or, equivalently, for the sequence of Riemannian metrics). Nevertheless, we will show that the sequence of associated measured metric spaces and the limit metric spaces (M_∞, d_∞) thus obtained satisfy many uniform geometrical properties.

3. OPTIMAL $L^{n/2}$ BOUNDS FOR THE SCALAR CURVATURE AND A_∞ WEIGHTS

In this section, we study sequences $(g_k = e^{2f_k}g_0)_{k \in \mathbb{N}}$ of conformal metrics with a uniform upper bound on the volumes, such that the $L^{n/2}$ -norms of the positive part of the scalar curvature on small spheres is uniformly bounded by the one of the standard sphere, and which moreover have some uniform $L^{n/2}$ bound for the total scalar curvature. We will show that the Riemannian volume densities $\{e^{nf_k}\}_{k \in \mathbb{N}}$ are then *uniformly A_∞ -weights* with respect to g_0 , see Definition 3.1.

To be uniformly A_∞ -weights has several nice geometrical consequences. First of all, the sequence of volume measures $(d\mu_{f_k})_{k \in \mathbb{N}}$ is uniformly doubling with respect to the initial distance d_0 . In particular, this rules out the kind of blowups presented in Example 1 in our introduction. More precisely, we will see that the A_∞ control implies that the Riemannian distances d_{f_k} are uniformly controlled in terms of the volumes $d\mu_{f_k}$, locally. We will show as well Gromov-Hausdorff precompactness for the set of conformal metrics satisfying a uniform A_∞ -weight bound, independently of any control on the curvature.

In the afore mentioned work of Gursky [23], a key step to get C^α compactness for a sequence of conformal metrics $(g_k = e^{2f_k}g_0)_{k \in \mathbb{N}}$ whose Riemann curvatures satisfy a uniform L^p bound ($p > n/2$) is to show that if g_0 is not conformal to the standard sphere, then the conformal factors in such sequences satisfy a uniform Harnack inequality: there exists $C > 1$ such that for all $k \in \mathbb{N}$,

$$\frac{\sup e^{f_k}}{\inf e^{f_k}} \leq C.$$

Uniform volume bounds and (even very small) $L^{n/2}$ bounds on the scalar curvature cannot ensure such Harnack inequality, as shows Example 2 in our introduction. A uniform A_∞ control on the weight e^{nf_k} is actually a local L^q integral version of this Harnack inequality for some $q > 1$.

3.1. A_∞ weights and conformal metrics. We now give a presentation of A_∞ weights on a closed Riemannian manifold. The reader can find a good exposition and more details in [36, chapter V]. Then, we will draw some geometric consequences for conformal metrics having such A_∞ control.

Let (M^n, g_0) be a closed Riemannian manifold. Throughout this document we will denote by

$\mathbb{B}(x, r)$: the g_0 -geodesic ball centered at x and of radius r .

We will sometimes write \mathbb{B} for a g_0 -geodesic ball whose radius will be denoted by $r(\mathbb{B})$. For $\theta > 0$, $\theta\mathbb{B}$ will be the ball with the same center and with radius $\theta r(\mathbb{B})$. D will denote the diameter of (M^n, g_0) . The measure μ_0 will still be the g_0 Riemannian volume. If $E \subset M$ is a measurable set and $f \in L^1(E, d\mu_0)$ the average of f over E will be denoted by

$$\int_E f d\mu_0 = \frac{1}{\mu_0(E)} \int_E f d\mu_0.$$

Definition 3.1. We say that a non negative function $w \in L^1(M, d\mu_0)$ is a A_∞ weight with respect to g_0 if one of the following equivalent properties is satisfied:

- i) There is a $q > 1$ and a constant C such that for any geodesic ball $\mathbb{B} \subset M$, the following reverse Hölder inequality with exponent q holds:

$$\left(\int_{\mathbb{B}} w^q d\mu_0 \right)^{\frac{1}{q}} \leq C \int_{\mathbb{B}} w d\mu_0.$$

- ii) There is $p > 1$ and a constant C such that for any geodesic ball $\mathbb{B} \subset M$:

$$\left(\int_{\mathbb{B}} w d\mu_0 \right) \left(\int_{\mathbb{B}} w^{-\frac{1}{p-1}} d\mu_0 \right)^{p-1} \leq C.$$

If w satisfies this condition, it is called A_p -weight.

- iii) There are constant $\delta, \varepsilon \in (0, 1)$ such that for any geodesic ball $\mathbb{B} \subset M$ and any $E \subset M$ if $E \subset \mathbb{B}$ satisfies $\mu_0(E) \leq \delta \mu_0(\mathbb{B})$ then $\int_E w d\mu_0 \leq \varepsilon \int_{\mathbb{B}} w d\mu_0$.
- iv) There are constants $\alpha > 1$ and $C > 0$ such that for any geodesic ball $\mathbb{B} \subset M$ and any $E \subset \mathbb{B}$:

$$\frac{1}{C} \left(\frac{\mu_0(E)}{\mu_0(\mathbb{B})} \right)^\alpha \leq \frac{\int_E w d\mu_0}{\int_{\mathbb{B}} w d\mu_0} \leq C \left(\frac{\mu_0(E)}{\mu_0(\mathbb{B})} \right)^{\frac{1}{\alpha}}.$$

Remark 3.2. The different constants in the different definitions are mutually controlled. In particular, we will use the following fact: If w is an A_p weight satisfying part ii) above for all $p > 1$, then it fulfills reverse Hölder inequality i) with exponent q for all $q > 1$, with mutually controlled constants. See [36, p.196-203] for a proof.

It is possible to show that the properties in Definition 3.1 hold for all geodesic balls if and only if they hold for geodesic balls of radius less than some fixed $R_0 > 0$. We will prove this statement for the Reverse Hölder inequality. In order to do that, we need the following general result.

Lemma 3.3. *If (X, d, μ) is a metric measure space whose balls B with radius less than R_0 satisfy the doubling condition:*

$$\mu(2B) \leq \theta \mu(B)$$

then for any $R_1 \geq R_0$ there is a constant θ' depending only on θ and R_1/R_0 such that any ball B with radius less than R_1 satisfies the doubling condition:

$$\mu(2B) \leq \theta' \mu(B)$$

Proof. Indeed let $B(x, R)$ a ball of radius $R \in [R_0, R_1]$, then from [11, Lemma 3.10], we have

$$\mu(B(x, 2R)) \leq \theta^{50+50\frac{2R}{R_0}} \mu_f(B(x, R_0)) \leq \theta^{50+50\frac{2R_1}{R_0}} \mu(B(x, R)).$$

□

Lemma 3.4. *Let $q > 1$, and $C, R_0 > 0$ be fixed. Let w be a uniformly (q, R_0, C) - A_∞ weight, i.e. every geodesic ball \mathbb{B} of radius $r \leq R_0$ satisfies:*

$$\left(\int_{\mathbb{B}} w^q d\mu_0 \right)^{\frac{1}{q}} \leq C \int_{\mathbb{B}} w d\mu_0.$$

Then the measure $w d\mu_0$ has the doubling volume property and there exists $C' = C'(q, C, R_0, g_0) > 0$ such that for all balls \mathbb{B} of radius $r \geq 0$, one has

$$(7) \quad \left(\int_{\mathbb{B}} w^q d\mu_0 \right)^{\frac{1}{q}} \leq C' \int_{\mathbb{B}} w d\mu_0.$$

Proof. Assume that w satisfies the hypotheses of the lemma for some $q > 1$ and $C, R_0 > 0$. We first show that $w d\mu_0$ has the doubling volume property. Let us first consider geodesic balls of radius less than $R_0 > 0$. We start by showing that there is $\theta > 0$ such that for the measure $d\omega = w d\mu_0$ we have the doubling property:

$$r(\mathbb{B}) \leq R_0/2 \implies \omega(2\mathbb{B}) \leq \theta \omega(\mathbb{B}).$$

Indeed using both Hölder and Reverse Hölder inequalities we easily get for any $\tau \in (0, 1)$:

$$\frac{\omega(\mathbb{B} \setminus \tau\mathbb{B})}{\omega(\mathbb{B})} \leq C \left(\frac{\mu_0(\mathbb{B} \setminus \tau\mathbb{B})}{\mu_0(\mathbb{B})} \right)^{1-\frac{1}{q}}.$$

The manifold M is compact hence there is a constant B such that for any geodesic ball :

$$\mu_0(\mathbb{B} \setminus \tau\mathbb{B}) \leq B(1-\tau)\mu_0(\mathbb{B}).$$

We choose $\tau \in (0, 1)$ such that $C(B(1-\tau))^{1-\frac{1}{q}} = \frac{1}{2}$ and we get $\omega(\mathbb{B}) \leq 2\omega(\tau\mathbb{B})$, therefore $\omega(\mathbb{B}) \leq 2^N \omega(\tau^N \mathbb{B})$. Choosing now N such that $\tau^N \leq \frac{1}{2} < \tau^{N-1}$ gives the result with $\theta = 2^N$.

As the diameter of (M, g_0) is finite, we deduce from the Lemma 3.3 that the measure ω is doubling: there is a constant θ such that for any ball $\mathbb{B} \subset M$:

$$\omega(2\mathbb{B}) \leq \theta \omega(\mathbb{B}).$$

We can now show that the reverse Hölder inequality holds for any ball. Let \mathbb{B} be a geodesic ball of radius $r \in (R_0, D]$, then we can find a minimal family of geodesic balls $\mathbb{B}_\alpha = \mathbb{B}(x_\alpha, R_0/2)$ of radius $R_0/2$ such that $\mathbb{B} \subset \cup_\alpha \mathbb{B}_\alpha$, and the balls $\mathbb{B}(x_\alpha, R_0/4)$ are

disjoint, hence include in $2\mathbb{B}$. Since w satisfies reverse Hölder on the balls $\mathbb{B}(x_\alpha, R_0/2)$, we get

$$\left(\int_{\mathbb{B}_\alpha} w^q d\mu_0 \right)^{\frac{1}{q}} \leq \frac{C\mu_0(\mathbb{B}_\alpha)^{\frac{1}{q}}}{\mu_0(\mathbb{B}_\alpha)} \int_{\mathbb{B}_\alpha} w d\mu_0 = C\mu_0(\mathbb{B}_\alpha)^{\frac{1}{q}-1} \int_{\mathbb{B}_\alpha} w d\mu_0.$$

From the doubling condition on (M, d_0, μ_0) there is a constant such that

$$\mu_0(\mathbb{B}) \leq C\mu_0(\mathbb{B}_\alpha).$$

Hence

$$\begin{aligned} \left(\int_{\mathbb{B}_\alpha} w^q d\mu_0 \right)^{\frac{1}{q}} &\leq C\mu_0(\mathbb{B})^{\frac{1}{q}-1} \int_{\mathbb{B}_\alpha} w d\mu_0 \\ \int_{\mathbb{B}} w^q d\mu_0 &\leq \sum_{\alpha} \int_{\mathbb{B}_\alpha} w^q d\mu_0 \\ &\leq C \sum_{\alpha} \mu_0(\mathbb{B}_\alpha)^{1-q} \left(\int_{\mathbb{B}_\alpha} w d\mu_0 \right)^q \\ &\leq C\mu_0(\mathbb{B})^{1-q} \sum_{\alpha} \left(\int_{\mathbb{B}_\alpha} w d\mu_0 \right)^q \\ &\leq C\mu_0(\mathbb{B})^{1-q} \theta^q \sum_{\alpha} \left(\omega\left(\frac{1}{2}\mathbb{B}_\alpha\right) \right)^q \\ &\leq C\mu_0(\mathbb{B})^{1-q} \theta^q \left(\sum_{\alpha} \omega\left(\frac{1}{2}\mathbb{B}_\alpha\right) \right)^q \\ &\leq C\mu_0(\mathbb{B})^{1-q} \theta^q \omega(2\mathbb{B})^q \leq C\mu_0(\mathbb{B})^{1-q} \theta^{2q} \omega(\mathbb{B})^q, \end{aligned}$$

from which the conclusion follows. \square

Let now $f \in C^\infty(M)$ and $g_f = e^{2f}g_0$ be a Riemannian metric conformal to g_0 . A key geometric consequence of having A_∞ control on the weight e^{nf} is given in the following lemma.

Lemma 3.5. *Let $f : M \rightarrow \mathbb{R}$ be a smooth map. Assume that $w = e^{nf}$ is an A_∞ weight satisfying the above reverse Hölder inequality with constants $C > 0$ and $q > 1$. Then there is a constant B , depending only on C, q and (M, g_0) , such that for any $x, y \in M$:*

$$d_f(x, y)^n \leq B \int_{\mathbb{B}(x, d_0(x, y))} e^{nf} d\mu_0.$$

This lemma has been shown on the Euclidean space by G. David and S. Semmes, see [17, inequality (1.2)] and [33, Proposition 3.12b].

Proof. By Sobolev inequality (see for instance [20] p.148), there is a constant $C > 0$ depending only on the geometry of (M, g_0) and q , such that for $p = qn$, we have for any balls $\mathbb{B} \subset M$ and any $\varphi \in W^{1,p}(2\mathbb{B})$ and $x, y \in \mathbb{B}$:

$$|\varphi(x) - \varphi(y)| \leq Cr(\mathbb{B})^{1-\frac{n}{p}} \left(\int_{2\mathbb{B}} |d\varphi|^p d\mu_0 \right)^{\frac{1}{p}}.$$

If we use this for the function $\varphi = d_f(x, \cdot)$ and \mathbb{B} a ball centered at some point m with $d_0(x, m) = d_0(m, y) = d_0(x, y)/2$ and with radius $\frac{3}{4}d_0(x, y)$. we will get

$$d_f(x, y) \leq Cr(\mathbb{B})^{1-\frac{n}{p}} \left(\int_{2\mathbb{B}} e^{qn_f} d\mu_0 \right)^{\frac{1}{qn}}$$

Then the conclusion follows from the reverse Hölder inequality and the fact that the μ_0 measure of $2\mathbb{B}$ is comparable to $r(\mathbb{B})^n$. \square

Let us show now that the space of conformal metrics with uniform A_∞ bounds is pre-compact in the Gromov-Hausdorff topology. We will not need this result in the sequel since Theorem 1.1 will already ensure Gromov-Hausdorff precompactness once we fix a suitable $L^{n/2}$ bound on the scalar curvature. Nevertheless, this is an easy consequence of the measure being A_∞ weights which seems to have remained unnoticed.

Proposition 3.6. *Let (M^n, g_0) be a closed Riemannian manifold let $C, V > 0$, $R_0 \in (0, \text{diam}(M, g_0)]$ and $q > 1$.*

Let $\mathcal{M}_{V, R_0, q, C}$ be the set of smooth functions such that $f \in \mathcal{M}_{V, R_0, q, C}$ satisfies the following conditions:

- i) *for the Riemannian metric $g_f = e^{2f}g_0$, $\text{Vol}(M, g_f) \leq V$;*
- ii) *$w = e^{nf}$ is (R_0, q, C) -uniformly A_∞ i.e. for any geodesic ball \mathbb{B} of radius less or equal than R_0 we have:*

$$\left(\int_{\mathbb{B}} w^q d\mu_0 \right)^{\frac{1}{q}} \leq C \int_{\mathbb{B}} w d\mu_0.$$

Then the set of distances $\{d_f, f \in \mathcal{M}_{V, R_0, q, C}\}$ is pre-compact in the C^α topology for all $\alpha \in (0, 1 - \frac{1}{q})$. In particular, the set of metric spaces $\{(M, d_f), f \in \mathcal{M}_{V, R_0, q, C}\}$ is precompact for the Gromov-Hausdorff topology.

Proof. By Lemma 3.4, there exists $C' > 0$ such that for all $f \in \mathcal{M}_{v, V, R_0, q, C}$ and all geodesic balls \mathbb{B} (of any radius),

$$\left(\int_{\mathbb{B}} w^q d\mu_0 \right)^{\frac{1}{q}} \leq C \int_{\mathbb{B}} w d\mu_0.$$

In particular, if the radius of \mathbb{B} is larger than the diameter of (M, g_0) , we get that for all $f \in \mathcal{M}_{v, V, R_0, q, C}$,

$$\int_M e^{nqf} \leq C' (\text{Vol}(M, g_0))^{\frac{1}{q}-1} V.$$

The conclusion follows then immediately from Proposition 2.2. \square

3.2. A_∞ control from $L^{n/2}$ bound on the scalar curvature. We now show that a $L^{n/2}$ bound on the scalar curvature, with $L^{n/2}$ bound on the positive part of the scalar curvature uniformly smaller than what it is for the standard sphere at some (even small) fixed scale, implies a uniform A_∞ control of the conformal factor. Once again, it follows from the Example 1 presented in our introduction that these integral bounds are optimal to get such A_∞ control. Theorem 3.7 below implies Theorem 1.2 of the introduction and explicit its result.

Theorem 3.7. *Let (M^n, g_0) be a closed Riemannian manifold, $\delta, \Lambda > 0$ and $R_0 > 0$. Then there exist $p = p(n, g_0, R_0, \Lambda) > 0$ and $C = C(n, g_0, R_0, \delta, \Lambda) > 0$ such that for any smooth function $f : M \rightarrow \mathbb{R}$ whose associate metric $g_f = e^{2f}g_0$ satisfies*

i) $\int_M |\text{Scal}_{g_f}|^{\frac{n}{2}} d\mu_f \leq \Lambda$, and

ii) for any $x \in M$: $\left(\int_{\mathbb{B}(x, R_0)} (\text{Scal}_{g_f})_+^{\frac{n}{2}} d\mu_f \right)^{\frac{2}{n}} \leq \alpha(n, 2) - \delta$,

we have that for any g_0 -geodesic ball \mathbb{B} of radius less than $R_0/2$,

$$\left(\int_{\mathbb{B}} e^{nf} d\mu_0 \right) \left(\int_{\mathbb{B}} e^{-\frac{nf}{p}} d\mu_0 \right)^p \leq C.$$

In particular, e^{nf} is an A_∞ -weight with respect to g_0 , with constants only depending on n, g_0, R_0, δ and Λ .

Proof. The proof of this theorem was directly inspired by the proof of the Harnack inequality for positive solution of second order elliptic equation, see [20, theorem 8.20] and its proof.

Let $R_0 \in (0, \text{diam}(M, g_0))$, $\delta, \Lambda > 0$ and let $f : M \rightarrow \mathbb{R}$ be a smooth function satisfying the hypotheses above. We write again $g_f = u^{\frac{4}{n-2}} g_0$, i.e. $e^f = u^{\frac{2}{n-2}}$. The proof is done in two steps.

Lemma 3.8. *Under the same hypothesis of Theorem 3.7, there exist $\varepsilon = \varepsilon(n, g_0, R_0, \Lambda) > 0$ and $C = C_1(n, g_0, R_0, \Lambda) > 0$ such that for any g_0 -geodesic ball \mathbb{B} of radius less than $R_0/2$,*

$$\left(\int_{\mathbb{B}} u^\varepsilon d\mu_0 \right) \left(\int_{\mathbb{B}} u^{-\varepsilon} d\mu_0 \right) \leq C.$$

Proof. Since we consider $g_f = u^{\frac{4}{n-2}} g_0$, let us recall equation 6:

$$\text{Scal}_{g_f} u^{\frac{4}{n-2}} u = \frac{4(n-1)}{n-2} \Delta_{g_0} u + \text{Scal}_{g_0} u$$

Set $V = \text{Scal}_{g_f} u^{\frac{4}{n-2}}$, $W = V - \text{Scal}_{g_0}$ and $\lambda_0 = \|\text{Scal}_{g_0}\|_{L^\infty}$. We have

$$\frac{4(n-1)}{n-2} \Delta_0 \ln(u) = W + \frac{4(n-1)}{n-2} |d \ln(u)|_{g_0}^2.$$

Now let \mathbb{B} and ξ be a Lipschitz cut-off function with support in $2\mathbb{B}$ such that $\xi = 1$ on \mathbb{B} and $|d\xi| \leq r(\mathbb{B})^{-1}$. Then

$$\begin{aligned} \int_M \xi^2 \left(W + \frac{4(n-1)}{n-2} |d \ln(u)|_{g_0}^2 \right) d\mu_0 &= \frac{4(n-1)}{n-2} \int_M 2\xi \langle d\xi, d \ln(u) \rangle d\mu_0 \\ &\leq 4 \int_M \xi^2 |d \ln(u)|_{g_0}^2 d\mu_0 + \frac{16(n-1)^2}{(n-2)^2} \int_M |d\xi|^2 d\mu_0 \end{aligned}$$

By Hölder inequality, we have

$$\int_M \xi^2 (-V) d\mu_0 \leq \left\| \text{Scal}_{g_f}^- \right\|_{L^{n/2}(2\mathbb{B})} \mu_0(2\mathbb{B})^{\frac{n-2}{2}},$$

Moreover, for g_0 there is an uniform constant ω such that for any $r \in (0, \text{diam}(M, g_0))$ and any $x \in M$:

$$(8) \quad \frac{1}{\omega} r^n \leq \mu_0(\mathbb{B}(x, r)) \leq \omega r^n.$$

Hence we get

$$\left(\frac{4(n-1)}{n-2} - 4 \right) \int_{\mathbb{B}} |d \ln(u)|_{g_0}^2 d\mu_0 \leq C \lambda_0 r(\mathbb{B})^n + C \left(1 + \left\| \text{Scal}_{g_f}^- \right\|_{L^{\frac{n}{2}}(2\mathbb{B})} \right) \mu_0(2\mathbb{B}) r(\mathbb{B})^{-2}.$$

In particular we have

$$(9) \quad r(\mathbb{B})^2 \int_{\mathbb{B}} |d \ln(u)|_{g_0}^2 d\mu_0 \leq C \left(1 + \left\| \text{Scal}_{g_f}^- \right\|_{L^{\frac{n}{2}}(2\mathbb{B})} \right).$$

There is a $r_0 > 0$ such that all the geodesic balls $\mathbb{B}(x, r_0)$ are almost Euclidean: i.e. there is a smooth map $\varphi_x : \mathbb{B}(0, r_0) \rightarrow \mathbb{B}(x, r_0)$ with

$$\frac{1}{2} \text{eucl} \leq \varphi_x^* g_0 \leq 2 \text{eucl},$$

where eucl is the Euclidean metric and $\mathbb{B}(0, r_0)$ the Euclidean ball of radius r_0 . Hence there are positive constants θ and λ that only depends on n such that if \mathbb{B} is a g_0 geodesic ball of radius $r(\mathbb{B}) \leq r_0/2$ then

$$\mu_0(2\mathbb{B}) \leq \theta \mu_0(\mathbb{B})$$

and

$$\forall \phi \in C^1(\mathbb{B}) : \int_{\mathbb{B}} \left(\phi - \int_{\mathbb{B}} \phi \right)^2 d\mu_0 \leq \lambda r(\mathbb{B})^2 \int_{\mathbb{B}} |d\phi|^2 d\mu_0.$$

We can assume that $R_0 \leq r_0$, hence the above estimation (9) and the above Poincaré inequality implies a BMO (bounded mean oscillation) estimate on the function $\ln(u)$. Moreover, we also get from (9) that for all balls \mathbb{B} with radius $R \leq R_0$,

$$\int_{\mathbb{B}} |d \ln(u)|_{g_0} d\mu_0 \leq CR^{n/2} \left(\int_{\mathbb{B}} |d \ln(u)|_{g_0}^2 d\mu_0 \right)^{1/2} \leq CR^{n-1}.$$

Therefore by John-Nirenberg inequality (cf [20, Theorem 7.21]), there exists $\varepsilon > 0$ such that for any geodesic ball \mathbb{B} of radius less than $R_0/2$,

$$\int_{\mathbb{B}} e^{\varepsilon |\ln u - (\ln u)_{\mathbb{B}}|} d\mu_0 \leq C,$$

where $(\ln u)_{\mathbb{B}} = \int_{\mathbb{B}} \ln(u) d\mu_0$. This implies that

$$\int_{\mathbb{B}} u^{\pm \varepsilon} d\mu_0 \leq C e^{\pm \varepsilon (\ln u)_{\mathbb{B}}}.$$

Therefore

$$\left(\int_{\mathbb{B}} u^{\varepsilon} d\mu_0 \right) \left(\int_{\mathbb{B}} u^{-\varepsilon} d\mu_0 \right) \leq C e^{\varepsilon (\ln u)_{\mathbb{B}}} e^{-\varepsilon (\ln u)_{\mathbb{B}}} \leq C.$$

□

So far we have only used an estimate of the negative part of the scalar curvature of g_f in $L^{n/2}$. However, we will need the $L^{n/2}$ bound on the positive part of the scalar curvature in order to get the following estimate.

Lemma 3.9. *Assume the same hypothesis of Theorem 3.7. For all $\varepsilon > 0$, there exists $C = C(n, g_0, R_0, \delta, \Lambda, \varepsilon) > 0$ such that for all g_0 -geodesic balls of radius less than R_0 ,*

$$\left(\int_{\frac{1}{2}\mathbb{B}} u^{\frac{2n}{n-2}} d\mu_0 \right) \leq C \left(\int_{\mathbb{B}} u^{\varepsilon} d\mu_0 \right)^{\frac{2n}{\varepsilon(n-2)}}.$$

Proof. The proof of this lemma is done in two stages, the first one being a localization of what we did in the proof of Proposition 2.3. Let $R, r \geq 0$ such that $r + R \leq R_0$ and let ξ be a cut-off function with support in $\mathbb{B}(x, r + R)$ such that

- $0 \leq \xi \leq 1$,

- $|d\xi|_{g_0} \leq \frac{1}{r}$
- $\xi = 1$ on $\mathbb{B}(x, R)$

We have the equality

$$\begin{aligned}
& \frac{4(n-1)}{n-2} \int_{\mathbb{B}(x, R+r)} |d(\xi u)|_{g_0}^2 d\mu_0 \\
&= \frac{4(n-1)}{n-2} \int_{\mathbb{B}(x, R+r)} (\Delta_{g_0} u) u \xi^2 d\mu_0 + \frac{4(n-1)}{n-2} \int_{\mathbb{B}(x, R+r)} |d\xi|_{g_0}^2 u^2 d\mu_0 \\
&= \int_{\mathbb{B}(x, R+r)} V u^2 \xi^2 d\mu_0 - \int_{\mathbb{B}(x, R+r)} \text{Scal}_{g_0} u^2 \xi^2 d\mu_0 \\
&\quad + \frac{4(n-1)}{n-2} \int_{\mathbb{B}(x, R+r)} |d\xi|_{g_0}^2 u^2 d\mu_0.
\end{aligned}$$

We use now the Hebey-Vaugon Sobolev inequality (5) and we get that

$$\begin{aligned}
(10) \quad \alpha(n, 2) \left(\int_{\mathbb{B}(x, R+r)} (\xi u)^{\frac{2n}{n-2}} d\mu_0 \right)^{1-\frac{2}{n}} &\leq \frac{4(n-1)}{n-2} \int_{\mathbb{B}(x, R+r)} |d(\xi u)|^2 d\mu_0 \\
&\quad + B \int_{\mathbb{B}(x, R+r)} (\xi u)^2 d\mu_0
\end{aligned}$$

Hence with Hölder inequality, we get

$$\begin{aligned}
(11) \quad \alpha(n, 2) \left(\int_{\mathbb{B}(x, R+r)} (\xi u)^{\frac{2n}{n-2}} d\mu_0 \right)^{1-\frac{2}{n}} &\leq \left(B + \lambda_0 + \frac{4(n-1)}{n-2} \frac{1}{r^2} \right) \int_{\mathbb{B}(x, R+r)} u^2 d\mu_0 \\
&\quad + \left(\int_{\mathbb{B}(R+r)} (\text{Scal}_{g_f})^{\frac{n}{2}} d\mu_f \right)^{\frac{2}{n}} \left(\int_{\mathbb{B}(x, R+r)} (\xi u)^{\frac{2n}{n-2}} d\mu_0 \right)^{1-\frac{2}{n}}.
\end{aligned}$$

We obtain hence that for a constant C that depends only on g_0, R_0 and n :

$$(12) \quad \left(\int_{\mathbb{B}(x, R)} u^{\frac{2n}{n-2}} d\mu_0 \right)^{1-\frac{2}{n}} \leq \frac{C}{\delta r^2} \int_{\mathbb{B}(x, R+r)} u^2 d\mu_0.$$

Therefore, if $\varepsilon \geq 2$, using again Hölder inequality and setting $r = R \leq R_0/2$ we get

$$\left(\int_{\mathbb{B}} u^{\frac{2n}{n-2}} d\mu_0 \right)^{\frac{n-2}{n}} \leq \frac{C}{\delta R^2} \left(\int_{2\mathbb{B}} u^\varepsilon d\mu_0 \right)^{\frac{2}{\varepsilon}} \mu_0(\mathbb{B})^{\frac{\varepsilon-2}{\varepsilon}}$$

which gives Lemma 3.9.

Assume now that $\varepsilon < 2$ and let us go to the second stage of the proof. We will use a trick of P. Li and R. Schoen [28, Theorem 2.1] in order to obtain the wanted conclusion. Assuming now that $r \leq R \leq R_0/2$, and using Hölder inequality with exponent

$$\alpha = 1 + \frac{4}{(2-\varepsilon)(n-2)}$$

we obtain

$$\left(\int_{\mathbb{B}(x, R)} u^{\frac{2n}{n-2}} d\mu_0 \right)^{1-\frac{2}{n}} \leq \frac{C}{\delta r^2} \left(\int_{B(x, R+r)} u^\varepsilon d\mu_0 \right)^{1-\frac{1}{\alpha}} \left(\int_{\mathbb{B}(x, R+r)} u^{\frac{2n}{n-2}} d\mu_0 \right)^{\frac{1}{\alpha}}.$$

Let us define now $\lambda = \frac{1}{\alpha} \frac{n}{n-2}$,

$$M(R) = \left(\int_{\mathbb{B}(x,R)} u^{\frac{2n}{n-2}} d\mu_0 \right)^{1-\frac{2}{n}} \quad \text{and} \quad \mathbf{I} = \left(\int_{B(x,2R)} u^\varepsilon d\mu_0 \right)^{1-\frac{1}{\alpha}}.$$

Notice that $\lambda \in (0, 1)$ since $\frac{1}{\lambda} = 1 + \frac{2\varepsilon}{n(2-\varepsilon)}$. Equation (12) shows hence that

$$M(R) \leq \frac{C\mathbf{I}}{\delta r^2} M(R+r)^\lambda.$$

Iterating this equation with $r_1 + r_2 + \dots + r_k \leq R/2$ leads to

$$M(R) \leq \left(\frac{C\mathbf{I}}{\delta} \right)^{1+\lambda+\dots+\lambda^{k-1}} \prod_{\ell=1}^k \frac{1}{r_\ell^{2\lambda^{\ell-1}}} M(R+r_1+r_2+\dots+r_k)^{\lambda^k}.$$

If we choose $r_\ell = R2^{-\ell-2}$, with $\beta = \frac{1}{1-\lambda}$, a little bit of arithmetic gives that

$$M(R) \leq C \frac{1}{R^{2\beta}} \mathbf{I}^\beta.$$

It follows that

$$\left(\int_{\mathbb{B}(x,R)} u^{\frac{2n}{n-2}} d\mu_0 \right) \leq C \left(\int_{\mathbb{B}(x,2R)} u^\varepsilon d\mu_0 \right)^{\frac{2n}{\varepsilon(n-2)}}$$

which concludes the proof of Lemma 3.9. \square

Let us now conclude the proof of Theorem 3.7. Let $\varepsilon > 0$ be given by Lemma 3.8, and set $p = \frac{2n}{\varepsilon(n-2)}$. Let us write $w = u^{\frac{2n}{n-2}} = e^{nf}$. It follows from Lemmas 3.8 and 3.9 that there exists $C = C(n, g_0, R_0, \delta, \Lambda, \varepsilon) > 0$ such that for all g_0 -balls \mathbb{B} with radius less than $R_0/2$,

$$\left(\int_{\frac{1}{2}\mathbb{B}} w d\mu_0 \right) \left(\int_{\mathbb{B}} w^{-\frac{1}{p}} d\mu_0 \right)^p \leq C.$$

Hence for any g_0 -ball \mathbb{B} with radius less than $R_0/4$:

$$\left(\int_{\mathbb{B}} w d\mu_0 \right) \left(\int_{\mathbb{B}} w^{-\frac{1}{p}} d\mu_0 \right)^p \leq \theta^p \left(\int_{\mathbb{B}} w d\mu_0 \right) \left(\int_{2\mathbb{B}} w^{-\frac{1}{p}} d\mu_0 \right)^p \leq \theta^p C,$$

where θ is the doubling constant for the volume measure g_0 . This is precisely characterization *ii*) of A_∞ weights, cf Definition 3.1. \square

4. STRONG A_∞ WEIGHTS AND BI-HÖLDER COMPACTNESS

We now introduce so-called *strong A_∞ weights*, also known as *metric doubling measures*, which are special cases of A_∞ weights with strong geometric controls, such as Sobolev, Poincaré and isoperimetric inequalities.

We could not produce examples of sequences of conformal deformations satisfying the hypotheses of Theorem 1.2 which would not be uniformly strong A_∞ . Our Theorem 1.3 shows that any conformal deformation is actually uniformly strong A_∞ provided the $L^{n/2}$ norm of the scalar curvature on small balls is pinched enough, which explains this fact.

4.1. Definitions and basic facts. Let (M, g_0) be a closed Riemannian manifold, μ_0 its Riemannian volume. If $g_f = e^{2f}g_0$ is a metric conformal to g_0 , we continue to denote by μ_f and by d_f the corresponding Riemannian volume and distance respectively.

4.1.1. *Definitions.*

Definition 4.1. Let $g_f = e^{2f}g_0$ be a conformal deformation of g_0 . We say that the volume density e^{nf} is a strong- A_∞ weight with respect to the Riemannian metric g_0 if there are positive constants η, θ such that

- (1) $x \in M, r \leq \eta \Rightarrow \mu_f(\mathbb{B}(x, 2r)) \leq \theta \mu_f(\mathbb{B}(x, r))$.
- (2) $x, y \in M, d_0(x, y) \leq \eta \Rightarrow \frac{d_f(x, y)^n}{\theta^n} \leq \mu_f(\mathbb{B}(x, d_0(x, y))) \leq \theta^n d_f(x, y)^n$.

In above definition, $\mathbb{B}(x, r)$ is a geodesic ball of radius r for the fixed $d_0 = d_{g_0}$ distance. Notice that the strong A_∞ condition is invariant under scaling: if $d\mu_f = e^{nf}d\mu_0$ is a strong- A_∞ weight with constants (θ, η) then for every $h > 0$, $h^n d\mu_f$ is also a strong- A_∞ weight with same constants (θ, η) .

These strong A_∞ weights were introduced in [17], in relation with the quasiconformal jacobian problem. The jacobian of a quasiconformal map is a strong A_∞ weight, as is any A_1 -weight (see Definition 3.1 ii). A strong A_∞ weight with constants (θ, η) is an A_∞ -weight with constants only depending on θ, η , see Proposition 3.4 of [33]. Nevertheless, not all A_∞ weights are strong A_∞ . Various characterizations of strong A_∞ weights can be found in [17] and [33], together with examples and details about their relationships with the quasiconformal jacobian problem and the bi-Lipschitz embedding problem.

It is easy to see that if the conditions in definition 4.1 are satisfied for some θ, η then for any other $\eta' \geq \eta$ there is a constant θ' depending on (θ, η) and on the geometry of g_0 such that the same conditions are satisfied with constant (θ', η') . We also remark that the doubling condition on μ_f implies that if $m \in M$ is a point such that

$$\rho = d_0(m, x) = d_0(m, y) = \frac{1}{2}d_0(x, y) \leq \frac{\eta}{2}$$

then the balls $\mathbb{B}(m, \rho)$, $\mathbb{B}(x, d_0(x, y))$, $\mathbb{B}(y, d_0(x, y))$ have comparable μ_f measure.

Note that when $d_0(x, y) \leq \frac{1}{2}\text{inj}(M, g_0)$, there is a unique point $m \in M$ such that

$$\rho = d_0(m, x) = d_0(m, y) = \frac{1}{2}d_0(x, y).$$

The ball $\mathbb{B}(m, \rho)$ will be denoted $\mathbb{B}_{x,y}$. We obtain thus the following alternative and equivalent definition of strong A_∞ weight.

Definition 4.2. If θ, η are positive number such that $\eta \leq \frac{1}{4}\text{inj}(M, g_0)$, we say that the volume density e^{nf} associated to the metric $g_f = e^{2f}g_0$ is a strong- A_∞ weight (with respect to g_0) with constant (θ, η) if

- (1) $x \in M, r \leq \eta \Rightarrow \mu_f(\mathbb{B}(x, 2r)) \leq \theta \mu_f(\mathbb{B}(x, r))$.
- (2) $x, y \in M, d_0(x, y) \leq 2\eta \Rightarrow \frac{d_f(x, y)^n}{\theta^n} \leq \mu_f(\mathbb{B}_{x,y}) \leq \theta^n d_f(x, y)^n$.

It follows from Lemma 3.5 that the left inequality in property (2) in definition 4.1 is satisfied for all A_∞ weights. The specificity of strong A_∞ weights relies hence in the other inequality.

Let $g_f = e^{2f}g_0$ be a conformal deformation of g_0 such that its volume density e^{nf} is a strong- A_∞ weight (with respect to g_0) with constants (θ, η) . We are going to show several basic controls on the geometry of (M, g_f) .

4.1.2. *Geometry of g_f balls.* In this subsection, we show that the strong A_∞ condition implies that the geodesic ball for g_f and g_0 are comparable.

Let $B_f(x, r)$ denote the g_f -geodesic ball of center x and radius r .

In this section and in what follows we use the following notation:

$$R(x, r) := \mu_f(\mathbb{B}(x, r))^{\frac{1}{n}},$$

where, as before, $\mathbb{B}(x, r)$ is the g_0 geodesic ball of center x and radius r .

Lemma 4.3. *There exists $\kappa = \kappa(\eta, \theta) > 0$ such that for any $r \leq \eta/\kappa$:*

$$B_f(x, R(x, r/\kappa)) \subset \mathbb{B}(x, r) \subset B_f(x, R(x, \kappa r)).$$

Proof. The doubling condition implies a reverse doubling estimate (see [22, Lemma 2.10]): there are positive constants α, ω depending only on η and θ such that

$$(13) \quad x \in M, 0 < s < r \leq \eta \Rightarrow \omega^{-n} \left(\frac{s}{r}\right)^{\frac{n}{\alpha}} \leq \frac{\mu_f(\mathbb{B}(x, s))}{\mu_f(\mathbb{B}(x, r))} \leq \omega^n \left(\frac{s}{r}\right)^{n\alpha}.$$

The doubling condition also implies that the μ_f -measure of g_0 -geodesic spheres is zero ([14, 38]), hence the function $r \mapsto R(x, r)$ is continuous and strictly increasing, in particular we have

$$R(x, s) < R(x, r) \iff s < r.$$

We have for $d_0(x, y) \leq \eta$

$$(14) \quad \theta^{-1} \leq \frac{d_f(x, y)}{R(x, d_0(x, y))} \leq \theta.$$

and

$$x \in M, 0 < s < r \leq \eta \Rightarrow \omega^{-1} \left(\frac{s}{r}\right)^{\frac{1}{\alpha}} \leq \frac{R(x, s)}{R(x, r)} \leq \omega \left(\frac{s}{r}\right)^\alpha.$$

With $\kappa := (\omega\theta)^{\frac{1}{\alpha}}$, and assuming without loss of generality that $\kappa > 1$, we get for any $r \leq \eta/\kappa$:

$$\theta R(x, r) \leq R(x, \kappa r) \text{ and } R(x, r/\kappa) \leq \theta^{-1} R(x, r).$$

Since by (14), we have

$$B_f(x, R(x, r)/\theta) \subset \mathbb{B}(x, r) \subset B_f(x, \theta R(x, r))$$

we get

$$B_f(x, R(x, r/\kappa)) \subset \mathbb{B}(x, r) \subset B_f(x, R(x, \kappa r)).$$

□

Note that the two radius $R(x, r/\kappa)$ and $R(x, \kappa r)$ are uniformly comparable:

$$\omega^{-1} \kappa^{\frac{2}{\alpha}} \leq \frac{R(x, \kappa r)}{R(x, r/\kappa)} \leq \omega \kappa^{2\alpha}.$$

4.1.3. *Volume doubling for (M, d_f, μ_f) .* As we mentioned above, if a weight is strong- A_∞ , it is A_∞ . We proved that A_∞ weights are doubling for the balls of the background metric. We show now that the metric measure space (M, d_f, μ_f) is doubling (for g_f -geodesic balls) with constants only depending on η and θ .

Lemma 4.4. *There exists $\delta = \delta(\eta, \theta) > 0$ and $k = k(\eta, \theta) > 0$ such that for all $R \in (0, \delta)$,*

$$\mu_f(\mathbb{B}_f(x, 2R)) \leq 2^k \mu_f(\mathbb{B}_f(x, R)),$$

where $\mathbb{B}_f(x, R)$ is the d_f -geodesic ball of radius R and center x .

Proof. Let D be the diameter of (M, g_0) , then according to [11, Lemma 3.10], the volume doubling condition for (M, d_0, μ_f) implies that for any $x \in M$ and any $r \leq \eta$:

$$(15) \quad \mu_f(M) \leq \theta^{50+50\frac{D}{r}} \mu_f(\mathbb{B}(x, r)).$$

Hence

$$R(x, s) \leq \left(\theta^{-50-50\frac{D}{r}} \mu_f(M) \right)^{\frac{1}{n}} \Rightarrow s \leq r.$$

Now we have for $r \leq \eta/\kappa$:

$$\begin{aligned} \mu_f(B_f(x, R(x, r))) &\leq \mu_f(\mathbb{B}(x, \kappa r)) \\ &\leq \theta^k \mu_f(\mathbb{B}(x, 2^{-k} r \kappa)) \\ &\leq \theta^k \mu_f(B_f(x, R(x, 2^{-k} r \kappa^2))) \end{aligned}$$

But

$$R(x, 2^{-k} r \kappa^2) \leq \omega (2^{-k} \kappa^2)^\alpha R(x, r)$$

Chosing k such that $\omega (2^{-k} \kappa^2)^\alpha \leq \frac{1}{2}$ we get

$$\mu_f(B_f(x, R(x, r))) \leq \theta^k \mu_f(B_f(x, R(x, r)/2)).$$

Hence with

$$\delta := \left(\theta^{-50-50\kappa\frac{D}{\eta}} \mu_f(M) \right)^{\frac{1}{n}},$$

this eventually implies that

$$R \leq 2\delta \Rightarrow \mu_f(B_f(x, 2R)) \leq 2^k \mu_f(B_f(x, R)).$$

□

4.1.4. *Estimate of the diameter of (M, g_f) .* We are going to show that the diameter of (M, g_f) is comparable to $\mu_f(M)^{\frac{1}{n}}$.

Lemma 4.5. *There is a constant C depending only on g_0, θ, η such that*

$$C^{-1} (\mu_f(M))^{\frac{1}{n}} \leq \text{diam}(M, g_f) \leq C (\mu_f(M))^{\frac{1}{n}}.$$

Proof. The upper bound for $\text{diam}(M, g_f)$ is an immediate consequence of Lemma 3.5. Let us show the lower bound.

It is always possible to assume moreover that $\eta \leq D$. Then there is always two points $x, y \in M$ with $d_0(x, y) = \eta$ and we have

$$\text{diam}(M, g_f) \geq d_f(x, y) \geq \theta^{-1} (\mu_f(\mathbb{B}(x, \eta)))^{\frac{1}{n}},$$

therefore by (15), we get

$$\text{diam}(M, g_f) \geq C(\theta, \eta, D) (\mu_f(M))^{\frac{1}{n}}.$$

□

4.1.5. *Bi-Hölder compactness for sequence of Strong A_∞ conformal metric.* Let $(g_k = e^{2f_k} g_0)_{k \in \mathbb{N}}$ be a sequence of conformal deformations of g_0 such that the weights e^{nf_k} are strong- A_∞ weight with uniform constant (θ, η) . As mentioned earlier, it follows from [33] that all the g_k are uniformly A_∞ . If the volumes of (M, g_k) are uniformly bounded, Proposition 3.6 implies then that the sequence of metric spaces $(M, d_{f_k})_{k \in \mathbb{N}}$ has a subsequence which converges in Gromov-Hausdorff topology to some limiting space (M_∞, d_∞) . Strong A_∞ weights will give the following stronger convergence result.

Proposition 4.6. *Let $(g_k = e^{2f_k} g_0)_{k \in \mathbb{N}}$ be a sequence of Riemannian metrics conformal to g_0 such that the weights e^{nf_k} are strong- A_∞ weights with uniform constant (θ, η) .*

- i) *If there are positive constants v, V such that $\forall k: v \leq \mu_{f_k}(M) \leq V$, then there is a subsequence that converges in the C^α and Gromov-Hausdorff topology to (M, d_∞) , where d_∞ is a distance on M which is bi-Hölder equivalent to d_0 .*
- ii) *If $\lim_{k \rightarrow \infty} \mu_{f_k}(M) = 0$, then the sequence of metric spaces $((M, d_{f_k}))_k$ converges in the Gromov-Hausdorff topology to a point.*
- iii) *If there is a positive constant v such that $\forall k: v \leq \mu_{f_k}(M)$, then for any $p \in M$, there is a subsequence of the sequence of pointed metric spaces $((M, d_{f_k}, p))_k$ that converges in the pointed Gromov-Hausdorff topology to a pointed metric space.*

Proof. We will only show i); the other cases follow from a straightforward scaling argument. As soon as e^{nf} is a strong A_∞ with constants η, θ , we have seen that if $d_0(x, y) \leq \eta$ then

$$\theta^{-1} R(x, d_0(x, y)) \leq d_f(x, y) \leq \theta R(x, d_0(x, y)).$$

Moreover by (13) we have

$$\omega^{-1} \left(\frac{d_0(x, y)}{\eta} \right)^{\frac{1}{\alpha}} \mu_f(\mathbb{B}(x, \eta))^{\frac{1}{n}} \leq R(x, d_0(x, y)) \leq \omega \left(\frac{d_0(x, y)}{\eta} \right)^\alpha \mu_f(\mathbb{B}(x, \eta))^{\frac{1}{n}}.$$

Hence if $v \leq \mu_f(M) \leq V$, there is a constant λ depending only on $v, V, \theta, \omega, \eta, \alpha$ such that

$$\frac{1}{\lambda} (d_0(x, y))^{\frac{1}{\alpha}} \leq d_f(x, y) \leq \lambda (d_0(x, y))^\alpha.$$

Hence the identity map

$$\text{Id}: (M, d_f) \rightarrow (M, d_0)$$

is uniformly bi-Hölder continuous

Therefore a uniform strong A_∞ estimate for the $(e^{nf_k})_{k \in \mathbb{N}}$ together with a uniform control on the volume $0 < v \leq \mu_{f_k}(M) \leq V$ implies that the sequence (d_{f_k}) is precompact in the Hölder topology and for any converging subsequence, the limiting map d_∞ is a distance on M which is uniformly bi-Hölder equivalent to d_0 . \square

4.1.6. *Sobolev, Poincaré and isoperimetric inequalities.* Let us now mention some other important geometric facts about strong A_∞ weights shown in [17].

Theorem 4.7 (David-Semmes (1990)). *Let (M, g_0) be a compact smooth Riemannian manifold, and let $\omega = e^{nf}$ be a strong A_∞ weight with respect to g_0 with constants η and θ , and let $g_f = e^{2f} g_0 = \omega^{\frac{2}{n}} g_0$. Then*

- i) *There is a constant $\theta = \theta(g_0, \eta, \theta)$ such that for any $x \in M$ and any g_f -geodesic $B(x, r)$ of radius less than $\text{diam}_{g_f}(M)$:*

$$\theta^{-1} r^n \leq \mu_f(B(x, r)) \leq \theta r^n.$$

ii) There is a constant $\gamma = \gamma(g_0, \eta, \theta) > 0$ such that for any smooth domain $\Omega \subset M$ with $\mu_f(\Omega) \leq \frac{1}{2}\mu_f(M)$, we have

$$\gamma \mu_f(\Omega)^{\frac{n-1}{n}} \leq \mu_f(\partial\Omega).$$

iii) There is a constant $\lambda = \lambda(g_0, \eta, \theta)$ such that for any g_f -geodesic $B(x, r)$ of radius less than $\text{diam}_{g_f}(M)$: and any $\varphi \in C^1(B(x, r))$ with $\int_{B(x, r)} \varphi d\mu_f = 0$ then

$$\int_{B(x, r)} |\varphi| d\mu_f \leq \lambda r \int_{B(x, r)} |d\varphi|_{g_f} d\mu_f$$

Sobolev inequalities for (M, g_f) are consequences of doubling and Poincaré inequality, see [17].

4.2. Q-curvature and Strong A_∞ bounds. Let us finish this section by presenting the results of S. Brendle [7] and Yi Wang [42] which have inspired our analysis of conformal metrics with $L^{n/2}$ pinching of the scalar curvature via A_∞ weights. These works are devoted to the study of conformal deformations with L^1 bounds for the Branson Q -curvature, which we present now.

Let (M, g) be a Riemannian manifold of dimension $n = 4$. The Branson Q -curvature of (M, g) is defined by

$$Q_g = \frac{1}{12} \left(\Delta_g \text{Scal}_g + \frac{1}{4} \text{Scal}_g^2 - 3 |E_g|_g \right),$$

where E_g is the traceless part of the Ricci tensor. This Q -curvature has a natural conformal invariance: if $g_f = e^{2f} g_0$, then

$$(16) \quad 2e^{4f} Q_{g_f} = P_{g_0} f + 2Q_{g_0},$$

where P_{g_0} is the Paneitz operator on (M, g_0) , which we shall not define here (see for instance [7]). This equation is the analogous for the Q -curvature or the Yamabe equation (6) for the scalar curvature. The existence of a Paneitz operator and a Q -curvature satisfying the conformal covariance property (16) has been generalized to all even dimension by Fefferman and Graham in [18]. Note that the leading term of P_{g_0} is always $(\Delta_{g_0})^{\frac{n}{2}}$, and $P_{g_0} = (\Delta_{g_0})^{\frac{n}{2}}$ when g_0 is a flat metric.

The following result was shown in [7], Proposition 1.4.

Theorem 4.8 (Brendle (2003)). *Let (\mathbb{S}^n, g_S) be the standard n -sphere and let $C, \delta > 0$ and $R_0 \in (0, \text{diam}(\mathbb{S}^n))$ be fixed. Let $(g_k = e^{2f_k} g_S)_{k \in \mathbb{N}}$ be a sequence of conformal deformations of g_S with constant volume such that:*

- $\int_{\mathbb{S}^n} Q_{g_k}^2 d\text{vol}_{g_k} \leq C$;
- for all $x \in \mathbb{S}^n$,

$$\int_{\mathbb{B}(x, R_0)} |Q_{g_k}| d\text{vol}_{g_k} \leq \frac{1}{2}(n-1)! \sigma_n - \delta,$$

where $\mathbb{B}(x, R_0)$ is the g_S -geodesic ball of radius x and center x .

Then $(f_k)_{k \in \mathbb{N}}$ is bounded in $\mathcal{H}^n = W^{2, n}$ with bounds only depending on C, δ, n and R_0 .

Note that on the round sphere, we have

$$\int_{\mathbb{S}^n} |Q_{g_S}| d\text{vol}_{g_S} = (n-1)! \sigma_n :$$

the critical constant in Brendle theorem is half of the total Q -curvature of the sphere. It was later shown by Wang in [42] that under an analogous L^1 pinching condition for the Q -curvature, conformal deformations of the euclidean metric on \mathbb{R}^n are uniformly strong A_∞ weights.

Theorem 4.9 (Wang (2013)). *Let $C, \delta > 0$ be fixed and let $(g_f = e^{2f} \text{eucl})_{k \in \mathbb{N}}$ be a conformal deformation of the euclidean plane $(\mathbb{R}^n, \text{eucl})$, which is normal metrics, and such that:*

- $\int_M |Q_{g_f}| d\mu_f \leq C;$
- $\int_M Q_{g_f}^+ d\mu_f \leq \frac{1}{2}(n-1)! \sigma_n - \delta.$

Then the Riemannian volume density $(e^{nf})_{k \in \mathbb{N}}$ is a strong A_∞ weight with respect to the standard Lebesgue measure on \mathbb{R}^n , with constants only depending on n, C and δ .

A normal metric on \mathbb{R}^n is a Riemannian metric whose behaviour at infinity is not too pathological, we refer to [42] for a precise definition. For instance, any smooth metric whose scalar curvature is non-negative at infinity is normal.

The previous theorem is a reformulation of Theorems 3.2 and 4.1 of [42]. The parallel between Brendle and Wang's assumptions and the assumptions of our Theorem 1.3 is obvious; indeed our approach was inspired by Wang's proof. Note that the constant $\frac{1}{2}(n-1)! \sigma_n$ is optimal to get a strong A_∞ weight, as can easily be shown by considering a sequence of metrics built from removing a ball in \mathbb{R}^n and gluing on its boundary a very long cylinder with a round cap. Such sequence (even smoothed) cannot be uniformly strong A_∞ , since it has no uniform isoperimetric inequality.

The proof of Wang can be localized using the same approach as we do in this paper, which gives following result.

Theorem 4.10. *Let (M, g_0) be a compact smooth manifold, and $C, \delta > 0$ and $R_0 \in (0, \text{diam}(M, g_0))$ be fixed. Let $(g_f = e^{2f} g_0)_{k \in \mathbb{N}}$ be a sequence of conformal deformations of g_0 , such that:*

- $\int_M |Q_{g_f}| d\mu_f \leq C;$
- for all $x \in M,$

$$\int_{\mathbb{B}(x, R_0)} Q_{g_f}^+ d\mu_f \leq \frac{1}{2}(n-1)! \sigma_n - \delta.$$

Then the Riemannian volume density $(e^{nf})_{k \in \mathbb{N}}$ is a strong A_∞ weight with respect to g_0 , with constants only depending on $(M, g_0), C$ and δ .

We will not detail here the proof of this result, which is a combination of our arguments with those of [42].

5. $L^{n/2}$ PINCHING FOR THE SCALAR CURVATURE AND STRONG A_∞ WEIGHTS

In Section 4 we explained that a uniform strong A_∞ control coupled with uniform volume bounds has important geometric consequences, which were gathered in Corollary 1.4. This section is devoted to the proof of Theorem 1.3, which is our second main result. In

order to motivate the sequel of our study, we start with the proof of Theorem 1.3 assuming Theorems 5.1 and 5.7 which will be stated and proved in Sections 5.2 and 5.3 respectively.

5.1. Proof of Theorem 1.3. Let $g_f = e^{2f}g_0$ be a conformal deformation such that for some Λ_0 small enough (only depending on g_0) and for some $R_0 > 0$, we have

$$\forall x \in M, \quad \int_{\mathbb{B}(x, R_0)} |\text{Scal}_{g_f}|^{\frac{n}{2}} d\mu_f \leq \Lambda_0.$$

By Yamabe's equation (6), writing $e^{2f} = u^{\frac{4}{n-2}}$, we have

$$\frac{4(n-1)}{n-2} \Delta_{g_0} u = \text{Scal}_{g_f} u^{\frac{4}{n-2}} u - \text{Scal}_{g_0} u.$$

This is a Schrödinger type equation of the form $\Delta_{g_0} u + q_f u = 0$, with

$$q_f = \frac{n-2}{4(n-1)} (\text{Scal}_{g_0} - \text{Scal}_{g_f} e^{2f}).$$

Moreover, for all $x \in M$,

$$\begin{aligned} \|q_f\|_{L^{\frac{n}{2}}(\mathbb{B}(x, R_0), g_0)} &\leq C(g_0)R_0^2 + \left(\int_{\mathbb{B}(x, R_0)} |\text{Scal}_{g_f}|^{\frac{n}{2}} d\text{vol}_{g_f} \right)^{\frac{2}{n}} \\ &\leq C(g_0)R_0^2 + \Lambda_0. \end{aligned}$$

Hence if one chose Λ_0 to be smaller than the half of the ϵ of the Theorem 5.1, then for R_0 small (so that $C(g_0)R_0^2 \leq \epsilon/2$) the Schrödinger operator $\Delta_{g_0} + q_f$ satisfies the hypothesis of the Theorem 5.1 and since u is a *positive* smooth solution of $(\Delta_{g_0} + q_f)u = 0$, the lowest eigenvalue of $\Delta_{g_0} + q_f$ is necessary zero, with eigenfunction u . By Theorem 5.1, we can decompose

$$f = h + w,$$

where $h \in W^{2, \frac{n}{2}}(M, g_0)$ and $w \in C^\alpha(M, d_0)$, with

$$\sup_{x, y \in M} \frac{|w(x) - w(y)|}{d_0(x, y)^\alpha} \leq C(g_0, R_0, \Lambda_0)$$

and

$$(17) \quad \|dh\|_{L^n} + \|\Delta h\|_{L^{\frac{n}{2}}} \leq C(g_0, R_0, \Lambda_0).$$

Let x_0 be a chosen base point in M . Since w is bounded in the C^α topology induced by d_0 , there exists $C > 0$ such that for all $x \in M$,

$$|w(x) - w(x_0)| \leq C \text{diam}(M, g_0)^\alpha.$$

Now, combining (17) with Theorem 5.7, there exists $\eta = \eta(g_0, R_0, \Lambda_0)$ and $\theta = \theta(g_0, R_0, \Lambda_0)$ such that $e^{\eta h}$ is a strong A_∞ weight with respect to (M, g_0) with constants η and θ . Since $w - w(x_0)$ is uniformly bounded from above and below, and since being Strong A_∞ weight is a scale invariant condition, this implies that $e^{\eta f} = e^{\eta(h+w-w(x_0))} e^{\eta w(x_0)}$ is also a strong A_∞ weight with constants η and θ . Moreover, by Proposition 5.9, for all $\alpha \in (0, 1)$, the distances d_f and d_0 are α -bi-Hölder with constants only depending on g_0, R_0, Λ_0 and α , which concludes the proof of Theorem 1.3.

5.2. On the ground state of Schrödinger operators with critical potential. The purpose of this section is to give a regularity result for the first eigenfunction of a Schrödinger operator whose potential has small $L^{n/2}$ norm on small balls, which is the first key step in the proof of Theorem 1.3. The results presented in this paragraph are valid for any compact Riemannian manifold (M, g) and of independent interest.

Theorem 5.1. *Let (M^n, g) be a closed Riemannian manifold of dimension $n > 2$. There are constants $\epsilon, \alpha > 0$ depending only of (M^n, g) such that if $V \in L^{\frac{n}{2}}$ satisfies:*

- (1) *the lowest eigenvalue of the Schrödinger operator $\Delta - V$ is zero;*
- (2) *there is some $\rho > 0$ such that for any $x \in M$:*

$$\int_{\mathbb{B}(x, \rho)} |V|^{\frac{n}{2}} < \epsilon.$$

Then for all positive C^2 solution $\varphi > 0$ of $\Delta\varphi - V\varphi = 0$, there are $f \in W^{2, \frac{n}{2}}$ and $w \in C^\alpha(M)$ such that

$$\varphi = e^{f+w}.$$

Moreover, writing $\mathbf{I} := \sup_x \left(\int_{\mathbb{B}(x, \rho)} |V|^{\frac{n}{2}} \right)^{\frac{2}{n}}$ we have

$$\|w\|_{C^\alpha} := \sup_{x, y \in M} \frac{|w(x) - w(y)|}{d_0(x, y)^\alpha} \leq C(g, \rho, \mathbf{I})$$

and

$$\|df\|_{L^n} + \|\Delta f\|_{L^{\frac{n}{2}}} \leq C(g, \rho)\mathbf{I}.$$

This theorem is a key step in showing that conformal deformations whose scalar curvature has small $L^{n/2}$ norm on small balls are strong A_∞ weights.

In order to prove Theorem 5.1, we need some preliminary tools. Let us start by recalling the following classical consequence of the fixed point theorem.

Proposition 5.2. *Let $(X, \|\bullet\|)$ be a Banach space, $\kappa \in (0, 1)$. If $S: \mathcal{B}(0, \rho) \rightarrow X$ is a κ -Lipschitz map with*

$$\|S(0)\| + \kappa\rho < \rho$$

then there is a unique $x \in \mathcal{B}(0, \rho)$ such that $S(x) = x$. Moreover we have $\|x\| \leq \frac{\|S(0)\|}{1 - \kappa}$.

Let $A(g)$ be the best constant in the following functional inequality on the closed Riemannian manifold (M, g) :

$$(18) \quad \forall \varphi \in C^\infty(M): \|d\varphi\|_{L^n} \leq A(g)\|\Delta\varphi\|_{L^{\frac{n}{2}}}.$$

Proposition 5.3. *Let (M, g) be a compact smooth Riemannian manifold such that $\text{Ric}_g \geq -(n-1)k^2g$, then the best constant $A(g)$ in (18) satisfies*

$$A(g) \leq e^{c(n)(1+k \text{diam}(M, g))} \frac{\text{diam}(M, g)}{\text{vol}^{\frac{1}{n}}(M, g)}.$$

The above estimate on $A(g)$ can be shown using a lower bound on the Ricci curvature, and upper bounds on the diameter and volume of (M, g) ; it follows from results of S. Gallot [19] and of D. Bakry [3, Théorème 4.1, Lemme 4.2 and Corollaire 4.3]. We will not show it here, since we will not need this expression.

Let us now show the following property of the ground state of the Schrödinger operator $\Delta - V$.

Proposition 5.4. *Let $V \in L^\infty(M) \cap L^{\frac{n}{2}}(M)$ and let λ_0 be the bottom of the spectrum of the Schrödinger operator $\Delta - V$. If*

$$8A(g)^2 \|V\|_{L^{\frac{n}{2}}} < 1$$

then for any $\varphi > 0$ satisfying $\Delta\varphi - V\varphi = \lambda_0\varphi$, we have

$$\|d \log \varphi\|_{L^n} \leq 2A(g) \|V\|_{L^{\frac{n}{2}}}.$$

Proof. We remark that if $\varphi = e^u$ satisfies

$$\Delta\varphi - V\varphi = \lambda_0\varphi,$$

then we have

$$\Delta u - |du|^2 = V + \lambda_0.$$

Moreover up to a scaling constant, there is a unique non trivial solution to the equation $\Delta\varphi - V\varphi = \lambda_0\varphi$. We introduce the Banach space

$$W_\perp^{1,n} := \{v \in W^{1,n}(M), \int_M v = 0\}$$

endowed with the norm

$$\|v\|_\perp = \|dv\|_{L^n}.$$

We introduce the operator $S: W_\perp^{1,n} \rightarrow W_\perp^{1,n}$ defined by

$$S(v) = \Delta^{-1}V + \Delta^{-1}|dv|^2.$$

where for all $f \in L^{\frac{n}{2}}$, we have denote $u := \Delta^{-1}f$ the solution of the equation

$$\begin{cases} \Delta u = f - f_M \\ \int_M u = 0. \end{cases}$$

By definition of $A(g)$, we have

$$\|S(0)\|_\perp \leq A(g) \|V\|_{L^{\frac{n}{2}}},$$

and

$$\|S(u) - S(w)\|_\perp \leq A(g) \|u - w\|_\perp (\|u\|_\perp + \|w\|_\perp).$$

As a consequence the restriction of S to the ball $\mathcal{B}(0, \rho) \subset W_\perp^{1,n}$, is $2A(g)\rho$ -Lipschitz and if

$$(19) \quad A(g) \|V\|_{L^{\frac{n}{2}}} + 2A(g)\rho^2 < \rho$$

then there is a unique $v \in \mathcal{B}(0, \rho)$ with $S(v) = v$ and

$$\|v\|_\perp \leq A(g) \frac{\|V\|_{L^{\frac{n}{2}}}}{1 - 2A(g)\rho}.$$

Choosing $\rho = (4A(g))^{-1}$, the condition (19) is satisfied as soon as

$$A(g) \|V\|_{L^{\frac{n}{2}}} < (8A(g))^{-1}.$$

We have obtained hence a function $v \in W_\perp^{1,n}$ such that $S(v) = v$ and $\|v\|_\perp \leq 2A(g) \|V\|_{L^{\frac{n}{2}}}$. Therefore there is a constant c such that

$$\Delta v - |dv|^2 = V + c,$$

therefore

$$\Delta e^v - V e^v = c e^v.$$

Since the only positive eigenfunction of $\Delta - V$ is associated to λ_0 , we have $c = \lambda_0$, $e^v = k\varphi$ for some $k > 0$ and $\|d \log \varphi\|_{L^n} = \|v\|_\perp \leq 2A(g) \|V\|_{L^{n/2}}$. \square

Let us now define the Sobolev constant $\beta = \beta(g_0) > 0$ of (M, g_0) to be the best constant in the inequality

$$(20) \quad \forall \varphi \in \mathcal{C}^\infty(M): \beta \|\varphi\|_{L^{\frac{2n}{n-2}}}^2 \leq \|d\varphi\|_{L^2}^2 + \|\varphi\|_{L^2}^2.$$

Note that this Sobolev inequality implies that

$$\beta \leq \text{Vol}(M, g_0)^{\frac{2}{n}}.$$

We also have that for any $\delta \in (0, 1)$:

$$\forall \varphi \in \mathcal{C}^\infty(M): \delta^2 \beta \|\varphi\|_{L^{\frac{2n}{n-2}}}^2 \leq \int_M [|d\varphi|_{g_0}^2 + \delta^2 |\varphi|^2] d\mu_0.$$

Hence we can easily prove the following lemma:

Lemma 5.5. *If $W \in L^{\frac{n}{2}}$ satisfies $\|W\|_{L^{\frac{n}{2}}} \leq \beta \delta^2$, then the Schrödinger operator $\Delta + \delta^2 + W$ is non negative.*

Proof. The Hölder inequality implies that if $\varphi \in \mathcal{C}^1(M)$:

$$-\int_M W \varphi^2 \leq \|W\|_{L^{\frac{n}{2}}} \|\varphi\|_{L^{\frac{2n}{n-2}}}^2 \leq \delta^2 \beta \|\varphi\|_{L^{\frac{2n}{n-2}}}^2 \leq \int_M [|d\varphi|_{g_0}^2 + \delta^2 |\varphi|^2] d\mu_0$$

□

Lemma 5.6. *Assume that $x_0 \in M$ and r_0 are chosen so that*

$$\text{Vol} \mathbb{B}(x_0, r_0) \leq \left(\frac{\beta}{2}\right)^{\frac{n}{2}}.$$

If $q \in L^{\frac{n}{2}}$ is such that

$$\text{supp}(q) \subset \mathbb{B}(x_0, r_0) \text{ and } \|q\|_{L^{\frac{n}{2}}} \leq \beta$$

then there is a constant c_0 satisfying :

$$|c_0| \leq \frac{2}{\beta} \|q\|_{L^{\frac{n}{2}}}$$

such that the lowest eigenvalue of the Schrödinger operator

$$\Delta + q + c_0 \mathbf{1}_{M \setminus \mathbb{B}(x_0, r_0)}$$

is zero.

Proof. Let $\lambda_0(c)$ be the lowest eigenvalue of the Schrödinger operator

$$\Delta + q + c \mathbf{1}_{M \setminus \mathbb{B}(x_0, r_0)}.$$

Using the constant function, we get that $\lambda_0(c) \leq 0$ as soon as

$$\int_M q + c \text{Vol}_{g_0}(M \setminus \mathbb{B}(x_0, r_0)) \leq 0.$$

Moreover, $\text{Vol} \mathbb{B}(x_0, r_0) \leq \frac{\text{Vol}(M, g_0)}{2}$, hence for $c_- = -\frac{2}{\text{Vol}(M, g_0)} \int_M |q|$, we have $\lambda_0(c_-) \leq 0$.

Using Lemma 5.5 for $W = q - c \mathbf{1}_{\mathbb{B}(x_0, r_0)}$, we also see that if $c > 0$ satisfies

$$\|q - c\|_{L^{\frac{n}{2}}(\mathbb{B}(x_0, r_0))} \leq \beta c$$

then $\lambda_0(c) \geq 0$. For instance for $c_+ = \frac{2}{\beta} \|q\|_{L^{\frac{n}{2}}}$, we have $\lambda_0(c_+) \geq 0$. Hence there is a $c_0 \in [c_-, c_+]$ such that $\lambda_0(c_0) = 0$.

Moreover, by Hölder inequality we have

$$|c_-| \leq \frac{2}{\text{Vol}_{g_0}(M)} (\text{Vol}_{g_0}(M))^{1-\frac{2}{n}} \|q\|_{L^{\frac{n}{2}}} \leq \frac{2}{\text{Vol}_{g_0}^{\frac{2}{n}}(M)} \|q\|_{L^{\frac{n}{2}}} \leq \frac{2}{\beta} \|q\|_{L^{\frac{n}{2}}}.$$

Therefore $|c_0| \leq \frac{2}{\beta} \|q\|_{L^{\frac{n}{2}}}$, which concludes the proof of Lemma 5.6. \square

We can now prove Theorem 5.1.

Proof of Theorem 5.1. We will use the constants $A(g_0)$ and β given by the estimate (18) and the Sobolev inequality (20). We can always assume that ρ also satisfies

$$\forall x \in M: \text{Vol}_{g_0} \mathbb{B}(x, \rho) \leq \left(\frac{\beta}{2}\right)^{\frac{2}{n}}.$$

We choose a finite cover

$$M = \cup_{i=1}^N \mathbb{B}(x_i, \rho/2)$$

such that for any $i \neq j: \mathbb{B}(x_i, \rho/4) \cap \mathbb{B}(x_j, \rho/4) = \emptyset$. If

$$\mathbf{I} := \sup_x \left(\int_{\mathbb{B}(x, \rho)} |V|^{\frac{n}{2}} \right)^{\frac{2}{n}} \leq \frac{\beta}{2}$$

then for any $i \in \{1, \dots, N\}$, using Lemma 5.6, we find c_i such that with

$$q_i = V \mathbf{1}_{\mathbb{B}(x_i, \rho)} + c_i \mathbf{1}_{M \setminus \mathbb{B}(x_i, \rho)},$$

the lowest eigenvalue of the Schrödinger operator $\Delta - q_i$ is zero. We have

$$\|q_i\|_{L^{\frac{n}{2}}} \leq \frac{3 \text{Vol}_{g_0}(M)^{\frac{2}{n}}}{\beta} \|V\|_{L^{\frac{n}{2}}(\mathbb{B}(x_i, \rho))}.$$

Hence using Proposition 5.4, we know that if

$$24 \frac{\text{Vol}_{g_0}(M)^{\frac{2}{n}}}{\beta} \mathbf{I} A^2(g_0) < 1$$

then we find $\varphi_i \in \ker(\Delta - q_i)$, $\varphi > 0$, with

$$\|d \log \varphi_i\|_{L^n} \leq C(g_0) \mathbf{I}.$$

On the other hand,

$$\Delta \log \varphi_i = q_i + |d \log \varphi_i|^2$$

hence

$$\|\Delta \log \varphi_i\|_{L^{\frac{n}{2}}} \leq C(g_0) \mathbf{I}.$$

We choose ϵ such that $24 \frac{\text{Vol}_{g_0}(M)^{\frac{2}{n}}}{\beta} A^2(g_0) \epsilon = \frac{1}{2}$. Let $\varphi: M \rightarrow (0, +\infty)$ be a global solution of $\Delta \varphi = V \varphi$. For all $i \in \{1, \dots, N\}$, we define $\psi_i: \mathbb{B}(x_i, \rho) \rightarrow \mathbb{R}$ by

$$\varphi = \varphi_i \psi_i.$$

Since on the ball $\mathbb{B}(x_i, \rho)$, we also have $\Delta \varphi_i = V \varphi_i$, we get

$$\Delta \psi_i - 2 \langle d \log \varphi_i, d \psi_i \rangle = 0.$$

Using the work of O. A. Ladyzhenskaya and N. N. Ural'tseva [27] and of M.V. Safonov [30], we know that ψ_i is Hölder continuous. There is some $\alpha = \alpha(g_0, \rho, \epsilon)$ such that

$$\|\psi_i\|_{\dot{C}^\alpha(\mathbb{B}(x_i, 3\rho/4))} := \sup_{x, y \in \mathbb{B}(x_i, 3\rho/4)} \frac{|\psi_i(x) - \psi_i(y)|}{d_0(x, y)^\alpha} \leq C(g_0, \rho, \mathbf{I}) \inf_{x \in \mathbb{B}(x_i, 3\rho/4)} \psi_i(x).$$

We let $\delta_i = \inf_{\mathbb{B}(x_i, 3\rho/4)} \psi_i$ and $\psi_i = \delta_i e^{w_i}$ and $\varphi_i = \frac{e^{f_i}}{\delta_i}$ and $\bar{f}_i = \int_{\mathbb{B}(x_i, 3\rho/4)} f_i$. We have

$$\|w_i\|_{L^\infty(\mathbb{B}(x_i, 3\rho/4))} + \|w_i\|_{\dot{C}^\alpha(\mathbb{B}(x_i, 3\rho/4))} \leq C(g_0, \rho, \mathbf{I})$$

and

$$\|df_i\|_{L^n} + \|\Delta f_i\|_{L^{\frac{n}{2}}} \leq C(g_0)\mathbf{I}.$$

Moreover, by Sobolev inequality

$$\|f_i - \bar{f}_i\|_{L^n} \leq C(g_0)\mathbf{I}.$$

Note that on all ball $\mathbb{B}(x_i, \rho)$, we have

$$(21) \quad \varphi = \varphi_i \psi_i = e^{w_i} e^{f_i}.$$

Let us fix now a partition of unity $(\chi_i)_{i \in \{1, \dots, N\}}$ with $\text{supp} \chi_i \subset \mathbb{B}(x_i, 3\rho/4)$ and $\chi_i = 1$ on $\mathbb{B}(x_i, \rho/4)$. This partition of unity only depends on (M, g_0) . Let us define

$$f := \sum_i \chi_i (f_i - \bar{f}_i) \quad \text{and} \quad w = \sum_i \chi_i (w_i + \bar{f}_i).$$

The functions f and w are well defined on M and it follows from (21) that $\varphi = e^f e^w$.

We have moreover $\|d \log f\|_{L^n} + \|\Delta f\|_{L^{\frac{n}{2}}} \leq C(g_0)\mathbf{I}$. We also have $\log \varphi = f + w$ and

$$\Delta \log \varphi - |d \log \varphi|^2 = V$$

Hence by Stokes formula,

$$\int_M |d \log \varphi|^2 d\mu_0 \leq \int_M |V| d\mu_0 \leq (\mu_0(M))^{1-\frac{2}{n}} \|V\|_{L^{\frac{n}{2}}} \leq C(g_0)\mathbf{I}.$$

Using the Poincaré inequality, one conclude that for $c = \int_M \log(\varphi) d\mu_0$:

$$\|f + w - c\|_{L^2} \leq C(g_0)\mathbf{I}.$$

On each ball $\mathbb{B}(x_i, \rho/4)$, the oscillation of $w_i = w - \bar{f}_i$ are controlled hence one get that

$$|\bar{f}_i - c| \leq C(g_0) \|f_i - c\|_{L^2(\mathbb{B}(x_i, \rho/4))} \leq C(g_0, \mathbf{I}).$$

Hence $w = \sum \chi_i (w_i + \bar{f}_i)$ is bounded in $\dot{C}^\alpha(M)$. \square

5.3. Strong A_∞ weights from $W^{1,n}$ bounds. In this section, we explain how the proof of Theorem 3.1 of [6] can be adapted for closed Riemannian manifolds, which gives the following result. This is the second key step in the proof of Theorem 1.3.

Theorem 5.7. *Let (M, g_0) be a closed Riemannian manifold. If $f: M \rightarrow \mathbb{R}$ is such that*

$$\|df\|_{L^n} \leq \mathbf{I},$$

for some \mathbf{I} . Then the weight e^{nf} is a strong A_∞ weight with respect to g_0 , with constants (θ, η) where η depends only on g_0 and θ depends only on g_0 and \mathbf{I} .

Note that, unlike Theorem 3.1 of [6], our statement and our proof are scale invariant.

5.3.1. *Presentation of the setting.* The estimates will depend on several geometric and analytic estimates. We continue to denote by μ_0 the Riemannian volume measure and by d_0 the Riemannian distance associated to g_0 . There is some $\eta > 0$ and positive constants θ, γ, C (that can be chosen depending only on the dimension n) such that for any ball $\mathbb{B} \subset M$ with radius $r(\mathbb{B}) \leq 2\eta$ we have

- i) as already mentioned in (8), $\theta^{-1}r(\mathbb{B})^n \leq \mu_0(\mathbb{B}) \leq \theta r(\mathbb{B})^n$;
- ii) as shown in [20], Lemma 7.16, for any Lipschitz function $\varphi: \mathbb{B} \rightarrow \mathbb{R}$ and any $x \in \mathbb{B}$:

$$|\varphi(x) - \varphi_{\mathbb{B}}| \leq C \int_{\mathbb{B}} \frac{|d\varphi|_{g_0}(z)}{d_0^{n-1}(x, z)} d\mu_0(z);$$

- iii) the following Adams-Moser-Trudinger inequality: for any Lipschitz function $\varphi: \mathbb{B} \rightarrow \mathbb{R}$ and any $x \in \mathbb{B}$:

$$\int_{\mathbb{B}} e^{\left(\frac{\gamma|\varphi(x) - \varphi_{\mathbb{B}}|}{\|d\varphi\|_{L^n(\mathbb{B})}}\right)^{\frac{n}{n-1}}} d\mu_0(x) \leq C.$$

For the Euclidean space \mathbb{R}^n , this inequality is due to N. Trudinger [40]. According to [20, Lemma 7.13], the first two geometric control i-ii) implies the Trudinger inequality.

We will use several times the following integration by parts formula

$$\begin{aligned} \int_{\mathbb{B}(x, R) \setminus \mathbb{B}(x, r)} \varphi(d_0(x, y)) f(y) d\mu_0(y) &= \varphi(R) \int_{\mathbb{B}(x, R)} f(y) d\mu_0(y) \\ &\quad - \varphi(r) \int_{\mathbb{B}(x, r)} f(y) d\mu_0(y) - \int_r^R \varphi'(r) \int_{\mathbb{B}(x, r)} f(y) d\mu_0(y) dr. \end{aligned}$$

5.3.2. *Estimate on $\mu_f(\mathbb{B})$.* We are going to prove that μ_f is an A_p weights for every $p > 1$ using characterization ii) in Definition 3.1. Let $\mathbb{B}(o, \rho) \subset M$ be such that its radius satisfies $r(\mathbb{B}) \leq \eta$. We have for any $x \in \mathbb{B}$ and any $\lambda \in \mathbb{R}$:

$$\lambda|f(x) - f_{\mathbb{B}}| \leq \frac{n-1}{n} \left(\frac{\gamma|f(x) - f_{\mathbb{B}}|}{\|df\|_{L^n(\mathbb{B})}} \right)^{\frac{n}{n-1}} + \frac{1}{n} \left(\frac{|\lambda|}{\gamma} \|df\|_{L^n(\mathbb{B})} \right)^n.$$

Hence, using Adams-Moser-Trudinger inequality, we get

$$\begin{aligned} \int_{\mathbb{B}} e^{\lambda f} d\mu_0 &\leq e^{\lambda f_{\mathbb{B}}} e^{\left(\frac{|\lambda|}{\gamma} \|df\|_{L^n(\mathbb{B})}\right)^n} \int_{\mathbb{B}} \exp\left(\left(\frac{\gamma|f(x) - f_{\mathbb{B}}|}{\|df\|_{L^n(\mathbb{B})}}\right)^{\frac{n}{n-1}}\right) d\mu_0(x) \\ &\leq C(g_0, |\lambda|, \mathbf{I}) e^{\lambda f_{\mathbb{B}}} \end{aligned}$$

Using the Cauchy-Schwarz inequality, this implies

$$1 \leq \int_{\mathbb{B}} e^{\lambda f} d\mu_0 \times \int_{\mathbb{B}} e^{-\lambda f} d\mu_0 \leq C(g_0, |\lambda|, \mathbf{I}) \int_{\mathbb{B}} e^{\pm \lambda f} d\mu_0 e^{\lambda f_{\mathbb{B}}},$$

and we eventually get

$$(22) \quad \frac{e^{\lambda f_{\mathbb{B}}}}{C(g_0, |\lambda|, \mathbf{I})} \leq \int_{\mathbb{B}} e^{\lambda f} d\mu_0 \leq C(g_0, |\lambda|, \mathbf{I}) e^{\lambda f_{\mathbb{B}}}.$$

This easily implies that for any $p > 1$ there is a constant C depending on g_0, \mathbf{I}, p such that

$$(23) \quad \int_{\mathbb{B}} e^{nf} d\mu_0 \left(\int_{\mathbb{B}} e^{-\frac{n}{p-1}f} d\mu_0 \right)^{p-1} \leq C.$$

Hence, e^{nf} is an A_p weight, in particular $d\mu_f$ is a doubling measure. Moreover, by Lemma 3.5, this implies that there exists a constant $C = C(g_0, \mathbf{I}) > 0$ such that

$$d_0(x, y) \leq 10\eta \Rightarrow d_f(x, y) \leq C (\mu_f(\mathbb{B}_{x,y}))^{\frac{1}{n}}.$$

5.3.3. *Distance estimate.* To show that $d\mu_f$ is a strong A_∞ weight, we are only left with showing that for all $x, y \in M$ and for some $C > 0$,

$$d_{g_f}(x, y) \geq C \mu_f(\mathbb{B}_{x,y})^{\frac{1}{n}}.$$

In this purpose, we will need the following lemma:

Lemma 5.8. *For any $\epsilon > 0$ there is a constant C depending on g_0, \mathbf{I} and $\epsilon > 0$ such that for any ball $\mathbb{B} \subset M$ with $r(\mathbb{B}) \leq \eta$ there is a set $E \subset \mathbb{B}$ such $E \subset \bigcup_\alpha \mathbb{B}_\alpha$ with*

$$\sum_\alpha r(\mathbb{B}_\alpha) \leq \epsilon r(\mathbb{B})$$

and

$$x \in \mathbb{B} \setminus E \Rightarrow |f(x) - f_{\mathbb{B}}| \leq C.$$

Proof. Assume that $x \in \mathbb{B}$ is such that for any $r \in (0, 2r(\mathbb{B}))$ we have

$$\int_{\mathbb{B}(x,r)} |df|^n \leq \lambda \frac{r}{r(\mathbb{B})}$$

then using $V(x, r) := \int_{\mathbb{B}(x,r)} |df(y)| d\mu_0(y)$ and one gets

$$\begin{aligned} |f(x) - f_{\mathbb{B}}| &\leq \int_{\mathbb{B}(x, 2r(\mathbb{B}))} \frac{C}{d_0^{n-1}(x, y)} |df(y)| d\mu_0(y) \\ &\leq \frac{C}{(2r(\mathbb{B}))^{n-1}} V(x, 2r(\mathbb{B})) + \int_0^{2r(\mathbb{B})} \frac{(n-1)C}{r^n} V(x, r) dr \end{aligned}$$

With Hölder inequality, one has

$$V(x, r) \leq \theta^{1-\frac{1}{n}} r^{n-1} \left(\int_{\mathbb{B}(x,r)} |df|^n \right)^{1/n} \leq \theta^{1-\frac{1}{n}} r^{n-1} \left(\lambda \frac{r}{r(\mathbb{B})} \right)^{1/n}.$$

Hence there is a constant depending only of g_0 such that

$$|f(x) - f_{\mathbb{B}}| \leq C \lambda^{\frac{1}{n}}.$$

If now E is the subset of point x where there is a $r_x \in (0, 2r(\mathbb{B}))$ such that

$$\int_{\mathbb{B}(x, r_x)} |df|^n > \lambda \frac{r_x}{r(\mathbb{B})}$$

Then using Vitali covering lemma, one gets a covering $E \subset \bigcup \mathbb{B}_\alpha$ where $\mathbb{B}_\alpha = 5\mathbb{B}(x_\alpha, r_\alpha)$ such that

$$\int_{\mathbb{B}(x_\alpha, r_\alpha)} |df|^n > \lambda \frac{r_\alpha}{r(\mathbb{B})}$$

and the balls $\mathbb{B}(x_\alpha, r_\alpha)$ are disjoint. In particular this implies that

$$\sum r(\mathbb{B}_\alpha) = \sum 5r_\alpha \leq \frac{5r(\mathbb{B})}{\lambda} \sum \int_{\mathbb{B}(x_\alpha, r_\alpha)} |df|^n \leq \frac{5r(\mathbb{B})}{\lambda} \int_M |df|^n.$$

Choosing $\lambda = \frac{5}{\epsilon \mathbf{I}^n}$, we get the desired result. \square

Let now $x, y \in M$ such that $\rho := d_0(x, y) \leq 10\eta$ and let $\gamma: [0, 1] \rightarrow M$ be a \mathcal{C}^1 curve joining x to y :

$$\gamma(0) = x \text{ and } \gamma(1) = y$$

Let $\tau > 0$ be the first time with $d_0(x, \gamma(\tau)) = \rho$ and let $\bar{y} = \gamma(\tau)$, $\mathbb{B} := \mathbb{B}_{x, \bar{y}} = \mathbb{B}(m, \rho/2)$. We have for any $t \in [0, \tau]$, $\gamma(t) \in 3\mathbb{B}$. Let $c(t) = \exp_m(tu)$ be the unit speed g_0 geodesic joining x to \bar{y} and $L := c([-3\rho/2, 3\rho/2])$ it is a diameter of $3\mathbb{B}$ and $\Pi: 3\mathbb{B} \rightarrow L$ the projection onto L associated to the Euclidean metric d_{eucl} induced by \exp_m . We know that Π is 1-Lipschitz for the Euclidean distance and for $x, y \in L$: $d_{\text{eucl}}(x, y) = d_0(x, y)$. Hence Π is κ -Lipschitz for the distance d_0 . According to Lemma 5.8 with

$$(24) \quad \epsilon = \frac{1}{6\kappa}$$

we can find $E \subset 3\mathbb{B}$ such that $E \subset \bigcup_\alpha \mathbb{B}_\alpha$ with

$$\sum_\alpha r(\mathbb{B}_\alpha) \leq \epsilon \frac{3}{2} \rho$$

and

$$x \in 3\mathbb{B} \setminus E \Rightarrow |f(x) - f_{\mathbb{B}}| \leq C.$$

We have

$$\begin{aligned} L_{g_f}(\gamma) &\geq L_{g_f}(\gamma|_{[0, \tau]}) \geq L_{g_f}(\gamma|_{[0, \tau]} \setminus E) \\ &\geq e^{-C} e^{f_{\mathbb{B}}} L_{g_0}(\gamma|_{[0, \tau]} \setminus E) \\ &\geq e^{-C} e^{f_{\mathbb{B}}} \kappa^{-1} L_{\text{eucl}}(\gamma|_{[0, \tau]} \setminus E) \\ &\geq e^{-C} e^{f_{\mathbb{B}}} \kappa^{-1} L_{\text{eucl}}(\Pi(\gamma|_{[0, \tau]} \setminus E)) \end{aligned}$$

Since $\Pi(\gamma|_{[0, \tau]})$ contains the geodesic segment joining x to \bar{y} ,

$$L_{\text{eucl}}(\Pi(\gamma|_{[0, \tau]})) \geq \rho.$$

whereas

$$L_{\text{eucl}}(\Pi(E)) \leq L_{\text{eucl}}(\Pi(\bigcup_\alpha \mathbb{B}_\alpha)) \leq \kappa \sum_\alpha 2r(\mathbb{B}_\alpha) \leq 2\kappa \epsilon \frac{3}{2} \rho.$$

With the choice of ϵ given by (24), we get

$$L_{\text{eucl}}(\Pi(E)) \leq \frac{1}{2} \rho$$

hence

$$L_{\text{eucl}}(\Pi(\gamma|_{[0, \tau]} \setminus E)) \geq \frac{1}{2} \rho.$$

Eventually using (22) we obtain:

$$L_{g_f}(\gamma) \geq C e^{f_{\mathbb{B}}} r \rho \geq C (\mu_f(\mathbb{B}_{x, y}))^{\frac{1}{n}}.$$

This concludes the proof of Theorem 5.7.

Let us now show that the previous proof implies that the distances d_f and d_0 are α -bi-Hölder for all $\alpha \in (0, 1)$.

Proposition 5.9. *Let (M, g_0) be a closed Riemannian manifold. If $f: M \rightarrow \mathbb{R}$ is such that*

$$\|df\|_{L^n} \leq \mathbf{I}$$

Then for each $\alpha \in (0, 1)$ there is a constant $C(g_0, \mathbf{I}, \alpha)$ such that for any $x, y \in M$

$$\frac{\mu_f(M)^{\frac{1}{n}}}{C(g_0, \mathbf{I}, \alpha)} d_0^{\frac{1}{\alpha}}(x, y) \leq d_f(x, y) \leq C(g_0, \mathbf{I}, \alpha) \mu_f(M)^{\frac{1}{n}} d_0^\alpha(x, y).$$

Proof. We have shown in Theorem 5.7 that the two weights e^{nf} and e^{-nf} are Strong A_∞ weight. In particular, there is a constant C depending on g_0 and \mathbf{I} such that for all $x, y \in M$:

$$\begin{aligned} \frac{1}{C} d_f^n(x, y) &\leq \mu_f(\mathbb{B}_{x,y}) \leq C d_f^n(x, y) \\ \frac{1}{C} d_{-f}^n(x, y) &\leq \mu_{-f}(\mathbb{B}_{x,y}) \leq C d_{-f}^n(x, y) \end{aligned}$$

The estimate (22) implies that for each $\lambda \in \mathbb{R}$ there is a constant C' depending on g_0, λ and \mathbf{I} such that for any x, y :

$$\frac{1}{C'} d_0^n(x, y) \leq \left(\int_{\mathbb{B}_{x,y}} e^{\lambda f} d\mu_0 \right) \left(\int_{\mathbb{B}_{x,y}} e^{-\lambda f} d\mu_0 \right) \leq C' d_0^n(x, y).$$

Hence, there is a constant Λ such that for any $x, y \in M$:

$$(25) \quad \frac{1}{\Lambda} d_0^{2n}(x, y) \leq d_f^n(x, y) d_{-f}^n(x, y) \leq \Lambda d_0^{2n}(x, y).$$

Notice that this implies that for some constant C :

$$\frac{1}{C} \leq \mu_f(M) \mu_{-f}(M) \leq C.$$

But with the Hölder inequality, we get that for any $p > 1$ and $x, y \in M$:

$$d_f^n(x, y) \leq C d_0^n(x, y) \left(\int_{\mathbb{B}_{x,y}} e^{\frac{p}{p-1} f} d\mu_0 \right)^{1-\frac{1}{p}},$$

and again the estimate (22) implies that

$$\left(\int_{\mathbb{B}_{x,y}} e^{\frac{p}{p-1} f} d\mu_0 \right)^{1-\frac{1}{p}} \leq C \frac{\mu_f(\mathbb{B}_{x,y})}{d_0^n(x, y)} \leq C \frac{\mu_f(M)}{d_0^n(x, y)}.$$

So that we get that for any $x, y \in M$ and any $p > 1$,

$$d_f^n(x, y) \leq C d_0^{\frac{n}{p}}(x, y) \mu_f(M)$$

which is the right-hand side inequality of Proposition 5.9. Moreover, since

$$d_{-f}^n(x, y) \leq C d_0^{\frac{n}{p}}(x, y) \mu_{-f}(M) \leq C d_0^{\frac{n}{p}}(x, y) (\mu_f(M))^{-1},$$

using the comparison (25) we get:

$$d_0^{2n}(x, y) \leq C d_0(x, y)^{\frac{n}{p}} d_f(x, y)^n (\mu_f(M))^{-1}$$

We get therefore for all $p > 1$,

$$d_0^{2n-\frac{n}{p}}(x, y) \leq C d_f^n(x, y) (\mu_f(M))^{-1},$$

which is the left-hand side inequality of Proposition 5.9. \square

6. CONFORMAL DEFORMATIONS OF \mathbb{R}^n WITH SCALAR CURVATURE IN $L^{n/2}$

In this section, we present some interesting application of our techniques to conformal deformations of $(\mathbb{R}^n, \text{eucl})$.

Let us first provide the following sufficient criterion to get strong A_∞ weights.

Theorem 6.1. *Let $f: \mathbb{R}^n \rightarrow \mathbb{R}$ be such that*

$$\|df\|_{L^n(\mathbb{R}^n, \text{eucl})} \leq \mathbf{I}$$

for some $\mathbf{I} > 0$. Then the weight e^{nf} is a strong A_∞ weight with respect to eucl, with constants (θ, η) , where η depends only on g_0 and θ depends only on g_0 and \mathbf{I} .

This slightly improves Theorem 3.1 of [6], since our statement is scale invariant. The proof of this result is an immediate adaptation of our proof of Theorem 5.7. We will not provide further details since it would be a mere repetition of the previous section.

Nevertheless, we show now that Theorem 6.1 implies Theorem 1.6, which provides a completely new criterion to get Strong A_∞ weights on $(\mathbb{R}^n, \text{eucl})$.

Proof of Theorem 1.6. Let $g = e^{2f}\text{eucl}$ be a conformal metric on the Euclidean space $\mathbb{R}^{n>2}$ such that

- i) $\text{Vol}(\mathbb{R}^n, g) = +\infty$;
- ii) $\int_{\mathbb{R}^n} |\text{Scal}_g|^{\frac{n}{2}}(x) \, d\text{vol}_g(x) < +\infty$.

We will show that e^{nf} is a strong A_∞ weight with respect to eucl.

The following lemma shows that our infinite volume assumption is often satisfied.

Lemma 6.2. *A complete conformal deformation $g = e^{2f}\text{eucl}$ of the Euclidean metric on $\mathbb{R}^{n>2}$ that satisfies $\int_{\mathbb{R}^n} |\text{Scal}_g|^{\frac{n}{2}}(x) \, d\text{vol}_g(x) < +\infty$ has necessary infinite volume.*

Proof. According to this hypothesis, the Riemannian manifold (\mathbb{R}^n, g) satisfies the Sobolev inequality ([12, Proposition 2.3]): for some $C > 0$,

$$\forall \varphi \in C_0^1(\mathbb{R}^n): \left(\int |\varphi|^{\frac{2n}{n-2}} \, d\text{vol}_g \right)^{1-\frac{2}{n}} \leq C \int |d\varphi|_g^2 \, d\text{vol}_g.$$

Hence (\mathbb{R}^n, g) has an infinite volume, since the volume of geodesic balls is then bounded from below ([1, 10]):

$$c(n, \mu)r^n \leq \text{Vol}_g(B(x, r)).$$

□

Now, our proof of Theorem 1.6 will be based on the following elementary lemma.

Lemma 6.3. *Let (M, g) be a complete Riemannian manifold that satisfies the elliptic Harnack inequality. Then harmonic functions which are bounded from below are constant.*

Proof. Harnack inequality establishes the existence of a constant $\gamma > 0$ such that for every geodesic ball $B(x, r) \subset M$ and every positive harmonic function h defined over $B(x, 2r)$ one has

$$\sup_{y \in B(x, r)} h(y) \leq \gamma \inf_{y \in B(x, r)} h(y).$$

Hence if $h: M \rightarrow \mathbb{R}$ is bounded from below, setting $\lambda = \inf_{y \in M} h(y)$, we get that for every geodesic ball

$$\sup_{y \in B(x, r)} (h(y) - \lambda) \leq \gamma \inf_{y \in B(x, r)} (h(y) - \lambda).$$

But as $r \rightarrow +\infty$, we have

$$\lim_{r \rightarrow +\infty} \inf_{y \in B(x, r)} h(y) = \lambda$$

therefore for all $y \in M$, $h(y) = \lambda$. \square

As an example that will be useful later, let us consider the case where $\bar{g} = e^{2u} \text{eucl}$ is a conformal metric on the Euclidean space \mathbb{R}^n and assume that

$$\int_{\mathbb{R}^n} |du|^n(x) dx < +\infty.$$

It follows from Theorem 6.1 that e^{nu} is a strong A_∞ weight and that the metric \bar{g} satisfies the Poincaré inequality and the doubling condition: there are constants θ, λ such that for all \bar{g} -geodesic balls $B(x, r)$, we have

$$\text{Vol}_{\bar{g}} B(x, 2r) \leq \theta \text{Vol}_{\bar{g}} B(x, r)$$

and, writing $\varphi_{B(x, r)} := \int_{B(x, r)} \varphi \, d\text{vol}_{\bar{g}}$,

$$\forall \varphi \in C^1(B(x, 2r)): \|\varphi - \varphi_{B(x, r)}\|_{L^2(B(x, r), \bar{g})}^2 \leq \|\nabla \varphi\|_{L^2(B(x, 2r), \bar{g})}^2.$$

Moreover, by the results of Section 4.1, the metric space $(\mathbb{R}^n, \bar{g}, d\text{vol}_{\bar{g}})$ is Alhfors regular: there is a constant η such that for every \bar{g} -geodesic ball $B(x, r)$:

$$\eta r^n \leq \text{Vol}_{\bar{g}} B(x, r) \leq r^n / \eta.$$

In particular by result in [21, 32], \bar{g} satisfies the parabolic and the elliptic Harnack inequalities and if $n \geq 3$, then (\mathbb{R}^n, \bar{g}) is a non parabolic manifold and has a positive Green kernel $G_{\bar{g}}(x, y)$ that satisfies the estimates

$$(26) \quad cd_{\bar{g}}^{2-n}(x, y) \leq G_{\bar{g}}(x, y) \leq Cd_{\bar{g}}^{2-n}(x, y)$$

or equivalently:

$$c (\text{Vol}_{\bar{g}}(\mathbb{B}_{x, y}))^{-1 + \frac{2}{n}} \leq G_{\bar{g}}(x, y) \leq C (\text{Vol}_{\bar{g}}(\mathbb{B}_{x, y}))^{-1 + \frac{2}{n}}.$$

Let us go back to our metric $g = e^{2f} \text{eucl}$ with

$$\text{Vol}(\mathbb{R}^n, g) = +\infty \quad \text{and} \quad \int_{\mathbb{R}^n} |\text{Scal}_g|^{\frac{n}{2}}(x) \, d\text{vol}_g(x) < +\infty.$$

Let Δ be the Euclidean Laplacian and let A_n be the norm of the operator $d\Delta^{-1}: L^{\frac{n}{2}} \rightarrow L^n$. Since the $L^{n/2}$ norm of Scal_g is finite, for R large enough we have

$$\frac{n-2}{4(n-1)} \left(\int_{\mathbb{R}^n \setminus \mathbb{B}(R)} |\text{Scal}_g|^{\frac{n}{2}}(x) \, d\text{vol}_g(x) \right)^{\frac{2}{n}} < \frac{1}{8A_n}.$$

Therefore, as was done in the proof of Proposition 5.4, we can find $v \in L^{\frac{n}{2}}(\mathbb{R}^n, dx)$ such that

$$v - |d\Delta^{-1}v|^2 = \frac{n-2}{4(n-1)} \text{Scal}_g e^{2f} \mathbf{1}_{\mathbb{R}^n \setminus \mathbb{B}(R)}.$$

Now, let $u \in W_{\text{loc}}^{2, \frac{n}{2}}(\mathbb{R}^n)$ be such that $\Delta u = v$, for instance u is a primitive of $d\Delta^{-1}v \in L^n$ or

$$u(x) = c_n \int_{\mathbb{R}^n} \left(\frac{1}{\|x-y\|^{n-2}} - \frac{1}{\|y\|^{n-2}} \right) v(y) dy.$$

We have

$$\Delta e^u = \frac{n-2}{4(n-1)} \text{Scal}_g e^{2f} \mathbf{1}_{\mathbb{R}^n \setminus \mathbb{B}(R)} e^u.$$

Writing $\phi = e^{\frac{n-2}{2}f}$, Yamabe equation (6) implies that

$$\Delta\phi = \frac{n-2}{4(n-1)} \text{Scal}_g e^{2f} \phi.$$

Hence if we define $\psi = \phi e^{-u}$ we get

$$\Delta\psi - \langle du, d\psi \rangle = \frac{n-2}{4(n-1)} \text{Scal}_g e^{2f} \mathbf{1}_{\mathbb{B}(R)} \psi.$$

As $du \in L^n$, the metric $\bar{g} = e^{\frac{4}{n-2}u} \text{eucl}$ induces a strong A_∞ weight, but

$$\Delta_{\bar{g}}\psi = e^{-\frac{4}{n-2}u} (\Delta\psi - \langle du, d\psi \rangle) = \frac{n-2}{4(n-1)} \text{Scal}_g e^{2f} \mathbf{1}_{\mathbb{B}(R)} e^{-\frac{4}{n-2}u} \psi.$$

As $\Delta_{\bar{g}}\psi$ has compact support and is in L^p for every $1 \leq p < +\infty$, we can find a solution of the equation

$$\Delta_{\bar{g}}\xi = \Delta_{\bar{g}}\psi,$$

for instance let us choose

$$\xi(x) = \int_{\mathbb{R}^n} G_{\bar{g}}(x, y) \Delta_{\bar{g}}\psi(y) d\text{vol}_{\bar{g}}(y).$$

Since $\text{supp } \Delta_{\bar{g}}\psi \subset K$, it follows from (26) that there exists $C > 0$ such that for all $x \in M \setminus K$,

$$\xi(x) \leq C \left(\int_K G_{\bar{g}}^q(x, y) d\text{vol}_{\bar{g}}(y) \right)^{\frac{1}{q}} \leq C d_{\bar{g}}(x, K)^{2-n}.$$

In particular, $\int_{\mathbb{R}^n} |\xi(y)|^{\frac{2n}{n-2}} d\text{vol}_{\bar{g}}(y) < +\infty$, the map ξ tends to zero at infinity and is in $W_{\text{loc}}^{2,p}(\mathbb{R}^n)$ for every $1 \leq p < +\infty$. Therefore ξ is a bounded function and $\psi - \xi$ is a \bar{g} -harmonic function that is bounded from below: it is hence a constant function. If this constant is 0 then $\psi = \xi$, and

$$\text{Vol}(\mathbb{R}^n, g) = \int_{\mathbb{R}^n} \psi(y)^{\frac{2n}{n-2}} d\text{vol}_{\bar{g}}(y) = \int_{\mathbb{R}^n} \xi^{\frac{2n}{n-2}}(y) dy < +\infty.$$

This is contradictory with our hypothesis hence there is a positive constant c such that

$$\psi = c + \phi,$$

hence there are positive constants α, A such that

$$\alpha \leq \psi \leq A.$$

This implies that the metric $g = \phi^{\frac{4}{n-2}} \text{eucl}$ and $\bar{g} = e^{2u} \text{eucl}$ are bi-Lipschitz, hence since e^{nu} is a strong A_∞ weight with respect to g_{eucl} , $e^{nf} = \phi^{\frac{2n}{n-2}}$ is also. \square

We emphasize again on the fact that this implies very strong geometric constraints on conformal deformations of $(\mathbb{R}^n, g_{\text{eucl}})$ with $\int_{\mathbb{R}^n} |\text{Scal}_g|^{\frac{n}{2}}(x) d\text{vol}_g(x) < +\infty$ and $\text{Vol}_g(\mathbb{R}^n, g) = +\infty$. It implies in particular Poincaré inequality, Euclidean-type isoperimetric inequality and Sobolev inequalities, which were unknown on such examples.

7. GROMOV-HAUSDORFF LIMIT AND CONFORMAL DISTANCES

Let us conclude by presenting a natural question about the limiting distances which arise in our Theorem 1.3, which we could not answer.

Let us first recall that to any strong A_∞ weight e^{nf_∞} (with respect to $d\text{vol}_{g_0}$) is naturally associated a distance d_f , defined as follows. For all $x, y \in M$, we write again \mathbb{B}_{xy} the largest g_0 -geodesic ball whose diameter is the g_0 -geodesic segment $[xy]$. Let us write

$$\delta_f(x, y) = \left(\frac{1}{\omega_n} \mu_f(\mathbb{B}_{xy}) \right)^{1/n} = \left(\frac{1}{\omega_n} \int_{\mathbb{B}_{xy}} e^{nf(x)} d\text{vol}_{g_0}(x) \right)^{1/n},$$

where ω_n is the Euclidean volume of the unit n -ball, and define

$$(27) \quad d_f(x, y) = \liminf_{\varepsilon \rightarrow 0} \left\{ \sum_{i=0}^N \delta_f(x_i, x_{i+1}) ; x_0 = x, x_{N+1} = y, d_0(x_i, x_{i+1}) \leq \varepsilon \right\}.$$

It can be easily shown that $e^{nf} \in L^1(M)$ is a strong A_∞ weight if and only if d_f is a distance (see [33] for more details). Moreover, an elementary approximation argument shows that if f is continuous, then d_f coincides with the classical Riemannian distance associated to the Riemannian metric $g_f = e^{2f} g_0$.

Let us now consider a sequence $(g_k = e^{2f_k} g_0)_{k \in \mathbb{N}}$ of conformal metrics satisfying the hypotheses of Corollary 1.4. Up to extracting a subsequence, the sequence of associated metric space (M, d_{g_k}) converges in Gromov-Hausdorff topology to some distance d_∞ on M , which is bi-Hölder equivalent to d_0 . Moreover, still up to extracting a subsequence, the Riemannian volume measure $\mu_k = d\text{vol}_{g_k} = e^{nf_k} d\text{vol}_{g_0}$ converge in the weak- $*$ topology to a positive measure μ_∞ on M . Since all the μ_k are uniformly strong A_∞ weights with respects to $d\text{vol}_{g_0}$, μ_∞ is also: there exists hence $e^{nf_\infty} \in L^1(M)$ which is still a strong A_∞ weight with respect to g_0 . This weight is therefore associated to a distance d_{f_∞} defined by (27). *Does d_∞ and d_{f_∞} coincide ?*

The reader will can check that in the Example 2 presented in our introduction, where the conformal factors converge in $W^{2, \frac{n}{2}}$ to the singular conformal factor given by 1) and the distances converge in the Gromov-Hausdorff topology, these distances indeed coincide.

One can easily show from semi-continuity arguments that $d_\infty \leq d_{f_\infty}$. Let us present now a striking example of pathological behaviour which shows that in general, for a sequence of (unifom) strong A_∞ weights $(e^{nf_k})_{k \in \mathbb{N}}$, the converse inequality is not true. The following examples of conformal deformations is due to D. Burago, in [8]

Let us consider the standard flat torus $(\mathbb{T}^n = \mathbb{R}^n / \Gamma, \text{eucl})$, where $\Gamma = (2\pi\mathbb{Z})^n$. For all $\ell > 1$, we consider the conformal deformation of the euclidean metric g_0 given by $g_\ell = e^{2f_\ell} \text{eucl}$, where

$$e^{nf_\ell}(x_1, \dots, x_n) = 1 - \frac{1}{2} \cos(\ell x_1).$$

Since e^{nf_ℓ} is uniformly bounded from above and below, all the metric space $(\mathbb{T}^n, d_\ell = d_{f_\ell})$ are uniformly bilipschitz to \mathbb{T}^n equiped with the Euclidean distance and the $(e^{nf_\ell})_{\ell \geq 1}$ are uniformly strong A_∞ weights with respect to the Lebesgue metric. Nevertheless, the distance associated to the limit of the volume measures and the limit of distances are radically distinct as shows the following result.

Theorem 7.1. *Under the above notations, as $\ell \rightarrow +\infty$,*

- *the sequence $(e^{nf_\ell})_\ell$ converges weakly in $L^1(\mathbb{T})$ to the function 1;*

- the sequence of metric spaces (\mathbb{T}^n, d_ℓ) , $\ell \in \mathbb{N}$ converges in the Gromov-Hausdorff topology to metric space (\mathbb{T}^n, d_∞) where d_∞ is the Finsler metric associated to the stable norm of the periodic metric on \mathbb{R}^n given by

$$\bar{g} = \left(1 - \frac{1}{2} \cos(x_1)\right)^{\frac{2}{n}} \text{ eucl.}$$

Proof. The first assertion is classical. As for the second assertion, let us recall that if \bar{d} is the Riemannian distance on \mathbb{R}^n associated to the metric \bar{g} then the associated stable norm $\|\bullet\|_*$ is defined for all $x \in \mathbb{R}^n$ by $\|x\|_* = \lim_{t \rightarrow +\infty} \frac{1}{t} \bar{d}(0, tx)$. Using the change of variable $\xi_i = \ell x_i$, $i = 1 \dots n$, we get that $(\mathbb{T}^n, d_\ell = d_{f_\ell})$ is isometric to the quotient $\mathbb{R}^n / (\ell\Gamma)$ with the metric $\frac{1}{\ell^2} \bar{g}$. More exactly let $\pi: \mathbb{R}^n \rightarrow \mathbb{T}^n = \mathbb{R}^n / \Gamma$ the quotient map. if $x, y \in \mathbb{R}^n$ then

$$d_\ell(\pi(x), \pi(y)) = \inf_{\gamma \in \Gamma} \frac{1}{\ell} d(x, y + \ell\gamma)$$

If $d_\infty(\pi(x), \pi(y)) = \inf_{\gamma \in \ell\Gamma} \|x - y - \gamma\|_*$, then we clearly have

$$\sup_{x, y \in \mathbb{R}^n} |d_\ell(\pi(x), \pi(y)) - d_\infty(\pi(x), \pi(y))| \leq \frac{C}{\ell}.$$

Hence the result. \square

We emphasize on the fact that in this example, $d_{f_\infty} = d_{eucl}$, whereas d_∞ is a Finsler non-Riemannian metric: it cannot in any weak sense be considered as a conformal metric for some limiting conformal factor.

The reader will easily notice than in the previous example, the volume of the conformal metrics are uniformly bounded, but the $L^{n/2}$ -norm of the scalar curvature blows up. Under the $L^{n/2}$ assumptions on the scalar curvature which ensures Theorem 1.3 and Corollary 1.4, we have a stronger control on the geometrical and analytical behaviour of the conformal factors than the information given by the strong A_∞ bounds. Nevertheless, we cannot show yet equality for d_∞ and d_{f_∞} under these hypotheses, nor give a counterexample.

REFERENCES

- [1] K. Akutagawa : Yamabe metrics of positive scalar curvature and conformally flat manifolds, *Differ. Geom. Appl.* **4** (1994) 4, 239–258.
- [2] T. Aubin, Problèmes isopérimétriques et espace de Sobolev, *J. Differential Geometry* **11** (1976), no. 4, 573–598.
- [3] D. Bakry, Étude des transformations de Riesz dans les variétés riemanniennes à courbure de Ricci minorée, in *Séminaire de Probabilités, XXI*, Lecture Notes in Math., vol. 1247, Springer, Berlin, 1987, p. 137–172.
- [4] A.L. Besse, *Einstein manifolds*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3), 10. Springer-Verlag, Berlin, 1987
- [5] M. Bonk, J. Heinonen, E. Saksman, Logarithmic potentials, quasiconformal flows, and Q-curvature, *Duke Math. J.* **142** (2) (2008) 197–239.
- [6] M. Bonk, J. Heinonen, E. Saksman, The quasiconformal Jacobian problem, *In the tradition of Ahlfors and Bers*, III, 77–96, Contemp. Math., 355, Amer. Math. Soc., Providence, RI, 2004
- [7] S. Brendle, Global existence and convergence for a higher order flow in conformal geometry, *Ann. of Math. (2)* **158** (2003), 323–343.
- [8] D. Burago, Periodic metrics, *Representation theory and dynamical systems*, 205–210, Adv. Soviet Math. 9 (1992)
- [9] D. Burago, Dmitri, Y. Burago, S. Ivanov, *A course in metric geometry*, Graduate Studies in Mathematics 33., American Math. Soc., Providence, RI, 2001
- [10] G. Carron : Inégalités isopérimétriques de FaberKrahn et conséquences, in: *Actes de la Table Ronde de Géométrie Différentielle (Luminy, 1992)*, Sémin. Congr., vol. 1, Soc. Math. France, Paris, 1996, 205–232.

- [11] G. Carron, Geometric inequalities for manifolds with Ricci curvature in the Kato class, arXiv:1612.03027
- [12] C. Carron and M. Herzlich: The Huber theorem for non-compact conformally flat manifolds, *Comment Math. Helv.* **77** (2002), 192–220.
- [13] S.-Y. A. Chang, M. Gursky, T. Wolff, Lack of compactness in conformal metrics with $L^{d/2}$ curvature, *J. Geom. Anal.* **4** (1994), no. 2, 143–153.
- [14] T. H. Colding, W. P. Minicozzi, II. Liouville theorems for harmonic sections and applications. *Comm. Pure Appl. Math.* **51** (1998), no. 2, 113–138.
- [15] S. Costea, Strong A_∞ -weights and scaling invariant Besov capacities, *Rev. Mat. Iberoam.* **23** (2007), no. 3, 1067–1114.
- [16] A. Chang, M. Gursky, T. Wolff, Lack of compactness in conformal metrics with $L^{d/2}$ curvature, *J. Geom. Anal.* **4** (1994), no. 2, 143–153.
- [17] G. David, S. Semmes, Strong A_∞ weights, Sobolev inequalities and quasiconformal mappings, in: *Analysis and Partial Differential Equations*, Lect. Notes Pure Appl. Math., vol. 122, Dekker, New York, 1990, pp. 101–111.
- [18] C. Fefferman, C. Graham, Q-curvature and Poincaré metrics. *Math. Res. Lett.* **9** (2002), no. 2-3, 139–151
- [19] S. Gallot, Inégalités isopérimétriques et analytiques sur les variétés riemanniennes, in *On the geometry of differentiable manifolds (Rome, 1986)*, Astérisque 163-164 (1988)
- [20] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Grundlehr. Math. Wiss., Vol. **224**, Springer, Berlin, 1977.
- [21] A. Grigor’yan: The heat equation on noncompact Riemannian manifolds (Russian). *Mat. Sb.* **182** (1991), no. 1, 55–87; translation in *Math. USSR-Sb.* **72** (1992), no. 1, 47–77.
- [22] A. Grigor’yan, L. Saloff-Coste, Stability results for Harnack inequalities, *Ann. Inst. Fourier* **55** (2005), 825–890.
- [23] M. Gursky, Compactness of conformal metrics with integral bounds on curvature, *Duke Math. J.* **72** (1993), 339–3671.
- [24] E. Hebey, M. Vaugon, Meilleures constantes dans le théorème d’inclusion de Sobolev. *Ann. Inst. H. Poincaré. Anal. Non Linéaire* **13** (1996) 57–93.
- [25] L. Hörmander, *The analysis of linear partial differential operator III*, Springer-Verlag, Berlin, 1985.
- [26] O.-E. Kansanen, R. Korte, Strong A_∞ -weights are A_∞ -weights on metric spaces. *Rev. Mat. Iberoam.* **27** (2011), no. 1, 335–354.
- [27] O. A. Ladyzhenskaya and N. N. Ural’tseva, Estimates for the Hölder constant for functions satisfying a uniformly elliptic or a uniformly parabolic quasilinear inequality with unbounded coefficients, *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov (LOMI)* **147** (1985), 72–94. [In Russian; English transl. in *J. Soviet Math.* **37** (1987), 837–851.]
- [28] P. Li, R. Schoen, L^p and mean value properties of subharmonic functions on Riemannian manifolds, *Acta Math.* **153** (1984) 279–301.
- [29] H. Matthiesen, Regularity of conformal metrics with large first eigenvalue, *Annales de la faculté des sciences de Toulouse*, **25** no. 5 (2016), 1079–1094
- [30] M.V. Safonov, Non-divergence Elliptic Equations of Second Order with Unbounded Drift, *Amer. Math. Soc. Transl. Ser. 2*, vol. 229, 2010, pp. 211–232.
- [31] T. Sakai, *Riemannian Geometry*, Translations of Mathematical Monographs **Vol 149** (1997).
- [32] L. Saloff-Coste: A note on Poincaré, Sobolev and Harnack inequalities. *Internat. Math. Res. Notices* **2** (1992), no. 2, 27–38.
- [33] S. Semmes, Bi-Lipschitz mappings and strong A_∞ weights, *Ann. Acad. Sci. Fenn. Ser. A I Math.* **18** (1993) no2, 211–248.
- [34] S. Semmes, On the nonexistence of bi-Lipschitz parameterizations and geometric problems about A_∞ -weights, *Rev. Mat. Iberoam.* **12** (1996) no2, 337–410.
- [35] S. Semmes, Some remarks about metric spaces, spherical mappings, functions and their derivatives. *Publ. Mat.* **40** (1996), 411–430.
- [36] E. M. Stein, *Harmonic Analysis : Real-variable methods, orthogonality and oscillatory integrals*, Princeton Univ. Press (1993).
- [37] J-O. Strömberg, A. Torchinsky, *Weighted Hardy spaces*. Lecture Notes in Mathematics 1381. Springer-Verlag, Berlin, 1989.
- [38] R. Tessera, Volumes of spheres in doubling measures metric spaces and groups of polynomial growth. *Bull. Soc. Math. France* **135** (2007), no. 1, 47–64.
- [39] A. Torchinsky, *Real-variable methods in harmonic analysis*, Pure and Applied Mathematics, vol. 123, Academic Press, Inc., Orlando, FL, 1986.

- [40] N. Trudinger, On imbeddings into Orlicz spaces and some applications. *J. Math. Mech.* **17** (1967) 473–483.
- [41] N. Trudinger, Remarks concerning the conformal deformation of Riemannian structures on compact manifolds, *Ann. Scuola Norm. Sup. Pisa* (3) **22** (1968) 265–274.
- [42] Yi Wang, The isoperimetric inequality and Q-curvature. *Adv. Math.* **281** (2015), 823–844.
- [43] H. Yamabe, On a deformation of Riemannian structures on compact manifolds. *Osaka Math. J.* 12 1960 21–37

E-mail address: clara.aldana@posteo.net

E-mail address: gilles.carron@univ-nantes.fr

E-mail address: samuel.tapie@univ-nantes.fr