



HAL
open science

COMPUTING THE GALOIS GROUP OF SOME PARAMETERIZED LINEAR DIFFERENTIAL EQUATION OF ORDER TWO

Thomas Dreyfus

► **To cite this version:**

Thomas Dreyfus. COMPUTING THE GALOIS GROUP OF SOME PARAMETERIZED LINEAR DIFFERENTIAL EQUATION OF ORDER TWO. Proceedings of the American Mathematical Society, 2014, 142 (4), pp.1193-1207. 10.1090/S0002-9939-2014-11826-0 . hal-01897284

HAL Id: hal-01897284

<https://hal.science/hal-01897284>

Submitted on 17 Oct 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

COMPUTING THE GALOIS GROUP OF SOME PARAMETERIZED LINEAR DIFFERENTIAL EQUATION OF ORDER TWO

THOMAS DREYFUS

ABSTRACT. We extend Kovacic's algorithm to compute the differential Galois group of some second order parameterized linear differential equation. In the case where no Liouvillian solutions could be found, we give a necessary and sufficient condition for the integrability of the system. We give various examples of computation.

CONTENTS

Introduction	1
1. Parameterized differential Galois theory	3
2. Computation of the parameterized differential Galois group	4
3. Examples	12
References	14

INTRODUCTION

Let us consider the linear differential equation

$$\begin{pmatrix} \partial_z Y(z) \\ \partial_z^2 Y(z) \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ r(z) & 0 \end{pmatrix} \begin{pmatrix} Y(z) \\ \partial_z Y(z) \end{pmatrix},$$

where $r(z)$ is a rational function with coefficients in \mathbb{C} . We have a Galois theory for this type of equation; see [vdPS03]. In particular, we can associate to this equation a group H , which we call the differential Galois group, that measures the algebraic relations of the solutions. In this case, this group can be viewed as a linear algebraic subgroup of $\mathrm{SL}_2(\mathbb{C})$. Kovacic in [Kov86] (see also [vdP99]) uses the classification of the linear algebraic subgroup of $\mathrm{SL}_2(\mathbb{C})$ to obtain an algorithm that determines the Liouvillian solutions, which are the solutions that involve exponentials, indefinite integrals and solutions of polynomial equations. In particular, four cases happen:

- (1) H is conjugated to a subgroup of $B = \left\{ \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix}, \text{ where } a \in \mathbb{C}^*, b \in \mathbb{C} \right\}$ and

there exists a Liouvillian solution of the form $e^{\int_0^z f(u)du}$, with $f(z) \in \mathbb{C}(z)$.

- (2) H is conjugated to a subgroup of

$$D_\infty = \left\{ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \cup \begin{pmatrix} 0 & b^{-1} \\ -b & 0 \end{pmatrix}, \text{ where } a, b \in \mathbb{C}^* \right\}$$

and there exists a Liouvillian solution of the form $e^{\int_0^z f(u)du}$, where $f(z)$ is algebraic over $\mathbb{C}(z)$ of degree two and $f(z) \notin \mathbb{C}(z)$.

- (3) H is finite and all the solutions are algebraic over $\mathbb{C}(z)$.
 (4) $H = \mathrm{SL}_2(\mathbb{C})$ and there are no Liouvillian solutions.

Date: September 13, 2013.

2010 *Mathematics Subject Classification.* Primary 34M15, 12H20, 34M03.

Work partially supported by ANR-06-JCJC-0028 and NFS CCF-0952591.

Various improvements of this algorithm have been made. See for example [DLR92, HvdP95, UW96, Zha95]. The case where H is finite has been totally solved in [SU93a, SU93b]; see also [vHW05].

Let $\{\partial_0, \partial_1, \dots, \partial_n\}$ be a set of $n + 1$ commuting derivations. In this article, we are interested in the parameterized linear differential equation of the form

$$\begin{pmatrix} \partial_0 Y \\ \partial_0^2 Y \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ r & 0 \end{pmatrix} \begin{pmatrix} Y \\ \partial_0 Y \end{pmatrix},$$

where r belongs in a suitable $(\partial_0, \partial_1, \dots, \partial_n)$ -differential field. The derivations $\partial_1, \dots, \partial_n$ should be thought of as derivations with respect to the parameters. We will denote by C its field of the ∂_0 -constants. In [Lan08] and [CS07, HS08], the authors develop a Galois theory for the parameterized linear differential equations. They define a parameterized differential Galois group that measures the $(\partial_1, \dots, \partial_n)$ -differential and algebraic relations between the solutions; see §1. This group can be seen as a differential group in the sense of Kolchin: this is a group of matrices whose entries lie in the differential field C and satisfy a set of polynomial differential equations with coefficients in C . In the case of the equation $\partial_0^2 Y = rY$, the Galois group will be a linear differential algebraic subgroup of $\mathrm{SL}_2(C)$. The goal of this paper is to extend the algorithm from Kovacic and compute the parameterized differential Galois group of the equation $\partial_0^2 Y = rY$.

The article is presented as follows. In the first section, we recall some basic facts about parameterized differential Galois theory. This theory needs to use a field of the ∂_0 -constants which is $(\partial_1, \dots, \partial_n)$ -differentially closed (see [CS07], Definition 3.2). We will make a stronger assumption on the field of the ∂_0 -constants C : we will assume that C is a universal $(\partial_1, \dots, \partial_n)$ -field (see §1). We use this assumption on C because a field $(\partial_1, \dots, \partial_n)$ -differentially closed is an abstract field which has no interpretation as a field of functions. We will see in §2 that a result of Seidenberg will allow us to identify the elements of the universal $(\partial_1, \dots, \partial_n)$ -field C which we will consider as meromorphic functions on a polydisc D of \mathbb{C}^n .

In the second section, we recall the result of Seidenberg which implies that the parameterized differential Galois group can be seen as a linear differential algebraic subgroup defined over a field of meromorphic functions on a polydisc D of \mathbb{C}^n . Since the original algorithm from [Kov86] can be applied if we consider rational functions having coefficients in an algebraically closed field, we apply Kovacic's algorithm for the field of rational functions having coefficients in C . We obtain Liouvillian solutions that can be interpreted as meromorphic functions. Then we explain how to compute the Galois group in the four cases of Kovacic's algorithm. In the case number 4, the Galois group is Zariski dense in SL_2 . We recall the definition of integrable systems and the link with integrable systems and equations with a Galois group that is Zariski dense in SL_2 . We decrease the number of integrability conditions by showing that this is enough to check the integrability condition for the pairs of derivations (∂_z, ∂) , where ∂ belongs in the vectorial space spanned by the derivations with respect to the parameters. Then, we obtain an effective way to compute the Galois group in the case number 4, see Proposition 2.8. We summarize the results of the section in Theorem 2.10.

In the last section we give various examples of computation.

After the submission of this paper, the authors in [GO12] has generalized Proposition 2.8 for equations with order more than two. Moreover, Carlos E Arreche has proved some other results in touch with parameterized Kovacic's algorithm. See [Arr12].

Acknowledgments. This article has been prepared during my thesis, which is supported by the region Île-de-France. I would like to thank my advisor, Lucia Di Vizio for sharing with me a part of her knowledge. I also want to thank Jacques-Arthur Weil and William Sit, for pointing me out some mistakes and inaccuracies in this paper. Thanks to the anonymous referee for the pertinent remarks.

1. PARAMETERIZED DIFFERENTIAL GALOIS THEORY

Let K be a differential field equipped with $n + 1$ commuting derivations $\partial_0, \dots, \partial_n$ and let $\Delta = \{\partial_1, \dots, \partial_n\}$. We will assume that its field of the ∂_0 -constants C is a universal (Δ) -field with characteristic 0; that is, a (Δ) -field such that for any (Δ) -field $C_0 \subset C$, (Δ) -finitely generated over \mathbb{Q} , and any (Δ) -finitely generated extension C_1 of C_0 , there is a (Δ) -differential C_0 -isomorphism of C_1 into C . See [Kol73], Chapter 3, Section 7, for more details. In particular, C is (Δ) -differentially closed. In this section, we will recall the result from [CS07] of Galois theory for the parameterized linear differential equation of the form

$$(1.1) \quad \begin{pmatrix} \partial_0 Y \\ \partial_0^2 Y \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ r & 0 \end{pmatrix} \begin{pmatrix} Y \\ \partial_0 Y \end{pmatrix},$$

with $r \in K$. A parameterized Picard-Vessiot extension of the equation (1.1) on K is a $(\partial_0, \dots, \partial_n)$ -differential field extension $\mathcal{K}|K$ generated over K by the entries of an invertible solution matrix (we will call it a fundamental solution) and such that the field of the ∂_0 -constants of \mathcal{K} is equal to C . We can apply [CS07], Theorem 9.5, for the equation (1.1), and deduce the existence and the uniqueness up to $(\partial_0, \dots, \partial_n)$ -differential isomorphism of the parameterized Picard-Vessiot extension $\mathcal{K}|K$. If $\Delta = \emptyset$, we recover the usual unparameterized Picard-Vessiot extension.

The parameterized (resp. unparameterized) differential Galois group G (resp. H) is the group of field automorphisms of the parameterized Picard-Vessiot extension (resp. the unparameterized Picard-Vessiot extension) of the equation (1.1), which induces the identity on K and commutes with all the derivations (resp. with the derivation ∂_0). Let U be a fundamental solution. In the unparameterized case,

$$\{U^{-1}\varphi(U), \varphi \in H\}$$

is a linear algebraic subgroup of $\mathrm{GL}_2(C)$. In the parameterized case we find that

$$\{U^{-1}\varphi(U), \varphi \in G\}$$

is a linear differential algebraic subgroup, that is, a subgroup of $\mathrm{GL}_2(C)$ which is the zero of a set of (Δ) -differential polynomials in 4 variables. See [CS07], Theorem 9.10, for a proof. Any other fundamental solution yields another differential algebraic subgroup of $\mathrm{GL}_2(C)$ which are all conjugated over $\mathrm{GL}_2(C)$. We will identify G (resp. H) with a linear differential algebraic subgroup of $\mathrm{GL}_2(C)$ (resp. with a linear algebraic subgroup of $\mathrm{GL}_2(C)$) for a chosen fundamental solution. The next lemma is a classical result.

Lemma 1.1 ([Kov86], Section 1.3). $G \subset \mathrm{SL}_2(C)$.

2. COMPUTATION OF THE PARAMETERIZED DIFFERENTIAL GALOIS GROUP

Until the end of the paper, C denotes a universal (Δ) -field equipped with n commuting derivations. Let $C(z)$ be the (∂_z, Δ) -differential field of rational functions in the indeterminate z , with coefficients in C , where z is a (Δ) -constant with $\partial_z z = 1$, C is the field of the ∂_z -constants and such that ∂_z commutes with all the derivations. Let us consider the parameterized linear differential equation

$$(2.1) \quad \begin{pmatrix} \partial_z Y(z) \\ \partial_z^2 Y(z) \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ r(z) & 0 \end{pmatrix} \begin{pmatrix} Y(z) \\ \partial_z Y(z) \end{pmatrix},$$

with $r(z) \in C(z)$. We want to apply Kovacic's algorithm for the parameterized linear differential equation (2.1). Let $G \subset \mathrm{SL}_2(C)$ be the parameterized differential Galois group. The algorithm from [Kov86] can be applied if the field of the ∂_z -constants is algebraically closed, which is the case here. The problem is that C is an abstract field which is not very convenient for the computations. In fact we have an interpretation of the elements of C as meromorphic functions. Let C_1 be the (Δ) -differential field generated over \mathbb{Q} by the z -coefficients of $r(z)$. Using the following result of Seidenberg (see [Sei58, Sei69]) with $K_0 = \mathbb{Q}$ and $K_1 = C_1$, we find the existence of a polydisc D of \mathbb{C}^n such that the z -coefficients of $r(z)$ can be considered as meromorphic functions on D .

Theorem 2.1 (Seidenberg). *Let $\mathbb{Q} \subset K_0 \subset K_1$ be finitely generated (Δ) -differential extensions of \mathbb{Q} and assume that K_0 consists of meromorphic functions on some domain Ω of \mathbb{C}^n . Then K_1 is isomorphic to the field K_1^* of meromorphic functions on $\Omega_1 \subset \Omega$ such that $K_0|_{\Omega_1} \subset K_1^*$, and the derivations in Δ can be identified with the derivations with respect to the coordinates on Ω_1 .*

Let $(\mathcal{M}_D, \partial_{t_1}, \dots, \partial_{t_n})$ denotes the $\Delta_t = \{\partial_{t_1}, \dots, \partial_{t_n}\}$ -differential field of meromorphic functions on D , a polydisc of \mathbb{C}^n . Let $t = (t_1, \dots, t_n)$. The discussion above tell us that the $r(z)$ of the equation (2.1) can be identified with $r(z, t)$, an element of $\mathcal{M}_D(z)^*$, where D is a polydisc of \mathbb{C}^n . We will consider the parameterized linear differential equation

$$\begin{pmatrix} \partial_z Y(z, t) \\ \partial_z^2 Y(z, t) \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ r(z, t) & 0 \end{pmatrix} \begin{pmatrix} Y(z, t) \\ \partial_z Y(z, t) \end{pmatrix},$$

with $r(z, t) \in \mathcal{M}_D(z)$. The group G is defined by a finite number of (Δ) -differential polynomials. Again, using the result of Seidenberg with the (Δ) -differential field generated over \mathbb{Q} by the coefficients of the (Δ) -differential polynomials that define G and the z -coefficients of $r(z)$, we deduce that G can be seen as a linear differential algebraic subgroup of $\mathrm{SL}_2(\mathcal{M}_D)$. Again using the result of Seidenberg, we remark that after shrinking D , we can assume that if G is conjugated over $\mathrm{SL}_2(C)$ to Q , we can identify Q and G as linear differential algebraic subgroups of $\mathrm{SL}_2(\mathcal{M}_D)$, and they are conjugated over $\mathrm{SL}_2(\mathcal{M}_D)$. Furthermore, we obtain that the Liouvillian solutions found are defined over the algebraic closure of $\mathcal{M}_D(z)$. We will compute G as a linear differential algebraic subgroup of $\mathrm{SL}_2(\mathcal{M}_D)$. We recall that we have four cases to consider:

- (1) There exists a Liouvillian solution of the form $g(z, t) = e^{\int_0^z f(u, t) du}$, with $f(z, t)$ belongs to $\mathcal{M}_D(z)$.
- (2) There exists a Liouvillian solution of the form $g(z, t) = e^{\int_0^z f(u, t) du}$, where $f(z, t)$ is algebraic over $\mathcal{M}_D(z)$ of degree two and $f(z, t) \notin \mathcal{M}_D(z)$.
- (3) All the solutions are algebraic over $\mathcal{M}_D(z)$.
- (4) There are no Liouvillian solutions.

* $\mathcal{M}_D(z)$ denotes the (∂_z, Δ_t) -differential field of rational functions with indeterminate z and with coefficients in \mathcal{M}_D such that $\partial_z z = 1$, z is a (Δ_t) -constant and the field \mathcal{M}_D is the field of the ∂_z -constants.

COMPUTING THE GALOIS GROUP OF SOME PARAMETERIZED
LINEAR DIFFERENTIAL EQUATION OF ORDER TWO

These correspond to the four cases recalled in the introduction. Proposition 6.26 of [HS08] says that if we take the same fundamental solution, the Zariski closure of G is the unparameterized differential Galois group. This means that in each case we are looking at the Zariski dense subgroups of the group given by usual Kovacic's algorithm.

2.1. We start with the case number 1. There exists a Liouvillian solution of the form:

$$g(z, t) = e^{\int_0^z f(u, t) du},$$

with $f(z, t) \in \mathcal{M}_D(z)$. The action of G on the solution $g(z, t)$ can be computed with the following lemma:

Lemma 2.2. *Let $\sigma \in G$.*

- (1) *Let $\alpha(t) \in \mathcal{M}_D$ and $p, q \in \mathbb{N}$, such that $\text{GCD}(p, q) = 1$. Then there exists $k \in \mathbb{N}$ such that $\sigma((z - \alpha(t))^{p/q}) = e^{\frac{2ik\pi}{q}}(z - \alpha(t))^{p/q}$.*
- (2) *Let $\alpha(t), \beta(t) \in \mathcal{M}_D$ and $\beta(t) \notin \mathbb{Q}$. Then there exists $a \in \mathbb{C}$ and $c \in \mathbb{C}^*$ such that $\sigma((z - \alpha(t))^{\beta(t)}) = ce^{a\beta(t)}(z - \alpha(t))^{\beta(t)}$.*
- (3) *Let $Q(z, t) \in \mathcal{M}_D(z)$. Then there exists $a \in \mathbb{C}^*$ such that $\sigma(e^{Q(z, t)}) = ae^{Q(z, t)}$.*

Proof. (1) We use the fact the elements of G are fields automorphisms that leave \mathcal{M}_D invariant.

(2) A computation shows that

$$\partial_{t_i}(z - \alpha(t))^{\beta(t)} = \left[\log(z - \alpha(t))\partial_{t_i}\beta(t) - \frac{\partial_{t_i}\alpha(t)\beta(t)}{z - \alpha(t)} \right] (z - \alpha(t))^{\beta(t)}.$$

The fact that σ commutes with all the derivations implies the existence of $a \in \mathbb{C}$ and $f(t) \in \mathcal{M}_D$ such that

$$\sigma\left(\log(z - \alpha(t))\right) = \log(z - \alpha(t)) + a \text{ and } \sigma\left((z - \alpha(t))^{\beta(t)}\right) = f(t)(z - \alpha(t))^{\beta(t)}.$$

Since $\partial_{t_i}\sigma = \sigma\partial_{t_i}$, we obtain that

$$\begin{aligned} & \left[\log(z - \alpha(t))\partial_{t_i}\beta(t) + a\partial_{t_i}\beta(t) - \frac{\partial_{t_i}\alpha(t)\beta(t)}{z - \alpha(t)} \right] f(t) \\ &= \partial_{t_i}f(t) + f(t) \left[\log(z - \alpha(t))\partial_{t_i}\beta(t) - \frac{\partial_{t_i}\alpha(t)\beta(t)}{z - \alpha(t)} \right]. \end{aligned}$$

Finally, $f(t)$ satisfies the parameterized linear differential equation

$$\partial_{t_i} \left(\frac{\partial_{t_i}f(t)}{f(t)a\partial_{t_i}\beta(t)} \right) = 0.$$

This means that $\frac{\partial_{t_i}f(t)}{f(t)a\partial_{t_i}\beta(t)} = c \in \mathbb{C}^*$ and $\log f(t) = a\beta(t) + \log(c)$. Then we deduce that $f(t) = ce^{a\beta(t)}$.

(3) We use the fact that

$$\partial_{t_i}\sigma\left(e^{Q(z, t)}\right) = \sigma\left(\partial_{t_i}\left(e^{Q(z, t)}\right)\right) = \sigma\left(\partial_{t_i}(Q(z, t))e^{Q(z, t)}\right) = \partial_{t_i}Q(z, t)\sigma\left(e^{Q(z, t)}\right).$$

The equation $\partial_{t_i}\sigma\left(e^{Q(z, t)}\right) = \partial_{t_i}Q(z, t)\sigma\left(e^{Q(z, t)}\right)$ admits $\sigma\left(e^{Q(z, t)}\right) = ae^{Q(z, t)}$ with $a \in \mathbb{C}^*$ as a solution. □

We deduce that the matrices of G are upper triangular. We will denote by $G_m \simeq \text{GL}_1(\mathcal{M}_D)$ the multiplicative group. The proof of the following proposition is inspired by the proof of [Sit75], Theorem 1.4. Let $p : G \rightarrow G_m$ that sends $\begin{pmatrix} m(t) & a(t) \\ 0 & m^{-1}(t) \end{pmatrix}$

on $m(t)$. Let M be the image of p and $A \subset \mathcal{M}_D$ such that

$$\left\{ \begin{pmatrix} 1 & a(t) \\ 0 & 1 \end{pmatrix}, \text{ where } a(t) \in A \right\}$$

is the kernel of p . We have already computed M with Lemma 2.2. For $m(t) \in M$, let $\Gamma_{m(t)}$ be the set of $\gamma_{m(t)} \in \mathcal{M}_D$ such that $\begin{pmatrix} m(t) & \gamma_{m(t)} \\ 0 & m(t)^{-1} \end{pmatrix} \in G$. We will identify \mathbb{C}^* with the field of constants elements of \mathcal{M}_D . If $\mathbb{C}^* \not\subset M$, because of Lemma 2.2, $g(z, t) \in \mathcal{M}_D(z)$, and we can compute explicitly $g(z, t) \int_{u=0}^z g(u, t)^{-2} du$, which is another solution. We obtain explicitly a fundamental solution and we can compute G . The next proposition explains how to compute G when $\mathbb{C}^* \subset M$.

Proposition 2.3. *Let us keep the same notations. Assume that $\mathbb{C}^* \subset M$. Then G is conjugated to*

$$\left\{ \begin{pmatrix} m(t) & a(t) \\ 0 & m(t)^{-1} \end{pmatrix}, \text{ where } m(t) \in M, a(t) \in A \right\}.$$

For the proof of the proposition, we will need the following lemmas.

Lemma 2.4. *Assume that $\mathbb{C}^* \subset M$. Let $m(t) \in M$ and $a(t) \in A$. Then $m(t)a(t) \in A$.*

Proof. With Lemma 2.2, we obtain that for all $m(t) \in M$, there exists $b(t) \in M$ such that $b(t)^2 = m(t)$. Let $m(t) \in M$, $b(t)^2 = m(t)$, $\gamma_{b(t)} \in \Gamma_{b(t)}$, and $a(t) \in A$. The computation

$$\begin{pmatrix} b(t) & \gamma_{b(t)} \\ 0 & b(t)^{-1} \end{pmatrix} \begin{pmatrix} 1 & a(t) \\ 0 & 1 \end{pmatrix} \begin{pmatrix} b(t) & \gamma_{b(t)} \\ 0 & b(t)^{-1} \end{pmatrix}^{-1} = \begin{pmatrix} 1 & m(t)a(t) \\ 0 & 1 \end{pmatrix}$$

shows that if $m(t) \in M$ and $a(t) \in A$, then $m(t)a(t) \in A$. \square

Lemma 2.5. *Assume that $\mathbb{C}^* \subset M$. Let $m(t) \in M$. Then $\gamma_{m(t)}, \gamma'_{m(t)} \in \Gamma_{m(t)}$ if and only if $(\gamma_{m(t)} - \gamma'_{m(t)}) \in A$.*

Proof. Let $\gamma_{m(t)}, \gamma'_{m(t)} \in \Gamma_{m(t)}$. The computation

$$\begin{pmatrix} m(t) & \gamma_{m(t)} \\ 0 & m(t)^{-1} \end{pmatrix} \begin{pmatrix} m(t) & \gamma'_{m(t)} \\ 0 & m(t)^{-1} \end{pmatrix}^{-1} = \begin{pmatrix} 1 & m(t)(\gamma_{m(t)} - \gamma'_{m(t)}) \\ 0 & 1 \end{pmatrix}$$

shows that $m(t)(\gamma_{m(t)} - \gamma'_{m(t)}) \in A$, and then $(\gamma_{m(t)} - \gamma'_{m(t)}) \in A$, because of Lemma 2.4. Conversely, if $(\gamma_{m(t)} - \gamma'_{m(t)}) \in A$ and $\gamma_{m(t)} \in \Gamma_{m(t)}$, then $m(t)(\gamma_{m(t)} - \gamma'_{m(t)}) \in A$, because of Lemma 2.4. The same computation shows that $\gamma'_{m(t)} \in \Gamma_{m(t)}$. \square

Lemma 2.6. *Assume that $\mathbb{C}^* \subset M$. Let $b \in \mathbb{C}^* \setminus \{\pm 1\}$ and $\gamma_b \in \Gamma_b$. Let*

$$\beta(t) = b(b^2 - 1)^{-1} \gamma_b.$$

Then, $\beta(t)(m(t) - m(t)^{-1}) \in \Gamma_{m(t)}$, for all $m(t) \in M$.

Proof. Let $m(t) \in M$ and $\gamma_{m(t)} \in \Gamma_{m(t)}$. The computation

$$\begin{aligned} & \begin{pmatrix} b & \gamma_b \\ 0 & b^{-1} \end{pmatrix} \begin{pmatrix} m(t) & \gamma_{m(t)} \\ 0 & m(t)^{-1} \end{pmatrix} \begin{pmatrix} b & \gamma_b \\ 0 & b^{-1} \end{pmatrix}^{-1} \begin{pmatrix} m(t) & \gamma_{m(t)} \\ 0 & m(t)^{-1} \end{pmatrix}^{-1} \\ &= \begin{pmatrix} 1 & (1 - m(t)^2)b\gamma_b - (1 - b^2)m(t)\gamma_{m(t)} \\ 0 & 1 \end{pmatrix} \end{aligned}$$

implies that $(1 - m(t)^2)b\gamma_b - (1 - b^2)m(t)\gamma_{m(t)} \in A$. Since $(1 - b^2)m(t) \in M$, Lemma 2.4 implies that

$$(1 - b^2)^{-1}m(t)^{-1}(1 - m(t)^2)b\gamma_b - \gamma_{m(t)} = \beta(t)(m(t) - m(t)^{-1}) - \gamma_{m(t)} \in A.$$

COMPUTING THE GALOIS GROUP OF SOME PARAMETERIZED
LINEAR DIFFERENTIAL EQUATION OF ORDER TWO

Therefore $\beta(t)(m(t) - m(t)^{-1}) \in \Gamma_{m(t)}$, because of Lemma 2.5. \square

Proof of Proposition 2.3. With Lemmas 2.5 and 2.6, we find that

$$G \simeq \left\{ \begin{pmatrix} m(t) & \beta(t)(m(t) - m(t)^{-1}) + a(t) \\ 0 & m(t)^{-1} \end{pmatrix}, \text{ where } m(t) \in M, a(t) \in A \right\}.$$

If we change the fundamental solution, i.e, if we conjugate G over $\text{GL}_2(\mathcal{M}_D)$, we can simplify the expression of G . After conjugation by the element $P = \begin{pmatrix} 1 & \beta(t) \\ 0 & 1 \end{pmatrix}$, we obtain that

$$PGP^{-1} \simeq \left\{ \begin{pmatrix} m(t) & a(t) \\ 0 & m(t)^{-1} \end{pmatrix}, \text{ where } m(t) \in M, a(t) \in A \right\}.$$

\square

We now want to compute G when $\mathbb{C}^* \subset M$. The computation of M has already been done in Lemma 2.2. We are now interested in the computation of A , which is a linear differential algebraic subgroup of $(\mathcal{M}_D, +)$. Cassidy classifies the linear differential algebraic subgroups of the additive group in [Cas72], Lemma 11. We define $\mathcal{M}_D[y_1, \dots, y_\nu]_{\Delta_t}$, as the ring of linear homogeneous differential polynomials. There exists $P_1, \dots, P_m \in \mathcal{M}_D[y]_{\Delta_t}$ such that

$$A = \{a(t) \in \mathcal{M}_D \mid P_1(a(t)) = \dots = P_m(a(t)) = 0\}.$$

We recall that $g(z, t) \int_{u=0}^z g(u, t)^{-2} du$ is another solution. We can choose $\beta(t) \in \mathcal{M}_D$ such that in the basis formed by the solutions $g(z, t)$ and $g(z, t) \int_{u=0}^z g(u, t)^{-2} du + \beta(t)g(z, t)$, G is equal to $\left\{ \begin{pmatrix} m(t) & a(t) \\ 0 & m(t)^{-1} \end{pmatrix}, \text{ where } m(t) \in M, a(t) \in A \right\}$. Let $G^g \subset G$ be the subfield of elements that fix $g(z, t)$ and let $\sigma \in G^g$. Let $a(t) \in A$ be such that

$$\begin{aligned} & \sigma \left(g(z, t) \int_{u=0}^z g(u, t)^{-2} du + \beta(t)g(z, t) \right) \\ &= \left(g(z, t) \int_{u=0}^z g(u, t)^{-2} du + \beta(t)g(z, t) \right) + a(t)g(z, t). \end{aligned}$$

Since

$$\sigma \left(g(z, t) \int_{u=0}^z g(u, t)^{-2} du + \beta(t)g(z, t) \right) = g(z, t) \left(\sigma \left(\int_{u=0}^z g(u, t)^{-2} du \right) + \beta(t) \right),$$

we deduce that

$$\sigma \left(\int_{u=0}^z g(u, t)^{-2} du \right) - \int_{u=0}^z g(u, t)^{-2} du = a(t) \in A.$$

Therefore, the differential polynomials P_i satisfy $\forall \sigma \in G^g$:

$$\begin{aligned} \sigma \left(P_i \left(\int_{u=0}^z g(u, t)^{-2} du \right) \right) &= P_i \left(\sigma \left(\int_{u=0}^z g(u, t)^{-2} du \right) \right) \\ &= P_i \left(\int_{u=0}^z g(u, t)^{-2} du + a(t) \right) \\ &= P_i \left(\int_{u=0}^z g(u, t)^{-2} du \right). \end{aligned}$$

Since $P_i \left(\int_{u=0}^z g(u, t)^{-2} du \right)$ is fixed by the elements of G^g , we deduce by the Galois correspondence in the parameterized differential Galois theory (see [CS07], Theorem 9.5) that

$$P_i(a(t)) = 0 \iff P_i \left(\int_{u=0}^z g(u, t)^{-2} du \right) \in \mathcal{M}_D(z) \langle g(z, t) \rangle_{\partial_z, \Delta_t},$$

where $\mathcal{M}_D(z) \langle g(z, t) \rangle_{\partial_z, \Delta_t}$ denotes the (∂_z, Δ_t) -differential field generated by $\mathcal{M}_D(z)$ and $g(z, t)$.

2.2. Let us consider the case number 2. There exists a Liouvillian solution of the form $e^{\int_0^z f(u,t)du}$, such that $f(z,t)$ satisfies

$$f(z,t)^2 + a(z,t)f(z,t) + b(z,t) = 0,$$

where $a(z,t), b(z,t) \in \mathcal{M}_D(z)$. There exists $\varepsilon \in \{\pm 1\}$ such that

$$f(z,t) = \frac{-a(z,t) + \varepsilon \sqrt{a(z,t)^2 - 4b(z,t)}}{2}.$$

By computing the action of G on $e^{\int_0^z \frac{-a(u,t) + \varepsilon \sqrt{a(u,t)^2 - 4b(u,t)}}{2} du}$, we find that $e^{\int_0^z \frac{-a(u,t) - \varepsilon \sqrt{a(u,t)^2 - 4b(u,t)}}{2} du}$ is another Liouvillian solution which is linearly independent of the first one. By computing the action of G on the second Liouvillian solution we find the existence of M , a linear differential algebraic subgroup of the multiplicative group G_m such that, in the basis formed by the two Liouvillian solutions

$$G \simeq \left\{ \begin{pmatrix} a(t) & 0 \\ 0 & a^{-1}(t) \end{pmatrix} \cup \begin{pmatrix} 0 & b^{-1}(t) \\ -b(t) & 0 \end{pmatrix}, \text{ where } a(t), b(t) \in M \right\}.$$

We are now interested in the computation of M . A direct computation shows that if there exists $\sigma \in G$ such that $\sigma \left(e^{\int_0^z f(u,t)du} \right) = \alpha(t) e^{\int_0^z f(u,t)du}$, then for all $i \leq n$, $\alpha(t)$ satisfies the parameterized differential equation

$$\partial_{t_i} \alpha(t) + \alpha(t) \left(\partial_{t_i} \int_0^z f(u,t)du \right) = \alpha(t) \sigma \left(\partial_{t_i} \int_0^z f(u,t)du \right).$$

Let $\widetilde{\partial}_{t_i} \alpha(t) = \frac{\partial_{t_i} \alpha(t)}{\alpha(t)}$ be the logarithm derivation. In [Cas72], Chapter 4, we see that there exist $P_1, \dots, P_k \in \mathcal{M}_D[y_1, \dots, y_n]_{\Delta_t}$ such that

$$M \simeq \left\{ \alpha(t) \mid P_1 \left(\widetilde{\partial}_{t_i} \alpha(t) \right) = \dots = P_k \left(\widetilde{\partial}_{t_i} \alpha(t) \right) = 0 \right\}.$$

The polynomial P_j satisfies, for all $\sigma \in G$, $P_j \left(\partial_{t_i} \int_0^z f(u,t)du \right) = \sigma \left(P_j \left(\partial_{t_i} \int_0^z f(u,t)du \right) \right)$ and then

$$P_j \left(\widetilde{\partial}_{t_i} \alpha(t) \right) = 0 \iff P_j \left(\partial_{t_i} \int_0^z f(u,t)du \right) \in \mathcal{M}_D(z).$$

2.3. In the third case, G is finite, because whose Zariski closure is finite. Since all finite linear differential algebraic subgroups of $\mathrm{SL}_2(\mathcal{M}_D)$ are finite linear algebraic subgroups of $\mathrm{SL}_2(\mathcal{M}_D)$, G is equal to the unparameterized differential Galois group. This is the same problem as in the unparameterized case. See [vHW05] for the computation of G .

2.4. We now consider the case where no Liouvillian solutions are found. We have seen in the introduction that in this case, the unparameterized differential Galois group is $\mathrm{SL}_2(\mathcal{M}_D)$. Therefore, G is Zariski dense in $\mathrm{SL}_2(\mathcal{M}_D)$.

The classification of the Zariski dense subgroup of $\mathrm{SL}_2(\mathcal{M}_D)$ has been made in [Cas72], Proposition 42. Let \mathbf{D} be the \mathcal{M}_D -vectorial space of derivations of the form

$$\left\{ \sum_{i=0}^n a_i(t) \partial_{t_i}, \text{ where } a_i(t) \in \mathcal{M}_D \right\},$$

and \mathbb{D} a vectorial subspace of \mathbf{D} . Let $\mathcal{M}_D^{\mathbb{D}}$ be the elements of \mathcal{M}_D that are constant for the derivations in \mathbb{D} . Remark that if $\mathbb{D} = \{0\}$, then $\mathcal{M}_D^{\mathbb{D}} = \mathcal{M}_D$. The linear differential algebraic subgroups of $\mathrm{SL}_2(\mathcal{M}_D)$ that are Zariski dense in $\mathrm{SL}_2(\mathcal{M}_D)$ are conjugated over $\mathrm{SL}_2(\mathcal{M}_D)$ to the groups of the form $\mathrm{SL}_2(\mathcal{M}_D^{\mathbb{D}})$, with \mathbb{D} a vectorial subspace of \mathbf{D} .

Let $\mathbb{D} \subset \mathbf{D}$ be such that G is conjugated over $\mathrm{SL}_2(\mathcal{M}_D)$ to $\mathrm{SL}_2(\mathcal{M}_D^{\mathbb{D}})$. We want to compute explicitly \mathbb{D} . This leads us to the notion of integrable systems. Let

COMPUTING THE GALOIS GROUP OF SOME PARAMETERIZED
LINEAR DIFFERENTIAL EQUATION OF ORDER TWO

$A_0(z, t), \dots, A_k(z, t)$, $m \times m$ matrices with entries in $\mathcal{M}_D(z)$ and $\partial'_{t_1}, \dots, \partial'_{t_k} \in \mathbf{D}$. The following system

$$[S] : \begin{cases} \partial_z Y(z, t) &= A_0(z, t)Y(z, t) \\ \partial'_{t_1} Y(z, t) &= A_1(z, t)Y(z, t) \\ &\vdots \\ \partial'_{t_k} Y(z, t) &= A_k(z, t)Y(z, t) \end{cases}$$

is integrable if and only if, for all $0 \leq i, j \leq k$,

$$\partial'_{t_j} A_i(z, t) - \partial'_{t_i} A_j(z, t) = A_j(z, t)A_i(z, t) - A_i(z, t)A_j(z, t),$$

where $\partial'_{t_0} = \partial_z$. We recall here [CS07], Proposition 6.3, which relates the integrable system and the parameterized differential Galois group in the case where the field of the ∂_z -constants is differentially closed.

Proposition 2.7. *Let $\{\partial'_{t_1}, \dots, \partial'_{t_k}\}$ be a commuting basis of \mathbb{D} , a vectorial subspace of \mathbf{D} . G is conjugated to $\mathrm{SL}_2(\mathcal{M}_{\mathbb{D}}^{\mathbb{D}})$ over $\mathrm{SL}_2(\mathcal{M}_{\mathbb{D}})$ if and only if there exist $A_1(z, t), \dots, A_k(z, t)$, $m \times m$ matrices with entries in $\mathcal{M}_D(z)^\dagger$, such that the following system is integrable:*

$$[S] : \begin{cases} \partial_z Y(z, t) &= A(z, t)Y(z, t) \\ \partial'_{t_1} Y(z, t) &= A_1(z, t)Y(z, t) \\ &\vdots \\ \partial'_{t_k} Y(z, t) &= A_k(z, t)Y(z, t). \end{cases}$$

We want to give simpler necessary and sufficient conditions for the integrability of the system in Proposition 2.7. First, we will write a necessary and sufficient condition for the integrability of

$$[S'] : \begin{cases} \partial_z Y(z, t) &= A(z, t)Y(z, t) \\ \partial' Y(z, t) &= A'(z, t)Y(z, t), \end{cases}$$

where $A'(z, t) = \begin{pmatrix} a(z, t) & b(z, t) \\ c(z, t) & d(z, t) \end{pmatrix}$ is an $m \times m$ matrix with entries in $\mathcal{M}_D(z)$ and $\partial' \in \mathbf{D}$.

The fact that $[S']$ is integrable is equivalent to the solution in $(\mathcal{M}_D(z))^\dagger$ of the parameterized differential system:

$$\begin{cases} \partial_z a(z, t) &= c(z, t) - b(z, t)r(z, t) \\ \partial_z b(z, t) &= d(z, t) - a(z, t) \\ \partial_z c(z, t) &= (a(z, t) - d(z, t))r(z, t) + \partial' r(z, t) \\ \partial_z d(z, t) &= b(z, t)r(z, t) - c(z, t) \end{cases}$$

$$\iff \begin{cases} \partial_z a(z, t) &= -\partial_z d(z, t) \\ \partial_z^2 b(z, t) &= 2\partial_z d(z, t) \\ \partial_z c(z, t) &= -\partial_z b(z, t)r(z, t) + \partial' r(z, t) \\ \frac{\partial_z^2 b(z, t)}{2} &= b(z, t)r(z, t) - c(z, t) \end{cases}$$

$$\iff \begin{cases} \partial_z a(z, t) &= -\partial_z d(z, t) \\ \partial_z^2 b(z, t) &= 2\partial_z d(z, t) \\ \partial_z c(z, t) &= -\partial_z b(z, t)r(z, t) + \partial' r(z, t) \\ \frac{\partial_z^3 b(z, t)}{2} &= 2\partial_z b(z, t)r(z, t) + b(z, t)\partial_z r(z, t) - \partial' r(z, t). \end{cases}$$

We can easily see that the existence of $b(z, t) \in \mathcal{M}_D(z)$ as a solution of

$$\frac{\partial_z^3 b(z, t)}{2} = 2\partial_z b(z, t)r(z, t) + b(z, t)\partial_z r(z, t) - \partial' r(z, t)$$

[†]Using the result of Seidenberg, we can identify the matrices as elements of $\mathrm{GL}_2(\mathcal{M}_D(z))$ because their entries involve a finite number of elements of the fields of the ∂_z -constants.

is equivalent to the fact that the system $[S']$ is integrable. There exists an algorithm to determine if such a system has a solution (see [vdPS03], p. 100). We obtain a necessary and sufficient condition on ∂' for the integrability condition of the system $[S']$. Let \mathbb{D} be the maximal vectorial subspace of \mathbf{D} such that for all derivations ∂' in \mathbb{D} , there exists $A'(z, t)$, $m \times m$ matrix with entries in $\mathcal{M}_D(z)$ such that the following system is integrable:

$$[S'] : \begin{cases} \partial_z Y(z, t) &= A(z, t)Y(z, t) \\ \partial' Y(z, t) &= A'(z, t)Y(z, t). \end{cases}$$

We want to prove that the parameterized differential Galois group of $\partial_z Y(z, t) = A(z, t)Y(z, t)$ is conjugated to $\mathrm{SL}_2(\mathcal{M}_D^{\mathbb{D}})$ over $\mathrm{SL}_2(\mathcal{M}_D)$. Assume that this is not the case. Then by Proposition 2.7, there exists $\mathbb{D}_1, \mathbb{D}_2 \subsetneq \mathbb{D}$, having at least dimension 1, with $\mathbb{D}_1 \neq \mathbb{D}_2$ such that G is conjugated to $\mathrm{SL}_2(\mathcal{M}_D^{\mathbb{D}_1})$ and $\mathrm{SL}_2(\mathcal{M}_D^{\mathbb{D}_2})$. In this case, $\mathrm{SL}_2(\mathcal{M}_D^{\mathbb{D}_1})$ is conjugated to $\mathrm{SL}_2(\mathcal{M}_D^{\mathbb{D}_2})$ over $\mathrm{SL}_2(\mathcal{M}_D)$. The fact that $\mathbb{D}_1 = \mathbb{D}_2$ is proved in [Sit75], Theorem 1.2, Chapter 2, but we will recall the proof here. Let $\alpha \in \mathcal{M}_D^{\mathbb{D}_1}$ and consider the diagonal matrix $M = \begin{pmatrix} \alpha & 0 \\ 0 & \alpha^{-1} \end{pmatrix} \in \mathrm{SL}_2(\mathcal{M}_D^{\mathbb{D}_1})$. Since similar matrices have the same set of eigenvalues and $\mathcal{M}_D^{\mathbb{D}_2}$ is algebraically closed, we obtain that $\alpha(t) \in \mathcal{M}_D^{\mathbb{D}_2}$. Therefore $\mathcal{M}_D^{\mathbb{D}_1} \subset \mathcal{M}_D^{\mathbb{D}_2}$ and, by symmetry, $\mathcal{M}_D^{\mathbb{D}_1} = \mathcal{M}_D^{\mathbb{D}_2}$. We then deduce $\mathbb{D}_1 = \mathbb{D}_2 = \mathbb{D}$. We have proved:

Proposition 2.8. *We have the following equivalences:*

- (1) G is conjugated to $\mathrm{SL}_2(\mathcal{M}_D^{\mathbb{D}})$ over $\mathrm{SL}_2(\mathcal{M}_D)$.
- (2) For all ∂' that belongs in a commuting basis of \mathbb{D} , the following parameterized differential equation has a solution in $\mathcal{M}_D(z)$:

$$\frac{\partial_z^3 b(z, t)}{2} = 2\partial_z b(z, t)r(z, t) + b(z, t)\partial_z r(z, t) - \partial' r(z, t).$$

- (3) For all $\partial' \in \mathbb{D}$, the following parameterized differential equation has a solution in $\mathcal{M}_D(z)$:

$$\frac{\partial_z^3 b(z, t)}{2} = 2\partial_z b(z, t)r(z, t) + b(z, t)\partial_z r(z, t) - \partial' r(z, t).$$

Remark 2.9. In the case where $n = 1$, i.e, there is only one parameter, the Zariski dense subgroups of $\mathrm{SL}_2(\mathcal{M}_D)$ are (up to conjugation over $\mathrm{SL}_2(\mathcal{M}_D)$) $\mathrm{SL}_2(\mathcal{M}_D)$ and $\mathrm{SL}_2(\mathbb{C})$. Then we only have to check whether

$$\frac{\partial_z^3 b(z, t)}{2} = 2\partial_z b(z, t)r(z, t) + b(z, t)\partial_z r(z, t) - \partial_t r(z, t),$$

has a solution in $\mathcal{M}_D(z)$.

2.5. We summarize in the next theorem the results of this section.

Theorem 2.10. *Let us consider $\partial_z^2 Y(z, t) = r(z, t)Y(z, t)$ with $r(z, t) \in \mathcal{M}_D(z)$ and let G be the parameterized differential Galois group, seen as a linear differential algebraic subgroup of $\mathrm{SL}_2(\mathcal{M}_D)$. There are four possibilities:*

- (1) *There exists a Liouvillian solution of the form $g(z, t) = e^{\int_0^z f(u, t) du}$, with $f(z, t)$ belongs to $\mathcal{M}_D(z)$. There are two possibilities:*
 - (a) *If $g(z, t) \in \mathcal{M}_D$, then we can compute explicitly another solution $g(z, t) \int_{u=0}^z g(u, t)^{-2} du$ which is linearly independent with $g(z, t)$. In this basis of solutions we can compute explicitly G .*
 - (b) *In the other case, G is conjugated to:*

$$\left\{ \begin{pmatrix} m(t) & a(t) \\ 0 & m(t)^{-1} \end{pmatrix}, \text{ where } m(t) \in M, a(t) \in A \right\},$$

where:

$$M = \{g(z, t)^{-1} \sigma(g(z, t)), \sigma \in G\},$$

$$A = \left\{ a(t) \in \mathcal{M}_D \left| \begin{array}{l} \forall P \in \mathcal{M}_D[y]_{\Delta_t}, \\ P(\int_{u=0}^z g(u, t)^{-2} du) \in \mathcal{M}_D(z) \langle g(z, t) \rangle_{\partial_z, \Delta_t} \iff P(a(t)) = 0 \end{array} \right. \right\}.$$

- (2) *There exists a Liouvillian solution of the form $g(z, t) = e^{\int_0^z f(u, t) du}$, where $f(z, t)$ is algebraic over $\mathcal{M}_D(z)$ of degree two and $f(z, t) \notin \mathcal{M}_D(z)$. In this case, G is conjugated to*

$$\left\{ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \cup \begin{pmatrix} 0 & b^{-1} \\ -b & 0 \end{pmatrix}, \text{ where } a, b \in M \right\}, \text{ where}$$

$$M = \left\{ f(t) \in \mathcal{M}_D \left| \begin{array}{l} \forall P \in \mathcal{M}_D[y_1 \dots, y_n]_{\Delta_t}, \\ P(\partial_{t_i} \int_{u=0}^z f(u, t) du) \in \mathcal{M}_D(z) \iff P(\tilde{\partial}_{t_i} f(t)) = 0 \end{array} \right. \right\}.$$

- (3) *G is finite. In this case, G is equal to the unparameterized differential Galois group.*
- (4) *There are no Liouvillian solutions. In this case, there exists \mathbb{D} , a \mathcal{M}_D -vectorial space of derivations spanned by Δ_t , such that G is conjugated to $\mathrm{SL}_2(\mathcal{M}_D^{\mathbb{D}})$. Moreover, $\partial'_t \in \mathbb{D}$ if and only if the following parameterized differential equation has a solution in $\mathcal{M}_D(z)$:*

$$\frac{\partial_z^3 b(z, t)}{2} = 2\partial_z b(z, t)r(z, t) + b(z, t)\partial_z r(z, t) - \partial'_t r(z).$$

Notice that the computation of the Liouvillian solutions and the unparameterized differential Galois group are already known. Our results compute the parameterized differential Galois group in the cases 1,2 and 4. The classification of the Zariski dense linear differential algebraic subgroup of $\mathrm{SL}_2(\mathcal{M}_D)$ and the link with integrable systems were already known (see [Cas72, CS07]), but we give here an effective way to compute the Galois group in the case number 4 and we decrease the number of integrability conditions.

3. EXAMPLES

In the following examples, we will consider equations having coefficients in $\mathcal{M}_D(z)$ and we will compute G as a linear differential algebraic subgroup of $\mathrm{SL}_2(\mathcal{M}_D)$. In the three first examples, we are in the case where no Liouvillian solutions are found. In the fourth example, we are in the case number 1 and in the last example, we are in the case number 2.

Example 3.1 (Schrodinger equation with rational potential of odd degree). Let us consider $r(z, t) = z^{2n+1} + \sum_{i=0}^{2n} t_i z^i$. There are no Liouvillian solutions. The parameterized linear differential equation

$$\frac{\partial_z^3 b(z, t)}{2} = 2\partial_z b(z, t)r(z, t) + b(z, t)\partial_z r(z, t) - \sum_{i=0}^{2n} a_i(t)z^i$$

has a rational solution if and only if there exists $c(t) \in \mathcal{M}_D$ such that

$$\begin{cases} a_{2n}(t) &= c(t)(2n+1) \\ i < 2n : &a_i(t) = c(t)(i+1)t_{i+1}. \end{cases}$$

Then

$$G \simeq \mathrm{SL}_2(\mathcal{M}_D^{\partial_{t'}}), \text{ where } \partial_{t'} = (2n+1)\partial_{t_{2n}} + \sum_{i=0}^{2n-1} (i+1)t_{i+1}\partial_{t_i}.$$

Example 3.2 (Bessel equation). Let $r(z, t) = \frac{4t^2-1}{4z^2} - 1$. In [Kov86], §4.2, Example 2, we see that if $t \notin \frac{1}{2} + \mathbb{Z}$, this parameterized linear differential equation has no Liouvillian solution. We can choose D such that $\{D \cap (\frac{1}{2} + \mathbb{Z})\} = \emptyset$. We obtain that G is Zariski dense in $\mathrm{SL}_2(\mathcal{M}_D)$. With Remark 2.9, we have to see whether the parameterized linear differential equation

$$(3.1) \quad \frac{\partial_z^3 b(z, t)}{2} = 2\partial_z b(z, t) \left(\frac{4t^2-1}{4z^2} - 1 \right) + b(z, t) \frac{1-4t^2}{2z^3} - \frac{2t}{z}$$

has a solution in $\mathcal{M}_D(z)$. Suppose that there exists $b(z, t) \in \mathcal{M}_D(z)$ satisfying such an equation. We can see directly that if $b(z, t)$ has a pole, then it is $z = 0$. Assume that $b(z, t)$ has a pole of order ν at $z = 0$ and let $0 \neq f(t) \in \mathcal{M}_D$ equal the value at $(0, t)$ of $z^\nu b(z, t)$. Since $b(z, t)$ satisfies the equation (3.1), we find for all $t \in D$:

$$\frac{-f(t)\nu(\nu-1)(\nu-2)}{2} = -f(t)\nu \frac{4t^2-1}{2} + f(t) \frac{1-4t^2}{2}.$$

For all ν , there is no $0 \neq f(t)$ satisfying this equality and we find that $b(z, t) \in \mathcal{M}_D[z]$. Let ν be its degree and $f(t)$ its leading term. The equation (3.1) has no constant solution, and we can assume $\nu > 1$. We find that for all $t \in D$,

$$0 = -2\nu f(t),$$

which implies that the equation (3.1) has no solutions in $\mathcal{M}_D(z)$ and then

$$G \simeq \mathrm{SL}_2(\mathcal{M}_D).$$

COMPUTING THE GALOIS GROUP OF SOME PARAMETERIZED
LINEAR DIFFERENTIAL EQUATION OF ORDER TWO

Example 3.3 (Harmonic oscillator). Let $r(z, t) = \frac{z^2}{4} + t$. There are no Liouvillian solutions. With Remark 2.9, we have to check whether the parameterized linear differential equation

$$\frac{\partial_z^3 b(z, t)}{2} = 2\partial_z b(z, t) \left(\frac{z^2}{4} + t \right) + b(z, t) \frac{z}{2} - 1$$

has a solution in $\mathcal{M}_D(z)$. We can see directly that if $b(z, t) \in \mathcal{M}_D(z)$ is a solution, then it has no poles, which means that $b(z, t) \in \mathcal{M}_D[z]$. Let ν be its degree and $0 \neq f(t)$ be its leading term. We find that $\frac{(\nu+1)f(t)}{2} = 0$, which admits no solution different from 0. Then

$$G \simeq \mathrm{SL}_2(\mathcal{M}_D).$$

Example 3.4. If $r(z, t) = \frac{t}{z^2}$, then we have two Liouvillian solutions

$$f_1(z, t) = \sqrt{z} z^{\frac{\sqrt{1+4t}}{2}} \text{ and } f_2(z, t) = \sqrt{z} z^{-\frac{\sqrt{1+4t}}{2}}.$$

We can compute the parameterized differential Galois group for the fundamental solution

$$\begin{pmatrix} f_1(z, t) & f_2(z, t) \\ \partial_z f_1(z, t) & \partial_z f_2(z, t) \end{pmatrix} :$$

$$G \simeq \left\{ \begin{pmatrix} \alpha e^{a(\sqrt{1+4t})} & 0 \\ 0 & \alpha^{-1} e^{-a(\sqrt{1+4t})} \end{pmatrix}, \text{ where } a \in \mathbb{C}, \alpha \in \mathbb{C}^* \right\}.$$

Viewed as a linear differential algebraic subgroup $\mathrm{GL}_2(\mathcal{M}_D)$,

$$G \simeq \left\{ \begin{pmatrix} \alpha(t) & 0 \\ 0 & \alpha^{-1}(t) \end{pmatrix}, \text{ where } \partial_t \left(\frac{\sqrt{1+4t} \partial_t \alpha(t)}{\alpha(t)} \right) = 0 \right\}.$$

Example 3.5. If $r(z, t) = \frac{t}{z} - \frac{3}{16z^2}$, then we have two Liouvillian solutions

$$f_1(z, t) = (z)^{1/4} e^{2(tz)^{1/2}} \text{ and } f_2(z, t) = (z)^{1/4} e^{-2(tz)^{1/2}}.$$

We can compute the parameterized differential Galois group for the fundamental solution

$$\begin{pmatrix} f_1(z, t) & f_2(z, t) \\ \partial_z f_1(z, t) & \partial_z f_2(z, t) \end{pmatrix} :$$

$$G \simeq \left\{ \begin{pmatrix} a(t) & 0 \\ 0 & a^{-1}(t) \end{pmatrix} \cup \begin{pmatrix} 0 & b^{-1}(t) \\ -b(t) & 0 \end{pmatrix}, \text{ where } a(t), b(t) \in \mathbb{C}^* \right\}.$$

We can remark that we have an integrable system

$$\begin{cases} \partial_z Y(z, t) & = A(z, t)Y(z, t) \\ \partial_t Y(z, t) & = B(z, t)Y(z, t) \end{cases}$$

with

$$A(z, t) = \begin{pmatrix} 0 & 1 \\ \frac{t}{z} - \frac{3}{16z^2} & 0 \end{pmatrix} \text{ and } B(z, t) = \begin{pmatrix} -\frac{1}{4t} & \frac{z}{t} \\ 1 - \frac{3}{16tz} & \frac{3}{4t} \end{pmatrix}.$$

REFERENCES

- [AH09] Primitivo B Acosta-Humanez. *Galoisian Approach to Supersymmetric Quantum Mechanics, Phd Dissertation*. PhD thesis, Universitat Politècnica de Catalunya, 2009.
- [AHMRW11] Primitivo B. Acosta-Humánez, Juan J. Morales-Ruiz, and Jacques-Arthur Weil. Galoisian approach to integrability of Schrödinger equation. *Rep. Math. Phys.*, 67(3):305–374, 2011.
- [Arr12] Carlos E. Arreche. Computing the differential galois group of a one-parameter family of second order linear differential equations. preprint available on arxiv. 2012.
- [Cas72] P. J. Cassidy. Differential algebraic groups. *Amer. J. Math.*, 94:891–954, 1972.
- [CS07] Phyllis J. Cassidy and Michael F. Singer. Galois theory of parameterized differential equations and linear differential algebraic groups. In *Differential equations and quantum groups*, volume 9 of *IRMA Lect. Math. Theor. Phys.*, pages 113–155. Eur. Math. Soc., Zürich, 2007.
- [DLR92] Anne Duval and Michèle Loday-Richaud. Kovačič’s algorithm and its application to some families of special functions. *Appl. Algebra Engrg. Comm. Comput.*, 3(3):211–246, 1992.
- [GO12] Sergey Gorchinskiy and Alexey Ovchinnikov. Isomonodromic differential equations and differential tannakian categories. to appear in journal de mathématiques pures et appliquées. 2012.
- [HS08] Charlotte Hardouin and Michael F. Singer. Differential Galois theory of linear difference equations. *Math. Ann.*, 342(2):333–377, 2008.
- [HvdP95] Peter A. Hendriks and Marius van der Put. Galois action on solutions of a differential equation. *J. Symbolic Comput.*, 19(6):559–576, 1995.
- [Kol73] E. R. Kolchin. *Differential algebra and algebraic groups*. Academic Press, New York, 1973. Pure and Applied Mathematics, Vol. 54.
- [Kov86] Jerald J. Kovacic. An algorithm for solving second order linear homogeneous differential equations. *J. Symbolic Comput.*, 2(1):3–43, 1986.
- [Lan08] Peter Landesman. Generalized differential Galois theory. *Trans. Amer. Math. Soc.*, 360(8):4441–4495, 2008.
- [Sei58] A. Seidenberg. Abstract differential algebra and the analytic case. *Proc. Amer. Math. Soc.*, 9:159–164, 1958.
- [Sei69] A. Seidenberg. Abstract differential algebra and the analytic case. II. *Proc. Amer. Math. Soc.*, 23:689–691, 1969.
- [Sit75] William Yu Sit. Differential algebraic subgroups of $SL(2)$ and strong normality in simple extensions. *Amer. J. Math.*, 97(3):627–698, 1975.
- [SU93a] Michael F. Singer and Felix Ulmer. Galois groups of second and third order linear differential equations. *J. Symbolic Comput.*, 16(1):9–36, 1993.
- [SU93b] Michael F. Singer and Felix Ulmer. Liouvillian and algebraic solutions of second and third order linear differential equations. *J. Symbolic Comput.*, 16(1):37–73, 1993.
- [UW96] Felix Ulmer and Jacques-Arthur Weil. Note on Kovacic’s algorithm. *J. Symbolic Comput.*, 22(2):179–200, 1996.
- [vdP99] Marius van der Put. Symbolic analysis of differential equations. In *Some tapas of computer algebra*, volume 4 of *Algorithms Comput. Math.*, pages 208–236. Springer, Berlin, 1999.
- [vdPS03] Marius van der Put and Michael F. Singer. *Galois theory of linear differential equations*, volume 328 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 2003.
- [vHW05] M. van Hoeij and J.-A. Weil. Solving second order linear differential equations with Klein’s theorem. In *ISSAC’05*, pages 340–347. ACM, New York, 2005.
- [Zha95] Alexey Zharkov. Coefficient fields of solutions in Kovacic’s algorithm. *J. Symbolic Comput.*, 19(5):403–408, 1995.

UNIVERSITÉ PARIS DIDEROT - INSTITUT DE MATHÉMATIQUES DE JUSSIEU,
 Current address: 4, place Jussieu 75005 Paris.
 E-mail address: tdreyfus@math.jussieu.fr.