



HAL
open science

Mean-field limit of a spatially-extended FitzHugh-Nagumo neural network

Joachim Crevat

► **To cite this version:**

Joachim Crevat. Mean-field limit of a spatially-extended FitzHugh-Nagumo neural network. *Kinetic and Related Models*, 2019, 12 (6), pp.1329-1358. hal-01897746v2

HAL Id: hal-01897746

<https://hal.science/hal-01897746v2>

Submitted on 6 Mar 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Mean-field limit of a spatially-extended FitzHugh-Nagumo neural network

Joachim Crevat^{*1}

¹Institut de Mathématiques de Toulouse ; UMR5219, Université de Toulouse ; UPS IMT, F-31062
Toulouse Cedex 9 France

March 1, 2019

Abstract

We consider a spatially-extended model for a network of interacting FitzHugh-Nagumo neurons without noise, and rigorously establish its mean-field limit towards a nonlocal kinetic equation as the number of neurons goes to infinity. Our approach is based on deterministic methods, and namely on the stability of the solutions of the kinetic equation with respect to their initial data. The main difficulty lies in the adaptation in a deterministic framework of arguments previously introduced for the mean-field limit of stochastic systems of interacting particles with a certain class of locally Lipschitz continuous interaction kernels. This result establishes a rigorous link between the microscopic and mesoscopic scales of observation of the network, which can be further used as an intermediary step to derive macroscopic models. We also propose a numerical scheme for the discretization of the solutions of the kinetic model, based on a particle method, in order to study the dynamics of its solutions, and to compare it with the microscopic model.

Keywords: Mean-field limit, neural network, FitzHugh-Nagumo, Wasserstein distance.

1 Introduction

The FitzHugh-Nagumo (FHN) model [21, 30] focuses on the evolution of the membrane electrical potential of a nerve cell depending on the input it receives. Such variations depend on the ion exchanges between the neuron and its environment through synapses, which were precisely described by the Hodgkin-Huxley model [27]. The FHN model was then developed as a simplification of the Hodgkin-Huxley model. Rather than precisely describe all the variations of activatory and inhibitory channels of ions, the FHN model gathers all the fast excitable dynamics in the variable v , which is still considered as the electrical membrane potential, and the slow refractory dynamics in a newly introduced adaptation variable w . It can be written as follows

^{*}joachim.crevat@math.univ-toulouse.fr

for $t > 0$:

$$\begin{cases} \frac{dv}{dt} = N(v) - w + I_{\text{ext}}, \\ \frac{dw}{dt} = \tau(v + a - bw), \end{cases} \quad (1.1)$$

where I_{ext} stands for the input current the neuron receives from its environment, $\tau \geq 0$, $a \in \mathbb{R}$ and $b \geq 0$ are some given constants, and $N(v)$ is a nonlinearity which models the cell excitability. A typical choice for the nonlinearity N (see for instance [2, 5, 29, 30]) is the cubic function

$$N : v \mapsto v(\alpha - \beta v^2) \quad (1.2)$$

for any $\alpha, \beta > 0$. Since the input I_{ext} is the result of the interactions between the nerve cell and its neighbours, we can replace it with an interaction term with other similar neurons, to consider an individual-based model for a network of n interacting neurons, where $n \in \mathbb{N}$. Biological observations seem to exclude the case of homogeneous interactions [32, 40], and show that the interactions inside a network of neurons are spatially structured [7]. Hence, as in [7, 28], we choose to modulate the neural connectivity with a spatial weight. Thus, we consider the following spatially-extended FHN system for n interacting neurons for $t > 0$:

$$\begin{cases} \frac{d\mathbf{x}_i}{dt} = 0, \\ \frac{dv_i}{dt} = N(v_i) - w_i - \frac{1}{n} \sum_{j=1}^n \Psi(\mathbf{x}_i, \mathbf{x}_j)(v_i - v_j), \\ \frac{dw_i}{dt} = \tau(v_i + a - bw_i). \end{cases} \quad (1.3)$$

21 where each neuron within the network is labelled by its index $i \in \{1, \dots, n\}$, $\mathbf{x}_i \in \mathbb{R}^d$ with $d \in \{1, 2, 3\}$
 22 is a parameter which stands for the position of the neuron i , and $(v_i, w_i) \in \mathbb{R}^2$ is the couple membrane
 23 potential and adaptation variable of the neuron i . Here, the connectivity weight $\Psi : \mathbb{R}^{2d} \rightarrow \mathbb{R}$ models
 24 the effect of spatial dependence on the strength of neuronal interactions which we assume to be of electric
 25 type. More precisely, for every neural cell i , each other neuron j contributes in the input current received
 26 by neuron i , and we write this interaction using Ohm's law *via* $\Psi(\mathbf{x}_i, \mathbf{x}_j)(v_i - v_j)$, which is then summed
 27 over the network. In doing so, we assumed that conductances from the neuron j to the neuron i , denoted
 28 $\Psi(\mathbf{x}_i, \mathbf{x}_j)$, are modulated by the positions of the two neurons i and j . Moreover, the scaling factor $1/n$ is
 29 introduced in order to renormalize the contributions of each neuron. From a biological viewpoint, since
 30 a neuron interacts with its closest neighbours, a reasonable choice for $\Psi(\mathbf{x}_i, \cdot)$ is the indicator function of
 31 a small ball centered in \mathbf{x}_i . Here, the framework considered in this paper admits only bounded Lipschitz
 32 continuous connectivity kernels.

33 The main purpose of this article is to rigorously derive a kinetic mean-field model of the FHN system (1.3)
 34 as the number of neurons n goes to infinity. The classical method to derive a mean-field description from
 35 an individually based model is to study the time evolution at each location of the probability of finding
 36 neurons characterized by a certain couple potential-adaptation variable. The framework of this approach is
 37 to work in a probability measure set independent on the number of neurons. Thus, we recall the following
 38 notion of empirical measure.

Definition 1.1 (Empirical measure). *To each n -tuple $\mathbf{X}_n = (\mathbf{x}_i)_{1 \leq i \leq n} \in (\mathbb{R}^d)^n$, $V_n = (v_i)_{1 \leq i \leq n} \in \mathbb{R}^n$ and $W_n = (w_i)_{1 \leq i \leq n} \in \mathbb{R}^n$, one associates its empirical measure:*

$$\mu_{(\mathbf{X}_n, V_n, W_n)} := \frac{1}{n} \sum_{j=1}^n \delta_{(\mathbf{x}_j, v_j, w_j)}. \quad (1.4)$$

Let us highlight that an empirical measure is a probability measure on \mathbb{R}^{d+2} . As the number of neurons n goes to infinity, the interaction term in (1.3) is expected to formally satisfy:

$$\frac{1}{n} \sum_{j=1}^n \Psi(\mathbf{x}, \mathbf{x}_j)(v - v_j) \rightarrow \int \Psi(\mathbf{x}, \mathbf{x}') (v - v') f(t, d\mathbf{x}', dv', dw'),$$

where $f(t, d\mathbf{x}, dv, dw)$ is the probability measure of finding neurons in an elementary volume of side $d\mathbf{x}$ with a potential membrane and an adaptation variable in an elementary interval respectively of length dv and dw at time $t \geq 0$ within the cortex. Our purpose is thus to prove the convergence of the empirical measures associated to the solutions of the FHN system (1.3) towards such a probability measure f satisfying the following nonlocal kinetic equation:

$$\partial_t f + \partial_v (f [N(v) - w - \mathcal{K}[f]]) + \partial_w (f A(v, w)) = 0, \quad t > 0, \quad (\mathbf{x}, v, w) \in \mathbb{R}^{d+2}, \quad (1.5)$$

where

$$\begin{cases} \mathcal{K}[f](t, \mathbf{x}, v) & := \int_{\mathbb{R}^{d+2}} \Psi(\mathbf{x}, \mathbf{x}') (v - v') f(t, d\mathbf{x}', dv', dw'), \\ A(v, w) & := \tau (v + a - b w). \end{cases}$$

The term $-\partial_v (f \mathcal{K}[f])$ describes nonlocal interactions through the whole network, and $\partial_v (f(N(v) - w)) + \partial_w (f A(v, w))$ accounts for the local reaction due to the excitability of nerve cells. Since this PDE can be written in a divergence form, we directly have the conservation of mass, which leads us to complement (1.5) with an initial condition:

$$f(0, \cdot) = f_0. \quad (1.6)$$

39 The derivation of the kinetic equation (1.5) joins a large literature in mathematical neuroscience and
40 kinetic theory. Indeed, the problem of the mean-field limit of systems of interacting particles towards
41 kinetic models has been widely investigated, varying the types of interactions. We mention the work of
42 Dobrushin [17], who introduced some classical methods to prove the mean-field limit of an individual-based
43 model of interacting particles with a bounded globally Lipschitz continuous interaction kernel towards a
44 Vlasov equation, using a stability result of the solutions to the kinetic model with respect to their initial
45 data, in a suitable topology of probability measures. Such a stability result is called Dobrushin's estimate.
46 See for instance [22] for a more recent review of this approach. Then, similar results have been proved for a
47 larger variety of interaction kernels. For example, in [25], the authors proved the mean-field limit towards
48 a Vlasov equation, in the case where the interaction kernel has a singularity. Besides, in [4], the authors
49 introduced an extension of the classical mean-field theory from [17] which also works for a certain class of
50 locally Lipschitz continuous interaction kernels.

51 Coming back to a neuroscience viewpoint, in [18, 38], the authors analysed the mean-field limit of a network
52 of neurons modeled by their firing rates, divided into a fixed number of populations, with random synaptic

53 weights whose probability distributions only depend upon the population indexes. They also studied the
 54 influence of the noise on collective behaviors in the network. Another perspective is to focus on the
 55 probability that each neuron releases an action potential. This kind of model is called time-elapsd neuron
 56 (see for instance [12–14, 33–35]). Here, in this article, we use another approach, with the derivation of a
 57 kinetic equation from the FHN model, which does not abstract the emission of action potentials, following
 58 the example of the Hodgkin-Huxley model.

59 The mean-field limit of the stochastic spatially-extended FHN model for interacting neurons has already
 60 been studied in the specific case of a compactly supported connectivity kernel in [28], and in the case
 61 when each cell interacts only with its p nearest neighbours in [28, 31]. Other approaches are considered
 62 in the literature to adjust the interactions with other quantities than space. For example, in [2, 5], the
 63 authors focused on the mean-field limit of a stochastic FHN model or Hodgkin-Huxley model of n neurons
 64 interacting through electro-chemical synapses towards a kinetic PDE similar to (1.5). In [29], the authors
 65 proved a similar result for a stochastic FHN model of a neural network with homogeneous conductance. We
 66 also mention works like [6, 36] which prove the synchronization of mean-field models of neural networks
 67 of respectively Hodgkin-Huxley and FHN type. The specificity of our present article compared to the
 68 literature is that we only consider spatially-weighted neural networks without noise.

69 The main contribution of this article is the rigorous justification of the kinetic model (1.5) we considered
 70 in [16] as the mean-field limit of the FHN system (1.3). It provides a mesoscopic description of the neural
 71 network, which can be used as an intermediary step for the derivation of macroscopic models from the FHN
 72 system (1.3), as in [16]. Here, we work with an initial data for (1.5) which is not compactly supported in
 73 general, but with finite exponential moments. Even though the main mathematical methods that we use
 74 were introduced in [4], to our knowledge, they have not been applied for the FHN system in a deterministic
 75 and spatially structured framework. Furthermore, we consider a numerical scheme for the kinetic equation
 76 (1.5), using a particle method, in order to display some numerical simulations of this model. It appears
 77 that for certain sets of parameters, we observe some dynamics usually expected for macroscopic models.
 78 We also compare this scheme with the FHN system (1.3), to get some numerical evidence of the relevance
 79 of the kinetic equation (1.5).

80 Here, in our framework the interaction kernel $(\mathbf{x}, \mathbf{y}, v) \mapsto \Psi(\mathbf{x}, \mathbf{y})v$ is locally Lipschitz continuous but
 81 not globally. Indeed, it is Lipschitz continuous on every bounded sets of \mathbb{R}^{d+1} but its Lipschitz constant
 82 goes to infinity with the diameter of the set considered. Therefore some terms of higher order arise in
 83 the computation of a Dobrushin’s estimate. Thus, we need additional assumptions to control these terms,
 84 and hence to compute a similar stability estimate. In the spirit of [4], we circumvent this issue with the
 85 estimate of exponential moments of the solution of the kinetic equation (1.5), and with an appropriate
 86 division of the set of integration in the nonlocal terms, to get a suitable estimate. We mention that in
 87 [10, 23], the authors tackled a similar difficulty for some kinetic models of collective motion, and they chose
 88 to work with compactly supported solutions specifically to overcome this problem, controlling the growth
 89 of velocity support to construct measure solutions.

90 2 Main result

91 This section is devoted to the statement of our main result on the mean-field limit from the solution of
 92 the microscopic model (1.3) towards the solution of the kinetic equation (1.5) as the number of neurons n

93 goes to infinity.

94 Before stating our main result, let us precisely define the notion of solution of the kinetic equation (1.5)
 95 we use in this article. In the following, we denote by $\mathcal{P}_2(\mathbb{R}^{d+2})$ the set of probability measures on \mathbb{R}^{d+2}
 96 with finite moments of order 2.

Definition 2.1 (Measure solution of (1.5)). *Consider an initial data $f_0 \in \mathcal{P}_2(\mathbb{R}^{d+2})$. Let $T > 0$. Then, f is said to be a measure solution of (1.5) with initial data f_0 if $f \in \mathcal{C}([0, T], \mathcal{P}_2(\mathbb{R}^{d+2}))$ and for all $t \in [0, T]$,*

$$f(t) = \mathcal{Z}_{f_0}(t, \cdot) \# f_0,$$

where $\#$ is our notation for the push-forward¹, and \mathcal{Z}_{f_0} is defined for all $t \in [0, T]$ and all $\mathbf{z} = (\mathbf{x}, v, w) \in \mathbb{R}^{d+2}$ through

$$\mathcal{Z}_{f_0}(t, \mathbf{z}) := (\mathbf{x}, \mathcal{V}_{f_0}(t, \mathbf{z}), \mathcal{W}_{f_0}(t, \mathbf{z})), \quad (2.1)$$

where $(\mathcal{V}_{f_0}, \mathcal{W}_{f_0})$ is a solution of the characteristic system associated to (1.5) for $t > 0$ and $\mathbf{z} = (\mathbf{x}, v, w) \in \mathbb{R}^{d+2}$:

$$\begin{cases} \partial_t \mathcal{V}(t, \mathbf{z}) = N(\mathcal{V}(t, \mathbf{z})) - \mathcal{W}(t, \mathbf{z}) - \int \Psi(\mathbf{x}, \mathbf{x}') (\mathcal{V}(t, \mathbf{z}) - \mathcal{V}(t, \mathbf{z}')) f_0(d\mathbf{x}', dv', dw'), \\ \partial_t \mathcal{W}(t, \mathbf{z}) = A(\mathcal{V}(t, \mathbf{z}), \mathcal{W}(t, \mathbf{z})), \\ \mathcal{V}(0, \mathbf{z}) = v, \quad \mathcal{W}(0, \mathbf{z}) = w. \end{cases} \quad (2.2)$$

97 **Remark 2.2.** *In the rest of this paper, for the sake of clarity, we will use the notation $\mathbf{z} = (\mathbf{x}, v, w) \in \mathbb{R}^{d+2}$
 98 and $\mathbf{z}' = (\mathbf{x}', v', w') \in \mathbb{R}^{d+2}$.*

As we can expect from this notion of measure solution to (1.5), the regularity of the solution to the characteristic system (2.2) is crucial. Therefore, we need to clearly define our framework for the connectivity kernel Ψ and the nonlinear function N to prove, on the one hand, the existence and uniqueness of solutions to the FHN system (1.3) and the kinetic equation (1.5), and on the other hand, to prove our result of mean-field limit. In the following, in the spirit of [4, 10], we choose a connectivity kernel $\Psi : \mathbb{R}^{2d} \rightarrow \mathbb{R}$ satisfying

$$\Psi \in \text{Lip}_b(\mathbb{R}^{2d}), \quad (2.3)$$

where $\text{Lip}_b(\mathbb{R}^{2d})$ is the set of bounded and globally Lipschitz continuous functions from \mathbb{R}^{2d} to \mathbb{R} . In the following, we note $L > 0$ the Lipschitz constant of the connectivity kernel Ψ , and

$$\|\Psi\|_\infty := \sup_{(\mathbf{x}, \mathbf{y})} |\Psi(\mathbf{x}, \mathbf{y})|.$$

Here, the Lipschitz continuity of the connectivity kernel is only a technical assumption. On the other hand, the choice of a bounded Ψ seems to be reasonable from a biological viewpoint, since currents transmitted

¹For all h , map from \mathbb{R}^{d+2} to itself, for all probability measure μ , the notation $\nu = h\#\mu$ is equivalent to

$$\int \mathbf{1}_B(z) \nu(dz) = \int \mathbf{1}_B(h(y)) \mu(dy)$$

for all B measurable subset of \mathbb{R}^{d+2} .

through synapses are bounded. We also choose a locally Lipschitz continuous nonlinearity $N : \mathbb{R} \rightarrow \mathbb{R}$ such that there exist two constants $\kappa, \tilde{\kappa} > 0$ satisfying:

$$\begin{cases} v N(v) \leq \kappa |v|^2 & \forall v \in \mathbb{R}, \\ (v - u)(N(v) - N(u)) \leq \tilde{\kappa} |v - u|^2 & \forall (v, u) \in \mathbb{R}^2. \end{cases} \quad (2.4)$$

99 Then, it remains to choose a suitable topology on $\mathcal{P}_2(\mathbb{R}^{d+2})$ to state our result of mean-field limit. The
 100 most convenient distance to describe the convergence of an empirical measure in such problems is the
 101 Wasserstein distance [4, 10, 22, 23], recalled in the next definition.

Definition 2.3 (Wasserstein distance of order 2). *Let μ and ν be two probability measures of \mathbb{R}^{d+2} with finite moments of second order. The Wasserstein distance of order 2 between μ and ν is defined by*

$$d_2(\mu, \nu) = \inf_{\pi \in \Lambda(\mu, \nu)} \left(\iint_{\mathbb{R}^{d+2} \times \mathbb{R}^{d+2}} \|(\mathbf{x} - \mathbf{x}', v - v', w - w')\|^2 \pi(d\mathbf{x}, dv, dw, d\mathbf{x}', dv', dw') \right)^{\frac{1}{2}},$$

where $\|(\mathbf{x}, v, w)\|$ stands for the euclidean norm of the vector (\mathbf{x}, v, w) in \mathbb{R}^{d+2} for all $\mathbf{x} \in \mathbb{R}^d$ and $(v, w) \in \mathbb{R}^2$, and $\Lambda(\mu, \nu)$ is the set of couplings of μ and ν , that is to say for all $\pi \in \Lambda(\mu, \nu)$, for all function $\phi \in \mathcal{C}(\mathbb{R}^{d+2})$ such that $\phi(\mathbf{z}) = \mathcal{O}(|\mathbf{z}|^2)$ as $|\mathbf{z}|$ goes to infinity,

$$\iint \phi(\mathbf{x}, v, w) \pi(d\mathbf{x}, dv, dw, d\mathbf{x}', dv', dw') = \int \phi(\mathbf{x}, v, w) \mu(d\mathbf{x}, dv, dw),$$

and

$$\iint \phi(\mathbf{x}', v', w') \pi(d\mathbf{x}, dv, dw, d\mathbf{x}', dv', dw') = \int \phi(\mathbf{x}', v', w') \nu(d\mathbf{x}', dv', dw').$$

102 In this paper, we choose to work with the Wasserstein distance of order 2 instead of 1 as in [10, 17, 22] to
 103 deal with the nonlinearity N , which naturally makes appear some moments of second order in v and w in
 104 the computation of the stability estimate.

Then, as explained in the introduction, to circumvent the issues caused by the class of interaction kernel we consider, even if it is not needed for the existence and uniqueness of a measure solution to (1.5), we make some assumptions on the initial data of the kinetic equation (1.5). Thus, we consider an initial data $f_0 \in \mathcal{P}_2(\mathbb{R}^{d+2})$ satisfying:

$$\int e^{\alpha_0 \langle v, w \rangle} f_0(d\mathbf{x}, dv, dw) < +\infty, \quad (2.5)$$

for some constant $\alpha_0 > 0$, using the notation for all $(v, w) \in \mathbb{R}^2$:

$$\langle v, w \rangle := (1 + |v|^2 + |w|^2)^{1/2}. \quad (2.6)$$

105 Now, we have all the tools we need to state our main theorem.

106 **Theorem 1** (Mean-field limit). *We consider a connectivity kernel Ψ satisfying (2.3), and a locally Lipschitz
 107 continuous nonlinearity N satisfying (2.4). Let $T > 0$.*

(i) Consider two initial data $f_{0,1}$ and $f_{0,2} \in \mathcal{P}_2(\mathbb{R}^{d+2})$ such that $d_2(f_{0,1}, f_{0,2}) < 1$. Assume that $f_{0,2}$ satisfies (2.5) for some constant $\alpha_0 > 0$. Further assume that there exist f_1 and $f_2 \in \mathcal{C}([0, T], \mathcal{P}_2(\mathbb{R}^{d+2}))$ two measure solutions of the kinetic equation (1.5) respectively with initial conditions $f_{0,1}$ and $f_{0,2}$. Then there exist two positive constants $C_T > 0$ and $K_T > 0$ such that for all $t \in [0, T]$,

$$d_2(f_1(t), f_2(t)) \leq K_T d_2(f_{0,1}, f_{0,2})^{\beta(t)}, \quad (2.7)$$

108 where $\beta(t) := e^{-C_T t}$.

(ii) For all $n \in \mathbb{N}$, consider the initial data $(\mathbf{x}_i, v_{0,i}, w_{0,i})_{1 \leq i \leq n} \in (\mathbb{R}^{d+2})^n$ and its associated empirical measure $f_{0,n}$. Consider an initial data $f_0 \in \mathcal{P}_2(\mathbb{R}^{d+2})$. Then, for all $n \in \mathbb{N}$, there exists a unique solution $(\mathbf{x}_i, v_i, w_i)_{1 \leq i \leq n} \in (\mathbb{R}^{d+2})^n$ of the FHN system (1.3) with initial data $(\mathbf{x}_i, v_{0,i}, w_{0,i})_{1 \leq i \leq n}$, and for all $t \in [0, T]$, we note its associated empirical measures $f_n(t)$. There also exists a unique measure solution $f \in \mathcal{C}([0, T], \mathcal{P}_2(\mathbb{R}^{d+2}))$ to (1.5) with initial data f_0 . Further assume that f_0 satisfies (2.5) for some positive constant $\alpha_0 > 0$ and that

$$\lim_{n \rightarrow +\infty} d_2(f_{0,n}, f_0) = 0. \quad (2.8)$$

Then, we get:

$$\lim_{n \rightarrow +\infty} \sup_{t \in [0, T]} d_2(f_n(t), f(t)) = 0. \quad (2.9)$$

109 The proof of Theorem 1 is postponed to Section 5. The first part (i) is the stability result of the measure
 110 solutions to (1.5) with respect to their initial data. Our approach follows the idea from [4, 10, 17, 22].
 111 Indeed, the main difficulty comes from the interaction kernel of the form $(\mathbf{x}, \mathbf{x}', v) \mapsto \Psi(\mathbf{x}, \mathbf{x}') v$, which is
 112 only locally Lipschitz continuous.

113 As for the second part (ii), the existence and uniqueness of the FHN system (1.3) follows from the Cauchy-
 114 Lipschitz Theorem, and the proof of the well-posedness of the kinetic equation (1.5) relies on the well-
 115 posedness of the characteristic system (2.2) using an classical arguments as in [22]. Then, the mean-field
 116 limit from (1.3) towards (1.5) is just a consequence of the first part (i).

117 **Remark 2.4.** We can extend the result of existence and uniqueness of the solution to (1.3) and the
 118 Definition 2.1 of measure solutions of (1.5) to the case where Ψ is only bounded and continuous with
 119 respect to its first variable uniformly relative to its second variable.

120 The rest of this paper is organized as follows. In Section 3, we prove an *a priori* estimate for the solution
 121 of the kinetic equation (1.5) which will be crucial for the proof of Theorem 1. Then, in Section 4, we
 122 present the proof of the existence and uniqueness of the measure solution of the kinetic equation (1.5).
 123 Furthermore, we prove our main result of mean-field limit in Section 5 and present as an application
 124 a stability result regarding monokinetic solutions in the following Section 6. Finally, in Section 7, we
 125 provide a numerical scheme for the kinetic equation (1.5) based on a particle method, and we display
 126 some numerical simulations, to illustrate the results established in Sections 5 and 6, and to show that this
 127 numerical scheme accurately reproduces the behavior of a large neural network.

3 *A priori* estimate

In this section, we prove an *a priori* estimate of the exponential moments of a solution of the kinetic equation (1.5). We will need to work with the characteristic system (2.2). In Section 4, we will prove that for all $T > 0$, the characteristics are well-defined in $\mathcal{C}([0, T], \mathcal{E})$, where

$$\mathcal{E} := \left\{ \mathcal{U} \in \mathcal{C}(\mathbb{R}^{d+2}, \mathbb{R}) \mid \|\mathcal{U}\|_{\mathcal{E}} < \infty \right\}, \quad (3.1)$$

with

$$\|\mathcal{U}\|_{\mathcal{E}} := \sup_{(\mathbf{x}, v, w) \in \mathbb{R}^{d+2}} \frac{|\mathcal{U}(\mathbf{x}, v, w)|}{\langle v, w \rangle}. \quad (3.2)$$

This choice of Banach space is justified since in general, the characteristics are not bounded, so we need to control some moments in v and w .

Lemma 3.1. *Let $T > 0$. We consider a connectivity kernel Ψ satisfying (2.3) and a locally Lipschitz continuous nonlinearity N satisfying (2.4). Let $p \geq 1$. Consider an initial data $f_0 \in \mathcal{P}_2(\mathbb{R}^{d+2})$ satisfying (2.5) for some $\alpha_0 > 0$ is a positive constant. Assume that there exists $f \in \mathcal{C}([0, T], \mathcal{P}_2(\mathbb{R}^{d+2}))$ a measure solution of (1.5) and a couple $(\mathcal{V}_{f_0}, \mathcal{W}_{f_0}) \in \mathcal{C}([0, T], \mathcal{E})^2$ solution of the characteristic system (2.2), such that if we define $\mathcal{Z}_{f_0} := (id_{\mathbb{R}^d}, \mathcal{V}_{f_0}, \mathcal{W}_{f_0})$, then for all $t \in [0, T]$,*

$$f(t) = \mathcal{Z}_{f_0}(t) \# f_0.$$

Then, there exists a constant $C_{f_0}^T > 0$ which depends only on the parameters of the equation (1.5), on T and on the moments of f_0 , and $\alpha_T := \alpha_0 e^{-p C_{f_0}^T T}$ such that for all $t \in [0, T]$,

$$\int e^{\alpha_T \langle v, w \rangle^p} f(t, d\mathbf{x}, dv, dw) \leq \int e^{\alpha_0 \langle v, w \rangle^p} f_0(d\mathbf{x}, dv, dw). \quad (3.3)$$

Proof. Consider $\alpha \in \mathcal{C}^1([0, T], \mathbb{R})$ a positive function to be determined later. Let $t \in [0, T]$. We have the following estimate:

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int e^{\alpha(t) \langle \mathcal{V}_{f_0}(t, \mathbf{z}), \mathcal{W}_{f_0}(t, \mathbf{z}) \rangle^p} f_0(d\mathbf{z}) &\leq \int \left[\frac{1}{2} \alpha'(t) \langle \mathcal{V}_{f_0}(t, \mathbf{z}), \mathcal{W}_{f_0}(t, \mathbf{z}) \rangle^p \right. \\ &\quad \left. + \frac{p}{2} \alpha(t) \langle \mathcal{V}_{f_0}(t, \mathbf{z}), \mathcal{W}_{f_0}(t, \mathbf{z}) \rangle^{p-2} (I_1 + I_2 + I_3) \right] e^{\alpha(t) \langle \mathcal{V}_{f_0}(t, \mathbf{z}), \mathcal{W}_{f_0}(t, \mathbf{z}) \rangle^p} f_0(d\mathbf{z}), \end{aligned}$$

where

$$\begin{cases} I_1 := \tau \mathcal{W}_{f_0}(t, \mathbf{z}) (\mathcal{V}_{f_0}(t, \mathbf{z}) + a - b \mathcal{W}_{f_0}(t, \mathbf{z})), \\ I_2 := \mathcal{V}_{f_0}(t, \mathbf{z}) (N(\mathcal{V}_{f_0}(t, \mathbf{z})) - \mathcal{W}_{f_0}(t, \mathbf{z})), \\ I_3 := -\mathcal{V}_{f_0}(t, \mathbf{z}) \int \Psi(\mathbf{x}, \mathbf{x}') [\mathcal{V}_{f_0}(t, \mathbf{z}) - \mathcal{V}_{f_0}(t, \mathbf{z}')] f_0(d\mathbf{z}'). \end{cases}$$

First of all, we can easily compute the first term I_1 with Young's inequality, which yields:

$$\begin{aligned} I_1 &\leq \frac{\tau}{2} |\mathcal{V}_{f_0}(t, \mathbf{z})|^2 + \tau |\mathcal{W}_{f_0}(t, \mathbf{z})|^2 + \frac{\tau a^2}{2} - \tau b |\mathcal{W}_{f_0}(t, \mathbf{z})|^2 \\ &\leq \tau \left(1 + \frac{a^2}{2} \right) |\langle \mathcal{V}_{f_0}(t, \mathbf{z}), \mathcal{W}_{f_0}(t, \mathbf{z}) \rangle|^2. \end{aligned}$$

Furthermore, we deal with the second term I_2 using the assumption on N (2.4) and Young's inequality:

$$\begin{aligned} I_2 &\leq \kappa |\mathcal{V}_{f_0}(t, \mathbf{z})|^2 + \frac{1}{2} (|\mathcal{V}_{f_0}(t, \mathbf{z})|^2 + |\mathcal{W}_{f_0}(t, \mathbf{z})|^2) \\ &\leq \left(\kappa + \frac{1}{2} \right) |\langle \mathcal{V}_{f_0}(t, \mathbf{z}), \mathcal{W}_{f_0}(t, \mathbf{z}) \rangle|^2. \end{aligned}$$

Then, we treat the third term I_3 using Young's inequality and then factorizing by $|\langle \mathcal{V}_{f_0}(t, \mathbf{z}), \mathcal{W}_{f_0}(t, \mathbf{z}) \rangle|^2$ as follows:

$$\begin{aligned} I_3 &\leq \frac{3}{2} |\mathcal{V}_{f_0}(t, \mathbf{z})|^2 \int |\Psi(\mathbf{x}, \mathbf{x}')| f_0(d\mathbf{z}') + \frac{1}{2} \int |\Psi(\mathbf{x}, \mathbf{x}')| |\mathcal{V}_{f_0}(t, \mathbf{z}')|^2 f_0(d\mathbf{z}') \\ &\leq \frac{1}{2} \|\Psi\|_\infty \left(3 |\mathcal{V}_{f_0}(t, \mathbf{z})|^2 + \int |\mathcal{V}_{f_0}(t, \mathbf{z}')|^2 f_0(d\mathbf{z}') \right) \\ &\leq \frac{1}{2} \|\Psi\|_\infty \left(3 + \sup_{s \in [0, T]} \|\mathcal{V}_{f_0}(s)\|_{\mathcal{E}}^2 \int |\langle v', w' \rangle|^2 f_0(d\mathbf{z}') \right) |\langle \mathcal{V}_{f_0}(t, \mathbf{z}), \mathcal{W}_{f_0}(t, \mathbf{z}) \rangle|^2. \end{aligned}$$

Finally, we get that there exists a positive constant $C_{f_0}^T > 0$ such that:

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \int e^{\alpha(t) \langle \mathcal{V}_{f_0}(t, \mathbf{z}), \mathcal{W}_{f_0}(t, \mathbf{z}) \rangle^p} f_0(d\mathbf{z}) \\ &\leq \int \left[\frac{1}{2} \alpha'(t) + \frac{p}{2} C_{f_0}^T \alpha(t) \right] \langle \mathcal{V}_{f_0}(t, \mathbf{z}), \mathcal{W}_{f_0}(t, \mathbf{z}) \rangle^p e^{\alpha(t) \langle \mathcal{V}_{f_0}(t, \mathbf{z}), \mathcal{W}_{f_0}(t, \mathbf{z}) \rangle^p} f_0(d\mathbf{z}). \end{aligned}$$

We choose for all $s \in [0, T]$, $\alpha(s) := \alpha_0 e^{-p C_{f_0}^T s}$, so that for all $s \in [0, T]$,

$$\alpha'(s) + p C_{f_0}^T \alpha(s) = 0, \quad \alpha(0) = \alpha_0.$$

Hence,

$$\frac{d}{dt} \int e^{\alpha(t) \langle \mathcal{V}_{f_0}(t, \mathbf{z}), \mathcal{W}_{f_0}(t, \mathbf{z}) \rangle^p} f_0(d\mathbf{z}) \leq 0.$$

To conclude the proof, it remains to define:

$$\alpha_T := \alpha(T),$$

and we deduce that for all $t \in [0, T]$,

$$\begin{aligned} \int e^{\alpha_T \langle \mathcal{V}_{f_0}(t, \mathbf{z}), \mathcal{W}_{f_0}(t, \mathbf{z}) \rangle^p} f_0(d\mathbf{z}) &\leq \int e^{\alpha(t) \langle \mathcal{V}_{f_0}(t, \mathbf{z}), \mathcal{W}_{f_0}(t, \mathbf{z}) \rangle^p} f_0(d\mathbf{z}) \\ &\leq \int e^{\alpha_0 \langle v, w \rangle^p} f_0(d\mathbf{z}). \end{aligned}$$

133

■

134 4 Proof of the well-posedness of the kinetic equation (1.5)

135 This section is devoted to the proof of existence and uniqueness of a measure solution to the kinetic equation
 136 (1.5), in the sense of Definition 2.1. Let $T > 0$ be a fixed final time and $f_0 \in \mathcal{P}_2(\mathbb{R}^{d+2})$. First, we focus on
 137 the well-posedness of the characteristic system (2.2), and then we will conclude by defining the measure
 138 solution to (1.5) as the push-forward of the initial data by the solution of the characteristic system.

139 **4.1 Proof of the well-posedness of the characteristic system (2.2)**

140 As a preliminary step, we establish the existence and uniqueness of the solution of the characteristic system
 141 (2.2) in $\mathcal{C}([0, T], \mathcal{E})$, where \mathcal{E} is defined with (3.1)-(3.2).

Proposition 4.1. *Let $T > 0$. Consider an initial data $f_0 \in \mathcal{P}_2(\mathbb{R}^{d+2})$. Then, there exists a unique couple $(\mathcal{V}_{f_0}, \mathcal{W}_{f_0})$ solution of (2.2) on $[0, T]$ such that*

$$\mathcal{V}_{f_0}, \mathcal{W}_{f_0} \in \mathcal{C}([0, T], \mathcal{E}).$$

142 **Proof.** Since we cannot directly conclude with the Cauchy-Lipschitz theorem because of the term resulting
 143 from the nonlocal interactions in (2.2), our approach is based on the construction of a Cauchy sequence
 144 $(\mathcal{V}_p, \mathcal{W}_p)_{p \in \mathbb{N}}$ in $\mathcal{C}([0, T], \mathcal{E})$, in order to circumvent this difficulty. Then, we will define the couple $(\mathcal{V}_{f_0}, \mathcal{W}_{f_0})$
 145 as its limit as p tends to infinity.

146 **Step 1: construction of the sequences**

147 First, we prove the following lemma, which yields the existence and uniqueness of the solution of a system
 148 of equations approximating (2.2), in which we consider the contribution of the interactions as a source
 149 term.

Lemma 4.2. *Let $T > 0$. Under the assumptions of Proposition 4.1, assume that there exists $\mathcal{U} \in \mathcal{C}([0, T], \mathcal{E})$ of class \mathcal{C}^1 in time. Then, there exists unique couple $(\mathcal{V}, \mathcal{W}) \in \mathcal{C}([0, T], \mathcal{E})^2$ solution of class \mathcal{C}^1 in time of the following system for $t > 0$ and $\mathbf{z} = (\mathbf{x}, v, w) \in \mathbb{R}^{d+2}$:*

$$\begin{cases} \partial_t \mathcal{V}(t, \mathbf{z}) = N(\mathcal{V}(t, \mathbf{z})) - \mathcal{W}(t, \mathbf{z}) - \int \Psi(\mathbf{x}, \mathbf{x}') (\mathcal{U}(t, \mathbf{z}) - \mathcal{U}(t, \mathbf{z}')) f_0(d\mathbf{z}'), \\ \partial_t \mathcal{W}(t, \mathbf{z}) = A(\mathcal{V}(t, \mathbf{z}), \mathcal{W}(t, \mathbf{z})), \\ \mathcal{V}(0, \mathbf{z}) = v, \quad \mathcal{W}(0, \mathbf{z}) = w. \end{cases} \quad (4.1)$$

The proof of Lemma 4.2 only relies on classical arguments, but for the sake of completeness, it is postponed to the Appendix A. Then, by induction, Lemma 4.2 implies the existence and uniqueness of a sequence $(\mathcal{V}_p, \mathcal{W}_p)_{p \in \mathbb{N}}$, such that $(\mathcal{V}_0, \mathcal{W}_0) := (0, 0)$, and for all $p \in \mathbb{N}$, for all $t \in [0, T]$ and $\mathbf{z} = (\mathbf{x}, v, w) \in \mathbb{R}^{d+2}$,

$$\begin{cases} \mathcal{V}_{p+1}(t, \mathbf{z}) = v + \int_0^t \left[N(\mathcal{V}_{p+1}(s, \mathbf{z})) - \mathcal{W}_{p+1}(s, \mathbf{z}) - \int \Psi(\mathbf{x}, \mathbf{x}') (\mathcal{V}_p(s, \mathbf{z}) - \mathcal{V}_p(s, \mathbf{z}')) f_0(d\mathbf{z}') \right] ds, \\ \mathcal{W}_{p+1}(s, \mathbf{z}) = w + \int_0^s A(\mathcal{V}_{p+1}(s, \mathbf{z}), \mathcal{W}_{p+1}(s, \mathbf{z})) ds. \end{cases} \quad (4.2)$$

150 **Step 2: Cauchy sequences**

Now, we want to prove that for all $t \in [0, T]$, $\{\mathcal{V}_p(t)\}_{p \in \mathbb{N}}$ and $\{\mathcal{W}_p(t)\}_{p \in \mathbb{N}}$ are two Cauchy sequences in \mathcal{E} . For all $p \in \mathbb{N}$, we define for all $\mathbf{z} \in \mathbb{R}^{d+2}$:

$$G_{p+1}(t) := \left(\|\mathcal{V}_{p+1}(t) - \mathcal{V}_p(t)\|_{\mathcal{E}}^2 + \|\mathcal{W}_{p+1}(t) - \mathcal{W}_p(t)\|_{\mathcal{E}}^2 \right)^{\frac{1}{2}},$$

and we want to prove by induction that this quantity is summable. Let $p \in \mathbb{N}$, $t \in [0, T]$ and $\mathbf{z} = (\mathbf{x}, v, w)$. Thus, we have:

$$\frac{1}{2} \left(|\mathcal{V}_{p+2}(t, \mathbf{z}) - \mathcal{V}_{p+1}(t, \mathbf{z})|^2 + |\mathcal{W}_{p+2}(t, \mathbf{z}) - \mathcal{W}_{p+1}(t, \mathbf{z})|^2 \right) \leq \int [\mathcal{T}_1(s) + \mathcal{T}_2(s) + \mathcal{T}_3(s) + \mathcal{T}_4(s)] ds,$$

where for all $s \in [0, t]$,

$$\left\{ \begin{array}{l} \mathcal{T}_1(s) := \tau (\mathcal{W}_{p+2}(s, \mathbf{z}) - \mathcal{W}_{p+1}(s, \mathbf{z})) [(\mathcal{V}_{p+2}(s, \mathbf{z}) - \mathcal{V}_{p+1}(s, \mathbf{z})) - b (\mathcal{W}_{p+2}(s, \mathbf{z}) - \mathcal{W}_{p+1}(s, \mathbf{z}))], \\ \mathcal{T}_2(s) := (\mathcal{V}_{p+2}(s, \mathbf{z}) - \mathcal{V}_{p+1}(s, \mathbf{z})) [N(\mathcal{V}_{p+2}(s, \mathbf{z})) - N(\mathcal{V}_{p+1}(s, \mathbf{z}))], \\ \mathcal{T}_3(s) := - (\mathcal{V}_{p+2}(s, \mathbf{z}) - \mathcal{V}_{p+1}(s, \mathbf{z})) (\mathcal{W}_{p+2}(s, \mathbf{z}) - \mathcal{W}_{p+1}(s, \mathbf{z})), \\ \mathcal{T}_4(s) := - (\mathcal{V}_{p+2}(s, \mathbf{z}) - \mathcal{V}_{p+1}(s, \mathbf{z})) \\ \quad \times \int \Psi(\mathbf{x}, \mathbf{x}') [(\mathcal{V}_{p+1}(s, \mathbf{z}) - \mathcal{V}_{p+1}(s, \mathbf{z}')) - (\mathcal{V}_p(s, \mathbf{z}) - \mathcal{V}_p(s, \mathbf{z}'))] f_0(d\mathbf{z}'). \end{array} \right.$$

151 Let $s \in [0, t]$. The first term $\mathcal{T}_1(s)$ is easily controlled using Young's inequality:

$$\begin{aligned} \mathcal{T}_1(s) &\leq \frac{\tau}{2} \left(|\mathcal{V}_{p+2}(s, \mathbf{z}) - \mathcal{V}_{p+1}(s, \mathbf{z})|^2 + |\mathcal{W}_{p+2}(s, \mathbf{z}) - \mathcal{W}_{p+1}(s, \mathbf{z})|^2 \right) \\ &\leq \frac{\tau}{2} |\langle v, w \rangle|^2 |G_{p+2}(s)|^2 \end{aligned} \quad (4.3)$$

Then, we treat the second term $\mathcal{T}_2(s)$ using assumption (2.4) satisfied by N :

$$\mathcal{T}_2(s) \leq \tilde{\kappa} |\mathcal{V}_{p+2}(s, \mathbf{z}) - \mathcal{V}_{p+1}(s, \mathbf{z})|^2 \leq \tilde{\kappa} |\langle v, w \rangle|^2 |G_{p+2}(s)|^2. \quad (4.4)$$

152 Furthermore, we can estimate the third term $\mathcal{T}_3(s)$ with Young's inequality:

$$\begin{aligned} \mathcal{T}_3(s) &\leq \frac{1}{2} \left(|\mathcal{V}_{p+2}(s, \mathbf{z}) - \mathcal{V}_{p+1}(s, \mathbf{z})|^2 + |\mathcal{W}_{p+2}(s, \mathbf{z}) - \mathcal{W}_{p+1}(s, \mathbf{z})|^2 \right) \\ &\leq \frac{1}{2} |\langle v, w \rangle|^2 |G_{p+2}(s)|^2. \end{aligned} \quad (4.5)$$

Finally, to deal with the nonlocal term $\mathcal{T}_4(s)$, using the boundedness of Ψ and Young's inequality, we can compute:

$$\begin{aligned} \mathcal{T}_4(s) &= (\mathcal{V}_{p+2}(s, \mathbf{z}) - \mathcal{V}_{p+1}(s, \mathbf{z})) \\ &\quad \times \int \Psi(\mathbf{x}, \mathbf{x}') [(\mathcal{V}_{p+1}(s, \mathbf{z}') - \mathcal{V}_p(s, \mathbf{z}')) - (\mathcal{V}_{p+1}(s, \mathbf{z}) - \mathcal{V}_p(s, \mathbf{z}))] f_0(d\mathbf{z}') \\ &\leq \frac{1}{2} |\mathcal{V}_{p+2}(s, \mathbf{z}) - \mathcal{V}_{p+1}(s, \mathbf{z})|^2 \\ &\quad + \frac{1}{2} \|\Psi\|_\infty^2 \int \left(|\mathcal{V}_{p+1}(s, \mathbf{z}') - \mathcal{V}_p(s, \mathbf{z}')|^2 + |\mathcal{V}_{p+1}(s, \mathbf{z}) - \mathcal{V}_p(s, \mathbf{z})|^2 \right) f_0(d\mathbf{z}'). \end{aligned}$$

153 Moreover, factorizing each term with $|\langle v, w \rangle|^2$, we get:

$$\begin{aligned} \mathcal{T}_4(s) &\leq \frac{1}{2} |\langle v, w \rangle|^2 \|\mathcal{V}_{p+2}(s) - \mathcal{V}_{p+1}(s)\|_{\mathcal{E}}^2 \\ &\quad + \frac{1}{2} \|\Psi\|_\infty^2 \left(\int |\langle v', w' \rangle|^2 f_0(d\mathbf{z}') + |\langle v, w \rangle|^2 \right) \|\mathcal{V}_{p+1}(s) - \mathcal{V}_p(s)\|_{\mathcal{E}}^2 \\ &\leq \frac{1}{2} |\langle v, w \rangle|^2 \left[|G_{p+2}(s)|^2 + \|\Psi\|_\infty^2 \left(\int |\langle v', w' \rangle|^2 f_0(d\mathbf{z}') + 1 \right) |G_{p+1}(s)|^2 \right]. \end{aligned} \quad (4.6)$$

Finally, (4.3), (4.4), (4.5) and (4.6) together yield that there exist two positive constants $C_1 > 0$ and $C_2 > 0$ independent of p such that:

$$\begin{aligned} |\mathcal{V}_{p+2}(t, \mathbf{z}) - \mathcal{V}_{p+1}(t, \mathbf{z})|^2 + |\mathcal{W}_{p+2}(t, \mathbf{z}) - \mathcal{W}_{p+1}(t, \mathbf{z})|^2 \\ \leq |\langle v, w \rangle|^2 \left[C_1 \int_0^t |G_{p+1}(s)|^2 ds + C_2 \int_0^t |G_{p+2}(s)|^2 ds \right], \end{aligned}$$

which implies, by dividing this inequality by $|\langle v, w \rangle|^2$ and then taking the supremum on \mathbb{R}^{d+2} :

$$|G_{p+2}(t)|^2 \leq C_1 \int_0^t |G_{p+1}(s)|^2 ds + C_2 \int_0^t |G_{p+2}(s)|^2 ds. \quad (4.7)$$

On the one hand, using Grönwall's lemma, we have for all $t \in [0, T]$:

$$|G_{p+2}(t)|^2 \leq C_1 e^{C_2 T} \int_0^t |G_{p+1}(s)|^2 ds. \quad (4.8)$$

On the other hand, for all $t \in [0, T]$,

$$|G_1(t)|^2 = \|\mathcal{V}_1(t)\|_{\mathcal{E}}^2 + \|\mathcal{W}_1(t)\|_{\mathcal{E}}^2 \leq C_T, \quad (4.9)$$

for some constant $C_T > 0$, since \mathcal{V}_1 and $\mathcal{W}_1 \in \mathcal{C}([0, T], \mathcal{E})$ according to Lemma 4.2. Hence, by induction, we can deduce from (4.8) and (4.9) that for all $p \in \mathbb{N}$ and all $t \in [0, T]$:

$$|G_{p+1}(t)|^2 \leq C_T \frac{(C_1 e^{C_2 T} t)^p}{p!}, \quad (4.10)$$

154 which is summable. Consequently, for all $t \in [0, T]$, $\{\mathcal{V}_p(t)\}_{p \in \mathbb{N}}$ and $\{\mathcal{W}_p(t)\}_{p \in \mathbb{N}}$ are Cauchy sequences
 155 in \mathcal{E} . Since \mathcal{E} is a Banach space, and since for all $p \in \mathbb{N}$, \mathcal{V}_p and $\mathcal{W}_p \in \mathcal{C}([0, T], \mathcal{E})$, there exist \mathcal{V}_{f_0}
 156 and $\mathcal{W}_{f_0} \in \mathcal{C}([0, T], \mathcal{E})$ such that for all $t \in [0, T]$, $\mathcal{V}_p(t, \cdot)$ (respectively \mathcal{W}_p) converges towards $\mathcal{V}_{f_0}(t, \cdot)$
 157 (respectively \mathcal{W}_{f_0}) uniformly in \mathcal{E} . Thus, passing to the limit $p \rightarrow +\infty$ in (4.2), we get that for all $\mathbf{z} \in \mathbb{R}^{d+2}$,
 158 $(\mathcal{V}_{f_0}(\cdot, \mathbf{z}), \mathcal{W}_{f_0}(\cdot, \mathbf{z}))$ is a solution of (2.2) of class \mathcal{C}^1 in time.

159 Step 3: Uniqueness

Now, we want to check that the solution of (2.2) is unique. Suppose that there exist $\mathcal{Z}_1 := (\text{id}_{\mathbb{R}^d}, \mathcal{V}_1, \mathcal{W}_1)$
 and $\mathcal{Z}_2 := (\text{id}_{\mathbb{R}^d}, \mathcal{V}_2, \mathcal{W}_2)$ such that $(\mathcal{V}_1, \mathcal{W}_1)$ and $(\mathcal{V}_2, \mathcal{W}_2)$ are two solutions of (2.2) in $\mathcal{C}([0, T], \mathcal{E})^2$. We
 define for all $t \in [0, T]$:

$$G(t) := \left(\|\mathcal{V}_1(t) - \mathcal{V}_2(t)\|_{\mathcal{E}}^2 + \|\mathcal{W}_1(t) - \mathcal{W}_2(t)\|_{\mathcal{E}}^2 \right)^{\frac{1}{2}}. \quad (4.11)$$

Thus, using similar computations as previously, we get that there exists a positive constant C such that
 for all $t \in [0, T]$:

$$|G(t)|^2 \leq C \int_0^t |G(s)|^2 ds.$$

160 Using Grönwall's inequality we get that $\mathcal{Z}_1 = \mathcal{Z}_2$.

161

■

162 **4.2 Construction of the measure solution to (1.5)**

We have proved so far that there exists a unique map $\mathcal{Z}_{f_0} := (\text{id}_{\mathbb{R}^d}, \mathcal{V}_{f_0}, \mathcal{W}_{f_0})$ such that $(\mathcal{V}_{f_0}, \mathcal{W}_{f_0})$ is a solution of the characteristic system (2.2) in $\mathcal{C}([0, T], \mathcal{E})$. We define:

$$f : t \mapsto \mathcal{Z}_{f_0}(t) \# f_0. \quad (4.12)$$

Thus, for all $t \in [0, T]$:

$$\begin{aligned} \int (|\mathbf{x}|^2 + |v|^2 + |w|^2) f(t, d\mathbf{z}) &= \int (|\mathbf{x}|^2 + |\mathcal{V}_{f_0}(t, \mathbf{z})|^2 + |\mathcal{W}_{f_0}(t, \mathbf{z})|^2) f_0(d\mathbf{z}) \\ &\leq \int |\mathbf{x}|^2 f_0(d\mathbf{z}) + (\|\mathcal{V}_{f_0}(t)\|_{\mathcal{E}}^2 + \|\mathcal{W}_{f_0}(t)\|_{\mathcal{E}}^2) \int |\langle v, w \rangle|^2 f_0(d\mathbf{z}), \end{aligned}$$

which is uniformly bounded. Hence, for all $t \in [0, T]$:

$$f(t) \in \mathcal{P}_2(\mathbb{R}^{d+2}).$$

Then, we want to prove that $f \in \mathcal{C}([0, T], \mathcal{P}_2(\mathbb{R}^{d+2}))$, where $\mathcal{P}_2(\mathbb{R}^{d+2})$ is equipped with the Wasserstein distance d_2 . Let t and $t' \in [0, T]$. Notice that the measure $(\mathcal{Z}_{f_0}(t) \times \mathcal{Z}_{f_0}(t')) \# f_0 \in \Lambda(f(t), f(t'))$. Therefore, we have:

$$\begin{aligned} d_2^2(f(t), f(t')) &= \inf_{\pi \in \Lambda(f(t), f(t'))} \iint |\mathbf{z}_1 - \mathbf{z}_2|^2 \pi(d\mathbf{z}_1 d\mathbf{z}_2) \\ &\leq \int |\mathcal{Z}_{f_0}(t, \mathbf{z}) - \mathcal{Z}_{f_0}(t', \mathbf{z})|^2 f_0(d\mathbf{z}) \\ &\leq (\|\mathcal{V}_{f_0}(t) - \mathcal{V}_{f_0}(t')\|_{\mathcal{E}}^2 + \|\mathcal{W}_{f_0}(t) - \mathcal{W}_{f_0}(t')\|_{\mathcal{E}}^2) \int |\langle v, w \rangle|^2 f_0(d\mathbf{z}). \end{aligned}$$

Since \mathcal{V}_{f_0} and $\mathcal{W}_{f_0} \in \mathcal{C}([0, T], \mathcal{E})$, and $f_0 \in \mathcal{P}_2(\mathbb{R}^{d+2})$, we have that:

$$f \in \mathcal{C}([0, T], \mathcal{P}_2(\mathbb{R}^{d+2})).$$

163 Hence, f is a measure solution of (1.5) in the sense of definition 2.1.

164 **5 Proof of Theorem 1**

165 This section is devoted to the proof of our main result Theorem 1. We start with the stability result (i).
 166 Our approach consists in using an estimate of the Wasserstein distance between two measure solutions of
 167 (1.5). Then, we will show how this stability result implies the mean-field limit result (ii).

168 **5.1 Proof of the stability result**

For $j \in \{1, 2\}$, we define the map \mathcal{Z}_j such that for all $t \in [0, T]$ and all $\mathbf{z} = (\mathbf{x}, v, w) \in \mathbb{R}^{d+2}$:

$$\mathcal{Z}_j(t, \mathbf{z}) := (\mathbf{x}, \mathcal{V}_j(t, \mathbf{z}), \mathcal{W}_j(t, \mathbf{z})),$$

where $(\mathcal{V}_j, \mathcal{W}_j)$ is the solution of the characteristic system (2.2) with initial data $f_{0,j}$. \mathcal{V}_j and \mathcal{W}_j are in $\mathcal{C}([0, T], \mathcal{E})$ according to Proposition 4.1. Let $\pi \in \Lambda(f_{0,1}, f_{0,2})$ be the optimal measure to compute $d_2(f_{0,1}, f_{0,2})$, that is

$$d_2(f_{0,1}, f_{0,2}) = \left(\iint \|\mathbf{z}_1 - \mathbf{z}_2\|^2 \pi(d\mathbf{z}'_1, d\mathbf{z}'_2) \right)^{\frac{1}{2}}.$$

The existence of such a minimizing measure is proved in [39]. First, we want to estimate the function $D[\pi]$ defined for all $t \in [0, T]$ with

$$D[\pi](t) := \left(\iint \|\mathcal{Z}_1(t, \mathbf{z}'_1) - \mathcal{Z}_2(t, \mathbf{z}'_2)\|^2 \pi(d\mathbf{z}'_1, d\mathbf{z}'_2) \right)^{\frac{1}{2}}.$$

Then, we will be able to conclude using the fact that for all $t \in [0, T]$,

$$d_2(f_1(t), f_2(t)) \leq D[\pi](t).$$

169 **Step 1: Estimate of $D[\pi]$**

Let $t \in [0, T]$, $\mathbf{z}_1 = (\mathbf{x}_1, v_1, w_1)$ and $\mathbf{z}_2 = (\mathbf{x}_2, v_2, w_2) \in \mathbb{R}^{d+2}$. We start by estimating the integrand $\|\mathcal{Z}_1(t, \mathbf{z}_1) - \mathcal{Z}_2(t, \mathbf{z}_2)\|^2$. Then, we will integrate with respect to the measure $\pi(d\mathbf{z}_1, d\mathbf{z}_2)$ in order to estimate $D[\pi]^2(t)$. In the following, for $j \in \{1, 2\}$, we use the shorthand notations $\mathcal{V}_j := \mathcal{V}_j(t, \mathbf{z}_j)$ and $\mathcal{V}'_j := \mathcal{V}_j(t, \mathbf{z}'_j)$, and the same for \mathcal{W}_j . First, we have:

$$\frac{1}{2} \frac{d}{dt} \|\mathcal{Z}_1(t, \mathbf{z}_1) - \mathcal{Z}_2(t, \mathbf{z}_2)\|^2 \leq \mathcal{T}_{l,1} + \mathcal{T}_{l,2} + \mathcal{T}_{nl}, \quad (5.1)$$

where

$$\begin{cases} \mathcal{T}_{l,1} := \tau(\mathcal{W}_1 - \mathcal{W}_2) [(\mathcal{V}_1 - \mathcal{V}_2) - b(\mathcal{W}_1 - \mathcal{W}_2)], \\ \mathcal{T}_{l,2} := (\mathcal{V}_1 - \mathcal{V}_2) [(N(\mathcal{V}_1) - N(\mathcal{V}_2)) - (\mathcal{W}_1 - \mathcal{W}_2)], \\ \mathcal{T}_{nl} := (\mathcal{V}_1 - \mathcal{V}_2) \left[\int \Psi(\mathbf{x}_1, \mathbf{x}'_1) (\mathcal{V}_1 - \mathcal{V}'_1) f_{0,1}(d\mathbf{z}'_1) - \int \Psi(\mathbf{x}_2, \mathbf{x}'_2) (\mathcal{V}_2 - \mathcal{V}'_2) f_{0,2}(d\mathbf{z}'_2) \right]. \end{cases}$$

By integrating (5.1) with respect to the measure $\pi(d\mathbf{z}_1, d\mathbf{z}_2)$, we get:

$$\frac{1}{2} \frac{d}{dt} D[\pi](t)^2 \leq \iint [\mathcal{T}_{l,1} + \mathcal{T}_{l,2} + \mathcal{T}_{nl}] \pi(d\mathbf{z}_1, d\mathbf{z}_2). \quad (5.2)$$

170 We easily treat the first local term $\mathcal{T}_{l,1}$ using Young's inequality:

$$\begin{aligned} \iint \mathcal{T}_{l,1} \pi(d\mathbf{z}_1, d\mathbf{z}_2) &\leq \frac{\tau}{2} \iint (|\mathcal{V}_1 - \mathcal{V}_2|^2 + |\mathcal{W}_1 - \mathcal{W}_2|^2) \pi(d\mathbf{z}_1, d\mathbf{z}_2) - \tau b \iint |\mathcal{W}_1 - \mathcal{W}_2|^2 \pi(d\mathbf{z}_1, d\mathbf{z}_2) \\ &\leq \frac{\tau}{2} D[\pi](t)^2. \end{aligned} \quad (5.3)$$

171 Then, we can estimate the second local term $\mathcal{T}_{l,2}$ using Young's inequality and the assumption on N (2.4)
172 as follows:

$$\begin{aligned} \iint \mathcal{T}_{l,2} \pi(d\mathbf{z}_1, d\mathbf{z}_2) &\leq \iint \left[\tilde{\kappa} |\mathcal{V}_1 - \mathcal{V}_2|^2 + \frac{1}{2} |\mathcal{V}_1 - \mathcal{V}_2|^2 + \frac{1}{2} |\mathcal{W}_1 - \mathcal{W}_2|^2 \right] \pi(d\mathbf{z}_1, d\mathbf{z}_2) \\ &\leq \left(\frac{1}{2} + \tilde{\kappa} \right) \iint \|\mathcal{Z}_1(t, \mathbf{z}_1) - \mathcal{Z}_2(t, \mathbf{z}_2)\|^2 \pi(d\mathbf{z}_1, d\mathbf{z}_2) \\ &= \left(\frac{1}{2} + \tilde{\kappa} \right) D[\pi](t)^2. \end{aligned} \quad (5.4)$$

Now, we deal with the nonlocal terms \mathcal{T}_{nl} . Since we have:

$$\begin{aligned} & \Psi(\mathbf{x}_1, \mathbf{x}'_1) (\mathcal{V}_1 - \mathcal{V}'_1) + \Psi(\mathbf{x}_2, \mathbf{x}'_2) (\mathcal{V}_2 - \mathcal{V}'_2) \\ &= \Psi(\mathbf{x}_1, \mathbf{x}'_1) [(\mathcal{V}_1 - \mathcal{V}_2) - (\mathcal{V}'_1 - \mathcal{V}'_2)] + (\Psi(\mathbf{x}_1, \mathbf{x}'_1) - \Psi(\mathbf{x}_2, \mathbf{x}'_2)) (\mathcal{V}_2 - \mathcal{V}'_2), \end{aligned}$$

we get:

$$\iint \mathcal{T}_{nl} \pi(d\mathbf{z}_1, d\mathbf{z}_2) \leq J_1 + J_2 + J_3,$$

where

$$\begin{cases} J_1 := \iiint \iint |\Psi(\mathbf{x}_1, \mathbf{x}'_1)| |\mathcal{V}_1 - \mathcal{V}_2|^2 \pi(d\mathbf{z}'_1, d\mathbf{z}'_2) \pi(d\mathbf{z}_1, d\mathbf{z}_2), \\ J_2 := \iiint \iint |\Psi(\mathbf{x}_1, \mathbf{x}'_1)| |\mathcal{V}_1 - \mathcal{V}_2| |\mathcal{V}'_1 - \mathcal{V}'_2| \pi(d\mathbf{z}'_1, d\mathbf{z}'_2) \pi(d\mathbf{z}_1, d\mathbf{z}_2), \\ J_3 := \iiint \iint |\Psi(\mathbf{x}_1, \mathbf{x}'_1) - \Psi(\mathbf{x}_2, \mathbf{x}'_2)| |\mathcal{V}_1 - \mathcal{V}_2| |\mathcal{V}_2 - \mathcal{V}'_2| \pi(d\mathbf{z}'_1, d\mathbf{z}'_2) \pi(d\mathbf{z}_1, d\mathbf{z}_2). \end{cases}$$

The two first terms J_1 and J_2 are easy to estimate, using the fact that Ψ is bounded and Young's inequality. We find:

$$\begin{cases} J_1 \leq \|\Psi\|_\infty D[\pi](t)^2, \\ J_2 \leq \|\Psi\|_\infty D[\pi](t)^2. \end{cases}$$

Furthermore, since Ψ satisfies the assumption (2.3), we have for all $(\mathbf{x}_1, \mathbf{x}'_1)$ and $(\mathbf{x}_2, \mathbf{x}'_2) \in \mathbb{R}^{2d}$:

$$|\Psi(\mathbf{x}_1, \mathbf{x}'_1) - \Psi(\mathbf{x}_2, \mathbf{x}'_2)| \leq \min \{L (|\mathbf{x}_1 - \mathbf{x}_2| + |\mathbf{x}'_1 - \mathbf{x}'_2|), 2\|\Psi\|_\infty\}. \quad (5.5)$$

Let $R(t) > 0$ to be determined later. We define the sets

$$\begin{cases} \varepsilon_{R(t)} := \{\mathbf{z} \in \mathbb{R}^{d+2}, |\mathcal{V}_2(t, \mathbf{z})| \leq R(t)\}, \\ \Theta_{R(t)} := (\mathbb{R}^{d+2} \times \varepsilon_{R(t)})^2. \end{cases} \quad (5.6)$$

Then, we can deal with the third term J_3 using (5.5) and then splitting the set of integration into $\Theta_{R(t)}$ and its complementary $\Theta_{R(t)}^C$:

$$\begin{aligned} J_3 &\leq \iiint \iint \min \{L (|\mathbf{x}_1 - \mathbf{x}_2| + |\mathbf{x}'_1 - \mathbf{x}'_2|), 2\|\Psi\|_\infty\} |\mathcal{V}_1 - \mathcal{V}_2| |\mathcal{V}_2 - \mathcal{V}'_2| \pi(d\mathbf{z}'_1, d\mathbf{z}'_2) \pi(d\mathbf{z}_1, d\mathbf{z}_2) \\ &\leq J_{31} + J_{32}, \end{aligned}$$

where

$$\begin{cases} J_{31} := L \iiint \iint_{\Theta_{R(t)}} |\mathcal{V}_1 - \mathcal{V}_2| (|\mathbf{x}_1 - \mathbf{x}_2| + |\mathbf{x}'_1 - \mathbf{x}'_2|) |\mathcal{V}_2 - \mathcal{V}'_2| \pi(d\mathbf{z}'_1, d\mathbf{z}'_2) \pi(d\mathbf{z}_1, d\mathbf{z}_2), \\ J_{32} := 2\|\Psi\|_\infty \iiint \iint_{\Theta_{R(t)}^C} |\mathcal{V}_1 - \mathcal{V}_2| |\mathcal{V}_2 - \mathcal{V}'_2| \pi(d\mathbf{z}'_1, d\mathbf{z}'_2) \pi(d\mathbf{z}_1, d\mathbf{z}_2). \end{cases}$$

We can treat the term J_{31} with the Cauchy-Schwarz inequality:

$$\begin{aligned}
J_{31} &\leq 2LR(t) \iint_{\Theta_{R(t)}} \iint |\mathcal{V}_1 - \mathcal{V}_2| (|\mathbf{x}_1 - \mathbf{x}_2| + |\mathbf{x}'_1 - \mathbf{x}'_2|) \pi(d\mathbf{z}'_1, d\mathbf{z}'_2) \pi(d\mathbf{z}_1, d\mathbf{z}_2) \\
&\leq 2LR(t) D[\pi](t)^2 + 2LR(t) \left(\iint |\mathcal{V}_1 - \mathcal{V}_2|^2 \pi(d\mathbf{z}_1, d\mathbf{z}_2) \right)^{\frac{1}{2}} \left(\iint |\mathbf{x}'_1 - \mathbf{x}'_2|^2 \pi(d\mathbf{z}'_1, d\mathbf{z}'_2) \right)^{\frac{1}{2}} \\
&\leq 4LR(t) D[\pi](t)^2.
\end{aligned}$$

Then, using the Young and Cauchy-Schwarz inequalities, we can estimate the second term $J_{32}(s)$ as follows:

$$\begin{aligned}
J_{32}(s) &\leq \|\Psi\|_\infty D[\pi](t)^2 + \|\Psi\|_\infty \iint_{\Theta_{R(t)}^C} \iint |\mathcal{V}_2 - \mathcal{V}'_2|^2 \pi(d\mathbf{z}'_1, d\mathbf{z}'_2) \pi(d\mathbf{z}_1, d\mathbf{z}_2) \\
&\leq \|\Psi\|_\infty D[\pi](t)^2 \\
&\quad + \|\Psi\|_\infty \left(\iint \iint |\mathcal{V}_2 - \mathcal{V}'_2|^4 \pi(d\mathbf{z}'_1, d\mathbf{z}'_2) \pi(d\mathbf{z}_1, d\mathbf{z}_2) \right)^{\frac{1}{2}} \left(\iint \iint \pi(d\mathbf{z}'_1, d\mathbf{z}'_2) \pi(d\mathbf{z}_1, d\mathbf{z}_2) \right)^{\frac{1}{2}} \\
&\leq \|\Psi\|_\infty D[\pi](t)^2 + C \left(\int_{\Theta_{R(t)}^C} |\mathcal{V}_2|^4 f_{0,2}(d\mathbf{z}_2) \right)^{\frac{1}{2}} \left(\int_{\Theta_{R(t)}^C} f_{0,2}(d\mathbf{z}_2) \right)^{\frac{1}{2}}.
\end{aligned}$$

On the one hand, according to Lemma 3.1, there exists a constant $\alpha_T > 0$ such that we have:

$$\int_{\Theta_{R(t)}^C} f_{0,2}(d\mathbf{z}_2) \leq \int_{\Theta_{R(t)}^C} \frac{e^{\alpha_T |\mathcal{V}_2|}}{e^{\alpha_T R(t)}} f_{0,2}(d\mathbf{z}_2) \leq e^{-\alpha_T R(t)} \int e^{\alpha_0 \langle v_2, w_2 \rangle} f_{0,2}(d\mathbf{z}_2).$$

On the other hand, there exists a constant $C > 0$ such that

$$\int |\mathcal{V}_2|^4 f_{0,2}(d\mathbf{z}_2) \leq C \int (1 + e^{\alpha_T |\mathcal{V}_2|}) f_{0,2}(d\mathbf{z}_2) < \infty,$$

according to Lemma 3.1. Finally, there exists a constant $C > 0$ such that

$$J_{32} \leq C(1 + R(t)) D[\pi](t)^2 + C e^{-\frac{\alpha_T}{2} R(t)}.$$

and consequently, there exists a constant $\tilde{C} > 0$ such that

$$\iint \mathcal{T}_{nl} \pi(d\mathbf{z}_1 d\mathbf{z}_2) \leq \tilde{C}(1 + R(t)) D[\pi](t)^2 + \tilde{C} e^{-\frac{\alpha_T}{2} R(t)}. \quad (5.7)$$

Now, we use (5.3), (5.4) and (5.7) to estimate respectively the local terms $\mathcal{T}_{l,1}$, $\mathcal{T}_{l,2}$, and the nonlocal terms \mathcal{T}_{nl} in (5.2). Finally, we get that there exists a constant $C_T > 0$ such that for all $t \in [0, T]$ and all $R(t) > 0$:

$$\frac{d}{dt} D[\pi](t)^2 \leq C_T(1 + R(t)) D[\pi](t)^2 + C_T e^{-\frac{\alpha_T}{2} R(t)}. \quad (5.8)$$

First, if we choose $R(t) = 1$ for all $t \in [0, T]$ in the inequality (5.8), Grönwall's lemma yields that there exists a constant $K_T > 0$ such that for all $t \in [0, T]$, $D[\pi](t)^2 < K_T$. Then, we define the function

$$u : t \mapsto \frac{D[\pi](t)^2}{e K_T},$$

so that for all $t \in [0, T]$ such that $u(t) > 0$, we have $1 \leq -\ln(u(t))$. Let $t \in [0, T]$ such that $u(t) > 0$. Then, we choose the quantity $R(t)$ in (5.8) as follows:

$$R(t) := -\frac{2}{\alpha_T} \ln(u(t)).$$

Hence, (5.8) becomes

$$\begin{aligned} u'(t) &\leq C_T \left(1 - \frac{2}{\alpha_T} \ln(u(t))\right) u(t) + \frac{C_T}{e K_T} u(t) \\ &\leq -C_T \left(1 + \frac{2}{\alpha_T}\right) u(t) \ln(u(t)) - \frac{C_T}{e K_T} u(t) \ln(u(t)) \\ &\leq -\tilde{C}_T u(t) \ln(u(t)), \end{aligned}$$

for some constant $\tilde{C}_T > 0$. With the convention $v \ln(v) = 0$ if $v = 0$, according to Osgood's lemma, we have for all $t \in [0, T]$:

$$\int_{u(0)}^{u(t)} \frac{-dy}{\tilde{C}_T y \ln(y)} \leq t.$$

This implies that

$$\ln(-\ln(u(t))) - \ln(-\ln(u(0))) \geq -\tilde{C}_T t.$$

Consequently, since $\ln(u(0)) < 0$, if we define the function

$$\beta : t \mapsto e^{-\tilde{C}_T t}, \tag{5.9}$$

we get that for all $t \in [0, T]$:

$$u(t) \leq u(0)^{\beta(t)},$$

173 and this remains true if $u(t) = 0$. Finally, we can conclude that for all $t \in [0, T]$

$$\begin{aligned} D[\pi](t) &\leq (e K_T)^{\frac{1-\beta(t)}{2}} D[\pi](0)^{\beta(t)} \\ &\leq \tilde{K}_T D[\pi](0)^{\beta(t)}, \end{aligned} \tag{5.10}$$

174 where $\tilde{K}_T > 0$ is a positive constant.

175 Step 2: Conclusion of the estimate of the Wasserstein distance

We note for all $j \in \{1, 2\}$ and all $t \in [0, T]$

$$f_j(t) := \mathcal{Z}_j(t) \# f_{0,j}.$$

Thus, for all $t \in [0, T]$ and for all $\pi \in \Lambda(f_{0,1}, f_{0,2})$,

$$(\mathcal{Z}_1(t) \times \mathcal{Z}_2(t)) \# \pi \in \Lambda(f_1(t), f_2(t)),$$

and therefore,

$$d_2(f_1(t), f_2(t)) \leq \left(\iint \|\mathcal{Z}_1(t, \mathbf{z}_1) - \mathcal{Z}_2(t, \mathbf{z}_2)\|^2 \pi(d\mathbf{z}_1 d\mathbf{z}_2) \right)^{\frac{1}{2}} = D[\pi](t).$$

Finally, we can conclude using the estimate (5.1):

$$\begin{aligned} d_2(f_1(t), f_2(t)) &\leq \tilde{K}_T D[\pi](0)^{\beta(t)} \\ &= \tilde{K}_T d_2(f_{0,1}, f_{0,2})^{\beta(t)}. \end{aligned}$$

Remark 5.1. *If we make the additional assumption that there exists $p > 1$ such that*

$$\int e^{\langle v, w \rangle^p} f_{0,2}(d\mathbf{z}) < \infty,$$

then instead of the estimate (5.7), we can conclude using a similar argument that there exists a constant $C_T > 0$ such that for all $t \in [0, T]$,

$$\frac{d}{dt} D[\pi](t)^2 \leq C_T (1 + R) D[\pi](t)^2 + C_T e^{-\frac{\alpha T}{2} R^p}.$$

Hence, Grönwall's lemma yields that for all $R > 0$ and all $t \in [0, T]$,

$$D[\pi](t)^2 \leq \left(D[\pi](0)^2 + \frac{e^{-\frac{\alpha T}{2} R^p}}{1 + R} \right) e^{C_T (1+R)t}.$$

Therefore, this implies that for all $t \in [0, T]$ and all $R > 0$,

$$d_2^2(f_1(t), f_2(t)) \leq D[\pi](t)^2 \leq \left(d_2^2(f_{0,1}, f_{0,2}) + \frac{e^{-\frac{\alpha T}{2} R^p}}{1 + R} \right) e^{C_T (1+R)t}. \quad (5.11)$$

Choosing for instance

$$R = -\frac{1}{2C_T T} \ln(d_2^2(f_{0,1}, f_{0,2})),$$

the estimate (5.11) is enough to prove the part (ii) of Theorem 1. Indeed, since $(f_{0,n})_{n \in \mathbb{N}}$ and f_0 satisfy assumption (2.8), we get

$$\lim_{n \rightarrow +\infty} \sup_{t \in [0, T]} d_2(f_n(t), f(t)) = 0.$$

176 5.2 Proof of the mean-field limit from (1.3) towards (1.5)

177 First, the existence and uniqueness of the solution to the FHN system is a direct consequence of the
178 Cauchy-Lipschitz theorem. Then, we have already proved the well-posedness of the kinetic equation (1.5)
179 in Section 4.

Now, let us conclude the proof using the stability result from the first part of Theorem 1. We notice that for all $n \in \mathbb{N}$, the empirical measure f_n is the measure solution of the kinetic equation (1.5) with initial condition $f_{0,n}$. Then, the part (i) of Theorem 1 yields that there exist two positive constants C_T and K_T independent of n such that for all $t \in [0, T]$ and for all $n \in \mathbb{N}$,

$$d_2(f_n(t), f(t)) \leq K_T d_2(f_{0,n}, f_0)^{\beta(t)},$$

where

$$\beta(t) := e^{-C_T t}.$$

Finally, using the assumption (2.8), we get that

$$\lim_{n \rightarrow +\infty} \sup_{t \in [0, T]} d_2(f_n(t), f(t)) = 0,$$

180 which concludes the proof of Theorem 1.

181 6 Application: stability of monokinetic solutions

One of the motivations to study the mean-field model is the analysis of the macroscopic quantities computed from a measure solution to (1.5), though the equation formally satisfied by the average membrane potential in the network is not closed. A way to study overcome this difficulty is to look for monokinetic solutions of (1.5), that is solutions f of the form

$$f(t, d\mathbf{x}, dv, dw) = \rho_0(\cdot) d\mathbf{x} \otimes \delta_{V(t, \cdot)}(dv) \otimes \delta_{W(t, \cdot)}(dw), \quad (6.1)$$

where ρ_0 is the average density of neurons, and (V, W) is the average couple membrane potential-adaptation variable in the network. Therefore, if f is a monokinetic solution of (1.5), then the couple (V, W) formally satisfies the nonlocal reaction-diffusion FHN system:

$$\begin{cases} \partial_t V(t, \mathbf{x}) - \int \Psi(\mathbf{x}, \mathbf{x}') (V(t, \mathbf{x}') - V(t, \mathbf{x})) \rho_0(\mathbf{x}') d\mathbf{x}' = N(V(t, \mathbf{x})) - W(t, \mathbf{x}), \\ \partial_t W(t, \mathbf{x}) = \tau (V(t, \mathbf{x}) + a - b W(t, \mathbf{x})). \end{cases} \quad (6.2)$$

In this subsection, we consider a connectivity kernel Ψ of the form:

$$\Psi(\mathbf{x}, \mathbf{y}) = \Phi(\mathbf{x} - \mathbf{y}),$$

182 where $\Phi : \mathbb{R}^d \rightarrow \mathbb{R}$ is symmetric, which means that the conductance between two neurons only depends on
 183 the distance between them. We want to use the stability result from Theorem 1 to prove that monokinetic
 184 solutions of the kinetic equation (1.5) are stable with respect to the Wasserstein distance d_2 . Under some
 185 additional assumptions, we can state the following well-posedness result.

Lemma 6.1. *Assume that Φ is a non-negative symmetric connectivity kernel in $L^1(\mathbb{R}^d)$, and N is the nonlinearity defined through:*

$$N : v \mapsto v - v^3. \quad (6.3)$$

Consider an initial data (ρ_0, V_0, W_0) satisfying

$$\rho_0 \in L^1 \cap L^\infty(\mathbb{R}^d), \quad \rho_0 \geq 0, \quad V_0, W_0 \in L^\infty(\mathbb{R}^d). \quad (6.4)$$

Then for any $T > 0$, there exists a unique couple (V, W) which is a classical solution to the nonlocal reaction-diffusion system (6.2) with initial data (V_0, W_0) , where

$$V, W \in \mathcal{C}^1([0, T], L^\infty(\mathbb{R}^d)).$$

186 The proof of Lemma 6.1 is based on a classical fixed point argument, and we refer to [16] for the details.
 187 Now, as a direct consequence of Theorem 1, we get the following result of stability of monokinetic solution
 188 of the equation (1.5).

Proposition 6.2 (Stability of monokinetic solutions). *Let $T > 0$. Assume that Φ is a non-negative symmetric connectivity kernel in $L^1(\mathbb{R}^d)$, and N is the nonlinearity defined through (6.3). Consider the initial data $f_0 \in \mathcal{P}_2(\mathbb{R}^{d+2})$, and (ρ_0, V_0, W_0) satisfying (6.4) and*

$$\int_{\mathbb{R}^d} \rho_0(\mathbf{x}) \, d\mathbf{x} = 1, \quad \int_{\mathbb{R}^d} \left(|\mathbf{x}|^2 + e^{\alpha_0 \langle V_0(\mathbf{x}), W_0(\mathbf{x}) \rangle} \right) \rho_0(\mathbf{x}) \, d\mathbf{x} < \infty, \quad (6.5)$$

for some positive constant $\alpha_0 > 0$. Let (V, W) be the solution of (6.2) with initial data (V_0, W_0) provided by Lemma 6.1, and let $f \in \mathcal{C}([0, T], \mathcal{P}_2(\mathbb{R}^{d+2}))$ be the measure solution of (1.5) with initial data f_0 . Then there exist two positive constants $C_T > 0$ and $K_T > 0$ such that for all $t \in [0, T]$,

$$d_2(f(t), \rho_0(\cdot) \, d\mathbf{x} \otimes \delta_{V(t, \cdot)}(dv) \otimes \delta_{W(t, \cdot)}(dw)) \leq K_T d_2(f_0, \rho_0 \, d\mathbf{x} \otimes \delta_{V_0}(dv) \otimes \delta_{W_0}(dw))^{\beta(t)}, \quad (6.6)$$

189 where $\beta(t) := e^{-C_T t}$.

Proof. According to the assumption (6.5), we have:

$$\int \left(|\mathbf{x}|^2 + e^{\alpha_0 \langle v, w \rangle} \right) \rho_0(\mathbf{x}) \, d\mathbf{x} \otimes \delta_{V_0(\mathbf{x})}(dv) \otimes \delta_{W_0(\mathbf{x})}(dw) < \infty,$$

190 so $\rho_0 \, d\mathbf{x} \otimes \delta_{V_0}(dv) \otimes \delta_{W_0}(dw) \in \mathcal{P}_2(\mathbb{R}^{d+2})$ and satisfies (2.5). Moreover, $\rho_0 \, d\mathbf{x} \otimes \delta_V(dv) \otimes \delta_W(dw) \in$
 191 $\mathcal{C}([0, T], \mathcal{P}_2(\mathbb{R}^{d+2}))$ is a measure solution of (1.5) since $(\rho_0, \rho_0 V, \rho_0 W)$ is a classical solution of (6.2).
 192 Thus, we can apply Theorem 1, which yields (6.6). ■

193 7 Numerical simulations

In this section, we approximate the solution f of the kinetic equation (1.5) for a one-dimensional network (*i.e.* $d = 1$), normalized to $[0, 1]$. There are few numerical methods specifically adapted to kinetic theory. In [1], the authors numerically approximate a mean-field model of neural network of FHN type using finite differences without considering any space dependence. On the contrary, we are particularly interested in the influence of space in the kinetic model (1.5). In order to approximate (1.5), we use a particle method. This kind of numerical scheme was first introduced by Harlow [24] for the numerical computation of specific problems in fluid dynamics, and precisely mathematically studied later [37]. Then, a large diversity of particle methods were introduced for simulations in fluid mechanics and plasma physics (see for instance [9, 20, 26] and references therein). Throughout this section, we fix $N(v)$ to be the following cubic nonlinearity:

$$N : v \mapsto v(1 - v)(v - 0.25). \quad (7.1)$$

194 7.1 Principle of the particle method

For $T > 0$, the standard particle method consists in approximating the solution f of the kinetic equation (1.5) on $[0, T]$ by a finite sum of Dirac masses

$$f_M(t, d\mathbf{x}, dv, dw) := \frac{1}{M} \sum_{i=1}^M \delta_{\mathbf{x}_i}(d\mathbf{x}) \otimes \delta_{v_i(t)}(dv) \otimes \delta_{w_i(t)}(dw),$$

where M is the number of distinct particles considered in the network, and for all $1 \leq i \leq M$, $(\mathcal{V}_i, \mathcal{W}_i)$ is the solution of the characteristic system (2.2) provided by Proposition 4.1 with initial condition $(\mathbf{x}_i, v_{0,i}, w_{0,i}) \in \mathbb{R}^3$ and initial measure $f_M(0, \cdot)$. In order to provide an approximation of the macroscopic quantities necessary to solve (2.2), we approach the Dirac masses by $\varphi_h := h^{-d} \varphi(\cdot/h)$, where $h > 0$ is a small fixed parameter, and

$$\varphi : z \mapsto \begin{cases} 0 & \text{if } |z| \geq 1, \\ 1 + z & \text{if } |z| \in [-1, 0], \\ 1 - z & \text{if } |z| \in [0, 1]. \end{cases}$$

Therefore, we can define the discrete densities on a mesh $(\mathbf{y}_i)_{1 \leq i \leq n_{\mathbf{y}}}$ of step size $h > 0$ of the considered interval $[0, 1]$, where $n_{\mathbf{y}} \in \mathbb{N}$ and $h > 0$ satisfy $h = 1/(n_{\mathbf{y}} - 1)$, with for all $1 \leq i \leq M$:

$$\begin{cases} \rho_{M,h}(\mathbf{y}_i) & := \frac{1}{M} \sum_{j=1}^M \varphi_h(\mathbf{y}_i - \mathbf{x}_j), \\ j_{M,h}(t, \mathbf{y}_i) & := \frac{1}{M} \sum_{j=1}^M \varphi_h(\mathbf{y}_i - \mathbf{x}_j) \mathcal{V}_j(t). \end{cases} \quad (7.2)$$

The whole point of the particle method is that $n_{\mathbf{y}} \ll M$. Now, for all $1 \leq i \leq M$, the couple $(\mathcal{V}_i, \mathcal{W}_i)$ is approached by a solution of an approximated characteristic system, noted (V_i, W_i) , in which we replace the nonlocal term in (2.2) with

$$\int \Psi(\mathbf{x}_i, \mathbf{x}') (\rho_{M,h}(\mathbf{x}') V_i(t) - j_{M,h}(t, \mathbf{x}')) d\mathbf{x}', \quad (7.3)$$

with the same initial data, and we approximate the integral terms in (7.3) with the rectangle method using the mesh $(\mathbf{y}_i)_{1 \leq i \leq n_{\mathbf{y}}}$. Finally, for the time discretization, we use a Runge-Kutta scheme of second order, with a time grid of step 0.01. In the following, we work in the case where the neurons in the network are homogeneously distributed in the interval $[0, 1]$. Consequently, we choose the parameters $(\mathbf{x}_i)_{1 \leq i \leq M}$ forming a regular mesh of the interval $[0, 1]$, and we fix for the rest of this section $M = 5001$ and $n_{\mathbf{y}} = 501$.

7.2 Numerical investigation of three different regimes

In this subsection, we display some numerical simulations of the kinetic model (1.5) to observe its dynamics for different sets of parameters. The dynamics of the microscopic system (1.3) when the number of neurons is large and of the kinetic model (1.5) are both rich, but not well known. Nevertheless, the model (1.1) for one isolated neuron has been extensively studied, and its asymptotic behaviors are perfectly predictable. Thus, we consider three different sets of parameters corresponding to three different regimes of the FHN model (1.1) with $I_{\text{ext}} = 0$.

- (i) Bistable regime: we remove the influence of the adaptation variable w , that is we consider $\tau = 0$. In this case, the equation (1.1) admits exactly two stable fixed points at $v = 0$ and $v = 1$ and one unstable at $v = 0.25$. Thus, the solution $v(t)$ of the equation (1.1) converges towards 0 as $t \rightarrow +\infty$ if $v(0) < 0.25$, or towards 1 if $v(0) > 0.25$.




Figure 7.1: **Bistable regime.** (a)-(b) Spatio-temporal evolution of the macroscopic function V_f computed from the solution f of the kinetic equation (1.5) with $\tau = 0$, and different values of the parameter ε , fixed at (a) 10^{-1} , (b) 10^{-3} . (c) Profile of the macroscopic function $V_f(t, \cdot)$ at different fixed times, computed with $\varepsilon = 10^{-3}$ and with $\tau = 0$.

- 211 (ii) Oscillatory regime: we choose $a = -0.25$ and $b = 3$, so that the system (1.1) admits a unique fixed
 212 point $(0.25, 0)$, which is unstable. We also consider $\tau = 0.02$. Consequently, in this setting, the
 213 solution of the system (1.1) converges towards a stable limit cycle if it is not initialized at the fixed
 214 point.
- 215 (iii) Excitable regime: we choose $a = 0$ and $b = 7$, so that the system (1.1) admits a unique fixed point
 216 $(0, 0)$, which is stable. All the solutions of (1.1) converge towards $(0, 0)$ as $t \rightarrow +\infty$. Moreover, we
 217 fix $\tau = 0.002$, so that (1.1) exhibits a slow/fast dynamics.

From now on, we define the connectivity kernel to be

$$\Psi(\mathbf{x}, \mathbf{y}) = G_\varepsilon(\mathbf{x} - \mathbf{y}) \quad \forall (\mathbf{x}, \mathbf{y}) \in \mathbb{R}^2 \quad \text{with} \quad G_\varepsilon(\mathbf{x}) := \frac{1}{\sqrt{2\pi\varepsilon}} \exp\left(-\frac{|\mathbf{x}|^2}{2\varepsilon}\right), \quad (7.4)$$

218 where $\varepsilon > 0$ is a rescaling parameter.

219 Let us discuss this choice of connectivity kernel in (7.4). First of all, since Ψ is non-negative, we consider
 220 a purely excitatory regime. Then, the parameter $\varepsilon > 0$ varies between 1 and very small values, in order
 221 to consider the regime of strong local interactions, which seems reasonable from a biological point of view
 222 since two neurons interact only through their contact point provided by their shared synapse.

223 We are interested in the dynamics of the average macroscopic couple (V_f, W_f) computed from the solution
 224 f of the kinetic model (1.5). In the previous section, we have proved the stability of monokinetic solutions
 225 (6.1) from well-prepared initial data (see Proposition 6.2). In the following, we present in the three regimes
 226 discussed above some numerical evidences that monokinetic solutions have a larger basin of attraction in
 227 the sense that the dynamics of (V_f, W_f) is found to be close to the dynamics of the solutions of the nonlocal
 228 reaction-diffusion system (6.2). To conclude this section, we will compare the dynamics between the FHN
 229 system (1.3), and the mean-field model (1.5) in the oscillatory regime for different values of n the number
 230 of neurons.




Figure 7.2: **Bistable regime.** Numerical approximation of the density function f solution of the kinetic equation (1.5) at fixed time (a) $t = 0$, (b) $t = 75$ and (c) $t = 150$, computed with the parameters $\varepsilon = 10^{-3}$ and $\tau = 0$.

231 Case (i) – Bistable regime.

We study the kinetic model (1.5) in the bistable regime with $\tau = 0$. As initial condition for our numerical scheme, we choose for all $1 \leq i \leq M$,

$$v_{0,i} = \frac{1}{2} \operatorname{erfc} \left(\sqrt{5000}(\mathbf{x}_i - 0.05) \right), \quad w_{0,i} = 0,$$

232 where erfc is the complementary error function.

233 In Figure 7.1, we show the spatio-temporal evolution of the macroscopic quantity V_f for different values
 234 of the parameter ε . First, in the case (a), $\varepsilon = 10^{-1}$ is large compared to the width of the considered
 235 interval $[0, 1]$. Thus, the space influence is almost homogenized, and the interactions between the particles
 236 of the neural network are expected to vanish after a few time. Consequently, for t large enough and for
 237 all fixed position $\mathbf{x} \in [0, 1]$, the macroscopic quantity $V_f(\cdot, \mathbf{x})$ is expected to behave as a solution of the
 238 one-neuron equation (1.1) in the same framework, that is to converge towards one of the two stable fixed
 239 points. Indeed, we can observe that it converges towards 0. Then, in the case (b), $\varepsilon = 10^{-3}$ is sufficiently
 240 small to observe another dynamic. Here, the behavior of the function V_f qualitatively looks like an invasion
 241 front, connecting the steady state 0 to 1, propagating at constant speed. Moreover, as shown in (c), after
 242 an initial transition phase, the shape of the front seems to be invariant and smooth.

243 We note that the qualitative behavior of the macroscopic function V_f when ε is small enough corresponds
 244 well to the dynamics of the nonlocal reaction-diffusion system (6.2) for which traveling front solutions are
 245 known to exist [3] when considered on the real line. Regarding the density function f , we show in Figure
 246 7.2 its temporal evolution for $\varepsilon = 10^{-3}$. We observe that the density f remains concentrated around the
 247 states $v = 1$ in an interval $[0, \mathbf{x}_0(t)]$ and then around $v = 0$ in the complementary interval $[\mathbf{x}_0(t), 1]$ for
 248 some $\mathbf{x}_0(t) \in (0, 1)$ propagating at a constant speed. This shows that f remains close to a monokinetic
 249 solution of the form $\rho_0 d\mathbf{x} \otimes \delta_{V_f(t, \cdot)}(dv)$, where the qualitative behavior of V_f is that of a traveling front, as
 250 previously detailed. This validates the fact monokinetic solutions seem to have a large basin of attraction.




Figure 7.3: **Oscillatory regime.** (a)-(b)-(c) Spatio-temporal evolution of the macroscopic function V_f computed from the solution f of the kinetic equation (1.5) with three different values of the parameter ε , fixed at (a) $\varepsilon = 10^{-1}$, (b) $\varepsilon = 10^{-3}$ and (c) $\varepsilon = 10^{-5}$. (d)-(e) Profile of the macroscopic function $V_f(t, \cdot)$ computed with $\varepsilon = 10^{-5}$ at time $t = 60$ and $t = 400$ respectively. (f) Trajectory in the phase space (v, w) of the couple (V_f, W_f) at fixed time $t = 400$ computed with $\varepsilon = 10^{-5}$. The other parameters are fixed at $a = -0.25$, $b = 3$, and $\tau = 0.02$.

251 Case (ii) – Oscillatory regime

In the oscillatory regime, we choose as initial data a perturbation of the steady state $(0.25, 0)$ concentrated around the position $\mathbf{x} = 0$:

$$v_{0,i} = 0.25 + 0.5 \exp(-5000 \mathbf{x}_i^2), \quad w_{0,i} = 0.$$

252

253 In Figure 7.3, we display the spatio-temporal evolution of the macroscopic function V_f computed from
 254 the solution f of the kinetic equation (1.5) with three different values of the variance of ε . In the first
 255 case (a), we fix the parameter $\varepsilon = 10^{-1}$. In that case, we expect that the space dependence of V_f to be
 256 suppressed and we indeed observe synchronized homogeneous oscillations. Then, we reduce the value of
 257 ε to localize the interactions and enforce the spatial dependence. For both $\varepsilon = 10^{-3}$ and $\varepsilon = 10^{-5}$, we




Figure 7.4: **Excitable regime.** (a) Spatio-temporal evolution of the macroscopic function V_f computed from the solution f of the kinetic equation (1.5) with $\varepsilon = 10^{-5}$. (b) Corresponding profile of the macroscopic function $V_f(t, \cdot)$ computed at different times. (c) Trajectory in the phase space (v, w) of the couple (V_f, W_f) at fixed time $t = 500$ computed with $\varepsilon = 10^{-5}$. The other parameters are fixed at $a = 0$, $b = 7$, and $\tau = 0.002$.

258 observe temporal oscillations whose phase is modulated spatially, similar to what is usually found for the
 259 local FHN reaction-diffusion system in the oscillatory regime [11]. More precisely, the dynamics is that of a
 260 modulated traveling wave propagating at constant speed from 0 towards the right, and leaving in the wake
 261 an oscillatory pattern with a constant frequency and amplitude. This is illustrated in Figure 7.3 panel (f)
 262 where the trajectory followed by the average couple (V_f, W_f) in the phase space at a given time converges
 263 towards a limit cycle.

264 Case (iii) – Excitable regime

We consider an initial condition that is a perturbation of the steady state $(0, 0)$ concentrated at the middle of the interval $[0, 1]$, that is for all $1 \leq i \leq M$:

$$v_{0,i} = \exp(-5000 |x_i - 0.5|^2), \quad w_{0,i} = 0.$$

265 We report in Figure 7.4, panel (a), the space-time representation of the macroscopic function V_f for
 266 $\varepsilon = 10^{-5}$. In that case, we see that the dynamics generates two counter-propagating traveling pulses.
 267 Once again, the behavior of the function V_f is qualitatively the same as the expected dynamics of the
 268 corresponding macroscopic model (6.2), where it is well known that the nonlocal reaction-diffusion FHN
 269 system supports traveling pulse solution [19]. We have drawn in Figure 7.4 panel (b) the profile of the
 270 v -component of the traveling pulse at different times and the corresponding trajectory in the phase plane
 271 in Figure 7.4 panel (c) where we recover that the profile of the traveling pulse is a homoclinic orbit to the
 272 stable fixed point $(0, 0)$. The study of traveling pulses in excitable media has received lots of interests in
 273 the past decades, especially for the local reaction-diffusion FHN system, and to our best knowledge, it is
 274 the first time that traveling pulses are reported for the FHN kinetic model.




Figure 7.5: Profile of the macroscopic function V_f computed from the solution f of the kinetic equation (1.5) with $\varepsilon = 10^{-4}$, and with the points $(\mathbf{x}_i, v_i)_{1 \leq i \leq n}$ from the solution of the FHN system (1.3), at fixed time $t = 225$. The other parameters are the same as in Figure 7.3.

7.3 Numerical comparison between the FHN system and the kinetic model

We consider a set of parameters corresponding to the oscillatory regime of the FHN model (1.1), that is the same parameters and initial condition as in paragraph above. We also fix $\varepsilon = 10^{-4}$, so that the average couple (V_f, W_f) computed from the solution to the kinetic equation (1.5) presents an oscillatory pattern modulated by a traveling wave propagating through space as previously.

In Figure 7.5, we display the profile of the macroscopic function V_f at fixed time $t = 225$ with respect to \mathbf{x} , together with the points $(\mathbf{x}_i, v_i)_{1 \leq i \leq n}$ standing for the solution of the FHN system (1.3) approximated with a Runge-Kutta scheme of second order, for different values of n . We choose $n = 50$ in the case (a), $n = 100$ in the case (b) and $n = 500$ in the case (c). In the case (a), n is large enough so that the oscillatory pattern emerges. Though, the frequency of these oscillations seems to be slightly higher than for the kinetic model (1.5). Then, we observe that for a relatively small number of neurons n in the network, the points $(\mathbf{x}_i, v_i)_{1 \leq i \leq n}$ match with the trajectory of $V(t, \cdot)$. Indeed, in the case (b), the oscillations of both models (1.3) and (1.5) are almost synchronized, and in the case (c), the points $(\mathbf{x}_i, v_i)_{1 \leq i \leq n}$ overlay the trajectory of V_f . It seems that the microscopic system (1.3) does not vary much any more for higher values of n , and quickly converges towards the solution of (1.5) as n tends to infinity.

On the one hand, the computations are obviously quicker than for the microscopic model (1.3) with a too large value of n . This is natural since in the numerical scheme of (1.5), we replace the sum over all the neurons in the network with a mean-field operator. On the other hand, the kinetic model (1.5) manages to accurately represent the behavior of a large neural network, and we still have access to more information than with a macroscopic model. Indeed, the kinetic model focuses on an approximation of the density of neurons rather than the average membrane potential. This shows the whole interest of the kinetic model (1.5) compared to the microscopic model (1.3) and to macroscopic models.

We mention that in the presence of noise as in [36], the authors showed some numerical simulations of the FHN system (1.3) with homogeneous interactions, where the finite number of neurons can cause the emergence of relaxation cycles near the transition from the excitable regime to the oscillatory regime, whereas the deterministic model still presents a unique stable fixed point.

301 8 Discussion

302 In this paper, we have proved the mean-field limit of the deterministic spatially-extended FHN model
 303 for neural networks towards a nonlocal kinetic equation as the number of neurons goes to infinity. Our
 304 approach is based on a stability estimate of solutions of the kinetic equation (1.5) with respect to their
 305 initial data. We have also proved the well-posedness of the kinetic equation in the space of probability
 306 measures with finite second moments, equipped with the Wasserstein distance of order 2. This mean-field
 307 limit provides a rigorous link between the microscopic scale of the neural network and a mesoscopic scale,
 308 which can then be used as an intermediary step for the derivation of macroscopic description of the neural
 309 network. Our microscopic model was obtained coupling the FHN model for a finite number of neurons,
 310 whose interactions with their neighbours are modulated only with their spatial position in the network
 311 with a connectivity kernel Ψ . Moreover, we have ignored the noise in the interactions, so that we only
 312 used deterministic tools. Finally, our numerical simulations showed that the kinetic model (1.5), with a
 313 sufficiently localized connectivity kernel, is robust enough to display some qualitative behaviors expected
 314 for macroscopic quantities in some specific frameworks, while retaining more information than macroscopic
 315 models.

316 Several extensions to this work seem natural. For example, taking inspiration from [8], a possibility could
 317 be to randomly choose the connectivity weights $\Psi_{i,j}$ in (1.3) with a probability law which depends only on
 318 the distance between the neurons i and j . A direct consequence of a mean-field limit result could be the
 319 propagation of chaos in the network as the number of neurons goes to infinity, that is the neurons become
 320 less and less correlated as their number gets large. Furthermore, another interesting extension comes from
 321 the recent work of Chiba and Medvedev [15] for the Kuramoto model, and to study the mean-field limit
 322 of the FHN model on various types of random graphs.

323 9 Acknowledgements

324 I thank Julien Chevallier for fruitful discussions during a poster session at the conference ICMNS 2018,
 325 and Grégory Faye and Francis Filbet for their help.

326 A Proof of Lemma 4.2

327 We start by proving that for all fixed $\mathbf{z} \in \mathbb{R}^{d+2}$, there exists a unique solution $(\mathcal{V}(\cdot, \mathbf{z}), \mathcal{W}(\cdot, \mathbf{z})) \in \mathcal{C}^1([0, T])^2$
 328 of (4.1) such that for all $t \in [0, T]$, $(\mathcal{V}(t, \cdot), \mathcal{W}(t, \cdot)) \in \mathcal{C}(\mathbb{R}^{d+2}, \mathbb{R})^2$. Then, we prove that \mathcal{V} and $\mathcal{W} \in$
 329 $\mathcal{C}([0, T], \mathcal{E})$, where \mathcal{E} is defined with (3.1)-(3.2).

330 Step 1: Existence and uniqueness of the solution of (4.1)

First of all, the function

$$\begin{pmatrix} v \\ w \end{pmatrix} \mapsto \begin{pmatrix} N(v) - w \\ A(v, w) \end{pmatrix}$$

is locally Lipschitz continuous in \mathbb{R}^2 . Moreover, since $\mathcal{U} \in \mathcal{C}([0, T], \mathcal{E})$, then for all $\mathbf{z} = (\mathbf{x}, v, w) \in \mathbb{R}^{d+2}$,

the function

$$t \mapsto \int \Psi(\mathbf{x}, \mathbf{x}') (\mathcal{U}(t, \mathbf{z}) - \mathcal{U}(t, \mathbf{z}')) f_0(d\mathbf{z}')$$

is continuous on $[0, T]$. Therefore, for all $\mathbf{z} \in \mathbb{R}^{d+2}$, the Cauchy-Lipschitz Theorem gives us the local existence and uniqueness of the solution $(\mathcal{V}(\cdot, \mathbf{z}), \mathcal{W}(\cdot, \mathbf{z}))$ of class \mathcal{C}^1 . Then, using an energy estimate, we want to prove that for all $\mathbf{z} \in \mathbb{R}^{d+2}$, the couple $(\mathcal{V}(\cdot, \mathbf{z}), \mathcal{W}(\cdot, \mathbf{z}))$ is well-defined on $[0, T]$. According to the definition of \mathcal{E} , we know that for all $(t, \mathbf{z}) \in [0, T] \times \mathbb{R}^{d+2}$:

$$|\mathcal{U}(t, \mathbf{z})| \leq \langle v, w \rangle \sup_{s \in [0, T]} \|\mathcal{U}(s)\|_{\mathcal{E}}. \quad (\text{A.1})$$

Let $\mathbf{z} \in \mathbb{R}^{d+2}$ and $t \in [0, T]$ small enough so that $(\mathcal{V}, \mathcal{W})$ exists in (t, \mathbf{z}) . We have the following energy estimate:

$$\frac{1}{2} \partial_t \left(|\mathcal{V}(t, \mathbf{z})|^2 + |\mathcal{W}(t, \mathbf{z})|^2 \right) = \mathcal{I}_1 + \mathcal{I}_2 + \mathcal{I}_3,$$

where

$$\begin{cases} \mathcal{I}_1 := \tau \mathcal{W}(t, \mathbf{z}) (\mathcal{V}(t, \mathbf{z}) + a - b \mathcal{W}(t, \mathbf{z})), \\ \mathcal{I}_2 := \mathcal{V}(t, \mathbf{z}) (N(\mathcal{V}(t, \mathbf{z})) - \mathcal{W}(t, \mathbf{z})), \\ \mathcal{I}_3 := -\mathcal{V}(t, \mathbf{z}) \int \Psi(\mathbf{x}, \mathbf{x}') (\mathcal{U}(t, \mathbf{z}) - \mathcal{U}(t, \mathbf{z}')) f_0(d\mathbf{z}'). \end{cases}$$

First of all, we treat the first term \mathcal{I}_1 using Young's inequality

$$\begin{aligned} \mathcal{I}_1 &\leq \frac{\tau}{2} |\mathcal{V}(t, \mathbf{z})|^2 + \tau |\mathcal{W}(t, \mathbf{z})|^2 + \frac{\tau a^2}{2} \\ &\leq \tau \left(|\mathcal{V}(t, \mathbf{z})|^2 + |\mathcal{W}(t, \mathbf{z})|^2 \right) + \frac{\tau a^2}{2}. \end{aligned}$$

Then, we estimate the second term \mathcal{I}_2 using Young's inequality and the assumption (2.4) satisfied by N :

$$\begin{aligned} \mathcal{I}_2 &\leq \kappa |\mathcal{V}(t, \mathbf{z})|^2 + \frac{1}{2} \left(|\mathcal{V}(t, \mathbf{z})|^2 + |\mathcal{W}(t, \mathbf{z})|^2 \right) \\ &\leq \left(\kappa + \frac{1}{2} \right) \left(|\mathcal{V}(t, \mathbf{z})|^2 + |\mathcal{W}(t, \mathbf{z})|^2 \right). \end{aligned}$$

Furthermore, we deal with the second term \mathcal{I}_3 using Young's inequality, the assumption that Ψ is bounded and the estimate (A.1):

$$\begin{aligned} \mathcal{I}_3 &\leq \frac{1}{2} |\mathcal{V}(t, \mathbf{z})|^2 + \frac{1}{2} \int |\Psi(\mathbf{x}, \mathbf{x}')|^2 \left(|\mathcal{U}(t, \mathbf{z})|^2 + |\mathcal{U}(t, \mathbf{z}')|^2 \right) f_0(d\mathbf{z}') \\ &\leq \frac{1}{2} |\mathcal{V}(t, \mathbf{z})|^2 + \frac{1}{2} \|\Psi\|_{\infty}^2 \left(|\mathcal{U}(t, \mathbf{z})|^2 + \int |\mathcal{U}(t, \mathbf{z}')|^2 f_0(d\mathbf{z}') \right) \\ &\leq \frac{1}{2} |\mathcal{V}(t, \mathbf{z})|^2 + \frac{1}{2} \|\Psi\|_{\infty}^2 \left(|\langle v, w \rangle|^2 + \int |\langle v', w' \rangle|^2 f_0(d\mathbf{z}') \right) \sup_{s \in [0, T]} \|\mathcal{U}(s)\|_{\mathcal{E}}^2 \\ &\leq \frac{1}{2} |\mathcal{V}(t, \mathbf{z})|^2 + \frac{1}{2} \|\Psi\|_{\infty}^2 \left(1 + \int |\langle v', w' \rangle|^2 f_0(d\mathbf{z}') \right) \sup_{s \in [0, T]} \|\mathcal{U}(s)\|_{\mathcal{E}}^2 |\langle v, w \rangle|^2. \end{aligned}$$

Hence, there exist two constants $C_1 > 0$ and $C_2 > 0$ such that for all $t \in [0, T]$ and $\mathbf{z} \in \mathbb{R}^{d+2}$,

$$\partial_t \left(|\mathcal{V}(t, \mathbf{z})|^2 + |\mathcal{W}(t, \mathbf{z})|^2 \right) \leq C_1 \left(|\mathcal{V}(t, \mathbf{z})|^2 + |\mathcal{W}(t, \mathbf{z})|^2 \right) + C_2 |\langle v, w \rangle|^2.$$

Therefore, Grönwall's inequality yields that for all $t \in [0, T]$,

$$\left(|\mathcal{V}(t, \mathbf{z})|^2 + |\mathcal{W}(t, \mathbf{z})|^2 \right) \leq (|v|^2 + C_2 t |\langle v, w \rangle|^2) e^{C_1 t}. \quad (\text{A.2})$$

This implies that $\mathcal{V}(\cdot, \mathbf{z})$ and $\mathcal{W}(\cdot, \mathbf{z})$ are defined on $[0, T]$, and thus $(\mathcal{V}(\cdot, \mathbf{z}), \mathcal{W}(\cdot, \mathbf{z})) \in \mathcal{C}^1([0, T])$. Moreover, if we divide (A.2) by $|\langle v, w \rangle|^2$ and if we take the supremum on \mathbb{R}^{d+2} , we conclude that for all $t \in [0, T]$,

$$\|\mathcal{V}(t)\|_{\mathcal{E}}^2 \leq (1 + C_2 t) e^{C_1 t}. \quad (\text{A.3})$$

Moreover, since Ψ is continuous with respect to its first variable uniformly relative to its second variable, and since $\mathcal{U} \in \mathcal{C}([0, T], \mathcal{E})$, we get that the function

$$(t, \mathbf{z}) = (t, \mathbf{x}, v, w) \mapsto \int \Psi(\mathbf{x}, \mathbf{x}') (\mathcal{U}(t, \mathbf{z}) - \mathcal{U}(t, \mathbf{z}')) f_0(d\mathbf{z}')$$

is continuous on $[0, T] \times \mathbb{R}^{d+2}$. Therefore, one can check that for all $t \in [0, T]$,

$$(\mathcal{V}(t, \cdot), \mathcal{W}(t, \cdot)) \in \mathcal{C}(\mathbb{R}^{d+2}, \mathbb{R}^2). \quad (\text{A.4})$$

Furthermore, (A.3) together with (A.4) yield that for all $t \in [0, T]$,

$$(\mathcal{V}(t, \cdot), \mathcal{W}(t, \cdot)) \in \mathcal{E}^2. \quad (\text{A.5})$$

331 Step 2: \mathcal{V} and \mathcal{W} are in $\mathcal{C}([0, T], \mathcal{E})$

It remains to prove that $(\mathcal{V}, \mathcal{W}) \in \mathcal{C}([0, T], \mathcal{E})^2$. Let us fix $t \in [0, T]$. Consider $\tilde{t} \in [0, T]$ and $\mathbf{z} = (\mathbf{x}, v, w) \in \mathbb{R}^{d+2}$. Thus, we have:

$$\frac{1}{2} \partial_t \left(|\mathcal{V}(t, \mathbf{z}) - \mathcal{V}(\tilde{t}, \mathbf{z})|^2 + |\mathcal{W}(t, \mathbf{z}) - \mathcal{W}(\tilde{t}, \mathbf{z})|^2 \right) = \mathcal{J}_1 + \mathcal{J}_2 + \mathcal{J}_3,$$

where

$$\left\{ \begin{array}{l} \mathcal{J}_1 := (\mathcal{W}(t, \mathbf{z}) - \mathcal{W}(\tilde{t}, \mathbf{z})) (A(\mathcal{V}(t, \mathbf{z}), \mathcal{W}(t, \mathbf{z})) - A(\mathcal{V}(\tilde{t}, \mathbf{z}), \mathcal{W}(\tilde{t}, \mathbf{z}))), \\ \mathcal{J}_2 := (\mathcal{V}(t, \mathbf{z}) - \mathcal{V}(\tilde{t}, \mathbf{z})) [(N(\mathcal{V}(t, \mathbf{z})) - N(\mathcal{V}(\tilde{t}, \mathbf{z}))) - (\mathcal{W}(t, \mathbf{z}) - \mathcal{W}(\tilde{t}, \mathbf{z}))], \\ \mathcal{J}_3 := -(\mathcal{V}(t, \mathbf{z}) - \mathcal{V}_{p+1}(\tilde{t}, \mathbf{z})) \\ \quad \times \int [\Psi(\mathbf{x}, \mathbf{x}') (\mathcal{U}(t, \mathbf{z}) - \mathcal{U}(t, \mathbf{z}')) + \Psi(\mathbf{x}, \mathbf{x}') (\mathcal{U}(\tilde{t}, \mathbf{z}) - \mathcal{U}(\tilde{t}, \mathbf{z}'))] f_0(d\mathbf{z}'). \end{array} \right.$$

We deal with the first term \mathcal{J}_1 using Young's inequality

$$\mathcal{J}_1 \leq \frac{\tau}{2} \left(|\mathcal{V}(t, \mathbf{z}) - \mathcal{V}(\tilde{t}, \mathbf{z})|^2 + |\mathcal{W}(t, \mathbf{z}) - \mathcal{W}(\tilde{t}, \mathbf{z})|^2 \right).$$

Then, we can estimate the second term \mathcal{J}_2 using Young's inequality and the assumption (2.4) satisfied by N :

$$\mathcal{J}_2 \leq \left(\tilde{\kappa} + \frac{1}{2} \right) |\mathcal{V}(t, \mathbf{z}) - \mathcal{V}(\tilde{t}, \mathbf{z})|^2 + \frac{1}{2} |\mathcal{W}(t, \mathbf{z}) - \mathcal{W}(\tilde{t}, \mathbf{z})|^2.$$

Then, to treat the last term \mathcal{J}_3 , we use Young's inequality and the assumption that Ψ is bounded:

$$\begin{aligned} \mathcal{J}_3 &= -(\mathcal{V}(t, \mathbf{z}) - \mathcal{V}(\tilde{t}, \mathbf{z})) \\ &\quad \times \left((\mathcal{V}(t, \mathbf{z}) - \mathcal{V}(\tilde{t}, \mathbf{z})) \int \Psi(\mathbf{x}, \mathbf{x}') f_0(d\mathbf{z}') + \int \Psi(\mathbf{x}, \mathbf{x}') (\mathcal{U}(t, \mathbf{z}') - \mathcal{U}(\tilde{t}, \mathbf{z}')) f_0(d\mathbf{z}') \right) \\ &\leq \frac{1}{2} |\mathcal{V}(t, \mathbf{z}) - \mathcal{V}(\tilde{t}, \mathbf{z})|^2 \\ &\quad + \frac{1}{2} \|\Psi\|_\infty^2 |\mathcal{U}(t, \mathbf{z}) - \mathcal{U}(\tilde{t}, \mathbf{z})|^2 + \frac{1}{2} \|\Psi\|_\infty^2 \|\mathcal{U}(t) - \mathcal{U}(\tilde{t})\|_{\mathcal{E}}^2 \int |\langle v', w' \rangle|^2 f_0(d\mathbf{z}'). \end{aligned}$$

Then, dividing \mathcal{J}_1 , \mathcal{J}_2 and \mathcal{J}_3 by $|\langle v, w \rangle|^2$ and taking the supremum on \mathbb{R}^{d+2} , we get that there exist two constants C_1 and C_2 such that:

$$\begin{aligned} \frac{d}{dt} \left(\|\mathcal{V}(t) - \mathcal{V}(\tilde{t})\|_{\mathcal{E}}^2 + \|\mathcal{W}(t) - \mathcal{W}(\tilde{t})\|_{\mathcal{E}}^2 \right) \\ \leq C_1 \left(\|\mathcal{V}(t) - \mathcal{V}(\tilde{t})\|_{\mathcal{E}}^2 + \|\mathcal{W}(t) - \mathcal{W}(\tilde{t})\|_{\mathcal{E}}^2 \right) + C_2 \|\mathcal{U}(t) - \mathcal{U}(\tilde{t})\|_{\mathcal{E}}^2. \end{aligned}$$

Using Grönwall's lemma the assumption $\mathcal{U} \in \mathcal{C}([0, T], \mathcal{E})$, we conclude that

$$\lim_{\tilde{t} \rightarrow t} \left(\|\mathcal{V}(t) - \mathcal{V}(\tilde{t})\|_{\mathcal{E}}^2 + \|\mathcal{W}(t) - \mathcal{W}(\tilde{t})\|_{\mathcal{E}}^2 \right) = 0,$$

and consequently,

$$(\mathcal{V}, \mathcal{W}) \in \mathcal{C}([0, T], \mathcal{E})^2.$$

References

- 332
- 333 [1] J. Baladron, D. Fasoli and O. Faugeras. *Three applications of GPU computing in neuroscience.*
334 Computing in science and engineering, 14 (2012), 40-47.
- 335 [2] J. Baladron, D. Fasoli, O. Faugeras and J. Touboul. *Mean-field description and propagation of chaos*
336 *in networks of Hodgkin-Huxley and FitzHugh-Nagumo neurons.* The Journal of Mathematical Neuro-
337 science, 2 (2012), 10.
- 338 [3] P.W. Bates, P.C. Fife, X. Ren and X. Wang *traveling Waves in a Convolution model for phase*
339 *transitions* Archive for rational Mechanics and Analysis, 138 (1997), 105–136.
- 340 [4] F. Bolley, J.A. Cañizo and J.A. Carrillo. *Stochastic mean-field limit: non-lipschitz forces and swarm-*
341 *ing.* Mathematical models and methods in applied sciences, 21, (2011), 2179-2210.
- 342 [5] M. Bossy, O. Faugeras and D. Talay. *Clarification and complement to "mean-field description and*
343 *propagation of chaos in networks of Hodgkin-Huxley and FitzHugh-Nagumo neurons".* The Journal of
344 Mathematical Neuroscience, 5 (2015), 1.
- 345 [6] M. Bossy, J Fontbona and H. Olivero. *Synchronization of stochastic mean field networks of Hodgkin-*
346 *Huxley neurons with noisy channels.* submitted (2018).
- 347 [7] P.C. Bressloff. *Spatially periodic modulation of cortical patterns by long-range horizontal connections.*
348 Physica D: Nonlinear Phenomena, 185 (2003), 131–157.

- 349 [8] T. Cabana and J. Touboul. *Large deviations for randomly connected neural networks: I. Spatially*
350 *extended systems*. Advances in Applied Probability, 50 (2018), 944-982.
- 351 [9] M. Campos Pinto, E. Sonnendrücker, A. Friedman, D.P. Grote, S.M. Lund. *Noiseless Vlasov-Poisson*
352 *simulations with linearly transformed particles*. Journal of computational Physics, 275 (2014), 236-256.
- 353 [10] J. A. Cañizo, J. A. Carrillo and J. Rosado. *A well-posedness theory in measures for some kinetic*
354 *models of collective motion*. Math. Mod. Meth. Appl. Sci., 21 (2011), 515-539.
- 355 [11] P. Carter and A. Scheel. *Wave train selection by invasion fronts in the FitzHugh-Nagumo equation*.
356 Nonlinearity, 31 (2018), 5536-5572.
- 357 [12] J. Chevallier, M.J. Caceres, M. Doumic and P. Reynaud-Bouret. *Microscopic approach of a time*
358 *elapsed neural model*. Mathematical Models and Methods in Applied Sciences, 25 (2015), 2669-2719.
- 359 [13] J. Chevallier. *Mean-field limit of generalized Hawkes processes*. Stochastic Processes and their Appli-
360 *cations*, 127 (2017), 3870-3912.
- 361 [14] J. Chevallier, A. Duarte, E. Löcherbach and G. Ost. *Mean-field limits for nonlinear spatially extended*
362 *hawkes processes with exponential memory kernels*. Stochastic Processes and their Applications, 129
363 (2018), 1-27.
- 364 [15] H. Chiba and G.S. Medvedev. *The mean field analysis for the Kuramoto model on graphs I. The*
365 *mean field equation and transition point formulas*. Discrete & Continuous Dynamical Systems - A, 39
366 (2019), 131-155.
- 367 [16] J. Crevat, G. Faye and F. Filbet. *Rigorous derivation of the nonlocal reaction-diffusion FitzHugh-*
368 *Nagumo system*. SIAM J. Math. Anal., to appear, (2019).
- 369 [17] R. Dobrushin. *Vlasov equations*. Funct. Anal. Appl., 13 (1979), 115-123.
- 370 [18] O. Faugeras, J. Touboul and B. Cessac. *A constructive mean-field analysis of multi population neural*
371 *networks with random synaptic weight and stochastic inputs*. Frontiers in computational neuroscience,
372 3 (2009), 1.
- 373 [19] G. Faye and A. Scheel. *Existence of pulses in excitable media with nonlocal coupling*. Advances in
374 Mathematics, 270 (2015), 400-456.
- 375 [20] F. Filbet, L.M. Rodrigues. *Asymptotically stable particle-in-cell methods for the Vlasov-Poisson system*
376 *with a strong external magnetic field*. SIAM J. Numer. Analysis, 542 (2016), 1120-1146.
- 377 [21] R. FitzHugh. *Impulses and physiological states in theoretical models of nerve membrane*. Biophysical
378 journal, 1 (1961), 445-466.
- 379 [22] F. Golse *On the dynamics of large particle systems in the mean field limit, in "Macroscopic and large*
380 *scale phenomena: coarse graining, mean field limits and ergodicity"*, Springer, 2016, 1-144.
- 381 [23] S.-Y. Ha and J.-G. Liu. *A simple proof of the Cucker-Smale flocking Dynamics and mean-field limit*.
382 Commun. Math. Sci., 7 (2009), 297-325.

- 383 [24] F.H. Harlow. *The particle-in-cell computing method for fluid dynamics*. Method in Computational
384 Physics, 3 (1964), 319–343.
- 385 [25] M. Hauray and P.-E. Jabin. *N-particles Approximation of the Vlasov Equations with Singular Poten-*
386 *tial*. Arch. Ration. Mech. Anal., 183 (2007), 489–524.
- 387 [26] D.W. Hewett. *Fragmentation, merging, and internal dynamics for PIC simulation with finite size*
388 *particles*. Journal of Computational Physics, 189 (2003), 390–426.
- 389 [27] A. L. Hodgkin and A. F. Huxley. *A quantitative description of membrane current and its application*
390 *to conduction and excitation in nerve*. The Journal of Physiology, 117 (1952), 500–544.
- 391 [28] E. Luçon and W. Stannat. *Mean-field limit for disordered diffusions with singular interactions*. The
392 *annals of applied probability*, 24 (2014), 1946–1993.
- 393 [29] S. Mischler, C. Quiñinao and J. Touboul. *On a kinetic FitzHugh-Nagumo model of neuronal network*.
394 *Comm. Math. Phys.*, 342 (2015), 1001–1042.
- 395 [30] J. Nagumo, S. Arimoto and S. Yoshizawa. *An active pulse transmission line simulating nerve axon*.
396 *Proceedings of the IRE*, 50 (1962), 2061–2070.
- 397 [31] I. Omelchenko, B. Riemenschneider, P. Hövel, Y. Maistrenko and F. Schöll. *Transition from spatial*
398 *coherence to incoherence in coupled chaotic systems*. Physical Review E, 85 (2012), 026212.
- 399 [32] David Parker. *Variable properties in a single class of excitatory spinal synapse*. The journal of
400 *Neuroscience*, 23 (2003), 3154–3163.
- 401 [33] K. Pakdaman, B. Perthame and D. Salort. *Dynamics of a structured neuron population*. Nonlinearity,
402 23 (2010), 55–75.
- 403 [34] K. Pakdaman, B. Perthame and D. Salort. *Relaxation and self-sustained oscillations in the time*
404 *elapsed neuron network model*. SIAM J. Appl. Math, 73 (2013), 1260–1279.
- 405 [35] K. Pakdaman, B. Perthame and D. Salort. *Adaptation and Fatigue Model for Neuron Networks and*
406 *Large Time Asymptotics in a Nonlinear Fragmentation Equation*. Journal of Mathematical Neuro-
407 *science*, 4 (2014), 14.
- 408 [36] C. Quiñinao, and J. Touboul. *Clamping and synchronization in the strongly coupled FitzHugh-Nagumo*
409 *model*. submitted (2018).
- 410 [37] P.A.Raviart. *An analysis of particle methods, in "Numerical methods in fluid dynamics"*. Springer,
411 243–324, (1983).
- 412 [38] J. Touboul, G. Hermann and O. Faugeras. *Noise-induced behaviors in neural mean field dynamics.1*
413 (2012), 49–81.
- 414 [39] C. Villani. *"Topics in Optimal Transportation"*, American Mathematical Society, 1–73, 2003.
- 415 [40] G. Wainrib and J. Touboul. *Topological and dynamical complexity of random neural networks*. Phys.
416 *Rev. Lett.*, 110 (2013), 118101.