

Micromechanics of root development in soil

Lx Dupuy, M Mimault, D Patko, Vincent Ladmiral, B. Ameduri, M P Macdonald, M Ptashnyk

▶ To cite this version:

Lx Dupuy, M Mimault, D Patko, Vincent Ladmiral, B. Ameduri, et al.. Micromechanics of root development in soil. Current Opinion in Genetics and Development, 2018, 51, pp.18 - 25. 10.1016/j.gde.2018.03.007. hal-01898743

HAL Id: hal-01898743

https://hal.science/hal-01898743

Submitted on 25 Apr 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

- 1 Micromechanics of root development in soil
- 2 Dupuy LX^{1*}, Mimault M¹, Patko D¹, Ladmiral V², Ameduri B², MacDonald M P³, Ptashnyk M⁴

4

- ¹The James Hutton Institute, Invergowrie, DD25DA, Dundee, United Kingdom
- ² ICGM, University of Montpellier, CNRS, ENSCM, 240 avenue du Professeur Emile Jeanbrau, 34296
- 6 Cedex 5 Montpellier, France
- 7 ³School of Science and Engineering, University of Dundee, DD1 4HN, Dundee, United Kingdom
- 8 ⁴School of Mathematical and Computer Sciences, Heriot-Watt University, EH14 4AS, United Kingdom
- 9 * for correspondence: lionel.dupuy@hutton.ac.uk
- 10 Keywords: plant roots, particle, granular media, biomechanics

11

12

13

Abstract

- 14 Our understanding of how root develop in soil may be at the eve of significant transformations. The
- 15 formidable expansion of imaging technologies enables live observations of the rhizosphere micro-pore
- architecture at unprecedented resolution. Granular matter physics provides ways to understand the
- 17 microscopic fluctuations of forces in soils, and the increasing knowledge of plant mechanobiology may
- shed new lights on how roots perceive soil heterogeneity. This opinion paper exposes how recent
- 19 scientific achievements may contribute to design a new theory for root growth in heterogeneous
- 20 environments.

Main text

- 22 Current knowledge of the biomechanics of plant root growth in soil is largely based on the extensive
- 23 work of plant biophysicists from the second half of the 20th century [1-3] . The view was that both

roots and soil must be considered as continua so that the description of root soil interactions can be achieved with continuous mathematical functions of macroscopic variables such as Young's modulus of root tissue, soil penetration stress, and pore water pressure [4]. Classical concepts from mechanics and physiology then provide a suitable framework to understand factors controlling tissue growth in its natural environment. The energy required to deform the root and surrounding soil, which originates from the photosynthetic chemical energy accumulated within the tissues, is converted into turgor pressure and mechanical energy [5]. Turgor pressure then overcome the resistance from cell wall to stretching, the resistance to movement of water across membranes, and the resistance to the displacement of the soil around the root [6]. This classical view of root-soil biomechanics has been central to identify the biophysical factors limiting growth in soil, but it is now challenged to predict morphologies and developmental patterns observed in natural conditions (Figure 1). If roots were to experience homogeneous mechanical stress from the soil, one would expect turgor pressure and Lockhart equation [1] to predict accurately growth arrest in soil. This is not the case and large discrepancies remain between measured turgor pressure (in the order of 1MPa [7]) and the levels of mechanical stresses at which growth is arrested (>5MPa [8]). Classical mechanics of continua is ill-equipped to explain the links between soil heterogeneity and stochasticity of plant development. The root tissue itself is heterogeneous and cell types have different roles in facilitating growth and penetration. Anchoring the base of the root for example, is necessary for cell elongation to produce apical movement and deformation of the soil [9]. The root cap and its associated border cells have also a fundamental role in reducing friction from the bulk soil. It was shown recently that wheat genotypes with sharper root tips are more efficient at soil penetration [10]. To establish a biomechanical framework that accounts for the complexity of root interactions with the granular medium, one must capture the microscopic nature of particle forces and the collective action they have on root tissues (Figure 1A). Kolb, et al. [11] proposed to categorise the nature of root mechanical responses to soil based on the scale of the soil heterogeneities. When the medium is

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

composed of small particles, individual variations in the force required to move them are not perceived by the root. The behaviour of roots and soil can be homogenised, and classical continuum mechanics usually applies (Box 1A) [12]. Soils also contain objects that are too large and or too rigid for a root to deform and displace, for example when roots grow in contact with stones, in cracks or pores [13,14]. Growth forces cannot displace the obstacle and the root usually combines tropic responses and mechanical buckling to avoid the obstacle (Box 1B) [15]. The behaviour of roots growing in soils with particles of intermediate sizes is more challenging to understand. A root can displace individual particles from the soil, but the forces exerted by each of the particles can also influence the course of root development (Box 1C). Although such growth environments are common for fine roots or due to the presence of aggregate and sand particles, growth patterns in such conditions are not well understood. How frequently does a root deflect from their growth trajectory? What are the magnitude of deflections? How does the distribution of particle forces modify the growth trajectory? Understanding the forces acting on a root during the elongation requires detailed knowledge of the physics of granular media. Granular media are assemblages of particles held by frictional and repulsive forces from adjacent particles. The forces holding particles together form chain-like networks that propagate at the contact points between neighbouring particles [16]. Because particles are disordered or have various sizes and shapes, large variations in magnitude and direction of particle forces arise [16,17]. Early theoretical work based on dry and static monodisperse particles showed that distribution of contact forces vary greatly and the overall force distribution follows an exponential decline [18,19]. Particles dynamics is better understood too. Contact forces in granular media propagate through complex waves [20] with appearance of macroscopic phenomenon such as clogging and arching, where particles spontaneously organise as vaults [21]. Solid, liquid and even gaseous phases may be observed in granular media depending on the external forces applied upon them [22]. Indeed, powerful techniques and hardware are available to examine theories in conditions that are nearly identical to experiments. 3D templates of the pore geometry together with description of the root and anatomical details can be obtained [23,24], and there are efficient computational

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

techniques that exploit the power of Graphical Processing Unit to simulate roots and soil at the particle and cell resolution. Discrete Element Modelling (DEM) for example uses Newton's second law to describe the motion of millions of interacting particles [25,26]. The models reproduce closely experimental observations, even in the case of biologically complex systems with detailed quantification of the force distribution surrounding growing roots [27,28]. Despite recent experimental and theoretical breakthroughs, granular matter physics has not transformed our understanding of the mechanics of root growth. Many current limitations are due to our lack of understanding of how roots respond to complex mechanical signals, and particularly how competition between multiple mechanical stimuli affects root responses. Cellular mechanisms involved in the response to physical obstacles have not been fully characterised, but a growing number of studies are now revealing the signalling and regulatory mechanisms involved in plant responses to mechanical force. Research in animal sciences have identified a multitude of proteins which binding domains are modified by mechanical forces [29] and their discovery in plants may follow. Large families of mechanosensitive ion channels have been identify in plants [30], with for example MCA calcium mechanosensitive channels being linked to growth response to hard gel layers [31]. Adaptation to mechanical forces are also well characterised, including the changes in cell division patterns, growth direction, cell differentiation and gene expression [32] A main difficulty, however, is to understand the nature of the mechanical signals perceived from the soil particles surrounding plant roots. It is central to develop capabilities to study not only the forces and displacement produced in the root soil system, but also the biological responses due to mechanical interactions with soil particles. Unfortunately, experimenting with natural soils is challenging because of its opacity. Rhizotron systems have been an extremely powerful tool to study root growth [33,34], glass interfaces introduce strong border effects and observations of biomechanical processes are often biased. X-ray imaging allows visualisation of interactions between roots and soil particles in situ in high resolution [35]. The technique allows time-lapse imaging for several weeks of growth. Improved images can be obtained with the application of contrasting agents.

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

For example, iodine perfused into plant leaves revealed the vascular structures of the roots and rhizobial nodules [36]. Root hairs can be resolved using synchrotron sources with resolution of up to 5µm and at temporal resolution sufficient for tracking particle movement due to root growth [37]. However, X-ray is an ionising radiation that affects biological processes especially meristematic regions where high cell division rates occurs [38], and despite the increase in resolutions, details of the inner cellular processes and biochemical activity have remained invisible [39]. Optics and microscopy in the visible range have thus remained the preferred approach to make observation of the biology and mechanics of the root. Confocal Laser Scanning Microscopes (CLSM) have provided the first live images of root-particle interaction in high resolution with details available on contact with particle surface, anatomical features at cell resolution and gene expression [40-42]. FRET imaging now allows tension sensors to record molecular forces at the piconewton scale [43]. However, CLSM has proved limited for long observations due to photo toxicity and photo bleaching. Because of the confined environment of the microscope, it has also remained limited to small plant samples. The field is now turning to different types of microscopes. Light Sheet Microscopy (LSM), in particular, has drastically reduced the light doses to the samples [44]. Illumination of the sample is planar and achieved orthogonal to the detection so that 2D images are generated instantaneously often using the new generation of scientific-CMOS cameras. By taking a whole 2D section in one "shot", volume scanning is accelerated, enabling small and fast developmental events to be tracked during development. The technique has considerably advanced our ability to observe living organisms both live and in situ with, for example, the ability to track cell growth, movement and divisions of entire embryos [45] or capturing the beating of a living heart [46]. Because axial resolution in light sheet systems is not dependent upon high numerical aperture imaging objectives, they allow larger fields of view and can easily accommodate microcosms and instruments for maintaining healthy growth conditions [47]. Details of the morphology and anatomy of tissues can be obtained without the use of markers [48,49] and recently dynamic light scattering (biospeckle) has been used to

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

127 enhance image contrast [50]. Light sheet imaging has also been used in granular matter physics for a 128 long time, although its application to root and soil is just emerging [50-52]. 129 Optics and microscopy also provides many ways to control and measure mechanical forces. Laser 130 ablation for example, has long been used to understand the distribution of forces within a tissue [53], 131 whilst optical trapping has been used to apply small localised forces [54]. Photoelastic materials have 132 been central to establishing the nature of the chains of forces and how they propagate within a 133 granular medium [55]. Kolb, et al. [56] used photoelasticity to characterise the forces created by root 134 growth within a pore, and Wendell, et al. [57] have successfully created a granular medium using a 135 photo elastic media where maximum growth forces and avoidance mechanisms could be observed 136 (Figure 1B). New cantilever-based optical sensors [58,59] have also been developed to measure 137 simultaneously growth forces generated by a root and three-dimensional strain rate in responses to 138 changes in external forces applied to the root. Bizet, et al. [59] for example, obtained stereoscopic data to decompose root response to axial mechanical forces into different phases (Figure 1C). 139 140 Hydrogels can also be combined with fluorescent dyes and light sheet imaging to reconstruct 141 interparticle forces within the granular medium [60]. 142 Techniques for mimicking soil physical conditions under a microscope are also emerging rapidly. 143 Transparent artificial media based on fluoropolymers that can mimic soil properties have been 144 developed [42]. The media reproduces the physical and chemical properties of soil through control of 145 the distribution of sizes and surface chemistry of the particles (Figure 1D). Because the particles are 146 made of fluoropolymers that have refractive index close to water, only small adjustment of refractive 147 indices, usually by adding a colloid to the nutrient solution, allows light to travel without refraction 148 through the substrate. Microfluidics techniques have also progressed significantly and are becoming 149 suitable to live and high resolution microscopy of roots and microbes [61,62]. Microfluidics allows 150 precise and repeatable control of liquids and this could be used, for example, to control water tension 151 and particle cohesion in soil during live experiments. The range of materials and fabrication techniques 152 has been considerably expanded with the use of 3D printing [63], photo lithography [64], etching technics [65] and the use of optically controlled fluidics [66,67]. It has been possible, for example, to produce chambers with physical heterogeneity, physical barriers and chemical gradients, with direct applications to root and soil studies [62,64,68].

The scientific community is better equipped than ever to make observations on the micromechanics of root development in soil. Experimental systems provide soil-like growth conditions and allow for observations, measurements and data generation with precision, accuracy and resolution. How then to transform the amount of information available to us into scientific breakthrough? The complexity of the root-particle interactions is a major challenge. At each growth step, a root is in contact with a new arrangement of particles that apply forces of varying magnitudes and orientations. Because there are countless numbers of possible arrangements, the forces applied on roots cannot be experimentally controlled. Measurements of granular forces *in situ* is required (Figure 2.1), and granular media physicist have achieved such measurements. There are now great opportunities to combine current knowledge of soil micromechanics with mechanobiology and propose a mechanistic framework that account for sensing and response to micro-scale heterogeneity (Figure 2.2). New theories must be developed to embrace stochasticity and explain responses to multiple mechanical stimuli (Figure 2.3-4). Major challenges remain, but a recent look at the literature indicates our thinking is evolving in the right way.

170 Figures





Figure 1: Growing roots interact mechanically with soil particles during growth. These interactions influence the morphology of the root, and the dynamics of development of the root system. A) Irregular growth of cortex cells is observed in hard or compacted soil [left, 69]. Resistance from the soil particles causes root diameter to increase and the root tip to buckle and bend towards the path of least resistance (middle, lentils roots grown at 2MPa confining pressure). At the scale of the root system, interactions causes growth trajectories to be stochastic as observed here on *Anthyllis vulneraria* grown on landslide soils (image courtesy Loïc Pagès). Technological developments now allow precise characterisation of mechanical interactions between a root and the growth substrate. These include for example, B) photoelastic discs for measurement of growth forces in soil pores [56] (images courtesy Evelyne Kolb), C) root growing on a cantilever sensor for measuring growth forces [59], D) transparent soil substrates that provide the physical structure of soil with the ability to carry out 3D live imaging [50], E) Dual flow microfluidic systems with microscale both physical and nutrient heterogeneity [68] and F) discrete element modelling for testing root responses to interactions with granular media [28] (image courtesy Mahmoud Fakih).



Box 1: Root primary growth is a local process where elongation of tissues is taking place at the root tip. Soil heterogeneity influences strongly how the tissue elongates and deforms (top), and local interactions taking place at the tip can have drastic effects on the morphology and development of the whole root system, and the resources available to the plant (bottom). Mathematical modelling provides a useful framework to explain how heterogeneity can affect the morphology of the root system.

(A) When roots grow in soil particles which representative volume is small compared to the diameter of the roots, the action of the particles can be averaged (top). In such conditions, it is unlikely for a plant to perceive the fluctuations of forces from individual particles. If the mechanical resistance of

the soil is not limiting, root trajectories follow smooth streamlines (bottom). Mathematically, this phenomenon has been described as the convection of root tips (density ρ) [70]. The growth velocity E (cm.d¹) and the rate of change in root angle due to gravitropism g (d¹) define the growth of the root system:

$$\partial_t \rho + \nabla \cdot \mathbf{F} + \partial_\theta g \rho = 0,$$

with $F = \rho E(\cos(\theta) + \sin(\theta))$ is the spatial flux of root tips and $g\rho$ is the angular flux of roots. In this case, growth and resource acquisition is optimal.

(B) When soil elements cannot be displaced, in the case of stones and rock for example, the root adopts avoidance behaviours. Optimal growth is not affected and remains similar to (A), until the obstacle is reached. Heterogeneities in this case define the boundaries within which convective growth is taking place. Using the same mathematical framework, presence of such boundaries can be modelled through boundary conditions, for example

$$\partial_n \rho = 0.$$

Large scale soil heterogeneities can be problematic because they may restrain access to pools of resources, e.g. deep water, even though the root growth in most parts of the soil domain is unaffected. They may also forms paths of least growth resistance, for example in the case of pores and cracks.

(C) Intermediate cases are more problematic to analyse. Roots are in contact with particles which apply forces of varying magnitudes and orientation. Although the root may overcome these forces, a single particle may be able to deflect the growth trajectory. Since particles have inhomogeneous distribution, root deflection occurs is stochastic. Mathematically, the phenomenon can be described by a convection, where the growth velocity $e < E(cm.d^{-1})$ and the rate of change in root angle due to gravitropism $g(d^{-1})$ and random fluctuations define the dynamics:

220
$$\partial_t \rho + \nabla \cdot \mathbf{F} + \partial_{\theta} (g\rho + D\partial_{\theta} \rho) = 0,$$

 $g\rho + D\partial_{\theta}\rho$ the angular flux of roots. The parameter D is the angular diffusion coefficient. Because D relates to the probability of roots to be deflected by a particle, and the magnitude of such deflection, there is a direct link between micro-mechanics of root particle interactions and the morphology of the root system. Diffusive growth makes root trajectories irregular, and limits the expansion of the root system, even when the elongation rate is not affected. Mathematical analysis of equation 3 reveals the conditions for which transitions from convective growth to diffusive growth occur, i.e. for Peclet number $P_e = \frac{g}{D} <<1$.




Figure 2: Dissecting the complexity of root particles mechanical interactions requires an elaborate research strategy. (1) First step is to better understand the nature of the forces applied to a root. This can be achieved, using photo elastic beads, imaging, or developing artificial roots equipped with sensors [71], but also by revisiting older techniques, for example by analysing micro penetrometer test and exploit force fluctuations [72]. (2) In the second step, it is essential to characterise how these forces (orange arrows) are perceived by plant roots. This could be achieved using e.g. modern LSM microscopes, artificial soils, calcium or FRET tension sensors to inform on the perception of forces induced by heterogeneous media [73]. (3) Finally, the mechanism of response to complex distribution of forces must be characterised. In this case, responses can be studied on simplified systems where position and magnitude of forces can be controlled accurately, using lab-on-chip device and more traditional developmental biology approaches. Experiments and data can then be used to formulate and test new concepts and biomechanical theories (black arrows). Computational models can test

- biomechanical theories in most realistic conditions using latest technologies, e.g. particle based simulations and computer hardware (4) and influence the design of new experiments (grey arrows).
- 246

247	Acknowledgements
248	This work was supported by the consolidator fellowship from the European Research Council ERC
249	SENSOILS-647857. We also thank the colleagues from the RoSoM project for inspiring discussions on
250	root biomechanics and discrete element modelling (Agropolis Fondation ID 1202-073; LabexAgro:
251	ANR-10-LABX- 001-01).
252	
253	

254 References

- 1. Lockhart JA: An analysis of irreversible plant cell elongation. Journal of Theoretical Biology 1965,
- **8**:264-275.
- 257 2. Barley K, Greacen E: Mechanical resistance as a soil factor influencing the growth of roots and
- underground shoots. *Advances in agronomy* 1967, **19**:1-43.
- 3. Silk WK, Erickson RO: **Kinematics of plant growth**. *Journal of Theoretical Biology* 1979, **76**:481-501.
- 4. Abdalla A, Hettiaratchi D, Reece A: **The mechanics of root growth in granular media**. *Journal of*
- 261 *Agricultural Engineering Research* 1969, **14**:236-248.
- 5. Silk WK, Wagner K: **Growth-sustaining water potential distributions in the primary corn root. a non**
- compartmented continuum model. *Plant Physiology* 1980, **66**:859-863.
- 6. Dexter A: **Mechanics of root growth**. *Plant and soil* 1987, **98**:303-312.
- 7. Clark L, Whalley W, Dexter A, Barraclough P, Leigh R: Complete mechanical impedance increases
- the turgor of cells in the apex of pea roots. *Plant, Cell & Environment* 1996, **19**:1099-1102.
- 8. Bengough A, Mullins C: Penetrometer resistance, root penetration resistance and root elongation
- rate in two sandy loam soils. *Plant and Soil* 1991, **131**:59-66.
- 9. Bengough AG, Loades K, McKenzie BM: Root hairs aid soil penetration by anchoring the root
- surface to pore walls. *Journal of Experimental Botany* 2016, **67**:1071-1078.
- 271 10. Colombi T, Kirchgessner N, Walter A, Keller T: Root tip shape governs root elongation rate under
- increased soil strength. *Plant Physiology* 2017, **174**:2289-2301.
- 273 11. Kolb E, Legue V, Bogeat-Triboulot M-B: Physical root-soil interactions. Physical Biology 2017,
- **14**:065004.
- 275 12. Faure A: Stress field developed by root growth: theoretical approach. *Journal of Agricultural*
- 276 Engineering Research 1994, **58**:53-67.
- 277 13. Jackson R, Moore L, Hoffmann W, Pockman W, Linder C: Ecosystem rooting depth determined
- 278 with caves and DNA. Proceedings of the National Academy of Sciences of the United States
- 279 *of America* 1999, **96**:11387-11392.

280 14. White RG, Kirkegaard JA: The distribution and abundance of wheat roots in a dense, structured 281 subsoil—implications for water uptake. *Plant, Cell & Environment* 2010, **33**:133-148. 282 15. Monshausen GB, Gilroy S: The exploring root—root growth responses to local environmental 283 conditions. *Current opinion in plant biology* 2009, **12**:766-772. 284 16. Mueth DM, Jaeger HM, Nagel SR: Force distribution in a granular medium. Physical Review E 285 1998, **57**:3164-3169. 17. Liu C, Nagel SR, Schecter D, Coppersmith S, Majumdar S: Force fluctuations in bead packs. Science 286 287 1995, **269**:513-515. 288 18. Coppersmith S, Liu C-h, Majumdar S, Narayan O, Witten T: Model for force fluctuations in bead 289 packs. *Physical Review E* 1996, **53**:4673-4685. 290 19. Hurley R, Hall S, Andrade J, Wright J: Quantifying interparticle forces and heterogeneity in 3D 291 granular materials. *Physical Review Letters* 2016, **117**:098005. 292 20. Zhang L, Lambert S, Nicot F: Discrete dynamic modelling of the mechanical behaviour of a granular 293 soil. International Journal of Impact Engineering 2017, 103:76-89. 294 21. Aranson IS, Tsimring LS: Patterns and collective behavior in granular media: Theoretical concepts. 295 *Reviews of Modern Physics* 2006, **78**:641-692. 296 22. Gnoli A, Lasanta A, Sarracino A, Puglisi A: Unified rheology of vibro-fluidized dry granular media: 297 From slow dense flows to fast gas-like regimes. Scientific Reports 2016, 6:38604. 298 23. Richard P, Philippe P, Barbe F, Bourlès S, Thibault X, Bideau D: Analysis by x-ray microtomography 299 of a granular packing undergoing compaction. *Physical Review E* 2003, **68**:020301-020301. 300 24. Vlahinić I, Andò E, Viggiani G, Andrade JE: Towards a more accurate characterization of granular 301 media: extracting quantitative descriptors from tomographic images. Granular Matter 2013, 302 **16**:9-21. 303 25. Guo Y, Curtis JS: Discrete Element Method Simulations for Complex Granular Flows. Annual 304 Review of Fluid Mechanics 2015, 47:21-46.

305	26. Nicot F, Xiong H, Wautier A, Lerbet J, Darve F: Force chain collapse as grain column buckling in
306	granular materials. <i>Granular Matter</i> 2017, 19 :18.
307	27. Bourrier F, Kneib F, Chareyre B, Fourcaud T: Discrete modeling of granular soils reinforcement by
308	plant roots. Ecological Engineering 2013, 61:646-657.
309	28. Fakih M, Delenne JY, Radjai F, Fourcaud T: Modeling root growth in granular soils: effects of root
310	stiffness and packing fraction. In EPJ Web of Conferences: EDP Sciences: 2017:14013.
311	29. Iskratsch T, Wolfenson H, Sheetz MP: Appreciating force and shape—the rise of
312	mechanotransduction in cell biology. Nature Reviews Molecular Cell Biology 2014, 15:825.
313	30. Hamilton ES, Schlegel AM, Haswell ES: United in diversity: mechanosensitive ion channels in
314	plants. Annual Review of Plant Biology 2015, 66:113-137.
315	31. Nakagawa Y, Katagiri T, Shinozaki K, Qi Z, Tatsumi H, Furuichi T, Kishigami A, Sokabe M, Kojima I,
316	Sato S: Arabidopsis plasma membrane protein crucial for Ca2+ influx and touch sensing in
317	roots. Proceedings of the National Academy of Sciences 2007, 104:3639-3644.
318	32. Mirabet V, Das P, Boudaoud A, Hamant O: The role of mechanical forces in plant morphogenesis.
319	Annual review of plant biology 2011, 62 :365-385.
320	33. Nagel KA, Putz A, Gilmer F, Heinz K, Fischbach A, Pfeifer J, Faget M, Blossfeld S, Ernst M, Dimaki
321	C: GROWSCREEN-Rhizo is a novel phenotyping robot enabling simultaneous measurements of
322	root and shoot growth for plants grown in soil-filled rhizotrons. Functional Plant Biology 2012,
323	39 :891-904.
324	34. Rellán-Álvarez R, Lobet G, Lindner H, Pradier P-L, Sebastian J, Yee M-C, Geng Y, Trontin C, LaRue
325	T, Schrager-Lavelle A: GLO-Roots: an imaging platform enabling multidimensional
326	characterization of soil-grown root systems. Elife 2015, 4.
327	35. Mooney SJ, Pridmore TP, Helliwell J, Bennett MJ: Developing X-ray computed tomography to non-
328	invasively image 3-D root systems architecture in soil. Plant and soil 2012, 352:1-22.

329	36. Keyes SD, Gostling NJ, Cheung JH, Roose T, Sinclair I, Marchant A: The Application of Contrast
330	Media for In Vivo Feature Enhancement in X-Ray Computed Tomography of Soil-Grown Plant
331	Roots. Microscopy and Microanalysis 2017, 23:538-552.
332	37. Keyes S, Cooper L, Duncan S, Koebernick N, Fletcher DM, Scotson C, Van Veelen A, Sinclair I,
333	Roose T: Measurement of micro-scale soil deformation around roots using four-dimensional
334	synchrotron tomography and image correlation. Journal of The Royal Society Interface 2017,
335	14 :20170560.
336	38. De Micco V, Arena C, Pignalosa D, Durante M: Effects of sparsely and densely ionizing radiation or
337	plants. Radiation and Environmental Biophysics 2011, 50:1-19.
338	39. Koebernick N, Daly KR, Keyes SD, George TS, Brown LK, Raffan A, Cooper LJ, Naveed M,
339	Bengough AG, Sinclair I, et al.: High-resolution synchrotron imaging shows that root hairs
340	influence rhizosphere soil structure formation. New Phytologist 2017, 216 :124-135.
341	40. Schloter M, Borlinghaus R, Bode W, Hartmann A: Direct identification, and localization of
342	Azospirillum in the rhizosphere of wheat using fluorescence-labelled monoclonal antibodies
343	and confocal scanning laser microscopy. <i>Journal of Microscopy</i> 1993, 171 :173-177.
344	41. Assmus B, Hutzler P, Kirchhof G, Amann R, Lawrence JR, Hartmann A: In situ localization of
345	Azospirillum brasilense in the rhizosphere of wheat with fluorescently labeled, rRNA-targeted
346	oligonucleotide probes and scanning confocal laser microscopy. Applied and Environmental
347	<i>Microbiology</i> 1995, 61 :1013-1019.
348	42. Downie H, Holden N, Otten W, Spiers AJ, Valentine TA, Dupuy LX: Transparent soil for imaging the
349	rhizosphere. <i>PLoS One</i> 2012, 7 :e44276.
350	43. Cost A-L, Ringer P, Chrostek-Grashoff A, Grashoff C: How to measure molecular forces in cells: a
351	guide to evaluating genetically-encoded FRET-based tension sensors. Cellular and molecular
352	bioengineering 2015, 8 :96-105.
353	44. Reynaud EG, Krzic U, Greger K, Stelzer EH: Light sheet-based fluorescence microscopy: more
354	dimensions, more photons, and less photodamage. HFSP Journal 2008, 2:266-275.

355	45. Rozbicki E, Chuai M, Karjalainen AI, Song F, Sang HM, Martin R, Knölker H-J, MacDonald MP,
356	Weijer CJ: Myosin II-mediated cell shape changes and cell intercalation contribute to primitive
357	streak formation. Nature Cell Biology 2015, 17:397.
358	46. Mickoleit M, Schmid B, Weber M, Fahrbach FO, Hombach S, Reischauer S, Huisken J: High-
359	resolution reconstruction of the beating zebrafish heart. <i>Nature Methods</i> 2014, 11 :919-922.
360	47. Reynaud EG, Peychl J, Huisken J, Tomancak P: Guide to light-sheet microscopy for adventurous
361	biologists. Nature Methods 2015, 12:30-34.
362	48. Reidt SL, O'Brien DJ, Wood K, MacDonald MP: Polarised light sheet tomography. Opt Express
363	2016, 24 :11239-11249.
364	49. Wilding D, Pozzi P, Soloviev O, Vdovin G, Sheppard CJ, Verhaegen M: Pupil filters for extending
365	the field-of-view in light-sheet microscopy. Optics Letters 2016, 41:1205-1208.
366	50. O'Callaghan FE, Braga RA, Neilson R, MacFarlane SA, Dupuy LX: New live screening of plant-
367	nematode interactions in the rhizosphere. Scientific Reports 2018, 8:1440.
368	51. Yang Z, Downie H, Rozbicki E, Dupuy LX, MacDonald MP: Light Sheet Tomography (LST) for in situ
369	imaging of plant roots. Optics Express 2013, 21:16239-16247.
370	52. Silverberg JL, Noar RD, Packer MS, Harrison MJ, Henley CL, Cohen I, Gerbode SJ: 3D imaging and
371	mechanical modeling of helical buckling in Medicago truncatula plant roots. Proc Natl Acad Sci
372	<i>USA</i> 2012, 109 :16794-16799.
373	53. Sampathkumar A, Krupinski P, Wightman R, Milani P, Berquand A, Boudaoud A, Hamant O,
374	Jönsson H, Meyerowitz EM: Subcellular and supracellular mechanical stress prescribes
375	cytoskeleton behavior in Arabidopsis cotyledon pavement cells. <i>Elife</i> 2014, 3 :e01967.
376	54. Mártonfalvi Z, Bianco P, Naftz K, Ferenczy GG, Kellermayer M: Force generation by titin folding.
377	Protein Science 2017.
378	55. Tordesillas A, Steer CAH, Walker DM: Force chain and contact cycle evolution in a dense granular
379	material under shallow penetration. Nonlinear Processes in Geophysics 2014, 21:505-519.

380	56. Kolb E, Hartmann C, Genet P: Radial force development during root growth measured by
381	photoelasticity. Plant and Soil 2012, 360:19-35.
382	57. Wendell DM, Luginbuhl K, Guerrero J, Hosoi AE: Experimental Investigation of Plant Root Growth
383	Through Granular Substrates. Experimental Mechanics 2011, 52 :945-949.
384	58. Ozoe K, Hida H, Kanno I, Higashiyama T, Notaguchi M: Early characterization method of plant root
385	adaptability to soil environments. In Micro Electro Mechanical Systems (MEMS), 2015 28th
386	IEEE International Conference on 18-22 Jan: IEEE: 2015:702-705.
387	59. Bizet F, Bengough AG, Hummel I, Bogeat-Triboulot MB, Dupuy LX: 3D deformation field in
388	growing plant roots reveals both mechanical and biological responses to axial mechanical
389	forces. Journal of Experimental Botany 2016, 67:5605-5614.
390	60. Brodu N, Dijksman JA, Behringer RP: Spanning the scales of granular materials through
391	microscopic force imaging. <i>Nature Communication</i> 2015, 6 :6361.
392	61. Massalha H, Korenblum E, Malitsky S, Shapiro OH, Aharoni A: Live imaging of root-bacteria
393	interactions in a microfluidics setup. Proceedings of the National Academy of Sciences of the
394	United States of America 2017, 114 :4549-4554.
395	62. Stanley CE, Grossmann G, Solvas XC, DeMello AJ: Soil-on-a-Chip: microfluidic platforms for
396	environmental organismal studies. Lab on a Chip 2016, 16:228-241.
397	63. Kitson PJ, Rosnes MH, Sans V, Dragone V, Cronin L: Configurable 3D-Printed millifluidic and
398	microfluidic 'lab on a chip'reactionware devices. Lab on a Chip 2012, 12:3267-3271.
399	64. Yanagisawa N, Sugimoto N, Arata H, Higashiyama T, Sato Y: Capability of tip-growing plant cells to
400	penetrate into extremely narrow gaps. Scientific Reports 2017, 7 :1403.
401	65. Anderson JR, Chiu DT, Wu H, Schueller O, Whitesides GM: Fabrication of microfluidic systems in
402	poly (dimethylsiloxane). Electrophoresis 2000, 21:27-40.
403	66. Neale SL, MacDonald MP, Dholakia K, Krauss TF: All-optical control of microfluidic components
404	using form birefringence. Nature Materials 2005, 4:530-533.

405	67. Baret J-C, Miller OJ, Taly V, Ryckelynck M, El-Harrak A, Frenz L, Rick C, Samuels ML, Hutchison JB,
406	Agresti JJ: Fluorescence-activated droplet sorting (FADS): efficient microfluidic cell sorting
407	based on enzymatic activity. Lab on a Chip 2009, 9:1850-1858.
408	68. Stanley CE, Shrivastava J, Brugman R, Heinzelmann E, Swaay D, Grossmann G: Dual-flow-RootChip
409	reveals local adaptations of roots towards environmental asymmetry at the physiological and
410	genetic levels. New Phytologist 2018, 217:1357-1369.
411	69. Lipiec J, Horn R, Pietrusiewicz J, Siczek A: Effects of soil compaction on root elongation and
412	anatomy of different cereal plant species. Soil and Tillage Research 2012, 121:74-81.
413	70. Kalogiros DI, Adu MO, White PJ, Broadley MR, Draye X, Ptashnyk M, Bengough AG, Dupuy LX:
414	Analysis of root growth from a phenotyping data set using a density-based model. Journal of
415	Experimental Botany 2016, 67 :1045-1058.
416	71. Sadeghi A, Mondini A, Del Dottore E, Mattoli V, Beccai L, Taccola S, Lucarotti C, Totaro M,
417	Mazzolai B: A plant-inspired robot with soft differential bending capabilities. Bioinspiration &
418	biomimetics 2016, 12 :015001.
419	72. Perfect E, Groenevelt P, Kay B, Grant C: Spatial variability of soil penetrometer measurements at
420	the mesoscopic scale. Soil and Tillage Research 1990, 16 :257-271.
421	73. Monshausen GB: Visualizing Ca 2+ signatures in plants. Current Opinion in Plant Biology 2012,
422	15 :677-682.