

Computation of edge metric dimension of barcycentric subdivision of Cayley graphs

Z S Mufti, M F Nadeem, Ali Ahmad, Z Ahmad

▶ To cite this version:

Z S Mufti, M F Nadeem, Ali Ahmad, Z Ahmad. Computation of edge metric dimension of barcycentric subdivision of Cayley graphs. 2018. hal-01902772

HAL Id: hal-01902772 https://hal.science/hal-01902772

Preprint submitted on 23 Oct 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Computation of edge metric dimension of barcycentric subdivision of Cayley graphs

Z. S. Mufti¹, M. F. Nadeem¹, Ali Ahmad², Z. Ahmad³

¹Department of Mathematics, COMSATS University Islamabad, Lahore Campus, Pakistan. zeeshansaleem0090gmail.com, mfaisalnadeem0ymail.com ²College of Computer Science & Information Systems, Jazan University, Jazan, Saudi Arabia. ahmadsms0gmail.com ³Khwaja Fareed University of Engineering and Information Technology Rahim Yar Khan, Pakistan zaheer0kfueit.edu.pk

Abstract

Let G = (V, E) be a connected graph, let $x \in V(G)$ be a vertex and $e = yz \in E(G)$ be an edge. The distance between the vertex x and the edge e is given by $d_G(x, e) = \min\{d_G(x, y), d_G(x, z)\}$. A vertex $t \in V(G)$ distinguishes two edges $e, f \in E(G)$ if $d_G(t, e) \neq d_G(t, f)$. A set $R \subseteq V(G)$ is an edge metric generator for G if every two edges of G are distinguished by some vertex of R. The minimum cardinality of R is called the edge metric dimension and is denoted by edim(G). In this paper, we compute the edge metric dimension of barcycentric subdivision of Cayley graphs $Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2)$.

Keywords: metric dimension, edge metric dimension, resolving set, barcycentric subdivision, Cayley graph

Mathematics Subject Classification: 05C12, 05C85, 05C90.

1 Introduction

The concept of metric dimension was introduced by Slater [25] and studied independently by Harary and Melter [12]. This problem has been investigated widely since then. The metric dimension has a lot of applications in different areas of science and technology. The concept of the edge metric dimension is a recent advancement in this line of research. Next we reveal some of the applications of metric dimension in various subjects. The metric dimension arises in many diverse areas, including navigation of robots [21], telecommunications [5], combinatorial optimization [24] and sonar and coast guard Loran [25] and applications to chemistry in [8, 9, 14]. Furthermore, this topic has some applications to problems of pattern recognition and image processing, some of which involve the use of hierarchical data structures [23]. Metric dimension of several interesting classes of graphs can be seen in [2, 1, 3, 4, 7, 15, 16, 17, 18, 22, 26].

Let G = (V, E) be a simple and connected graph. For a vertex $x \in V(G)$ distinguishes two vertices $y, z \in V(G)$ if $d_G(y, x) \neq d_G(z, x)$, where $d_G(x, y)$ denotes the length of the shortest path between the vertices x and y in G. A vertex set $R_1 \subseteq V(G)$ is a metric generator for G, if any pair of vertices of G is distinguished by at least one vertex of R_1 and R_1 is the resolving set of G. The minimum cardinality of any metric generator for G is the metric dimension of G, denoted by dim(G). Let $R_1 = \{r_1, r_2, \ldots, r_s\}$ be an ordered set of vertices of G and let $x \in V(G)$, then the representation $r(x|R_1)$ of x with respect to R_1 is the s-tuple $(d_G(x, r_1), d_G(x, r_2), d_G(x, r_3), \ldots, d_G(x, r_s))$. Since the set R_1 has the minimum cardinality, therefore this is also known as the basis of G, and its cardinality is called the metric dimension or location number [6].

Similarly, for $x \in V(G)$ be a vertex and $e = yz \in E(G)$ be an edge. The distance between the vertex x and the edge e is given by $d_G(x, e) = \min\{d_G(x, y), d_G(x, z)\}$. A vertex $t \in V(G)$ distinguishes two edges $e, f \in E(G)$ if $d_G(t, e) \neq d_G(t, f)$. A set $R \subseteq V(G)$ is an edge metric generator for G if every two edges of G are distinguished by some vertex of R. The minimum cardinality of R is called the edge metric dimension and is denoted by edim(G) [19]. Let $R = \{r_1, r_2, \ldots, r_t\}$ be an ordered set of vertices of G and let $e \in E(G)$, then the representation r(e|R) of e with respect to R is the t-tuple $(d_G(e, r_1), d_G(e, r_2), d_G(e, r_3), \ldots, d_G(e, r_t))$.

In addition, combined (mixed) form of these two parameters depicted above is of fascinate. A vertex $x \in V(G)$ distinguishes two elements (vertices or edges) $u, v \in V(G) \cup E(G)$ if $d_G(x, u) \neq d_G(x, v)$. A set $R^m \subseteq V(G)$ is a mixed metric generator for G if every two distinct elements (vertices or edges) of G are distinguished by some vertex of R^m . The smallest cardinality of R^m is the mixed metric dimension and is denoted by mdim(G) [20].

Geometrically, an operation that splits an edge into two edges by inserting a new vertex into the interior of an edge is known as subdividing an edge. If we are performing a sequence of edge-subdivision operations, then it is called *Subdividing a graph* G and resulting graph is called a *subdivision of the graph* G. The subdivision of graph can be used to convert a general graph into a simple graph. If we subdividing each edge of the graph G, then this subdivision is called the *barycentric subdivision* of G. Gross and Yellen [11] proved the results that the barycentric subdivision of any graph is a simple and bipartite graph.

A graph G is *planar* if it can be drawn in the plane without edge crossings. Subdivision of graphs play a very important role in characterization of planar graphs. A graph Gis planar if and only if every subdivision of G is planar. Two graphs are said to be homeomorphic if they are subdivisions of same graph G. The next theorem gives a nice characterization of planar graphs.

Theorem 1.1. [11] A graph is planar if and only if it does not contain a subdivision of K_5 or $K_{3,3}$.

In this paper, we study the edge metric dimension of barcycentric subdivision of Cayley graphs $Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2)$. We prove that these subdivisions of Cayley graphs have constant edge metric dimension and only three vertices chosen appropriately suffice to resolve all the vertices of these subdivision of Cayley graphs $Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2)$.

2 Results and Discussions

As expressed, there are a several graphs in which metric generator and edge metric generator are same. In this sense, one could believe that most likely any edge metric generator R is likewise a standard metric generator. In any case, this is again further far from the truth, despite the fact that there are a few families of graphs in which such actualities happen. Kelenc *et al.* [19] explained some comparison between the edge metric generator and standard metric generator in detailed. We show a few results concerning the edge metric dimension of graphs. The first importance result about the complexity is as follows:

Theorem 2.1. [19] Computing the edge metric dimension of graphs is NP -hard.

The edge metric dimension of Cartesian product of two paths P_r and P_t with r and t vertices is determined in the following proposition.

Proposition 2.2. [19] Let G be the grid graph $G = P_r \Box P_t$, with $r \ge t \ge 2$. Then edim(G) = dim(G) = 2.

Kelenc *et al.* [19] proved in the next proposition that the edge metric dimension of wheel graphs and observe it is strictly larger than the value for the metric dimension, except in the case $W_{1,3}$.

The wheel graph $W_{1,n}$ is the graph obtained from a cycle $C_n, n \ge 3$ by joining all vertices of C_n to an additional vertex. In [19], they determined the edge metric dimension of wheel graph $W_{1,n}$ in the following proposition:

Proposition 2.3. [19] Let $W_{1,n}$ be a wheel graph. Then

$$edim(W_{1,n}) = \begin{cases} n, & for n = 3, 4\\ n-1, & for n \ge 5 \end{cases}$$

The fan graph $F_{1,n}$ is the graph obtained by joining each vertices of a path P_n , to an additional vertex. In the next proposition, the edge metric dimension of fan graph $F_{1,n}$ is determined.

Proposition 2.4. [19] Let $F_{1,n}$ be a fan graph. Then

$$edim(F_{1,n}) = \begin{cases} n, & forn = 1, 2, 3\\ n-1, & forn \ge 4 \end{cases}$$

Kelenc *et al.* [19] also determined the edge metric dimension of path, cycle, complete graph, complete bipartite, cartesian product of cycles and bounds for some families of graphs.

3 The edge metric dimension of barcycentric subdivision of Cayley graphs $Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2)$

Let G be a semigroup, and let H be a nonempty subset of G. The Cayley graph Cay(G, H) of G relative to H is defined as the graph with vertex set G and edge set E(H) consisting of those ordered pairs (a, b) such that ha = b for some $h \in H$. Cayley graphs of groups are significant both in group theory and in constructions of interesting graphs with nice properties. The Cayley graph Cay(G, H) of a group G is symmetric or undirected if and only if $H = H^{-1}$.

The Cayley graphs $Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2), n \geq 3$, is a 3-regular graph which is also known as the cartesian product $C_n \Box P_2$ of a cycle of order n with a path of order 2. The Cayley graphs $Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2), n \geq 3$ consists of an inner n-cycle $a_1a_2a_3...a_n$, an outer n-cycle $x_1x_2x_3...x_n$ and n spokes $a_ix_i, 1 \leq i \leq n$. This implies that the order and size of $Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2)$ is 2n and 3n, respectively. The metric dimension of Cayley graphs $Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2)$ has been determined in [7] while the metric dimension of Cayley graphs $Cay(\mathbb{Z}_n : H)$ for all $n \geq 7$ and $H = \{\pm 1, \pm 3\}$ has been determined in [13].

The barcycentric subdivision graph $BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))$ can be obtained by splitting edges $a_i a_{i+1}$ by inserting a new vertices b_i , splitting edges $a_i x_i$ by inserting a new vertices c_i splitting edges $x_i x_{i+1}$ by inserting a new vertices y_i . From this we observe that, $BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))$ contains 5n vertices among of these 3n vertices of degree 2 and 2n vertices of degree 3 and 6n edges. In the next theorem, we prove that the metric dimension of the barcycentric subdivision $BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))$ of is constant and only three vertices appropriately chosen suffice to resolve all the vertices of the $BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))$.

Theorem 3.1. Let $BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))$ be the barcycentric subdivision of Cayley graphs $(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))$; then $edim(BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))) = 3$ for every $n \ge 6$.

Proof. We will prove the above equality by double inequalities.

Case 1. When n is even.

Let $R = \{a_1, a_{\frac{n}{2}+1, a_n}\} \subset V(BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2)))$, we show that R is a resolving set for $BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))$ in this case. For this we give representations of any edge of $E(BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2)))$ with respect to R. Representations for the edges of $BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))$ are

$$r(a_i b_i | R) = \begin{cases} (2i - 2, n - 2i + 1, 2i), & \text{for } 1 \le i \le \frac{n}{2} - 1\\ (n - 2, 1, n - 1), & \text{for } i = \frac{n}{2}\\ (2n - 2i + 1, 2i - n - 2, 2n - 2i - 1), & \text{for } \frac{n}{2} + 1 \le i \le n - 1\\ (1, n - 2, 0), & \text{for } i = n \end{cases}$$

and

$$r(b_{i}a_{i+1}|R) = \begin{cases} (2i-1, n-2i, 2i+1), & \text{for } 1 \le i \le \frac{n}{2} - 1\\ (n-1, 0, n-2), & \text{for } i = \frac{n}{2}\\ (2n-2i, 2i-n-1, 2n-2i-2), & \text{for } \frac{n}{2} + 1 \le i \le n-1\\ (0, n-1, 1), & \text{for } i = n \end{cases}$$

Representations for the set of interior edges of $BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))$ are $r(a_ic_i|R) = \begin{cases} (2i-2, n-2i+2, 2i), & \text{for } 1 \leq i \leq \frac{n}{2} \\ (2n-2i+2, 2i-n-2, 2n-2i), & \text{for } \frac{n}{2}+1 \leq i \leq n \end{cases}$ and

$$r(c_i x_i | R) = \begin{cases} (2i - 1, n - 2i + 3, 2i + 1), & \text{for } 1 \le i \le \frac{n}{2} \\ (2n - 2i + 3, 2i - n - 1, 2n - 2i + 1), & \text{for } \frac{n}{2} + 1 \le i \le n \end{cases}$$

Representations for the edges on the outer cycle of $BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))$ are

$$r(x_i y_i | R) = \begin{cases} (2i, n - 2i + 3, 2i + 2), & \text{for } 1 \le i \le \frac{n}{2} - 1\\ (n, 3, n + 1), & \text{for } i = \frac{n}{2}\\ (2n - 2i + 3, 2i - n, 2n - 2i + 1), & \text{for } \frac{n}{2} + 1 \le i \le n - 1\\ (3, n, 2), & \text{for } i = n \end{cases}$$

and

$$r(y_i x_{i+1}|R) = \begin{cases} (2i+1, n-2i+2, 2i+3), & \text{for } 1 \le i \le \frac{n}{2} - 1\\ (n+1, 2, n), & \text{for } i = \frac{n}{2}\\ (2n-2i+2, 2i-n+1, 2n-2i), & \text{for } \frac{n}{2} + 1 \le i \le n-1\\ (2, n+1, 3), & \text{for } i = n \end{cases}$$

We note that there are no two edges having the same representations implying that $edim(BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))) \leq 3$.

On the other hand, we show that $edim(BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))) \geq 3$. Suppose on contrary that $edim(BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))) = 2$, then there are the following possibilities to be discussed.

(1) Both vertices are in the inner cycle. Here are the following subcases.

• Both vertices belong to the set $\{a_i : 1 \le i \le n\}$. Without loss of generality, we can suppose that one resolving vertex is a_1 . Suppose that the second resolving vertex is a_k $(2 \le k \le \frac{n}{2}+1)$. Then for $2 \le k \le \frac{n}{2}$, we have $r(a_1c_1|\{a_1, a_k\}) = r(a_1b_n|\{a_1, a_k\}) = (0, 2k-2)$, and for $k = \frac{n}{2} + 1$, we have $r(a_1b_1|\{a_1, a_{\frac{n}{2}+1}\}) = r(a_1b_n|\{a_1, a_{\frac{n}{2}+1}\}) = (0, n-1)$, a contradiction.

• Both vertices belong to the set $\{b_i : 1 \leq i \leq n\}$. Without loss of generality, we can suppose that one resolving vertex is b_1 . Suppose that the second resolving vertex is b_k $(2 \leq k \leq \frac{n}{2} + 1)$. Then for $2 \leq k \leq \frac{n}{2}$, we have $r(a_1c_1|\{b_1, b_k\}) = r(a_1b_n|\{(b_1, b_k\}) = (1, 2k-1)$, and for $k = \frac{n}{2} + 1$, we have $r(a_1b_1|\{b_1, b_{\frac{n}{2}+1}\}) = r(a_2b_1|\{b_1, b_{\frac{n}{2}+1}\}) = (0, n-1)$, a contradiction.

• One vertex belong to the set $\{a_i : 1 \le i \le n\}$ and the second vertex belong to the set $\{b_i : 1 \le i \le n\}$. Without loss of generality, we can suppose that one resolving vertex is a_1 . Suppose that the second resolving vertex is b_k $(1 \le k \le \frac{n}{2} + 1)$. Then for $1 \le k \le \frac{n}{2}$, we have $r(a_1b_n|\{a_1, b_k\}) = r(a_1c_1|\{a_1, b_k\}) = (0, 2k - 1)$, and for $k = \frac{n}{2} + 1$, we have $r(a_1b_1|\{a_1, b_{\frac{n}{2}+1}\}) = r(a_1c_1|\{a_1, b_{\frac{n}{2}+1}\}) = (0, n - 1)$, a contradiction.

(2) Both vertices are in the interior vertices. Without loss of generality, we can suppose that one resolving vertex is c_1 . Suppose that the second resolving vertex is c_k $(2 \le k \le \frac{n}{2}+1)$. Then for $2 \le k \le \frac{n}{2}+1$, we have $r(x_1c_1|\{c_1,c_k\}) = r(a_1c_1|\{c_1,c_k\}) = (0,2k-1)$, a contradiction.

(3) Both vertices are in the outer cycle. Due to the symmetry of the graph, this case is analogous to case (1).

(4) One vertex is in the inner cycle and the other one is in the set of interior vertices. Here are the two subcases.

• One vertex is in the set $\{a_i : 1 \leq i \leq n\}$ and the other one is in the set of interior vertices $\{c_i : 1 \leq i \leq n\}$. Without loss of generality, we can suppose that one resolving vertex is a_1 . Suppose that the second resolving vertex is c_k $(1 \leq k \leq \frac{n}{2} + 1)$. Then for k = 1, we have $r(a_1b_1|\{a_1,c_1\}) = r(a_1b_n|\{a_1,c_1\}) = (0,1)$. For $2 \leq k \leq \frac{n}{2}$, we have $r(a_1b_n|\{a_1,c_k\}) = r(a_1c_1|\{a_1,c_k\}) = (0,2k-1)$ and for $k = \frac{n}{2} + 1$, we have $r(a_1b_n|\{a_1,c_{\frac{n}{2}+1}\}) = r(a_1b_1|\{a_1,c_{\frac{n}{2}+1}\}) = (0,2n)$, a contradiction.

• One vertex is in the set $\{b_i : 1 \leq i \leq n\}$ and the other one is in the set of interior vertices $\{c_i : 1 \leq i \leq n\}$. Without loss of generality, we can suppose that one resolving vertex is b_1 . Suppose that the second resolving vertex is c_k $(1 \leq k \leq \frac{n}{2} + 1)$. Then for k = 1, we have $r(x_1y_1|\{b_1, c_1\}) = r(x_1y_n|\{b_1, c_1\}) = (3, 1)$. For $2 \leq k \leq \frac{n}{2}$, we have $r(a_1c_1|\{b_1, c_k\}) = r(a_1b_n|\{b_1, c_k\}) = (1, 2k - 1)$ and for $k = \frac{n}{2} + 1$, we have $r(c_2x_2|\{b_1, c_{\frac{n}{2}+1}\}) = r(a_nb_n|\{b_1, c_{\frac{n}{2}+1}\}) = (2, n - 1)$, a contradiction.

(5) One vertex is in the outer cycle and the other one is in the set of interior vertices. Due to the symmetry of the graph, this case is analogous to case (4).

(6) One vertex is in the inner cycle and the other one is in the outer cycle. Here are the following subcases.

• One vertex is in the set $\{a_i : 1 \le i \le n\}$ and the other one is in the set $\{x_i : 1 \le i \le n\}$. Without loss of generality, we can suppose that one resolving vertex is a_1 . Suppose that the second resolving vertex is x_k $(1 \le k \le \frac{n}{2} + 1)$. Then for k = 1, we have $r(a_1b_1|\{a_1, x_1\}) = r(a_1b_n|\{a_1, x_1\}) = (0, 2)$. For $2 \le k \le \frac{n}{2} + 1$, we have $r(a_1b_1|\{a_1, x_k\}) = r(a_1c_1|\{a_1, x_k\}) = (0, 2k - 1)$, a contradiction.

• One vertex is in the set $\{a_i : 1 \leq i \leq n\}$ and the other one is in the set $\{y_i : 1 \leq i \leq n\}$. Without loss of generality, we can suppose that one resolving vertex is a_1 . Suppose that the second resolving vertex is y_k $(1 \leq k \leq \frac{n}{2} + 1)$. Then for k = 1, we have $r(a_1b_1|\{a_1, y_1\}) = r(a_1b_n|\{a_1, y_1\}) = (0, 3)$. For $2 \leq k \leq \frac{n}{2}$,

we have $r(a_1b_1|\{a_1, y_k\}) = r(a_1c_1|\{a_1, y_k\}) = (0, 2k)$ and for $k = \frac{n}{2} + 1$, we have $r(a_1b_1|\{a_1, y_{\frac{n}{2}+1}\}) = r(a_1c_1|\{a_1, y_{\frac{n}{2}+1}\}) = (0, n + 1)$, a contradiction. • One vertex is in the set $\{b_i : 1 \le i \le n\}$ and the other one is in the set $\{y_i : 1 \le i \le n\}$.

• One vertex is in the set $\{b_i : 1 \le i \le n\}$ and the other one is in the set $\{y_i : 1 \le i \le n\}$. Without loss of generality, we can suppose that one resolving vertex is b_1 . Suppose that the second resolving vertex is y_k $(1 \le k \le \frac{n}{2} + 1)$. Then for k = 1, we have $r(a_1b_1|\{b_1, y_1\}) = r(a_2b_1|\{b_1, y_1\}) = (0, 3)$. For k = 2, we have $r(x_1y_1|\{b_1, y_2\}) = r(a_3c_3|\{b_1, y_2\}) = (3, 2)$. For $3 \le k \le \frac{n}{2} + 1$, we have $r(x_2y_2|\{b_1, y_k\}) = r(a_3c_3|\{b_1, y_k\}) = (3, 2k - 4)$, a contradiction.

Hence from above it follows that there is no resolving set with two vertices for $V(BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2)))$ implying that $edim(BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))) \neq 2$ in this case. Therefore, $edim(BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))) = 3$.

Case 2. When n is odd.

Let $R = \{a_1, b_{\lceil \frac{n}{2} \rceil}, a_n\} \subset V(BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2)))$, we show that R is a resolving set for $BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))$ in this case. For this we give representations of any edge of $E(BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2)))$ with respect to R.

Representations for the edges of the inner cycle of $BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))$ are

$$r(a_i b_i | R) = \begin{cases} (2i - 2, n + 1 - 2i, 2i), & \text{for } 1 \le i \le \lceil \frac{n}{2} \rceil - 1\\ (2i - 2, n + 1 - 2i, 2n - 2i - 1), & \text{for } i = \lceil \frac{n}{2} \rceil\\ (2n - 2i + 1, 2i - n - 2, 2n - 2i - 1), & \text{for } \lceil \frac{n}{2} \rceil + 1 \le i \le n - 1\\ (1, n - 2, 0), & \text{for } i = n. \end{cases}$$

and

$$r(b_{i}a_{i+1}|R) = \begin{cases} (2i-1, n-2i, 2i+1), & \text{for } 1 \le i \le \left\lceil \frac{n}{2} \right\rceil - 2\\ (2i-1, n-2i, 2n-2i-2), & \text{for } i = \left\lceil \frac{n}{2} \right\rceil - 1\\ (2n-2i, 2i-n-1, 2n-2i-2), & \text{for } \left\lceil \frac{n}{2} \right\rceil \le i \le n-1\\ (0, n-1, 1), & \text{for } i = n \end{cases}$$

Representations for the edges on the outer cycle of $BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))$ are

$$r(x_i y_i | R) = \begin{cases} (2i, n+3-2i, 2i+2), & \text{for } 1 \le i \le \lceil \frac{n}{2} \rceil - 1\\ (2i, n-2i+3, 2n-2i+1), & \text{for } i = \lceil \frac{n}{2} \rceil\\ (2n-2i+3, 2i-n, 2n-2i+1), & \text{for } \lceil \frac{n}{2} \rceil + 1 \le i \le n-1\\ (3, n, 2), & \text{for } i = n \end{cases}$$

and

$$r(y_i x_{i+1}|R) = \begin{cases} (2i+1, n-2i+2, 2i+3), & \text{for } 1 \le i \le \lceil \frac{n}{2} \rceil - 2\\ (2i+1, n-2i+2, 2n-2i), & \text{for } i = \lceil \frac{n}{2} \rceil - 1\\ (2n-2i+2, 3, 2n-2i), & \text{for } i = \lceil \frac{n}{2} \rceil\\ (2n-2i+2, 2i-n+1, 2n-2i), & \text{for } \lceil \frac{n}{2} \rceil + 1 \le i \le n-1\\ (2, n+1, 3), & \text{for } i = n \end{cases}$$

Representations for the set of interior edges of $BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))$ are

$$r(a_i c_i | R) = \begin{cases} (2i - 2, n - 2i + 2, 2i), & \text{for } 1 \le i \le \lceil \frac{n}{2} \rceil - 1\\ (2i - 2, n - 2i + 2, 2n - 2i), & \text{for } i = \lceil \frac{n}{2} \rceil\\ (2n - 2i + 2, 2i - n - 2, 2n - 2i), & \text{for } \lceil \frac{n}{2} \rceil + 1 \le i \le n \end{cases}$$

$$r(c_i x_i | R) = \begin{cases} (2i - 1, n - 2i + 3, 2i + 1), & \text{for } 1 \le i \le \lceil \frac{n}{2} \rceil - 1\\ (2i - 1, n - 2i + 3, 2n - 2i + 1), & \text{for } i = \lceil \frac{n}{2} \rceil\\ (2n - 2i + 3, 2i - n - 1, 2n - 2i + 1), & \text{for } i = \lceil \frac{n}{2} \rceil + 1 \le i \le n \end{cases}$$

Again we see that there are no two vertices having the same representations which implies that $edim(BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))) \leq 3$.

On the other hand, suppose that $edim(BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))) = 2$, then there are the same possibilities as in case (1) and contradictions can be deduced analogously. This implies that $edim(BS(Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2))) = 3$ in this case, which completes the proof.

References

- A. Ahmad, M. Baca, S. Sultan, Minimal doubly resolving sets of Necklace graph, Mathematical report, 20(70). No. 2,(2018).
- [2] A. Ahmad, M. Imran, O. Al-Mushayt and S.A.H. Bokhary, On the metric dimension of barcycentric subdivision of Cayley graphs $Cay(\mathbb{Z}_n \oplus \mathbb{Z}_2)$, *Miskolc Mathematical Notes*, **16(2)**(2015), 637-646.
- [3] R.F. Bailey, P.J. Cameron, Basie size, metric dimension and other invariants of groups and graphs, Bull. London Math. Soc., 43(2011) 209–242.
- [4] R.F. Bailey, K. Meagher, On the metric dimension of Grassmann graphs, Discret. Math. Theo. Comput. Sci., 13(4)(2011) 97–104.
- [5] Z. Beerloiva, F. Eberhard, T. Erlebach, A. Hall, M. Hoffmann, M. Mihalk, L. Ram, Network discovery and verification, *IEEE J. Sel. Area Comm.*, 24 (2006) 2168–2181.
- [6] P. S. Buczkowski, G. Chartrand, C. Poisson and P. Zhang, On k-dimensional graphs and their bases, *Periodica Math.Hung.*, 46(1) (2003) 9–15.
- [7] J. Cceres, C. Hernando, M. Mora, I.M. Pelayoe, M.L. Puertas, C. Seara, D.R. Wood, On the metric dimension of cartesian products of graphs, *SIAM J. Discrete Math.*, 21(2007) 423–441.
- [8] G. Chartrand, L. Eroh, M. A. Johnson, and O. R. Oellermann, Resolvability in graphs and the metric dimension of a graph, *Discrete Applied Mathematics*, **105** (2000) 99– 113.
- [9] G. Chartrand, C. Poisson, and P. Zhang, Resolvability and the upper dimension of graphs, Computers and Mathematics with Applications, 39 (2000) 19–28.

- [10] M. Fehr, S. Gosselin, O. Oellermann, The metric dimension of Cayley digraphs, *Discrete Math.*, **306**(2006) 31–41.
- [11] J. L. Gross and J. Yellen, Graph theorey and its applications, Chapman & Hall/CRC, New York, 2006.
- [12] F. Harary, R.A. Melter, On the metric dimension of a graph, Ars. Combin., 2(1976) 191–195.
- [13] I. Javaid, M. N. Azhar and M. Salman, Metric dimension and determining number of Cayley graphs, World Applied Sciences Journal, 18(12)(2012) 1800–1812.
- [14] M. A. Johnson, Structure-activity maps for visualizing the graph variables arising in drug design, Journal of Biopharmaceutical Statistics, 3 (1993) 203–236.
- [15] M. Imran, S.A.H. Bokhary, A.Ahmad and A. Semaničová–Feňovčíková, On classes of regular graphs with constant metric dimension, *Acta Mathematica Scientia, Series* B, 33 (1), (2013), 187-206.
- [16] M. Imran, A. Q. Baig, A. Ahmad, Families of plane graphs with constant metric dimension, Utilitas Math., 88(2012), 43–57.
- [17] M. Imran, S.A.H. Bokhary and A. Ahmad, On the metric dimension of two classes of convex polytopes, *JCMCC*, 77 (2011), 51–63.
- [18] M. Imran, H.M.A. Siddiqui, Computing the metric dimension of convex polytopes generated by the wheel related graphs, Acta Math. Hungar., 149(2016) 10–30.
- [19] A. Kelenc, N. Tratnik, and I. G. Yero, Uniquely identifying the edges of a graph: the edge metric dimension, Manuscript, (2016).
- [20] A. Kelenc, D. Kuziak, A. Taranenko, and I. G. Yero, On the mixed metric dimension of graphs, *Applied Mathematics and Computation*, **314** (2017) 429–438.
- [21] S. Khuller, B. Raghavachari, A. Rosenfeld, Landmarks in graphs, Discrete Appl. Math., 70 (1996) 217–229.
- [22] J. Kratica, V. Kovacevic-Vujcic, M. Cangalovic, M. Stojanovic, Minimal doubly resolving sets and the strong metric dimension of some convex polytopes, *Appl. Math. Comput.*, **218**(2012) 9790–9801.
- [23] R. A. Melter, I. Tomescu, Metric bases in digital geometry, Computer Vision, Graphics, and Image Processing 25 (1) (1984) 113–121.
- [24] A. Sebö and E. Tannier, On metric generators of graphs, Math. Oper. Res., 29(2004) 383–393.
- [25] P.J. Slater, Leaves of trees, Proc. 6th Southeastern Conf. on Combinatorics, Graph Theory, and Computing, Congr. Numer. (1975) 549–559.

- [26] T. Vetrik and A. Ahmad, Computing the metric dimension of the categorial product of graphs, *International Journal of Computer Mathematics*, **94(2)** (2017), 363–371.
- 10