



Full Waveform Inversion Adjoint Studies MATHIAS 2018

Pierre Jacquet, Hélène Barucq, Julien Diaz, Henri Calandra

► To cite this version:

Pierre Jacquet, Hélène Barucq, Julien Diaz, Henri Calandra. Full Waveform Inversion Adjoint Studies MATHIAS 2018. MATHIAS 2018 - Computational Science Engineering & Data Science by TOTAL, Oct 2018, Serris, France. hal-01907804

HAL Id: hal-01907804

<https://hal.science/hal-01907804>

Submitted on 29 Oct 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Full Waveform Inversion Adjoint Studies

MATHIAS 2018

Pierre Jacquet

pierre.jacquet@inria.fr

Barucq Hélène, Diaz Julien, Calandra Henri

First year PhD Student
Inria - Magique 3D - DIP

Pau, FRANCE



Outline

The acoustic model

The continuous model

The discretized problem

Assets of Bernstein polynomials

Adjoint Studies

FWI Introduction

Adjoint then Discretized

Discretize then Adjoint

Some Results

Consistency of the Adjoint Solution

FWI Preliminary test

Qualitative Cost Function Gradient Study

Outline



The acoustic model

The continuous model

The discretized problem

Assets of Bernstein polynomials

Adjoint Studies

FWI Introduction

Adjoint then Discretized

Discretize then Adjoint

Some Results

Consistency of the Adjoint Solution

FWI Preliminary test

Qualitative Cost Function Gradient Study

The continuous model

Continuous Problem:

$$\begin{cases} \frac{1}{\rho_0 v_p^2} \frac{\partial p}{\partial t} + \frac{\partial v}{\partial x} = f_p & \text{in } \Omega \\ \rho_0 \frac{\partial v}{\partial t} + \frac{\partial p}{\partial x} = f_v & \text{in } \Omega \end{cases}$$

$$\begin{cases} p(t=0) = 0 \\ v(t=0) = 0 \\ \frac{\partial p}{\partial t} + v_p \frac{\partial p}{\partial x} \cdot \mathbf{n} = 0 \quad \text{on } \Gamma \end{cases}$$




Figure: 1D Domain Model

The discretized model

Discretized Problem:

$$\begin{cases} \frac{\partial \bar{\mathbf{P}}}{\partial t} = A_{pv} \bar{\mathbf{V}} + A_{pp} \bar{\mathbf{P}} + \bar{\mathbf{F}}_p \\ \frac{\partial \bar{\mathbf{V}}}{\partial t} = A_{vp} \bar{\mathbf{P}} \end{cases}$$




Figure: 1D Discretized Domain

- ▶ Discontinuous Galerkin space discretization
- ▶ Different time-schemes (RK4,AB3)
- ▶ Two polynomial basis (Lagrange and Bernstein)
- ▶ Constant velocity (v_p) per cells
- ▶ Constant density (ρ_0) per cells

Bernstein/DG Properties

Bernstein formulation:

$$B_{ijkl}^N = C_{ijkl}^N \lambda_0^i \lambda_1^j \lambda_2^k \lambda_3^l \quad \text{with: } C_{ijkl}^N = \frac{N!}{i! j! k! l!}$$



$P[X^5]$ Bernstein basis


Bernstein/DG Properties

Bernstein formulation:

$$B_{ijkl}^N = C_{ijkl}^N \lambda_0^i \lambda_1^j \lambda_2^k \lambda_3^l \quad \text{with: } C_{ijkl}^N = \frac{N!}{i! j! k! l!}$$

Easy Derivative expression:

$$\frac{\partial B_\alpha^N}{\partial \lambda_p} = N B_{\alpha - e_p}^{N-1} \quad \text{with: } \alpha = (i, j, k, l)$$



$P[X^5]$ Bernstein basis

Bernstein/DG Properties

Bernstein formulation:


$$B_{ijkl}^N = C_{ijkl}^N \lambda_0^i \lambda_1^j \lambda_2^k \lambda_3^l \quad \text{with: } C_{ijkl}^N = \frac{N!}{i! j! k! l!}$$

Easy Derivative expression:

$$\frac{\partial B_\alpha^N}{\partial \lambda_p} = N B_{\alpha-e_p}^{N-1} \quad \text{with: } \alpha = (i, j, k, l)$$

Sparse Degree Elevation operator:


$$B_\alpha^{N-1} = \sum_{p=0}^d \frac{\alpha_p + 1}{N} B_{\alpha+e_p}^N$$




$P[X^5]$ Bernstein basis

Bernstein/DG Properties

Unique boundary condition values :




$P(X^5)$ Lagrange basis




$P(X^5)$ Bernstein basis

⇒ Same Flux Management

Derivative-Operator Analysis



3D Lagrange D matrix



3D Bernstein D matrix

[1] Chan J. and Warburton T.

GPU-Accelerated Bernstein Bézier Discontinuous Galerkin Methods for Wave Problems
SIAM Journal on Scientific Computing 2017

1D Results




Figure: Operators NZVs as a function of the order

Outline



The acoustic model

- The continuous model

- The discretized problem

- Assets of Bernstein polynomials

Adjoint Studies

- FWI Introduction

- Adjoint then Discretized

- Discretize then Adjoint

Some Results

- Consistency of the Adjoint Solution


- FWI Preliminary test

- Qualitative Cost Function Gradient Study

FWI Introduction





10




FWI Introduction



10



Data and Simulation



$$J(p) = \frac{1}{2} \|R\mathbf{p} - \text{data}\|^2$$

► Find v_p and ρ_0 to minimize J

Adjoint Studies



*Continuous
Direct Problem*


Adjoint Studies




11




Adjoint Studies



Adjoint Studies



Adjoint Studies



AtD : Adjoint then Discretized Strategy

$$J(p) = \frac{1}{2} \|Rp - data\|^2$$

$$\begin{cases} \frac{1}{\rho_0 \mathbf{v}_p^2} \frac{\partial p}{\partial t} + \nabla \cdot \mathbf{v} = f_p \\ \rho_0 \frac{\partial \mathbf{v}}{\partial t} + \nabla p = 0 \\ p(t=0) = 0 \\ \mathbf{v}(t=0) = 0 \\ \frac{\partial p}{\partial t} + \mathbf{v}_p \nabla p \cdot \mathbf{n} = 0 \quad \text{on } \Gamma \end{cases}$$

$$\begin{cases} \frac{1}{\rho_0 \mathbf{v}_p^2} \frac{\partial \lambda_1}{\partial t} + \nabla \cdot \lambda_2 = \frac{\partial J}{\partial p} \\ \rho_0 \frac{\partial \lambda_2}{\partial t} + \nabla \lambda_1 = 0 \\ \lambda_1(t=T) = 0 \\ \lambda_2(t=T) = 0 \\ \frac{\partial \lambda_1}{\partial t} + \mathbf{v}_p \nabla \lambda_1 \cdot \mathbf{n} = 0 \quad \text{on } \Gamma \end{cases}$$

$$t \in [0, T]$$


$$t \in [T, 0]$$

AtD : Adjoint then Discretized Strategy

$$J(\bar{\mathbf{P}}) = \frac{1}{2} ||R\bar{\mathbf{P}} - data||^2$$

$$\begin{cases} \frac{\partial \bar{\mathbf{P}}^n}{\partial t} = A_{pv} \bar{\mathbf{V}}^n + A_{pp} \bar{\mathbf{P}}^n + \bar{\mathbf{F}}_p^n \\ \frac{\partial \bar{\mathbf{V}}^n}{\partial t} = A_{vp} \bar{\mathbf{P}}^n \end{cases}$$

$$\begin{cases} \frac{\partial \bar{\Lambda}_1^n}{\partial t} = +A_{pv} \bar{\Lambda}_2^n + A_{pp} \bar{\Lambda}_1^n + \bar{\mathbf{D}}_p^n \\ \frac{\partial \bar{\Lambda}_2^n}{\partial t} = A_{vp} \bar{\Lambda}_1^n \end{cases}$$



DtA : Discretize then Adjoint Strategy

$$\frac{\partial \bar{\mathbf{U}}^n}{\partial t} = A \bar{\mathbf{U}}^n + \bar{\mathbf{F}}^n \quad \text{With : } \bar{\mathbf{U}} = \begin{pmatrix} \bar{\mathbf{P}} \\ \bar{\mathbf{V}} \end{pmatrix}, A = \begin{pmatrix} A_{pp} & A_p \\ A_v & 0 \end{pmatrix}, \bar{\mathbf{F}} = \begin{pmatrix} \bar{\mathbf{F}}_p \\ 0 \end{pmatrix}$$

All time scheme can be summed-up such as :

$$L \bar{\mathbf{U}} = E \bar{\mathbf{F}}$$

We are looking for a Discrete Adjoint state satisfying :

$$L^* \bar{\Lambda} = -R^*(R \bar{\mathbf{U}} - \text{data})$$

DtA : Discretize then Adjoint Strategy

Example with RK4

RK4 time-scheme leads to :

$$\bar{\mathbf{U}}^{n+1} = B\bar{\mathbf{U}}^n + C_0\bar{\mathbf{F}}^n + C_{\frac{1}{2}}\bar{\mathbf{F}}^{n+\frac{1}{2}} + C_1\bar{\mathbf{F}}^{n+1}$$

$$L\bar{\mathbf{U}} = E\bar{\mathbf{F}} = \bar{\mathbf{G}}$$
$$\begin{pmatrix} I & & & \\ -B & I & & \\ & -B & I & \\ & & \ddots & \ddots \\ & & & -B & I \end{pmatrix} \begin{pmatrix} \bar{\mathbf{U}}^0 \\ \bar{\mathbf{U}}^1 \\ \bar{\mathbf{U}}^2 \\ \vdots \\ \bar{\mathbf{U}}^n \end{pmatrix} = \begin{pmatrix} \bar{\mathbf{G}}^0 \\ \bar{\mathbf{G}}^1 \\ \bar{\mathbf{G}}^2 \\ \vdots \\ \bar{\mathbf{G}}^n \end{pmatrix}$$

So :

$$L^* = \begin{pmatrix} I & -B^* & & & \\ & I & -B^* & & \\ & & \ddots & \ddots & \\ & & & I & -B^* \\ & & & & I \end{pmatrix}$$

Adjoint test



$$\langle L\bar{\mathbf{U}}, \bar{\boldsymbol{\lambda}} \rangle = \langle \bar{\mathbf{U}}, L^*\bar{\boldsymbol{\lambda}} \rangle$$

Adjoint test

$$\langle L\bar{\mathbf{U}}, \bar{\boldsymbol{\Lambda}} \rangle = \langle \bar{\mathbf{U}}, L^*\bar{\boldsymbol{\Lambda}} \rangle$$

$$\begin{cases} L\bar{\mathbf{U}} = E\bar{\mathbf{F}} = \bar{\mathbf{G}} \\ \bar{\mathbf{U}}(t=0) = 0 \end{cases}$$

$$\begin{cases} L^*\bar{\boldsymbol{\Lambda}} = -R^*(R\bar{\mathbf{U}} - \text{data}) = \bar{\mathbf{D}} \\ \bar{\boldsymbol{\Lambda}}(t=T) = 0 \end{cases}$$



Adjoint test

$$\langle L\bar{\mathbf{U}}, \bar{\boldsymbol{\lambda}} \rangle = \langle \bar{\mathbf{U}}, L^*\bar{\boldsymbol{\lambda}} \rangle$$

$$\begin{cases} L\bar{\mathbf{U}} = E\bar{\mathbf{F}} = \bar{\mathbf{G}} \\ \bar{\mathbf{U}}(t=0) = 0 \end{cases}$$

$$\begin{cases} L^*\bar{\boldsymbol{\lambda}} = -R^*(R\bar{\mathbf{U}} - \text{data}) = \bar{\mathbf{D}} \\ \bar{\boldsymbol{\lambda}}(t=T) = 0 \end{cases}$$



$$\langle E\bar{\mathbf{F}}, \bar{\boldsymbol{\lambda}} \rangle = \langle \bar{\mathbf{U}}, -R^*(R\bar{\mathbf{U}} - \text{data}) \rangle$$

Adjoint test

$$\langle L\bar{\mathbf{U}}, \bar{\boldsymbol{\Lambda}} \rangle = \langle \bar{\mathbf{U}}, L^*\bar{\boldsymbol{\Lambda}} \rangle$$

$$\begin{cases} L\bar{\mathbf{U}} = E\bar{\mathbf{F}} = \bar{\mathbf{G}} \\ \bar{\mathbf{U}}(t=0) = 0 \end{cases}$$

$$\begin{cases} L^*\bar{\boldsymbol{\Lambda}} = -R^*(R\bar{\mathbf{U}} - \text{data}) = \bar{\mathbf{D}} \\ \bar{\boldsymbol{\Lambda}}(t=T) = 0 \end{cases}$$



$$\langle E\bar{\mathbf{F}}, \bar{\boldsymbol{\Lambda}} \rangle = \langle \bar{\mathbf{U}}, -R^*(R\bar{\mathbf{U}} - \text{data}) \rangle$$

$$\langle \bar{\mathbf{G}}, \bar{\boldsymbol{\Lambda}} \rangle = \langle \bar{\mathbf{U}}, \bar{\mathbf{D}} \rangle$$

Adjoint test succeeds \iff operator L^* well established

Outline



The acoustic model

- The continuous model

- The discretized problem

- Assets of Bernstein polynomials

Adjoint Studies

- FWI Introduction

- Adjoint then Discretized

- Discretize then Adjoint

Some Results

- Consistency of the Adjoint Solution

- FWI Preliminary test

- Qualitative Cost Function Gradient Study

Adjoint test



Adjoint test passed for :

- ▶ Lagrange Operators
- ▶ Bernstein Operators
- ▶ Runge Kutta 4 time-scheme
- ▶ Adams Bashforth 3
time-scheme
- ▶ With a canonical space
inner-product
 $(\langle u, v \rangle_x = \sum_i u_i v_i)$
- ▶ With a M-space inner
product
 $(\langle u, v \rangle_X^M = \langle Mu, v \rangle_x)$

Adjoint test

Adjoint test passed for :

- ▶ Lagrange Operators
- ▶ Bernstein Operators
- ▶ Runge Kutta 4 time-scheme
- ▶ Adams Bashforth 3 time-scheme
- ▶ With a canonical space inner-product
 $\langle u, v \rangle_x = \sum_i u_i v_i$
- ▶ With a M-space inner product
 $\langle u, v \rangle_X^M = \langle Mu, v \rangle_x$


```
./run
--- Adjoint test ----
inner product UP/DUDP 553123.57586755091
inner product GPGU/QPQU 553123.57586756046

./run
--- Adjoint test ----
inner product UP/DUDP -75077.332007383695
inner product GPGU/QPQU -75077.332007386358


./run
--- Adjoint test ----
inner product UP/DUDP 125669.89223600870
inner product GPGU/QPQU 125669.89223600952

./run
--- Adjoint test ----
inner product UP/DUDP -132852.64215701097
inner product GPGU/QPQU -132852.64215701059
```

Non consistency of the Adjoint solution



With the AtD strategy




With the DtA strategy using the canonical inner-product (Lagrange+RK4)

Adjoint test succeeds !


[1] Sei Alain and Symes William

A Note on Consistency and Adjointness for Numerical Schemes
1997

FWI Preliminary test (for all strategies)




Target model




Initial model (iter=0)


FWI Preliminary test (for all strategies)




Target model



Initial model (iter=0)



Intermediate model (iter=20)



Final model (iter=50)

Qualitative Cost Function Gradient Study




Figure: Comparison between a Reference Gradient and the FWI Gradient with AtD strategy (Bernstein elements and RK4 time scheme)

Qualitative Cost Function Gradient Study




Figure: Comparison between a Reference Gradient and the FWI Gradient with DtA strategy (Bernstein elements and RK4 time scheme)

Qualitative Cost Function Gradient Study




Figure: Comparison between a Reference Gradient and the FWI Gradient with AtD strategy (Lagrange elements and RK4 time scheme)

Qualitative Cost Function Gradient Study




Figure: Comparison between a Reference Gradient and the FWI Gradient with DtA strategy (Lagrange elements and RK4 time scheme)

Conclusion and Perspectives



Conclusion :

- ▶ Adjoint then Discretized strategy works
- ▶ Discretized then Adjoint strategy has unexpected results
(Gradient formulation ? Bug ?)
- ▶ The adjoint state is not consistent by using the Discretized and
Adjoint strategy (but Adjoint test succeeds)

Conclusion and Perspectives



25

Conclusion :

- ▶ Adjoint then Discretized strategy works
- ▶ Discretized then Adjoint strategy has unexpected results
(Gradient formulation ? Bug ?)
- ▶ The adjoint state is not consistent by using the Discretized and
Adjoint strategy (but Adjoint test succeeds)

Perspectives :

- ▶ Complementary 1D tests
- ▶ 2D FWI + tests
- ▶ 3D FWI + tests
- ▶ Coupling SEM/DG elements (Aurélien Citrain's thesis)