



HAL
open science

Fatty acids and stable isotopes as tracers of food assimilation in fish-oyster-macroalgae integrated multitrophic aquaculture ponds

Sarah Nahon, Goncalo Villa de Brito, Hugo Quental-Ferreira, Joël Aubin,
Maria Emilia Cunha

► **To cite this version:**

Sarah Nahon, Goncalo Villa de Brito, Hugo Quental-Ferreira, Joël Aubin, Maria Emilia Cunha. Fatty acids and stable isotopes as tracers of food assimilation in fish-oyster-macroalgae integrated multitrophic aquaculture ponds. World Aquaculture 2018, Aug 2018, Montpellier, France. 848 p., 2018, Abstract Book. hal-01913008

HAL Id: hal-01913008

<https://hal.science/hal-01913008>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

FATTY ACIDS AND STABLE ISOTOPES AS TRACERS OF FOOD ASSIMILATION IN FISH-OYSTER-MACROALGAE INTEGRATED MULTITROPHIC AQUACULTURE PONDS

Sarah Nahon*, Goncalo Villa de Brito, Hugo Quental-Ferreira, Joel Aubin, Maria Emilia Cunha

UMR 1419, Nutrition, Métabolisme et Aquaculture NuMÉA, AquaPôle INRA
64310 Saint Pée-sur-Nivelle, France
sarah.nahon@inra.fr

Integrated MultiTrophic Aquaculture (IMTA) is based on the production of fed species (*e.g.*, finfish) along with extractive species which re-use the inorganic (*e.g.*, seaweeds) and organic (*e.g.*, suspension-feeders) nutrients excess. To increase the development of IMTA practices, it is important to understand the trophic relationship within the system. In Mediterranean earthen ponds, meagre (*Argyrosomus regius*), white seabream (*Diplodus sargus*) and mullet (*Mugil chepalus*) are three fish species of different trophic levels that are good candidates to be cultured with oysters (*Crassostrea gigas*) and sea lettuce (*Ulva flexuosa*) but their trophic behavior are not well known under polyculture. The aim of this study was to identify the food sources used by fish and oysters using fatty acids (FA) as well as carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) stable isotopes.

Within the Era Net COFASP project IMTA-Effect, experiments were conducted in 6 earthen ponds at IPMA's Estação de Piscicultura de Olhão, south of Portugal. Similar biomass and proportions of meagre, white seabream and mullet were introduced in each pond and three different combinations of organisms were tested: fish-oysters, fish-macroalgae, fish-oysters-macroalgae. According to the combination, oyster spats were introduced while autochthonous macroalgae were let grown or removed from the ponds. Fish were fed daily with a commercial diet. At the end of the experiment, fish and oyster tissues were collected as well as all potential food sources present in ponds: commercial pellet, macroalgae, suspended particulate organic matter (SPOM) and polychetes. FA composition as well as $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of samples were determined.

The combinations of organisms had no effect on FA composition of fish and oysters. FA composition of fish were significantly different between species. FA composition of meagre and white seabream was close to those of feed. FA composition of mullet was farther than those of feed with lower proportion of C18:1 \square 9 and C18:2 \square 6 and higher proportion of C22:6 \square 3. Compared to feed, polychetes also have lower proportion of C18:1 \square 9 and C18:2 \square 6 and higher proportion of C22:6 \square 3. FA markers of macroalgae were not found in fish. FA composition of oysters was similar to those of SPOM and FA markers of commercial feed were not detected. The combinations of species had no effect on $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of fish. Meagre, white seabream and mullet were slightly enriched in ^{13}C and ^{15}N in respect to feed. However, mullet were more ^{13}C -enriched than the other fish species and closer to polychetes. Oysters cultured in the fish-oysters-macroalgae combination were ^{13}C and ^{15}N depleted compared to those from the fish-oysters combination. Similar results were found for SPOM.

Our results highlighted that meagre and white seabream only consumed commercial feed while mullets consumed feed and probably, some polychetes whatever the combination of organisms. Macroalgae were not consumed by fish but influenced inorganic C and N recycling in ponds. Oysters fed on SPOM but not on particulate feed. FA and stable isotope analysis are two complementary trophic tracers powerful to study trophic transfers between species co-cultivated in IMTA ponds.

AQUA 2018



Montpellier France
August 25-29 2018



#We R Aquaculture

We are the producers, the investors, the suppliers, the processors, the vendors, the scientists, the educators, the students and the consumers of farmed aquatic products.

gold sponsor



silver sponsor



eas premium sponsor



was premier sponsors



conference sponsors



industry sponsors

