

1 **Annex (glossary) : Advances in computational modeling approaches of pituitary gonadotropin**
2 **signaling**

3

4 **I. Ordinary Differential equations**

5 1) *Ordinary Differential Equation*. Mathematical equation that links the time derivative of a
6 function y at time t with its current value at that time. In general, such an equation has
7 the form

8
$$\frac{dy(t)}{dt} = f(t, y(t)), \forall t \geq 0, \quad y(t = 0) = y_0. \quad (\text{Equation 1})$$

10 Given initial condition y_0 at time $t = 0$, and function f , appropriate analytical and/or
11 numerical tools allow one to solve equation 1 to obtain the unknown function $t \mapsto y(t)$.
12 This formalism is extensively used in mathematical physiology [1].

13 2) *(Non) Autonomous dynamical systems*. Equation 1 is said to be autonomous when the
14 function f does not explicitly depend on time t . Non-autonomous dynamical systems are
15 widely used in mathematical physiology to represent the impact of external rhythms (or
16 inputs) not specifically included in the model. The same dichotomy autonomous / non-
17 autonomous applies to all dynamical systems considered in this note.

18 3) *(Piecewise) Linear ODE*. Equation 1 is said to be linear if function f is a linear function
19 in y . Classical one-dimensional examples yield to the Malthusian exponential growth
20 model or the radioactive exponential decay (first order or one-step processes). The ODE
21 is piecewise linear if the function f is piecewise linear (its graph consist of line
22 segments).

23 4) *(Un)Damped harmonic oscillator*. The harmonic oscillator is a two-dimensional linear
24 ODE, giving rise to oscillatory solution (like the pendulum or mass connected to
25 springs). The harmonic oscillator is said to be damped if the amplitude of the oscillations

1 vanishes as time increases (decay to equilibrium position) or undamped otherwise
2 (periodic solution).


- 3 5) *Goodwin model*. The Goodwin model describes negative feedback oscillators in cellular
4 systems (such as lactose production in bacteria); its simplest form is given by a three-
5 dimensional ODE

$$\begin{cases} \frac{dy_1}{dt} = \gamma_1 (g(y_3) - y_1) , \\ \frac{dy_2}{dt} = \gamma_2 (y_1 - y_2) , \\ \frac{dy_3}{dt} = \gamma_3 (y_2 - y_3) , \end{cases} \quad (\text{Equation 2})$$

7 where function g represents the feedback mechanism.

- 8 6) *Excitable systems*. Excitability refers to a phenomenon where a system has a single
9 stable attractor, and two ways to return to equilibrium after perturbations. For small
10 perturbations away from equilibrium, the return is monotonic; however, for
11 perturbations beyond a threshold, the return is not monotonic, and undergoes a large
12 excursion before settling down.
- 13 7) *The FitzHugh-Nagumo model*. The FitzHugh-Nagumo equation is a two-dimensional
14 ODE that describes the essential features of the firing of a nerve. It is able to reproduce
15 qualitative features of neural spikes, which are a typical example of excitability. It can
16 also exhibit periodic solutions.
- 17 8) *Stiff nonlinear ODE with several timescales*. A system of linear ODEs is said stiff if the
18 ratio between the largest and the smallest eigenvalue is large. By extension, for nonlinear
19 systems, an ODE is said stiff if it involves several times scales of various different
20 orders of magnitude, and requires dedicated numerical solvers to deal with.
- 21 9) *Biochemical reaction network*. A biochemical reaction network is defined by
22 • *Species*. A finite set $\mathcal{S} = \{S_1, \dots, S_d\}$ of $d \geq 1$ species.

1 • *Reactions.* A finite set $\mathcal{R} = \{R^1, \dots, R^n\}$ of $n \geq 1$ relations between linear
2 combinations of species, that is



4 where α_j^i, β_j^i are non-negative integers and called respectively reactant and product
5 stoichiometric coefficients for reaction i and species j . The set of reactions defines a
6 directed graph between linear combinations of species.

7 10) *Law of mass action.* A deterministic mass-action model associated with the biochemical
8 reaction network $(\mathcal{S}, \mathcal{R})$ is given by the set of ordinary differential equations

$$9 \quad \frac{dy_j}{dt} = \sum_{i=1}^n (\beta_j^i - \alpha_j^i) k_i \prod_{l=1}^d y_l^{\alpha_l^i}, \quad j = 1, 2, \dots, d. \quad (\text{Equation 3})$$

10 The law of mass-action assumes that the rate of reaction R^i is proportional to the
11 concentration of its reactants to the power of their stoichiometry. Constant k_i is called
12 the rate constant of reaction R^i .

14 II. Ordinary Differential equations with event or jumps

- 15 1) *Reset decision functions.* A function that resets the value of the solution of an ODE when
16 a specific condition is met (for instance when the solution exceeds a given threshold).
- 17 2) *Stepwise functions.* Functions that are defined piecewisely. The right-hand side of the
18 ODE can change according to some specific conditions (for instance when the solution
19 exceeds a given threshold).
- 20 3) *Threshold mediated transition.* Formalism that includes generic jumps for the value of
21 the solution of an ODE when a specific condition is met (for instance when the solution
22 exceeds a given threshold).

- 1 4) *Impulse ordinary differential equations*. Class of differential equations combining ODE
2 with discrete jumps of the solution when specific conditions are met.
- 3 5) *Deterministic jump discontinuities*. Modification of the current value of the solution
4 ruled by a non-continuous jump, which is pre-defined deterministically (no randomness
5 in the instant of jump nor the magnitude of the jump).
- 6 6) *Dirac Delta functions*. Mathematical formalism representing impulses. A Dirac delta
7 function can be seen as a function that equals zero everywhere except in one point where
8 it is equal to infinity.

9 **III. Algebraic-integro-differential equations**

- 10 1) *Convolution integral*. Function given by an integral combining two functions in a very
11 specific way. The convolution integral of two functions f and g is given by

$$12 \qquad I(t) = \int_0^t f(x)g(t-x)dx . \qquad \text{(Equation 4)}$$

13 In a pharmacokinetic context, if f is the rate of production of a chemical substance, and
14 g the survival probability of that substance up to a time t , then $I(t)$ gives its current level
15 at time t .

- 16 2) *Algebraic-integro-differential equation*. Equation involving algebraic relations,
17 derivatives and integrals of functions. Algebraic relations give constraints to the values
18 of one or several functions; differential terms involve the net rate of change of the
19 functions; integrals typically involve the cumulative effect of the history of the values of
20 a function.

21 **IV. Delay differential equations**

22 Mathematical equations linking the time derivative of a function y at time t with its current
23 value at that time, together with the values at anterior times. For DDE involving a single

1 fixed time delay, the equation has the form

$$2 \quad \frac{dy(t)}{dt} = f(t, y(t), y(t - t_d)), \forall t \geq 0, \quad y(s) = y_0(s), s \in [-t_d, 0). \quad (\text{Equation 5})$$

3 Given initial condition y_0 on times $[-t_d, 0)$, and function f , appropriate analytical and/or
4 numerical tools allow one to solve equation 5 to obtain the unknown function $t \mapsto y(t)$. This
5 formalism is extensively used in mathematical physiology [1]. The delay may for instance
6 represent the time needed to synthesize a molecule (which is not modeled in great details).

7 **V. Partial differential equations**

8 Mathematical equations that link the partial derivatives of a function y of several variables
9 (typically time and space, or other structuring variables like age, maturity) with its current
10 value at that time.

11 1) *(Nonlinear) diffusion equations.* The linear diffusion equation, also called the heat
12 equation, describes the constant isotropic diffusion of particles (or variation of
13 temperature), and is written in its simplest one-dimensional form as

$$14 \quad \frac{\partial u(t, x)}{\partial t} = \frac{\partial}{\partial x} \left(D \frac{\partial u(t, x)}{\partial x} \right) \quad (\text{Equation 6})$$

15 When D is non constant and depends on the solution itself ($D = D(x, u)$), the latter
16 equation is called a nonlinear diffusion equation, and can represent crowding and/or
17 anisotropic effects.

18 2) *Reaction diffusion equation.* A class of PDE combining diffusion equation 6 and
19 biochemical reaction model 3.

20 3) *Structured cell populations.* Class of models involving PDE to describe the time
21 evolution of cell populations together with some of their characteristics (like age,
22 volume etc.).

23 **VI. Stochastic differential equations**

1 Mathematical equation that links the derivatives of a function y at time t with its current
2 value at that time, perturbed by random dynamical systems (like Brownian motion for
3 instance). The unknown function then becomes a random function with a given probability
4 distribution, and is generally called a stochastic process.

5 1) *Stochastic point process, intensity*. A class of stochastic processes that is a collection of
6 points randomly located on some underlying mathematical space. Point processes can be
7 used as mathematical models of phenomena representable as points in some type of
8 space (like discrete events in time). Roughly, for a point process, the intensity refers to
9 the function ruling the (mean) rate of apparition of new points. In molecular biology, the
10 transcription process (synthesis of new mRNA upon stimulation) is typically modeled
11 using such formalism.

12 2) *(Weibull) renewal process*. A renewal process is a point process for which the time
13 between any two successive events only depends on the value of the process at the
14 latest event that has occurred. A Weibull renewal process is a particular kind of renewal
15 process for which the interevent times follow a Weibull distribution.

16 3) *Generalized gamma density*. A specific probability distribution with three parameters
17 (generalizing the exponential, the Weibull and the Gamma distribution), generally used
18 to represent the complete duration of a sequence of coordinated random events (like the
19 cell cycle for instance).

20 4) *Ornstein-Uhlenbeck process*. Stochastic process describing the velocity of a massive
21 Brownian particle under the influence of friction.

22 **VII. Stability and Bifurcation for dynamical systems**

23 Stability and bifurcation theory for dynamical systems aims to characterize the long time
24 behavior of a dynamical system, and in particular to know if the system reaches an
25 equilibrium or not as time goes to infinity.

1) *Locally asymptotically stable*. An equilibrium for a dynamical system is locally asymptotically stable if any trajectory starting from a small perturbation away from that equilibrium converges to the same equilibrium as time goes to infinity.

2) *(Hopf) bifurcation*. Critical point where a system's stability switches: in a (supercritical) Hopf bifurcation, a stable equilibrium point becomes unstable and a periodic solution arises.

VIII. Statistical tools

1) *Deconvolution methods*. Statistical algorithms aiming to recovering a signal that has been convoluted by some other signal. For instance, recording the signal I from equation 4, one wishes to recover f (knowing some properties of g).

2) *Regularization scheme*. Regularization is a process of introducing additional information in order to solve an ill-posed problem or to prevent overfitting. In deconvolution methods, this consists typically in imposing some restriction to the unknown signal in order to be able to better identify its parameters.

IX. Pharmacology

1) *Physiologically-based pharmacokinetic model*. Mathematical modeling technique for predicting the absorption, distribution, metabolism and excretion of synthetic or natural chemical substances in humans and other animal species. These models typically consists of system of (ordinary) differential equations.

2) *Equilibrium pharmacology model*. Class of models aiming to describe dose-response relationship (without kinetics). These models can often be derived as equilibrium of deterministic mass-action models like equation 3.

3) *Operational model*. A classical example of pharmacology models describing agonist-induced dose-response relationships. The model was first introduced by [2].

1 **REFERENCES**

- 2 [1] Keener, J. P. and Sneyd, J. (1998). *Mathematical Physiology*. Number 8 in *Interdisciplinary Applied Mathematics*.
3 Springer-Verlag New York, springer-verlag new york edition.
- 4 [2] Black, J. W. and Leff, P. (1983). Operational models of pharmacological agonism. *Proceedings of the Royal society*
5 *of London. Series B. Biological sciences*, 220(1219):141–162.

6