



**HAL**  
open science

## Recovery of Mediterranean steppe vegetation after cultivation: legacy effects on plant composition, soil properties and functional traits

Juliane Helm, Thierry Dutoit, Arne Saatkamp, Solveig Franziska Bucher, Matthias Leiterer, Christine Römermann

### ► To cite this version:

Juliane Helm, Thierry Dutoit, Arne Saatkamp, Solveig Franziska Bucher, Matthias Leiterer, et al.. Recovery of Mediterranean steppe vegetation after cultivation: legacy effects on plant composition, soil properties and functional traits. Applied Vegetation Science, 2019, Applied Vegetation Science, 22, pp.71-84. 10.1111/avsc.12415 . hal-01915741

**HAL Id: hal-01915741**

**<https://hal.science/hal-01915741>**

Submitted on 8 Nov 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Recovery of Mediterranean steppe vegetation after cultivation: legacy effects on  
2 plant composition, soil properties and functional traits

3

4 Juliane Helm<sup>1,2</sup>, Thierry Dutoit<sup>3</sup>, Arne Saatkamp<sup>4</sup>, Solveig Franziska Bucher<sup>1</sup>, Matthias Leierer<sup>5</sup> &  
5 Christine Römermann<sup>1,6</sup>

6

7 **Helm, J.** (Corresponding author, juliane.helm@bgc-jena.mpg.de)<sup>1,2</sup>

8 **Dutoit, T.** (thierry.dutoit@imbe.fr)<sup>3</sup>

9 **Saatkamp, A.** (arne.saatkamp@imbe.fr)<sup>4</sup>

10 **Bucher, S. F.** (solveig.franziska.bucher@uni-jena.de)<sup>1</sup>

11 **Leierer, M.** (Matthias.Leierer@tll.thueringen.de)<sup>5</sup>

12 **Römermann, C.** (christine.roemermann@uni-jena.de)<sup>1,6</sup>

13

14 <sup>1</sup>Institute of Ecology and Evolution, Department of Plant Biodiversity, Friedrich-Schiller-University  
15 Jena, Philosophenweg 16, 07743 Jena, Germany

16 <sup>2</sup>Current address: Max-Planck Institute for Biogeochemistry, Department of Biogeochemical Processes,  
17 Hans Knöll Str. 10, 07745 Jena, Germany

18 <sup>3</sup>Institut Méditerranéen de Biodiversité et d'Ecologie, Avignon Univ, Aix Marseille Univ, CNRS, IRD,  
19 IMBE, IUT site Agroparc, 337 Chemin des Meinajaries BP 61207, F-84911 Avignon cedex 09, France

20 <sup>4</sup>Institut Méditerranéen de Biodiversité et d'Ecologie, Aix Marseille Univ, CNRS, IRD, Avignon Univ,  
21 IMBE, Facultés St Jérôme case 421, F-13397 Marseille cedex 20, France

22 <sup>5</sup>Thüringer Landesanstalt für Landwirtschaft, Naumburger Str. 98, 07702 Jena, Germany

23 <sup>6</sup>German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, 04103 Leipzig,  
24 Germany

25

26

27

28

29 **Abstract**

30 **Questions:** Are the vegetation structure and soil properties of fallows similar to adjacent  
31 species-rich Mediterranean steppe communities 35 years after abandonment? Is there a  
32 spontaneous re-development towards the original steppe vegetation after re-introduction of the  
33 traditional grazing system? Can differences in functional trait composition be identified  
34 according to different land-use types? Do traits of selected species affect the ability to re-  
35 colonize fallow land?

36

37 **Location:** Mediterranean steppe of La Crau (Southeastern France)

38

39 **Methods:** We conducted 80 vegetation surveys and soil analyses in the steppe and adjacent  
40 fallow land in six locations in La Crau in 2015; for three locations, data from 2001 was included.  
41 To compare the functional composition of steppe vs. fallow land, community-weighted means  
42 (CMW) of specific leaf area (SLA), leaf dry matter content (LDMC) and canopy height were  
43 calculated. To test whether trait variability is associated with the re-colonization success of 12  
44 selected target species, we measured SLA, LDMC, canopy height, plant width and aboveground  
45 biomass on eight replicate individuals per site and species.

46

47 **Results:** Fallow land was characterized by lower species richness and elevated phosphorus and  
48 potassium content in the soil. Comparing vegetation relevés between 2001 and 2015 revealed  
49 that the fallows were developing towards steppe vegetation. CWMs of SLA were significantly  
50 lower and CWM of LDMC higher in steppe than in fallow land. There was no similarity in trait  
51 composition between steppe species that successfully re-colonized the fallow-land community,  
52 and no evidence that intraspecific trait variability is associated with re-colonization success of  
53 these species.

54

55 **Conclusions:** Even after decades of abandonment and re-introduction of the grazing system,  
56 the vegetation structure, soil properties and plant functional trait composition of fallows differ  
57 from those of adjacent steppe communities. These results illustrate the need for active  
58 restoration to accelerate the recovery of former arable land to the target steppe communities.

59

60 **Keywords:** Community trait means; Cultivation; Dry Grassland; Grazing; Intraspecific trait  
61 variability; Leaf dry matter content; Natural recovery; Restoration; Specific leaf area

62

63 **Nomenclature:** Jäger & Werner (2002); Pavon (2014)

64

65 **Abbreviations:** CV = coefficient of variation; CWM = community-weighted mean; ITV =  
66 intraspecific trait variability; LDMC = leaf dry matter content; SLA = specific leaf area

67

68 **Running head:** Recovery of Mediterranean steppe vegetation

69

## 70 **Introduction**

71 Semi-natural grasslands including the steppes of the Mediterranean basin are among  
72 Europe's most species-rich ecosystems (Willems 1983; WallisDeVries et al. 2002).  
73 Nevertheless, the 20<sup>th</sup> century's agricultural intensification caused a dramatic decrease in steppe  
74 grasslands worldwide and led to fragmentation as well as eutrophication of these habitats,  
75 leading to a decrease in plant diversity (Poschlod et al. 1998; White et al. 2000; Karlik &  
76 Poschlod 2009). To protect the species diversity and to better connect semi-natural grasslands,  
77 grassland remnants need to be conserved or, if already altered, restored. For the restoration of  
78 these species rich grasslands, several techniques have already been proposed (Kiehl et al. 2010  
79 for review) as the natural colonization of characteristic grassland species into former intensively  
80 cultivated areas is usually very slow (Römermann et al. 2005, Buisson et al. 2006). It has  
81 already been shown that it might take several decades or centuries until the species composition  
82 and diversities of grasslands return (Forey & Dutoit 2012); accordingly, active restoration  
83 measures are typically needed.

84 To evaluate restoration success, traditional restoration studies assess the floristic similarity  
85 between restored and original (target) vegetation, focussing on species composition as well as  
86 diversity (Jaunatre et al., 2013). However, these indicators are highly dependent on site  
87 characteristics that determine the local species-pool (Stuble et al. 2017). A complementary way  
88 of assessing restoration success is to focus on the restoration of the functional composition  
89 (Laughlin 2014, Engst et al. 2016), which is especially relevant for those sites where original  
90 species are lost from the local species pool. The analyses of plant functional traits can provide  
91 insights into how land-use changes affect vegetation dynamics and function (e.g. Römermann  
92 et al. 2009, Breitschwerdt et al. 2015), but this trait-based approach has rarely been examined  
93 in connection with restoration success in Mediterranean dry grasslands.

94 Various habitat filters (e.g., environmental factors such as climate conditions, soil properties,  
95 or disturbance) allow only species with certain trait values to establish (Keddy 1992; Weiher &  
96 Keddy 1995; Diaz et al. 1998; Venn et al. 2011). The species that pass these filters are  
97 necessarily similar, which eventually leads to trait convergence (Diaz et al. 1998; Wilson 2007;  
98 Laughlin et al. 2012). But, species also need to be dissimilar enough to avoid competition,  
99 according to the limiting similarity hypothesis, which eventually leads to trait divergence  
100 (MacArthur & Levins 1967; Wilson 2007; Cornwell & Ackerly 2009; Laughlin et al. 2012).  
101 Comparing the functional composition of original vs. changed plant communities and analyzing  
102 the trait composition of target species that have already successfully re-established in the  
103 restoration sites seem promising as this approach complements the taxonomic evaluation of  
104 restoration success, therewith leading to a better understanding of the factors preventing  
105 successful restoration.

106 The aim of this study was to evaluate whether the taxonomic and functional composition of  
107 fallow lands can develop into the direction of original steppe communities without having  
108 established active restoration measures such as the addition of propagules.

109 We performed vegetation relevés to assess whether fallow lands re-developed into the  
110 direction of the target steppe vegetation with respect to the species composition and diversity.  
111 To elucidate the potential natural development of fallow land vegetation towards the original  
112 steppe vegetation we analyzed temporal trends in vegetation changes by comparing vegetation  
113 relevés and soil analyses from 2001 with results for 2015.

114 We explored the functional composition (community-weighted mean = CWM) of traits  
115 related to plant performance and growth rate and as indicators of competitive ability and  
116 environmental tolerance of species (Garnier 1992; Reich et al. 1997; Poorter et al. 2009;  
117 Lauterbach et al. 2013; Pérez-Harguindeguy et al. 2013). Nutrient-rich environments such as  
118 former agricultural land typically favor plants with high specific leaf area (SLA) and low leaf  
119 dry matter content (LDMC). Vice versa, species with trait values indicating nutrient

120 conservation strategies, slow growth rates and high longevity should dominate in nutrient poor  
121 environments (Chapin et al. 1993; Garnier 1992; Lavorel & Garnier 2002; Pérez-Harguindeguy  
122 et al. 2013; Poorter & Garnier 1999; Poorter & Remkes 1990; Römermann et al. 2009; Villar  
123 et al. 2005). Canopy height reflects the competitive ability of plant communities through better  
124 light assessment (Gaudet & Keddy 1988; Westoby et al. 2002), yet the investment in height  
125 incurs costs for construction and maintenance of the stem (Falster & Westoby 2003) and thus  
126 should be positively related to soil nutrient availability.

127 Intraspecific variation patterns in plant functional traits are a major determinant of the ability  
128 to respond and evolve with climate change and changing environmental conditions (Nicotra et  
129 al. 2010; Laforest-Lapointe et al. 2014; Bucher et al. 2017; Bucher et al. 2018). In some cases,  
130 trait variability can even be more important than particular trait values: species showing high  
131 functional variability may successfully re-establish because they manage to adapt to changing  
132 environmental conditions (Albert et al. 2011; Jung et al. 2010). In addition to the above  
133 mentioned traits we included additional plant width and biomass to test whether high  
134 intraspecific trait variability is related to re-establishment success. Both traits reflect  
135 competitive strength and should thus relate positively to soil nutrients (Pérez-Harguindeguy et  
136 al. 2013).

137 To summarize, this study combines a taxonomic with a trait-based approach to i) identify  
138 the effects of former land uses on present vegetation composition, species diversity and  
139 physical-chemical soil parameters, and ii) investigate whether re-development towards the  
140 original steppe vegetation can be detected after a further 14 years of recovery under traditional  
141 grazing conditions (2001 vs. 2015). Moreover, we iii) investigate whether differences in CWM  
142 of plant height, SLA and LDMC between fallow land and original steppe vegetation are still  
143 observable and attempt to iv) determine whether the traits of successful colonizers show higher  
144 functional variability than those of species that have not re-established on fallow lands yet.

## 145 **Materials and Methods**

### 146 *Study sites*

147 The plain of La Crau (Fig. 1) is the last remaining xeric steppe (Devaux et al. 1983) in  
148 southeastern France (43°33'N, 4°52'W). The area was formed by ancient river deposits during  
149 the Pleistocene period between 2 000 000 BP and 30 000 BP (Molliex et al. 2013). It has a  
150 Mediterranean climate with a frequent, strong northwesterly wind, the Mistral. Siliceous  
151 pebbles cover approximately 50% of La Crau, and plant roots are prevented from reaching the  
152 alluvial water table by an impermeable conglomerate lying 40–60 cm deep (Colomb & Roux  
153 1978). Traditionally, the land has been used for itinerant sheep grazing, probably since early  
154 Neolithic times (Buisson & Dutoit 2006). Industry, military activities, quarries, highways and  
155 intensive agriculture led to fragmentation and reduction of the semi-natural steppe vegetation  
156 from approximately 60 000 ha to 11 500 ha in the 20<sup>th</sup> century (Buisson & Dutoit 2006; Deverre  
157 1996; Etienne et al. 1998). During the period of melon cultivation (1965–1985), intensive  
158 fertilization, herbicides and deep ploughing changed the soil structure and degraded the  
159 conglomerate (Römermann et al. 2004, 2005). In contrast, cereals (1950–1966) were cultivated  
160 without deep ploughing and using less fertilizer (Devaux et al. 1983; Masip 1991). At the time  
161 of abandonment, no active restoration operations were realized. Grazing, representing an  
162 important factor for the maintenance of these grasslands, was re-introduced immediately when  
163 cultivation ceased (Dureau & Bonnefon 1998). The recovery of fallow land in our study area  
164 was thus not a strictly passive restoration technique. The flocks of sheep link remaining steppe  
165 patches (the target) and fallow lands and potentially facilitate the dispersal and establishment  
166 of steppe species into fallow lands (Fischer et al. 1996).

### 167 *Vegetation surveys*

168 To investigate whether type of cultivation and time since abandonment play a role in  
169 restoration success on fallow land, six locations containing a total of 16 study sites differing in

170 their land-use histories were examined (Table 1, Fig. 1). One long-abandoned and one recently  
171 abandoned cereal field were included, to represent different time spans.

172 We conducted vegetation relevés on 20 m x 20 m sites, placing five replicate 2 m x 2 m  
173 quadrats at each corner and in the middle (hereafter referred to as 'plots'). Species cover was  
174 estimated using the scale of Braun-Blanquet (1964). At each location, at least two sites  
175 consisting of one undisturbed steppe plot and one adjacent fallow land plot grazed by the same  
176 flock of sheep were studied. The locations Petit Abondoux, Poulagère and Couliès were  
177 sampled in May 2014 (Tanet 2014; Fig. 1), and the locations Peau de Meau, Grand Abondoux  
178 and Valigne in April 2015 (Fig. 1).

179 Percentages of vegetation cover, stone cover and bare ground were estimated on the five  
180 plots at the time of vegetation sampling. To estimate the standing crop, we cut the total  
181 aboveground biomass on three 0.5 m x 0.5 m randomly chosen quadrat within each site. The  
182 leaf area index (LAI), the ratio of foliage area to ground surface area (Watson 1947), of each of  
183 the five plots was measured using the LAI-2200C Plant Canopy Analyzer. Species richness  $S$   
184 representing the total number of species, Shannon diversity index ( $H_s$ ) and Evenness ( $E$ ) were  
185 calculated for each plot (Mühlenberg 1993).

### 186 ***Soil sampling***

187 To identify the effect of former land use on physical-chemical soil parameters, three replicate  
188 mixed soil samples of the upper soil layer (0–10 cm) were collected randomly within each site.  
189 After drying and sieving with a 2 mm mesh sieve, soil samples were analyzed at the INRA soil  
190 analysis laboratory (*Institut National de la Recherche Agronomique*, Arras, France) according  
191 to standardized protocols. Organic carbon (C in g kg<sup>-1</sup>) and total nitrogen content (N in g kg<sup>-1</sup>)  
192 were analyzed via the dry combustion method using a CHN elemental analyzer (Vario EL  
193 Elemental Analyzer, Elementar). Phosphorus content (P in g kg<sup>-1</sup>) was measured by the Olsen  
194 method (Olsen 1954). Calcium (Ca in g kg<sup>-1</sup>), magnesium (Mg in g kg<sup>-1</sup>) and potassium (K in  
195 g kg<sup>-1</sup>) were determined by the Metson method (Metson 1957). Additionally, soil pH was



196 measured in a 1:1 soil-water solution, after stirring once followed by a 24 h rest period. Carbon-  
197 nitrogen ratio (C:N) and organic matter (OM in g kg<sup>-1</sup>) were calculated using total C and N  
198 results. The amount of clay (> 0.002 mm), fine silt (0.002–0.02 mm), coarse silt (0.02–  
199 0.05 mm), fine sand (0.05–0.2 mm) and coarse sand (0.2–2 mm) in g kg<sup>-1</sup> were determined  
200 according to the Robinson method (Robinson 1922) without previous decarbonation (Baize  
201 2000).

### 202 ***Comparison of vegetation composition between 2001 and 2015***

203 To investigate whether the vegetation composition of the fallow land had re-developed  
204 towards the vegetation of unaltered steppes over the previous 14 years, we compared old  
205 vegetation relevés from Römermann et al (2004, 2005) to recent vegetation relevés for the  
206 locations Peau de Meau (steppe, cereal fallow, melon fallow; Table 1), Petit Abondoux (steppe,  
207 cereal fallow; Table 1) and Valigne (steppe, melon fallow; Table 1). Ten replicate relevés were  
208 carried out per site in 2001 and five in 2015.

### 209 ***Trait data and community-weighted means (CWM)***

210 We compiled trait data for SLA (the ratio of fresh leaf area to dry mass in g cm<sup>-2</sup>), LDMC  
211 (the ratio of dry mass to fresh mass in mg g<sup>-1</sup>) and canopy height (cm) for all species observed  
212 in the vegetation relevés using the trait databases TRY (Kattge et al. 2011) and BROT (Paula  
213 & Pausas 2013). This dataset was then supplemented with our own measurements for species  
214 missing from the databases. For our own measurements, five individuals per species were  
215 collected at the site where they were most abundant, following the approach described in  
216 Rosbakh et al. (2015). Plants were collected and measured from May to July 2015, following  
217 standardized protocols (Pérez-Harguindeguy et al. 2013). For each species, two healthy sun  
218 leaves were collected, wrapped in moist paper and stored in a plastic bag in a cool box until the  
219 fresh leaf weight was measured. The fresh leaf area was determined at 300 dpi on a flatbed  
220 scanner. Leaf area was measured on scanned fresh leaves with the Leaf Trait package for R

221 (Bernhardt-Römermann, unpubl.). Then, the leaves were dried at 70° C to constant mass and  
 222 subsequently weighed. For each vegetation relevé, we then calculated the CWM for SLA,  
 223 LDMC and canopy height according to:

Formula 1: 
$$CWM_c = \sum_{i=1}^s p_{ic} \times T_{ic}$$

224 where the mean value of trait T for species *i* in the community *c* was weighted by species  
 225 relative abundance  $p_{ic}$  in the community, with a total of *S* species (Garnier et al. 2004; Violle et  
 226 al. 2007).

### 227 ***Species selection and trait measurements for intraspecific trait variability (ITV)***

228 Species selection was based on vegetation relevés from 2001 (Römermann et al. 2004, 2005)  
 229 and 2014 (Tanet 2014). The final selection was based on indicator species analysis conducted  
 230 with the package ‘indicspecies’ (Cáceres & Legendre 2009) supplemented by expert knowledge.  
 231 We identified three successful colonizers as those occurring originally in the steppe but also in  
 232 the fallow land (*Carthamus lanatus* L., *Lobularia maritima* L. Desv., *Sideritis romana* L.). Five  
 233 unsuccessful colonizers were limited to the steppe (*B. retusum*, *Hypochaeris glabra* L., *Linum*  
 234 *trigynum* L., *Stipa capillata* L., *Trifolium campestre* Schreb.). Four species whose major  
 235 distribution was in fallow land were added for comparison (*Crepis foetida* L., *Dactylis*  
 236 *glomerata* L., *Galactites tomentosus* Moench., *Trifolium subterraneum* L.). We sampled a  
 237 maximum of eight individuals per species and site if abundant: for details, see Table 2 and  
 238 Table 3. Besides SLA, LDMC and canopy height, we expanded our trait set to include plant  
 239 width (cm), aboveground biomass (g), seed mass (mg) and leaf carbon (leaf C in mg kg<sup>-1</sup>),  
 240 nitrogen (leaf N in mg kg<sup>-1</sup>) and phosphorus content (leaf P in mg kg<sup>-1</sup>; Table 3). Plant width was  
 241 measured with a measuring tape at maximum diameter without extending the plant, biomass  
 242 was dried at 70° C to constant mass and subsequently weighed. We determined average seed  
 243 mass by weighing five batches of 20–100 ripe seeds per species, depending on the species. Seed

244 sampling was not possible for *B. retusum*, *H. glabra* and *S. capillata* as seed production was  
245 not sufficient in 2015. Due to insufficient plant material, leaf samples of the eight individuals  
246 were pooled per site prior to nutrients analyses. Thus, ITV could not be calculated thereof but  
247 leaf nutrients could be compared between the different land-use types (see Appendix S3).

248 To characterize the differences in trait composition between individuals measured in the  
249 steppe and the fallow land sites, a principle component analysis (PCA) was conducted (matrix:  
250 854 individuals of 9 species x 9 traits; *H. glabra*, *B. retusum*, *T. subterraneum* were excluded  
251 because of data constraints – insufficient biomass and seeds available). Prior to analyses, data  
252 was standardized to zero mean and unit variance.

253 To characterize ITV, the coefficient of variation (CV) was calculated as the standard  
254 deviation divided by the mean for all measured traits (Pérez-Harguindeguy et al. 2013) based  
255 on the eight individuals per species and site. ITV was calculated for all species occurring in the  
256 steppe sites to assess whether species with higher trait variability were more likely to pass  
257 through the environmental filter and thus manage to re-establish in fallow land. Therefore, we  
258 included four unsuccessful colonizers three successful colonizers and two fallow land species  
259 in our CV calculation (Table 2).

## 260 ***Statistical analyses***

### 261 *Vegetation composition and soil properties*

262 To test for the effect of land-use history on vegetation structure (S, H<sub>s</sub>, E, standing crop,  
263 LAI, vegetation cover/stone cover/bare ground) and soil properties (clay, silt, sand, organic  
264 matter, total C, total N, C:N, pH, P, Ca, Mg and K), one-way ANOVA followed by a Tukey  
265 Honest Significant Difference post-hoc test was performed for normally distributed data with  
266 homogeneous variances. Any data not showing normal distribution and homogeneous variances  
267 were analyzed by the nonparametric Kruskal–Wallis test followed by the pairwise Wilcoxon  
268 test with *P*-value adjustment according to Holm.

269 To assess associations between differences in vegetation composition and environmental  
270 parameters, a detrended correspondence analysis (DCA) was conducted. We used species  
271 abundances of the five vegetation relevés per site as the main matrix (a total of 12 species x 16  
272 sites). Then, environmental parameters were correlated with the axes using mean values  
273 calculated per site for each parameter. The environmental factors considered were five physical  
274 soil variables (clays, fine and coarse silt, fine and coarse sand) and eight chemical soil variables  
275 (total C, total N, C:N, pH, P, Ca, Mg and K) as well as percentage cover of vegetation, stones  
276 and bare ground and time since abandonment as deduced from the fallow land cultivation period  
277 (Table 1). All parameters were standardized to zero mean unit variance prior to analyses.

#### 278 *Comparison of vegetation composition between 2001 and 2015*

279 As the 2001 data were obtained on 2m x 2m plots via the frequency method (Pfadenhauer et  
280 al. 1986), we harmonized the data for 2001 and 2015 into presence/ absence data to avoid any  
281 observer effects in the estimation of the frequencies and for the non-metric multidimensional  
282 scaling (NMDS). Species that occurred just once across the overall dataset were removed to  
283 avoid discriminating the dataset too strongly by their presence alone instead of discriminating  
284 the gradient by their frequencies.

#### 285 *Differences in trait composition: Community-weighted means (CWM) and intraspecific trait* 286 *variability (ITV)*

287 To analyze differences between CWM of the three land-use types and CVs, a one-way  
288 ANOVA followed by a Tukey Honest Significant Difference post-hoc test was performed for  
289 normally distributed data with homogeneous variances. Any data not showing normal  
290 distribution and homogeneous variances were analyzed by the nonparametric Kruskal–Wallis  
291 test followed by the pairwise Wilcoxon test with *P*-value adjustment according to Holm.

292 All statistical analyses were carried out in *R* (version 3.3.1, R Development Core Team 2015,  
293 Vienna, AT). Multivariate analyses (DCA, PCA, NMDS) were performed using the *R* packages  
294 ‘vegan’ (Oksanen et al. 2008) and ‘MASS’ (Venables & Ripley 2002).

## 295        **Results**

### 296        *Vegetation and soil attributes*

297        Vegetation structure and soil properties of fallow lands differed from those of adjacent  
298        steppe (Table). The steppe showed highest values in species richness, Shannon diversity and  
299        Evenness; cereal fallows typically had intermediate positions (Table 4). Melon fallows had  
300        significantly higher values in soil Ca, P and pH. The C/N ration was highest in steppes.

301        The DCA (Fig. 2) showed a trend towards very homogeneous vegetation composition in the  
302        steppe plots at different sites, whereas the fallow lands tended to be more heterogeneous in  
303        vegetation composition. The first axis separated steppe from fallow land and was negatively  
304        correlated with total soil C, vegetation cover and soil C:N ratio. Percentages of bare ground,  
305        clays, pH and P were positively correlated with fallow land.

### 306        *Comparison of vegetation composition between 2001 and 2015*

307        In 2001, steppe and fallow lands had distinct species composition trends indicated by the  
308        separated polygons of the NMDS (Fig. 3). Fourteen years later, the fallow land had developed  
309        towards the original steppe vegetation, while there had been little change in steppe vegetation  
310        composition. The steppe communities of both 2001 and 2015 were characterized by perennial  
311        species, such as the dominant steppe species *B. retusum* and *S. capillata* and the annual species  
312        *Aira cupaniana* Guss. and *Linum strictum* L.. Cereal fallows in 2001 commonly contained  
313        *Euphorbia falcata* L. and *Reseda phyteuma* L., whereas *Medicago monspeliaca* L. and *Poa*  
314        *bulbosa* L. were identified as late colonizers of these cereal fallows. Melon fallows in 2001  
315        were characterized by *Bromus rubens* L., *Bromus intermedius* L., *Hordeum murinum* L. and  
316        *Rostraria cristata* (L.) Tzvelev. In 2015, these species had declined and *Plantago lagopus* L.  
317        and *Catapodium rigidum* (L.) C.E.Hubb. had become the typical melon fallow species. For  
318        details on species distribution see Fig. 3.

### 319 ***Differences in trait composition***

320 As shown in Table 4, CWM of SLA was significantly lower in steppe than in fallow-land  
321 communities ( $\chi^2 = 54.19, P < 0.001$ ) while CWM of LDMC was significantly higher in steppe  
322 communities ( $F_{2,77} = 52.38, P < 0.001$ ). Canopy height differed among all the land-use types.  
323 The highest values were observed in the steppe and the lowest in cereal fallows ( $F_{2,77} = 5.19,$   
324  $P < 0.01$ ). For individual species information see Appendix S1.

325 The PCA (Appendix S2) partly separated the species according to their trait composition  
326 measured in the steppe and the fallow land. *C. foetida*, *D. glomerata*, *L. trigynum* and  
327 *S. romana* showed a tendency towards a distinction between the steppe and fallow lands and  
328 thus changes in plant trait values due to land-use change, as indicated by the black arrow.

329 The CV did not reveal any differences between successful and unsuccessful colonizers (Fig.  
330 4). Overall, significant differences between species were found for SLA ( $\chi^2 = 27.44, P > 0.001$ ),  
331 LDMC ( $\chi^2 = 31.75, P > 0.001$ ) and plant width ( $\chi^2 = 36.91, P > 0.001$ ). *L. trigynum*, as a typical  
332 steppe species, varied more in SLA than the fallow land species *D. glomerata* and the successful  
333 colonizer *S. romana*. Regarding LDMC values, *L. trigynum* displayed the highest variability,  
334 while *B. retusum* and *D. glomerata* had significantly lower variability. *B. retusum* had a  
335 significantly lower CV in plant width than *L. trigynum* and *S. romana*. *S. romana* showed  
336 significantly higher variation in plant width than *C. lanatus* (successful colonizer).

### 337 **Discussion**

338 The recovery of fallow land in our study area implemented grazing and was thus not a strictly  
339 passive restoration technique. The fact that more than 35 years after cultivation has ceased the  
340 studied vegetation did not fully transform to steppe calls for more active intervention, like  
341 seeding and planting. Notwithstanding these results, the comparison of vegetation relevés from  
342 2001 with results for 2015 indicated a (though slow) conversion of fallow land towards steppe  
343 vegetation. Furthermore, there was no particular similarity in trait composition between steppe

344 species that successfully re-colonized fallow lands and the fallow-land community, and no  
345 evidence that intraspecific trait variability is associated with re-colonization success of these  
346 species.

### 347 ***Effect of former cultivation on vegetation attributes and soil parameters***

348 Former cultivation practices still influence present vegetation attributes. The species  
349 composition of recently abandoned cereal fallows can be distinguished from the species  
350 composition of the steppe. However, in the long-abandoned cereal fallow, there is a visible  
351 vegetation development towards the original steppe community.

352 Our results show that the dominant perennial steppe grass *B. retusum* has so far been unable  
353 to re-establish on the fallow land; this is not only directly explained by habitat change due to  
354 deep soil ploughing and high fertilization, but also indirectly by competition with the common  
355 fallow land species (Buisson et al. 2015; Coiffait et al. 2012). Moreover, dispersal limitation  
356 and the mainly vegetative reproduction of this steppe grass (Dureau & Bonnefon 1998) have  
357 prevented successful re-establishment. The decline of C:N ratios on the fallow land can be  
358 explained by organic matter mineralization and the absence of restitution from perennial grass  
359 species unable to re-colonize fallow land (Römermann et al. 2004, 2005).

360 The high percentage of bare ground is characteristic of fallow land, where stones were  
361 removed for cultivation purposes (Buisson & Dutoit 2006). However, stones create important  
362 micro-habitats for the typical steppe vegetation: they preserve humidity, serve as shelter and  
363 create safe sites for rhizomes of e.g., *B. retusum* (Bourrelly 1984; Caturla et al. 2000; Buisson  
364 et al. 2015).

365 According to Janssens et al. (1998), plant diversity is limited by high soil fertility, especially  
366 by plant-available phosphorus. The fallow land remains distinguished from the steppe by its  
367 elevated phosphorus and potassium content due to former fertilization, as also observed by  
368 Gough and Marrs (1990) on abandoned fields in the UK. Because of different cultivation regime  
369 intensities, i.e., deep ploughing, fertilization, the application of pesticides in melon-cultivated

370 areas (Masip 1991) versus less intensive cereal cultivation, phosphorus content was higher in  
371 melon fallows than in cereal fallows. Deep ploughing for melon cultivation destroyed the upper  
372 part of the geological conglomerate 40 cm deep in the soil, bringing some pieces of the matrix  
373 (composed of  $\text{CaCO}_3$ ) to the surface. Hence, Ca content increased and higher pH was measured.  
374 The effect of ploughing also appears in the physical soil parameters. As a result of soil  
375 disturbance through ploughing, clays from the steppe subsoil layer (20–40 cm) were transported  
376 to the topsoil, leading to a higher clay content for the fallow land (10–20 cm; Duclos 1994).  
377 The opposite effect is observed for the coarse silt of the steppe, which was buried in the subsoil  
378 as a result of ploughing (Bulot et al. 2017; Chenot et al. 2017).

379 Öster et al. (2009) found that in semi-natural grassland communities in Sweden, species  
380 richness and similarity were still lower in fallow lands more than 50 years after cultivation.  
381 Differences in plant attributes between calcareous grassland and fallow land after 60–100 years  
382 were also shown by Fagan et al. (2008), Karlik & Poschlod (2009) and Forey and Dutoit (2012).  
383 Remnants of fertilization persist in the soil of fallow land for many years and still affect plant  
384 communities and the vegetation structure decades after abandonment (Willems et al. 1993;  
385 Austrheim & Olsson 1999; Lepš 1999).

### 386 ***Comparison of vegetation structure between 2001 and 2015***

387 The resurvey of the research site in 2015 shows a trajectory of the former cultivated areas  
388 towards the original steppe community, even though differences in vegetation and soil structure  
389 still exist. Although common in 2001, ruderal species *H. murinum* and *R. cristata* were not  
390 recorded on melon fallows in 2015. Similarly, *Bromus* species- typical species in the first year  
391 of abandonment- were less common on fallow land in 2015. In cereal fallows, not only weed  
392 species but also some typical steppe species had established by 2015. The steppe vegetation did  
393 not change over time because the disturbance regime (i.e., extensive grazing) remained stable  
394 between 2001 and 2015 (Sauguet 2013). The more rapid development from cereal fallow  
395 vegetation towards steppe vegetation was probably due to the fact that less fertilizer was applied



396 during cultivation and that there was no deep ploughing (Römermann et al. 2005, Buisson et al.  
397 2006).

### 398 *Effect of former cultivation on plant functional traits*

399 This study confirms the relevance of the leaf traits SLA and LDMC at community level to  
400 explain the response of ecosystems to land-use change, as proposed by previous studies (Díaz  
401 et al. 2007; de Bello et al. 2010; Vandewalle et al. 2010). The steppe community was  
402 characterized by smaller and denser leaves, typical of plants following a conservative strategy  
403 (Grime 1973; Saatkamp et al. 2010; Pérez-Harguindeguy et al. 2013). Our study demonstrates  
404 that the fallow land is still a nutrient-rich environment favoring communities with high SLA  
405 and low LDMC. This is in line with findings demonstrating the association of low SLA with  
406 efficient nutrient conservation strategies, nutrient-poor habitats, slow growth rates and high  
407 longevity, as outlined above (Poorter & Remkes 1990; Garnier 1992; Chapin et al. 1993;  
408 Poorter & Garnier 1999; Lavorel & Garnier 2002; Pérez-Harguindeguy et al. 2013). Our  
409 analyses showing that the cereal fallows occupy an intermediate position with respect to trait  
410 composition also reveal that type of cultivation plays an important role in re-development  
411 towards the unaltered steppe vegetation. Plant height was greater in steppe communities than  
412 in fallow land, contradicting our initial hypothesis. However, it is in line with findings from  
413 Australian ecosystems (Falster & Westoby 2003) that showed positive relationships between  
414 plant height and LAI. Our finding of higher plants in the steppe may be related to the higher  
415 fodder quality in nutrient-rich fallow land, which could lead to preferential grazing by  
416 herbivores, reducing plant height (Falster & Westoby 2003; Saatkamp et al. 2010). This  
417 differential selection by herbivores of recently impacted sites due to their higher fodder quality  
418 could also have implications for restoration (Yates et al. 2000). Tests of plant and soil mineral  
419 content of our data revealed a significantly positive correlation between soil phosphorus and  
420 leaf phosphorus content ( $r = 0.67$ ,  $P < 0.001$ ), soil nitrogen and leaf nitrogen content ( $r = 0.35$ ,  
421  $P < 0.02$ , analyses not shown).

422 We found no evidence of particular similarity between the trait values of successful  
423 colonizers and those of the species typically found in fallow-land communities. We thus have  
424 to reject the idea that colonizers simply “fit into” the assembly of recent fallow-land  
425 communities with initial assembly conditions of high nutrient levels and recently-moved soil.  
426 Alternatively, the data can be understood as indicating that successful colonizers exploit the  
427 diverging conditions created during longer grazing periods with decreased nutrient status, and  
428 that these colonizers may be better adapted to grazing than typical fallow-land species  
429 (Saatkamp et al. 2010). However, our species pool was quite small with differences in sample  
430 size, and further studies should consider a more extensive species set covering more families.

431 Furthermore, our data show higher functional variability to be highly species- and trait-  
432 specific, with no clear pattern revealed for the trait values of successful or unsuccessful  
433 colonizers. This supports the view that plants in fallow lands are assembled according to their  
434 adaptability to specific grazed fallow-land conditions, rather than reflecting a wide intraspecific  
435 trait variability that ensures their survival in both fallow-land and steppe communities. Hence,  
436 our study suggests that plants with specifically high ITV are not favored in the assembly process  
437 after abandonment and during grazing.

### 438 ***Conclusions and implications for management***

439 Our study demonstrates that natural recovery of abandoned fallow land, notwithstanding its  
440 close proximity to the target vegetation, can take more than 35 years, even when the traditional  
441 grazing system is re-introduced just after cultivation abandonment. Regular monitoring of  
442 further vegetation development in the abandoned fallow land would enable the potential for  
443 natural recovery after a period of cultural land-use to be assessed.

444 In addition to monitoring natural recovery over the long term, new grazing systems could be  
445 introduced to hasten recovery or more active restoration implemented by seeding nurse or  
446 structuring species (e.g. *B. retusum*; Coiffait-Gombault et al., 2012; Buisson et al., 2015).  
447 Future analyses could cover an extended set of species and include further traits to better

448 elucidate assembly processes. In particular, they should consider traits closely related to soil  
449 water conditions, such as leaf venation traits and water-use efficiency, and traits related to soil  
450 stoichiometry, such as root traits, mycorrhizal type (Jaunatre et al. 2016) and seed nutrient  
451 content. Future projects need to identify project-specific restoration targets in order to  
452 implement a trait-based approach into their restoration protocol.

## 453 **Acknowledgments**

454 We would like to thank the *Conservatoire d'espaces naturels de Provence-Alpes-Côte d'Azur*  
455 (CEN PACA) and the *Réserve Naturelle Nationale des Coussouls de Crau* for their support  
456 during field work. Thanks to Julie Tanet and Adeline Bulot working at the Mediterranean  
457 Institute of Biodiversity and Ecology of Marseille for realizing the botanical relevés in 2014.  
458 We acknowledge valuable comments of Karen Russel, Viktoria Wagner, Anna Johnson and an  
459 anonymous referee. This study was supported by the TRY initiative on plant traits  
460 (<http://www.trydb.org>) hosted, developed and maintained by J. Kattge and G. Bönisch (Max-  
461 Planck-Institute for Biogeochemistry, Jena, Germany). TRY is/was supported by  
462 DIVERSITAS, IGBP, the Global Land Project, the UK Natural Environment Research Council  
463 (NERC) through its program QUEST (Quantifying and Understanding the Earth System), the  
464 French Foundation for Biodiversity Research (FRB), and GIS "Climat, Environnement et  
465 Société" France. CR acknowledges financial support by the German Research Foundation  
466 (project RO 3842/3-1). The authors declare no conflict of interest.

467

## 468 **References**

- 469 Albert, C.H., Grassein, F., Schurr, F.M., Vieilledent, G. & Violle, C. 2011. When and how should  
470 intraspecific variability be considered in trait-based plant ecology? *Perspectives in Plant*  
471 *Ecology, Evolution and Systematics* 13: 217-225.
- 472 Austrheim, G. & Olsson, E.G.A. 1999. How does continuity in grassland management after ploughing  
473 affect plant community patterns? *Plant Ecology* 145: 59-74.

- 474 Baize, D. 2000. *Guide des analyses en pédologie: 2e édition, revue et augmentée*. Editions Quae, FR.
- 475 Bucher, S.F., König, P., Menzel, A., Migliavacca, M., Ewald, J. & Römermann, C. 2018. Traits and  
476 climate are associated with first flowering day in herbaceous species along elevational gradients.  
477 *Ecology and Evolution* 8: 1147-1158.
- 478 Bourrelly, M. 1984. *Contribution à l'étude écologique de la Crau (Bouches-du-Rhône). Caractéristiques  
479 floristiques, dynamique annuelle, production primaire de la Crau steppique*. Ph.D. thesis, Aix-  
480 Marseille University, Marseille, FR.
- 481 Braun-Blanquet, J. 1964. *Pflanzensoziologie : Grundzüge der Vegetationskunde*, Springer-Verlag, Wien.  
482 AT.
- 483 Breitschwerdt, E., Jandt, U. & Bruelheide, H. 2015. Do newcomers stick to the rules of the residents?  
484 Designing trait-based community assembly tests. *Journal of Vegetation Science* 26: 219-232.
- 485 Bucher, S. F., Auerswald, K., Grün-Wenzel, C., Higgins, S. I., Garcia Jorge, J. & Römermann, C. 2017.  
486 Stomatal traits relate to habitat preferences of herbaceous species in a temperate climate. *Flora*  
487 229: 107-115.
- 488 Buisson, E., Corcket, E. & Dutoit, T. 2015. Limiting processes for perennial plant reintroduction to  
489 restore dry grasslands. *Restoration Ecology* 23: 947-954.
- 490 Buisson, E. & Dutoit, T. 2004. Colonisation by native species of abandoned farmland adjacent to a  
491 remnant patch of Mediterranean steppe. *Plant Ecology* 174: 371-384.
- 492 Buisson, E. & Dutoit, T. 2006. Creation of the natural reserve of La Crau: implications for the creation  
493 and management of protected areas. *Journal of Environmental Management* 80: 318-326.
- 494 Buisson, E., Dutoit, T., Torre, F., Römermann, C. & Poschlod, P. 2006. The implications of seed rain  
495 and seed bank patterns for plant succession at the edges of abandoned fields in Mediterranean  
496 landscapes. *Agriculture, Ecosystems & Environment* 115: 6-14.
- 497 Bulot, A., Potard, K., Bureau, F., Bérard, A. & Dutoit, T. 2017. Ecological restoration by soil transfer:  
498 impacts on restored soil profiles and topsoil functions. *Restoration Ecology* 25: 354-366.
- 499 Cáceres, M.D. & Legendre, P. 2009. Associations between species and groups of sites: indices and  
500 statistical inference. *Ecology* 90: 3566-3574.
- 501 Caturla, R.N., Raventós, J., Guàrdia, R. & Vallejo, V.R. 2000. Early post-fire regeneration dynamics of  
502 *Brachypodium retusum* Pers.(Beauv.) in old fields of the Valencia region (eastern Spain). *Acta*  
503 *Oecologica* 21: 1-12.
- 504 Chapin, F.S., Autumn, K. & Pugnaire, F. 1993. Evolution of suites of traits in response to environmental  
505 stress. *American Naturalist*: S78-S92.
- 506 Chenot, J., Jaunatre, R., Buisson, E. & Dutoit, T. 2017. Long-term effects of topsoil transfer assessed  
507 thirty years after rehabilitation of dry alluvial quarries in Southeastern France. *Ecological*  
508 *Engineering* 99: 1-12.
- 509 Coiffait, C., Buisson, E. & Dutoit, T. 2012. Using a two-phase sowing approach in restoration – sowing  
510 foundation species to restore and subordinate species to evaluate restoration success. *Applied*  
511 *Vegetation Science* 15: 277-289.
- 512 Colomb, E. & Roux, R. 1978. La Crau, données nouvelles et interprétations. *Géologie méditerranéenne*  
513 5: 303-324.
- 514 Cornwell, W.K. & Ackerly, D.D. 2009. Community assembly and shifts in plant trait distributions across  
515 an environmental gradient in coastal California. *Ecological Monographs* 79: 109-126.

- 516 de Bello, F., Lavorel, S., Díaz, S., Harrington, R., Cornelissen, J.H., Bardgett, R.D., Berg, M.P., Cipriotti,  
517 P., Feld, C.K. & Hering, D. 2010. Towards an assessment of multiple ecosystem processes and  
518 services via functional traits. *Biodiversity and Conservation* 19: 2873-2893.
- 519 Devaux, J., Archiloque, A., Borel, L., Bourrelly, M. & Louis-Palluel, J. 1983. Notice de la carte phyto-  
520 écologique de la Crau (Bouches du Rhône). *Biologie et écologie méditerranéenne* 10: 5-54.
- 521 Deverre, C. 1996. La nature mise au propre dans la steppe de Crau et la forêt du Var. *Études rurales*  
522 141-142: 45-61.
- 523 Diaz, S., Cabido, M. & Casanoves, F. 1998. Plant functional traits and environmental filters at a regional  
524 scale. *Journal of Vegetation Science* 9: 113-122.
- 525 Dureau, R. & Bonnefon, O. 1998. Etude des pratiques de gestion pastorale des coussouls. *Patrimoine*  
526 *nature et pratiques pastorales en Crau*. Ed. CEEP Ecomusée de Crau: 61-89.
- 527 Dutoit, T., Buisson, E., Fadda, S., Henry, F., Coiffait-Gombault, C., Jaunatre, R., Alignan, J.F., Masson,  
528 S. & Bulot, A. 2013. The pseudo-steppe of La Crau (South-Eastern France): origin, management  
529 and restoration of a Mediterranean rangeland. In: Traba J. Morales M (eds.) *Steppe Ecosystems:*  
530 *Biological Diversity, Management and Restoration*, pp. 287-301. Nova Science Publishers,  
531 Hauppauge, New York, US.
- 532 Engst, K., Baasch, A., Erfmeier, A., Jandt, U., May, K., Schmiede, R., & Bruelheide, H. 2016.  
533 Functional community ecology meets restoration ecology: Assessing the restoration success of  
534 alluvial floodplain meadows with functional traits. *Journal of Applied Ecology* 53: 751-764.
- 535 Etienne, M., Aronson, J. & Le Floc'h, E. 1998. Abandoned lands and land use conflicts in southern  
536 France. Piloting ecosystem trajectories and redesigning outmoded landscapes in the 21st century.  
537 In: Rundel P.W., Montenegro G. and Jaksic F.M. (eds.) *Landscape disturbance and biodiversity*  
538 *in Mediterranean-type ecosystems*, pp. 127-140. Ecological Studies, Springer-Verlag, Berlin,  
539 DE.
- 540 Fagan, K.C., Pywell, R.F., Bullock, J.M. & Marrs, R.H. 2008. Do restored calcareous grasslands on  
541 former arable fields resemble ancient targets? The effect of time, methods and environment on  
542 outcomes. *Journal of Applied Ecology* 45: 1293-1303.
- 543 Falster, D.S. & Westoby, M. 2003. Plant height and evolutionary games. *Trends in Ecology & Evolution*  
544 18: 337-343.
- 545 Fischer, S.F., Poschlod, P. & Beinlich, B. 1996. Experimental studies on the dispersal of plants and  
546 animals on sheep in calcareous grasslands. *Journal of Applied Ecology*: 1206-1222.
- 547 Forey, E. & Dutoit, T. 2012. Vegetation, soils and seed banks of limestone grasslands are still impacted  
548 by former cultivation one century after abandonment. *Community Ecology* 13: 194-202.
- 549 Garnier, E. 1992. Growth analysis of congeneric annual and perennial grass species. *Journal of Ecology*  
550 80: 665-675.
- 551 Garnier, E., Cortez, J., Billès, G., Navas, M.-L., Roumet, C., Debussche, M., Laurent, G., Blanchard,  
552 A., Aubry, D. & Bellmann, A. 2004. Plant functional markers capture ecosystem properties  
553 during secondary succession. *Ecology* 85: 2630-2637.
- 554 Gaudet, C.L. & Keddy, P.A. 1988. A comparative approach to predicting competitive ability from plant  
555 traits. *Nature* 334: 242-243.
- 556 Gough, M. & Marrs, R. 1990. A comparison of soil fertility between semi-natural and agricultural plant  
557 communities: Implications for the creations of species-rich grassland on abandoned agricultural  
558 land. *Biological Conservation* 51: 83-96.

- 559 Grime, J.P. 1973. Competitive exclusion in herbaceous vegetation. *Nature, UK* 242: 344-347.
- 560 Jäger, E.J. & Werner, K. 2002. Rothmaler, Exkursionsflora von Deutschland. *Gefäßpflanzen: Kritischer*  
561 *Band 9*. Springer Spektrum, Berlin Heidelberg, DE.
- 562 Janssens, F., Peeters, A., Tallowin, J., Bakker, J., Bekker, R., Fillat, F. & Oomes, M. 1998. Relationship  
563 between soil chemical factors and grassland diversity. *Plant and Soil* 202: 69-78.
- 564 Jaunatre, R., Fonvieille, N., Spiegelberger, T., Buisson, E., & Dutoit T. 2016. Recovery of arbuscular  
565 mycorrhizal fungi root colonization after severe anthropogenic disturbance: four species  
566 assessed in old-growth Mediterranean grassland. *Folia Geobotanica* 51: 319-332.
- 567 Jung, V., Violle, C., Mondy, C., Hoffmann, L. & Muller, S. 2010. Intraspecific variability and trait-based  
568 community assembly. *Journal of Ecology* 98: 1134-1140.
- 569 Karlik, P. & Poschod, P. 2009. History or abiotic filter: which is more important in determining the  
570 species composition of calcareous grasslands? *Preslia* 81: 321-340.
- 571 Kattge, J., Diaz, S., Lavorel, S., Prentice, I., Leadley, P., Bönlisch, G., Garnier, E., Westoby, M., Reich,  
572 P.B., (...) & Wright, I. 2011. TRY—a global database of plant traits. *Global Change Biology* 17:  
573 2905-2935.
- 574 Keddy, P.A. 1992. Assembly and response rules: two goals for predictive community ecology. *Journal*  
575 *of Vegetation Science* 3: 157-164.
- 576 Kiehl, K., Kirmer, A., Donath, T. W., Rasran, L., & Hölzel, N. 2010. Species introduction in restoration  
577 projects—Evaluation of different techniques for the establishment of semi-natural grasslands in  
578 Central and Northwestern Europe. *Basic and Applied Ecology* 11: 285-299.
- 579 Kuznetsova, A., Brockhoff, P.B. & Christensen, R.H.B. 2015. *lmerTest: Tests in Linear Mixed Effects*  
580 *Models*. R package version 2.0-29.
- 581 Laforest-Lapointe, I., Martínez-Vilalta, J. & Retana, J. 2014. Intraspecific variability in functional traits  
582 matters: case study of Scots pine. *Oecologia* 175: 1337-1348.
- 583 Laughlin, D.C. 2014. Applying trait-based models to achieve functional targets for theory-driven  
584 ecological restoration. *Ecology Letters* 17: 771-784.
- 585 Laughlin, D.C., Joshi, C., Bodegom, P.M., Bastow, Z.A. & Fulé, P.Z. 2012. A predictive model of  
586 community assembly that incorporates intraspecific trait variation. *Ecology Letters* 15: 1291-  
587 1299.
- 588 Lauterbach, D., Römermann, C., Jeltsch, F. & Ristow, M. 2013. Factors driving plant rarity in dry  
589 grasslands on different spatial scales: a functional trait approach. *Biodiversity and Conservation*  
590 22: 2337-2353.
- 591 Lavorel, S. & Garnier, E. 2002. Predicting changes in community composition and ecosystem  
592 functioning from plant traits: revisiting the Holy Grail. *Functional ecology* 16: 545-556.
- 593 Lenth, R.V. 2016. *Ismeans: Least-Squares Means*. R Package version 2.22.
- 594 Lepš, J. 1999. Nutrient status, disturbance and competition: an experimental test of relationships in a  
595 wet meadow. *Journal of Vegetation Science* 10: 219-230.
- 596 MacArthur, R. & Levins, R. 1967. The limiting similarity, convergence, and divergence of coexisting  
597 species. *American Naturalist* 101: 377-385.
- 598 Masip, A. 1991. *Le peuplement végétal de la Réserve de Peau de Meau. Données pour la gestion*. Ph.D.  
599 thesis, University of Barcelona. Barcelona, ES.

- 600 Metson, A.J. 1957. Methods of chemical analysis for soil survey samples. *Soil Science* 83: 245.
- 601 Molliex, S., Siame, L.L., Bourlès, D.L., Bellier, O., Braucher, R. & Clauzon, G. 2013. Quaternary  
602 evolution of a large alluvial fan in a periglacial setting (Crau Plain, SE France) constrained by  
603 terrestrial cosmogenic nuclide (<sup>10</sup>Be). *Geomorphology* 195: 45-52.
- 604 Mühlenberg, M. 1993. *Freilandökologie*. UTB für Wissenschaft, 3<sup>rd</sup> ed. Quelle & Meyer, Heidelberg,  
605 Wiesbaden, DE.
- 606 Nicotra, A.B., Atkin, O.K., Bonser, S.P., Davidson, A.M., Finnegan, E., Mathesius, U., Poot, P.,  
607 Purugganan, M.D., Richards, C. & Valladares, F. 2010. Plant phenotypic plasticity in a changing  
608 climate. *Trends in Plant Science* 15: 684-692.
- 609 Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, M.H.H., Oksanen, M.J. & Suggests, M. 2007.  
610 *Vegan: Community ecology package*. R package version 2.0-10.
- 611 Olsen, S.R. 1954. *Estimation of available phosphorus in soils by extraction with sodium bicarbonate*.  
612 United States Department of Agriculture, Washington, DC, US.
- 613 Öster, M., Ask, K., Cousins, S.A. & Eriksson, O. 2009. Dispersal and establishment limitation reduces  
614 the potential for successful restoration of semi-natural grassland communities on former arable  
615 fields. *Journal of Applied Ecology* 46: 1266-1274.
- 616 Paula, S. & Pausas, J. 2013. *BROT: a plant trait database for Mediterranean Basin species*. Version  
617 2013.06. [http:// www.uv.es/jgpausas/brot.htm](http://www.uv.es/jgpausas/brot.htm).
- 618 Pavon, D. 2014. Mémento pour l'aide à la détermination de la flore vasculaire du département des  
619 Bouches-du-Rhône. *Bulletin de la Société linnéenne de Provence*. Numero spécial 18.
- 620 Pérez-Harguindeguy, N., Díaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., Bret-Harte,  
621 M., Cornwell, W., Craine, J., (...) & Cornelissen, J.H.C. 2013. New handbook for standardised  
622 measurement of plant functional traits worldwide. *Australian Journal of Botany* 61: 167-234.
- 623 Pfadenhauer, J., Poschlod, P. & Buchwald, R. 1986. Überlegungen zu einem Konzept geobotanischer  
624 Dauerbeobachtungsflächen für Bayern. Teil 1. Methodik der Anlage und Aufnahme. *Bayerische  
625 Akademie für Naturschutz und Landschaftspflege* 10: 41-60.
- 626 Poorter, H. & Garnier, E. 1999. Ecological significance of inherent variation in relative growth rate and  
627 its components. *Handbook of functional Plant Ecology* 20: 81-120.
- 628 Poorter, H., Niinemets, Ü., Poorter, L., Wright, I.J. & Villar, R. 2009. Causes and consequences of  
629 variation in leaf mass per area (LMA): a meta-analysis. *New Phytologist* 182: 565-588.
- 630 Poorter, H. & Remkes, C. 1990. Leaf area ratio and net assimilation rate of 24 wild species differing in  
631 relative growth rate. *Oecologia* 83: 553-559.
- 632 Poschlod, P., Kiefer, S., Tränkle, U., Fischer, S. & Bonn, S. 1998. Plant species richness in calcareous  
633 grasslands as affected by dispersability in space and time. *Applied Vegetation Science* 1: 75-91.
- 634 Reich, P.B., Walters, M.B. & Ellsworth, D.S. 1997. From tropics to tundra: global convergence in plant  
635 functioning. *Proceedings of the National Academy of Sciences* 94: 13730-13734.
- 636 Robin, V., Nelle, O., Talon T., Poschlod, P., Schwartz, D., Bal M.C., Allée, P., Vernet, J.L. & Dutoit T.  
637 2018. A comparative review of soil charcoal data: spatiotemporal patterns of origin and long-  
638 term dynamics of Western European nutrient-poor grasslands. *The Holocene (in press)*
- 639 Römermann, C., Bernhardt-Römermann, M., Dutoit, T., Poschlod, P. & Rolando, C. 2004. Histoire  
640 culturale de la Crau: potentialités de ré-établissement des espèces caractéristiques du coussous  
641 après abandon. *Ecologia Mediterranea* 30: 47-70.

- 642 Römermann, C., Dutoit, T., Poschlod, P. & Buisson, E. 2005. Influence of former cultivation on the  
643 unique Mediterranean steppe of France and consequences for conservation management.  
644 *Biological Conservation* 121: 21-33.
- 645 Römermann, C., Bernhardt-Römermann, M., Kleyer, M. & Poschlod, P. 2009. Substitutes for grazing  
646 in semi-natural grasslands – do mowing or mulching represent valuable alternatives to maintain  
647 vegetation structure? *Journal of Vegetation Science* 20: 1086-1098.
- 648 Römermann, C., Bucher, S. F., Hahn, M. & Bernhardt-Römermann, M. 2016. Plant functional traits -  
649 fixed facts or variable depending on the season? *Folia Geobotanica* 51: 143-159.
- 650 Robinson, G. (1922). A new method for the mechanical analysis of soils and other dispersions. *The*  
651 *Journal of Agricultural Science*, 12(3), 306-321.
- 652 Rosbakh, S., Römermann, C. & Poschlod, P. 2015. Specific leaf area correlates with temperature: new  
653 evidence of trait variation at the population, species and community levels. *Alpine Botany* 125:  
654 79-86.
- 655 Saatkamp, A., Römermann, C. & Dutoit, T. 2010. Plant functional traits show non-linear response to  
656 grazing. *Folia Geobotanica*, 45: 239-252.
- 657 Sauguet, F. 2013. L'élevage ovin en Crau: situation économique, enjeux et perspectives. In: *Écologie et*  
658 *conservation d'une steppe méditerranéenne. La plaine de Crau*, pp. 192-205. Editions Quae,  
659 FR.
- 660 Tanet, J. 2014. *Les fourmis sont-elles de bons indicateurs de la régénération spontanée des pelouses*  
661 *sèches méditerranéennes après dégradation culturale?* Master Thesis. Aix-Marseille University,  
662 Marseille, FR.
- 663 Vandewalle, M., De Bello, F., Berg, M.P., Bolger, T., Dolédec, S., Dubs, F., Feld, C.K., Harrington, R.,  
664 Harrison, P.A., (...) & Woodcock, B. A.2010. Functional traits as indicators of biodiversity  
665 response to land use changes across ecosystems and organisms. *Biodiversity and Conservation*  
666 19: 2921-2947.
- 667 Venables, W. & Ripley, B. 2002. *Modern Applied Statistics with S*. 4<sup>th</sup> ed. Springer-Verlag, New York,  
668 USA.
- 669 Venn, S.E., Green, K., Pickering, C.M. & Morgan, J.W. 2011. Using plant functional traits to explain  
670 community composition across a strong environmental filter in Australian alpine snowpatches.  
671 *Plant Ecology* 212: 1491-1499.
- 672 Villar, R., Maranon, T., Quero, J.L., Panadero, P., Arenas, F. & Lambers, H. 2005. Variation in relative  
673 growth rate of 20 *Aegilops* species (Poaceae) in the field: the importance of net assimilation rate  
674 or specific leaf area depends on the time scale. *Plant and Soil* 272: 11-27.
- 675 Violle, C., Navas, M.L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I. & Garnier, E. 2007. Let the  
676 concept of trait be functional! *Oikos* 116: 882-892.
- 677 WallisDeVries, M.F., Poschlod, P. & Willems, J.H. 2002. Challenges for the conservation of calcareous  
678 grasslands in northwestern Europe: integrating the requirements of flora and fauna. *Biological*  
679 *Conservation* 104: 265-273.
- 680 Watson, D.J. 1947. Comparative physiological studies on the growth of field crops: I. Variation in net  
681 assimilation rate and leaf area between species and varieties, and within and between years.  
682 *Annals of Botany* 11: 41-76.
- 683 Weiher, E. & Keddy, P.A. 1995. Assembly rules, null models, and trait dispersion: new questions from  
684 old patterns. *Oikos* 74: 159-164.



- 685 Weiher, E., Werf, A., Thompson, K., Roderick, M., Garnier, E. & Eriksson, O. 1999. Challenging  
686 Theophrastus: a common core list of plant traits for functional ecology. *Journal of Vegetation*  
687 *Science* 10: 609-620.
- 688 Westoby, M. 1998. A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant and Soil* 199: 213-  
689 227.
- 690 Westoby, M., Falster, D.S., Moles, A.T., Vesk, P.A. & Wright, I.J. 2002. Plant ecological strategies:  
691 some leading dimensions of variation between species. *Annual review of Ecology and*  
692 *Systematics* 33: 125-159.
- 693 White, R.P., Murray, S., Rohweder, M., Prince, S. & Thompson, K. 2000. *Grassland ecosystems*. World  
694 Resources Institute, Washington, DC, US.
- 695 Willems, J. 1983. Species composition and above ground phytomass in chalk grassland with different  
696 management. *Vegetatio* 52: 171-180.
- 697 Willems, J., Peet, R.K. & Bik, L. 1993. Changes in chalk-grassland structure and species richness  
698 resulting from selective nutrient additions. *Journal of Vegetation Science* 4: 203-212.
- 699 Wilson, J.B. 2007. Trait-divergence assembly rules have been demonstrated: Limiting similarity lives!  
700 A reply to Grime. *Journal of Vegetation Science* 18: 451-452.
- 701 Yates, C.J., Norton, D.A. & Hobbs, R.J. 2000. Grazing effects on plant cover, soil and microclimate in  
702 fragmented woodlands in south-western Australia: implications for restoration. *Austral Ecology*  
703 25: 36-47.
- 704 Yulin, L., Johnson, D.A., Yongzhong, S., Jianyuan, C. & Zhang, T. 2005. Specific leaf area and leaf dry  
705 matter content of plants growing in sand dunes. *Botanical Bulletin of Academia Sinica* 46: 127-  
706 134.
- 707

## 708 **Supporting Information**

709

710 **Appendix S1.** Trait values of specific leaf area (SLA), leaf dry matter content (LDMC) and canopy  
711 height for all species.

712 **Appendix S2.** Principal component analysis (PCA) of the species–trait matrix (9 species × 9 traits).

713 **Appendix S3.** Leaf nutrient content of carbon (C), nitrogen (N) and phosphorus (P) compared between  
714 original steppe and fallow lands.

715

716

**Table 1.** Overview of the land-use history of the study sites in La Crau studied in 2015. \*Comparison between 2001 and 2015.

Location	Study sites	Land-use type	Abbreviation	Cultivation period
Peau de Meau	steppe	-	PM_CSa*	-
	steppe	-	PM_CSb	-
	steppe	-	PM_CSc	-
	cereal fallow	melons, cereals	PM_MCa*	1974–1977
	cereal fallow	melons, cereals	PM_MCb	1969–1970
	melon fallow	melons	PM_ME*	1965–1968, 1984
Petit Abondoux	steppe	-	PA_CS	-
	cereal fallow*	melons, cereals	PA_MC*	1979–1984
Grand Abondoux	steppe	-	GA_CS	-
	melon fallow	melons	GA_ME	1980–1981
Valigne	steppe	-	VA_CS*	-
	melon fallow	melons	VA_ME*	1978, 1984
Poulagère	steppe	-	PO_CS	-
	old cereal fallow	cereals	PO_OC	< 1900
Couliès	steppe	-	CO_CS	-
	recent cereal fallow	cereals	CO_RC	1994, 2000

\*comparison between 2001 and 2015

**Table 2.** List of the 12 species, classified into unsuccessful colonizers, fallow-land species and successful colonizers, on which trait measurements were performed on the study sites (Table 1) in 2015.


Species group	Species	Abbreviation	Family	Individuals sampled per species
Unsuccessful colonizers	<i>Brachypodium retusum</i> (Pers.) P. Beauv.	Bra_ret	Poaceaea	64
	<i>Hypochaeris glabra</i> L.	Hyp_gla	Asteraceae	15
	<i>Linum trigynum</i> L.	Lin_tri	Linaceae	120
	<i>Stipa capillata</i> L.	Sti_cap	Poaceae	110
	<i>Trifolium campestre</i> Schreb.	Tri_cam	Fabaceae	59
Fallow-land species	<i>Crepis foetida</i> L.	Cre_foe	Asteraceae	128
	<i>Dactylis glomerata</i> L.	Dac_glo	Poaceae	128
	<i>Galactites tomentosus</i> Moench.	Gal_tom	Asteraceae	52
	<i>Trifolium subterraneum</i> L.	Tri_sub	Fabaceae	16
Successful colonizers	<i>Carthamus lanatus</i> L.	Car_lan	Asteraceae	125
	<i>Lobularia maritima</i> L. Desv.	Lob_mar	Brassicaceae	128
	<i>Sideritis romana</i> L.	Sid_rom	Lamiaceae	128

**Table 3.** Plant functional traits measured on 12 focal species (Table 2). \*These plant traits are not part of the main study, but available in Supporting Information (Appendix S2, S3).


Organ	Plant trait	Abbreviation	Unit	Measurements per site
Whole plant	Canopy height		cm	8 individuals
	Plant width		cm	8 individuals
	Aboveground biomass		g	8 individuals
Leaf	Specific leaf area	SLA	g cm <sup>-2</sup>	8 individuals
	Leaf dry matter content	LDMC	mg g <sup>-1</sup>	8 individuals
	Leaf carbon content*	Leaf C	mg g <sup>-1</sup>	pooled
	Leaf nitrogen content*	Leaf N	mg g <sup>-1</sup>	pooled
	Leaf phosphorus content*	Leaf P	mg g <sup>-1</sup>	pooled
Seed	Seed mass*		mg	5 individuals

**Table 4.** Summary of mean  $\pm$  standard error (SE) of the measured vegetation and soil parameters and community-weighted means (CWM) for the three land-use types. The statistical test values are given. Different letters indicate significant differences between the three land-use types as extracted from pairwise tests. \*\*\*  $P < 0.001$ ; \*\*  $P < 0.01$ ; \*  $P < 0.05$ ; n.s.  $P > 0.05$


	Steppe	Cereal	Melon	Statistical test values	
	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE		
Vegetation	Species richness	35.3 $\pm$ 0.1 a	29.0 $\pm$ 1.6 b	22.7 $\pm$ 1.5 c	$\chi^2 = 27.5^{***}$
	Shannon diversity index	2.6 $\pm$ 0.06 a	2.3 $\pm$ 0.1 a	1.9 $\pm$ 0.09 b	$\chi^2 = 24.7^{***}$
	Evenness	0.7 $\pm$ 0.01 a	0.7 $\pm$ 0.02 a	0.6 $\pm$ 0.02 b	$\chi^2 = 15.9^{***}$
	Standing crop	42.8 $\pm$ 1.2 a	17.9 $\pm$ 0.7 b	18.7 $\pm$ 0.8 b	F = 289.5 <sup>***</sup>
	Leaf area index	3.0 $\pm$ 0.1 a	1.3 $\pm$ 0.2 b	1.0 $\pm$ 0.1 b	F = 57.2 <sup>***</sup>
	Vegetation cover	60.3 $\pm$ 1.9 a	50.2 $\pm$ 3.2 b	43.7 $\pm$ 3.4 b	F = 10.2 <sup>***</sup>
	Stone cover	46.5 $\pm$ 1.2 a	36.6 $\pm$ 2.5 b	45.0 $\pm$ 2.0 a	F = 10.7 <sup>***</sup>
	Bare ground	9.9 $\pm$ 0.7 b	16.2 $\pm$ 1.2 a	18.8 $\pm$ 2.5 a	F = 13.6 <sup>***</sup>
Soil	Clays	192.7 $\pm$ 2.0 c	202.2 $\pm$ 3.4 b	213.4 $\pm$ 2.6 a	F = 20.0 <sup>***</sup>
	Fine silt	192.1 $\pm$ 1.7	191.4 $\pm$ 1.1	194.3 $\pm$ 2.3	n.s.
	Coarse silt	161.5 $\pm$ 4.0 a	140.1 $\pm$ 1.2 b	144.6 $\pm$ 2.1 b	F = 12.0 <sup>***</sup>
	Fine sand	210.8 $\pm$ 4.8	203.7 $\pm$ 3.2	214.4 $\pm$ 5.4	n.s.
	Coarse sand	242.9 $\pm$ 6.2 b	262.7 $\pm$ 4.2 a	233.2 $\pm$ 7.8 b	F = 7.2 <sup>**</sup>
	Total Carbon	16.2 $\pm$ 0.3	15.6 $\pm$ 0.2	15.4 $\pm$ 0.6	n.s.
	Total Nitrogen	1.4 $\pm$ 0.03	1.5 $\pm$ 0.02	1.5 $\pm$ 0.05	n.s.
	Carbon-Nitrogen ratio	11.0 $\pm$ 0.06 a	10.6 $\pm$ 0.04 b	10.5 $\pm$ 0.09 b	F = 25.0 <sup>***</sup>
	Organic matter	28.0 $\pm$ 0.5	26.9 $\pm$ 0.4	26.59 $\pm$ 1.1	n.s.
	pH	6.9 $\pm$ 0.04 c	7.1 $\pm$ 0.03 b	7.25 $\pm$ 0.05 a	F = 20.6 <sup>***</sup>
	Phosphorus	0.009 $\pm$ 0.0003 b	0.03 $\pm$ 0.006 a	0.04 $\pm$ 0.007 a	F = 22.0 <sup>***</sup>
	Calcium	1.4 $\pm$ 0.03 b	1.5 $\pm$ 0.03 ab	1.6 $\pm$ 0.05 a	F = 7.2 <sup>**</sup>
	Magnesium	0.9 $\pm$ 0.009	0.9 $\pm$ 0.02	0.9 $\pm$ 0.03	n.s.
	Potassium	0.3 $\pm$ 0.01 c	0.4 $\pm$ 0.03 b	0.4 $\pm$ 0.03 a	$\chi^2 = 24.7^{***}$
CWM traits	SLA	15.2 $\pm$ 0.3 b	21.4 $\pm$ 0.7 a	21.4 $\pm$ 0.9 a	$\chi^2 = 54.2^{***}$
	LDMC	333.9 $\pm$ 5.7 a	252.4 $\pm$ 6.3 b	276.0 $\pm$ 5.5 b	F = 52.4 <sup>***</sup>
	Canopy height	32.6 $\pm$ 0.5 a	29.3 $\pm$ 1.2 b	29.5 $\pm$ 1.1 ab	F = 5.2 <sup>**</sup>



**Figure 1.** Map of La Crau area in France with the six selected locations. Modified from Sylvain Fadda, IMBE (In light grey, the area of steppe remnants ). At every location, we chose at least two study sites that represented one undisturbed steppe and at least one fallow land.




**Figure 2.** Detrended correspondence analysis (DCA) of all plots. Main matrix: vegetation data (16 sites with 5 vegetation relevés; 136 species) and second matrix: environmental factors (data on 19 parameters for 16 sites). In total, plots from eight steppe sites (white), five cereal fallows (light grey; rectangle: long-abandoned field (since 1900); triangle: recently-abandoned field (since 2000) and three melon fallows (dark grey) are shown. Vectors were significantly related to the axes ( $P < 0.05$ ).



**Figure 3.** Results of non-metric multidimensional scaling (NMDS) carried out on the presence/ absence vegetation data for 2001 and 2015. The color of polygons (convex hulls) indicates the land- use types (white: steppe, light grey: cereal fallows, dark grey: melon fallows) while the structure represents the time of assessment (plain: assessed in 2001 with  $n = 10$ , dotted: assessed in 2015 with  $n = 5$ ). Vegetation relevés from Peau de Meau, Valigne and Petit Abondoux were included. Final stress = 0.19. For clarity, only 56 species are displayed (out of 130).





**Figure 4.** Coefficient of variation (CV) of (a) specific leaf area (SLA), (b) leaf dry matter content (LDMC), (c) canopy height, (d) plant width and (e) biomass for selected species occurring in the steppe sites. The scales of the y-axis differ among panels. In the right-hand corner, the total CV of plant traits is given, represented by a dashed line in the boxplots. Kruskal–Wallis tests were applied followed by the pairwise Wilcoxon test with P-value adjustment according to Holm (\*\*\*)  $P < 0.001$ ; \*\*  $P < 0.01$ ; \*  $P < 0.05$ ; n.s.  $P > 0.05$ ). Different letters indicate significant differences among species. For abbreviations of species names, see Table 2.


### Supporting information

Helm, J. et al. Recovery of Mediterranean steppe vegetation after cultivation: legacy effects on plant composition, soil properties and functional traits, Applied Vegetation Science.

**Appendix S1.** Trait values of specific leaf area (SLA), leaf dry matter content (LDMC) and canopy height for all species. Mean  $\pm$  standard errors are given for the measured trait values per species and land-use type. Different letters indicate significant differences between steppe, cereal and melon fallow. \*\*\*  $P < 0.001$ ; \*\*  $P < 0.01$ ; \*  $P < 0.05$ ; n.s.  $P > 0.05$

	Species	SLA				LDMC				Canopy height			
		Steppe	Cereal	Melon	Statistical test values	Steppe	Cereal	Melon	Statistical test values	Steppe	Cereal	Melon	Statistical test values
Unsuccessful colonizers	Bra_ret	9.5 $\pm$ 0.3	-	-	-	512.8 $\pm$ 7.1	-	-	-	15.6 $\pm$ 0.5	-	-	-
	Hyp_gla	22.7 $\pm$ 1.6	-	-	-	152.3 $\pm$ 13.5	-	-	-	2.0 $\pm$ 0.4	-	-	-
	Lin_tri	17.8 $\pm$ 0.7 a	16.2 $\pm$ 0.6 ab	15.0 $\pm$ 1.0 b	$\chi^2 = 7.4^*$	330.9 $\pm$ 11.9	327.4 $\pm$ 13.2	324.2 $\pm$ 18.0	n.s.	11.9 $\pm$ 0.4	11.3 $\pm$ 0.8	10.7 $\pm$ 0.7	n.s.
	Sti_cap	5.1 $\pm$ 0.1	4.9 $\pm$ 0.1	5.0 $\pm$ 0.3	n.s.	569.9 $\pm$ 7.2	575.6 $\pm$ 16.9	558.0 $\pm$ 9.1	n.s.	32.7 $\pm$ 1.0 a	26.4 $\pm$ 1.5 b	25.3 $\pm$ 1.6 b	F = 9.8***
	Tri_cam	27.7 $\pm$ 0.7	-	-	-	281.7 $\pm$ 7.0	-	-	-	6.1 $\pm$ 0.4	-	-	-
Fallow-land species	Cre_foe	22.2 $\pm$ 0.6	19.9 $\pm$ 0.6	20.1 $\pm$ 0.7	n.s.	192.7 $\pm$ 4.3 a	181.4 $\pm$ 5.5 b	186.7 $\pm$ 5.2 ab	$\chi^2 = 6.6^*$	4.4 $\pm$ 0.3 ab	4.0 $\pm$ 0.3 a	5.3 $\pm$ 0.4 b	$\chi^2 = 9.7^{**}$
	Dac_glo	20.8 $\pm$ 0.4	20.8 $\pm$ 0.5	22.0 $\pm$ 0.6	n.s.	361.0 $\pm$ 4.7	360.8 $\pm$ 5.5	360.5 $\pm$ 5.1	n.s.	14.8 $\pm$ 0.6 a	11.7 $\pm$ 0.6 b	11.9 $\pm$ 0.7 b	$\chi^2 = 11.8^{**}$
	Gal_tom	11.3 $\pm$ 0.3 a	10.9 $\pm$ 0.3 a	9.6 $\pm$ 0.4 b	F = 5.5**	228.9 $\pm$ 8.2	233.2 $\pm$ 10.8	228.8 $\pm$ 19.1	n.s.	28.3 $\pm$ 1.9	32.2 $\pm$ 2.0	27.9 $\pm$ 2.2	n.s.
Successful colonizers	Tri_sub	-	19.4 $\pm$ 0.5	-	-	-	313.5 $\pm$ 10.8	-	-	-	2.3 $\pm$ 0.2	-	-
	Car_lan	7.4 $\pm$ 0.2 a	6.5 $\pm$ 0.2 b	7.7 $\pm$ 0.3 a	$\chi^2 = 15.9^{***}$	394.3 $\pm$ 8.5 a	341.8 $\pm$ 11.1 b	366.7 $\pm$ 15.1 ab	F = 4.3*	23.0 $\pm$ 1.1 a	15.8 $\pm$ 1.0 b	25.0 $\pm$ 1.5 a	$\chi^2 = 15.1^{***}$
	Lob_mar	11.3 $\pm$ 0.3	10.3 $\pm$ 0.4	10.2 $\pm$ 0.5	n.s.	303.1 $\pm$ 7.6	323.7 $\pm$ 9.8	333.1 $\pm$ 12.1	n.s.	8.9 $\pm$ 0.3 a	6.3 $\pm$ 0.4 b	6.2 $\pm$ 0.3 b	F = 22.5***
	Sid_rom	24.0 $\pm$ 0.5	22.5 $\pm$ 0.6	23.1 $\pm$ 0.6	n.s.	224.8 $\pm$ 3.9	230.0 $\pm$ 6.5	222.9 $\pm$ 7.7	n.s.	7.0 $\pm$ 0.4 a	3.3 $\pm$ 0.3 c	5.4 $\pm$ 0.6 b	$\chi^2 = 29.6^{***}$

**Appendix S2.** Principal component analysis (PCA) of the species–trait matrix (854 Individuals of 9 species × 9 traits; without *Brachypodium retusum* and *Trifolium subterraneum* due to data constraints). Symbols represent the location of individuals in the multidimensional trait space. Species groups are indicated by colors (white: steppe species; light grey: fallow land species; dark grey: common species). Origins of measurements are distinguished: steppe (triangle) and fallow lands (circle). As a conceptual summary, the black arrow indicates the direction of changes in plant trait values due to land-use change (from steppe to fallow land). For abbreviations of species, see Table 2. For abbreviations of functional traits, see Table 3.



**Appendix S3.** Leaf nutrient content of carbon (C), nitrogen (N) and phosphorus (P) compared between original steppe and fallow lands (eight individuals per site pooled). Mean  $\pm$  standard errors (SE) are shown. P was measured with inductively-coupled plasma optical emission spectrometry (IPC-OES, Optima 3000, Fa. Perkin Elmer) at the TLL (Thüringer Landesanstalt für Landwirtschaft, Jena). Leaf N and leaf P content were analyzed with a CHN elemental analyzer (Vario EL Elemental Analyzer, Elementar). Leaf nutrient content could not be measured for *Hypochaeris glabra*, *Trifolium campestre* and *Trifolium subterraneum* due to insufficient plant material.

	<b>Steppe</b>	<b>Cereal</b>	<b>Melon</b>	<b>Statistical test values</b>
	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	
Leaf carbon content	43.24 $\pm$ 0.25 a	41.40 $\pm$ 0.34 b	42.13 $\pm$ 0.45 ab	$\chi^2 = 13.0^{**}$
Leaf nitrogen content	1.70 $\pm$ 0.08 b	2.15 $\pm$ 0.14 a	1.94 $\pm$ 0.18 ab	$\chi^2 = 6.7^*$
Leaf phosphorus content	1.26 $\pm$ 0.01 c	2.86 $\pm$ 0.07 b	3.39 $\pm$ 0.10 a	$\chi^2 = 58.1^{***}$