



HAL
open science

DTM-based filtrations

Hirokazu Anai, Frédéric Chazal, Marc Glisse, Yuichi Ike, Hiroya Inakoshi,
Raphaël Tinarrage, Yuhei Umeda

► **To cite this version:**

Hirokazu Anai, Frédéric Chazal, Marc Glisse, Yuichi Ike, Hiroya Inakoshi, et al.. DTM-based filtrations. 2018. hal-01919562v1

HAL Id: hal-01919562

<https://hal.science/hal-01919562v1>

Preprint submitted on 12 Nov 2018 (v1), last revised 22 May 2020 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DTM-BASED FILTRATIONS

Hirokazu ANAI[†], Frédéric CHAZAL^{*}, Marc GLISSE^{*}, Yuichi IKE[†], Hiroya INAKOSHI[†], Raphaël TINARRAGE^{*}, Yuhei UMEDA[†]

^{*}Datashape, Inria Paris-Saclay

[†] Fujitsu Laboratories, AI Lab

Abstract. Despite strong stability properties, the persistent homology of filtrations classically used in Topological Data Analysis, such as, e.g. the Čech or Vietoris-Rips filtrations, are very sensitive to the presence of outliers in the data from which they are computed. In this paper, we introduce and study a new family of filtrations, the DTM-filtrations, built on top of point clouds in the Euclidean space which are more robust to noise and outliers. The approach adopted in this work relies on the notion of distance-to-measure functions [1], and extends some previous work on the approximation of such functions [2, 3].

Contents

1	Introduction	1
2	Filtrations and interleaving distance	3
3	Weighted Čech filtrations	5
3.1	Definition	5
3.2	Stability	7
3.3	Weighted Vietoris-Rips filtrations	8
4	DTM-filtrations	10
4.1	The distance to measure (DTM)	10
4.2	DTM-filtrations	11
4.3	Stability when $p = 1$	13
4.4	Stability when $p > 1$	16
4.5	Proof of Proposition 4.8	19
5	Conclusion	23
	References	24
A	Supplementary results for Section 3	24
B	Supplementary results for Section 4	25

1 Introduction

The inference of relevant topological properties of data represented as point clouds in Euclidean spaces is a central challenge in Topological Data Analysis (TDA).

Given a (finite) set of points X in \mathbb{R}^d , persistent homology provides a now classical and powerful tool to construct persistence diagrams whose points can be interpreted as homological features of X at different scales. These persistence diagrams are obtained from *filtrations*, i.e. nested families of subspaces or simplicial complexes, built on top of X . Among the many filtrations available to the user, unions of growing balls $\cup_{x \in X} B(x, t)$ (sublevel sets of distance functions), $t \in \mathbb{R}^+$, and their nerve, the Čech complex filtration, or its usually easier to compute version, the Vietoris-Rips filtration, are widely used. The main theoretical advantage of these filtrations is that they have been shown to produce persistence diagrams that are stable with respect to perturbations of X in the Hausdorff metric [4].

Unfortunately, the Hausdorff distance turns out to be very sensitive to noise and outliers, preventing the direct use of distance functions and classical Čech or Vietoris-Rips filtrations to infer relevant topological properties from real noisy data. Several attempts have been made in the recent years to overcome this issue. Among them, the filtration defined by the sublevel sets of the distance-to-measure (DTM) function introduced in [1], and some of its variants [5], have been proven to provide relevant information about the geometric structure underlying the data. Unfortunately, from a practical perspective, the computation of the sublevel sets filtration of the DTM and its persistent homology turns out to be far too expensive in most cases. To address this problem, [2] introduces a variant of the DTM function, the k -witness distance, whose persistence is easier to compute and proves that the k -witness distance approximates the DTM persistence up to a fixed additive constant. In [3, 6], a weighted version of the Vietoris-Rips complex filtration is introduced to approximate the persistence of the DTM function, and several stability and approximation results, comparable to the ones of [2], are established.

In this paper, we introduce and study a new family of filtrations based on the notion of DTM. Our contributions are the following:

- Given a set $X \subset \mathbb{R}^d$, a weight function f defined on \mathbb{R}^d and a real number $p \geq 1$, we introduce the weighted Čech and Rips filtrations that extend the notion of sublevel set filtration of power distances of [3]. Using classical results we show that these filtrations are stable with respect to perturbations of X in the Hausdorff metric and perturbations of f with respect to the sup norm (Propositions 3.2 and 3.3).
- As for a general function f , the stability results for the weighted Čech and Rips filtrations are not adapted to deal with noisy data or data containing outliers, we consider the case where f is the DTM-function associated to the input point cloud. In this case, we show that, given two point clouds $X, Y \subseteq \mathbb{R}^d$, the closeness between the persistence diagrams of the resulting filtrations relies on the existence of a subset of X which is both close to X and Y in the Wasserstein metric—here the point clouds are seen as empirical measures—see Theorems 4.5 and 4.9.

Practical motivations. Even though this aspect is not considered in this paper, it is interesting to mention that the DTM filtration was first experimented in the setting of an industrial research project whose goal was to address an anomaly detection problem from inertial sensor data in bridge and building monitoring [7]. In this problem, the input

data comes as time series measuring the acceleration of devices attached to the monitored bridge/building. Using sliding windows and time-delay embedding, these times series are converted into a series of fixed size point clouds in \mathbb{R}^d . Filtrations are then built on top of these point clouds and their persistence is computed, giving rise to a time-dependent sequence of persistence diagrams that are then used to detect anomalies or specific features occurring along the time [8, 9]. In this practical setting it turned out that the DTM filtrations reveals to be not only more resilient to noise but also able to better highlight topological features in the data than the standard Vietoris-Rips filtrations, as illustrated on a basic synthetic example on Figure 1. One of the goal of the present work is to provide theoretical foundations to these promising experimental results by studying the stability properties of the DTM filtrations.




Figure 1: A synthetic example comparing Vietoris-Rips filtration to DTM filtration. The first row represents two time series with very different behavior and their embedding into \mathbb{R}^3 (here a series (x_1, x_2, \dots, x_n) is converted in the 3D point cloud $\{(x_1, x_2, x_3), (x_2, x_3, x_4), \dots, (x_{n-2}, x_{n-1}, x_n)\}$). The second row shows the persistence diagrams of the Vietoris-Rips filtration built on top of the two point clouds (red and green points represent respectively the 0-dimensional 1-dimensional diagrams); one observes that the diagrams do not clearly ‘detect’ the different behavior of the time series. The third row shows the persistence diagrams of the DTM filtration built on top of the two point clouds; a red point clearly appear away from the diagonal in the second diagram that highlight the rapid shift occurring in the second time series.

Organisation of the paper. Preliminary definitions, notations, and basic notions on filtrations and persistence modules are recalled in Section 2. The weighted Čech and Vietoris-Rips filtrations are introduced in Section 3 where their stability properties are established. The DTM-filtrations are introduced in Section 4. Their main stability property are established in Theorems 4.5 and 4.9, and their relation with the sublevel set filtration of the DTM-functions is established in Proposition 4.6. For the clarity of the paper, the proof of several lemmas have been postponed to the appendices.

Acknowledgements. This work was partially supported by a collaborative research agreement between Inria and Fujitsu, and the Advanced Grant of the European Research Council GUDHI (Geometric Understanding in Higher Dimensions).

2 Filtrations and interleaving distance

In the sequel, we consider interleavings of filtrations, interleavings of persistence modules and their associated pseudo-distances. Their definitions, restricted to the setting of the paper, are briefly recalled in this section.

Let $T = \mathbb{R}^+$ and $E = \mathbb{R}^d$ be endowed with the standard Euclidean norm.

Filtrations of sets and simplicial complexes. A family of subsets $(V^t)_{t \in T}$ of $E = \mathbb{R}^d$ is a *filtration* if it is non-decreasing for the inclusion, i.e. for any $s, t \in T$, if $s \leq t$ then $V^s \subseteq V^t$. Given $\epsilon \geq 0$, two filtrations $(V^t)_{t \in T}$ and $(W^t)_{t \in T}$ of E are ϵ -*interleaved* if, for every $t \in T$, $V^t \subseteq W^{t+\epsilon}$ and $W^t \subseteq V^{t+\epsilon}$. The interleaving pseudo-distance between $(V^t)_{t \in T}$ and $(W^t)_{t \in T}$ is defined as the infimum of such ϵ :

$$d_i((V^t)_{t \in T}, (W^t)_{t \in T}) = \inf\{\epsilon : (V^t) \text{ and } (W^t) \text{ are } \epsilon\text{-interleaved}\}.$$

Filtrations of simplicial complexes and their interleaving distance are similarly defined: given a set X and S an abstract simplex with vertex set X , a *filtration of S* is a non-decreasing family $(S^t)_{t \in T}$ of subcomplexes of S . The interleaving pseudo-distance between two filtrations $(S_1^t)_{t \in T}$ and $(S_2^t)_{t \in T}$ of S is the infimum of the $\epsilon \geq 0$ such that they are ϵ -interleaved, i.e. for any $t \in T$, $S_1^t \subseteq S_2^{t+\epsilon}$ and $S_2^t \subseteq S_1^{t+\epsilon}$.

Notice that the interleaving distance is only a pseudo-distance, as two distinct filtrations may have zero interleaving distance.

Persistence modules. Let k be a field. A persistence module \mathbb{V} over $T = \mathbb{R}^+$ is a pair $\mathbb{V} = ((\mathbb{V}^t)_{t \in T}, (v_s^t)_{s \leq t \in T})$ where $(\mathbb{V}^t)_{t \in T}$ is a family of k -vector spaces, and $(v_s^t : \mathbb{V}^s \rightarrow \mathbb{V}^t)_{s \leq t \in T}$ a family of linear maps such that:

- for every $t \in T$, $v_t^t : \mathbb{V}^t \rightarrow \mathbb{V}^t$ is the identity map,
- for every $r, s, t \in T$ such that $r \leq s \leq t$, $v_s^t \circ v_r^s = v_r^t$.

An isomorphism between two persistence modules \mathbb{V} and \mathbb{W} is a family of isomorphisms of vector spaces $(\phi_t : \mathbb{V}^t \rightarrow \mathbb{W}^t)_{t \in T}$ such that the following diagrams commute for every $s \leq t \in T$

$$\begin{array}{ccc} \mathbb{V}^s & \xrightarrow{v_s^t} & \mathbb{V}^t \\ \downarrow \phi_s & & \downarrow \phi_t \\ \mathbb{W}^s & \xrightarrow{w_s^t} & \mathbb{W}^t \end{array}$$

Given $\epsilon \geq 0$, an ϵ -interleaving between two persistence modules \mathbb{V} and \mathbb{W} is a pair of families of linear maps $(\phi_t : \mathbb{V}^t \rightarrow \mathbb{W}^{t+\epsilon})_{t \in T}$ and $(\psi_t : \mathbb{W}^t \rightarrow \mathbb{V}^{t+\epsilon})_{t \in T}$ such that the following diagrams commute for every $t \in T$:

$$\begin{array}{ccc}
\mathbb{V}t & \xrightarrow{v_t^{t+2\epsilon}} & \mathbb{V}t+2\epsilon \\
& \searrow \phi_t & \nearrow \psi_{t+\epsilon} \\
& & \mathbb{W}t+\epsilon
\end{array}
\qquad
\begin{array}{ccc}
& & \mathbb{V}t+\epsilon \\
& \nearrow \psi_t & \searrow \phi_{t+\epsilon} \\
\mathbb{W}t & \xrightarrow{w_t^{t+2\epsilon}} & \mathbb{W}t+2\epsilon
\end{array}$$

The interleaving pseudo-distance between \mathbb{V} and \mathbb{W} is defined as

$$d_i(\mathbb{V}, \mathbb{W}) = \inf\{\epsilon \geq 0, \mathbb{V} \text{ and } \mathbb{W} \text{ are } \epsilon\text{-interleaved}\}.$$

In some cases, the proximity between persistence modules is expressed with a function. Let $\eta : T \rightarrow T$ be a non-decreasing function such that for any $t \in T$, $\eta(t) \geq t$. A η -interleaving between two persistence modules \mathbb{V} and \mathbb{W} is a pair of families of linear maps $(\phi_t : \mathbb{V}^t \rightarrow \mathbb{W}^{\eta(t)})_{t \in T}$ and $(\psi_t : \mathbb{W}^t \rightarrow \mathbb{V}^{\eta(t)})_{t \in T}$ such that the following diagrams commute for every $t \in T$:

$$\begin{array}{ccc}
\mathbb{V}t & \xrightarrow{v_t^{\eta(\eta(t))}} & \mathbb{V}\eta(\eta(t)) \\
& \searrow \phi_t & \nearrow \psi_{\eta(t)} \\
& & \mathbb{W}\eta(t)
\end{array}
\qquad
\begin{array}{ccc}
& & \mathbb{V}\eta(t) \\
& \nearrow \psi_t & \searrow \phi_{\eta(t)} \\
\mathbb{W}t & \xrightarrow{v_t^{\eta(\eta(t))}} & \mathbb{W}\eta(\eta(t))
\end{array}$$

When η is $t \mapsto t + c$ for some $c \geq 0$, it is called an additive c -interleaving and corresponds with the previous definition. When η is $t \mapsto ct$ for some $c \geq 1$, it is called a multiplicative c -interleaving.

A persistent module \mathbb{V} is said to be q -tame if for every $s, t \in T$ such that $s < t$, the map v_s^t is of finite rank. The q -tameness of a persistence module ensures that we can define a notion of persistence diagram—see [10]. Moreover, given two q -tame persistence modules \mathbb{V}, \mathbb{W} with persistence diagrams $D(\mathbb{V}), D(\mathbb{W})$, the so-called isometry theorem states that $d_b(D(\mathbb{V}), D(\mathbb{W})) = d_i(\mathbb{V}, \mathbb{W})$ ([10, Theorem 4.11]) where $d_b(\cdot, \cdot)$ denotes the bottleneck distance between diagrams.

Relation between filtrations and persistence modules. Applying the homology functor to a filtration gives rise to a persistence module where the linear maps between homology groups are induced by the inclusion maps between sets (or simplicial complexes). As a consequence, if two filtrations are ϵ -interleaved then their associated homology persistence modules are also ϵ -interleaved, the interleaving homomorphism being induced by the interleaving inclusion maps. Moreover, if the considered modules are q -tame, then the bottleneck distance between their persistence diagrams is upperbounded by ϵ .

The filtrations considered in this paper are obtained as union of growing balls. Their associated persistence module is the same as the persistence module of a filtered simplicial complex via the persistent nerve lemma ([11, Lemma 3.4]). Indeed, consider a filtration $(V^t)_{t \in T}$ of E and assume that there exists a family of points $(x_i)_{i \in I} \in E^I$ and a family of non-decreasing functions $r_i : T \rightarrow \mathbb{R}^+ \cup \{-\infty\}$ such that, for every $t \in T$, V^t is equal to the union of closed balls $\bigcup_I \overline{B}(x_i, r_i(t))$, with the convention $\overline{B}(x_i, -\infty) = \emptyset$. For every $t \in T$, let \mathcal{V}^t denotes the cover $\{\overline{B}(x_i, r_i(t)), i \in I\}$ of V^t , and S^t be its nerve. Let \mathbb{V} be the persistence module associated with the filtration $(V^t)_{t \in T}$, and $\mathbb{V}_{\mathcal{N}}$ the one associated with the simplicial filtration $(S^t)_{t \in T}$. Then \mathbb{V} and $\mathbb{V}_{\mathcal{N}}$ are isomorphic persistence modules. In particular, if \mathbb{V} is q -tame, \mathbb{V} and $\mathbb{V}_{\mathcal{N}}$ have the same persistence diagrams.

3 Weighted Čech filtrations

In order to define the DTM-filtrations, we go through an intermediate and more general construction, namely the weighted Čech filtrations. It generalizes the usual notion of Čech filtration of a subset of \mathbb{R}^d , and shares comparable regularity properties.

3.1 Definition

In the following, we consider the Euclidean space $E = \mathbb{R}^d$, the index set $T = \mathbb{R}^+$, a function $f : E \rightarrow \mathbb{R}^+$ and a real number $p \geq 1$. For every $x \in E$ and $t \in T$, we define

$$r_x(t) = \begin{cases} -\infty & \text{if } t < f(x), \\ (t^p - f(x)^p)^{\frac{1}{p}} & \text{otherwise.} \end{cases}$$

We denote by $\bar{B}_f(x, t) = \bar{B}(x, r_x(t))$ the closed Euclidean ball of center x and radius $r_x(t)$. By convention, an Euclidean ball of radius $-\infty$ is the empty set. For $p = \infty$, we also define

$$r_x(t) = \begin{cases} -\infty & \text{if } t < f(x), \\ t & \text{otherwise,} \end{cases}$$

and the balls $\bar{B}_f(x, t) = \bar{B}(x, r_x(t))$. Some of these radius functions are represented in Figure 2.




Figure 2: Graph of $t \mapsto r_x(t)$ for $f(x) = 1$ and several values of p .

Definition 3.1. Let $X \subseteq E$. For every $t \in T$, we define the following set:

$$V^t[X, f] = \bigcup_{x \in X} \bar{B}_f(x, t).$$

The family $V[X, f] = (V^t[X, f])_{t \geq 0}$ is a filtration of E . It is called the *weighted Čech filtration with parameters* (X, f, p) . We denote by $\mathbb{V}[X, f]$ its persistent (singular) homology module.

Introduce $\mathcal{V}^t[X, f] = \{\bar{B}_f(x, t)\}_{x \in X}$. It is a cover of $V^t[X, f]$ by closed Euclidean balls. Let $\mathcal{N}(\mathcal{V}^t[X, f])$ be the nerve of the cover $\mathcal{V}^t[X, f]$. It is a simplicial complex over the vertex set X . The family $\mathcal{N}(\mathcal{V}[X, f]) = (\mathcal{N}(\mathcal{V}^t[X, f]))_{t \geq 0}$ is a filtered simplicial complex.

We denote by $\mathbb{V}_{\mathcal{N}}[X, f]$ its persistent (simplicial) homology module. As a consequence of the persistent nerve theorem ([11], Lemma 3.4), $\mathbb{V}[X, f]$ and $\mathbb{V}_{\mathcal{N}}[X, f]$ are isomorphic persistent modules.

When $f = 0$, $V[X, f]$ does not depend on $p \geq 1$, and it is the filtration of E by the sublevel sets of the distance function to X . In the sequel, we denote it by $V[X, 0]$. The corresponding filtered simplicial complex, $\mathcal{N}(V[X, 0])$, is known as the usual Čech complex of X .

When $p = 2$, the filtration value of $y \in E$, i.e. the infimum of the t such that $y \in V^t[X, f]$, is called the power distance of y associated to the weighted set (X, f) in [3, Definition 4.1]. The filtration $V[X, f]$ is called the weighted Čech filtration ([3, Definition 4.4]).

Example. Consider the point cloud X drawn on the right (black). It is a 200-sample of the uniform distribution on $[-1, 1]^2 \subseteq \mathbb{R}^2$. We choose f to be the distance function to the Cassini oval (magenta). Let $t = 0.2$. Figure 3 represents the sets $V^t[X, f]$ for several values of p .





Figure 3: The sets $V^t[X, f]$ for $t = 0.2$ and several values of p .

The following proposition states the regularity of the persistent module $\mathbb{V}[X, f]$.

Proposition 3.1. *If $X \subseteq E$ is finite and f is any function, then $\mathbb{V}[X, f]$ is a pointwise finite-dimensional persistence module.*

More generally, if X is any bounded subset of E and f is continuous, then $\mathbb{V}[X, f]$ is q -tame.

Proof. First, suppose that X is finite. Then $\mathcal{N}(V[X, f])$ is a filtration of a finite simplicial complex, and thus $\mathbb{V}_{\mathcal{N}}[X, f]$ is pointwise finite-dimensional. It is also the case for $\mathbb{V}[X, f]$ since it is isomorphic to $\mathbb{V}_{\mathcal{N}}[X, f]$.

Secondly, suppose that X is bounded and f is continuous. Consider the ‘filtration value’ function:

$$t_X : E \longrightarrow \mathbb{R}^+ \\ y \longmapsto \inf \{ t \in \mathbb{R}^+, \exists x \in X, y \in \overline{B}_f(x, t) \}$$

For every $y \in E$, $x \in X$ and $t \geq 0$ the assertion $y \in \overline{B}_f(x, t)$ is equivalent to $(\|x - y\|^p + f(x)^p)^{\frac{1}{p}} \leq t$. Therefore the function t_X can be written as follows:

$$t_X(y) = \inf \{ (\|x - y\|^p + f(x)^p)^{\frac{1}{p}}, x \in X \}.$$

It is 1-Lipschitz as it is the infimum of the set of the 1-Lipschitz functions $y \mapsto (\|x - y\|^p + f(x)^p)^{\frac{1}{p}}$. It is also proper as X is bounded.

Let \tilde{V} be the filtration of E defined for all $t \geq 0$ by $\tilde{V}^t = t_X^{-1}([-\infty, t])$. Let $\tilde{\mathbb{V}}$ be its persistent homology module. The last two properties of t_X (continuous and proper) imply that $\tilde{\mathbb{V}}$ is q -tame ([10], Corollary 3.34).

Notice that, since X may not be compact, $V^t[X, f]$ may not be equal to \tilde{V}^t . However, it follows from the definition of t_X that $V[X, f]$ and \tilde{V} are ϵ -interleaved for every $\epsilon > 0$. Therefore, $\mathbb{V}[X, f]$ also is q -tame. \square

3.2 Stability

We still consider a function $f : E \rightarrow \mathbb{R}^+$ and a subset $X \subseteq E$. Using the fact that two ϵ -interleaved filtrations induce ϵ -interleaved persistence modules, we immediately obtain stability results for the persistence module $\mathbb{V}[X, f]$.

The following proposition relates the stability of the filtration $V[X, f]$ with respect to f .

Proposition 3.2. *Let $g : E \rightarrow \mathbb{R}^+$ be a function such that $\|f - g\|_\infty = \sup_{x \in E} |f(x) - g(x)| \leq \epsilon$. Then the filtrations $V[X, f]$ and $V[X, g]$ are ϵ -interleaved.*

Proof. By symmetry, it suffices to show that, for every $t \geq 0$, $V^t[X, f] \subseteq V^{t+\epsilon}[X, g]$. Let $t \geq 0$. Choose $y \in V^t[X, f]$, and $x \in X$ such that $y \in \overline{B}_f(x, t)$, i.e. $(\|x - y\|^p + f(x)^p)^{\frac{1}{p}} \leq t$. Let us prove that $y \in \overline{B}_g(x, t + \epsilon)$, i.e. $(\|x - y\|^p + g(x)^p)^{\frac{1}{p}} \leq t + \epsilon$. From $g(x) \leq f(x) + \epsilon$, we obtain $(\|x - y\|^p + g(x)^p)^{\frac{1}{p}} \leq (\|x - y\|^p + (f(x) + \epsilon)^p)^{\frac{1}{p}}$. Now, consider the function $\eta \mapsto (\|x - y\|^p + (f(x) + \eta)^p)^{\frac{1}{p}}$. Its derivative is $\eta \mapsto \left(\frac{f(x) + \eta}{\|x - y\|^p + (f(x) + \eta)^p} \right)^{p-1}$. It is consequently 1-Lipschitz on \mathbb{R}^+ . The Lipschitz property implies that

$$(\|x - y\|^p + (f(x) + \epsilon)^p)^{\frac{1}{p}} \leq (\|x - y\|^p + f(x)^p)^{\frac{1}{p}} + \epsilon$$

Hence $(\|x - y\|^p + g(x)^p)^{\frac{1}{p}} \leq (\|x - y\|^p + (f(x) + \epsilon)^p)^{\frac{1}{p}} \leq (\|x - y\|^p + f(x)^p)^{\frac{1}{p}} + \epsilon \leq t + \epsilon$. \square

The following proposition states the stability of $V[X, f]$ with respect to X . It generalizes [3, Proposition 4.3] (case $p = 2$).

Proposition 3.3. *Suppose that f is c -Lipschitz, $c \geq 0$. Let $Y \subseteq E$. Suppose that X and Y are compact, and that the Hausdorff distance $d_H(X, Y) \leq \epsilon$. Then the filtrations $V[X, f]$ and $V[Y, f]$ are k -interleaved with $k = \epsilon(1 + c^p)^{\frac{1}{p}}$.*

Proof. It suffices to show that for every $t \geq 0$, $V^t[X, f] \subseteq V^{t+k}[Y, f]$. Let $t \geq 0$. Choose $z \in V^t[X, f]$, and $x \in X$ such that $z \in \overline{B}_f(x, t)$, i.e. $\|x - z\| \leq r_x(t)$. From the hypothesis $d_H(X, Y) \leq \epsilon$, there exists $y \in Y$ such that $\|y - x\| \leq \epsilon$. Let us prove that $z \in \overline{B}_f(y, t + k)$, i.e. $\|z - y\| \leq r_y(t + k)$.

By triangle inequality, $\|z - y\| \leq \|z - x\| + \|x - y\| \leq r_x(t) + \epsilon$. It is enough to show that $r_x(t) + \epsilon \leq r_y(t + k)$, i.e.

$$\underbrace{((t + k)^p - f(y)^p)^{\frac{1}{p}}}_{r_y(t+k)} - \underbrace{(t^p - f(x)^p)^{\frac{1}{p}}}_{r_x(t)} \geq \epsilon.$$

The left-hand side of this expression is decreasing in $f(y)$. Moreover, since f is c -Lipschitz, $f(y)$ is at most $f(x) + c\epsilon$. Therefore:

$$\begin{aligned} & ((t+k)^p - f(y)^p)^{\frac{1}{p}} - (t^p - f(x)^p)^{\frac{1}{p}} \\ \geq & ((t+k)^p - (f(x) + c\epsilon)^p)^{\frac{1}{p}} - (t^p - f(x)^p)^{\frac{1}{p}}. \end{aligned}$$

It is now enough to prove that this last expression is not less than ϵ , which is the content of Lemma A.1. \square

Notice that the bounds in Proposition 3.2 and 3.3 are tight. In the first case, consider for example $E = \mathbb{R}$, the set $X = \{0\}$, and the functions $f = 0$ and $g = \epsilon$. For every $t < \epsilon$, we have $V^t[Y, f] = \emptyset$, while $V^t[X, f] \neq \emptyset$. Regarding the second proposition, consider $E = \mathbb{R}$, $f : x \mapsto cx$, $X = \{0\}$ and $Y = \{\epsilon\}$. We have, for every $t \geq 0$, $V^t[X, f] = \overline{B}(0, t)$ and $V^t[Y, f] = \overline{B}(\epsilon, (t^p - (c\epsilon)^p)^{\frac{1}{p}})$. For every $t < \epsilon(1 + c^p)^{\frac{1}{p}}$, we have $(t^p - (c\epsilon)^p)^{\frac{1}{p}} < \epsilon$, hence $0 \notin V^t[Y, f]$. In comparison, $\forall t \geq 0$, $0 \in V^t[X, f]$.

When considering data with outliers, the observed set X may be very distant from the underlying signal Y in Hausdorff distance. Therefore, the tight bound in Proposition 3.3 may be unsatisfactory. Moreover, a usual choice of f would be d_X , the distance function to X . But the bound in Proposition 3.2 then becomes $\|d_X - d_Y\|_\infty = d_H(X, Y)$. We address this issue in Section 4 by considering an outliers-robust function f , the so-called distance-to-measure function (DTM).

3.3 Weighted Vietoris-Rips filtrations

Rather than computing the persistence of the Čech filtration of a point cloud $X \subseteq E$, one sometimes consider the corresponding Vietoris-Rips filtration, which is usually easier to compute.

If G is a graph with vertex set X , its corresponding clique complex is the simplicial complex over X consisting of the sets of vertices of cliques of G . If S is a simplicial complex, its corresponding flag complex is the clique complex of its 1-skeleton. We denote it $\text{Rips}(S)$.

Recall that $\mathcal{N}(\mathcal{V}^t[X, f])$ denotes the nerve of $\mathcal{V}^t[X, f]$, where $\mathcal{V}^t[X, f]$ is the cover $\{\overline{B}_f(x, t)\}_{x \in X}$ of $V^t[X, f]$.

Definition 3.2. We denote by $\text{Rips}(\mathcal{V}^t[X, f])$ the flag complex of $\mathcal{N}(\mathcal{V}^t[X, f])$, and by $\text{Rips}(\mathcal{V}[X, f])$ the corresponding filtered simplicial complex. It is called the *weighted Rips complex with parameters* (X, f, p) .

The following proposition states that the filtered simplicial complexes $\mathcal{N}(\mathcal{V}[X, f])$ and $\text{Rips}(\mathcal{V}[X, f])$ are 2-interleaved multiplicatively, generalizing the classical case of the Čech and Vietoris-Rips filtrations (case $f = 0$).

Proposition 3.4. For every $t \geq 0$,

$$\mathcal{N}(\mathcal{V}^t[X, f]) \subseteq \text{Rips}(\mathcal{V}^t[X, f]) \subseteq \mathcal{N}(\mathcal{V}^{2t}[X, f])$$

Proof. Let $t \geq 0$. The first inclusion follows from that $\text{Rips}(\mathcal{V}^t[X, f])$ is the clique complex of $\mathcal{N}(\mathcal{V}^t[X, f])$. To prove the second one, choose a simplex $\omega \in \text{Rips}(\mathcal{V}^t[X, f])$. It means that for every $x, y \in \omega$, $\overline{B}_f(x, t) \cap \overline{B}_f(y, t) \neq \emptyset$, i.e. $\overline{B}(x, r_x(t)) \cap \overline{B}(y, r_y(t)) \neq \emptyset$. We have to prove that $\omega \in \mathcal{N}(\mathcal{V}^{2t}[X, f])$, i.e. $\bigcap_{x \in \omega} \overline{B}(x, r_x(2t)) \neq \emptyset$.

For every $x \in \omega$, one has $r_x(2t) \geq 2r_x(t)$. Indeed,

$$\begin{aligned} r_x(2t) &= \left((2t)^p - f(x)^p \right)^{\frac{1}{p}} \\ &= 2 \left(t^p - \left(\frac{f(x)}{2} \right)^p \right)^{\frac{1}{p}} \\ &\geq 2 \left(t^p - f(x)^p \right)^{\frac{1}{p}} = 2r_x(t) \end{aligned}$$

Using the fact that doubling the radius of pairwise intersecting balls is enough to make them intersect globally, we obtain that $\omega \in \mathcal{N}(\mathcal{V}^{2t}[X, f])$. \square

Note that weighted Rips filtration shares the same stability properties as the weighted Čech filtration. Indeed, the proofs of Proposition 3.2 and 3.3 immediately extend to this case.

In order to compute the flag complex $\text{Rips}(\mathcal{V}^t[X, f])$, it is enough to know the filtration values of its 0- and 1-simplices. The following proposition describes these values.

Proposition 3.5. *Let $p < +\infty$. The filtration value of a vertex $x \in X$ is given by $t_X(\{x\}) = f(x)$.*

The filtration value of an edge $\{x, y\} \subseteq E$ is given by

$$t_X(\{x, y\}) = \begin{cases} \max\{f(x), f(y)\} & \text{if } \|x - y\| \leq |f(x)^p - f(y)^p|^{\frac{1}{p}}, \\ t & \text{otherwise,} \end{cases}$$

where t is the only positive root of

$$\|x - y\| = (t^p - f(x)^p)^{\frac{1}{p}} + (t^p - f(y)^p)^{\frac{1}{p}} \quad (1)$$

When $\|x - y\| \geq |f(x)^p - f(y)^p|^{\frac{1}{p}}$, the positive root of Equation (1) does not always admit a closed form. We give some particular cases for which it can be computed.

- For $p = 1$, the root is $t_X(\{x, y\}) = \frac{f(x) + f(y) + \|x - y\|}{2}$,
- for $p = 2$, it is $t_X(\{x, y\}) = \frac{\sqrt{((f(x) + f(y))^2 + \|x - y\|^2)((f(x) - f(y))^2 + \|x - y\|^2)}}{2\|x - y\|}$,
- for $p = \infty$, the condition reads $\|x - y\| \geq \max\{f(x), f(y)\}$, and the root is $t_X(\{x, y\}) = \frac{\|x - y\|}{2}$. In either case, $t_X(\{x, y\}) = \max\{f(x), f(y), \frac{\|x - y\|}{2}\}$.

Proof. The filtration value of a vertex $x \in X$ is, by definition of the nerve, $t_X(\{x\}) = \inf\{s \in T, \overline{B}_f(x, s) \neq \emptyset\}$. It is equal to $f(x)$.

Also by definition, the filtration value of an edge $\{x, y\}$, with $x, y \in X$ and $x \neq y$, is given by

$$t_X(\{x, y\}) = \inf\{s \in \mathbb{R}, \overline{B}_f(x, s) \cap \overline{B}_f(y, s) \neq \emptyset\}$$

Two cases may occur: the balls $\overline{B}_f(x, t(\{x, y\}))$ and $\overline{B}_f(y, t(\{x, y\}))$ have both positive radius, or one of these is a singleton. In the last case, $t(\{x, y\}) = \max\{f(x), f(y)\}$. In the first case, we have $\|x - y\| = r_x(t) + r_y(t)$, i.e. $\|x - y\| = (t^p - f(x)^p)^{\frac{1}{p}} + (t^p - f(y)^p)^{\frac{1}{p}}$. Notice that Equation (1) admits only one solution since the function $t \mapsto (t^p - f(x)^p)^{\frac{1}{p}} + (t^p - f(y)^p)^{\frac{1}{p}}$ is strictly increasing on $[\max\{f(x), f(y)\}, +\infty)$. \square

We close this subsection by discussing the influence of p on the weighted Čech and Rips filtrations. Let $D_0(\mathcal{N}(\mathcal{V}[X, f, p]))$ be the persistence diagram of the 0th-homology of $\mathcal{N}(\mathcal{V}[X, f, p])$. We say that a point (b, d) of $D_0(\mathcal{V}[X, f, p])$ is non-trivial if $b \neq d$. Let $D_0(\text{Rips}(\mathcal{V}[X, f, p]))$ be the persistence diagram of the 0th-homology of $\text{Rips}(\mathcal{V}[X, f, p])$.

Proposition 3.6. *The number of non-trivial points in $D_0(\mathcal{N}(\mathcal{V}[X, f, p]))$ is non-increasing with respect to $p \in [1, +\infty)$. The same holds for $D_0(\text{Rips}(\mathcal{V}[X, f, p]))$.*

Proof. Since the persistence diagram of the 0th-homology of a filtered simplicial complex only depends on the filtration of its 1-skeleton, it is enough to show the lemma for $D_0(\text{Rips}(\mathcal{V}[X, f, p]))$. As before, denote by t_X the ‘filtration value’ function of $\text{Rips}(\mathcal{V}[X, f, p])$.

The number of points in $D_0(\text{Rips}(\mathcal{V}[X, f, p]))$ is equal to the cardinal of X . Any $p \geq 1$ being fixed, we can pair every $x \in X$ with some edge $\{y, z\} \in \text{Rips}(\mathcal{V}[X, f, p])$ such that the points of $D_0(\text{Rips}(\mathcal{V}[X, f, p]))$ are of the form $(t_X(\{x\}), t_X(\{y, z\}))$.

Notice that the filtration values of the points in X do not depend on p : for all $p \geq 1$ and $x \in X$, $t_X(\{x\}) = f(x)$. Moreover, the filtration values of the edges in $\text{Rips}(\mathcal{V}[X, f, p])$ are non-increasing with respect to p . Indeed, for all $\{y, z\} \in \text{Rips}(\mathcal{V}[X, f, p])$ with $y \neq z$, according to Proposition 3.5, the filtration value $t_X(\{y, z\})$ is either $\max\{f(x), f(y)\}$ if $\|x - y\| \leq |f(x)^p - f(y)^p|^{\frac{1}{p}}$, or is the only positive root of Equation (1) otherwise. Note that the positive root of Equation (1) is greater than $\max\{f(x), f(y)\}$ and decreasing in p . Besides, the term $|f(x)^p - f(y)^p|^{\frac{1}{p}}$ is non-decreasing in p .

These two facts ensure that for every $x \in X$, the point of $D_0(\text{Rips}(\mathcal{V}[X, f, p]))$ created by x has an ordinate which is non-increasing with respect to p . In particular, the number of non-trivial points in $D_0(\text{Rips}(\mathcal{V}[X, f, p]))$ is non-increasing with respect to p . \square

Figure 8 in Subsection 4.4 illustrates the previous proposition in the case of the DTM-filtrations.

Now, consider $k > 0$, and let $D_k(\mathcal{N}(\mathcal{V}[X, f, p]))$ be the persistence diagram of the k th-homology of $\mathcal{N}(\mathcal{V}[X, f, p])$. In this case, one can easily build examples showing that the number of non-trivial points of $D_k(\mathcal{N}(\mathcal{V}[X, f, p]))$ does not have to be non-increasing with respect to p . The same holds for $D_k(\text{Rips}(\mathcal{V}[X, f, p]))$.

4 DTM-filtrations

The results of previous section suggests that in order to construct a weighted Čech filtration $V[X, f]$ that is robust to outliers, it is necessary to choose a function f that depends on X and that is itself robust to outliers. The so-called distance-to-measure function (DTM) satisfies such properties, motivating the introduction of the DTM-filtrations in this section.

4.1 The distance to measure (DTM)

Let μ be a probability measure over $E = \mathbb{R}^d$, and $m \in [0, 1)$ a parameter. For every $x \in \mathbb{R}^d$, let $\delta_{\mu, m}$ be the function defined on E by $\delta_{\mu, m}(x) = \inf\{r \geq 0, \mu(\overline{B}(x, r)) > m\}$.

Definition 4.1. Let $m \in [0, 1[$. The DTM μ of parameter m_0 is the function:

$$d_{\mu, m} : E \longrightarrow \mathbb{R} \\ x \longmapsto \sqrt{\frac{1}{m} \int_0^m \delta_{\mu, t}^2(x) dt}$$

When m is fixed - which is the case in the sequel - and when there is no risk of confusion, we write d_μ instead of $d_{\mu,m}$.

Notice that when $m = 0$, $d_{\mu,m}$ is the distance function to $\text{supp}(\mu)$, the support μ .

Proposition 4.1 ([1], Corollary 3.7). *For every probability measure μ and $m \in [0, 1[$, $d_{\mu,m}$ is 1-Lipschitz.*

A fundamental property of the DTM is its stability with respect to the probability measure μ in the Wasserstein metric. Recall that given two probability measures μ and ν over E , a transport plan between μ and ν is a probability measure π over $E \times E$ whose marginals are μ and ν . The *Wasserstein distance with quadratic cost between μ and ν* is defined as $W_2(\mu, \nu) = \left(\inf_{\pi} \int_{E \times E} \|x - y\|^2 d\pi(x, y) \right)^{\frac{1}{2}}$, where the infimum is taken over all the transport plans. When $\mu = \mu_X$ and $\nu = \mu_Y$ are the empirical measures of the finite point clouds X and Y , i.e the normalized sums of the Dirac measures on the points of X and Y respectively, we write $W_2(X, Y)$ instead of $W_2(\mu_X, \mu_Y)$.

Proposition 4.2 ([1], Theorem 3.5). *Let μ, ν be two probability measures, and $m \in [0, 1[$. Then*

$$\|d_{\mu,m} - d_{\nu,m}\|_\infty \leq m^{-\frac{1}{2}} W_2(\mu, \nu).$$

Notice that for every $x \in E$, $d_\mu(x)$ is not lower than the distance from x to $\text{supp}(\mu)$, the support of μ . This remark, along with the propositions 4.1 and 4.2, are the only properties of the DTM that will be used to prove the results in the sequel of the paper.

In practice, the DTM can be computed. If X is a finite subset of E of cardinal n , we denote by μ_X its empirical measure. Assume that $m = \frac{k_0}{n}$, with k_0 an integer. In this case, d_{μ_X} reformulates as follows: for every $x \in E$,

$$d_{\mu_X}^2(x) = \frac{1}{k_0} \sum_{k=1}^{k_0} \|x - p_k(x)\|^2,$$

where $p_1(x), \dots, p_{k_0}(x)$ are a choice of k_0 -nearest neighbors of x in X .

4.2 DTM-filtrations

In the following, the two parameters $p \in [1, +\infty]$ and $m \in [0, 1)$ are fixed.

Definition 4.2. Let $X \subseteq E$ be a finite point cloud, μ_X the empirical measure of X , and d_{μ_X} the corresponding DTM of parameter m . The weighted Čech filtration $V[X, d_{\mu_X}, p]$, as defined in Section 3, is called the *DTM-filtration associated with the parameters (X, m, p)* . It is denoted by $W[X, m, p]$ or $W[X]$ when there is no risk of confusion. The corresponding persistence module is denoted by $\mathbb{W}[X]$.

Let $\mathcal{W}^t[X, f] = \{\overline{B}_f(x, t)\}_{x \in X}$ denote the cover of $W^t[X, f]$ by the balls $\overline{B}_f(x, t)$ and let $\mathcal{N}(\mathcal{W}^t[X, f])$ be its nerve. The family $\mathcal{N}(\mathcal{W}[X, f]) = (\mathcal{N}(\mathcal{W}^t[X, f]))_{t \geq 0}$ is a filtered simplicial complex, and its persistent (simplicial) homology module is denoted by $\mathbb{W}_{\mathcal{N}}[X, f]$. By the persistent nerve lemma, the persistence modules $\mathbb{W}[X, f]$ and $\mathbb{W}_{\mathcal{N}}[X, f]$ are isomorphic.

As in Definition 3.2, $\text{Rips}(\mathcal{W}^t[X, f])$ denotes the flag complex of $\mathcal{N}(\mathcal{W}^t[X, f])$, and $\text{Rips}(\mathcal{W}[X, f])$ the corresponding filtered simplicial complex.

Example. Consider the point cloud X drawn on the right. It is the union of \tilde{X} and Γ , where \tilde{X} is a 50-sample of the uniform distribution on $[-1, 1]^2 \subseteq \mathbb{R}^2$, and Γ is a 300-sample of the uniform distribution on the unit circle. We consider the weighted Čech filtrations $V[\Gamma, 0]$ and $V[X, 0]$ for $p = 1$, and the DTM-filtration $W[X]$ for $p = 1$ and $m = 0.1$. They are represented in Figure 4.2 for the value $t = 0.3$.





Figure 4: The sets $V^t[\Gamma, 0]$, $V^t[X, 0]$ and $W^t[X]$ for $p = 1$, $m = 0.1$ and $t = 0.3$.

Because of the outliers \tilde{X} , the value of t from which the sets $V^t[X, 0]$ are contractible is small. On the other hand, we observe that the set $W^t[X]$ does not suffer too much from the presence of outliers.

We plot in Figure 5 the persistence diagrams of the persistence modules associated to $\text{Rips}(\mathcal{V}[\Gamma, 0])$, $\text{Rips}(\mathcal{V}[X, 0])$ and $\text{Rips}(\mathcal{W}[X])$ ($p = 1$, $m = 0.1$).




Figure 5: Persistence diagrams of some simplicial filtrations. Points in red (resp. green) represent the persistent homology in dimension 0 (resp. 1).

Observe that the diagrams $D(\text{Rips}(\mathcal{V}[\Gamma, 0]))$ and $D(\text{Rips}(\mathcal{W}[X]))$ appear to be close to each other, while $D(\text{Rips}(\mathcal{V}[X, 0]))$ does not.

Applying the results of Section 3, we immediately obtain the following proposition.

Proposition 4.3. *Let X and Y be two finite subsets of E . Consider the DTM-filtrations $W[X]$ and $W[Y]$. Then*

$$d_i(W[X], W[Y]) \leq m^{-\frac{1}{2}} W_2(X, Y) + 2^{\frac{1}{p}} d_H(X, Y),$$

Proof. By definition of the DTM-filtrations, we have $W[X] = V[X, d_{\mu_X}]$ and $W[Y] = V[Y, d_{\mu_Y}]$. We use the triangle inequality for the interleaving distance:

$$d_i(V[X, d_{\mu_X}], V[Y, d_{\mu_Y}]) \leq \underbrace{d_i(V[X, d_{\mu_X}], V[Y, d_{\mu_X}])}_{(1)} + \underbrace{d_i(V[Y, d_{\mu_X}], V[Y, d_{\mu_Y}])}_{(2)}.$$

(1) : From Proposition 3.3, we have $d_i(V[X, d_{\mu_X}], V[Y, d_{\mu_X}]) \leq (1 + c^p)^{\frac{1}{p}} d_H(X, Y)$, where c is the Lipschitz constant of $d_{X,m}$. According to Proposition 4.1, $c = 1$. We obtain $d_i(V[X, d_{\mu_X}], V[Y, d_{\mu_X}]) \leq 2^{\frac{1}{p}} d_H(X, Y)$.

(2) : From Proposition 3.2, we have $d_i(V[Y, d_{\mu_X}], V[Y, d_{\mu_Y}]) \leq \|d_{\mu_X} - d_{\mu_Y}\|_{\infty}$. According to Proposition 4.2, $\|d_{\mu_X} - d_{\mu_Y}\|_{\infty} \leq m^{-\frac{1}{2}} W_2(X, Y)$. \square

Note that this stability result is worse than the stability of the usual Čech filtrations, which only involves the Hausdorff distance. However, the term $W_2(X, Y)$ is inevitable, as shown in the following example.

Let $E = \mathbb{R}$, $Z = \{0, 1\}$, and $\epsilon > 0$. Pick two finite subsets X and Y of E close to Z in Hausdorff distance, and such that μ_X is close to $\epsilon\delta_0 + (1 - \epsilon)\delta_1$ in Wasserstein distance, and μ_Y close to $(1 - \epsilon)\delta_0 + \epsilon\delta_1$. If $m \geq \epsilon$, then $d_{\mu_Y}(0)$ is close to 0, while $d_{\mu_X}(0)$ is close to $\sqrt{1 - \frac{\epsilon}{m}}$. If $p = 1$, the interleaving distance $d_i(W[X], W[Y])$ is then close to $\sqrt{1 - \frac{\epsilon}{m}}$ (as in Proposition 3.2).

In comparison, the interleaving distance between the usual Čech filtrations is close to 0. In this case, it would be more robust to consider these usual Čech filtrations.

However, in the case where the Hausdorff distance $d_H(X, Y)$ is large, the usual Čech filtrations may be very distant. The DTM-filtrations however may still be close, as we discuss in the next subsection.

4.3 Stability when $p = 1$

We first consider the case $p = 1$, for which the proofs are simpler and results are stronger.

We fix $m \in [0, 1)$. If μ is a probability measure on E with compact support $\text{supp}(\mu)$, we define

$$c(\mu, m) = \sup_{\text{supp}(\mu)} (d_{\mu, m}).$$

If $\mu = \mu_{\Gamma}$ is the empirical measure of a finite set $\Gamma \subseteq E$, we denote it $c(\Gamma, m)$.

Proposition 4.4. *Let μ be a probability measure on E with compact support Γ . Let d_{μ} be the corresponding DTM. Consider a set $X \subseteq E$ such that $\Gamma \subseteq X$. The weighted Čech filtrations $V[\Gamma, d_{\mu}]$ and $V[X, d_{\mu}]$ are $c(\mu, m)$ -interleaved.*

Moreover, if $Y \subseteq E$ is another set such that $\Gamma \subseteq Y$, $V[Y, d_{\mu}]$ and $V[X, d_{\mu}]$ are $c(\mu, m)$ -interleaved.

In particular, if Γ is a finite set, we obtain that $W[\Gamma]$ and $V[X, d_{\mu_{\Gamma}}]$ are $c(\Gamma, m)$ -interleaved.

Proof. Let $c = c(\Gamma, m)$. Since $\Gamma \subseteq X$, we have $V^t[\Gamma, d_{\mu}] \subseteq V^t[X, d_{\mu}]$ for every $t \geq 0$.

Let us show that, for every $t \geq 0$, $V^t[X, d_{\mu}] \subseteq V^{t+c}[\Gamma, d_{\mu}]$. Let $x \in X$, and choose $\gamma \in \Gamma$ a projection of x on the compact set Γ , i.e. one of the closest points to $x \in \Gamma$. By definition of the DTM, we have that $d_{\mu}(x) \geq \|x - \gamma\|$. Together with $d_{\mu}(\gamma) \leq c$, we obtain

$$t + c - d_{\mu}(\gamma) \geq t \geq t - d_{\mu}(x) + \|x - \gamma\|,$$

which means that $B_{d_\mu}(x, t) \subseteq B_{d_\mu}(\gamma, t + c)$. The inclusion $V^t[X, d_\mu] \subseteq V^{t+c}[\Gamma, d_\mu]$ follows.

If Y is another set containing Γ , we obtain $V^t[X, d_\mu] \subseteq V^{t+c}[\Gamma, d_\mu] \subseteq V^{t+c}[Y, d_\mu]$ for every $t \geq 0$. \square

Theorem 4.5. *Consider two finite sets $X, Y \subseteq E$. For every $\Gamma \subseteq X$ and $\Omega \subseteq Y$, we have*

$$d_i(W[X], W[Y]) \leq m^{-\frac{1}{2}}W_2(X, \Gamma) + m^{-\frac{1}{2}}W_2(Y, \Omega) + m^{-\frac{1}{2}}W_2(\Gamma, \Omega) + c(\Gamma, m) + c(\Omega, m).$$

In particular, with $\Omega = Y$, we obtain

$$d_i(W[X], W[\Omega]) \leq m^{-\frac{1}{2}}W_2(X, \Gamma) + m^{-\frac{1}{2}}W_2(\Gamma, \Omega) + c(\Gamma, m) + c(\Omega, m).$$

Proof. Let $d_X = d_{\mu_X}$, $d_Y = d_{\mu_Y}$, $d_\Gamma = d_{\mu_\Gamma}$ and $d_\Omega = d_{\mu_\Omega}$. Recall that the DTM filtrations $W[X]$ and $W[Y]$ are equal to the weighted Čech filtration $V[X, d_X]$ and $V[Y, d_Y]$. We prove the assertion by introducing the following filtrations between $V[X, d_X]$ and $V[Y, d_Y]$:

$$V[X, d_X] \longleftrightarrow V[X, d_\Gamma] \longleftrightarrow V[\Gamma \cup \Omega, d_\Gamma] \longleftrightarrow V[\Gamma \cup \Omega, d_\Omega] \longleftrightarrow V[Y, d_\Omega] \longleftrightarrow V[Y, d_Y].$$

We have:

- $d_i(V[X, d_X], V[X, d_\Gamma]) \leq m^{-\frac{1}{2}}W_2(X, \Gamma)$ (Propositions 4.2 and 3.2),
- $d_i(V[X, d_\Gamma], V[\Gamma \cup \Omega, d_\Gamma]) \leq c(\Gamma, m)$ (Proposition 4.4),
- $d_i(V[\Gamma \cup \Omega, d_\Gamma], V[\Gamma \cup \Omega, d_\Omega]) \leq m^{-\frac{1}{2}}W_2(\Gamma, \Omega)$ (Propositions 4.2 and 3.2),
- $d_i(V[\Gamma \cup \Omega, d_\Omega], V[Y, d_\Omega]) \leq c(\Omega, m)$ (Proposition 4.4),
- $d_i(V[Y, d_\Omega], V[Y, d_Y]) \leq m^{-\frac{1}{2}}W_2(\Omega, Y)$ (Propositions 4.2 and 3.2).

\square

The second inequality of Theorem 4.5 can be seen as an approximation result. Indeed, suppose that Ω is some underlying set of interest, and X is a sample of it with, possibly, noise or outliers. If one can find a subset Γ of X such that X and Γ are close to each other—in the Wasserstein metric—and such that Γ and Ω are also close, then the filtrations $W[X]$ and $W[\Omega]$ are close. Their closeness depends on the constants $c(\Gamma, m)$ and $c(\Omega, m)$.

For any probability measure μ of support $\Gamma \subseteq E$, The constant $c(\mu, m)$ might be seen as a bias term, expressing the behaviour of the DTM over Γ . It relates to the concentration of μ on its support. Recall that a measure μ with support Γ is said to be (a, b) -standard, with $a, b \geq 0$, if for all $x \in \Gamma$ and $r \geq 0$, $\mu(\overline{B}(x, r)) \geq \min\{ar^b, 1\}$. For example, the Hausdorff measure associated to a compact b -dimensional submanifold of E is (a, b) -standard for some $a > 0$. In this case, a simple computation shows that there exists a constant C , depending only on a and b , such that for all $x \in \Gamma$, $d_{\mu, m}(x) \leq Cm^{\frac{1}{b}}$. Therefore, $c(\mu, m) \leq Cm^{\frac{1}{b}}$.

Regarding the first inequality of Theorem 4.5, suppose for the sake of simplicity that one can choose $\Gamma = \Omega$. The bound of Theorem 4.5 then reads

$$d_i(W[X], W[Y]) \leq m^{-\frac{1}{2}}W_2(X, \Gamma) + m^{-\frac{1}{2}}W_2(Y, \Gamma) + 2c(\Gamma, m).$$

Therefore, the DTM-filtrations $W[X]$ and $W[Y]$ are close to each other if μ_X and μ_Y are both close to a common measure μ_Γ . This would be the case if X and Y are noisy samples of Γ . This bound, expressed in terms of Wasserstein distance rather than Hausdorff distance, shows the robustness of the DTM-filtration to outliers.

Notice that, in practice, for finite data sets X, Y and for given Γ and Ω , the constants $c(\Gamma, m)$ and $c(\Omega, m)$ can be explicitly computed, as it amounts to evaluating the DTM on Γ and Ω , as well as the bounds of Theorem 4.5.

Example. Consider the set $X = \tilde{X} \cup \Gamma$ as defined in the example page 12. Figure 6 displays the sets $W^t[X]$, $V^t[X, d_{\mu_\Gamma}]$ and $W^t[\Gamma]$ for the values $p = 1$, $m = 0.1$ and $t = 0.4$ and the persistence diagrams of the corresponding weighted Rips filtrations, illustrating the stability properties of Proposition 4.4 and Theorem 4.5.




Figure 6: Filtrations for $t = 0.4$, and their corresponding persistence diagrams.

The following proposition relates the DTM-filtration to the filtration of E by the sublevel sets of the DTM.

Proposition 4.6. *Let μ be a probability measure on E with compact support K . Let $m \in [0, 1)$ and denote by V the sublevel sets filtration of $d_{\mu, m}$. Consider a finite set $X \subseteq E$. Then*

$$d_i(V, W[X]) \leq c(\mu, m) + 2^{\frac{1}{p}}\epsilon + m^{-\frac{1}{2}}W_2(\mu, \mu_X),$$

with $\epsilon = d_H(K \cup X, X)$.

Proof. First, notice that $V = V[E, d_\mu]$. Indeed, for every $t \geq 0$, $V^t \subseteq V^t[E, d_\mu]$ by the definition of the weighted Čech complex. To prove $V^t[E, d_\mu] \subseteq V^t$, let $x \in V^t[E, d_\mu]$, and $y \in E$ such that $x \in B_{d_\mu}(y, t)$. We have $\|x - y\| \leq t - f(y)$. For d_μ is 1-Lipschitz, we deduce $f(x) \leq f(y) + \|x - y\| \leq f(y) + t - f(y) \leq t$. Hence $x \in V^t$.

Then we compute:

$$\begin{aligned} d_i(V, W[X]) &= d_i(V[E, d_\mu], V[X, d_{\mu_X}]) \\ &\leq d_i(V[E, d_\mu], V[X \cup K, d_\mu]) + d_i(V[X \cup K, d_\mu], V[X, d_\mu]) + d_i(V[X, d_\mu], V[X, d_{\mu_X}]) \\ &\leq c(\mu, m) + 2^{\frac{1}{p}}\epsilon + m^{-\frac{1}{2}}W_2(\mu, \mu_X), \end{aligned}$$

using Proposition 4.2 for the first term, Proposition 3.3 for the second one, and Proposition 3.2 and Proposition 4.4 for the third one. \square

4.4 Stability when $p > 1$

Let now assume that $p > 1$, $m \in [0, 1)$ being still fixed. In this case, stability properties turn out to be more difficult to establish. For small values of t , Lemma 4.10 gives a tight non-additive interleaving between the filtrations. However, for large values of t , the filtrations are poorly interleaved. To overcome this issue we consider stability at the homological level, i.e. between the persistence modules associated to the DTM filtrations.

Let first show why one cannot expect a similar result as Proposition 4.4. Consider the ambient space $E = \mathbb{R}$ and the sets $\Gamma = \{0\}$ and $X = \Gamma \cup \{1\}$. We have $d_{\mu_\Gamma}(1) = 1$ and, for all $t \geq 1$, $W^t[\Gamma] = \overline{B}(0, t)$ and $V^t[X, d_{\mu_\Gamma}] = \overline{B}(0, t) \cup \overline{B}(1, (t^p - 1)^{\frac{1}{p}})$. The sets $V^t[X, d_{\mu_\Gamma}]$ are represented in Figure 7 for $t = 1.5$, $t = 5$ and several values of p .

For $p = 1$, the ball $\overline{B}(1, (t^p - 1)^{\frac{1}{p}})$ is contained in $\overline{B}(0, t)$. Whereas for $p > 1$, the radius $(t^p - 1)^{\frac{1}{p}}$ is asymptotically equal to $t + o_{t \rightarrow +\infty}(\frac{1}{t^{p-1}})$. Therefore, an $\epsilon \geq 0$ for which the ball $\overline{B}(1, (t^p - 1)^{\frac{1}{p}})$ would be included in $\overline{B}(0, t + \epsilon)$ for all $t \geq 0$ should not be lower than $1 = d_H(\Gamma, X)$. Therefore, $d_i(W[\Gamma], V[X, d_{\mu_\Gamma}]) = 1 = d_H(\Gamma, X)$.




Figure 7: Some sets $V^t[X, d_{\mu_\Gamma}]$ for $t = 1.5$ (first row) and $t = 5$ (second row).

Even though the filtrations $W[\Gamma]$ and $V[X, d_{\mu_\Gamma}]$ are distant, the set $V^t[X, d_{\mu_\Gamma}]$ is contractible for all $t \geq 0$, and therefore the interleaving distance between the persistence modules $\mathbb{W}[\Gamma]$ and $\mathbb{V}[X, d_{\mu_\Gamma}]$ is 0.

In general, and in the same spirit as Proposition 4.4, we can obtain a bound on the interleaving distance between the persistence modules $\mathbb{W}[\Gamma]$ and $\mathbb{V}[X, d_{\mu_\Gamma}]$ which does not depend on X —see Proposition 4.7.

For Γ a finite subset of E , we define

$$c(\Gamma, m, p) = \sup_{\Gamma} (d_{\mu_\Gamma, m}) + \kappa(p)t(\Gamma),$$

with $\kappa(p) = 1 - \frac{1}{p}$ and $t(\Gamma)$ is the filtration value of the simplex Γ in $\mathcal{N}(\mathcal{W}[\Gamma])$.

Proposition 4.7. *Let Γ be a finite subset of E , and d_{μ_Γ} be the corresponding DTM of parameter m . Consider a finite set $X \subseteq E$ such that $\Gamma \subseteq X$. The persistence modules $\mathbb{W}[\Gamma]$ and $\mathbb{V}[X, d_{\mu_\Gamma}]$ are $c(\Gamma, m, p)$ -interleaved.*

Moreover, if $Y \subseteq E$ is another finite set such that $\Gamma \subseteq Y$, $\mathbb{V}[X, d_{\mu_\Gamma}]$ and $\mathbb{V}[Y, d_{\mu_\Gamma}]$ are $c(\Gamma, m, p)$ -interleaved.

The proof relies on the following proposition, whose proof is postponed to Subsection 4.5.

Proposition 4.8. *Let $\Gamma \subseteq X \subseteq E$ be finite sets. Consider a map $\pi : X \rightarrow \Gamma$ which maps every $x \in X$ to one of its closest points in Γ . Denote the inclusion map $j : \Gamma \rightarrow X$ and the identity map $v : X \rightarrow X$. Consider the filtered simplicial complexes $\mathcal{N}(\mathcal{V}[X, d_{\mu_\Gamma}])$ and $\mathcal{N}(\mathcal{W}[\Gamma])$.*

For every $t \geq 0$, the map π extends to a simplicial map $\mathcal{N}(\mathcal{V}^t[X, d_{\mu_\Gamma}]) \xrightarrow{\pi^t} \mathcal{N}(\mathcal{W}^{t+c}[\Gamma])$.

We obtain the following diagram between simplicial complexes:

$$\begin{array}{ccccc} & & \xrightarrow{v^t} & & \\ & \swarrow & & \searrow & \\ \mathcal{N}(\mathcal{V}^t[X, d_{\mu_\Gamma}]) & \xrightarrow{\pi^t} & \mathcal{N}(\mathcal{W}^{t+c}[\Gamma]) & \xrightarrow{j^{t+c}} & \mathcal{N}(\mathcal{V}^{t+c}[X, d_{\mu_\Gamma}]) \end{array}$$

where v^t is the simplicial map induced by v and j^{t+c} the one induced by j . This diagram commutes at the homology level, i.e. the induced homomorphism $(j^{t+c} \circ \pi^t)_*$ and i_*^t between simplicial homology groups are equal.

Proof of Proposition 4.7. Choose a map $\pi : X \rightarrow \Gamma$ which maps every $x \in X$ to one of its closest points in Γ . For every $t \geq 0$, let $\pi^t : \mathcal{N}(\mathcal{V}^t[X, d_{\mu_\Gamma}]) \rightarrow \mathcal{N}(\mathcal{W}^{t+c}[\Gamma])$ be the simplicial map given by Proposition 4.8. Denote by $j^t : \mathcal{N}(\mathcal{W}^t[\Gamma]) \rightarrow \mathcal{N}(\mathcal{V}^t[X, d_{\mu_\Gamma}])$ the inclusion map. Let us show that the pair $((\pi_*^t)_{t \geq 0}, (j_*^t)_{t \geq 0})$ define a $c(\Gamma, m, p)$ -interleaving between $\mathbb{W}_{\mathcal{N}}[\Gamma]$ and $\mathbb{V}_{\mathcal{N}}[X, d_{\mu_\Gamma}]$.

Let $t \geq 0$. Denote $c = c(\Gamma, m, p)$, and by $v^t : \mathcal{N}(\mathcal{V}^t[X, d_{\mu_\Gamma}]) \rightarrow \mathcal{N}(\mathcal{V}^{t+c}[X, d_{\mu_\Gamma}])$ and $w^t : \mathcal{N}(\mathcal{W}^t[\Gamma]) \rightarrow \mathcal{N}(\mathcal{W}^{t+c}[\Gamma])$ the inclusion maps. Let us show that the following diagrams commute:

$$\begin{array}{ccc} \mathbb{V}_{\mathcal{N}}^t[X, d_{\mu_\Gamma}] & \xrightarrow{v_*^t} & \mathbb{V}_{\mathcal{N}}^{t+c}[X, d_{\mu_\Gamma}] \\ & \searrow \pi_*^t & \uparrow j_*^{t+c} \\ & & \mathbb{W}_{\mathcal{N}}^{t+c}[\Gamma] \end{array} \quad \begin{array}{ccc} \mathbb{V}_{\mathcal{N}}^t[X, d_{\mu_\Gamma}] & & \\ \uparrow j_*^t & \searrow \pi_*^t & \\ \mathbb{W}_{\mathcal{N}}^t[\Gamma] & \xrightarrow{w_*^t} & \mathbb{W}_{\mathcal{N}}^{t+c}[\Gamma] \end{array}$$

According to Proposition 4.8, the following diagram commutes at the homological level.

$$\begin{array}{ccccc} & & \xrightarrow{v^t} & & \\ & \swarrow & & \searrow & \\ \mathcal{N}(\mathcal{V}^t[X, d_{\mu_\Gamma}]) & \xrightarrow{\pi^t} & \mathcal{N}(\mathcal{W}^{t+c}[\Gamma]) & \xrightarrow{j^{t+c}} & \mathcal{N}(\mathcal{V}^{t+c}[X, d_{\mu_\Gamma}]) \end{array}$$

Hence we obtain the first diagram by applying the simplicial homology functor. Similarly, the second diagram results from the following one, which is clearly commutative since $\pi^{t+c} \circ j^t$ is the identity map.

$$\begin{array}{ccc} \mathcal{N}(\mathcal{W}^t[\Gamma]) & \xrightarrow{j^t} & \mathcal{N}(\mathcal{V}^t[X, d_{\mu_\Gamma}]) & \xrightarrow{\pi^{t+c}} & \mathcal{N}(\mathcal{W}^{t+c}[\Gamma]) \\ & & \searrow & \nearrow & \\ & & & & w^t \end{array}$$

This shows that $\mathbb{W}[\Gamma]$ and $\mathbb{V}[X, d_{\mu_\Gamma}]$ are c -interleaved.

If Y is another set containing Γ , let $\theta : Y \rightarrow \Gamma$ be a projection, and $k : \Gamma \rightarrow Y$ and $u : Y \rightarrow Y$ the inclusion maps. For all $t \geq 0$, define θ^t , k^t and u^t as before. We can compose the previous diagrams to obtain the following:

$$\begin{array}{ccccc} \mathbb{V}_{\mathcal{N}}^t[X, d_{\mu_\Gamma}] & \xrightarrow{v_*^t} & \mathbb{V}_{\mathcal{N}}^{t+c}[X, d_{\mu_\Gamma}] & \xrightarrow{v_*^{t+c}} & \mathbb{V}_{\mathcal{N}}^{t+2c}[X, d_{\mu_\Gamma}] \\ & \searrow \pi_*^t & \uparrow j_*^{t+c} & \searrow \pi_*^{t+c} & \uparrow j_*^{t+2c} \\ & & \mathbb{W}_{\mathcal{N}}^{t+c}[\Gamma] & \xrightarrow{w_*^{t+c}} & \mathbb{W}_{\mathcal{N}}^{t+2c}[\Gamma] \\ & & \downarrow k_*^{t+c} & \nearrow \theta_*^{t+c} & \\ & & \mathbb{V}_{\mathcal{N}}^{t+c}[Y, d_{\mu_\Gamma}] & & \end{array}$$

Since all the triangles commute, so does the following:

$$\begin{array}{ccc} \mathbb{V}_{\mathcal{N}}^t[X, d_{\mu_\Gamma}] & \xrightarrow{v_*^{t+2c}} & \mathbb{V}_{\mathcal{N}}^{t+2c}[X, d_{\mu_\Gamma}] \\ & \searrow k_*^{t+c} \pi_*^t & \nearrow j_*^{t+2c} \theta_*^{t+c} \\ & & \mathbb{V}_{\mathcal{N}}^{t+c}[Y, d_{\mu_\Gamma}] \end{array}$$

We can obtain the same interchanging X and Y . Therefore, by definition, the persistence modules $\mathbb{V}_{\mathcal{N}}[X, d_{\mu_\Gamma}]$ and $\mathbb{V}_{\mathcal{N}}[Y, d_{\mu_\Gamma}]$ are c -interleaved, with the interleaving $((k_*^{t+c} \pi_*^t)_{t \geq 0}, (j_*^{t+c} \theta_*^t)_{t \geq 0})$. \square

Theorem 4.9. *Consider two finite sets $X, Y \subseteq E$. For every $\Gamma \subseteq X$ and $\Omega \subseteq Y$, we have*

$$d_i(\mathbb{W}[X], \mathbb{W}[Y]) \leq m^{-\frac{1}{2}} W_2(X, \Gamma) + m^{-\frac{1}{2}} W_2(Y, \Omega) + m^{-\frac{1}{2}} W_2(\Gamma, \Omega) + c(\Gamma, m, p) + c(\Omega, m, p).$$

In particular, with $\Omega = Y$, we obtain

$$d_i(\mathbb{W}[X], \mathbb{W}[\Gamma]) \leq m^{-\frac{1}{2}} W_2(X, \Gamma) + m^{-\frac{1}{2}} W_2(\Gamma, \Omega) + c(\Gamma, m, p) + c(\Omega, m, p).$$

Proof. The proof is the same as Theorem 4.5, using Proposition 4.7 instead of Proposition 4.4. \square

Notice that when $p = 1$, the constant $c(\Gamma, m, p)$ is equal to the constant $c(\Gamma, m)$ defined in Subsection 4.3, and we recover Theorem 4.5.

As an illustration of these results, we represent in Figure 8 the persistence diagrams associated to $\text{Rips}(\mathcal{W}[X])$ for several values of p . The point cloud X is the one defined in the example page 12. Observe that, as stated in Proposition 3.6, the number of red points (homology in dimension 0) is non-increasing with respect to p .




Figure 8: Persistence diagrams of the simplicial filtrations $\text{Rips}(\mathcal{W}[X])$ for several values of p .

4.5 Proof of Proposition 4.8

We first state the following lemma, which gives a (non-additive) interleaving between the filtrations $W[\Gamma]$ and $V[X, d_{\mu_\Gamma}]$.

Lemma 4.10. *Let X, Γ and π be as defined in Proposition 4.8. Let $x \in X$, and $\gamma = \pi(x)$. For every $t \geq 0$,*

$$\overline{B}_f(x, t) \subseteq \overline{B}_f(\gamma, 2^{1-\frac{1}{p}}t + d_{\mu_\Gamma}(\gamma)).$$

Let $\phi : t \mapsto 2^{1-\frac{1}{p}}t + \sup_\Gamma d_{\mu_\Gamma}$. Then for every $t \geq 0$,

$$W^t[\Gamma] \subseteq V^t[X, d_{\mu_\Gamma}] \subseteq W^{\phi(t)}[\Gamma].$$

Proof. Define $d = d_{\mu_\Gamma}(\gamma)$. Let $u \in E$. Remind that

$$\begin{cases} u \in \overline{B}_f(\gamma, t) & \iff t \geq (\|u - \gamma\|^p + d_{\mu_\Gamma}(\gamma)^p)^{\frac{1}{p}}, \\ u \in \overline{B}_f(x, t) & \iff t \geq (\|u - x\|^p + d_{\mu_\Gamma}(x)^p)^{\frac{1}{p}}. \end{cases}$$

We shall only use

$$\begin{cases} u \in \overline{B}_f(\gamma, t) & \iff t \geq \|u - \gamma\| + d, \\ u \in \overline{B}_f(x, t) & \implies t \geq (\|u - x\|^p + \|x - \gamma\|^p)^{\frac{1}{p}}. \end{cases}$$

Let $u \in \overline{B}_f(x, t)$. Let us prove that $u \in \overline{B}_f(\gamma, 2^{1-\frac{1}{p}}t + d)$. If $\|u - \gamma\| \leq \|\gamma - x\|$, then $t \geq \|u - \gamma\|$, and we deduce $u \in \overline{B}_f(\gamma, t + d) \subseteq \overline{B}_f(\gamma, 2^{1-\frac{1}{p}}t + d)$.

Else, we have $\|u - \gamma\| \geq \|\gamma - x\|$. Consider the line segment $[\gamma, u]$ and the sphere $S(\gamma, \|\gamma - x\|)$ of center γ and radius $\|\gamma - x\|$. The intersection $S(\gamma, \|\gamma - x\|) \cap [\gamma, u]$ is a singleton. Call its element x' . The situation is represented in Figure 9.




Figure 9: Definition of the point x' .

We have $\|u - x'\| \leq \|u - x\|$ and $\|\gamma - x'\| = \|\gamma - x\|$. Therefore

$$\left(\|u - x'\|^p + \|x' - \gamma\|^p\right)^{\frac{1}{p}} \leq \left(\|u - x\|^p + \|x - \gamma\|^p\right)^{\frac{1}{p}}.$$

We also have $\|\gamma - u\| = \|\gamma - x'\| + \|x' - u\|$ and $\left(\|u - x\|^p + \|x - \gamma\|^p\right)^{\frac{1}{p}} \leq t$. Thus it follows from the last inequality that

$$\left(\|u - x'\|^p + (\|u - \gamma\| - \|u - x'\|)^p\right)^{\frac{1}{p}} \leq t.$$

The left-hand term of this inequality is not lower than $2^{\frac{1}{p}-1}\|u - \gamma\|$. Indeed, consider the function $s \mapsto \left(s^p + (\|u - \gamma\| - s)^p\right)^{\frac{1}{p}}$ defined for $s \in [0, \|u - \gamma\|]$. One shows directly, by computing its derivative, that its minimum is $2^{\frac{1}{p}-1}\|u - \gamma\|$, attained at $s = \frac{\|u - \gamma\|}{2}$.

We deduce that $2^{\frac{1}{p}-1}\|u - \gamma\| \leq t$, and $\|u - \gamma\| \leq 2^{1-\frac{1}{p}}t$. Thus $u \in \overline{B}_f(\gamma, 2^{1-\frac{1}{p}}t + d)$. \square

Notice that the previous lemma gives a tight bound, as we can see with the following example. Consider set $\Gamma = \{0\} \subset \mathbb{R}$, $L > 0$, and $X = \Gamma \cup \{x\}$ with $x = \frac{L}{2}$. Let $m < \frac{1}{2}$, and $f = d_{\mu_\Gamma}$, which is the function distance to Γ . For all $t \geq 2^{\frac{1}{p}-1}L$, we have $L \in \overline{B}_f(x, t)$. Indeed, $r_x(2^{\frac{1}{p}-1}L) = \left((2^{\frac{1}{p}-1}L)^p - \left(\frac{L}{2}\right)^p\right)^{\frac{1}{p}} = \frac{L}{2}$. In comparison, for every $t < L$, $L \notin \overline{B}_f(0, t)$.

Following this example, we can find a lower bound on the interleaving distance between the persistence modules $\mathbb{W}[\Gamma]$ and $\mathbb{V}[X, d_{\mu_\Gamma}]$. Consider $L > 0$, the set $\Gamma = \{0, 2L\} \subset \mathbb{R}$, $x = \frac{L}{2}$, and $X = \Gamma \cup \{x, 2L - x\}$. Let $m < \frac{1}{2}$, and $f = d_{\mu_\Gamma}$. The persistence diagram of the 0th-homology of $W[\Gamma]$ consists of two points, $(0, +\infty)$ and $(0, L)$. Regarding $V[X, f]$, the point of finite ordinate in the persistence diagram of its 0th-homology is $(0, 2^{\frac{1}{p}-1}L)$. Indeed, for $t = 2^{\frac{1}{p}-1}L$, we have $L \in \overline{B}_f(x, t)$ and $L \in \overline{B}_f(L - x, t)$, hence the set $V^t[X, d_{\mu_\Gamma}]$ is connected. We deduce that these persistent modules are at most $(1 - 2^{\frac{1}{p}-1})L$ -interleaved.

In comparison, the upper bound we prove in Proposition 4.7 is $(1 - \frac{1}{p})L$.

We now prove the proposition stated in the previous subsection.

Proof of Proposition 4.8. Let $t_X : \mathcal{P}(X) \rightarrow \mathbb{R}^+$ be the ‘filtration value’ function of $\mathcal{N}(\mathcal{V}[X, d_{\mu_\Gamma}])$, which maps every simplex $\sigma \subseteq X$ to $\inf\{t \geq 0, \sigma \in \mathcal{N}(\mathcal{V}^t[X, d_{\mu_\Gamma}])\}$. Note that the restriction of t_X to the set $\mathcal{P}(\Gamma) \subseteq \mathcal{P}(X)$ is equal to the ‘filtration value’ function of $\mathcal{N}(\mathcal{W}[\Gamma])$. Denote $f = d_{\mu_\Gamma}$ and $c = c(\Gamma, m, p)$.

Let $t \geq 0$. We decompose the proof in two steps.

First step: $\pi^t : \mathcal{N}(\mathcal{V}^t[X, d_{\mu_\Gamma}]) \rightarrow \mathcal{N}(\mathcal{W}^{t+c}[\Gamma])$ is simplicial.

Let $\sigma \in \mathcal{N}(\mathcal{V}^t[X, d_{\mu_\Gamma}])$, and $\tau = \pi^t(\sigma)$. We have to show that $\tau \in \mathcal{N}(\mathcal{W}^{t+c}[\Gamma])$.

If $t_X(\tau) \leq t$, i.e. $\tau \in \mathcal{N}(\mathcal{V}^t[X, d_{\mu_\Gamma}])$, then $\tau \in \mathcal{N}(\mathcal{W}^t[\Gamma]) \subseteq \mathcal{N}(\mathcal{W}^{t+c}[\Gamma])$.

Else, we have $t_X(\tau) \geq t \geq t_X(\sigma)$. Let $x \in \sigma$, and define $\gamma = \pi(x)$. According to Lemma 4.10, $\overline{B}_f(x, t_X(\sigma)) \subseteq \overline{B}_f(\gamma, 2^{1-\frac{1}{p}}t_X(\sigma) + d_{\mu_\Gamma}(\gamma))$. Hence

$$\bigcap_{x \in \sigma} \overline{B}_f(x, t_X(\sigma)) \subseteq \bigcap_{\gamma \in \tau} \overline{B}_f(\gamma, 2^{1-\frac{1}{p}}t_X(\sigma) + \sup_{\tau} d_{\mu_\Gamma}).$$

Since $\bigcap_{x \in \sigma} \overline{B}_f(x, t_X(\sigma)) \neq \emptyset$, we obtain $\bigcap_{\gamma \in \tau} \overline{B}_f(\gamma, 2^{1-\frac{1}{p}}t_X(\sigma) + \sup_{\tau} d_{\mu_\Gamma}) \neq \emptyset$. By definition of the nerve, this means that $t_X(\tau) \leq 2^{1-\frac{1}{p}}t_X(\sigma) + \sup_{\tau} d_{\mu_\Gamma}$. We conclude by observing that

$$\begin{aligned} t_X(\tau) &\leq 2^{1-\frac{1}{p}}t_X(\sigma) + \sup_{\tau} d_{\mu_\Gamma} = t_X(\sigma) + (2^{1-\frac{1}{p}} - 1)t_X(\sigma) + \sup_{\tau} d_{\mu_\Gamma} \\ &\leq t + (2^{1-\frac{1}{p}} - 1)t_X(\Gamma) + \sup_{\Gamma} d_{\mu_\Gamma} \\ &\leq t + (1 - \frac{1}{p})t_X(\Gamma) + \sup_{\Gamma} d_{\mu_\Gamma} \\ &= t + c \end{aligned}$$

where, for the second line, we used $t_X(\sigma) \leq t_X(\tau) \leq t_X(\Gamma)$ and $2^{1-\frac{1}{p}} - 1 \leq 1 - \frac{1}{p}$ (Lemma B.1).

Notice that, still in the case $t_X(\tau) \geq t \geq t_X(\sigma)$, the same argument gives $\bigcap_{y \in \sigma \cup \tau} \overline{B}_f(y, 2^{1-\frac{1}{p}}t(\sigma) + \sup_{\tau} d_{\mu_\Gamma}) \neq \emptyset$, and $t_X(\sigma \cup \tau) \leq t + c$.

Second step: the induced homomorphism $(j^{t+c} \circ \pi^t)_*$ and i_*^t are equal.

According to the last paragraph, we have the following diagram of simplicial maps:

$$\begin{array}{ccccc} & & \xrightarrow{v^t} & & \\ & \searrow & & \nearrow & \\ \mathcal{N}(\mathcal{V}^t[X, d_{\mu_\Gamma}]) & \xrightarrow{\pi^t} & \mathcal{N}(\mathcal{W}^{t+c}[\Gamma]) & \xrightarrow{j^{t+c}} & \mathcal{N}(\mathcal{V}^{t+c}[X, d_{\mu_\Gamma}]) \end{array}$$

In order to prove that it commutes at the homological level, we distinguish the cases $t \leq t_X(\Gamma)$ and $t \geq t_X(\Gamma)$.

First, assume that $t \leq t_X(\Gamma)$. Let us show that the simplicial maps $j^{t+c} \circ \pi^t$ and i^t are contiguous. Let $\sigma \in \mathcal{N}(\mathcal{V}^t[X, d_{\mu_\Gamma}])$, and define $\tau = \pi^t(\sigma)$. We have seen in the last paragraph that $t_X(\sigma \cup \tau) \leq t + c$. Hence the simplex $\sigma \cup \tau$ is in the simplicial complex $\mathcal{N}(\mathcal{V}^{t+c}[X, d_{\mu_\Gamma}])$. This proves that the simplicial maps $j^{t+c} \circ \pi^t$ and v^t are contiguous, therefore they induce equal homomorphisms at homological level.

Secondly, assume that $t \geq t_X(\Gamma)$. In the remainder of the proof, we say that a homology group $H_n(\cdot)$ is trivial if it is of rank 0 when $n > 0$, or if it is of rank 1 when $n = 0$. We say that a homomorphism between homology groups $H_n(\cdot) \rightarrow H_n(\cdot)$ is trivial if the homomorphism is of rank 0 when $n > 0$, or of it is of rank 1 when $n = 0$.

The homomorphism $(j^{t+c} \circ \pi^t)_*$ is trivial. Indeed, $\mathcal{N}(\mathcal{W}^{t+c}[\Gamma])$ is the whole simplex on Γ , hence its homology groups are trivial, and $(j^{t+c} \circ \pi^t)_*$ factorizes through these. Let us show that v_*^t also is trivial.

We introduce an intermediate set between $V^t[X, d_{\mu_\Gamma}]$ and $V^{t+c}[X, d_{\mu_\Gamma}]$:

$$\begin{cases} V^t[X, d_{\mu_\Gamma}] &= \bigcup_{x \in X \setminus \Gamma} \overline{B}_f(x, t) \cup \bigcup_{\gamma \in \Gamma} \overline{B}_f(\gamma, t), \\ \tilde{V}^t &:= \bigcup_{x \in X \setminus \Gamma} \overline{B}_f(x, t) \cup \bigcup_{\gamma \in \Gamma} \overline{B}_f(\gamma, t+c), \\ V^{t+c}[X, d_{\mu_\Gamma}] &= \bigcup_{x \in X \setminus \Gamma} \overline{B}_f(x, t+c) \cup \bigcup_{\gamma \in \Gamma} \overline{B}_f(\gamma, t+c). \end{cases}$$

Let $O_\Gamma \in \bigcap_{\gamma \in \Gamma} \overline{B}_f(\gamma, t_X(\Gamma))$. Since $t \geq t_X(\Gamma)$, we have $O_\Gamma \in \tilde{V}^t$. Let us show that \tilde{V}^t is star-shaped around O_Γ .

Let $x \in X$ and consider $\gamma = \pi(x) \in \Gamma$. We first prove that $\overline{B}_f(x, t) \cup \overline{B}_f(\gamma, t+c)$ is star-shaped around O_Γ . Let $y \in \overline{B}_f(x, t)$. We have to show that the line segment $[y, O_\Gamma]$ is a subset of $\overline{B}_f(x, t) \cup \overline{B}_f(\gamma, t+c)$. Let D be the affine line going through y and O_Γ , and denote by q the orthogonal projection on D . We have $[y, O_\Gamma] \subset [y, q(x)] \cup [q(x), O_\Gamma]$. The first line segment $[y, q(x)]$ is a subset of $\overline{B}_f(x, t)$. Regarding the second line segment $[q(x), O_\Gamma]$, let us show that $q(x) \in \overline{B}_f(\gamma, t+c)$, and $[q(x), O_\Gamma] \subset \overline{B}_f(\gamma, t+c)$ will follow. The situation is pictured in Figure 10.




Figure 10: Construction of an intermediate set \tilde{V}^t .

Denote by $q(\gamma)$ the orthogonal projection of γ on D . According to Lemma B.2,

$$\|\gamma - q(x)\|^2 \leq \|x - \gamma\|^2 + \|x - q(x)\|(2\|\gamma - q(\gamma)\| - \|x - q(x)\|).$$

Let $d = \|x - q(x)\|$. Since $d = \|x - q(x)\| \leq (t^p - d_{\mu_\Gamma}(x)^p)^{\frac{1}{p}} \leq (t^p - \|x - \gamma\|^p)^{\frac{1}{p}}$, we have $\|x - \gamma\| \leq (t^p - d^p)^{\frac{1}{p}}$. Moreover, $\|\gamma - q(\gamma)\| \leq \|\gamma - O_\Gamma\| \leq t_X(\Gamma)$. The last inequality then gives

$$\|\gamma - q(x)\|^2 \leq (t^p - d^p)^{\frac{2}{p}} + d(2t_X(\Gamma) - d).$$

According to Lemma B.3, we obtain that $\|\gamma - q(x)\|$ is not greater than $t + \kappa(p)t_X(\Gamma)$. Therefore, we have the inequality

$$((t + \kappa(p)t_X(\Gamma) + f(\gamma))^p - f(\gamma)^p)^{\frac{1}{p}} \geq (t + \kappa(p)t_X(\Gamma) + f(\gamma)) - f(\gamma) \geq \|\gamma - q(x)\|,$$

and we deduce $q(x) \in \overline{B}_f(\gamma, t + \kappa(p)t_X(\Gamma) + f(\gamma)) \subset \overline{B}_f(\gamma, t+c)$.

In conclusion, $[y, O_\Gamma] \subset \overline{B}_f(x, t) \cup \overline{B}_f(\gamma, t + c)$. This being true for every $y \in \overline{B}_f(x, t)$, and obviously true for $y \in \overline{B}_f(\gamma, t + c)$, we deduce that $\overline{B}_f(x, t) \cup \overline{B}_f(\gamma, t + c)$ is star-shaped around O_Γ . Finally, since $O_\Gamma \in \bigcap_{\gamma \in \Gamma} \overline{B}_f(\gamma, t_X(\Gamma))$, we have that \widetilde{V}^t is star-shaped around O_Γ .

Let $u^t : V^t[X, d_{\mu_\Gamma}] \rightarrow V^{t+c}[X, d_{\mu_\Gamma}]$ be the inclusion map between sets. Consider the diagram of inclusions:

$$V^t[X, d_{\mu_\Gamma}] \longrightarrow \widetilde{V}^t \longrightarrow V^{t+c}[X, d_{\mu_\Gamma}].$$

$\xrightarrow{\quad u^t \quad}$

Applying the singular homology functor, we obtain

$$\mathbb{V}^t[X, d_{\mu_\Gamma}] \longrightarrow H_*(\widetilde{V}^t) \longrightarrow \mathbb{V}^{t+c}[X, d_{\mu_\Gamma}].$$

$\xrightarrow{\quad u_*^t \quad}$

Since \widetilde{V}^t is star-shaped, $H_*(\widetilde{V}^t)$ is trivial, and so is u_*^t . Moreover, according to the persistent nerve lemma ([11, Lemma 3.4]), we have the following commutative diagram

$$\begin{array}{ccc} \mathbb{V}_{\mathcal{N}}^t[X, d_{\mu_\Gamma}] & \xrightarrow{v_*^t} & \mathbb{V}_{\mathcal{N}}^{t+c}[X, d_{\mu_\Gamma}] \\ \wr \uparrow & & \wr \uparrow \\ \mathbb{V}^t[X, d_{\mu_\Gamma}] & \xrightarrow{u_*^t} & \mathbb{V}^{t+c}[X, d_{\mu_\Gamma}] \end{array}$$

where the vertical arrows are isomorphisms. Therefore v_*^t is trivial, as claimed. \square

5 Conclusion

In this paper we have introduced the DTM-filtrations that depend on a parameter $p \geq 1$. This new family of filtrations extends the filtration introduced in [3] that corresponds to the case $p = 2$.

The established stability properties are, as far as we know, of a new type: the closeness of two DTM-filtrations associated to two data sets relies on the existence of a well-sampled underlying object that approximate both data sets in the Wasserstein metric. This makes the DTM filtrations robust to outliers. Even though large values of p lead to persistence diagrams with less points in the 0th homology, the choice of $p = 1$ gives the strongest stability results. When $p > 1$, the interleaving bound is less significant since it involves the diameter of the underlying object but, the obtained bound is consistent with the case $p = 1$ as it converges to the bound for $p = 1$ as p goes to 1.

It is interesting to notice that the proofs rely on only a few properties of the DTM. As a consequence, the results should extend to other weight functions, such that the DTM with a parameter different from 2, or kernel density estimators. Some variants concerning the radius functions in the weighted Čech filtration, are also worth considering. The analysis shows that their asymptotic behaviour should look like the one of the case $p = 1$. In the same spirit as in [12, 3] where sparse-weighted Rips filtrations were considered, it would also be interesting to consider sparse versions of the DTM-filtrations and to study their stability properties.

Last, the obtained stability results, depending on the choice of underlying sets, open the way to the statistical analysis of the persistence diagrams of the DTM-filtrations, a problem that will be addressed in a further work.

References

- [1] F. Chazal, D. Cohen-Steiner, and Q. Mérigot. Geometric inference for probability measures. *Journal on Found. of Comp. Mathematics*, 11(6):733–751, 2011.
- [2] Leonidas Guibas, Dmitriy Morozov, and Quentin Mérigot. Witnessed k-distance. *Discrete & Computational Geometry*, 49(1):22–45, 2013.
- [3] Mickaël Buchet, Frédéric Chazal, Steve Y Oudot, and Donald R Sheehy. Efficient and robust persistent homology for measures. *Computational Geometry*, 58:70–96, 2016.
- [4] Frédéric Chazal, Vin De Silva, and Steve Oudot. Persistence stability for geometric complexes. *Geometriae Dedicata*, 173(1):193–214, 2014.
- [5] J. Phillips, B. Wang, and Y Zheng. Geometric inference on kernel density estimates. In *Proc. 31st Annu. Sympos. Comput. Geom (SoCG 2015)*, pages 857–871, 2015.
- [6] Mickaël Buchet. *Topological inference from measures*. PhD thesis, Paris 11, 2014.
- [7] Fujitsu Laboratories. Estimating the degradation state of old bridges—fujitsu supports ever-increasing bridge inspection tasks with ai technology. *Fujitsu Journal*, March 2018.
- [8] Lee M Seversky, Shelby Davis, and Matthew Berger. On time-series topological data analysis: New data and opportunities. In *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition Workshops*, pages 59–67, 2016.
- [9] Yuhei Umeda. Time series classification via topological data analysis. *Transactions of the Japanese Society for Artificial Intelligence*, 32(3):D–G72_1, 2017.
- [10] Frédéric Chazal, Vin de Silva, Marc Glisse, and Steve Oudot. *The Structure and Stability of Persistence Modules*. SpringerBriefs in Mathematics, 2016.
- [11] Frédéric Chazal and Steve Yann Oudot. Towards persistence-based reconstruction in euclidean spaces. In *Proceedings of the twenty-fourth annual symposium on Computational geometry*, SCG '08, pages 232–241, New York, NY, USA, 2008. ACM.
- [12] Donald R. Sheehy. Linear-size approximations to the Vietoris-Rips filtration. *Discrete & Computational Geometry*, 49(4):778–796, 2013.

A Supplementary results for Section 3

Lemma A.1. *Let c, ϵ and x be non-negative real numbers, and $t \geq a$. Define $\alpha = (1+c^p)^{\frac{1}{p}}$ and $k = \epsilon\alpha$. Then $t + k \geq a + c\epsilon$, and*

$$((t + k)^p - (x + c\epsilon)^p)^{\frac{1}{p}} - (t^p - x^p)^{\frac{1}{p}} \geq \epsilon$$

Proof. Let $\mathcal{D} = \{(t, x), t \geq x \geq 0\} \subseteq \mathbb{R}^2$. Let us find the minimum of

$$\begin{aligned} \Phi : \mathcal{D} &\longrightarrow \mathbb{R} \\ (t, x) &\longmapsto \left((t + \alpha\epsilon)^p - (x + c\epsilon)^p \right)^{\frac{1}{p}} - (t^p - x^p)^{\frac{1}{p}} \end{aligned}$$

An $x > 0$ being fixed, we study $\phi : t \mapsto \Phi(t, x)$ on the interval $(x, +\infty)$. Its derivative is

$$\phi'(t) = \frac{(t + \alpha\epsilon)^{p-1}}{\left((t + \alpha\epsilon)^p - (x + c\epsilon)^p \right)^{1-\frac{1}{p}}} - \frac{t^{p-1}}{(t^p - x^p)^{1-\frac{1}{p}}}$$

We solve:

$$\begin{aligned} \phi'(t) = 0 &\iff (t + \alpha\epsilon)^{p-1}(t^p - x^p)^{1-\frac{1}{p}} = t^{p-1}\left((t + \alpha\epsilon)^p - (x + c\epsilon)^p \right)^{1-\frac{1}{p}} \\ &\iff \underbrace{(t + \alpha\epsilon)^p}_{\text{}} (t^p - x^p) = \underbrace{t^p}_{\text{}} \left((t + \alpha\epsilon)^p - (x + c\epsilon)^p \right) \\ &\iff (t + \alpha\epsilon)^p x^p = t^p (x + c\epsilon)^p \\ &\iff \frac{t + \alpha\epsilon}{t} = \frac{x + c\epsilon}{x} \\ &\iff t = \frac{\alpha}{c}x \end{aligned}$$

We obtain the second line by raising the equality to the power of $\frac{p}{p-1}$. Hence the derivative of ϕ vanishes only at $t = \frac{\alpha}{c}x$. Together with $\lim_{+\infty} \phi = +\infty$, we deduce that ϕ attains its minimum at $t = x$ or $t = \frac{\alpha}{c}x$.

Let us show that $\phi\left(\frac{\alpha}{c}x\right) = \epsilon$.

$$\begin{aligned} \phi\left(\frac{\alpha}{c}x\right) = \Phi\left(\frac{\alpha}{c}x, x\right) &= \left(\left(\frac{\alpha}{c}x + \alpha\epsilon\right)^p - (x + c\epsilon)^p \right)^{\frac{1}{p}} - \left(\left(\frac{\alpha}{c}x\right)^p - x^p \right)^{\frac{1}{p}} \\ &= \left(\left(\frac{\alpha}{c}\right)^p (x + c\epsilon)^p - (x + c\epsilon)^p \right)^{\frac{1}{p}} - x \left(\left(\frac{\alpha}{c}\right)^p - 1 \right)^{\frac{1}{p}} \\ &= (x + c\epsilon) \left(\left(\frac{\alpha}{c}\right)^p - 1 \right)^{\frac{1}{p}} - x \left(\left(\frac{\alpha}{c}\right)^p - 1 \right)^{\frac{1}{p}} \\ &= c\epsilon \left(\left(\frac{\alpha}{c}\right)^p - 1 \right)^{\frac{1}{p}} \end{aligned}$$

Using $\alpha = (1 + c^p)^{\frac{1}{p}}$, one obtains that $c \left(\left(\frac{\alpha}{c}\right)^p - 1 \right)^{\frac{1}{p}} = 1$. Therefore, $\phi\left(\frac{\alpha}{c}x\right) = \epsilon$.

Secondly, consider Φ on the interval $\{(x, x), x \geq 0\}$.

The function $t \mapsto \Phi(x, x) = \left((x + \alpha\epsilon)^p - (x + c\epsilon)^p \right)^{\frac{1}{p}}$ is increasing. Its minimum is $\Phi(0, 0) = \left((\alpha\epsilon)^p - (c\epsilon)^p \right)^{\frac{1}{p}} = \epsilon(\alpha^p - c^p)^{\frac{1}{p}} = \epsilon$.

In conclusion, on every interval $(x, +\infty) \times \{x\} \subseteq \mathcal{D}$, Φ admits ϵ as a minimum. Therefore, ϵ is the minimum of Φ on \mathcal{D} . \square

B Supplementary results for Section 4

Lemma B.1. For all $p \geq 1$, $2^{1-\frac{1}{p}} - 1 \leq 1 - \frac{1}{p}$.

Proof. The convexity property of the function $x \mapsto 2^x$ gives, for all $x \in [0, 1]$, $2^x \leq x + 1$. Hence $2^{1-\frac{1}{p}} - 1 \leq 1 - \frac{1}{p}$. \square

Lemma B.2. Let $\gamma, x \in E$, D an affine line, and $q(\gamma), q(x)$ the projections of γ and x on D . Then

$$\|\gamma - q(x)\|^2 \leq \|x - \gamma\|^2 + \|x - q(x)\| (2\|\gamma - q(\gamma)\| - \|x - q(x)\|)$$

Proof. We first study the case where γ, x and D lie in the same affine plane. If γ and x are on opposite sides of D , the result is obvious. Otherwise, the points $\gamma, x, q(\gamma)$ and $q(x)$ form a right trapezoid (see Figure 11).




Figure 11: The points $\gamma, x, q(\gamma)$ and $q(x)$ form a right trapezoid.

Using the Pythagorean theorem on the orthogonal vectors $\gamma - q(\gamma)$ and $q(\gamma) - q(x)$, and on $(\gamma - q(\gamma)) - (x - q(x))$ and $q(\gamma) - q(x)$, we obtain

$$\begin{cases} \|\gamma - q(\gamma)\|^2 + \|q(\gamma) - q(x)\|^2 = \|\gamma - q(x)\|^2, \\ \|(\gamma - q(\gamma)) - (x - q(x))\|^2 + \|q(\gamma) - q(x)\|^2 = \|\gamma - x\|^2. \end{cases}$$

Using that $\|(\gamma - q(\gamma)) - (x - q(x))\| = |\|\gamma - q(\gamma)\| - \|x - q(x)\||$, the second equality rephrases as $\|q(\gamma) - q(x)\|^2 = \|\gamma - x\|^2 - (\|\gamma - q(\gamma)\| - \|x - q(x)\|)^2$. Combining these two equalities give

$$\begin{aligned} \|\gamma - q(x)\|^2 &= \|\gamma - q(\gamma)\|^2 + \|q(\gamma) - q(x)\|^2 \\ &= \|\gamma - q(\gamma)\|^2 + \|\gamma - x\|^2 - (\|\gamma - q(\gamma)\| - \|x - q(x)\|)^2 \\ &= \|\gamma - x\|^2 + \|x - q(x)\|(2\|\gamma - q(\gamma)\| - \|x - q(x)\|) \end{aligned}$$

Now, if γ, x and D do not lie in the same affine plane, denote by P the affine plane containing D and x . Let $\tilde{\gamma}$ the point of P such that $\|\gamma - q(\gamma)\| = \|\tilde{\gamma} - q(\gamma)\|$ and $\|\gamma - q(x)\| = \|\tilde{\gamma} - q(x)\|$. Using the previous result on $\tilde{\gamma}$ and the inequality $\|\gamma - x\| \geq \|\tilde{\gamma} - x\|$, we obtain the result. \square

Lemma B.3. *Let $a, b, d \geq 0$ such that $b \leq a$ and $d \leq a$. Then*

$$(a^p - d^p)^{\frac{2}{p}} + d(2b - d) \leq (a + \kappa b)^2,$$

with $\kappa = 1 - \frac{1}{p}$.

Proof. The equation being homogeneous with respect to a , it is enough to show that

$$(1 - d^p)^{\frac{2}{p}} + d(2b - d) \leq (1 + \kappa b)^2$$

with $b \leq 1$ and $d \leq 1$. We actually show that $(1 - d^p)^{\frac{2}{p}} + d(2b - d) \leq 1 + 2\kappa b$

Note that this is true when $d \leq \kappa$. Indeed, $(1 - d^p)^{\frac{2}{p}} + d(2b - d) \leq 1 + 2db \leq 1 + 2\kappa b$. Now, notice that it is enough to show the inequality for $b = 1$. Indeed, it is equivalent

to $(1 - d^p)^{\frac{2}{p}} - 1 - d^2 \leq 2\kappa b - 2db = 2b(\kappa - d)$. For every $d \geq \kappa$, the right-hand side of this inequality is nonpositive, hence the worst case happens when $b = 1$. What is left to show is the following: $\forall d \in [\kappa, 1]$,

$$(1 - d^p)^{\frac{2}{p}} + d(2 - d) \leq 1 + 2\kappa.$$

The function $x \mapsto (1 - x)^{\frac{1}{p}}$ being concave on $[0, 1]$, we have $(1 - x)^{\frac{1}{p}} \leq 1 - \frac{1}{p}x$ for all $x \in [0, 1]$. Therefore, $(1 - d^p)^{\frac{1}{p}} \leq 1 - \frac{1}{p}d^p$. Consider the function

$$\phi : d \mapsto (1 - \frac{1}{p}d^p)^2 + d(2 - d).$$

Let us show that $\forall d \in [0, 1]$, $\phi(d) \leq 1 + 2\kappa$.

This inequality is obvious for $d = 0$. It is also the case for $d = 1$, since we obtain $(1 - \frac{1}{p}d^p)^2 + d(2 - d) = (1 - \frac{1}{p})^2 + 1 = \kappa^2 + 1$. In the interval $[0, 1]$, the derivative of ϕ is $\phi'(d) = \frac{2}{p}d^{2p-1} - 2d^{p-1} - 2d + 2$. Let d_* be such that $\phi'(d_*) = 0$. Multiplying $\phi'(d_*)$ by $\frac{d_*}{2}$ gives the relation $\frac{1}{p}d_*^{2p} - d_*^p - d_*^2 + d_* = 0$. Subtracting this equality in $\phi(d_*)$ gives $\phi(d_*) = 1 - (\frac{1}{p} - \frac{1}{p^2})d_*^{2p} + (1 - \frac{2}{p})d_*^p + d_*$. We shall show that the following function ψ , defined for all $d \in [0, 1]$, is not greater than $1 + 2\kappa$.

$$\psi : d \mapsto 1 - \frac{1}{p}(1 - \frac{1}{p})d^{2p} + (1 - \frac{2}{p})d^p + d.$$

We consider the cases $p \geq 2$ and $p \leq 2$ separately. In each case, $1 - \frac{1}{p} \geq 0$. Assume that $p \geq 2$. Then $d^p \leq 1$ and $1 - \frac{2}{p} \geq 0$. Therefore $(1 - \frac{2}{p})d^p \leq 1 - \frac{2}{p}$, and we obtain

$$\begin{aligned} \psi(d) &\leq 1 + (1 - \frac{2}{p})d^p + d \\ &\leq 1 + (1 - \frac{2}{p})d + d \\ &= 1 + 2(1 - \frac{1}{p}) \end{aligned}$$

Now assume that $p \leq 2$. We have the following inequality: $d - d^p \leq p - 1$. Indeed, by considering its derivative, one shows that the application $d \mapsto d - d^p$ is maximum for $d = p^{-\frac{1}{p-1}}$, for which

$$\begin{aligned} d - d^p &= d(1 - d^{p-1}) = p^{-\frac{1}{p-1}}(1 - p^{-1}) \\ &= p^{-\frac{1}{p-1}-1}(p - 1) \\ &= p^{-\frac{p}{p-1}}(p - 1) \leq p - 1. \end{aligned}$$

Using $(\frac{2}{p} - 1) \geq 0$ and $d^p \geq d - (p - 1)$, we obtain $(\frac{2}{p} - 1)d^p \geq (\frac{2}{p} - 1)d - (\frac{2}{p} - 1)(p - 1)$. Going back to $\psi(d)$, we have

$$\begin{aligned} \psi(d) &= 1 - \frac{1}{p}(1 - \frac{1}{p})d^{2p} - (\frac{2}{p} - 1)d^p + d \\ &\leq 1 - \frac{1}{p}(1 - \frac{1}{p})d^{2p} - (\frac{2}{p} - 1)d + (\frac{2}{p} - 1)(p - 1) + d \\ &= 1 - \frac{1}{p}(1 - \frac{1}{p})d^{2p} + (2 - \frac{2}{p})d + (\frac{2}{p} - 1)(p - 1). \end{aligned}$$

Let us verify that $d \mapsto 1 - \frac{1}{p}(1 - \frac{1}{p})d^{2p} + 2(1 - \frac{1}{p})d + (\frac{2}{p} - 1)(p - 1)$ is increasing. Its derivative is

$$\begin{aligned} -2p\frac{1}{p}(1 - \frac{1}{p})d^{2p-1} + 2(1 - \frac{1}{p}) &\geq -2p\frac{1}{p}(1 - \frac{1}{p}) + 2(1 - \frac{1}{p}) \\ &= 0 \end{aligned}$$

We deduce that $\psi(d) \leq \psi(1)$ for all $d \in [0, 1]$. The value $\psi(1)$ is $1 - \frac{1}{p}(1 - \frac{1}{p}) + 2(1 - \frac{1}{p}) + (\frac{2}{p} - 1)(p - 1)$. Moreover, we have $-\frac{1}{p}(1 - \frac{1}{p}) + (\frac{2}{p} - 1)(p - 1) \leq 0$. Indeed, $-\frac{1}{p}(1 - \frac{1}{p}) + (\frac{2}{p} - 1)(p - 1) = -\frac{(p-1)^3}{p^2}$. Therefore $\psi(1) \leq 1 + 2(1 - \frac{1}{p})$. □