



Mitochondrial protection by the mixed muscarinic/ $\alpha 1$ ligand ANAVEX2-73, a tetrahydrofuran derivative, in $A\beta 25-35$ peptide-injected mice, a nontransgenic Alzheimer's disease model

Valentine Lahmy, Romain Long, Didier Morin, Vanessa Villard, Tanguy Tm Maurice

► To cite this version:

Valentine Lahmy, Romain Long, Didier Morin, Vanessa Villard, Tanguy Tm Maurice. Mitochondrial protection by the mixed muscarinic/ $\alpha 1$ ligand ANAVEX2-73, a tetrahydrofuran derivative, in $A\beta 25-35$ peptide-injected mice, a nontransgenic Alzheimer's disease model. *Frontiers in Cellular Neuroscience*, 2015, 8, 10.3389/fncel.2014.00463 . hal-01922607

HAL Id: hal-01922607

<https://hal.science/hal-01922607>

Submitted on 27 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Mitochondrial protection by the mixed muscarinic/ σ_1 ligand ANAVEX2-73, a tetrahydrofuran derivative, in A β_{25-35} peptide-injected mice, a nontransgenic Alzheimer's disease model

Valentine Lahmy^{1,2}, Romain Long^{3,4}, Didier Morin^{3,4}, Vanessa Villard² and Tangui Maurice^{1,2*}

¹ Inserm U 710, University of Montpellier 2, Montpellier, France

² Amylgen, Montferrier-sur-Lez, France

³ Inserm U 955, Team 03, Créteil, France

⁴ Faculty of Medicine, Université Paris-Est, Unité Mixte de Recherche S955, Université Paris-Est Créteil Val-de-Marne, Créteil, France

Edited by:

Fabio Blandini, National Institute of Neurology C. Mondino Foundation, Italy

Reviewed by:

Rafael Linden, Federal University of Rio de Janeiro, Brazil

Mohan Pabba, Center for Addiction and Mental Health, Canada

*Correspondence:

Tangui Maurice, Inserm U 710, University of Montpellier, cc 105, place Eugène Bataillon, 34095 Montpellier cedex 5, France
e-mail: maurice@univ-montp2.fr

Alzheimer's disease (AD), the most prevalent dementia in the elderly, is characterized by progressive synaptic and neuronal loss. Mitochondrial dysfunctions have been consistently reported as an early event in AD and appear before A β deposition and memory decline. In order to define a new neuroprotectant strategy in AD targeting mitochondrial alterations, we develop tetrahydro-N,N-dimethyl-2,2-diphenyl-3-furanmethanamine (ANAVEX2-73, AE37), a mixed muscarinic receptor ligand and a sigma-1 receptor (σ_1 R) agonist. We previously reported that ANAVEX2-73 shows anti-amnesic and neuroprotective activities in mice injected intracerebroventricular (ICV) with oligomeric amyloid- β_{25-35} peptide (A β_{25-35}). The σ_1 R is present at mitochondria-associated endoplasmic reticulum (ER) membranes, where it acts as a sensor/modulator of ER stress responses and local Ca²⁺ exchanges with the mitochondria. We therefore evaluated the effect of ANAVEX2-73 and PRE-084, a reference σ_1 R agonist, on preservation of mitochondrial integrity in A β_{25-35} -injected mice. In isolated mitochondria from hippocampus preparations of A β_{25-35} injected animals, we measured respiration rates, complex activities, lipid peroxidation, Bax/Bcl-2 ratios and cytochrome *c* release into the cytosol. Five days after A β_{25-35} injection, mitochondrial respiration in mouse hippocampus was altered. ANAVEX2-73 (0.01–1 mg/kg IP) restored normal respiration and PRE-084 (0.5–1 mg/kg IP) increased respiration rates. Both compounds prevented A β_{25-35} -induced increases in lipid peroxidation levels, Bax/Bcl-2 ratio and cytochrome *c* release into the cytosol, all indicators of increased toxicity. ANAVEX2-73 and PRE-084 efficiently prevented the mitochondrial respiratory dysfunction and resulting oxidative stress and apoptosis. The σ_1 R, targeted selectively or non-selectively, therefore appears as a valuable target for protection against mitochondrial damages in AD.

Keywords: Alzheimer's disease, mitochondrial damages, ANAVEX2-73, sigma-1 receptor, cytoprotection

INTRODUCTION

Alzheimer's disease (AD), the most common form of dementia in the elderly is defined histologically by the presence of the two hallmarks: extracellular senile plaques constituted with amyloid- β (A β) proteins and intracellular inclusions of hyperphosphorylated microtubule-associated Tau protein, referred as neurofibrillary tangles (Selkoe, 2004). Clinically, the disease is characterized by progressive cognitive decline associated with synaptic and neuronal loss in regions critical for memory process, e.g., hippocampus, entorhinal and frontal cortex (Crews and Masliah, 2010). Mitochondria, which are the main energy provider in the cell, are particularly enriched in synapses where energy is critical for synaptic transmission (Verstreken

et al., 2005). Mitochondrial dysfunctions have been consistently reported as an early event in AD pathophysiology and appear before A β deposition and memory deficits in AD patients and transgenic mice (Maurer et al., 2000; Caspersen et al., 2005; Mosconi et al., 2008). Indeed, during AD pathogenesis, A β oligomers accumulate in mitochondria, resulting in disrupted energy metabolism, increased oxidative stress and apoptosis (Lustbader et al., 2004; Caspersen et al., 2005; Manczak et al., 2006). Studies in AD patients and transgenic mice have reported consistent decreases in tricarboxylic acid cycle enzymes and cytochrome *c* oxidase activities (Yates et al., 1990; Maurer et al., 2000; Caspersen et al., 2005; Leuner et al., 2007). Moreover, studies in isolated mitochondria from transgenic mice brains

or in isolated mitochondria exposed to A β oligomers showed decreased respiration rates and ATP production and increased oxidative stress, suggesting A β -related defects in mitochondrial respiratory chain (Casley et al., 2002a; Aleardi et al., 2005; Caspersen et al., 2005; Clementi et al., 2005; Leuner et al., 2007). Disruption in mitochondrial respiratory chain results in increased production of oxidative stress by mitochondria, a key element in AD physiopathology (Smith et al., 1996; Leuner et al., 2012). One other aspect of mitochondrial failure is the opening of mitochondrial permeability transition pore (mPTP) which can be triggered by different effectors, e.g., mitochondrial Ca²⁺ overload and oxidative stress (Kroemer and Reed, 2000). mPTP opening could also be triggered by the translocation to the mitochondrial outer membrane of the cytosolic proapoptotic members of the Bcl-2 family proteins. Among them is the proapoptotic protein Bax which, in proapoptotic conditions, translocates to mitochondria, dimerizes and constitutes a pore in the mitochondrial membrane and/or contributes to the formation of mPTP (Jürgensmeier et al., 1998; Marzo et al., 1998; Narita et al., 1998). Bax activity could be antagonized by heterodimerization with antiapoptotic Bcl-2 family proteins, including Bcl-2 itself (Hanada et al., 1995; Yin et al., 1995; Yang et al., 1997). mPTP opening allows liberation of mitochondrial protein, such as cytochrome c, which promotes apoptosis, binding to the apoptosis protease activation factor (APAF-1) and forming a complex indicated as “apoptosome” (Liu et al., 1996; Yang et al., 1997).

In the present study, we used the intracerebroventricular (ICV) injection of A β _{25–35} fragment in oligomeric form in mice as a rapid, standardized pharmacological model of AD toxicity, suitable to investigate the impact of mitochondrial alteration. Indeed, ICV A β _{25–35} injection results in neuroinflammation and reactive gliosis, pro-apoptotic caspase activity, oxidative stress, reduction in the number of neurons measured in hippocampal pyramidal cell layers, loss of cholinergic neurons, and memory deficits (Maurice et al., 1996; Delobette et al., 1997; Meunier et al., 2006; Villard et al., 2009, 2011; Zussy et al., 2011). *In vitro* evidences have been accumulated suggesting that the peptide also altered mitochondrial physiology, after direct application on isolated mitochondria or in mitochondria isolated from cultured neurons exposed to A β _{25–35} (Canevari et al., 1999; Casley et al., 2002a,b; Aleardi et al., 2005; Clementi et al., 2005; Dong et al., 2010; Ren et al., 2011). Isolated mitochondria from rat brain subjected to subchronic ICV injection of A β _{25–35} or A β _{1–40} produced significantly more H₂O₂ compared to control mice after three days of treatment, suggesting mitochondrial dysfunction in this model (Kaminsky and Kosenko, 2008). However, to our knowledge, the functionality of isolated mitochondria has never been explored *ex vivo* after ICV administration of A β _{25–35} in rodents.

Tetrahydro-N,N-dimethyl-2,2-diphenyl-3-furanmethanamine hydrochloride (ANAVEX2-73, AE37) is a novel ligand, acting at muscarinic acetylcholine (mAChR) and σ_1 receptors (σ_1 R) with affinities in the low micromolar range (Espallergues et al., 2007). We previously reported that this compound had anti-amnesic and neuroprotective potential in the AD model

in mice induced by the ICV injection of A β _{25–35} peptide. In particular, ANAVEX2-73 attenuated cellular loss, A β _{1–42} seeding, Tau hyperphosphorylation and learning and memory deficits observed several days after A β _{25–35} injection (Villard et al., 2009, 2011; Lahmy et al., 2013). The drug acts synergistically on its two main targets, as an antagonist of M2 mAChR and as an agonist of σ_1 R. The σ_1 R is an endoplasmic reticulum (ER)-resident molecular chaperone (Hayashi and Su, 2007), also expressed at the mitochondrial membrane (Klouz et al., 2002), particularly at mitochondria-associated ER membranes (MAM). Its activity regulates Ca²⁺ exchange from ER to mitochondria (Hayashi and Su, 2007), known to be critical for mitochondrial bioenergetics (Cárdenas et al., 2010). Moreover, the σ_1 R is involved in regulation of the mitochondrial membrane potential, reactive oxygen species (ROS) production and apoptosis (Tsai et al., 2009; Meunier and Hayashi, 2010) and σ_1 R agonists have been shown to protect mitochondrial function *in vivo* against ischemic damage (Klouz et al., 2008; Tagashira et al., 2013).

In this study, we analyzed whether the neuroprotective activity of ANAVEX2-73 involves mitochondrial protection. We analyzed the respiratory activity in mitochondria isolated from the mouse hippocampus, five days after ICV injection of oligomerized A β _{25–35}. We measured respiration rates of freshly extracted mitochondria and the activity of the respiratory chain complexes I to IV. We then analyzed biochemical markers of mitochondrial damage, including Bax, Bcl-2 protein expression and cytochrome c release from the mitochondria into the cytosol. PRE-084 was used as a reference σ_1 R agonist.

MATERIAL AND METHODS

ANIMALS

Male Swiss OF-1 mice (Depré, St Doulchard, France), aged 7–9 weeks and weighing 32 \pm 2 g were used. They were housed in plastic cages in groups with free access to food and water, except during behavioral experiments. They were kept in a regulated environment (23 \pm 1°C, 50–60% humidity) under a 12 h light/dark cycle (light on at 8:00 am). Behavioral experiments were carried out between 10:00 am and 4:00 pm, in an experimental room within the animal facility. All animal procedures were conducted in adherence with the 2010/63 EU Directive.

DRUGS AND ADMINISTRATION PROCEDURES

The amyloid- β [25–35] (A β _{25–35}) and scrambled A β _{25–35} (Sc. A β) peptides were purchased from Genepep (Saint Jean-de-Védas, France). They were solubilized in sterile distilled water at a concentration of 3 mg/ml and stored at –20°C until use. Before injection, peptides were aggregated by incubation at 37°C for 4 days (Maurice et al., 1996). This procedure is currently standardized in the laboratory and previous analyses using electronic microscopy, IR spectroscopy, and photon correlation spectroscopy techniques showed that the aggregated A β _{25–35} solution is mainly composed of a mixture of soluble oligomeric amyloid species, which sizes extended from 52.8 to 295.3 nm (98%) (Zussy et al., 2011, 2013). They were administered intracerebroventricularly (ICV) in a final volume of 3 μ l per mouse, as previously described (Maurice et al., 1996;

Meunier et al., 2006). Tetrahydro-*N,N*-dimethyl-2,2-diphenyl-3-furanmethanamine hydrochloride (ANAVEX2-73, AE37) was provided by Dr Alexandre Vamvakides (Anavex Life Science, Pallini, Greece). PRE-084 was a gift from Dr Tsung-Ping Su (NIDA/NIH, Baltimore, MD, USA). 2-(3,4-Dichlorophenyl)-*N*-(2-dimethylaminoethyl)-*N*-methylethanamine dihydrobromide (BD1047) was from Sigma-Aldrich (St Quentin-Fallavier, France). Drugs were solubilized in physiological saline at the concentration of 5 mg/ml and then diluted to the appropriate concentrations. They were injected intraperitoneally (IP) in a volume of 100 μ l/20 g body weight. Drugs were injected once, 20 min before the A β _{25–35} peptide, also injected once. Animals were used 1, 3, 5, or 7 days after injections, as indicated in the figure legends.

LIPID PEROXIDATION

Mice were killed by decapitation and brains were rapidly removed, the hippocampus dissected out, weighed, and kept in liquid nitrogen until assayed. After thawing, the hippocampus was homogenized in cold methanol (1/10 w/v), centrifuged at 1,000 g for 5 min and the supernatant collected. Homogenate was added to a solution containing 1 mM FeSO₄, 0.25 M H₂SO₄, 1 mM xylenol orange, and incubated for 30 min at room temperature. Absorbance was measured at 580 nm (A₅₈₀₁), and 10 ml of 1 mM cumene hydroperoxide (CHP; Sigma-Aldrich) was added to the sample and incubated for 30 min at room temperature, to determine the maximal oxidation level. Absorbance was measured at 580 nm (A₅₈₀₂). The level of lipid peroxidation was determined as CHP equivalents according to: CHP eq. $1/4 A_{5801}/A_{5802}$ [CHP (nmol)] dilution, and expressed as CHP eq. *per wet tissue weight*.

RESPIRATION RATES

Five days after peptide injection, mice were sacrificed, and their hippocampus rapidly removed on ice and homogenized in 10 ml of ice-cold homogenization buffer (220 mM mannitol, 70 mM sucrose, 10 mM 2-[4-(2-hydroxyethyl)piperazin-1-yl]ethanesulfonic acid (HEPES; Sigma-Aldrich), 2 mM ethylene glycol tetraacetic acid (EGTA; Sigma-Aldrich), pH 7.4) using a Teflon potter homogenizer. Homogenate was centrifuged at 1,000 g for 5 min. The supernatant was then centrifuged at 10,000 g for 10 min. Pellet was collected and resuspended in 50 μ l of ice-cold isolation buffer (220 mM mannitol, 70 mM sucrose, 10 mM HEPES, 0.01 mM EGTA, pH 7.4). Mitochondrial protein content was determined by spectrophotometry. O₂ consumption was measured with a Clarke type oxygen electrode (Hansatech, Cergy, France). Mitochondria (0.8 mg/ml) were loaded in the chamber filled up with respiration buffer maintained at 30°C (50 mM sucrose, 100 mM KCl, 10 mM HEPES, 5 mM KH₂PO₄, 0.5 mM MgCl₂, pH 7.4). State 2 respiration was measured after addition of 5 mM pyruvate-malate and state 3 respiration was triggered by addition of 0.5 mM adenosine diphosphate (ADP; Sigma-Aldrich). State 4 respiration was then measured after addition of 4 μ M carboxyatractyloside (CAT; Sigma-Aldrich), a blocker of the adenosine triphosphate (ATP)-ADP carrier, and uncoupled state with 1 μ M tyrphostine (uncoupling agent). The mean of two measures of respiration rates for the four states were calculated for each sample. The control

ratio was calculated as the ratio between state 3 and state 4 respiratory rates.

In a series of experiment, the effects of direct application of the drugs (ANAVEX2-73 and PRE-084, in the 10^{−8} to 10^{−4} M concentration range) were examined. Isolated mitochondria (0.8 mg/ml) were loaded in the chamber. Drugs were applied and after 5 min, the respiration was examined.

COMPLEX ACTIVITIES

Five days after peptide injection, mice were sacrificed, and their hippocampus rapidly removed on ice. Mitochondria were isolated in the same way as for respiration rates. Mitochondrial protein content was determined using a BCA assay (Pierce Biotechnology, Rockford, IL, USA). Mitochondria were frozen/thawed four times to break mitochondrial membranes. Complex activities were measured at 30°C, using a Jasco V-530 spectrophotometer (Jasco, Nantes, France). Complex activities are calculated as nmol/min/mg of proteins and expressed as percentage of Sc. A β -treated group data.

Complex I (NADH ubiquinone reductase)

The oxidation of nicotinamide adenine dinucleotide (NADH; Sigma-Aldrich) by complex I was measured using decylubiquinone (DUQ; Sigma-Aldrich), an ubiquinone analog, as electron acceptor. 200 μ M NADH and 100 μ M DUQ were added to the assay medium (25 mM KH₂PO₄, 5 mM MgCl₂, 250 μ M KCN, 1 mg/ml bovine serum albumin (BSA; Sigma-Aldrich), pH 7.4) in a final volume of 1.5 ml. Enzyme activity was measured by initiating the reaction with 0.15 mg of mitochondrial protein. The decrease in absorbance due to NADH oxidation was measured at 340 nm.

Complex II (succinate dehydrogenase)

The activity of complex II was measured indirectly by record of the reduction of 2,6-dichlorophenolindo-phenol (DCIP; Sigma-Aldrich) following the oxidation of succinate. 6 mM succinate was incubated with 0.05 mg of mitochondrial proteins for 5 min at 30°C in the assay medium (10 mM KH₂PO₄, 2 mM ethylenediaminetetraacetic acid (EDTA; Sigma-Aldrich), 2 μ M rotenone, 1 mg/ml BSA) in a final volume of 1.5 ml. 100 μ M DUQ and 80 μ M DCIP were then added and the decrease in absorbance at 600 nm, corresponding to the reduction of DCIP, was measured.

Complex III (ubiquinol cytochrome c reductase)

The oxidation of DUQH₂ by complex III was determined using cytochrome *c*(III) as an electron acceptor. 0.25 mg of mitochondrial protein was added in the assay medium (10 mM KH₂PO₄, 2 mM EDTA, 1 mg/ml BSA, 2 μ M rotenone, 250 μ M KCN) in a final volume of 1.5 ml. The reaction was started with the addition of 100 μ M DUQH₂ (Sigma-Aldrich) and the increase in absorbance at 550 nm, corresponding to reduction of cytochrome *c* was measured. Because cytochrome *c*(III) could also be reduced by DUQH₂ independently of complex III, the reduction of cytochrome *c*(III) was measured in the presence of 1 μ M antimycin (an inhibitor of complex III activity) and deducted from the total reduction measured before.

Complex IV (cytochrome c oxidase)

0.025 mg of mitochondrial protein were incubated for 1 min at 30°C in the assay medium (10 mM KH_2PO_4 , 10 mM EDTA, 2 mM MgCl_2 , 1 mg/ml BSA) in a final volume of 1.5 ml. The reaction was started when 33 μM of cytochrome *c*(II) was added to the assay medium and the decreased in absorbance at 550 nm, corresponding to oxidation of cytochrome *c*(II), was measured.

WESTERN BLOTTING

Mice were sacrificed at indicated days after injections and the hippocampus rapidly dissected on ice and kept at -80°C until used. For cytochrome *c* release experiments, the hippocampus were homogenized with a glass Dounce homogenizer in ice-cold homogenization buffer (250 μM sucrose, 10 mM HEPES, pH 7.4), including a protease and phosphatase inhibitor cocktail (Roche Diagnostics, Meylan, France) in a final volume of 250 μl . Homogenates were centrifuged at 600 g for 5 min and the supernatant collected and centrifuged again at 10,300 g for 20 min. The supernatant, corresponding to the cytosolic fraction (C), and the pellet, corresponding to the crude mitochondrial fraction (M), were separated. The mitochondrial fraction was resuspended in 50 μl of ice-cold isolation buffer (250 mM mannitol, 5 mM HEPES, 0.5 mM EGTA, pH 7.4).

For Bax and Bcl-2 experiments, the hippocampus were homogenized by sonication in an ice-cold lysis buffer (125 mM Tris HCl pH 6.8, 4% sodium dodecyl sulfate (SDS; Sigma-Aldrich), 20% glycerol) including a protease and phosphatase inhibitor cocktail (Roche Diagnostics, Meylan, France). Homogenates were heated at 70°C for 10 min and centrifuged at 16,000 g for 30 min. The supernatant was collected.

For all samples, protein concentration was determined using a BCA assay (Pierce Biotechnology, Rockford, IL, USA) according to the manufacturer's instructions.

Proteins, 20 μg per lane, were resolved on a 12% SDS-polyacrylamide gel and transferred to a polyvinylidene fluoride (PVDF) membrane (GE Healthcare, Orsay, France). After 1 h blocking in 5% non-fat dry milk in a 20 mM Tris-buffered saline pH 7.5 buffer containing 0.1% Tween-20 (TBS-T), membranes were incubated overnight at 4°C with the following primary antibodies: mouse anti-cytochrome *c* (CytC, dilution 1/1000; BioLegend, San Diego, CA, USA), mouse anti-oxphos-complex IV subunit I (Oxphos, 1/1000; Invitrogen Life Technologies, St Aubin, France), rabbit anti-Bax (1/2000; Cell Signaling Technology, Ozyme, St Quentin-en-Yvelines, France), mouse anti-Bcl-2 antibody (1/1000; Santa Cruz Biotechnology; Heidelberg, Germany). After brief washes, membranes were incubated for 1 h at room temperature with corresponding secondary antibody: goat anti-mouse IgG peroxidase conjugate (1/2000; Sigma-Aldrich) or goat anti-rabbit IgG peroxidase conjugate (1/2000; Sigma-Aldrich). The immunoreactive bands were visualized with the enhanced chemiluminescence reagent (ECL, Millipore, Molsheim, France) using an Odyssey[®] Fc fluorescent imaging system (Li-Cor, Eurobio, Courtaboeuf, France). Then, membranes were stripped using the Restore Western Blot

Stripping Buffer (Pierce Biotechnology) and reprobed with anti- β -tubulin antibody (1/2000) (Sigma-Aldrich). The intensity of peroxidase activity was quantified using the Odyssey[®] Fc software (Li-Cor).

STATISTICAL ANALYSES

Data were expressed as mean \pm S.E.M. The number of animals is indicated in the figure legends. Data were analyzed using one-way ANOVA (*F* values), followed by the Dunnett's *post-hoc* multiple comparison test. For reading clarity, ANOVA values were all reported in the figure legends. The level of statistical significance was $p < 0.05$.

RESULTS

ICV administration of $\text{A}\beta_{25-35}$ in mice induced a robust (+50–70%) increase in lipid peroxidation level in the hippocampus of the animals, starting at 5–7 days after injection (Figure 1A). The IP pretreatment with ANAVEX2-73, in the 0.01–1 mg/kg dose-range dose-dependently prevented the $\text{A}\beta_{25-35}$ -induced increase in lipid peroxidation with highly significant effect measured at 0.3 and 1 mg/kg (Figure 1B). The reference $\sigma_1\text{R}$ agonist PRE-084 similarly blocked $\text{A}\beta_{25-35}$ -induced increase in lipid peroxidation at 0.5 and 1 mg/kg (Figure 1C). Note that both drug effects at the highest doses were completely blocked by the reference $\sigma_1\text{R}$ antagonist BD1047, confirming the $\sigma_1\text{R}$ selectivity (Figures 1B,C). These initial observations confirmed that ICV injection of $\text{A}\beta_{25-35}$ induces a marked oxidative stress in mice and that the two compounds act as anti-oxidant drugs. The prominent role of mitochondrial dysfunction in ROS generation and oxidative stress led us to analyze the impact of the drug treatments on mitochondrial respiration.

O_2 consumption was analyzed in mice, five days after Sc. $\text{A}\beta$ and $\text{A}\beta_{25-35}$ injection, and following ANAVEX2-73 or PRE-084 treatment (Figure 2). The $\text{A}\beta_{25-35}$ ICV treatment failed to affect state 2 or state 4 O_2 consumption (Figure 2A), but significantly altered by -23% and -22% the state 3 or uncoupled state O_2 consumption, respectively (Figure 2B). As a consequence the peptide injection significantly altered the respiratory control ratio by -17% (Figure 2C). The ANAVEX2-73 treatment (0.3 mg/kg IP) significantly prevented the $\text{A}\beta_{25-35}$ -induced decreases in O_2 consumption (Figure 2B). However, since the drug increased in fact significantly O_2 consumption at all states (Figure 2A), there was no significant impact on the respiratory control ratio (Figure 2C). A similar profile was observed with PRE-084 (0.5 mg/kg IP) in $\text{A}\beta_{25-35}$ -treated mice (Figures 2A,B) but the increase in state 3 O_2 consumption was more marked resulting in a restoration of the respiratory control ratio value, *i.e.*, a significant prevention of the $\text{A}\beta_{25-35}$ -induced alteration (Figure 2C). Note that neither ANAVEX2-73 nor PRE-084 affected O_2 consumption in Sc. $\text{A}\beta$ -treated animals. In addition, we analyzed whether the drugs have a direct effect on mitochondrial respiration. The drugs were both applied at increasing concentrations, from 10^{-8} M to 10^{-4} M, on isolated mitochondria and respiration analyzed. State 3 O_2 consumption and respiratory control ratio are presented (Figures 3A,B). The drugs




FIGURE 1 | ANAVEX2-73 and PRE-084 prevented oxidative stress in Aβ₂₅₋₃₅-treated mice. Mice were administered ICV with Sc. Aβ or Aβ₂₅₋₃₅ peptide (9 nmol) and sacrificed after 1, 3, 5 or 7 days for lipid peroxidation measures (A). $n = 6$ per group, * $p < 0.05$, *** $p < 0.001$ vs. the (Sc. Aβ + V)-treated group at the same timepoint; t -test. Mice were then administered IP with (B) ANAVEX2-73 (0.01–1 mg/kg) or (C) PRE-084 (0.5–1 mg/kg) 20 min

before Aβ₂₅₋₃₅ (9 nmol). In two groups, BD1047 (10 mg/kg) was administered simultaneously with the highest dose of each agonist. Mice were sacrificed after 7 days for lipid peroxidation measures. One-way ANOVA: $F_{(7,70)} = 22.5$, $p < 0.0001$, $n = 6$ –14 per group in (B); $F_{(4,69)} = 31.9$, $p < 0.0001$, $n = 6$ –22 in (C). ** $p < 0.01$, *** $p < 0.001$ vs. the (Sc. Aβ+V)-treated group; ### $p < 0.001$ vs. the (Aβ₂₅₋₃₅+V)-treated group; Dunnett's test.

affected both parameters only at the highest concentrations tested.

We analyzed the nature of the Aβ₂₅₋₃₅-induced alteration of respiration (Figure 4). The measure of each complex activity showed that only the complex IV activity was significantly decreased by −14% (Figure 4A). The ANAVEX2-73 treatment prevented the Aβ₂₅₋₃₅-induced decrease in complex IV activity (Figure 4B).

Among the different biochemical markers that can be examined to evidence the damage resulting from impaired mitochondrial functionality, we first addressed cytochrome *c* release from the mitochondrial membrane into the cytosol (Figure 5A). The time-course analysis showed that significant cytochrome *c*

release can be evidenced at day 5 after the peptide injection (Figure 5B). At this timepoint, both the ANAVEX2-73 treatment (Figure 5C) and PRE-084 treatment (Figure 5D) completely prevented the Aβ₂₅₋₃₅-induced cytochrome *c* release in the mouse hippocampus.

We then measured Bax and Bcl-2 protein expression, by a western blot approach, in order to confirm that the impact of the treatment on mitochondrial damage readily transfer into anti-apoptosis protection (Figure 6). The Aβ₂₅₋₃₅ treatment resulted, between 1 to 7 days after injection, in a progressive but non-significant increase in Bax level in the hippocampus (Figure 6A), accompanied by a trend to diminution of Bcl-2 level (Figure 6B). As a consequence, the Bax/Bcl-2 ratio increased, and significantly




FIGURE 2 | ANAVEX2-73 (AN2-73) and PRE-084 (PRE) protected against Aβ₂₅₋₃₅-induced alteration of mitochondrial respiration in mice. Mice were treated with ANAVEX2-73 (0.3 mg/kg IP), PRE-084 (0.5 mg/kg IP) or vehicle, before injection of Sc. Aβ or Aβ₂₅₋₃₅ (9 nmol ICV). **(A–C)** Mitochondria (0.8 mg/ml) were loaded in the chamber with appropriate buffer at 30°C. State 2 respiration was activated by addition of pyruvate-malate (5 mM). State 3 respiration was induced with ADP. State 4 respiration was provoked by the addition of carboxyatractylide (CAT), a blocker of ATP-ADP

carrier. Finally, addition of tyrophostine, an uncoupling agent, activated the uncoupled respiration. The respiratory control ratio is the state 3/state 4 ratio. $n = 4-8$ per group, $F_{(5,38)} = 7.11$, $p < 0.0001$ for state 2 and $F_{(5,38)} = 8.43$, $p < 0.0001$ for state 4 in **(A)**; $F_{(5,38)} = 13.1$, $p < 0.0001$ for state 3 and $F_{(5,38)} = 8.43$, $p < 0.0001$ for state 3 in **(B)**; $F_{(5,38)} = 2.58$, $p < 0.05$ in **(C)**. * $p < 0.05$, ** $p < 0.01$ vs. Sc. Aβ/Veh; # $p < 0.05$, ## $p < 0.01$, ### $p < 0.001$ vs. Aβ₂₅₋₃₅/Veh. Dunnett's test. **(D)** Typical trace of O₂ consumption for Sc. Aβ and Aβ₂₅₋₃₅-injected mice.





FIGURE 3 | Direct application of ANAVEX2-73 or PRE-084 affected mitochondrial respiration only at high concentrations. Mitochondria (0.8 mg/ml) were loaded in the chamber with appropriate buffer at 30°C. The drugs (10^{-8} to 10^{-4} M) were bath applied 5 min before the respiration measures. State 2 respiration was activated by addition of pyruvate-malate (5 mM). State 3 respiration was induced with ADP. State 4 respiration was

provoked by the addition of carboxyatractylide (CAT), a blocker of ATP-ADP carrier. The respiratory control ratio is the state 3/state 4 ratio. $n = 4-8$ per group ($n = 14$ for the no-drug group), $F_{(5,38)} = 3.08$, $p < 0.05$ for ANAVEX2-73 and $F_{(5,31)} = 5.19$, $p < 0.01$ for PRE-084 in **(A)**; $F_{(5,38)} = 22.1$, $p < 0.0001$ for ANAVEX2-73 and $F_{(5,31)} = 7.53$, $p < 0.001$ for PRE-084 in **(B)**. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ vs. no-drug data; Dunnett's test.



at days 5 and 7 after the peptide injection (Figure 5C). ANAVEX2-73 was tested at 0.1–1 mg/kg. A dose-dependent decrease in Bax level was measured at the highest dose tested (Figure 6D). The drug impacted marginally Bcl-2 levels (Figure 6E), but the data examined as Bax/Bcl-2 ratio clearly showed a beneficial effect of the compound, particularly significant at 0.3 mg/kg (Figure 6F).

DISCUSSION

In this study, we showed that mitochondrial functionality was altered several days after an ICV injection of the A β_{25-35} peptide

in mouse. It was shown by reduced respiration rates, decreased complex IV activity and increased oxidative stress and proapoptotic markers. Five days after a single injection of oligomeric A β_{25-35} peptide, respiration was reduced in isolated mitochondria from mouse hippocampus, as compared with control Sc. A β -treated mice. Basal respiration (State 2 and State 4) was unaffected by A β_{25-35} , but ADP-stimulated respiration (State 3) was significantly decreased. Reduced State 3 respiration resulted in a decreased control ratio in A β_{25-35} -injected mice. Uncoupled state was also altered, suggesting a defect in respiratory chain complex I-IV rather than a defect in ATP synthase complex activity. The reduced respiratory rates and control ratio observed in this *in vivo* model are coherent with previous studies exploring the impact of A β peptides, and particularly A β_{25-35} , on mitochondrial respiration *in vitro*. Application of A β_{25-35} on isolated mitochondria decreased respiration rates (Casley et al., 2002a; Aleardi et al., 2005; Clementi et al., 2005). Interestingly, in these studies, basal respiration (State 4) was also altered after A β_{25-35} treatment. In the same way, mitochondria isolated from cultured neurons and exposed to A β_{25-35} showed decreased ATP production, suggesting a reduced mitochondrial respiration (Casley et al., 2002b; Dong et al., 2010). In search for an explanation to this decreased mitochondrial respiration after A β_{25-35} , we measured the respiratory chain complex I-IV activities. We found that complex IV activity was significantly reduced 5 days after A β_{25-35} . In most of the *in vitro* studies, application of A β_{25-35} on cultured neurons or directly on isolated mitochondria resulted in reduced activities of complexes I and IV (Casley et al., 2002a,b; Aleardi et al., 2005; Clementi et al., 2005; Dong et al., 2010). This appears to be a difference between *in vitro* and *in vivo* studies with the A β_{25-35} peptide. It could explain the decreased state 4 respiration observed *in vitro*. But, consistent with our results, are data in transgenic mice expressing human mutant amyloid precursor protein (hAPPm). A β peptide has been found to accumulate in mitochondria and respiratory rates and complex IV activity were also decreased in these animals (Caspersen et al., 2005; Rhein et al., 2009; Du et al., 2010). Moreover, Canevari et al. (1999) described a specific and concentration-dependent decrease in complex IV activity after A β_{25-35} application in isolated mitochondria. Disruption of complex IV activity induced by A β peptides could be due to a direct interaction between A β and the enzymatic subunits (Crouch et al., 2006; Hernandez-Zimbron et al., 2012) but also to the direct induction of oxidative stress by A β peptides, and particularly A β_{25-35} . Indeed, membrane lipid peroxidation induced by ROS overproduction might alter complex IV subunits and disrupt complex IV activity (Paradies et al., 1998; Bobba et al., 2013), as we confirmed the marked increase in oxidative stress in mouse hippocampus, already described in this model (Meunier et al., 2006). The release of cytochrome *c* from mitochondria into the cytosol that was increased 5 days after A β_{25-35} injection can also contribute to the dysfunction of the mitochondrial respiratory chain and thus to the oxidative stress. Concomitantly to the cytochrome *c* release, the ratio Bax/Bcl-2 was significantly increased in mouse hippocampus, 5 and 7 days after A β_{25-35} injection and this is likely related to the cellular response to the high level of oxidative




FIGURE 5 | ANAVEX2-73 and PRE-084 prevented Aβ₂₅₋₃₅-induced cytochrome c release. (A) Typical blots of mitochondrial (M) and cytosolic (C) fractions with cytochrome c, oxphos and β-tubulin antibodies. **(B)** Cytochrome c content (cytosol to mitochondrial content ratios). Mice were injected with Sc. Aβ or Aβ₂₅₋₃₅ (9 nmol ICV) and sacrificed at indicated days. *n* = 9–12 per

group. * *p* < 0.05 vs. Sc. Aβ. Student's *t*-test. **(C, D)** Mice were treated with ANAVEX2-73 (0.3 mg/kg IP) or PRE-084 (0.5 mg/kg IP) before Aβ₂₅₋₃₅ and sacrificed after 5 days. *n* = 8–17, *F*_(2,37) = 5.96, *p* < 0.01 in **(C)**; *n* = 8–14, *F*_(2,33) = 8.19, *p* < 0.01 in **(D)**. **p* < 0.05, ***p* < 0.01 vs. Sc. Aβ, #*p* < 0.05, ##*p* < 0.01 vs. Aβ₂₅₋₃₅; Dunnett's test.

stress and decreased mitochondrial respiration (Mattson et al., 2008).

The aim of the study was to establish whether the localization of the σ_1 R on mitochondria (Klouz et al., 2002) and more precisely at MAM (Hayashi and Su, 2007) is related to its particular neuroprotective activity. Indeed, both ANAVEX2-73 and PRE-084, prevented Aβ₂₅₋₃₅-induced oxidative stress and increase in pro-apoptotic markers. The results are coherent with previously reported observations, since the σ_1 R has been involved in regulation of oxidative stress (Pal et al., 2012; Tsai et al., 2012). Moreover, σ_1 R agonists prevented oxidative stress-induced apoptosis *in vitro* through upregulation of the anti-apoptotic Bcl-2 protein expression (Meunier and Hayashi, 2010). The σ_1 R activation also prevented Bax increase and caspase-3 activation, linked to cytochrome *c* release, after glutamate insult in retinal ganglionic cells (Tchedre and Yorino, 2008). Another study also showed a decreased caspase-3 activation after PRE-084 treatment in a Huntington's disease model (Hyrskyluoto et al., 2013). ANAVEX2-73 and PRE-084 both

fully reversed state 3 and uncoupled state respiration deficits. Surprisingly both drugs also increased basal respiration (state 2 and state 4) and PRE-084, but not ANAVEX2-73 increased state 3 and uncoupled state respiration, compared to control mice. These differences in ANAVEX2-73 and PRE-084 effects could be explained by the higher affinity of the later for the σ_1 receptor. Although state 3 respiration raised to the control level after ANAVEX2-73 treatment, CR was not different from the Aβ₂₅₋₃₅ group because state 4 respiration also increased. However, the high increased of both state 3 and state 4 respiration after PRE-084 treatment increased the control ratio to the control level. Such increased respiration after stimulation of σ_1 R has never been described and is not observed in Sc. Aβ-treated control mice. An attempt to explain this phenomenon is challenging. The increased state 3 and uncoupled state respiration rates observed after PRE-084 treatment were clearly not an artifact or due to experimental conditions as all the samples were prepared in strictly the same way and measures realized randomly among treatment groups and in

several experimental sessions. Few studies have examined the involvement of σ_1 R in mitochondrial respiration. BHDP is a selective ligand with nanomolar affinity for the σ_1 receptor (Klouz et al., 2003). A three days pretreatment with BHDP in rats prevented the decrease in CR in mitochondria isolated from rat liver after ischemia-reperfusion (Klouz et al., 2008). We previously suggested that protective effect on mitochondrial function of σ_1 agonist could be attributable to the presence of σ_1 R to the mitochondrial membrane (Klouz et al., 2002). Indeed, σ_1 R are present in lipid rafts and modify their composition and particularly the ganglioside content (Hayashi and Su, 2005). In mitochondrial membrane, lipid rafts microdomains, and particularly the glycosphingolipid GD3, are involved in alteration of mitochondrial membrane potential and apoptosis (Malorni et al., 2007). Therefore, an explanation of increased mitochondrial respiration by σ_1 ligand could be through the capacity of the protein to modulate lipid rafts gangliosids at mitochondrial membrane. Another explanation could be the chaperone activity of the σ_1 protein on InsP₃ receptor at the ER membrane (Hayashi and Su, 2007). The InsP₃ receptor stabilization allows proper Ca²⁺ exchange between ER and mitochondria, which are critical for mitochondrial bioenergetics (Cárdenas et al., 2010). Cutamesine (SA4503), another selective σ_1 agonist, increased ATP production in cardiomyocytes, even without pathological conditions, suggesting an increase in mitochondrial respiration after σ_1 R stimulation (Tagashira et al., 2013). The increased mitochondrial bioenergetics after σ_1 agonist treatment is largely attributable to ER Ca²⁺ mobilization into mitochondria, because treatment with xestospongine C, an antagonist of the InsP₃ receptor abolished the effects of the σ_1 agonist on ATP production. However, in our model, this mechanism did not seem to be involved in the σ_1 protein-induced increased mitochondrial respiration because the agonist was injected only once, five days before the mitochondrial isolation. Moreover, Tsai et al. (2009) described Ca²⁺-independent perturbations of mitochondrial functions in hippocampal neurons after σ_1 receptor knockdown with siRNA. To note, another interesting consequence of σ_1 R activity is the modulation of different genes expression. Some of them are involved in oxidative stress regulation, e.g., NAD(P)H quinone oxidoreductase (Nqo1), Superoxide dismutase (SOD), Atf-4 and Prdx6 (Pal et al., 2012; Tsai et al., 2012). It could be worth investigating whether the PRE-084-induced increased respiratory efficacy could be due to the modification of expression of specific genes.

In conclusion, we provided here the clear evidence that the neuroprotective activity of σ_1 receptor agonists, including the mixed muscarinic ligand/ σ_1 agonist ANAVEX2-73, involves mitochondrial protection in AD models. Indeed, the drug protected mitochondria against A β _{25–35} insult. Mitochondrial dysfunction in AD is a critical hallmark of the pathology, as mitochondria are, at the same time, a target and promoter of the physiopathological process (for review, see Eckert et al., 2010; Selfridge et al., 2013). Mitochondrial dysfunction and, notably, resulting oxidative stress are responsible for the increased A β production (Dyrks et al., 1992; Leuner et al., 2007) and Tau hyperphosphorylation (Melov et al., 2007), the two biochemical




FIGURE 6 | ANAVEX2-73 protected against A β _{25–35}-induced increase in Bax/Bcl-2 ratio in the mouse hippocampus. (A–C) Mice were sacrificed at 1, 3, 5 and 7 days after A β _{25–35} injection and Bax and Bcl-2 contents determined by western blots. **(D–F)** Effect of ANAVEX2-73 (0.1–1 mg/kg IP) on Bax and Bcl-2 levels, 7 days after A β _{25–35} injection. Typical blots are represented above the bar graphs. Mice were injected with ANAVEX2-73 or vehicle before A β _{25–35} peptide (9 nmol). $n = 5–15$ per groups. $F_{(4,44)} = 1.98$, $p > 0.05$ in **(A)**; $F_{(4,44)} = 1.95$, $p > 0.05$ in **(B)**; $F_{(4,44)} = 2.66$, $p < 0.05$ in **(C)**; $F_{(4,49)} = 3.01$, $p < 0.05$ in **(D)**; $F_{(4,49)} = 0.89$, $p > 0.05$ in **(E)**, $F_{(4,49)} = 3.09$, $p < 0.05$ in **(F)**. * $p < 0.05$, ** $p < 0.01$ vs. Sc. A β ; # $p < 0.05$ vs. A β _{25–35}; Dunnett's test.

signatures of AD. Thus, by preserving mitochondrial function ANAVEX2-73 could expectedly be more than a symptomatic drug and block the physiopathological amplification of the pathology.

ACKNOWLEDGMENTS

This work was supported by collaboration contracts between Anavex Life Sciences (New York, NY, USA) and Amylgen (Montferrier-sur-Lez, France) and between Amylgen and the University of Montpellier (France). The companies had no role in study design or analysis. Valentine Lahmy is recipient of a CIFRE PhD funding (ANRT, Paris, France).

REFERENCES

- Aleardi, A. M., Benard, G., Augereau, O., Malgat, M., Talbot, J. C., Mazat, J. P., et al. (2005). Gradual alteration of mitochondrial structure and function by beta-amyloids: importance of membrane viscosity changes, energy deprivation, reactive oxygen species production and cytochrome c release. *J. Bioenerg. Biomembr.* 37, 207–225. doi: 10.1007/s10863-005-6631-3
- Bobba, A., Amadoro, G., Valenti, D., Corsetti, V., Lassandro, R., and Atlante, A. (2013). Mitochondrial respiratory chain Complexes I and IV are impaired by β -amyloid via direct interaction and through Complex I-dependent ROS production, respectively. *Mitochondrion* 13, 298–311. doi: 10.1016/j.mito.2013.03.008
- Canevari, L., Clark, J. B., and Bates, T. E. (1999). β -Amyloid fragment 25–35 selectively decreases complex IV activity in isolated mitochondria. *FEBS Lett.* 457, 131–134. doi: 10.1016/S0014-5793(99)01028-5
- Cárdenas, C., Miller, R. A., Smith, I., Bui, T., Molgó, J., Müller, M., et al. (2010). Essential regulation of cell bioenergetics by constitutive InsP₃ receptor Ca²⁺ transfer to mitochondria. *Cell* 142, 270–283. doi: 10.1016/j.cell.2010.06.007
- Casley, C. S., Canevari, L., Land, J. M., Clark, J. B., and Sharpe, M. A. (2002b). β -Amyloid inhibits integrated mitochondrial respiration and key enzyme activities. *J. Neurochem.* 80, 91–100. doi: 10.1046/j.0022-3042.2001.00681.x
- Casley, C. S., Land, J. M., Sharpe, M. A., Clark, J. B., Duchon, M. R., and Canevari, L. (2002a). β -Amyloid fragment 25–35 causes mitochondrial dysfunction in primary cortical neurons. *Neurobiol. Dis.* 10, 258–267. doi: 10.1006/nbdi.2002.0516
- Caspersen, C., Wang, N., Yao, J., Sosunov, A., Chen, X., Lustbader, J. W., et al. (2005). Mitochondrial A β : a potential focal point for neuronal metabolic dysfunction in Alzheimer's disease. *FASEB J.* 19, 2040–2041. doi: 10.1096/fj.05-3735fje
- Clementi, M. E., Marini, S., Coletta, M., Orsini, F., Giardina, B., and Misi, F. (2005). A β 31–35 and A β 25–35 fragments of amyloid beta-protein induce cellular death through apoptotic signals: role of the redox state of methionine-35. *FEBS Lett.* 579, 2913–2918. doi: 10.1016/j.febslet.2005.04.041
- Crews, L., and Masliah, E. (2010). Molecular mechanisms of neurodegeneration in Alzheimer's disease. *Hum. Mol. Genet.* 19, R12–R20. doi: 10.1093/hmg/ddq160
- Crouch, P. J., Barnham, K. J., Duce, J. A., Blake, R. E., Masters, C. L., and Trounce, I. A. (2006). Copper-dependent inhibition of cytochrome c oxidase by A β 1–42 requires reduced methionine at residue 35 of the A β peptide. *J. Neurochem.* 99, 226–236. doi: 10.1111/j.1471-4159.2006.04050.x
- Delobette, S., Privat, A., and Maurice, T. (1997). In vitro aggregation facilitates β -amyloid peptide-(25–35)-induced amnesia in the rat. *Eur. J. Pharmacol.* 319, 1–4. doi: 10.1016/s0014-2999(96)00922-3
- Dong, W., Huang, F., Fan, W., Cheng, S., Chen, Y., Zhang, W., et al. (2010). Differential effects of melatonin on amyloid- β peptide 25–35-induced mitochondrial dysfunction in hippocampal neurons at different stages of culture. *J. Pineal Res.* 48, 117–125. doi: 10.1111/j.1600-079x.2009.00734.x
- Du, H., Guo, L., Yan, S., Sosunov, A. A., McKhann, G. M., and Yan, S. S. (2010). Early deficits in synaptic mitochondria in an Alzheimer's disease mouse model. *Proc. Natl. Acad. Sci. U S A* 107, 18670–18675. doi: 10.1073/pnas.1006586107
- Dyrks, T., Dyrks, E., Hartmann, T., Masters, C., and Beyreuther, K. (1992). Amyloidogenicity of β A4 and β A4-bearing amyloid protein precursor fragments by metal-catalyzed oxidation. *J. Biol. Chem.* 267, 18210–18217.
- Eckert, A., Schulz, K. L., Rhein, V., and Götz, J. (2010). Convergence of amyloid- β and tau pathologies on mitochondria in vivo. *Mol. Neurobiol.* 41, 107–114. doi: 10.1007/s12035-010-8109-5
- Espallargues, J., Lapalud, P., Christopoulos, A., Avlani, V. A., Sexton, P. M., Vamvakides, A., et al. (2007). Involvement of the sigma₁ (σ_1) receptor in the anti-amnesic, but not antidepressant-like, effects of the aminotetrahydrofuran derivative ANAVEX1–41. *Br. J. Pharmacol.* 152, 267–279. doi: 10.1038/sj.bjp.0707386
- Hanada, M., Aimé-Sempé, C., Sato, T., and Reed, J. C. (1995). Structure-function analysis of Bcl-2 protein. Identification of conserved domains important for homodimerization with Bcl-2 and heterodimerization with Bax. *J. Biol. Chem.* 270, 11962–11969. doi: 10.1074/jbc.270.20.11962
- Hayashi, T., and Su, T. P. (2005). The potential role of σ_1 receptors in lipid transport and lipid raft reconstitution in the brain: implication for drug abuse. *Life Sci.* 77, 1612–1624. doi: 10.1016/j.lfs.2005.05.009
- Hayashi, T., and Su, T. P. (2007). Sigma-1 receptor chaperones at the ER-mitochondrion interface regulate Ca²⁺ signaling and cell survival. *Cell* 131, 596–610. doi: 10.1016/j.cell.2007.08.036
- Hernandez-Zimbron, L. F., Luna-Muñoz, J., Mena, R., Vazquez-Ramirez, R., Kubli-Garfias, C., Cribbs, D. H., et al. (2012). Amyloid- β peptide binds to cytochrome C oxidase subunit 1. *PLoS One* 7:e42344. doi: 10.1371/journal.pone.0042344
- Hyrskyluoto, A., Pulli, I., Törnqvist, K., Ho, T. H., Korhonen, L., and Lindholm, D. (2013). Sigma₁ receptor agonist PRE084 is protective against mutant huntingtin-induced cell degeneration: involvement of calpastatin and the NF- κ B pathway. *Cell Death Dis.* 4:e646. doi: 10.1038/cddis.2013.170
- Jürgensmeier, J. M., Xie, Z., Deveraux, Q., Ellerby, L., Bredesen, D., and Reed, J. C. (1998). Bax directly induces release of cytochrome c from isolated mitochondria. *Proc. Natl. Acad. Sci. USA* 95, 4997–5002. doi: 10.1073/pnas.95.9.4997
- Kaminsky, Y. G., and Kosenko, E. A. (2008). Effects of amyloid- β peptides on hydrogen peroxide-metabolizing enzymes in rat brain in vivo. *Free Radic. Res.* 42, 564–573. doi: 10.1080/10715760802159057
- Klouz, A., Said, D. B., Ferchichi, H., Kourda, N., Ouane, L., Lakhal, M., et al. (2008). Protection of cellular and mitochondrial functions against liver ischemia by N-benzyl-N'-(2-hydroxy-3,4-dimethoxybenzyl)-piperazine (BHDP), a σ_1 ligand. *Eur. J. Pharmacol.* 578, 292–299. doi: 10.1016/j.ejphar.2007.09.038
- Klouz, A., Sapena, R., Liu, J., Maurice, T., Tillement, J. P., Papadopoulos, V., et al. (2002). Evidence for σ_1 -like receptors in isolated rat liver mitochondrial membranes. *Br. J. Pharmacol.* 135, 1607–1615. doi: 10.1038/sj.bjp.0704626
- Klouz, A., Tillement, J. P., Boussard, M. F., Wierzbicki, M., Berezowski, V., Cecchelli, R., et al. (2003). [³H]BHDP as a novel and selective ligand for σ_1 receptors in liver mitochondria and brain synaptosomes of the rat. *FEBS Lett.* 553, 157–162. doi: 10.1016/s0014-5793(03)01011-1
- Kroemer, G., and Reed, J. C. (2000). Mitochondrial control of cell death. *Nat. Med.* 6, 513–519. doi: 10.1038/74994
- Lahmy, V., Meunier, J., Malmström, S., Naert, G., Givalois, L., Kim, S. H., et al. (2013). Blockade of Tau hyperphosphorylation and A β 1–42 generation by the aminotetrahydrofuran derivative ANAVEX2–73, a mixed muscarinic and σ_1 receptor agonist, in a nontransgenic mouse model of Alzheimer's disease. *Neuropsychopharmacology* 38, 1706–1723. doi: 10.1038/npp.2013.70
- Leuner, K., Hauptmann, S., Abdel-Kader, R., Scherping, I., Keil, U., Strosznajder, J. B., et al. (2007). Mitochondrial dysfunction: the first domino in brain aging and Alzheimer's disease?. *Antioxid. Redox Signal.* 9, 1659–1675. doi: 10.1089/ars.2007.1763
- Leuner, K., Müller, W. E., and Reichert, A. S. (2012). From mitochondrial dysfunction to amyloid β formation: novel insights into the pathogenesis of Alzheimer's disease. *Mol. Neurobiol.* 46, 186–193. doi: 10.1007/s12035-012-8307-4
- Liu, X., Kim, C. N., Yang, J., Jemmerson, R., and Wang, X. (1996). Induction of apoptotic program in cell-free extracts: requirement for dATP and cytochrome c. *Cell* 86, 147–157. doi: 10.1016/s0092-8674(00)80085-9
- Lustbader, J. W., Cirilli, M., Lin, C., Xu, H. W., Takuma, K., Wang, N., et al. (2004). ABAD directly links A β to mitochondrial toxicity in Alzheimer's disease. *Science* 304, 448–452. doi: 10.1126/science.1091230
- Malorni, W., Giammaroli, A. M., Garofalo, T., and Sorice, M. (2007). Dynamics of lipid raft components during lymphocyte apoptosis: the paradigmatic role of GD3. *Apoptosis* 12, 941–949. doi: 10.1007/s10495-007-0757-1
- Manczak, M., Anekonda, T. S., Henson, E., Park, B. S., Quinn, J., and Reddy, P. H. (2006). Mitochondria are a direct site of a beta accumulation in Alzheimer's disease neurons: implications for free radical generation and oxidative damage in disease progression. *Hum. Mol. Genet.* 15, 1437–1449. doi: 10.1093/hmg/ddl066
- Marzo, I., Brenner, C., Zamzami, N., Jürgensmeier, J. M., Susin, S. A., Vieira, H. L., et al. (1998). Bax and adenine nucleotide translocator cooperate in the mitochondrial control of apoptosis. *Science* 281, 2027–2031. doi: 10.1126/science.281.5385.2027
- Mattson, M. P., Gleichmann, M., and Cheng, A. (2008). Mitochondria in neuroplasticity and neurological disorders. *Neuron* 60, 748–766. doi: 10.1016/j.neuron.2008.10.010
- Maurer, I., Zierz, S., and Möller, H. J. (2000). A selective defect of cytochrome c oxidase is present in brain of Alzheimer disease patients. *Neurobiol. Aging* 21, 455–462. doi: 10.1016/s0197-4580(00)00112-3
- Maurice, T., Lockhart, B. P., and Privat, A. (1996). Amnesia induced in mice by centrally administered β -amyloid peptides involves cholinergic dysfunction. *Brain Res.* 706, 181–193. doi: 10.1016/0006-8993(95)01032-7
- Melov, S., Adlard, P. A., Morten, K., Johnson, E., Golden, T. R., Hinerfeld, D., et al. (2007). Mitochondrial oxidative stress causes hyperphosphorylation of tau. *PLoS One* 2:e536. doi: 10.1371/journal.pone.0000536

- Meunier, J., and Hayashi, T. (2010). Sigma-1 receptors regulate Bcl-2 expression by reactive oxygen species-dependent transcriptional regulation of nuclear factor κ B. *J. Pharmacol. Exp. Ther.* 332, 388–397. doi: 10.1124/jpet.109.160960
- Meunier, J., Ieni, J., and Maurice, T. (2006). The anti-amnesic and neuroprotective effects of donepezil against amyloid β 25–35 peptide-induced toxicity in mice involve an interaction with the σ_1 receptor. *Br. J. Pharmacol.* 149, 998–1012. doi: 10.1038/sj.bjp.0706927
- Mosconi, L., Pupi, A., and De Leon, M. J. (2008). Brain glucose hypometabolism and oxidative stress in preclinical Alzheimer's disease. *Ann. N Y Acad. Sci.* 1147, 180–195. doi: 10.1196/annals.1427.007
- Narita, M., Shimizu, S., Ito, T., Chittenden, T., Lutz, R. J., Matsuda, H., et al. (1998). Bax interacts with the permeability transition pore to induce permeability transition and cytochrome c release in isolated mitochondria. *Proc. Natl. Acad. Sci. U S A* 95, 14681–14686. doi: 10.1073/pnas.95.25.14681
- Pal, A., Fontanilla, D., Gopalakrishnan, A., Chae, Y. K., Markley, J. L., and Ruoho, A. E. (2012). The σ_1 receptor protects against cellular oxidative stress and activates antioxidant response elements. *Eur. J. Pharmacol.* 682, 12–20. doi: 10.1016/j.ejphar.2012.01.030
- Paradies, G., Ruggiero, F. M., Petrosillo, G., and Quagliariello, E. (1998). Peroxidative damage to cardiac mitochondria: cytochrome oxidase and cardiolipin alterations. *FEBS Lett.* 424, 155–158. doi: 10.1016/s0014-5793(98)00161-6
- Ren, R., Zhang, Y., Li, B., Wu, Y., and Li, B. (2011). Effect of β -amyloid (25–35) on mitochondrial function and expression of mitochondrial permeability transition pore proteins in rat hippocampal neurons. *J. Cell. Biochem.* 112, 1450–1457. doi: 10.1002/jcb.23062
- Rhein, V., Song, X., Wiesner, A., Ittner, L. M., Baysang, G., Meier, F., et al. (2009). Amyloid- β and tau synergistically impair the oxidative phosphorylation system in triple transgenic Alzheimer's disease mice. *Proc. Natl. Acad. Sci. U S A* 106, 20057–20062. doi: 10.1073/pnas.0905529106
- Selfridge, J. E., E. L., Lu, J., and Swerdlow, R. H. (2013). Role of mitochondrial homeostasis and dynamics in Alzheimer's disease. *Neurobiol. Dis.* 51, 3–12. doi: 10.1016/j.nbd.2011.12.057
- Selkoe, D. J. (2004). Cell biology of protein misfolding: the examples of Alzheimer's and Parkinson's diseases. *Nat. Cell Biol.* 6, 1054–1061. doi: 10.1038/ncb1104-1054
- Smith, M. A., Perry, G., Richey, P. L., Sayre, L. M., Anderson, V. E., Beal, M. F., et al. (1996). Oxidative damage in Alzheimer's. *Nature* 382, 120–121. doi: 10.1038/382120b0
- Tagashira, H., Zhang, C., Lu, Y. M., Hasegawa, H., Kanai, H., Han, F., et al. (2013). Stimulation of σ_1 -receptor restores abnormal mitochondrial Ca^{2+} mobilization and ATP production following cardiac hypertrophy. *Biochim. Biophys. Acta* 1830, 3082–3094. doi: 10.1016/j.bbagen.2012.12.029
- Tchedre, K. T., and Yorio, T. (2008). σ_1 receptors protect RGC-5 cells from apoptosis by regulating intracellular calcium, Bax levels and caspase-3 activation. *Invest. Ophthalmol. Vis. Sci.* 49, 2577–2588. doi: 10.1167/iovs.07-1101
- Tsai, S. Y., Hayashi, T., Harvey, B. K., Wang, Y., Wu, W. W., Shen, R. F., et al. (2009). σ_1 Receptors regulate hippocampal dendritic spine formation via a free radical-sensitive mechanism involving Rac1xGTP pathway. *Proc. Natl. Acad. Sci. U S A* 106, 22468–22473. doi: 10.1073/pnas.0909089106
- Tsai, S. Y., Rothman, R. K., and Su, T. P. (2012). Insights into the σ_1 receptor chaperone's cellular functions: a microarray report. *Synapse* 66, 42–51. doi: 10.1002/syn.20984
- Verstreken, P., Ly, C. V., Venken, K. J., Koh, T. W., Zhou, Y., and Bellen, H. J. (2005). Synaptic mitochondria are critical for mobilization of reserve pool vesicles at Drosophila neuromuscular junctions. *Neuron* 47, 365–378. doi: 10.1016/j.neuron.2005.06.018
- Villard, V., Espallergues, J., Keller, E., Alkam, T., Nitta, A., Yamada, K., et al. (2009). Antiamnesic and neuroprotective effects of the aminotetrahydrofuran derivative ANAVEX1–41 against amyloid β 25–35-induced toxicity in mice. *Neuropsychopharmacology* 34, 1552–1566. doi: 10.1038/npp.2008.212
- Villard, V., Espallergues, J., Keller, E., Vamvakides, A., and Maurice, T. (2011). Anti-amnesic and neuroprotective potentials of the mixed muscarinic receptor/sigma $_1$ (σ_1) ligand ANAVEX2–73, a novel aminotetrahydrofuran derivative. *J. Psychopharmacol.* 25, 1101–1117. doi: 10.1177/0269881110379286
- Yang, J., Liu, X., Bhalla, K., Kim, C. N., Ibrado, A. M., Cai, J., et al. (1997). Prevention of apoptosis by Bcl-2: release of cytochrome c from mitochondria blocked. *Science* 275, 1129–1132. doi: 10.1126/science.275.5303.1129
- Yates, C. M., Butterworth, J., Tennant, M. C., and Gordon, A. (1990). Enzyme activities in relation to pH and lactate in postmortem brain in Alzheimer-type and other dementias. *J. Neurochem.* 55, 1624–1630. doi: 10.1111/j.1471-4159.1990.tb04948.x
- Yin, X. M., Oltvai, Z. N., and Korsmeyer, S. J. (1995). Heterodimerization with Bax is required for Bcl-2 to repress cell death. *Curr. Top. Microbiol. Immunol.* 194, 331–338. doi: 10.1007/978-3-642-79275-5_38
- Zussy, C., Brureau, A., Delair, B., Marchal, S., Keller, E., Ixart, G., et al. (2011). Time-course and regional analyses of the physiopathological changes induced after cerebral injection of an amyloid β fragment in rats. *Am. J. Pathol.* 179, 315–334. doi: 10.1016/j.ajpath.2011.03.021
- Zussy, C., Brureau, A., Keller, E., Marchal, S., Blayo, C., Delair, B., et al. (2013). Alzheimer's disease related markers, cellular toxicity and behavioral deficits induced six weeks after oligomeric amyloid- β peptide injection in rats. *PLoS One* 8:e53117. doi: 10.1371/journal.pone.0053117

Conflict of Interest Statement: Valentine Lahmy and Vanessa Villard are employees of Amylgen. Tanguy Maurice is a member of the scientific advisory board of Anavex Life Sciences and scientific director of Amylgen. The companies had no role in the present research, except providing drug and funding. Other authors declare no conflict of interest.

Received: 29 October 2014; accepted: 20 December 2014; published online: 20 January 2015.

Citation: Lahmy V, Long R, Morin D, Villard V and Maurice T (2015) Mitochondrial protection by the mixed muscarinic/ σ_1 ligand ANAVEX2-73, a tetrahydrofuran derivative, in $\text{A}\beta_{25-35}$ peptide-injected mice, a nontransgenic Alzheimer's disease model. *Front. Cell. Neurosci.* 8:463. doi: 10.3389/fncel.2014.00463

This article was submitted to the journal *Frontiers in Cellular Neuroscience*.

Copyright © 2015 Lahmy, Long, Morin, Villard and Maurice. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution and reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.