



HAL
open science

Effect of salting process on the histological structure of salmon flesh

Olivier Loison, Weijunlang Jiang, Annie Venien, Oulyana Gaubin, Chantal Cazevieille, Thierry Astruc

► **To cite this version:**

Olivier Loison, Weijunlang Jiang, Annie Venien, Oulyana Gaubin, Chantal Cazevieille, et al.. Effect of salting process on the histological structure of salmon flesh. International Symposium on Food Nutrition and Health, May 2017, Dalian, China. hal-01924390

HAL Id: hal-01924390

<https://hal.science/hal-01924390v1>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Effect of salting process on the histological structure of salmon flesh

Olivier Loison¹, Weijunlang Jiang¹, Annie Vénien¹, Oulyana Gaubain¹, Chantal Cazevielle², Thierry Astruc^{1*}

¹INRA de Theix, UR370 QuaPA, F-63122 St-Genès-Champanelle (France)

²MRI-COMET, Hôpital Saint Eloi F-34091 Montpellier (France)

Abstract:

(Less than 300 words with one page, Times New Rome, No. 12)

Atlantic Salmon , *Salmo Salar*, is composed of approximately 70% water, 19% protein, 10% lipid and 1% small nutrients (vitamins, glycogen, pigments ...). Smoked salmon comes from the processing of fresh salmon: the fillets are removed from the fish, salted and then smoked. Salting can be carried out with dry salt or by brine injection. The objective of the study was to compare the evolution of the cell structure and ultrastructure of the salmon muscle subjected to salting with dry salt and salting by brine injection.

The experiment was carried out on 6 salmons. For each salmon, 1 fillet was salted with dry salt and the other fillet was additionally injected with brine. The fillets were smoked at low temperature (20-22 ° C). Samples were taken for histological analyzes (cryofixation, preparation of 10 µm thick histological sections, staining and observation by optical microscopy) and ultrastructural analyzes (chemical fixation, dehydration, resin inclusion, ultrathin sections, staining and observation by electron microscopy).

The results show a decrease in the size of the extracellular spaces after injection of brine, which is probably related to a swelling of the muscle cells that accumulate water. Dry salt salting, on the other hand, shows no difference in the extracellular size compared to the unsalted muscle.

The salting substantially degrades the ultrastructure of the muscle with in particular a significant solubilization of the Z-lines. Salting by injection of brine leads to swelling of the myofibrils and almost complete solubilization of the Z lines.

Note: Please send abstract to the referred email address (nercsf@163.com) before April 23, 2017.