



**HAL**  
open science

## Multistage Matchings

Evripidis Bampis, Bruno Escoffier, Michael Lampis, Vangelis Th. Paschos

► **To cite this version:**

Evripidis Bampis, Bruno Escoffier, Michael Lampis, Vangelis Th. Paschos. Multistage Matchings. 16th Scandinavian Symposium and Workshops on Algorithm Theory (SWAT 2018), Jun 2018, Malmo, Sweden. pp.7:1–7:13, 10.4230/LIPIcs.SWAT.2018.7. hal-01926860

**HAL Id: hal-01926860**

**<https://hal.science/hal-01926860>**

Submitted on 19 Nov 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# 1 Multistage Matchings

2 **Evrpidis Bampis**

3 Sorbonne Université, CNRS, Laboratoire d'Informatique de Paris 6, LIP6, Paris, France  
4 evripidis.bampis@lip6.fr

5 **Bruno Escoffier**

6 Sorbonne Université, CNRS, Laboratoire d'Informatique de Paris 6, LIP6, Paris, France  
7 bruno.escoffier@lip6.fr

8 **Michael Lampis**

9 Université Paris-Dauphine, PSL Research University, CNRS, LAMSADE, Paris, France  
10 michail.lampis@dauphine.fr

11 **Vangelis Th. Paschos**

12 Université Paris-Dauphine, PSL Research University, CNRS, LAMSADE, Paris, France  
13 vangelis.paschos@dauphine.fr

---

## 14 **Abstract**

15 We consider a *multistage* version of the PERFECT MATCHING problem which models the  
16 scenario where the costs of edges change over time and we seek to obtain a solution that achieves  
17 low total cost, while minimizing the number of changes from one instance to the next. Formally,  
18 we are given a sequence of edge-weighted graphs on the same set of vertices  $V$ , and are asked to  
19 produce a perfect matching in each instance so that the total edge cost plus the transition cost  
20 (the cost of exchanging edges), is minimized. This model was introduced by Gupta et al. (ICALP  
21 2014), who posed as an open problem its approximability for bipartite instances. We completely  
22 resolve this question by showing that *Minimum Multistage Perfect Matching* (MIN-MPM) does  
23 not admit an  $n^{1-\epsilon}$ -approximation, even on bipartite instances with only two time steps.

24 Motivated by this negative result, we go on to consider two variations of the problem. In  
25 *Metric Minimum Multistage Perfect Matching* problem (METRIC-MIN-MPM) we are promised  
26 that edge weights in each time step satisfy the triangle inequality. We show that this problem  
27 admits a 3-approximation when the number of time steps is 2 or 3. On the other hand, we  
28 show that even the metric case is APX-hard already for 2 time steps. We then consider the  
29 complementary maximization version of the problem, *Maximum Multistage Perfect Matching*  
30 problem (MAX-MPM), where we seek to maximize the total profit of all selected edges plus the  
31 total number of non-exchanged edges. We show that MAX-MPM is also APX-hard, but admits  
32 a constant factor approximation algorithm for any number of time steps.

33 **2012 ACM Subject Classification** Theory of Computation → Design and Analysis of Algorithms  
34 → Approximation Algorithms Analysis

35 **Keywords and phrases** Perfect Matching; Temporal Optimization; Multistage Optimization

36 **Digital Object Identifier** 10.4230/LIPIcs.SWAT.2018.7

37 **Acknowledgements** This work benefited from the support of the FMJH program “Gaspard  
38 Monge in optimization and Operation Research” and from the support to this program from  
39 EDF, via the project 2016-1760H/C16/1507 “Stability versus Optimality in Dynamic Environ-  
40 nment Algorithmics”.



© Evripidis Bampis, Bruno Escoffier, Michael Lampis, and Vangelis Th. Paschos;  
licensed under Creative Commons License CC-BY

16th Scandinavian Symposium and Workshops on Algorithm Theory (SWAT 2018).

Editor: David Eppstein; Article No. 7; pp. 7:1–7:14

Leibniz International Proceedings in Informatics



LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

41 **1 Introduction**

42 In classical Combinatorial Optimization, given an instance of a problem the goal is to find a  
 43 solution optimizing the value of the objective function. However, in many applications the  
 44 instance may change over time and the goal is to find a tradeoff between the quality of the  
 45 solution in each time step and the stability of the solution in consecutive time steps. As  
 46 an example, consider an instance of an assignment problem, where the goal is to compute  
 47 the best assignment of tasks to workers, assuming that we know the cost  $c_{ij}$  of performing  
 48 task  $j$  by worker  $i$ . In the classical setting, it is possible to choose the assignment that  
 49 minimizes the total cost in polynomial time. When the costs change over time (as for  
 50 instance when a worker is not able to do some long task on a very busy day (infinite cost))  
 51 the optimal solutions of each time step may differ, inducing a transition cost for setting new  
 52 task-worker pairs between two consecutive solutions. Hence, the naïve approach of finding a  
 53 new optimal solution in each time step has the drawback that it does not take care of the  
 54 penalty (transition cost) that is induced by the changes in the solution.

55 In this paper we study a multistage version of the *Perfect Matching problem* that follows  
 56 this motivation and was originally introduced by Gupta, Talwar, and Wieder [11]. In this  
 57 problem we are given a time horizon:  $t = 1, 2, \dots, T$  where for each time  $t$  we are given an  
 58 instance  $G_t$  of *Perfect Matching* (that is, an edge-weighted graph) on the same set of vertices  
 59  $V$ . The goal is to determine a sequence of solutions  $S = (M_1, M_2, \dots, M_T)$  that both (1) are  
 60 near-optimal (*quality*), and (2) induce small transition costs (*stability*). In other words, the  
 61 goal is to determine a sequence of perfect matchings, one for each stage (time step)  $t$ , such  
 62 that their total cost is small and the solution does not change too radically from one step to  
 63 the next.

64 It was shown in [11] that this multistage problem is significantly harder than classical  
 65 *Perfect Matching*. In fact, it is NP-hard to even approximate the optimal solution within  
 66  $n^{1-\epsilon}$ , for instances with only 8 times steps. Gupta et al. then posed as an explicit question  
 67 whether the problem becomes easier for bipartite instances. Their work suggests also the  
 68 question whether this hardness also applies for fewer than 8 steps. The bipartite restriction  
 69 is especially interesting because Gupta et al. showed that related matroid-based optimization  
 70 problems remain tractable for  $T = 2$ , and bipartite *Perfect Matching* can be seen as a matroid  
 71 intersection problem. One could therefore hope that the matroid structure might make the  
 72 bipartite case tractable for some small values of  $T$ , or at least approximable.

73 Our main contribution in this paper is to settle this question from [11] in the negative:  
 74 we show that *Minimum Multistage Perfect Matching* (MIN-MPM) is  $n^{1-\epsilon}$ -inapproximable,  
 75 even for  $T = 2$  time steps, unless  $P = NP$ . Motivated by this very negative result, we  
 76 then investigate two other version of the problem: the *Metric Minimum Multistage Perfect*  
 77 *Matching* problem (METRIC-MIN-MPM), where the input is guaranteed to satisfy the triangle  
 78 inequality, and the *Maximum Multistage Perfect Matching* problem (MAX-MPM), where we  
 79 consider the complementary optimization objective.

80 **Problem definition.** Formally, the MIN-MPM problem is defined as follows: We are given  
 81 a sequence  $G_1, \dots, G_T$  of  $T$  undirected graphs, on the same set of vertices  $V$ . At each time  
 82 step  $1 \leq t \leq T$ , the graph  $G_t$  is given with a cost function  $c_t$  on edges:  $c_t(e) \in \mathbb{Q}_{\geq 0} \cup \{+\infty\}$ .  
 83 We are also given a transition cost  $M \geq 0$ . A solution is a sequence  $S = (M_1, \dots, M_t)$  where  
 84  $M_t$  is a perfect matching of  $G_t$ . Each solution (sequence) has two costs: a *matching cost*  
 85  $c(S)$  and a *transition cost*  $D(S)$ . The goal is to minimize  $c(S) + D(S)$ . A matching  $M_t$  has  
 86 a matching cost  $c_t(M_t)$  which is equal to the sum of the costs of the edges of the perfect

87 matching. The *matching cost* of  $S$  is  $c(S) = \sum_{t=1}^T c_t(M_t)$ . The *transition cost* is defined  
 88 as  $D(S) = \sum_{t=1}^{T-1} D_t$ , where  $D_t = M \cdot |M_{t+1} \setminus M_t|$  is proportional to the number of edges  
 89 removed between time  $t$  and  $t + 1$  – which is equal to the number of added edges since the  
 90 matchings are perfect. Notice that by allowing infinite cost on edges we may assume w.l.o.g.  
 91 the graphs to be complete.

92 In the METRIC-MIN-MPM, at each stage  $c_t$  obeys the triangle inequality:  $c_t(u, v) +$   
 93  $c_t(v, w) \geq c_t(u, w)$ . Finally, in the MAX-MPM version, we consider that  $c_t(e)$  is the *profit*  
 94 obtained by taking edge  $e$  (at time  $t$ ). Then a solution sequence  $S$  has a *matching profit*  
 95  $c(S) = \sum_t c_t(M_t)$ . We define the *transition profit*  $D(S)$  as  $D(S) = \sum_{t \leq T-1} D_t$  where  
 96  $D_t = M \cdot |M_{t+1} \cap M_t|$  is proportional to the number of edges that *remain* between time  $t$   
 97 and  $t + 1$ . The goal now is to maximize  $c(S) + D(S)$ . Notice that in MAX-MPM, we may  
 98 no longer assume that the graphs are complete, since this assumption modifies the problem  
 99 (we get profit by maintaining an edge, even of profit 0, from one time step to the next one).

100 **Related work.** A model that is close to our setting is the reoptimization model of Schieber  
 101 et al. [15]. In their work, they are given a starting solution and a new instance and the goal  
 102 is to minimize the sum of the cost of the new instance and of the transition cost. The model  
 103 of multistage optimization that we use in this work has been studied earlier by Buchbinder et  
 104 al. [5] and Buchbinder, Chen and Naor [4] for solving a set of fractional problems. Eisenstat  
 105 et al. [7] studied a similar multistage optimization model for facility location problems. Their  
 106 main result was a logarithmic approximation algorithm, which was later improved to a  
 107 constant factor approximation by An et al. [1]. More broadly, many classical optimization  
 108 problems have been considered in online or semi-online settings, where the input changes over  
 109 time and the algorithm tries to adjust the solution (re-optimize) by making as few changes  
 110 as possible. We refer the reader to [2, 3, 6, 10, 13, 14] and the references therein.

111 As mentioned, Gupta et al. [11] studied the Multistage Maintenance Matroid problem for  
 112 both the offline and the online settings. Their main result was a logarithmic approximation  
 113 algorithm for this problem, which includes as a special case a natural multistage version of  
 114 SPANNING TREE. The same paper also introduced the study of MIN-MPM, which is the  
 115 main problem we study here. They showed that the problem becomes hard to approximate  
 116 even for a constant number of stages. More precisely, they showed the following result ( $n$   
 117 denotes the number of vertices in the graphs).

118 ► **Theorem 1** ([11]). *For any  $\epsilon > 0$ , MIN-MPM is not  $n^{1-\epsilon}$ -approximable unless  $P = NP$ .  
 119 This holds even when the costs are in  $\{0, \infty\}$ ,  $M = 1$ , and the number of time steps is a  
 120 constant.*

121 Theorem 1 is proved for  $T = 8$ , starting from the fact that 3-colorability is NP-hard in  
 122 graphs of maximum degree 4 [8]. The authors leave as an open question the approximability of  
 123 the problem in bipartite graphs, and ask for subcases with better approximability behavior.

124 **Our contribution.** We answer the open question of [11] by showing that the problem is  
 125 hard to approximate even for bipartite graphs and for the case of two steps ( $T = 2$ ). Then,  
 126 we focus on the case where the edge costs are metric within every time step (METRIC-MIN-  
 127 MPM). On the negative side, we prove that the problem remains APX-hard even if  $T = 2$ .  
 128 On the positive side, we show that METRIC-MIN-MPM admits a 3-approximation algorithm  
 129 for two and three stages. Finally, for the maximization version of the problem, MAX-MPM,  
 130 we prove that it admits a constant factor approximation algorithm but is APX-hard.

131 **2** MIN-MPM for bipartite graphs

132 We answer the open question of [11] about the approximability of bipartite MIN-MPM.

133 ► **Theorem 2.** *For any  $\epsilon > 0$ , MIN-MPM cannot be approximated within a factor of  $n^{1-\epsilon}$ ,*  
 134 *even if the input has  $T = 2$  time steps, the input graphs are bipartite,  $M = 1$  and the costs*  
 135 *of edges are in  $\{0, \infty\}$ , unless  $P=NP$ .*

136 Using infinite costs, the same result immediately holds for bipartite complete graphs, as well  
 137 as for complete graphs.

138 **Proof.** We give a gap-introducing reduction from Perfect 3DM (3-Dimensional Matching),  
 139 known to be NP-complete [9]. We are given an instance of Perfect 3DM which consists of  
 140 three sets  $X, Y, Z$ , with  $|X| = |Y| = |Z| = n$ , and a set  $Q$  of elements of  $X \times Y \times Z$ , with  
 141  $|Q| = m \leq n^3$ . We are whether there exists a subset of  $n$  pair-wise disjoint elements of  $Q$ , or  
 142 not.

143 We construct an instance of our problem as follows: first, we create four sets of vertices  
 144  $A, B, C, D$  with  $|A| = |B| = n$  and  $|C| = |D| = m$ . To ease notation suppose that the ele-  
 145 ments of our sets  $X, Y, Z, Q, A, B, C, D$  are labeled as  $\{x_1, \dots, x_n\}, \{y_1, \dots, y_n\}, \{z_1, \dots, z_n\},$   
 146  $\{q_1, \dots, q_m\}, \{a_1, \dots, a_n\}, \{b_1, \dots, b_n\}, \{c_1, \dots, c_m\}$ , and  $\{d_1, \dots, d_m\}$  respectively.

147 For any  $j \in \{1, \dots, m\}$  we construct a set of  $2n^{\lceil \frac{4}{\epsilon} \rceil}$  new vertices. We connect  $c_j$  to  $d_j$   
 148 through a path traversing all these vertices (thus this is a path from  $c_j$  to  $d_j$  with  $2n^{\lceil \frac{4}{\epsilon} \rceil} + 2$   
 149 vertices). We set the cost of all the internal edges of these paths for both time-steps to 0.

150 For all  $i \in \{1, \dots, n\}, j \in \{1, \dots, m\}$  we do the following: if  $x_i \in q_j$  we set the cost of the  
 151 edge  $(a_i, c_j)$  to 0 in time step 1; if  $y_i \in q_j$  we set the cost of the edge  $(a_i, c_j)$  to 0 in time  
 152 step 2; if  $z_i \in q_j$  we set the cost of the edge  $(b_i, d_j)$  to 0 in both time steps. All other edge  
 153 costs are set to  $\infty$  (or some other sufficiently large value). This completes the construction.  
 154 Observe that the new graph has  $5n + 3m + 2mn^{\lceil \frac{4}{\epsilon} \rceil}$  vertices, so at most  $C \cdot n^{\frac{4}{\epsilon} + 4}$  (for some  
 155 constant  $C$ ) since  $m \leq n^3$ . Note also that the new graph is bipartite because the paths  
 156 that we added from  $c_j$  to  $d_j$  have odd lengths, hence the bipartition  $(A \cup D, B \cup C)$  can be  
 157 extended to a bipartition of the whole graph.

158 Suppose that the original instance has a set  $Q' \subseteq Q$  such that  $|Q'| = n$  and no element of  
 159  $X \cup Y \cup Z$  appears in two elements of  $Q'$ . We obtain a multistage matching as follows: For  
 160 each  $q_j \in Q'$  such that  $q_e = (x_{i_1}, y_{i_2}, z_{i_3})$  we use the edge  $(a_{i_1}, c_j)$  in step 1, the edge  $(a_{i_2}, c_j)$   
 161 in step 2, and the edge  $(b_{i_3}, d_j)$  in both time steps. Note that this fully specifies how the  
 162 vertices of  $A \cup B$  are matched. We now complete the matching by selecting a set of edges  
 163 from the paths connecting each  $c_j$  to  $d_j$ : if  $q_j \in Q'$ , then both  $c_j, d_j$  have been matched  
 164 to  $A \cup B$  in both time steps, and we select in both time steps the unique perfect matching  
 165 of the path connecting them; if  $q_j \notin Q'$ , then neither  $c_j, d_j$  is matched to  $A \cup B$  in either  
 166 time step, so we select the perfect matching on the path from  $c_j$  to  $d_j$ , including these two  
 167 vertices. Observe that the cost of all edges we use is 0, while we only change at most  $n$  edges  
 168 from one time step to the other, hence the total transition cost is at most  $nM$ .

169 Suppose that the original instance does not have a solution and consider any multistage  
 170 matching in the new instance. We will show that it must make at least  $n^{\frac{4}{\epsilon}}$  changes from one  
 171 time step to the other. We will say that  $q_j \in Q$  is selected in time step 1, if in that time step  
 172  $c_j$  is matched to an element of  $A$ . If  $q_j$  is selected in time step 1, then  $d_j$  is matched to an  
 173 element of  $B$  in that time step, otherwise it would be impossible to have a perfect matching  
 174 on the path connecting  $c_j$  to  $d_j$ . If some  $q_j$  is selected in time step 1, but not in time step 2,  
 175 then the solution must change all internal edges on the perfect matching on the path from  
 176  $c_j$  to  $d_j$ , hence it makes at least  $n^{\frac{4}{\epsilon}}$  changes, and we are done. What remains therefore to

show is that if the solution maintains the set of selected  $q_j$  in the two time steps, then we can construct a solution to the original instance. Indeed, since all of  $A \cup B$  is matched, we have  $n$  selected  $q_j$ 's. Each element of  $C \cup D$  has at most one edge connecting it to  $A \cup B$  in each step, hence if it is selected this edge must be used. But if we select  $q_{j_1}, q_{j_2}$  that overlap, then two selected elements will have a common neighbor in  $A \cup B$  and will therefore not be matched, contradiction.

Since the new graph has  $N$  vertices with  $n^{\frac{1}{\epsilon}} \leq N \leq Cn^{\frac{4}{\epsilon}+4}$  vertices, it is NP-hard to distinguish if the optimal is at most  $nM \leq N^\epsilon M$  or at least  $n^{\frac{4}{\epsilon}} M \geq N^{1-\epsilon} M/C$ . ◀

### 3 METRIC-MIN-MPM

We consider in this section that  $c_t$  obeys the triangle inequality:  $c_t(u, v) + c_t(v, w) \geq c_t(u, w)$ . In particular, the graph is complete. As seen before, the problem is hard to approximate even if there are only 2 time steps with general costs. We show here that while the problem is APX-hard in the metric case even with only 2 time steps (Section 3.1), it admits a 3-approximation algorithm in this case (2 time steps), see Section 3.2. We then extend this last result to the case of 3 time steps in Section 3.3.

#### 3.1 APX-hardness for 2 time steps

In the case of 2 time steps the following result is proved.

► **Theorem 3.** METRIC-MIN-MPM is APX-hard, even if the input has  $T = 2$  time steps.

**Proof.** We give a gap-preserving reduction from Max 3DM. We are given an instance of Max 3DM which consists of three sets  $X, Y, Z$ , with  $|X| = |Y| = |Z| = n$ , a set  $Q$  of elements of  $X \times Y \times Z$ , with  $|Q| = m$ , and an integer  $k$ . We are asked if there exists a subset of  $k$  pair-wise disjoint elements of  $Q$ . We assume that  $n, m$  and  $k$  are even (if not simply make two independent copies of the initial instance). This problem is APX-hard even if the occurrence of each element is bounded above by a constant  $C = 3$  [12]. Note that in this case the optimum value is at least  $m/7$  (greedy algorithm; at most 6 incompatible triplets are removed when a triplet is chosen). So  $m, n$  and  $k$  are linearly related ( $3n \geq m \geq k \geq m/7 \geq n/21$ ).

We construct an instance of METRIC-MIN-MPM as follows: first, we create five sets of vertices  $X, Y, Z, G, D$  with  $X = \{x_1, \dots, x_n\}$ ,  $Y = \{y_1, \dots, y_n\}$ ,  $Z = \{z_1, \dots, z_n\}$ ,  $G = \{g_1, \dots, g_m\}$  and  $D = \{d_1, \dots, d_m\}$ .


The graph is complete, and we set the following costs:

- At time step 1,  $Z$  is seen as a single point very far from the rest of the graph:  $(z_i, z_j)$  has cost 0 for  $z_i, z_j \in Z$ , and  $(z_i, v)$  has infinite cost for  $z_i \in Z, v \notin Z$ .
- The same is done for  $X$  at time 2.
- The  $m$  edges  $(g_i, d_i)$  have cost 1 at both time steps.
- For each triplet  $q_i = (x_j, y_p, z_s)$ : at time 1 edges  $(x_j, g_i)$  and  $(d_i, y_p)$  have cost  $a$  ( $a$  is a sufficiently large constant, to be specified later), and, for the triangle inequality to hold,  $(x_j, d_i)$  and  $(g_i, y_p)$  have cost  $a + 1$ . Similarly at time 2:  $(z_s, g_i)$  and  $(d_i, y_p)$  have cost  $a$ , and, for the triangle inequality to hold,  $(z_s, d_i)$  and  $(g_i, y_p)$  have cost  $a + 1$ .

All non yet defined costs are equal to  $2a$ . The transition cost is  $M = 1$ . Figure 1 gives an illustration of the construction.

Note that the triangle inequality holds in both time steps.

We show that (1) if there is a 3DM of size  $k$  then there exists a solution of METRIC-MIN-MPM whose total cost is at most  $2m + 4an - k/2$ , and (2) conversely from a solution of the



■ **Figure 1** An illustration of the reduction at time  $t = 1$ , without representing  $Z$  - the construction is symmetric for time step  $t = 2$ . The third element of  $X$  is in the first, third and last triplet of  $Q$ . The second element of  $Y$  is in the second and third triplet. The dashed edges have costs  $a + 1$ . Not represented edges have cost  $2a$ .

220 multistage problem of total cost  $z$  we can construct a 3DM of size at least  $2(2m + 4an - z)$ .  
 221 This proves  $APX$ -hardness since  $a$  is a constant, and  $m$ ,  $n$  and  $k$  are linearly related.

222 Let us first prove (1), and suppose that we have a 3DM of size  $k$ , say (for ease of notation)  
 223  $q_1, \dots, q_k$  where  $q_i = (x_i, y_i, z_i)$ . Then we define a solution  $S$  of the multistage matching as  
 224 follows:

- 225 ■ We take the  $(m - k)$  edges of triplets  $(g_j, d_j)$  *not* in the 3DM, at both time steps 1 and 2;
- 226 ■ For  $q_i, 1 \leq i \leq k$ : we take edges  $(x_i, g_i)$  at time 1,  $(z_i, g_i)$  at time 2, and  $(y_i, d_i)$  at time 1  
 227 and 2.
- 228 ■ We match together the  $(n - k)$  remaining vertices of  $Y$ , choosing the same  $\frac{n-k}{2}$  edges at  
 229 both time steps.
- 230 ■ We match together the  $(n - k)$  remaining vertices of  $X$  at time 1. At time 2 we keep  
 231 these  $\frac{n-k}{2}$  edges and match the remaining  $k$  vertices of  $X$  together.
- 232 ■ We do the same for  $Z$ .

233 We get a solution  $(M_1, M_2)$  whose costs are:

- 234 ■ At time 1, the matching cost is  $(m - k) + 2ak + 2a\frac{n-k}{2} + 2a\frac{n-k}{2} = m + 2an - k$ ;
- 235 ■ The matching cost at time 2 is the same.
- 236 ■ The number of modifications is  $3k/2$ :  $k$  edges  $(x_i, g_i)$  become  $(z_i, g_i)$ , and  $k/2$  edges in  $Z$   
 237 disappear at time 1 ( $k/2$  edges appear in  $X$  at time 2).

238 In all,  $(M_1, M_2)$  has cost  $2m + 4an - k/2$ .

239

240 Conversely, suppose that we have a solution  $(M_1^0, M_2^0)$  of total cost  $z$  for the instance  
 241 of METRIC-MIN-MPM. We first structure this solution using local modifications, and then  
 242 show how to derive a matching from it.

- 243 ■ *Replacement 1.* First, suppose that  $M_1^0$  takes (at time 1) an edge  $(x_j, g_i)$  of cost  $2a$  - so  
 244  $x_j$  is not in the  $i$ -th triplet  $q_i$  of  $Q$ . Then  $d_i$  is matched with a vertex  $v$  with an edge  
 245 of cost at least  $a$ . By replacing (at time 1)  $(x_j, g_i)$  and  $(d_i, v)$  by  $(x_j, v)$  and  $(g_i, d_i)$  we  
 246 get a matching cost for these two edges at most  $2a + 1$  instead of (at least)  $3a$ . Even

247 considering that the transition cost may have increased by two, this replacement does  
 248 not increase the cost of the solution for  $a \geq 3$ . The same argument applies for an edge  
 249  $(x_j, d_i)$  (time step 1), an edge  $(y_j, d_i)$  or  $(y_j, g_i)$  (time step 1 or 2) and for an edge  $(z_j, g_i)$   
 250 or  $(z_j, d_i)$  in  $M_2^0$ .

251 ■ *Replacement 2.* Now, suppose that  $M_1^0$  takes an edge of cost  $2a$  in  $G \cup D$ , say  $(g_i, g_j)$   
 252 with  $i \neq j$  (the very same argument works for the 2 other cases  $(g_i, d_j)$  and  $(d_i, d_j)$ ).  
 253 Let  $v$  and  $w$  be the neighbors of  $d_i$  and  $d_j$  in  $M_1^0$ . By replacing the three edges  $(g_i, g_j)$ ,  
 254  $(d_i, v)$  and  $(d_j, w)$  by  $(g_i, d_i)$ ,  $(g_j, d_j)$  and  $(v, w)$ , we get a matching cost at most  $(2a + 2)$   
 255 instead of (at least)  $4a$ . Even considering that the transition cost may have increased by  
 256 three, this replacement does not increase the cost of the solution for  $a \geq 5/2$ . The same  
 257 holds for  $M_2^0$ .

258 ■ *Replacement 3.* Last, suppose that edges  $(y_j, g_i)$  and  $(y_s, d_i)$  are both taken at time 1  
 259 and 2. This costs  $2(a + a + 1) = 4a + 2$ . Then we can take instead edges  $(g_i, d_i)$  and  
 260  $(y_j, y_s)$  at both time steps, with the same cost  $2 + 2(2a) = 4a + 2$ .

261 In this way, we transform  $(M_1^0, M_2^0)$  into a solution  $(M_1, M_2)$  of cost at most  $z$  such that:

- 262 ■ No  $g_i$  (and no  $d_i$ ) is matched using an edge of cost  $2a$  (replacements 1 and 2).
- 263 ■  $g_i$  and  $d_i$  cannot be both matched to the same vertices at time 1 and 2, unless they are  
 264 matched together (replacement 3).

265 We now show how to find a 3DM from this solution  $(M_1, M_2)$ . Let:

266 ■  $N_x$  and  $N_z$  be respectively the number of edges in  $X \times (G \cup D)$  at time 1 and in  $Z \times (G \cup D)$   
 267 at time 2.

268 ■  $N_y^1$  and  $N_y^2$  be respectively the number of edges in  $Y \times (G \cup D)$  at time 1 and time 2,  
 269 among which  $\lambda_1$  (resp.,  $\lambda_2$ ) are of cost  $a + 1$ .

270 ■  $N_y$  be the number of edges in  $Y \times (G \cup D)$  that are taken at both times 1 and 2.

271 At time 1, besides these  $N_x + N_y^1$  edges and the  $n/2$  edges of cost 0 (vertices of  $Z$ ), the  
 272 other edges of  $(M_1, M_2)$  have cost either 1 (edges  $(g_i, d_i)$ ) or  $2a$ . Since  $N_x + N_y^1$  vertices in  
 273  $G \cup D$  are already matched at time 1, there are at most  $\frac{2m - N_x - N_y^1}{2}$  edges of cost 1 at time 1.

274 Similarly, there are at most  $\frac{2m - N_z - N_y^2}{2}$  edges of cost 1 at time 2.

275 Then, computing the matching cost of  $(M_1, M_2)$  we have

$$\begin{aligned}
 276 \quad c(M_1, M_2) &\geq a(N_x + N_z + N_y^1 + N_y^2) + \lambda_1 + \lambda_2 + \frac{4m - N_x - N_z - N_y^1 - N_y^2}{2} \\
 277 &\quad + 2a \left( \frac{n - N_x + n - N_y^1 + n - N_z + n - N_y^2}{2} \right) \\
 278 &\geq 2m + 4na + \lambda_1 + \lambda_2 - \frac{N_x + N_z + N_y^1 + N_y^2}{2}.
 \end{aligned}$$

279 Now, note that at time 1 at least  $N_x + N_y^1 - N_y + \frac{N_z}{2}$  edges disappear, so  $D(M_1, M_2) \geq$   
 280  $N_x + N_y^1 - N_y + \frac{N_z}{2}$ . Similarly, at least  $N_z + N_y^2 - N_y + \frac{N_x}{2}$  edges appear at time 2. So  
 281  $D(M_1, M_2) \geq N_z + N_y^2 - N_y + \frac{N_x}{2}$ . Then,

$$D(M_1, M_2) \geq \frac{N_x + N_z + N_y^1 + N_y^2}{2} - N_y + \frac{N_x + N_z}{4}.$$

282 This gives:

$$283 \quad z \geq c(M_1, M_2) + D(M_1, M_2) \geq 2m + 4na + \lambda_1 + \lambda_2 - N_y + \frac{N_x + N_z}{4}.$$



## 7:8 Multistage Matchings

284 Now, consider the set of indices  $i$  such that edge  $(y_j, d_i)$  is taken at both time steps, or  
 285 edge  $(y_j, g_i)$  is taken at both time steps. Since, thanks to the preprocessing, for a given  $i$   
 286 this cannot concern both  $d_i$  or  $g_i$ , we know that there are exactly  $N_y$  such indices (edges).  
 287 Since there are  $\lambda_1 + \lambda_2$  edges of cost  $a + 1$  between  $Y$  and  $G \cup D$ , among these  $N_y$  indices at  
 288 least  $N_y - (\lambda_1 + \lambda_2)$  are such that: (1) edge  $(d_i, y_j)$  is used at both time steps (2) an edge  
 289  $(x_s, g_i)$  of cost  $a$  is used at time 1 (since no edge of cost  $2a$  is used for vertices in  $G$ ) and (3)  
 290 an edge  $(z_p, g_i)$  of cost  $a$  is used at time 2.

In other words these at least  $N_y - (\lambda_1 + \lambda_2)$  indices correspond to triplets of a 3DM. So  
 we have a 3DM of size (at least)  $k = N_y - (\lambda_1 + \lambda_2)$ . Then,  $N_x \geq N_y - (\lambda_1 + \lambda_2) = k$  and  
 similarly  $N_z \geq k$ , so  $\frac{N_x + N_z}{4} \geq \frac{k}{2}$ . All together, we get

$$z \geq 2m + 4an - k + \frac{k}{2} = 2m + 4an - \frac{k}{2}.$$

291

### 292 3.2 A 3-approximation algorithm for 2 time steps

293 We now devise an approximation algorithm. Informally, this algorithm first guesses the  
 294 number  $k$  of edges that an optimal solution keeps between steps 1 and 2. Then it computes  
 295 a set of  $k$  edges with low matching cost that it maintains between time 1 and 2. Finally,  
 296 it completes this set of  $k$  edges into two perfect matchings, in such a way that, using the  
 297 triangle inequality, the matching cost does not increase too much.

298 Formally, the algorithm **Metric2** runs the following procedure for  $k$  from 0 to  $n/2$ .

- 299 1. Let  $G_{1+2}$  be the graph where the edge costs are  $c(u, v) = c_1(u, v) + c_2(u, v)$ . Compute a  
 300 minimum cost matching  $M^k$  of size exactly  $k$  in  $G_{1+2}$ .
- 301 2. Compute a minimum cost perfect matching  $M_1$  in  $G_1$ , and a minimum cost perfect  
 302 matching  $M_2$  in  $G_2$ .
- 303 3. Consider the symmetric difference of the two matchings  $M^k$  and  $M_1$  in  $G_1$ . This is a  
 304 (vertex disjoint) set of paths  $P_1, \dots, P_p$  and cycles. Define  $M_1^k$  as  $M^k$  plus the  $p$  edges  
 305 linking the first vertex and last vertex of each path  $P_j$ .
- 306 4. Do the same to get  $M_2^k$ .
- 307 5. Consider  $S^k = (M_1^k, M_2^k)$ .

308 **Metric2** outputs the best solution  $S^k$ .

309

310 ► **Theorem 4.** *Metric2* is a (polytime) 3-approximation algorithm for METRIC-MIN-MPM  
 311 when  $T = 2$ .

312 **Proof.** We first prove that  $S^k$  is a feasible solution, i.e.,  $M_i^k$  is a perfect matching of  $G_i$ .  
 313 Since  $M_i$  is a perfect matching, in all paths  $P_j$  the first and last edges belong to  $M_i$ . Hence  
 314 the first and last vertices are not covered by  $M^k$ , so  $M_1^k$  is a matching. Every other vertex is  
 315 covered by  $M^k$ , so the matching is perfect.

316 Now, let us prove the claimed approximation ratio. Let us denote  $S^* = (M_1^*, M_2^*)$  be an  
 317 optimal solution, and consider  $S^k$  where  $k = |M_1^* \cap M_2^*|$ .

318 Since at least  $M^k$  is common between  $M_1^k$  and  $M_2^k$ , at least  $k$  edges are maintained  
 319 between time 1 and 2 in  $S^k$ , as in  $S^*$ . So:

$$320 \quad D(S^k) \leq D(S^*). \tag{1}$$

321 Now, let us prove that:

$$322 \quad c_1(M_1^k) + c_2(M_2^k) \leq 3c_1(M_1^*) + 3c_2(M_2^*). \quad (2)$$

323 Thanks to the triangle inequality, in a path  $P = (v_0, v_1, \dots, v_t)$ ,  $c_i(v_0, v_t) \leq \sum_j c_i(v_j, v_{j+1})$ :  
 324 when adding edges  $(v_0, v_t)$  we add in total at most the total length of the paths, hence at  
 325 most  $c_i(M_i) + c_i(M^k)$ . So  $c_i(M_i^k) \leq c_i(M_i) + 2c_i(M^k)$ . Using that  $c_i(M_i) \leq c_i(M_i^*)$ , we get:

$$c_1(M_1^k) + c_2(M_2^k) \leq c_1(M_1^*) + c_2(M_2^*) + 2(c_1(M^k) + c_2(M^k)).$$

326 By optimality of  $M^k$  and since  $S^*$  has  $k$  common edges between times 1 and 2, these  $k$   
 327 common edges induce a cost in  $S^*$  at least  $c_1(M^k) + c_2(M^k)$ . Then:

$$c_1(M_1^k) + c_2(M_2^k) \leq c_1(M_1^*) + c_2(M_2^*) + 2(c_1(M_1^*) + c_2(M_2^*))$$

and Equation 2 follows. From Equations 1 and 2 we derive:

$$c(S) + D(S) \leq 3c(S^*) + D(S^*).$$

328 The result immediately follows. ◀

### 329 3.3 A 3-approximation algorithm for 3 time steps

330 We now extend the previous result to the case of  $T = 3$ . As previously, if an optimal solution  
 331 preserves in total  $k$  edges (operates in total  $n - k$  modifications between time steps 1 and 2,  
 332 and 2 and 3) we would like to first compute a set of  $k$  ‘preserved’ edges inducing a low cost,  
 333 and then to complete this set as perfect matchings in each of the time steps. Now things get  
 334 more complex since an edge can be preserved between steps 1 and 2, between steps 2 and 3,  
 335 or during the whole process. It seems hard to mimic an optimal solution on these 3 types of  
 336 edges (while inducing a low matching cost), but this difficulty can be overcome as follows.

337 Let  $G$  be the graph with edge cost  $w = \min\{c_1 + c_2 + c_3, c_1 + c_2 + M, c_2 + c_3 + M\}$ . If  
 338 the minimum is  $c_1 + c_2 + c_3$  (resp.,  $c_1 + c_2 + M$ ,  $c_2 + c_3 + M$ ) we say that the edge is of type  
 339 1 (resp., 2, 3). Intuitively, edges of type 1 will be taken in steps 1, 2 and 3, edges of type  
 340 2 (resp., 3) will be taken in steps 1 and 2 (resp., 2 and 3). We present a 3-approximation  
 341 algorithm **Metric3**. It runs the following procedure for  $k$  from 0 to  $n/2$ .  
 342

- 343 1. Compute a minimum cost matching  $M^k$  of size exactly  $k$  in  $G$ . Denote  $M_1^k$  the set of  
 344 edges of  $M^k$  of type 1 or 2,  $M_2^k = M^k$  and  $M_3^k$  the set of edges of  $M^k$  of type 1 or 3.
- 345 2. Compute a minimum cost perfect matching  $M_i$  in  $G_i$ ,  $i = 1, 2, 3$ .
- 346 3. Consider the symmetric difference of the two matchings  $M_i^k$  and  $M_i$  in  $G_i$ . This is a  
 347 (vertex disjoint) set of paths  $P_1, \dots, P_p$  and cycles. Define  $M_i'^k$  as the set of  $p$  edges  
 348 linking the first vertex and last vertex of each path  $P_j$ .
- 349 4. Consider  $S^k = (M_1^k \cup M_1'^k, M_2^k \cup M_2'^k, M_3^k \cup M_3'^k)$ .

350 Then **Metric3** outputs the best solution  $S^k$ .

351

352 ▶ **Theorem 5.** *Metric3* is a (polytime) 3-approximation algorithm for METRIC-MIN-MPM  
 353 when  $T = 3$ .

## 7:10 Multistage Matchings

354 **Proof.** We first note that, as in the case for  $T = 2$  time steps,  $M_i^k \cup M_i'^k$  is a perfect matching  
 355 of  $G_i$ , so  $S^k$  is a feasible solution.

356 Now let us deal with the approximation ratio. Let  $S^* = (M_1^*, M_2^*, M_3^*)$  be an optimal  
 357 solution. Let us consider the set  $H = (M_1^* \cap M_2^*) \cup (M_2^* \cap M_3^*)$  of edges in  $S^*$  that are in (at  
 358 least) two consecutive steps. Note that  $H$  is a matching (it is included in  $M_2^*$ ). Consider  $S^k$   
 359 where  $k = |H|$ . We now prove the following result:

360 ► **Lemma 6.**  $D(S^k) + \sum_i c_i(M_i^k) \leq D(S^*) + c(S^*)$ .

361 **Proof.** To prove this, let  $k_1 = |M_1^* \cap M_2^* \cap M_3^*|$  be the number of edges in  $S^*$  that are taken  
 362 at each of the 3 time steps. Hence,  $k - k_1$  edges are taken at (only) 2 consecutive time steps.  
 363 So there are  $(n/2 + n/2 - 2k_1 - (k - k_1))$  modifications in total, and:

$$364 \quad D(S^k) = M(n - k - k_1). \quad (3)$$

365 Recall that in  $G$ ,  $w = \min\{c_1 + c_2 + c_3, c_1 + c_2 + M, c_2 + c_3 + M\}$ .  $k_1$  edges of  $H$  are present  
 366 on the 3 time steps (matching cost  $c_1 + c_2 + c_3$ ), while  $k - k_1$  are present in two consecutive  
 367 time steps (matching cost  $c_1 + c_2$  or  $c_2 + c_3$ ).

$$368 \quad w(H) \leq c(S^*) + M(k - k_1). \quad (4)$$

369 Similarly, let  $\lambda_1$  be the number of edges of type 1 in  $M^k$ . There are  $(k - \lambda_1)$  edges of  
 370 type 2 or 3, hence

$$371 \quad w(M^k) = \sum_i c_i(M_i^k) + M(k - \lambda_1). \quad (5)$$

372 Indeed, in  $G$  cost  $c_1$  applies to edges of type 1 and 2 ( $c_1(M_1^k)$ ), cost  $c_2$  applies to all edges of  
 373  $M^k$  ( $c_2(M_2^k)$ ), cost  $c_3$  applies to edges of type 1 and 3 ( $c_3(M_3^k)$ ), and cost  $M$  to the  $(k - \lambda_1)$   
 374 edges of type 2 and 3.

375 Also, the number of preserved edges in  $S^k$  is at least  $k + \lambda_1$ , so:

$$376 \quad D(S^k) \leq M(n - k - \lambda_1). \quad (6)$$

377 Since  $H$  is a matching, in  $G$  we have  $w(H) \geq w(M^k)$ . This gives using Equations 4 and 5:

$$\sum_i c_i(M_i^k) + M(k - \lambda_1) \leq c(S^*) + M(k - k_1)$$

378 so  $\sum_i c_i(M_i^k) \leq c(S^*) + M(\lambda_1 - k_1)$ . Then using Equations 3 and 6 we get:

$$\begin{aligned} \sum_i c_i(M_i^k) + D(S^k) &\leq c(S^*) + M(\lambda_1 - k_1) + M(n - k - \lambda_1) = c(S^*) + M(n - k - k_1) \\ &= c(S^*) + D(S^*) \end{aligned}$$

379 which concludes the proof of Lemma 6. ◀

380 Now, by triangle inequality, and the fact that  $c_i(M_i) \leq c_i(M_i^*)$ , we know that:

$$381 \quad c_i(M_i'^k) \leq c_i(M_i^*) + c_i(M_i^k). \quad (7)$$

382 Then, from Lemma 6 and Equation 7 we get:

$$\begin{aligned}
 383 \quad c(S^k) + D(S^k) &= \sum_i (c_i(M_i^k) + c_i(M_i'^k)) + D(S^k) \leq \sum_i (2c_i(M_i^k) + c_i(M_i^*)) + D(S^k) \\
 384 \quad &\leq c(S^*) + 2 \left( \sum_i c_i(M_i^k) + D(S^k) \right) \leq 3c(S^*) + 2D(S^*).
 \end{aligned}$$

385 The result follows. ◀

## 386 **4** MAX-MPM

387 In the maximization version, we consider that  $c_t(e)$  is the *profit* obtained by taking edge  $e$  (at  
 388 time  $t$ ). Then a solution sequence  $S$  has a matching profit  $c(S) = \sum_t c_t(M_t)$ . We define the  
 389 transition profit  $D(S)$  as  $D(S) = \sum_{t \leq T-1} D_t$  where  $D_t = M \cdot |M_{t+1} \cap M_t|$  is proportional  
 390 to the number of edges that *remain* between time  $t$  and  $t+1$ . The goal now is to maximize  
 391  $c(S) + D(S)$ . Recall that in the maximization version we may no longer assume that the  
 392 graphs are complete.

### 393 **4.1 APX-hardness for 2 time steps**

394 We first show that MAX-MPM, even in the case of 2 time steps is APX-hard.

395 ▶ **Theorem 7.** MAX-MPM is APX-hard even if  $T = 2$ .

396 **Proof.** As previously, we consider the maximum 3DM problem in the case where the  
 397 occurrence of each element is bounded by 3, hence the optimal value, the number of triplets  
 398 and the size of the ground sets are linearly related.

399 Given three sets  $X, Y, Z$  each of size  $n$ , and  $m$  triplets  $q_i$  of  $X \times Y \times Z$ , we build two  
 400 graphs  $G_1$  and  $G_2$  with  $n' = 2m + 4n$  vertices:

- 401 ■ 4 sets  $D, E, F, G$  of size  $n$ ;
- 402 ■ 2 sets  $A = \{a_1, \dots, a_m\}$  and  $B = \{b_1, \dots, b_m\}$  of size  $m$ .

403 Vertices of  $D$  will represent elements of  $X$ , vertices of  $E$  and  $F$  elements of  $Y$  (twice), vertices  
 404 of  $G$  elements of  $Z$ . Each triplet  $q_i$  is represented by one edge  $(a_i, b_i)$  in both graphs. It has  
 405 cost 0.

406 If a triplet  $q_i$  is  $(x_j, y_k, z_l)$  then:

- 407 ■ In  $G_1$  we put edges  $(d_j, a_i)$  and  $(b_i, e_k)$ , both with cost  $M'$ ;
- 408 ■ In  $G_2$  we put edges  $(f_k, a_i)$  and  $(b_i, z_l)$ , both with cost  $M'$ .

409 Note that vertices in  $F, G$  have degree 0 in  $G_1$ , vertices in  $D, E$  have degree 0 in  $G_2$ .

410 We fix  $M' = \frac{M+1}{4}$ , and  $M \geq 3$ .

411 Let us show that there is a 3DM of size (at least)  $k$  if and only if there is a solution of  
 412 profit at least  $Mm + k$ .

413 Suppose first that there is a set  $S$  of  $k$  independent triplets. Then we build matchings  
 414  $(M_1, M_2)$  as follows:

- 415 ■ if  $q_i$  is not in  $S$ , we take  $(a_i, b_i)$  both in  $M_1$  and  $M_2$ . This gives transition profit  $M(m-k)$ .
- 416 ■ if  $q_i = (x_j, y_k, z_l)$  is in  $S$ , then we take in  $M_1$  the two edges  $(d_j, a_i)$  and  $(b_i, e_k)$ , and in  
 417  $M_2$  the two edges  $(f_k, a_i)$  and  $(b_i, z_l)$ . This gives a matching profit  $4kM'$ .

418 Note that since any element of  $X, Y, Z$  is in at most one triplet of  $S$ , vertices in  $D, E, F, G$   
 419 are adjacent to at most one chosen edge. In other words  $M_1$  and  $M_2$  are matchings.

420 The profit of the solution is  $4kM' + M(m-k) = k(M+1) + M(m-k) = Mm + k$ .

421 Suppose now that there is a solution  $(M_1, M_2)$  of profit at least  $Mm + k$ . Suppose first  
 422 that there is an edge  $(a_i, b_i)$  which is in  $M_1$  but not in  $M_2$ . Then we get no transition profit  
 423 for this edge. In  $M_2$  we have taken at most one edge incident to  $a_i$ , and one edge incident  
 424 to  $b_i$ , with matching profit at most  $2M'$ . Since these edges are not in  $G_1$  they cannot give  
 425 transition profit. So we can put in  $M_2$  the edge  $(a_i, b_i)$  and remove the edges incident to  $a_i$   
 426 and  $b_i$  (if any). The profit increases by  $M - 2M' = M/2 - 1/2 \geq 0$ .

427 So we can assume that  $M_1$  and  $M_2$  have the same set of edges between  $A$  and  $B$ .  
 428 Suppose now that there are two edges  $(a_i, b_i)$  and  $(a_s, b_s)$  both not in  $M_1$  (equiv. not in  $M_2$ )  
 429 corresponding to two intersecting triplets. Suppose for instance that  $x_j$  is in both triplets.  
 430 This means that in  $M_1$  we cannot take both edges  $(c_j, a_i)$  and  $(c_j, a_s)$ , for instance  $(c_j, a_s)$  is  
 431 not in  $M_1$ . Then we can add  $(a_s, b_s)$  to  $M_1$  and  $M_2$ , and remove the (at most) 3 incident  
 432 edges. This increases profit by  $M - 3M' \geq 0$ .

433 So, the set of edges  $(a_i, b_i)$  not in  $M_1$  (or not in  $M_2$ ) corresponds to a set of independent  
 434 triplets. Let  $t$  the number of such edges. Since  $M_1$  is a matching, besides these edges between  
 435  $A$  and  $B$ , there is at most two edges for each  $(a_i, b_i)$  not in  $M_1$ . Similarly, there is at most  
 436 two edges in  $M_2$  for each  $(a_i, b_i)$  not in  $M_2$ . So the matching profit is at most  $4tM'$ , and  
 437 the transition profit is  $M(m - t)$ . The profit is  $M(m - t) + 4tM' = Mm + t \geq Mm + k$ . So  
 438  $t \geq k$ . ◀

## 439 4.2 Constant factor approximation algorithms

440 ► **Theorem 8.** MAX-MPM is  $1/2$ -approximable. If  $T = 2$  it is  $2/3$ -approximable, if  $T = 3$   
 441 it is  $3/5$ -approximable.

442 **Proof.** Note that if the graphs are assumed to be complete (bipartite complete) then the  
 443 ratio  $1/2$  is easily achievable. Indeed, consider two solutions:

- 444 ■ The first one  $S_1$  consisting of the same perfect matching  $M_0$  at all time steps;
- 445 ■ The second one  $S_2$  consisting of a matching  $\hat{M}_t$  of maximum profit on  $G_t$  for each  $t$ .

446 Output the best one.

447 Let  $S^* = (M_1^*, \dots, M_T^*)$  be an optimal solution. Clearly the profit of  $S_1$  is at least the  
 448 transition profit  $D(S^*)$  of  $S^*$ . Also,  $c(M_t^*) \leq c(\hat{M}_t)$  so the matching profit of  $S^*$  is at most  
 449 the one of  $S_2$ . The ratio  $1/2$  follows.

450 If the graphs are not assumed to be complete things get harder since one cannot trivially  
 451 optimize the transition profit by keeping a perfect matching along the multistage process.

452 Let us consider three consecutive time steps  $t - 1, t, t + 1$ . Let us consider the graph  $G'_t$   
 453 which is the same as  $G_t$  up to the profit on edges, which is now  $c'_t(e)$  where:

- 454 1.  $c'_t(e) = c_t(e) + 2M$  if  $e$  is in  $G_{t-1}$  and  $G_{t+1}$ ;
- 455 2. otherwise,  $c'_t(e) = c_t(e) + M$  if  $e$  is in  $G_{t-1}$  or  $G_{t+1}$ ;
- 456 3. otherwise  $c'_t(e) = c_t(e)$ .

457 Let us consider a matching  $M'_t$  of maximum profit in  $G'_t$ .

458 ► **Lemma 9.**  $c'_t(M'_t) \geq D_{t-1}(S^*) + c_t(M_t^*) + D_t(S^*)$ .

459 **Proof.** Let us consider the profit of  $M'_t$  on  $G'_t$ . Since the set of edges preserved from time  
 460  $t - 1$  to time  $t$  is included in  $M_t^*$ , the profit  $D_{t-1}(S^*)$  appears in the profit of  $M'_t$  on  $G'_t$   
 461 ( $+M$  on each common edges between the two consecutive graphs). This is also the case  
 462 for  $D_t(S^*)$ , for the same reason. Of course, the profit  $c_t(e)$  appears as well. Since  $M'_t$  is of  
 463 maximum profit, the Lemma follows. ◀

464 Because of Lemma 9, choosing the matching  $M'_t$  at time steps  $t - 1$ ,  $t$  and  $t + 1$  in a  
 465 solution generates a profit at least  $D_{t-1}(S^*) + c_t(M'_t) + D_t(S^*)$ .

466 Note that, with similar arguments, if two times steps  $t, t + 1$  are involved, we can compute  
 467 a matching  $H_i$  that we take at time steps  $t, t + 1$  generating a profit at least  $c_t(M'_t) + D_t(S^*)$ .  
 468 Symmetrically, we can compute a matching  $H'_i$  that we take at time steps  $t, t + 1$  generating  
 469 a profit at least  $c_{t+1}(M'_t) + D_t(S^*)$ .

470

471 Now we consider the following 2 solutions:

472 ■  $S_1$  consists of choosing  $H_1$  at steps 1, 2,  $H_3$  at step 3, 4, . . . . If  $T$  is even then we are  
 473 done, otherwise we take an optimal matching  $\hat{M}_T$  at step  $T$ .

474 ■  $S_2$  consisting of choosing an optimal matching  $\hat{M}_1$  at step 1, then  $H_2$  at steps 2, 3,  $H_4$  at  
 475 steps 4, 5, . . . . If  $T$  is even we take an optimal matching  $\hat{M}_T$  at step  $T$ .

476 Output the best of these two solutions. Then:  $S_1$  covers the transition profit of an optimal  
 477 solution  $D_t$  for  $t$  odd, plus the matching profits for  $t$  odd.  $S_2$  covers the transition profit  
 478 of an optimal solution  $D_t$  for  $t$  even, plus the matching profits for  $t$  even. The ratio  $1/2$  follows.  
 479

480 *Improvement for  $T = 3$ .* The previous solutions  $S_1$  and  $S_2$  have profit (respectively) at  
 481 least  $c_1(S^*) + D_1(S^*) + c_3(S^*)$  and  $c_1(S^*) + D_2(S^*) + c_2(S^*)$ .  $S_3$  takes  $\hat{M}_1$  at step 1 and  $H'_2$   
 482 at time steps 2 and 3, with profit at least  $c_1(S^*) + D_2(S^*) + c_3(S^*)$ ;  $S_4$  takes  $H'_1$  at steps 1  
 483 and 2, and  $\hat{M}_3$  at step 3, with profit at least  $D_1(S^*) + c_2(S^*) + c_3(S^*)$ .  $S_5$  uses  $M'_2$  at the 3  
 484 steps with profit at least  $D_1(S^*) + c_2(S^*) + D_2(S^*)$  (thanks to Lemma 9). Take the best of  
 485 these 5 solutions, and the ratio follows.

486 *Improvement for  $T = 2$ .* Simply take 3 solutions:  $S_1$  is defined as previously, with profit  
 487 at least  $c_1(S^*) + D_1(S^*)$ .  $S_2$  takes  $H'_1$  at both steps with profit at least  $D_1(S^*) + c_2(S^*)$ .  $S_3$   
 488 consists of one optimal matching at step 1, and an optimal matching at step 2, with profit at  
 489 least  $c_1(S^*) + c_2(S^*)$ . The ratio  $2/3$  follows.

490

## 491 5 Concluding remarks

492 Following the results of Section 3, we leave as an open question the existence of a constant  
 493 factor approximation algorithm for the metric case for a number of time steps bigger than 3.  
 494 Also, we considered here an off-line version of the problem where the whole set of instances  
 495 is known in advance. It would be worth investigating the on-line case where data are not  
 496 known in advance.

## 497 References

- 498 1 Hyung-Chan An, Ashkan Norouzi-Fard, and Ola Svensson. Dynamic facility location via  
 499 exponential clocks. *ACM Trans. Algorithms*, 13(2):21:1–21:20, 2017.
- 500 2 Barbara M. Anthony and Anupam Gupta. Infrastructure leasing problems. In *IPCO*,  
 501 volume 4513 of *Lecture Notes in Computer Science*, pages 424–438. Springer, 2007.
- 502 3 Nicolas K. Blanchard and Nicolas Schabanel. Dynamic sum-radii clustering. In *WALCOM*,  
 503 volume 10167 of *Lecture Notes in Computer Science*, pages 30–41. Springer, 2017.
- 504 4 Niv Buchbinder, Shahar Chen, and Joseph Naor. Competitive analysis via regularization.  
 505 In *SODA*, pages 436–444. SIAM, 2014.
- 506 5 Niv Buchbinder, Shahar Chen, Joseph Naor, and Ohad Shamir. Unified algorithms for  
 507 online learning and competitive analysis. *Math. Oper. Res.*, 41(2):612–625, 2016.

- 508 **6** Edith Cohen, Graham Cormode, Nick G. Duffield, and Carsten Lund. On the tradeoff  
509 between stability and fit. *ACM Trans. Algorithms*, 13(1):7:1–7:24, 2016.
- 510 **7** David Eisenstat, Claire Mathieu, and Nicolas Schabanel. Facility location in evolving  
511 metrics. In *ICALP (2)*, volume 8573 of *Lecture Notes in Computer Science*, pages 459–470.  
512 Springer, 2014.
- 513 **8** M. R. Garey, David S. Johnson, and Larry J. Stockmeyer. Some simplified NP-complete  
514 problems. In *STOC*, pages 47–63. ACM, 1974.
- 515 **9** Michael R. Garey and David S. Johnson. *Computers and Intractability: A Guide to the*  
516 *Theory of NP-Completeness*. W. H. Freeman, 1979.
- 517 **10** Albert Gu, Anupam Gupta, and Amit Kumar. The power of deferral: Maintaining a  
518 constant-competitive steiner tree online. *SIAM J. Comput.*, 45(1):1–28, 2016.
- 519 **11** Anupam Gupta, Kunal Talwar, and Udi Wieder. Changing bases: Multistage optimization  
520 for matroids and matchings. In *ICALP (1)*, volume 8572 of *Lecture Notes in Computer*  
521 *Science*, pages 563–575. Springer, 2014.
- 522 **12** Viggo Kann. Maximum bounded 3-dimensional matching is MAX SNP-complete. *Inf.*  
523 *Process. Lett.*, 37(1):27–35, 1991.
- 524 **13** Nicole Megow, Martin Skutella, José Verschae, and Andreas Wiese. The power of recourse  
525 for online MST and TSP. *SIAM J. Comput.*, 45(3):859–880, 2016.
- 526 **14** Chandrashekar Nagarajan and David P. Williamson. Offline and online facility leasing.  
527 *Discrete Optimization*, 10(4):361–370, 2013.
- 528 **15** Baruch Schieber, Hadas Shachnai, Gal Tamir, and Tami Tamir. A theory and algorithms  
529 for combinatorial reoptimization. *Algorithmica*, 80(2):576–607, 2018.