



HAL
open science

Quantitative study of open channels by digital holography

Alexey Brodoline, Nicolas Verrier, Ludovic Depreter, Didier Felbacq, Michel Gross

► **To cite this version:**

Alexey Brodoline, Nicolas Verrier, Ludovic Depreter, Didier Felbacq, Michel Gross. Quantitative study of open channels by digital holography. Conference on Lasers and Electro-Optics Europe European Quantum Electronics Conference (CLEO/Europe-EQEC), Jun 2017, München, Germany. pp.EG_2_5, 10.1109/cleo-eqec.2017.8087591 . hal-01927223

HAL Id: hal-01927223

<https://hal.science/hal-01927223>

Submitted on 19 Nov 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Quantitative Study of open channels by digital holography

Alexey Brodoline¹, Nicolas Verrier², Ludovic Depreter¹, Didier Felbacq¹ and Michel Gross^{1*}

¹Laboratoire Charles Coulomb - UMR 5221 CNRS-Université Montpellier Place Eugène Bataillon 34095 Montpellier (FRANCE)

²Laboratoire MIPS - EA 2332 IUT de Mulhouse (GEII) 61 rue Albert Camus 68200 Mulhouse (FRANCE)

*michel.gross@umontpellier.fr

Citation: A. Brodoline, N. Verrier, L. Depreter, D. Felbacq, and M. Gross, "Quantitative Study of open channels by digital holography," in *2017 European Conference on Lasers and Electro-Optics and European Quantum Electronics Conference*, (Optical Society of America, 2017), paper EG_2_5.

European Quantum Electronics Conference 2017 : Munich Germany 25–29 June 2017

ISBN: 978-1-5090-6736-7 DOI : [10.1109/CLEOE-EQEC.2017.8087591](https://doi.org/10.1109/CLEOE-EQEC.2017.8087591)

From the session [Complex Electromagnetic Fields \(EG_2\)](#)

Open channels are peculiar incident modes of the optical field that are transmitted by a highly diffusing medium with high efficiency (100 % in the ideal case). The existence of these modes has been predicted by Dorokhov et al. [1] more than 25 years ago. Although many experiments were made to prepare these modes [2,3], the maxima of transmission obtained in experiments, are ever much lower than 100 %. To test the theory, we have considered here another Dorokhov's prediction: the average transmission of the modes that sustain the transmitted field must be $T_T=66\%$, where T_T is defined by $N_2 T_T = N_1 T$ (where T is the averaged transmission of energy, and N_1 and N_2 the number of incident and transmitted modes i.e. the size of the incident and transmitted mode basis).




Fig.1 (a) Holographic experimental setup. (b) Residual correlation of the transmitted field. The horizontal axis is SQ' . The sample transverse size is $L=70 \mu\text{m}$ red, $50 \mu\text{m}$ green, $35 \mu\text{m}$ blue.... The horizontal grey line is the residual correlation expected for $N_1=N_2$.

To verify this prediction, we have measured N_2 on a slab sample of $Z_{\text{N}}\text{O}$ powder (transverse size $L \leq 70 \mu\text{m}$, thickness $20 \mu\text{m}$) [4], by assuming that N_1 is given by the sample geometry. The experimental setup is seen in Fig. 1(a). The incident fields are random (there are scrambled by a diffusing liquid DL) and digital holography is used to measure the complex transmitted field $E(x,y)$ for the two polarisations. N_2 is measured by analysing the residual correlations of the transmitted field measured for uncorrelated incident fields.

Since the average transmission is $T=1/25$, we expect $N_2=N_1/16$. To get this optimal result, the transverse size of sample L must be infinite and the defocusing SQ' of the microscope objective MO2 must be zero (where S is the sample outgoing surface and Q' the plane imaged by MO2). Figure 1 (b) shows the residual correlation (which is proportional to $1/N_2$) as a function these two parameters. We get $N_2=N_1/12$ without extrapolation, and $N_2=N_1/16$ with extrapolation to L infinite. Here, the open channel effect is huge (factor 16) and the agreement with the Dorokhov's prediction is excellent.

We acknowledge Labex Numev (convention ANR-10-LABX-20) for funding.

References

1. O.N. Dorokhov. On the coexistence of localized and extended electronic states in the metallic phase. *Solid state communications*, 51(6): 381–384, (1984).

2. Hyeonseung Yu, Timothy R Hillman, Wonshik Choi, Ji Oon Lee, Michael S Feld, Ramachandra R Dasari, and YongKeun Park. Measuring large optical transmission matrices of disordered media. *Physical review letters*, 111(15):153902, (2013).
3. SM Popoff, A Goetschy, SF Liew, A Douglas Stone, and Hui Cao. Coherent control of total transmission of light through disorder *Physical review letters*, 112(13), 133903. (2014)
4. Verrier, N., Depraeter, L., Felbacq, D., & Gross, M. Measuring enhanced optical correlations induced by transmission open channels in a slab geometry. *Physical Review B*, 93(16), 161114. (2016)