



HAL
open science

Models projecting the fate of fish populations under climate change need to be based on valid physiological mechanisms

Sjannie Lefevre, David J. Mckenzie, Göran E. Nilsson

► **To cite this version:**

Sjannie Lefevre, David J. Mckenzie, Göran E. Nilsson. Models projecting the fate of fish populations under climate change need to be based on valid physiological mechanisms. *Global Change Biology*, 2017, 23 (9), pp.3449–3459. 10.1111/gcb.13652 . hal-01928499

HAL Id: hal-01928499

<https://hal.science/hal-01928499v1>

Submitted on 22 Nov 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

OPINION

Models projecting the fate of fish populations under climate change need to be based on valid physiological mechanisms

SJANNIE LEFEVRE¹ , DAVID J. MCKENZIE² and GÖRAN E. NILSSON¹

¹Department of Biosciences, Faculty of Mathematics and Natural Sciences, University of Oslo, Blindernveien 31, Postbox 1066 Blindern, Oslo NO-0316, Norway, ²Centre for Marine Biodiversity Exploitation and Conservation, UMR 9190 MARBEC (CNRS, IRD, IFREMER, UM), Place E. Bataillon cc 093, 34095 Montpellier, France

Abstract

Some recent modelling papers projecting smaller fish sizes and catches in a warmer future are based on erroneous assumptions regarding (i) the scaling of gills with body mass and (ii) the energetic cost of ‘maintenance’. Assumption (i) posits that insurmountable geometric constraints prevent respiratory surface areas from growing as fast as body volume. It is argued that these constraints explain allometric scaling of energy metabolism, whereby larger fishes have relatively lower mass-specific metabolic rates. Assumption (ii) concludes that when fishes reach a certain size, basal oxygen demands will not be met, because of assumption (i). We here demonstrate unequivocally, by applying accepted physiological principles with reference to the existing literature, that these assumptions are not valid. Gills are folded surfaces, where the scaling of surface area to volume is not constrained by spherical geometry. The gill surface area can, in fact, increase linearly in proportion to gill volume and body mass. We cite the large body of evidence demonstrating that respiratory surface areas in fishes reflect metabolic needs, not *vice versa*, which explains the large interspecific variation in scaling of gill surface areas. Finally, we point out that future studies basing their predictions on models should incorporate factors for scaling of metabolic rate and for temperature effects on metabolism, which agree with measured values, and should account for interspecific variation in scaling and temperature effects. It is possible that some fishes will become smaller in the future, but to make reliable predictions the underlying mechanisms need to be identified and sought elsewhere than in geometric constraints on gill surface area. Furthermore, to ensure that useful information is conveyed to the public and policymakers about the possible effects of climate change, it is necessary to improve communication and congruity between fish physiologists and fisheries scientists.

Keywords: aerobic scope, gill surface area, growth, metabolism, oxygen consumption, respiration, scaling

Received 10 November 2016; revised version received 28 January 2017 and accepted 30 January 2017

Introduction

Global change has become alarmingly rapid and the scientific community has a responsibility to provide society with valid and reliable information about potential consequences. For a planet whose surface is overwhelmingly oceans, the threats to marine ecosystems are particularly worrying, with negative impacts on fishes and fisheries that could become a pressing concern for humans (Brander, 2015). Physiology can provide a mechanistic, cause and effect, understanding of how animals might respond to a changing environment, which can be used to improve the predictive capacity of models (Jørgensen *et al.*, 2012; Peck *et al.*, 2016). We argue, however, that physiologists must play

a more active role in developing such models, to ensure that they are based upon valid underlying mechanistic assumptions.

One major projection, of obvious significance for ecosystem productivity and human food security, is that oceanic warming will lead to a systematic global decline in the size of fishes (Cheung *et al.*, 2013a). The modelling in that particular study predicted that over 2000 fish species would become up to 24% smaller by 2050. In an earlier paper based upon similar modelling, Cheung *et al.* (2011) estimated that there would be severely reduced fish catch potentials in 2050 relative to 2005. These projections have been used to explain empirical observations that adults of some marine fishes are indeed becoming smaller (Baudron *et al.*, 2014), and to suggest that climate warming will drive evolution towards smaller body sizes in fishes (Waples & Audzijonyte, 2016), and they have recently been

Correspondence: Sjannie Lefevre, tel. +4748364745, fax +4722854726, e-mail: sjannie.lefevre@imbv.uio.no

enshrined in a report from the International Union for Conservation of Nature (IUCN) (Cheung & Pauly, 2016). These model projections are, however, based on the assumption that 'growth and maximum body size in marine fish and invertebrates are determined primarily by availability of oxygen; the latter is related partly to the availability of respiratory surfaces'. We are now obliged to explain, beyond any further debate (Brander *et al.*, 2013; Cheung *et al.*, 2013b), why the physiological principles on which the modelling is based are erroneous. Most physiologists would be surprised by the suggestion that increases in oxygen demand, due to elevated future water temperatures, would constrain fishes to smaller adult sizes because their gills could not supply enough oxygen for a larger body. Here, therefore, we aim to clarify the physiological principles of gill function, metabolic rate and growth in fishes.

Gills are folded surfaces, not spheres

The key physiological assumptions about gills and metabolic rate adopted by Cheung *et al.* (2011, 2013a) were originally posited by Pauly (1981, 1998, 2010), who stated that 'Geometric constraints prevent surfaces to grow as fast as volumes' and that 'Gills, being a surface area, cannot grow, for insurmountable geometric reasons, as fast as the volume they are meant to supply with oxygen' (Pauly, 1998). As we will demonstrate, the second statement is not true: fish gills do not adhere to the geometric surface-area-to-volume relationship that is referred to in the first statement. That faulty assumption is then used to argue that gill surface areas cannot grow as fast as body mass, and that this results in lowered food conversion efficiency, when body mass increases. In fact, it is proposed that geometric constraints on the gills are what underlie the well-known allometric decrease in mass-specific oxygen uptake in relation to body mass in fishes.

The geometric relationship that prevents the surface area of an object from growing as fast as its volume would, for example, apply to oxygen uptake over a body surface. The underlying reason for this geometrical relationship is that the volume of objects like spheres increases with the cube of their radius (volume $\propto r^3$), while their surface area only increases with the square (area $\propto r^2$). The same applies to structures similar to spheres, which includes many animal bodies. As a consequence, the surface area of an animal's skin scales to the volume (or mass) of that animal with an exponent (b) approximating 0.67 (i.e. 2/3), or -0.333 ($b - 1$) for the scaling of mass-specific surface area to mass ratio (Fig. 1a). This relationship was already known to biologists by the 19th century (reviewed by Schmidt-Nielsen, 1984) and Rubner (1883) used it in a

famous attempt to explain why smaller dogs have higher mass-specific metabolic rates than larger ones, ascribing it to greater heat loss over the skin.

This simple geometric relationship does not, however, apply to fish gills. These have evolved into very large surface areas of delicate respiratory epithelium, where gas exchange takes place over many small sheets (called lamellae) attached to gill filaments that, in turn, are attached to the gill arches. Thus, gills have folded surfaces (e.g. J. W. Price, 1930) and are one of many examples in nature where the 2/3-exponent relationship between surface area and volume is overcome by folding the surface. Other examples include lungs, intestinal membranes and inner mitochondrial membranes. As each single lamella is a sheet with a half-circle to rectangular shape that grows by increasing the radius or length of the sides, rather than the thickness, it has a surface-to-mass scaling exponent (b) that approaches 1.0 ($b - 1 = 0$ for mass-specific exponent; Fig. 1a, green line). When gills grow by adding more and more lamellae, the surface-to-mass scaling exponent essentially becomes 1.0 (Fig. 1a, blue line; Fig. 1c). Thus, in morphometric studies where both total lamellar area and gill mass have been measured, a linear scaling relationship (scaling exponent of 1.0) has been found in fishes (Gehrke, 1987) as well as bivalves (Scott, 2005). Consequently, there is no geometric constraint that prevents an increase in body size (mass or volume) from being accompanied by a corresponding increase in gill mass and hence respiratory surface area. In other words, gill surface area can scale proportionally with body mass and, if it does not do so, it is because oxygen demands are reduced with body size, as discussed further below.

A simple way to explain the geometry of gills is by analogy to a book, where pages correspond to lamellae. It is easy to grasp that if you double the thickness of a book (thereby doubling its volume and mass), you can fit in twice the number of pages and, therefore, the total surface area of the pages is also doubled. Moreover, increasing the number of books (analogous to increasing the number of gill filaments) leads to a corresponding increase in page area (analogous to respiratory surface area). Thus, with the structural arrangement of books and gills, the surface area will scale to volume with an exponent of 1.0, a linear correlation. The analogy can be expanded by considering the height or width of the book, where any increase will lead to a corresponding increase in total page area. So, increasing the area of each lamella will lead to a corresponding increase in the volume occupied by the gills: still a scaling exponent of 1.0 (Fig. 1c).

An important prerequisite of the book analogy is that the distance between the lamellae, determining the




Fig. 1 Scaling of relative surface area with volume of different shapes. The volume-specific surface area (SA) of different shapes is shown as a function of volume (V) in (a). The relative surface area of the sphere (grey line) decreases fast, due to the geometry ($SA_{\text{sphere}} = 4\pi r_{\text{sphere}}^2$ and $V_{\text{sphere}} = \frac{4}{3}\pi r^3$, where r increases linearly). In a disc of the same volume as the sphere (with larger radius than height, resembling the shape of a gill lamella; green line), but with unchanging thickness and increasing radius ($r_{\text{disc}} = \sqrt{\frac{0.5V_{\text{sphere}}}{\pi h}}$), the rate of decrease in surface area ($SA_{\text{disc}} = 2\pi r_{\text{disc}}h + 2\pi r_{\text{disc}}^2$) actually falls as the radius grows. In the third scenario (blue line), when the volume is increased by increasing the number of discs, and maintaining the original thickness and radius, the surface-area-to-volume ratio is constant. For both the single disc and multiple discs, it has been taken into account that only half of the volume will be occupied (as the secondary lamellae of gills have some space between them). In (b), as the fish and, hence, the volume of the gill and the radius of the sphere grow, the surface-area-to-mass ratio decreases, assuming that the volume of the gills relative to the body mass is the same in a big compared to a small fish (at least it is not bigger). This means that larger fish have a relatively smaller respiratory surface area at their disposal, if the gills are assumed to be shaped like spheres. In (c), it is assumed, for simplicity, that as the fish grows, so does the number of respiratory 'units', the lamellae, because the thickness and distance between lamellae does not change consistently with body size (see text). This means that a larger fish could in fact have a relative surface area that is roughly the same as a smaller fish. But because there are trade-offs associated with having large respiratory surfaces, the gill surface-area-to-mass ratio scales to body mass with a mass-specific exponent ($b - 1$) slightly less than 0, guided by the oxygen demand of the fish, rather than the other way around. The curves in (a) were created using GRAPHPAD PRISM 6 (GraphPad Software Inc., La Jolla, CA, USA) after calculation of values according to the equations above, and drawings of shapes and fish were made in Microsoft PowerPoint 2010.

number of lamellae per mm of filament, is not forced by physical constraints to increase with increasing body size. There is good evidence for this assumption: Gray (1954) found no clear relationship between body mass and number of lamellae per mm filament in 31 species of marine fishes ranging in mass from 71 to 6392 g, while lifestyle had a clear influence on how lamella were packed, with 31 lamellae per mm of filament in mackerel *Scomber scombrus* compared to only 11 in the sluggish toadfish *Opsanus tau*. Likewise, there was no increase in lamellar spacing with mass in carp *Cyprinus carpio* ranging from 2 to 2000 g (Oikawa & Itazawa, 1985), and similar results were reported for porgy *Pagrus major*, a marine teleost (Oikawa *et al.*, 1999). In Nile tilapia *Oreochromis niloticus*, a small increase in lamellar spacing with body mass was associated with a relatively minor decrease in gill surface area per gill mass in larger individuals, but the scaling exponent was approximately 0.88 (Kisia & Hughes, 1992), which is far from the 0.67 predicted if a sphere-like surface-area-to-volume relationship was at play. Furthermore, there were either no, or very minor, intraspecific increases in lamellar spacing over a tenfold increase in body size in several species of tunas and billfishes (Wegner *et al.*, 2010), and Palzenberger & Pohla (1992) saw a similar pattern in freshwater fishes. In species where the interlamellar distance does increase with body mass, it could be one of the mechanisms by which gill surface area is matched to metabolic needs, but there is no evidence that it is due to physical constraints. It should also be pointed out that the thickness of the cell layers making up the diffusion barrier between water and blood does not change with body mass in ectothermic vertebrates such as fishes (most recently reviewed by Gillooly *et al.*, 2016).

Finally, if there were an upper limit for gill surface area that was reached as fishes grew, one would expect that the gill mass in relation to body mass would increase with body mass until this limit was reached, to compensate for the increasing 'geometric constraints' of the respiratory surface area. As far as we can see there is no support for this in the literature, rather the opposite. Thus, there is some evidence that gill mass, in relation to body mass, is generally smaller in larger individuals: in Nile tilapia gill mass falls from 5.8% to 2.9% of the body mass when the fish grows from 1 to 1000 g (Kisia & Hughes, 1992), and similar data have been obtained for carp (Oikawa & Itazawa, 1984) and tunas (Brill *et al.*, 2001) while, in the porgy, gill mass scales nearly linearly with body mass (Oikawa *et al.*, 1992). A likely explanation for relatively smaller gills in larger fishes is the general reduction in metabolic demands with body mass, which we will discuss further below.

Geometric constraints do not explain scaling of metabolic rate

The idea that insurmountable geometric constraints on the size of the gills could determine the metabolic rate of fishes has never, as far as we know, been pursued as a valid hypothesis among respiratory physiologists. It is, for example, not mentioned in Schmidt-Nielsen's book 'Animal Physiology: Adaptation and Environment' (1997) or in Evans and Claiborne's 'The Physiology of Fishes' (Evans & Claiborne, 2006), two books widely used as sources for overviews of animal and fish physiology. In our field, it is generally accepted that a species' oxygen demand determines the size of their respiratory surface area, not the other way around. Moreover, as mentioned by Pauly (2010), there are trade-offs involved in having large surface areas, such as passive losses of water or ions, or risk of infections. Therefore, the respiratory surface areas of fishes are in fact smaller than geometric constraints would allow. For example, maintaining acid-base and ion homeostasis will be easier for a fish with smaller gill area, because the rate of passive exchange of ions is directly proportional to the area of the surface over which the exchange occurs. The cost may be in the range of 4–10% of resting metabolic rate (Ern *et al.*, 2014). It will also cost more to ventilate a larger surface area, because of a larger resistance. The cost of ventilation is estimated to be 5–15% under resting, normoxic conditions (e.g. Holeton, 1980; Steffensen & Lomholt, 1983; Scheid, 1987; Steffensen, 1993). Still, none of these factors would be expected to increase disproportionately to gill size as a fish grows. There is, instead, very good evidence that fishes have gills sizes that are matched to their metabolic needs, which in turn are determined by factors such as habitat and life style (Gray, 1954; Kisia & Hughes, 1992; Palzenberger & Pohla, 1992; Chapman *et al.*, 2000; Killen *et al.*, 2016). Thus, fast-swimming pelagic fishes have much larger gills than sluggish benthic ones, and hypoxia-tolerant species have larger gills than less hypoxia-tolerant ones with a similar activity level. It is now also clear that gills are quite dynamic structures where the physical size of the respiratory surface can be regulated within days to months, in response to ambient factors such as oxygen levels and temperature (Chapman *et al.*, 2000; Nilsson *et al.*, 2012; Bowden *et al.*, 2014). This comes in addition to the immediate physiological responses that are available to fishes to boost oxygen uptake when required, including upregulating ventilation and cardiac output, and increasing blood oxygen affinity, the number of perfused lamellae and the number of circulating red blood cells (Perry & Gilmour, 2010). Such minute-to-minute adjustments allow fishes

to at least double their oxygen uptake, usually much more than that, with some species capable of greater than tenfold increases in uptake (Killen *et al.*, 2016; see also Fig. 2d).

Another striking argument against the suggestion that geometric constraints on gill surface area make fishes in warm water smaller than fishes in cold water

is the fact that the very largest teleosts, rays and sharks occur in tropical waters (masses given below are from fishbase.org). These are the sunfish *Mola mola*, the giant manta ray *Manta birostris* and the whale shark *Rhincodon typus*, weighing up to 2300, 3000 and 34 000 kg, respectively. The latter two are almost exclusively tropical. When it comes to highly active fishes with a great




Fig. 2 Scaling of mass-specific oxygen demand to body mass at different temperatures. The mass-specific basal oxygen demand (or 'maintenance metabolism', measured as minimum oxygen uptake (MO_{2min}) (a), decreases with a mass-specific scaling exponent of -0.13 ± 0.05 (95% confidence interval, CI), and fish at warmer temperatures have a higher MO_{2min} , at all body masses. The mass-specific maximum capacity for oxygen supply, estimated as maximum oxygen uptake (MO_{2max}) (b), decreases with body mass with an exponent of -0.12 ± 0.04 (95% CI), and fish at warmer temperatures have a higher MO_{2max} , at all body masses. The absolute scope for activities beyond basic maintenance needs, measured as absolute aerobic scope (AAS) (c) also decreases slightly with body mass, with an exponent of -0.11 ± 0.04 (95% CI). Because MO_{2max} and MO_{2min} both have a similar scaling exponent, the relative ability to increase oxygen uptake, measured as the factorial aerobic scope (FAS) (d), is maintained as body mass increases, showing that the capacity for oxygen supply does not become increasingly limited as a fish grows. Overall, the data indicate that the mass-specific scaling exponent for oxygen uptake and hence aerobic metabolism in fishes is close to -0.1 . Note the large variability in all the measures (approximately a tenfold difference at a given body mass), which is partly due to temperature effects but also to differences in species lifestyle (Killen *et al.*, 2016). The data have been sourced from the supplementary material of Lefevre (2016) and Killen *et al.* (2016), where references for all data points can be found. The colours of the points reflect the acclimation temperature of the individuals used in a given study. Green lines are fitted power relationships, while grey shades indicate 95% confidence interval for the fit, with equations and R^2 indicated in each panel. Graphs were created using the package `GGPLOT2` (Wickham, 2009) in R version 3.3.2 (R Core Team, 2016).

oxygen demand and thereby need for large gills, there are also very large tropical species, including blue marlin *Makaira nigricans* that reaches 636 kg, black marlin *Istiompax indica* (750 kg) and tiger shark *Galeocerdo cuvier* (800 kg). It has been questioned whether marlins really are tropical (Pauly, 1998), but temperature and depth loggers attached to them in tropical waters reveal a preference for warm surface waters at 26–30 °C in both Pacific and Atlantic oceans (Block *et al.*, 1992; Horodysky *et al.*, 2007; Prince *et al.*, 2005). Other gigantic tropical teleosts include the Queensland grouper *Epinephelus lanceolatus* (400 kg) and the Atlantic goliath grouper *E. itajara* (455 kg). The list of very large tropical fishes goes on and, in our view, is impossible to reconcile with any constraints on oxygen uptake as they grow.

Oxygen demand vs. oxygen supply and how they scale with body mass

Before dealing with the second assumption concerning the scaling of ‘maintenance metabolism’, it is necessary to point out some principles that most fish physiologists agree upon. First of all, the ATP used by fishes under steady state conditions is assumed to be derived mainly from aerobic metabolism, and hence, oxygen uptake is used as a proxy for metabolic rate (Nelson, 2016). Traditionally, physiologists consider two boundaries to oxygen uptake: the minimum (basal or standard) oxygen uptake and maximum oxygen uptake. Minimum oxygen uptake in fishes is measured over several hours (ideally 24–48 h, depending on the species) in a respirometer (e.g. Svendsen *et al.*, 2016), and care is taken to ensure that the oxygen uptake has stabilized before measurements are considered reliable. The fish should be in an unfed state (postabsorptive, but not starving), meaning that there is no contribution of direct costs associated with digestion and anabolism (the specific dynamic action response, discussed below), and the fish should display minimal levels of activity. Minimum oxygen uptake can also be estimated using a swim-tunnel respirometer, by extrapolating oxygen consumption at different swimming speeds down to a notional swimming speed of zero (Brett, 1964). This is particularly useful for obligatory ram-ventilating fishes, such as tunas and several sharks, which ventilate their gills by swimming with their mouth open (e.g. Brown & Muir, 1970; Wegner *et al.*, 2013). The rates of oxygen uptake obtained by these methods are considered to reflect the basal amount of oxygen needed simply to stay alive, which comprises the costs of ventilation, circulation, neuronal activity and membrane transport – generally speaking the processes that contribute to homeostasis (e.g. Chabot

et al., 2016). The minimum oxygen uptake would therefore be considered, by physiologists, to be equivalent to maintenance metabolism as defined by Pauly (1979, 2010). Maximum capacity for oxygen uptake, on the other hand, is typically estimated during swimming at high speed in a swim-tunnel or immediately after exhaustive exercise (e.g. Norin & Clark, 2016), possibly in combination with feeding (Jourdan-Pineau *et al.*, 2010). The difference between the maximum and minimum oxygen uptake is commonly referred to as the net or absolute aerobic scope (AAS), which is a measure of the oxygen that can be provided to support activities beyond basal maintenance, such as swimming, growth and reproduction.

An average interspecific scaling exponent for resting metabolism of fishes between 0.8 and 0.9 (= a mass-specific scaling exponent of –0.1 to –0.2) has now been obtained by various authors for more than half a century (e.g. Winberg, 1960; White *et al.*, 2006). We have taken the metadata on both minimum and maximum oxygen uptake, absolute aerobic scope and factorial aerobic scope (FAS, i.e. maximum divided by minimum) collected by Killen *et al.* (2016) and Lefevre (2016) to illustrate how these measures scale with body mass (Fig. 2). As expected, minimum oxygen demand scales with an exponent of 0.87 (Fig. 2a), while maximum supply capacity scales with an exponent of 0.88 (Fig. 2b), and AAS scales with an exponent of 0.89 (Fig. 2c). Consequently, FAS is maintained with increasing body mass because maximum oxygen supply capacity scales with virtually the same exponent as minimum demand (Fig. 2d). At the same time, it is clear that there is profound variation in FAS among species, which was shown by Killen *et al.* (2016) to correlate with ecology and life style. If there were insurmountable geometric constraints on the size of the gills that limited oxygen uptake in larger fishes, this would by necessity show up as a drop in FAS with increasing body mass. While the present analysis includes different species, a similar pattern has also been observed within species (Killen *et al.*, 2007; Norin & Malte, 2011, 2012; Clark *et al.*, 2012; Huang *et al.*, 2013; Mager *et al.*, 2014; Zhang *et al.*, 2014; Luo *et al.*, 2015; Tirsgaard *et al.*, 2015; Messmer *et al.*, 2016).

In contrast to these observed scaling values, the model presented in the IUCN report (Cheung & Pauly, 2016), which ultimately derives from Pauly (1979), assumes that maintenance metabolism, or more specifically the oxygen needed for maintenance, scales in direct proportion to body mass. That is, the mass-specific ‘cost of living’ is assumed to be the same for a small fish and a big fish. This is clearly not the case, as shown by the negative scaling exponent of mass-specific minimum oxygen demand, which we would argue reflects

the maintenance metabolism of fishes (Fig. 2a). Pauly (1979, 2010) offers no explanation for assuming a constant mass-specific cost for maintenance, merely stating that *'The amount of body substance degraded per unit of time, however, increases in direct proportion to body weight'* and that *'"oxygen demand" refers to the amount required or "needed" by a fish body, not to the observed O₂ consumption (or "metabolism") which is here referred to as "supply"'*. There is no mention of fishes' ability to increase their oxygen uptake in response to their demand, which of course they can do when required, as revealed by their aerobic scope.

The two assumptions – (i) that oxygen needed for maintenance, referred to as maintenance metabolism (Pauly, 2010; Cheung *et al.*, 2013a; Cheung & Pauly, 2016) or routine metabolism (Pauly, 1998), is the same for big and small fishes (as described by the hypothetical, horizontal line in Fig. 3a) and (ii) that the ability to supply oxygen falls with body mass due to mass-dependent geometric constraints on the gills – led to the conclusion that oxygen supply becomes limiting as fishes grow. Apparently the claim that maintenance (routine) metabolism scales to body mass with an exponent of 1.0 comes from the theories of von Bertalanffy, stating that *'catabolism occurs in all living cells of a fish and is therefore directly proportional to the mass of the fish's body'* (quoted by Pauly, 2010). This idea, however, has been completely discarded, as already noted by Enberg *et al.* (2008) and Brander *et al.* (2013). The oxygen consumption of different tissues in an animal vary by more than one order of magnitude and fishes are no exception (e.g. Vernberg, 1954; Itazawa & Oikawa, 1983), and the relative sizes of the various tissues and organs differ in small and large animals. Moreover, cells from small animals have higher rates of oxygen uptake than those from large ones (at least in mammals, see Porter & Brand, 1995) and, even if such measurements have yet to be made for fishes, the activity of oxidative enzymes falls with body mass in fishes (Davies & Moyes, 2007). Most importantly, the primary function of catabolism (i.e. the breakdown of proteins, fat and carbohydrates) in resting, unfed animals is to provide energy for maintenance metabolism and, as we have already explained, maintenance metabolism measured as mass-specific oxygen consumption falls with body size in resting, unfed fishes (Fig. 2a).

It bears pointing out that Weatherley *et al.* (1987) and Blier *et al.* (1997) had already concluded that there was no evidence that capacity for gas exchange or gill surface area could limit growth performance in fishes, and their analysis remains valid today. Feeding leads to a transient increase in oxygen demand in fishes (and other animals), the so-called specific dynamic action (SDA) response, which reflects the metabolic costs of

digesting and assimilating the meal, hence 'growing' (McCue, 2006). In most fishes, maximum rates of oxygen uptake exceed the highest oxygen uptake observed during an SDA response (Blier *et al.*, 1997; Fig. 2b). In salmonids, for example, peak SDA when feeding at maximum ration is about half the maximum rate of oxygen uptake during sustained aerobic exercise (Brett & Groves, 1979; McKenzie *et al.*, 2007). The database is limited; therefore, the peak SDA response in species that consume infrequent large meals may, conceivably, approach their maximum capacity for oxygen uptake (Soofiani & Hawkins, 1982; Blier *et al.*, 1997; Wang *et al.*, 2012). There is no reason to believe, however, that such constraints would become more severe as individuals increase in body mass.

The claims that there is a direct proportional increase in oxygen demand for maintenance metabolism with body mass (Fig. 3a) and that oxygen supply, in turn, cannot keep up are the two critical assumptions that underpin the conclusion by Cheung & Pauly (2016) and Cheung *et al.* (2013a): that fishes will be smaller in a warmer future. None of these assumptions are supported by current data and the principles generally recognized by physiologists. On the contrary, the evidence shows that both mass-specific minimum oxygen uptake (i.e. for maintenance) and maximum capacity for oxygen supply scale with the same exponent (ca. -0.12) and that relative scope for activity and growth are independent of body size. In conclusion, large fishes have no less capacity to increase their rate of oxygen consumption and meet the costs of growth than small fishes. Therefore, based on their respiratory capacities, we would not predict a change in the size of fishes in a warmer world (as summarized in Fig. 3b).

The importance of exponents and coefficients

In the modelling exercises by Cheung *et al.* (2011, 2013a), a scaling exponent of 0.7 for anabolic metabolism (i.e. -0.3 for the mass-specific relationship) underlying growth is used, after referring to Pauly (1981, 2010). This exponent appears to have been derived from the scaling of respiratory surface area to mass, which mirrors the exponents for minimum and maximum oxygen uptake (discussed above). A low value of this exponent in the equations underlying the model will lead to a large effect of temperature on growth and future body size (Fig. 3a). As we have argued, oxygen supply by the gills is unlikely to put constraints on metabolism and growth but, nonetheless, we would like to point out that a scaling exponent of 0.7 is considerably lower than those reported in the literature. Muir (1969) put the average exponent at 0.8–0.9 for the




Fig. 3 Predicted scaling of mass-specific oxygen supply and demand to body mass under different assumptions. In (a), which is based on the reasoning in Cheung *et al.* (2013a) and Pauly (2010), it is assumed that mass-specific ‘maintenance’ metabolism (equal to basal oxygen demand), is maintained with body mass ($MO_2 = a \cdot M_b^0$; solid purple line), while maximum capacity for oxygen supply decreases with a mass-specific exponent ($b - 1$) of -0.3 ($MO_{2supply} = c \cdot M_b^{-0.3}$; solid black line). This means that when a certain body size is reached, oxygen demand for maintenance exceeds capacity for supply and the rate of growth is constrained (the growth limit; solid blue arrow). Note that decreasing the exponent from -0.3 to -0.2 (dotted black line) has a pronounced effect on the body mass at which the growth limit is reached (dotted blue arrow). In a warmer future ($\Delta T = +2^\circ C$), MO_2 is assumed to increase with a Q_{10} of 2.4 ($MO_{2future} = MO_2 \cdot Q_{10}^{(\Delta T/10)}$; dashed purple line). It is also assumed that $MO_{2supply}$ remains the same, and hence, maintenance oxygen demand surpasses oxygen supply at a smaller body size (the future growth limit; solid red arrow). In (b) – which we argue is the condition supported by physiological evidence – it is the oxygen demand that determines the capacity for oxygen uptake, not vice versa, as assumed in (a). In (b), the mass-specific minimum oxygen demand is assumed to decrease with an exponent of -0.2 ($MO_{2min} = a_2 \cdot M_b^{-0.2}$; solid purple line) (although it is likely to be closer to -0.1 , see text and Fig. 2a), while maximum oxygen uptake is x times higher and scales with the same exponent ($MO_{2max} = c \cdot M_b^{-0.2}$ or $x \cdot MO_{2min}$; where x = factorial aerobic scope; solid orange line). Here, $MO_{2supply}$ is determined by the respiratory surface area, which in turn is guided by the maximum oxygen demand and, hence, it is slightly higher but scales with the same exponent as MO_{2max} ($MO_{2supply} = (c+d) \cdot M_b^{-0.2}$; solid black line). In this scenario, the oxygen supply that can be allocated to anabolism (growth) is somewhere between basal demands for maintenance (MO_{2min}) and maximum capacity for supply (i.e. within the aerobic scope). In a warmer future (dashed lines), MO_{2min} will increase ($MO_{2min,future} = MO_{2min} \cdot Q_{10}^{(\Delta T/10)}$), and may or may not cause a decrease in the scope for aerobic metabolism, depending on the ability of the species to increase oxygen supply ($MO_{2supply} = (c+e) \cdot M_b^{-0.2}$). Regardless, the aerobic scope will never become zero – oxygen supply is not limiting and there is no oxygen-constrained growth limit. Graphs were created in GRAPHPAD PRISM 6 using the equations above.

scaling of gill area, and subsequent estimates range from 0.76 to 0.9 (see Nilsson & Östlund-Nilsson, 2008). We expect that, had an exponent of 0.8–0.9 been used

in the modelling in Cheung *et al.* (2011, 2013a), rather than 0.7, it would have significantly reduced the future temperature effects proposed in those studies.

Moreover, we would question the use of a single temperature coefficient (Q_{10}) by Cheung *et al.* (2011, 2013a). The Q_{10} describes the factorial increase in metabolic rate with temperature. A Q_{10} of 2 means that metabolic rate doubles with a 10 °C increase in temperature, a Q_{10} of 1 means that there is no temperature effect, and values lower than 1 imply that metabolism falls with temperature. The models of Cheung and co-workers adopt a single Q_{10} of 2.4, based on the median Q_{10} of 14 studies reported in a review of Q_{10} values in fishes by Clarke & Johnston (1999). Q_{10} of many more fishes are now available, not least because of the current focus on global warming. Clarke & Johnston (1999) pointed out that Q_{10} varied from 0.45 to 3.41 among species, and they did not report how long the fishes had been allowed to acclimate. Many fishes have the ability to acclimate to a rise in temperature if given enough time, which will reduce their Q_{10} considerably. For example, in shorthorn sculpin *Myoxocephalus scorpius* exposed to a rise in temperature from 10 to 16 °C, Q_{10} fell from 2.7 to 1.0 when the acclimation time was extended from 1 to 8 weeks (Sandblom *et al.*, 2014). That is, acclimation completely abolished any temperature effect on aerobic metabolic rate. Although all fishes may not show such perfect compensation, it illustrates the problem of using a single, high Q_{10} of 2.4, as it will cause all species to have elevated metabolic rates at warmer temperatures, causing an exaggeration of unknown magnitude in the model, for an unknown number of species. For models that include Q_{10} in the parameterization, it would be pertinent to make an updated survey of published Q_{10} values, to at least estimate what proportion of species are likely to show an elevation of metabolic rate, or not, when exposed to warmer water.

Conclusions

We fully appreciate that large-scale modelling studies have focussed much-needed attention on potential effects of climate change on fishes, and we do not dispute that global warming may lead to reductions in average body size and size-at-age of fishes (see Munday *et al.*, 2008; Daufresne *et al.*, 2009; Baudron *et al.*, 2014). It is essential, however, that the correct underlying mechanisms be investigated and identified and that projections of the effects on fish populations be modelled using sound physiological knowledge and principles. The temperature effects on body size in bacteria and plankton observed by Daufresne *et al.* (2009) may very well relate to the 2/3 exponential relationship between body mass and surface area, as these organisms lack respiratory organs with a folded surface and will rely largely on their body surface for exchange of gasses with the environment. However, other

mechanisms must be at play in the French river fishes (Daufresne *et al.*, 2009) and some marine fishes (Baudron *et al.*, 2014), and this should be food for thought for physiologists and fishery biologists.

Physiologists have failed to take an active role in ensuring that influential reports, such as the current one from IUCN (Laffoley & Baxter, 2016), have a sound basis for what they promulgate as universal, physiological paradigms. This includes not just the incorrect notion that oxygen uptake limits growth in fishes, but also the hypothesis of 'Oxygen- and Capacity-Limited Thermal Tolerance' (OCLTT) (Pörtner, 2010), whose universality is far from accepted (Lefevre, 2016). Incidentally, Cheung *et al.* (2013a) refer to the OCLTT hypothesis to support their claim that '*the capacity for growth is limited by oxygen in aquatic water-breathing ectotherms*', although this hypothesis does not embrace the notion that the size of a fish is relevant for its response to warming. To ensure that accurate and useful information is conveyed to the public, about the possible consequences of climate change, there is an urgent need to improve communication and congruity between fish physiologists and fisheries scientists. If fishes are indeed becoming smaller, it is crucial that researchers from different fields collaborate to identify and understand the underlying causes, to then search for relevant solutions.

Acknowledgements

The authors would like to thank Philip L. Munday for valuable comments on an earlier draft of the manuscript. The authors also acknowledge the importance of the COST action 'Conservation Physiology of Marine Fishes' (FA1004) in providing a forum for discussion of physiology and modelling.

Conflict of interest

The authors declare no conflict of interests.

Author contributions

All authors contributed equally.

References

- Baudron AR, Needle CL, Rijnsdorp AD, Tara Marshall C (2014) Warming temperatures and smaller body sizes: synchronous changes in growth of North Sea fishes. *Global Change Biology*, **20**, 1023–1031.
- Blier PU, Pelletier D, Dutil JG (1997) Does aerobic capacity set a limit on fish growth rate? *Reviews in Fisheries Science*, **5**, 323–340.
- Block BA, Booth DT, Carey FG (1992) Depth and temperature of the blue marlin, *Makaira nigricans*, observed by acoustic telemetry. *Marine Biology*, **114**, 175–183.
- Bowden AJ, Gardiner NM, Couturier CS, Stecyk JAW, Nilsson GE, Munday PL, Rummer JL (2014) Alterations in gill structure in tropical reef fishes as a result of elevated temperatures. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology*, **175**, 64–71.

- Brander K (2015) Improving the reliability of fishery predictions under climate change. *Current Climate Change Reports*, **1**, 40–48.
- Brander K, Neuheimer A, Andersen KH, Hartvig M (2013) Overconfidence in model projections. *ICES Journal of Marine Science: Journal du Conseil*, **70**, 1065–1068.
- Brett JR (1964) The respiratory metabolism and swimming performance of young sockeye salmon. *Journal of the Fisheries Research Board of Canada*, **21**, 1183–1226.
- Brett JR, Groves TDD (1979) Physiological energetics. In: *Bioenergetics and Growth*, (eds Hoar WS, Randall DJ, Brett JR), pp. 279–352. Academic Press Inc., New York.
- Brill R, Swimmer Y, Taxboel C, Cousins K, Lowe T (2001) Gill and intestinal Na⁺-K⁺ ATPase activity, and estimated maximal osmoregulatory costs, in three high-energy-demand teleosts: yellowfin tuna (*Thunnus albacares*), skipjack tuna (*Katsuwonus pelamis*), and dolphin fish (*Coryphaena hippurus*). *Marine Biology*, **138**, 935–944.
- Brown CE, Muir BS (1970) Analysis of ram ventilation of fish gills with application to Skipjack Tuna (*Katsuwonus pelamis*). *Journal of the Fisheries Research Board of Canada*, **27**, 1637–1652.
- Chabot D, Steffensen JF, Farrell AP (2016) The determination of standard metabolic rate in fishes. *Journal of Fish Biology*, **88**, 81–121.
- Chapman LG, Galis F, Shinn J (2000) Phenotypic plasticity and the possible role of genetic assimilation: hypoxia-induced trade-offs in the morphological traits of an African cichlid. *Ecology Letters*, **3**, 387–393.
- Cheung WWL, Pauly D (2016) Impacts and effects of ocean warming on marine fishes. In: *Explaining ocean warming: Causes, Scale, Effects and Consequences* (eds Laffoley D, Baxter JM), pp. 239–253. International Union for Conservation of Nature (IUCN), Gland, Switzerland.
- Cheung WWL, Dunne J, Sarmiento JL, Pauly D (2011) Integrating ecophysiology and plankton dynamics into projected maximum fisheries catch potential under climate change in the Northeast Atlantic. *ICES Journal of Marine Science: Journal du Conseil*, **68**, 1008–1018.
- Cheung WWL, Sarmiento JL, Dunne J *et al.* (2013a) Shrinking of fishes exacerbates impacts of global ocean changes on marine ecosystems. *Nature Climate Change*, **3**, 254–258.
- Cheung WWL, Pauly D, Sarmiento JL (2013b) How to make progress in projecting climate change impacts. *ICES Journal of Marine Science: Journal du Conseil*, **70**, 1069–1074.
- Clark TD, Donaldson MR, Pieperhoff S *et al.* (2012) Physiological benefits of being small in a changing world: responses of Coho Salmon (*Oncorhynchus kisutch*) to an acute thermal challenge and a simulated capture event. *PLoS ONE*, **7**, e39079.
- Clarke A, Johnston NM (1999) Scaling of metabolic rate with body mass and temperature in teleost fish. *Journal of Animal Ecology*, **68**, 893–905.
- Daufresne M, Lengfellner K, Sommer U (2009) Global warming benefits the small in aquatic ecosystems. *Proceedings of the National Academy of Sciences*, **106**, 12788–12793.
- Davies R, Moyes CD (2007) Allometric scaling in centrarchid fish: origins of intra- and inter-specific variation in oxidative and glycolytic enzyme levels in muscle. *Journal of Experimental Biology*, **210**, 3798–3804.
- Enberg K, Dunlop ES, Jørgensen C (2008) Fish growth. *Encyclopedia of Ecology*, (ed Goldstein MI), pp. 1564–1572. Academic Press, Oxford.
- Ern R, Huong DTT, Cong NV, Bayley M, Wang T (2014) Effect of salinity on oxygen consumption in fishes: a review. *Journal of Fish Biology*, **84**, 1210–1220.
- Evans DH, Claiborne JB (2006) *The Physiology of Fishes*. CRC Press Taylor & Francis Group, Boca Raton, FL, USA.
- Gehrke PC (1987) Cardio-respiratory morphometrics of spangled perch, *Leiopotherapon unicolor* (Günther, 1859), (Percoidae, Teraponidae). *Journal of Fish Biology*, **31**, 617–623.
- Gillooly JF, Gomez JP, Mavrodiev EV, Rong Y, McIlmore ES (2016) Body mass scaling of passive oxygen diffusion in endotherms and ectotherms. *Proceedings of the National Academy of Sciences*, **113**, 5340–5345.
- Gray IE (1954) Comparative study of the gill area of marine fishes. *Biological Bulletin*, **107**, 219–225.
- Holeton GF (1980) Oxygen as an environmental factor of fishes. In: *Environmental Physiology of Fishes* (ed. Ali MA), pp. 7–32. Springer Science + Business Media, LLC, New York.
- Horodysky AZ, Kerstetter DW, Latour RJ, Graves JE (2007) Habitat utilization and vertical movements of white marlin (*Tetrapturus albidus*) released from commercial and recreational fishing gears in the western North Atlantic Ocean: inferences from short duration pop-up archival satellite tags. *Fisheries Oceanography*, **16**, 240–256.
- Huang Q, Zhang Y, Liu S, Wang W, Luo Y (2013) Intraspecific scaling of the resting and maximum metabolic rates of the Crucian Carp (*Carassius auratus*). *PLoS ONE*, **8**, e82837.
- Itazawa Y, Oikawa S (1983) Metabolic rates in excised tissues of carp. *Experientia*, **39**, 160–161.
- Jørgensen C, Peck MA, Antognarelli F *et al.* (2012) Conservation physiology of marine fishes: advancing the predictive capacity of models. *Biology Letters*, **8**, 900–903.
- Jourdan-Pineau H, Dupont-Prinet A, Claireaux G, McKenzie DJ (2010) An investigation of metabolic prioritization in the European sea bass, *Dicentrarchus labrax*. *Physiological and Biochemical Zoology*, **83**, 68–77.
- Killen SS, Costa J, Brown JA, Gamperl AK (2007) Little left in the tank: metabolic scaling in marine teleosts and its implications for aerobic scope. *Proceedings of the Royal Society B: Biological Sciences*, **274**, 431–438.
- Killen SS, Glazier DS, Rezende EL, Clark TD, Atkinson D, Willener AST, Halsey LG (2016) Ecological influences and morphological correlates of resting and maximal metabolic rates across teleost fish species. *The American Naturalist*, **187**, 592–606.
- Kisia SM, Hughes GM (1992) Estimation of oxygen-diffusing capacity in the gills of different sizes of a tilapia, *Oreochromis niloticus*. *Journal of Zoology*, **227**, 405–415.
- Laffoley D, Baxter JM (2016) *Explaining Ocean Warming: Causes, Scale, Effects and Consequences*. International Union for Conservation of Nature (IUCN), Gland, Switzerland.
- Lefevre S (2016) Are global warming and ocean acidification conspiring against marine ectotherms? A meta-analysis of the respiratory effects of elevated temperature, high CO₂ and their interaction. *Conservation Physiology*, **4**, 1–31.
- Luo Y, He D, Li G, Xie H, Zhang Y, Huang Q (2015) Intraspecific metabolic scaling exponent depends on red blood cell size in fishes. *The Journal of Experimental Biology*, **218**, 1496–1503.
- Mager EM, Esbaugh AJ, Stieglitz JD *et al.* (2014) Acute embryonic or juvenile exposure to deepwater horizon crude oil impairs the swimming performance of Mahi-Mahi (*Coryphaena hippurus*). *Environmental Science and Technology*, **48**, 7053–7061.
- McCue MD (2006) Specific dynamic action: a century of investigation. *Comparative Biochemistry and Physiology A - Molecular and Integrative Physiology*, **144**, 381–394.
- McKenzie DJ, Pedersen PB, Jokumsen A (2007) Aspects of respiratory physiology and energetics in rainbow trout (*Oncorhynchus mykiss*) families with different size-at-age and condition factor. *Aquaculture*, **263**, 280–294.
- Messmer V, Pratchett MS, Hoey AS, Tobin AJ, Coker DJ, Cooke SJ, Clark TD (2016) Global warming will disproportionately affect larger adults in a predatory coral reef fish. *Global Change Biology*. doi:10.1111/gcb.13552.
- Muir BS (1969) Gill Dimensions as a function of fish size. *Journal of the Fisheries Research Board of Canada*, **26**, 165–170.
- Munday PL, Kingsford M, O'callaghan M, Donelson J (2008) Elevated temperature restricts growth potential of the coral reef fish *Acanthochromis polyacanthus*. *Coral Reefs*, **27**, 927–931.
- Nelson JA (2016) Oxygen consumption rate v. rate of energy utilization of fishes: a comparison and brief history of the two measurements. *Journal of Fish Biology*, **88**, 10–25.
- Nilsson GE, Östlund-Nilsson S (2008) Does size matter for hypoxia tolerance in fish? *Biological Reviews*, **83**, 173–189.
- Nilsson GE, Dymowska A, Stecyk JAW (2012) New insights into the plasticity of gill structure. *Respiratory Physiology and Neurobiology*, **184**, 214–222.
- Norin T, Clark TD (2016) Measurement and relevance of maximum metabolic rate in fishes. *Journal of Fish Biology*, **88**, 122–151.
- Norin T, Malte H (2011) Repeatability of standard metabolic rate, active metabolic rate and aerobic scope in young brown trout during a period of moderate food availability. *The Journal of Experimental Biology*, **214**, 1668–1675.
- Norin T, Malte H (2012) Intraspecific variation in aerobic metabolic rate of fish: relations with organ size and enzyme activity in brown trout. *Physiological and Biochemical Zoology: Ecological and Evolutionary Approaches*, **85**, 645–656.
- Oikawa S, Itazawa Y (1984) Relative growth of organs and parts of the carp, *Cyprinus carpio*, with special reference to the metabolism-size relationship. *Copeia*, **1984**, 800–803.
- Oikawa S, Itazawa Y (1985) Gill and body surface areas of the carp in relation to body mass, with special reference to the metabolism-size relationship. *Journal of Experimental Biology*, **117**, 1–14.
- Oikawa S, Takemori M, Itazawa Y (1992) Relative growth of organs and parts of a marine teleost, the porgy, *Pagrus major*, with special reference to metabolism-size relationships. *Japanese Journal of Ichthyology*, **39**, 243–249.
- Oikawa S, Hirata M, Kita J, Itazawa Y (1999) Ontogeny of respiratory area of a marine teleost, porgy, *Pagrus major*. *Ichthyological Research*, **46**, 233–244.
- Palzenberger M, Pöhla H (1992) Gill surface area of water-breathing freshwater fish. *Reviews in Fish Biology and Fisheries*, **2**, 187–216.

- Pauly D (1979) Gill size and temperature as governing factors in fish growth: a generalization of von Bertalanffy's growth formula. Unpublished PhD Thesis, Christian-Albrechts-Universität Kiel, Kiel.
- Pauly D (1981) The relationships between gill surface-area and growth-performance in fish - a generalization of von Bertalanffy theory of growth. *Meeresforschung - Reports on Marine Research*, **28**, 251–282.
- Pauly D (1998) Tropical fishes: patterns and propensities. *Journal of Fish Biology*, **53**, 1–17.
- Pauly D (2010) *Gasping Fish and Panting Squids: Oxygen, Temperature and the Growth of Water-Breathing Animals*. International Ecology Institute Oldendorf, Germany.
- Peck MA, Arvanitidis C, Butenschön M *et al.* (2016) Projecting changes in the distribution and productivity of living marine resources: a critical review of the suite of modelling approaches used in the large European project VECTORS. *Estuarine, Coastal and Shelf Science*. doi:10.1016/j.ecss.2016.05.019.
- Perry SF, Gilmour K (2010) Oxygen uptake and transport in water breathers. In: *Respiratory Physiology of Vertebrates: Life with and Without Oxygen* (ed. Nilsson GE), pp. 49–94. Cambridge University Press, Cambridge.
- Porter RK, Brand MD (1995) Cellular oxygen consumption depends on body mass. *American Journal of Physiology - Regulatory, Integrative and Comparative Physiology*, **269**, R226–R228.
- Pörtner HO (2010) Oxygen- and capacity-limitation of thermal tolerance: a matrix for integrating climate-related stressor effects in marine ecosystems. *Journal of Experimental Biology*, **213**, 881–893.
- Price JW (1930) Growth and Gill Development in the Small-Mounted Black Bass, *Micropterus dolomieu*, Lacepede. Unpublished PhD Thesis, Ohio State University.
- Prince ED, Cowen RK, Orbesen ES, Luthy SA, Llopiz JK, Richardson DE, Serafy JE (2005) Movements and spawning of white marlin (*Tetrapturus albidus*) and blue marlin (*Makaira nigricans*) off Punta Cana, Dominican Republic. *Fishery Bulletin*, **103**, 659–669.
- R Core Team (2016) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available at: <https://www.R-project.org/> (accessed 31 October 2016).
- Rubner M (1883) Über den Einfluss der Körpergröße auf Stoff und Kraftwechsel. *Zeitschrift für Biologie*, **19**, 535–562.
- Sandblom E, Gräns A, Axelsson M, Seth H (2014) Temperature acclimation rate of aerobic scope and feeding metabolism in fishes: implications in a thermally extreme future. *Proceedings of the Royal Society of London B: Biological Sciences*, **281**, 1–9.
- Scheid P (1987) Cost of breathing in water- and air-breathers. In: *Comparative Physiology: Life in Water and on Land* (eds Dejours P, Bolis L, Taylor CR, Weibel ER), pp. 83–92. Springer Verlag, Berlin, Liviana Press, Padova, Italy.
- Schmidt-Nielsen K (1984) *Scaling: Why is Animal Size So Important?* Cambridge University Press, Cambridge.
- Schmidt-Nielsen K (1997) Energy metabolism. In: *Animal Physiology: Adaptation and Environment* (ed. Duke JB), pp. 169–216. Cambridge University Press, Cambridge.
- Scott KM (2005) Allometry of gill weights, gill surface areas, and foot biomass $\delta^{13}\text{C}$ values of the chemoautotroph-bivalve symbiosis *Solemya velum*. *Marine Biology*, **147**, 935–941.
- Soofiani NM, Hawkins AD (1982) Energetic costs at different levels of feeding in Juvenile Cod, *Gadus morhua* L. *Journal of Fish Biology*, **21**, 577–592.
- Steffensen JF (1993) Ventilatory and respiratory responses in fish: adaptations to the environment. In: *The Vertebrate Gas Transport Cascade: Adaptations to Environment and Mode of Life* (ed. Bicudo JEPW), pp. 60–71. CRC Press Taylor and Francis Group, Boca Raton.
- Steffensen JF, Lomholt JP (1983) Energetic cost of active branchial ventilation in the shark sucker, *Echeneis naucrates*. *Journal of Experimental Biology*, **103**, 185–192.
- Svendsen MBS, Bushnell PG, Steffensen JF (2016) Design and setup of intermittent-flow respirometry system for aquatic organisms. *Journal of Fish Biology*, **88**, 26–50.
- Tirsgaard B, Behrens JW, Steffensen JF (2015) The effect of temperature and body size on metabolic scope of activity in juvenile Atlantic cod *Gadus morhua* L. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology*, **179**, 89–94.
- Vernberg FJ (1954) The respiratory metabolism of tissues of marine teleosts in relation to activity and body size. *Biological Bulletin*, **106**, 360–370.
- Wang Q, Wang W, Huang Q, Zhang Y, Luo Y (2012) Effect of meal size on the specific dynamic action of the juvenile snakehead (*Channa argus*). *Comparative Biochemistry and Physiology - Part A: Molecular and Integrative Physiology*, **161**, 401–405.
- Waples RS, Audzijonyte A (2016) Fishery-induced evolution provides insights into adaptive responses of marine species to climate change. *Frontiers in Ecology and the Environment*, **14**, 217–224.
- Weatherley AH, Gill HS, Casselman JM (1987) Metabolism and growth. In: *The Biology of Fish Growth* (eds Weatherley AH, Gill HS, Casselman JM), pp. 51–100. Academic Press, Cambridge, MA.
- Wegner NC, Sepulveda CA, Bull KB, Graham JB (2010) Gill morphometrics in relation to gas transfer and ram ventilation in high-energy demand teleosts: scombrids and billfishes. *Journal of Morphology*, **271**, 36–49.
- Wegner NC, Sepulveda CA, Aalbers SA, Graham JB (2013) Structural adaptations for ram ventilation: gill fusions in scombrids and billfishes. *Journal of Morphology*, **274**, 108–120.
- White CR, Phillips NF, Seymour RS (2006) The scaling and temperature dependence of vertebrate metabolism. *Biology Letters*, **2**, 125–127.
- Wickham H (2009) *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag, New York.
- Winberg G (1960) Rate of metabolism and food requirements of fishes. *Fisheries Research Board of Canada Translation Series*, **194**, 1–202.
- Zhang Y, Huang Q, Liu S, He D, Wei G, Luo Y (2014) Intraspecific mass scaling of metabolic rates in grass carp (*Ctenopharyngodon idellus*). *Journal of Comparative Physiology B*, **184**, 347–354.